Lavinia Gambelli[1,2], Benjamin Meyer[3], Mathew McLaren[1,2], Kelly Sanders[1,4], Tessa E.F. Quax[5],

Vicki Gold[1,4], Sonja-Verena Albers[5] and Bertram Daum[1,4]*

[1] Living Systems Institute, University of Exeter, Exeter, United Kingdom

[2] College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, United Kingdom

[3] Molecular Enzymology, Faculty for Chemistry, University of Duisburg-Essen, Germany

[4] College of Life and Environmental Sciences, University of Exeter, Exeter, United Kingdom

[5] Institute of Biology II, Molecular Biology of Archaea, University of Freiburg, Freiburg, Germany

*Corresponding author

12 Architecture and modular assembly of the *Sulfolobus* S-layers

13 revealed by electron cryo-tomography

15 Abstract

Surface protein layers (S-layers) often form the only structural component of the archaeal cell wall

and are therefore important for cell survival. S-layers have a plethora of cellular functions including

maintenance of cell shape, osmotic and mechanical stability, the formation of a semi-permeable

protective barrier around the cell, cell-cell interaction, as well as surface adhesion. Despite the

central importance of the S-layer for archaeal life, their three-dimensional architecture is still

poorly understood. Here we present the first detailed 3D electron cryo-microscopy maps of

archaeal S-layers from three different *Sulfolobus* strains. We were able to pinpoint the positions

and determine the structure of the two subunits SlaA and SlaB. We also present a model

describing the assembly of the mature S-layer.

26 Introduction

Many bacteria and archaea are surrounded by an outermost layer – the S-layer – which is

composed of glycosylated surface proteins. These proteins arrange into flexible, porous, yet

highly stable lattices that form cage-like coats around the plasma membrane. In bacteria, S-layers

are anchored to the peptidoglycan or the outer membrane. In archaea, S-layers can be either

incorporated into the periplasmatic polysaccharide layers such as pseudomurein and

methanochondroitin, or simply integrated into the cytoplasmic membrane. In most cases, S-layers

form ordered 2-dimensional arrays and serve a variety of functions, which are thought to be

specific to genera or groups of organisms sharing the same environment (Albers & Meyer, 2011;

Engelhardt, 2007; Rodrigues-Oliveira, Belmok, Vasconcellos, Schuster, & Kyaw, 2017).

For archaea, S-layers are of particular importance as they often comprise the only cell wall

component. They therefore define cellular shape and provide osmotic, thermal and mechanical

stability (Engelhardt, 2007). In addition, *in vitro* experiments have shown that S-layers change the

physical and biochemical properties of lipid layers, rendering them less flexible, less fluid, more

stable and heat-resistant, and possibly more resistant to hydrostatic pressure. Furthermore, it has

been suggested that S-layers provide protection against immunological defence systems and

viruses, act as pathogenic virulence factors, serve as phage receptors, promote surface adhesion,

establish a quasi-periplasmic space, provide anchoring scaffolds for membrane proteins,

sequester ions and facilitate biomineralization (Engelhardt, 2007). S-layers are intrinsically

capable of self-assembly *in vitro*, resulting in tube-like, spherical two-dimensional crystals

(Dietmar Pum & Sleytr, 2014). S-layers are therefore highly interesting for various applications in

nanotechnology (Sleytr et al., 2014).

Sulfolobus is a genus of hyperthermophilic acidophilic archaea, which grow in low-pH terrestrial

hot springs at 75–80°C all around the globe. They are well-established model organisms, as they

can be relatively easily cultured in the laboratory and the genomes of a number of strains have

been sequenced (Chen et al., 2005; Kawarabayasi et al., 2001; She et al., 2009). The S-layer of

all *Sulfolobus* species consists of two proteins, SlaA and SlaB. Biochemical analysis of SlaA and

SlaB from different *Sulfolobus* species revealed molecular masses ranging from 120kDa - 180kDa

or 40kDa - 45kDa, respectively (Grogan, 2010; Veith et al., 2009). Comparative sequence

analysis and molecular modelling of SlaB revealed that it exists in two species-dependent

variants. In *S. ambivalens*, *S. acidocaldarius*, *S. tokodaii* and *S. sedula,* SlaB is comprised of an

N-terminal Sec-dependent signal sequence, followed by three consecutive beta-sandwich

domains, an alpha-helical coiled coil domain and one C-terminal transmembrane helix. In

contrast, the sequences of *S. solfataricus* and *S. islandicus* are shorter by one beta-sandwich

domain (Veith et al., 2009). Interestingly, it was recently shown by negative stain electron

microscopy (EM) that SlaB knockout strains of *S. islandicus* still assemble partial S-layers (Zhang

et al., 2018).

In contrast, SlaA was predicted to be a soluble protein rich in beta-strands and to form the outer
region of the *Sulfolobus* S-layer (Veith et al., 2009). Deletion mutants of SlaA led to deformed
cells without a distinctive cell envelope (Zhang et al, 2018).

So far, the structure of the *Sulfolobus* S-layer has only been inferred by early 2D crystallography
of negatively stained and isolated S-layers. These data suggested that Sulfolobus *S-layers* adopt
a structurally conserved lattice with P3 symmetry, which encompasses 4,5 nm triangular and 8
nm hexagonal pores at a 21 nm distance (Baumeister & Lembcke, 1992; Taylor, Deatherage, &
Amos, 1982). However, as detailed 3D maps were so-far unavailable, it has been unclear how
SlaA und SlaB assemble into the final S-layer structure.

Using electron cryo-tomography (cryoET) and sub-tomogram averaging (STA), we obtained the
first 3D cryoET maps of S-layers from three *Sulfolobus* species at unprecedented resolution.
Through difference maps of fully-assembled and SlaB-depleted S-layers, we were able to
unambiguously pinpoint the positions of the component subunits SlaA and SlaB. Based on these
experiments, we present a 3D model describing their assembly. In addition, our data reveal that
strain-specific variants of SlaB lead to marked differences in the outward-facing S-layer surface.
We also find that SlaB is not required for SlaA S-layer assembly *in vitro*, which has important
implications about the function of both S-layer components.

Results

In situ structure of the *Sulfolobus* S-layer

To obtain a first detailed understanding of the molecular architecture of the archaeal S-layer, we
chose the *Sulfolobus* species *S. islandicus (Sisl)*, *S. solfataricus (Ssol)* and *S. acidocaldarius*
(Saci). In *Sisl* and *Ssol* the S-layer subunit proteins SlaA and SlaB show share 87.4% and 87.7%
sequence identity (Fig. 1S1) and are thus virtually identical. In contrast, SlaA and SlaB from *Saci*
show only ~24% or ~25 % identity with *Sisl / Ssol* (Fig. 1S2) and are thus far less conserved
between those species. Moreover, SlaB from *Saci and Sisl/Ssol* have previously been proposed
to represent two different structural "families" with respect to SlaB, which exists as a long form in
Saci and a short form in *Ssol* and *Sisl* (Veith et al., 2009). We hypothesised that the two different
variants of SlaA and SlaB may cause distinct S-layer geometries in *Saci* when compared to *Sisl*
or *Ssol*.

To prepare cells for cryoET, cellular suspensions were plunge-frozen on holy carbon grids and
investigated in the electron microscope. The majority of cells were surrounded by intact S-layers

and membranes and showed various degrees of cytoplasmic density, which may either be a result
of different metabolic states of slight cytosolic leakage during the sample preparation procedure.
Tomographic tilt series were collected of cells with low cytosolic density, which were more
transparent to the electron beam and thus resulted in tomograms with better signal-to-noise ratio
(Fig. 1A).
105
In tomographic reconstructions, cells appeared disk-shaped with a diameter of up to 2 μm and a
thickness of 250-300 nm (Fig. 1A). Since *Sulfolobus* cells are usually roughly spherical, this
suggests that the cells had been compressed due to surface tension of the buffer during the
plunge freezing procedure. In tomographic cross-sections, two layers confining the cells were
distinguished (Fig. 1A). Whereas the inner layer, the membrane, was smooth, the outer S-layer
had a corrugated appearance (Fig. 1B).
Tomographic sections parallel to the plane of the S-layers clearly showed that they indeed form
regular two-dimensional arrays, as confirmed by power spectra of the respective tomographic
slices. These power spectra showed clear spots up to the third order and indicated a lattice with
P3 symmetry (Fig. 1S3). The fuzzy spots indicated that the S-layers were not perfectly crystalline.
This was expected, as roughly round shapes cannot be contained in a hexagonal lattice unless
defects are included (D Pum, Messner, & Sleytr, 1991). Moreover, gaps in the lattice are needed
to accommodate surface filaments such as archaella (Daum et al., 2017) or to allow the cells to
grow and divide. In tomographic slices perpendicular to the membrane plane, S-layers formed
regular, corrugated canopy-like arrays at a centre-to-centre distance of ~30 nm from the
membrane (Fig. 1 B).
122
To obtain structural information of these S-layers in situ, 1809 subvolumes (*Sisl*) and 2068
subvolumes (*Saci)* were cut from tomograms, in which the cell surface was clearly resolved.
Subsequently, subvolumes were aligned and averaged in PEET. This resulted in 3D maps at ~30
Å and 28 Å resolution for *Sisl* and *Saci*, respectively (Fig. 1S5). Both S-layer maps revealed a
perforated two-dimensional protein lattice with P3 symmetry (Fig. 1; 1S4; 1S5). For *Sisl*, unit cell
dimensions were ~21.9 x 20.9 nm including an angle of 120° and for *Saci* the unit cell measured
~23.9 x 23.6 nm with an angle of 120°. The unit cell of the *Saci* S-layer is therefore roughly 10 %
larger than that of *Sisl.* The total height of the maps was ~25 nm for *Sisl* and ~28 nm for *Saci*,
respectively (Fig. 1 D-G).
132

Our 3D maps revealed structurally conserved and unique features for both S-layers (Fig. 1 D-G).
In particular, the organisation of the inward-facing domains of the S-layer of both species are
generally the same. In either case, they consist of trimeric stalks (Fig. 1 blue and yellow). These
stalks support an outer canopy-like assembly (Fig. 1 D-G, green and red). At their membrane-
proximal parts, each of the trimeric stalks merge into a single protrusion, giving them a tulip-like
shape (Fig. 1 E, G blue and yellow). This protrusion most likely forms the membrane anchor of
the S-layer. Distal to the membrane, the stalks are connected to the S-layer canopy via their
trimeric domains.
141
The outer canopy forms the outward-facing side of the S-layer and resides on top of the array of
the trimeric stalks (Fig. 1 D-G). In *Sisl* and *Saci* the base of the canopy appears to be cross-linked
by a protein network, which has roughly the same in-plane geometry in both species (Fig. 1 D-G,
green). In cross-section perpendicular to plane of the S-layers, this network forms a continuous
band throughout the entire structure and includes repeating triangular and hexagonal pores. Two
types of triangular pores can be distinguished, of which one is situated on top of a trimeric stalk
(Fig. 1, E, G blue) while the other is not. The triangular pores with and without stalk alternate
throughout the lattice (Fig. 1 D-G).
150
151   The outward-facing part of the canopy is distinct in both species. In *Sisl,* it is composed of an
152   array of hexameric cone-shaped protrusions (Fig. 1 G, red). Each of the cones encloses a
153   hexagonal pore within the membrane. At their base, each cone is surrounded by six neighbouring
154   cones. The spaces between three cones contain the triangular pores (Fig 1 G).
155
156   In *Saci*, the most distal part of the outer layer is composed of elongated, petal-like structures (Fig.
157   1 D, red). As in *Saci* these structures do not protrude as far from the general plane of the canopy,
158   the outer S-layer face appears rather smooth compared to that of *Sisl*. As seen in Fig. 1 D, each
159   of these structures show a 2-fold symmetry around their long axis, which suggests that they are
160   dimers.
161

Structural dissection of *Sulfolobus* S-layers
To obtain higher resolution maps for *Sulfolobus* S-layers and to investigate if the cell context is
important to maintain correct S-layer assembly, we isolated S-layers – this time from *Saci* and
Ssol, the latter of which was by sequence alignments predicted to have the same shortened
domain structure as *Sisl.* Tomograms were recorded at optimised imaging conditions and sub-

tomogram averaging resulted in more detailed maps (Fig. 2) at 21 Å (*Saci*) and 16 Å (*Ssol*)
resolution (Fig. 1S5). The lattice parameters of the *Saci* S-layer remained unchanged, showing
that their integrity does not crucially depend on the cellular context (Fig. 2 A). Moreover, the map
of the *Ssol* S-layer was virtually identical to the one of *Sisl* in terms of lattice dimensions and outer
canopy topology (Fig. 2 B), indicating that 87 % sequence identity conserves the general
structural features.

To further investigate the structural differences between *Saci* and *Ssol*, we segmented each map
into units composed of three stalks and the adjacent canopy (Fig 2 C, D). As with *Sisl*, the S-layer
of *Ssol* clearly showed a smaller unit size of 10 % when compared to the one of *Saci*. Furthermore,
structural differences surrounding the hexagonal pores were resolved more clearly. Whereas in
Saci each hexagonal pore is flanked by six inward-curled protein domains, it is surrounded by six
outward projecting densities in *Ssol,* giving rise to the cone-like assemblies*.*

Pinpointing SlaA and SlaB
In order to locate SlaA and SlaB within the *Sulfolobus* S-layer, we first removed the subunit SlaB
from isolated S-layers using the detergent N-laurylsarcosine (Fig. 3 A). After 3-4 repeated
washing steps, this subunit ceased to be detectable by SDS PAGE. CryoET and sub-tomogram
averaging of the SlaB-depleted S-layer resulted in a map that lacked the trimeric stalks (Fig 3 C),
leaving behind the porous canopy. This clearly indicates that SlaB constitutes the trimeric stalks
that anchor the S-layer in the membrane. Consequently, the canopy must be formed by SlaA,
which is in line with previous predictions (Veith et al., 2009).
To visualise the location and architecture of SlaB unambiguously, we calculated a difference map
by subtracting the SlaB-depleted from the fully assembled map (Fig. 3 D). This revealed that SlaB
adopts a tripod-like shape, which is consistent with earlier sequence-based predictions that
suggested that SlaB foms a trimer (Veith et al., 2009). In our structure, the three branches of each
SlaB trimer are buried inside the SlaA canopy, whilst the "monomeric" stalk projects away from it,
towards the membrane plane. In multitude, these pillars act to raise the SlaA canopy above the
membrane. Interestingly, individual SlaB units are not in contact, but are linked to each other via
the SlaA canopy network. Notably, the canopy structure of the SlaB-depleted S-layer was less
well resolved than in the fully-assembled control sample (Fig. 3 B). This suggests that SlaB may
act in reinforcing the stability of the SlaA network.

Role of SlaB in S-layer assembly

As the S-layer structure is maintained after removal of SlaB, we asked if SlaB proteins merely function as pillars for the SlaA lattice or if they also aide correct assembly of the outer canopy. To investigate this, we isolated S-layers from both *Saci* and *Ssol*, disassembled them by transfer into pH 10 buffer and subsequently performed recrystallisation by dialysis against water at pH 7 and incubation for 120 h. Inspection of the reaction containing both subunits in the electron microscope revealed patches of reassembled S-layers (Fig. 4 A, D). Sub-tomogram averaging showed that these S-layers had retained their original lattice dimensions (Fig. 4 B, E) and that the SlaA canopy as well as SlaB stalks could be identified (Fig. 4 C, F).

Interestingly, electron microscopy of the SlaB-depleted samples also revealed newly-formed S-layers (Fig. 4 G, H, J, K). Again, sub-tomogram averaging confirmed that the reassembled S-layers had adopted their original lattice parameters, however this time without the SlaB stalks (Fig. 4 I and L). These observations indicate that SlaB is not involved in assembly process of the SlaA canopy and suggests that it mainly functions as a membrane anchor and distance ruler that determines the width of the periplasmic space.

Assembly model of the *Sulfolobus* S-layer

It has previously been shown biochemically and confirmed by us that SlaA forms a stable ~250 kDa dimer in solution (Veith et al., 2009). In conjunction with early electron microscopy, it has also been predicted that this dimer forms the primary *Sulfolobus* S-layer building block (Taylor et al., 1982; Veith et al., 2009). This repeating dimer-array is apparent in our sub-tomogram averages of negatively-stained SlaA-only S-layers (Fig. 4 I). To build a cryoEM model of the *Saci* and *Ssol* S-layers, we first segmented the SlaB-depleted sub-tomogram averaging maps by the water-shedding method in Chimera (Pettersen et al., 2004). For guidance, we used the constraints of repeating dimeric SlaA densities that roughly fitted the molecular weight of 250 kDa, as well as our negative stain map in which the individual dimers were clearly visible (Fig. 4 I). We then superimposed this segmentation with the fully-assembled S-layer, which provided us with the positions of SlaB trimers.

The resulting model shows the location of SlaA dimers and SlaB trimers within the assembled S-layers of *Saci* (Fig. 5 A-C) and *Ssol* (Fig. 5 D-F). In both cases, individual SlaA dimers adopt the shape of a boomerang. The elbow of each boomerang interacts with one of the three branches of a SlaB trimer, whereas both SlaA arms project outward. Thus, each SlaB trimer is bound to

three SlaA dimers (SlaB$_3$/3SlaA$_2$), which intersect to form a triangular pore above each SlaB

trimer. Three of these flower-shaped SlaB$_3$/3SlaA$_2$ surround one hexagonal pore, formed by six

SlaA dimers. Herein, each of the three SlaB trimers contributes 2 SlaA dimers to the hexameric

ring. At the same time, each SlaA dimer spans two neighbouring hexameric pores. The second

triangular pore type that does not coincide with a SlaB trimer is formed by three intersecting SlaA

dimers at the interface of three hexagonal pores. In total, each SlaA dimer interacts with four SlaA

dimers and 3 SlaB trimers. Therefore, SlaA dimers interlink into a tightly woven S-layer canopy

that is stabilised by multiple interactions.

SlaA determines S-layer geometry and topology

To understand how *Saci* and *Ssol* assemble S-layers with similar geometry but different topology

and unit cell size, we compared the structures of the SlaA proteins from each species. To this

end, we extracted individual dimer densities from our segmented maps shown in Fig. 5.

This comparison revealed that while the overall shape of the dimers is the same, there are marked

differences with respect to their horizontal and vertical dimensions, as well as the angles between

both arms of each boomerang (Fig 6). *Saci* SlaA measures 23 nm along its long axis, which

corresponds with the size of the unit cell of the S-layer (Fig 6A). This is to be expected, as this

molecule spans to neighbouring hexagonal S-layer pores (Fig. 6C). The corresponding portion of

Ssol SlaA measures ~20 nm (Fig. 6B), which is again in accordance with the 10% smaller

assembled array (Fig 6D). The length of the SlaA dimer therefore determines the size of the unit

cell. The differences in length appear to be established by the angle between both arms of the

molecule, which is 102° for the longer *Saci* SlaA and 96° for the shorter *Ssol* homolog.

Interestingly, both SlaA variants also differ in an apical domain, which is curled inwards in *Saci*

and projects upwards in *Ssol*. In assembly, these domains are responsible for the differences in

the topology of both S-layers, as it is these domains that delineate the shallow hexagonal pores

in S*aci* and the conical protrusions in *Ssol* (Fig. 6 C, D). Taken together, the strain-specific

dimensions and surface morphology of *Sulfolobus* S-layers are determined by unique structural

characteristics of SlaA, in particular its apical domains.

Discussion

Roles of SlaA and SlaB in S-layer assembly

In this study, we describe the first cryoET-based molecular maps of archaeal S-layers. These maps unambiguously show that the *Sulfolobus* S-layer consists of trimeric SlaB proteins that adopt tripod-like structures and act as pillars that anchor the outer S-layer canopy in the cell membrane. The building block of the S-layer canopy is a dimer of SlaA, which assembles into a porous array of P3 symmetry. These observations are consistent with earlier predictions based on sequence analysis, biochemistry and early negative-stain electron microscopy maps (Baumeister & Lembcke, 1992; Taylor et al., 1982; Veith et al., 2009).

While X-ray structures for neither of the component proteins are available, sequence predictions indicated that SlaA and SlaB contain Sec-dependent signal peptides and are thus translocated through the membrane by the general secretory pathway. In addition, glycoproteomic analysis showed that both proteins contain several sites for N- and O-linked glycosylation and are heavily glycosylated prior to assembly (Palmieri et al., 2013; Peyfoon et al., 2010) (Fig. 7). Moreover, SlaB has been predicted to consist of an N-terminal membrane-anchor, followed by coiled-coil domain including a serine and threonine-rich highly-glycosylated region (ST linker) and 2-3 C-terminal beta sandwich domains (Veith et al., 2009) (Fig. 7). Upon trimerisation of SlaB, three C-termini presumably form a trimeric coiled-coil, which most likely corresponds to the stalk-like base of each SlaB tripod (Fig. 7). The SlaB stalks in our maps appear shorter than expected for a coiled-coil of roughly 100 amino acids (Veith et al., 2009) in length. This is likely due to high flexibility in this region, which is thus mostly averaged out during our sub-tomogram averaging procedure. The three protrusions of each SlaB tripod are likely formed by the predicted C-terminal beta-sandwich domains (Veith et al., 2009) of the three SlaB proteins in the trimer, which therefore also form the interface with the overarching SlaA lattice (Fig. 7). However due to the limited resolution of our maps, we were unable to distinguish between the length differences in the *Saci* and *Ssol* SlaB beta-sandwich domains.

Our reassembly experiments reveal that SlaB is not required for SlaA self-assembly *in vitro*. This is consistent with recent whole-cell negative stain EM data, which showed that *Sisl* SlaB knockout strains still assemble partial S-layer like coats (Zhang et al., 2018). We therefore conclude that the SlaA protein is the driving factor of S-layer assembly. In addition, our data shows that strain-specific S-layer surface features including lattice dimensions and topology, are determined by the shape of SlaA.

In contrast, SlaB's main function is to serve as a membrane anchor, distance ruler and to create
the periplasmic space. This space has been proposed to have important roles in archaeal cell
biology, including the coordination of membrane-embedded molecular machines, such as the
archaellum (Banerjee et al., 2015; Daum et al., 2017) and various other archaeal surface
structures (Chaudhury, Quax, & Albers, 2018). Secondarily, we observe that SlaA-only S-layers
are slightly less ordered than their fully-assembled counterparts, indicating that SlaB also
increases the stability of the S-layer, presumably by crosslinking $3SlaA_2$ subcomplexes with each
other.

The functional significance of the different S-layer topologies is largely unknown and awaits
further exploration. It may be speculated that distinct S-layer surface structures have evolved to
modulate the cell-adhesive properties of different archaeal strains. In addition, S-layers may act
as receptors in cell-cell recognition or phage infection (Plaut et al., 2014) and we could show
previously that archaeal viruses evolved elaborate cell entry and egress strategies to overcome
the S-layer barrier (Daum et al, 2013; Quemin et al, 2014; Quax & Daum 2018). It is therefore
conceivable that distinct S-layer topologies provide unique recognition tags for strain-specific
interaction and communication and that the evolution of new structures is a manifestation of the
cell's strategy to avoid viral infection (Tschitschko et al, 2015). SlaB, in contrast, appears
structurally more conserved than SlaA. Indeed, it has been shown previously that SlaB proteins
have lower sequence variability than SlaA and their molecular masses differ less across different
Sulfolobus strains (Veith et al., 2009). This is likely due to the fact that SlaB interacts less with the
extracellular medium and is thus less prone to environmentally-related evolutionary pressures.

Recently, an additional paralogous SlaB gene product (M164_1049, SSO1175 or
WP_011278654.1 in *SisI*, *Ssol,* or *Saci*, respectively) was suggested to form another membrane-
anchored protein component of the *Sulfolobus* S-layer (Zhang et al, 2018). Double M164_1049
and SlaB mutants showed that no S-layer is present in *SisI*, as opposed to single SlaB mutants
where a partial coat of SlaA was observed. In addition, a single M164_1049 deletion mutant
showed virtually no difference compared to the wild type. It was concluded that M164_1049 may
aid in anchoring SlaA to the cytoplasmic membrane (Zhang et al., 2018). Based on our maps, we
cannot confirm the presence of this protein. As it is membrane-anchored, it likely resides near the
membrane plane, the region which is averaged out in our data due to the highly flexible nature of
the SlaB coiled-coil domain.

In contrast to *Sulfolobus,* many archaeal strains possess only one S-layer protein. For example,

in many haloarchaea, a single S-layer glycoprotein (gene product *csg1* / SLG protein in *Haloferax*

volcanii) combines membrane-anchoring and canopy-forming functions into one protein.

Moreover, this protein is not membrane-integral but instead inserted to the bilayer via a C-terminal

lipid anchor (Fig. 7) (Pohlschroder, Pfeiffer, Schulze, & Halim, 2018). While employing only one

S-layer protein might be energetically more favorable, it is likely that using two increases the

adaptability of the S-layer surface (SlaA) without compromising the membrane anchor (SlaB).

Function of S-layer pores

One of the most striking features of the *Saci* and *Ssol* S-layers are the circular apertures of ~4.5

or ~8 nm in diameter that repeat throughout the array. Similar porous, skeleton-like architectures

have been reported for other S-layers such as that of the bacterium *Caulobacter cresentus*

(Bharat et al., 2017), highlighting that S-layer porosity is a highly conserved trait throughout

bacterial and archaeal species and thus provides a significant evolutionary advantage. This poses

the question what function these openings may convey. It is likely that – similar to chainmail - this

skeleton-like structure enhances S-layer flexibility, enabling it to coat cell bodies of various shape

and allowing the cell to morph and divide. It is also tantalising to assume that the particular shape

and size of the pores has evolved to accommodate particular cellular filaments, such as adhesive

pili or archaella. In contrast to this notion, we find that most known archaeal surface filaments,

such as adhesive (AAP) pili (Braun et al., 2016; Henche et al., 2012) and archaella (Daum et al.,

2017; Poweleit et al., 2016) are over 10 nm in diameter and thus too wide to be accommodated

by S-layer pores. Thus, the S-layer will have to be partially disassembled or adopt a different local

geometry wherever these filaments emerge from the cell body (Daum et al., 2017). Finally, it is

safe to assume that due to their porosity, S-layers provide a semi-permeable barrier, similar to

the outer membrane of Gram-negative bacteria, albeit with a more liberal molecular weight cut-

off. At the first glance, this cut-off appears to be determined by the pore diameter (<8nm), which

would allow a large variety of solutes, macromolecules and even large proteins to pass. However,

this notion may be deceptive, as both SlaA and SlaB are highly glycosylated (Palmieri et al., 2013;

Peyfoon et al., 2010). While (likely due to their high flexibility) these posttranslational modifications

are averaged out in our cryoEM maps, they are thought to cover much of the S-layer surface and

thus possibly also project into the pores (Fig. 7). It is conceivable that these glycans would

significantly lower the permeability of the S-layer pores to macromolecules, similar to the hydrogel

found in nuclear pore complexes (D'Angelo & Hetzer, 2008).

Conclusions

We present the first detailed 3D models of S-layers of three different *Sulfolobus* strains and pinpoint the location and organisation of their component subunits SlaA and SlaB. We find that the structure of the SlaA dimer determines the unit cell size and topology of the S-layer, whilst SlaB anchors the S-layer in the plasma membrane and defines a pseudo-periplasmic space. Different S-layer topologies between species are likely the result of various evolutionary adaptations, including cell-cell interactive specificity and adhesive properties. Despite being an ubiquitous hallmark in archaea, many functional aspects of S-layers still need to be clarified through further structural investigation.

Materials and Methods

Strains and growth conditions

The strains *Ssol* and *Saci* MW001 were grown in basal Brock medium* at pH 3 (Brock, 1972). The medium was supplemented with 0.1 % (w/v) NZ-amine and 0.2 % (w/v) dextrin just before inoculation. For growth of *Saci* MW001, 10 µg/ml uracil was added to the medium. *Sulfolobus* cultures were grown at 75°C, 150 rpm, until an OD600 of > 0.6 was reached. Cells were harvested by centrifugation at 5000 x g (Sorvall ST 8R) for 30 min and stored at -20°C for subsequent use.

* Brock media contains (amount per litre): 1.3 g $(NH_4)_2SO_4$, 0.28 g KH_2PO_4, 0.25 g $MgSO_4 \times 7H_2O$, 0.07 g $CaCl_2 \times 2H_2O$, 0.02 g $FeCl_2 \times 4H_2O$, 1.8 mg $MnCl_2 \times 4H_2O$, 4.5 mg $Na_2B_4O_7 \times 10H_2O$, 0.22 mg $ZnSO_4 \times 7H_2O$, 0.05 mg $CuCl_2 \times 2H_2O$, 0.03 mg $NaMoO_4 \times 2H_2O$, 0.03 mg $VOSO_4 \times 2H_2O$ and 0.01 mg $CoSO_4 \times 7H_2O$.

S-layer isolation

Cell pellets of frozen cells from a 50 ml culture were incubated and inverted on a rotator at 40 rpm (Stuart SB3) for 45 min at 37°C, in 40 ml of buffer A (10 mM NaCl, 1 mM phenylmethylsulfonyl fluoride, 0.5 % sodium lauroylsarcosine), with the addition of 10 µg/ml DNAse I just prior to use. The samples were pelleted by centrifugation at 18.000 x g (Sorvall Legend XTR) for 30 min, and subsequently resuspended in 1.5 ml buffer A, before further incubation and inversion at 37°C, for 30 min. After centrifugation at 14.000 rpm for 30 min (Sorvall ST 8R), the pellet was purified by resuspension and incubation in 1.5 ml buffer B (10 mM NaCl, 0.5 mM $MgSO_4$, 0.5% SDS) and rotated for 20 min at 37°C, 40 rpm. To retain both SlaA and SlaB, only one wash was performed.

402 To remove SlaB, washing with buffer B was repeated a further three times. Purified S-layer
403 proteins were washed once with distilled water and stored at 4°C. The removal of SlaB was
404 confirmed by SDS-PAGE analysis.
405

406 **Disassembly and reassembly of S-layers**
407 S-layers were disassembled by increasing the pH with the addition of 20 mM $NaCO_3$, pH 10, 10
408 mM $CaCl_2$ and incubation at 60°C, 600 rpm (Thermomixer F1.5, Eppendorf) for 1h. Disassembly
409 was verified by SDS-PAGE analysis.
410

411 Reassembly was achieved by buffer exchange using 0.5 ml concentrators according to the
412 manufacturer's instructions (Pierce), with dH_2O to reduce the pH to 7, followed by the addition of
413 10mM $CaCl_2$ and incubation at 60°C for 120 hours.
414

Negative stain tomography
A 3 µl sample of S-layers with and without SlaB was placed on glow-discharged carbon-coated
copper grids (Agar Scientific) and incubated at room temperature for 1 min. The excess sample
was removed by blotting using filter paper (Whatman, GH healthcare). The specimens were
stained using ammonium molybdate for 1 min. Grids were blotted dry with filter paper and let air
dry completely. Single axis tilt series (-60 to +60) were recorded on a FEI T12 electron microscope
operated at 120 kV, using DigitalMicrograph (Gatan Inc.).
Tilt series were reconstructed using the IMOD package (Kremer et al, 1996) and tomograms were
generated using the weighted back-projection algorithm.
424

Electron cryo-tomography
3 µl of S-layer suspension were applied to glow-discharged 300 mesh copper Quantifoil grids
(R2/2, Quantifoil, Jena, Germany), blotted for 3 – 5 seconds and rapidly injected into liquid ethane
using a homemade plunge-freezer. Tomograms were recorded using a Polara G2 Tecnai
(Thermo-Fisher, Eindhoven, The Netherlands) or a Jeol 3200 FSC TEM (JEOL, Tokyo, Japan),
both based at the Max Planck Institute of Biophysics in Frankfurt, Germany. The microscopes
were operated at 300 kV. Data were collected using a 4 x 4 k CCD camera or a 4x4 k K2 Summit
direct electron detector (Gatan Inc., Pleasanton, USA) running in counting mode. Inelastically
scattered electrons were removed using a Gatan Tridiem energy filter (Gatan Inc., Pleasanton,
USA) for the Polara and an in-column energy filter for the JEOL microscope. Tilt series were
collected in zero-loss mode using Digital Micrograph (Gatan, Pleasonton, USA) from -62° to +62°

and in steps of 2°. For tomograms of whole cells, the Polara was used. The magnification was set

to 41.000 x (resulting in a final pixel size of 5.4 Å on the final image) and defocus values of 6-8

µm were applied. For isolated S-layers, the magnification used was 72.000 (final image pixel of

2.8 Å) on the Polara and 10.000 x (final pixel size 3.35) on the JEOL. Defocus values of 2-4 µm

were used. Whole-cell tomograms were recorded using a CCD camera with a total dose of 100

e⁻ Å⁻². Tomograms of isolated S-layers were recorded using the K2 and in dose-fractionation mode

at a dose rate of 8-10 e⁻ px⁻¹ s⁻¹ and a maximum total dose of 60 e⁻ Å⁻². Dose-fractionated tilt

images were aligned using an in-house script based on IMOD (Kremer et al, 1996) programmes,

CTF corrected and reconstructed into tomograms using the IMOD software package (Kremer et

al, 1996).

Sub-tomogram averaging

To average the S-layer, a random grid of points was applied over the S-layer within a tomograms

of *Sulfolobus* cells or isolated S-layers, which were previously binned twofold. Whole-cell

tomograms were filtered by NAD (Frangakis and Hegerl, 2001) to enhance the contrast. Using

PEET (Nicastro et al, 2006; Heumann et al, 2011), subvolumes were extracted, aligned and

averaged. For the *in situ* S-layer average, 1809 subvolumes (*SisI*) and 2068 subvolumes (*Saci)*

of 60x60x60 pixels in dimension were used. For isolated S-layers, 325 subvolumes (*Saci)* and

5561 subvolumes (*Ssol)* of 100 x 100 x 80 were used. P3 symmety was applied to the final

averages. S-layers were visualised and segmented using UCSF Chimera (Pettersen et al, 2004).

The resolution of all averages was estimated based the reflections in their respective power

spectra calculated by IMOD (Kremer et al, 1996). For the *in situ* structures this suggested a

resolution of 28 Å for *Saci* and 30 Å for *SisI*. For the maps of the isolated S-layers, 16 Å were

measured for *Ssol* and 21 Å for *Saci* (Fig. 1S5).

Difference maps and assembly models

The difference maps were calculated and assembly models built in UCSF Chimera (Pettersen et

al, 2004). To calculate the difference map between the fully assembled and SlaB-depleted S-

layers, the *vop subtract* command was used. This resulted in clear densities for SlaB trimers. To

build the assembly model of the SlaA canopy regions, semi-automated water-shedding

segmentation was initially used on the sub-tomogram average of the SlaB-depleted S-layers. This

approach segmented the map in equal-sized parts, which were then grouped to generate a P3

lattice of dimers. Individual SlaB and SlaA subunits were extracted as separate densities from the

map and fitted into the fully assembled *Ssol and Saci* S-layers to yield the complete SlaA/SlaB
assembly model.
471
472 **Sequence alignments**
473 Protein sequences were sourced online using Uniprot (https://www.uniprot.org/) and multiple
474 sequence alignments performed with the Praline Server
475 (http://www.ibi.vu.nl/programs/pralinewww/)
476

Acknowledgements

478 We thank Werner Kühlbrandt and Deryck Mills (MPI of Biophysics in Germany) for allowing us to
479 use the Polara electron microscope. This project was funded by the Max Planck Society (BD), the
480 University of Exeter Research Fellow's Startup grant (BD) and the ERC starting grant
481 "MICROROBOTS" (BD, LG, MM, KS).

References

Albers, S. V., & Meyer, B. H. (2011). The archaeal cell envelope. *Nature Reviews*
Microbiology. https://doi.org/10.1038/nrmicro2576

Banerjee, A., Tsai, C. L., Chaudhury, P., Tripp, P., Arvai, A. S., Ishida, J. P., … Albers, S.
V. (2015). FlaF is a beta-sandwich protein that anchors the archaellum in the archaeal
cell envelope by binding the S-layer protein. *Structure*, *23*(5), 863–872.
https://doi.org/10.1016/j.str.2015.03.001

Baumeister, W., & Lembcke, G. (1992). Structural features of archaebacterial cell
envelopes. *Journal of Bioenergetics and Biomembranes*.
https://doi.org/10.1007/BF00762349

Bharat, T. A. M., Kureisaite-Ciziene, D., Hardy, G. G., Yu, E. W., Devant, J. M., Hagen, W.
J. H., … Löwe, J. (2017). Structure of the hexagonal surface layer on Caulobacter
crescentus cells. *Nature Microbiology*. https://doi.org/10.1038/nmicrobiol.2017.59

Braun, T., Vos, M. R., Kalisman, N., Sherman, N. E., Rachel, R., Wirth, R., … Egelman, E.
H. (2016). Archaeal flagellin combines a bacterial type IV pilin domain with an Ig-like
domain. *Proceedings of the National Academy of Sciences*, *113*(37), 10352–10357.
https://doi.org/10.1073/pnas.1607756113

Chaudhury, P., Quax, T. E. F., & Albers, S. V. (2018). Versatile cell surface structures of
archaea. *Molecular Microbiology*. https://doi.org/10.1111/mmi.13889

Chen, L., Brügger, K., Skovgaard, M., Redder, P., She, Q., Torarinsson, E., … Garrett, R.
A. (2005). The genome of Sulfolobus acidocaldarius, a model organism of the
Crenarchaeota. *Journal of Bacteriology*. https://doi.org/10.1128/JB.187.14.4992-
4999.2005

D'Angelo, M. A., & Hetzer, M. W. (2008). Structure, dynamics and function of nuclear pore
complexes. *Trends in Cell Biology*. https://doi.org/10.1016/j.tcb.2008.07.009

Daum, B., Vonck, J., Bellack, A., Chaudhury, P., Reichelt, R., Albers, S.-V., … Kühlbrandt,
W. (2017). Structure and *in situ* organisation of the *Pyrococcus furiosus* archaellum
machinery. *ELife*, *6*. https://doi.org/10.7554/eLife.27470

Daum, B., Quax, T.E.F., Sachse, M., Mills, D., Reimann, J., Yildiz, Ö., Häder, S., Saveanu,
C., Forterre, P., Albers, S.V., Kühlbrandt, W. and Prangishvili, D. (2014). Self-
assembly of the universal membrane-remodeling protein PVAP into 7-fold virus-
associated pyramids. Proceedings of the National Academy of Sciences of the United
States of America, 111(10):3829-34. https://doi.org/10.1073/pnas.1319245111

Engelhardt, H. (2007). Are S-layers exoskeletons? The basic function of protein surface
layers revisited. *Journal of Structural Biology*, *160*(2), 115–124.
https://doi.org/10.1016/j.jsb.2007.08.003

Grogan, D. W. (2010). Organization and interactions of cell envelope proteins of the
extreme thermoacidophile Sulfolobus acidocaldarius . *Canadian Journal of*
Microbiology. https://doi.org/10.1139/m96-148

Henche, A. L., Ghosh, A., Yu, X., Jeske, T., Egelman, E., & Albers, S. V. (2012). Structure
and function of the adhesive type IV pilus of Sulfolobus acidocaldarius. *Environmental*
Microbiology. https://doi.org/10.1111/j.1462-2920.2012.02898.x

Kawarabayasi, Y., Hino, Y., Horikawa, H., Jin-No, K., Takahashi, M., Sekine, M., …
Kikuchi, H. (2001). Complete genome sequence of an aerobic thermoacidophilic
crenarchaeon, Sulfolobus tokodaii strain. *DNA Research*.
https://doi.org/10.1093/dnares/8.4.123

Palmieri, G., Balestrieri, M., Peter-Katalinić, J., Pohlentz, G., Rossi, M., Fiume, I., &
Pocsfalvi, G. (2013). Surface-exposed glycoproteins of hyperthermophilic sulfolobus
solfataricus P2 show a common N- glycosylation profile. *Journal of Proteome*
Research. https://doi.org/10.1021/pr400123z

Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E.
C., & Ferrin, T. E. (2004). UCSF Chimera: A visualization system for exploratory
research and analysis. *Journal of Computational Chemistry*, *25*(13), 1605–1612.
https://doi.org/10.1002/jcc.20084

Peyfoon, E., Meyer, B., Hitchen, P. G., Panico, M., Morris, H. R., Haslam, S. M., … Dell, A.
(2010). The S-layer glycoprotein of the crenarchaeote Sulfolobus acidocaldarius is
glycosylated at multiple sites with chitobiose-linked N-glycans. *Archaea*.
https://doi.org/10.1155/2010/754101

Plaut, R. D., Beaber, J. W., Zemansky, J., Kaur, A. P., George, M., Biswas, B., …
Sozhamannan, S. (2014). Genetic evidence for the involvement of the S-Layer protein
gene Sap and the sporulation genes spo0A, spo0B, and spo0F in phage AP50c
infection of Bacillus anthracis. *Journal of Bacteriology*.
https://doi.org/10.1128/JB.00739-13

Pohlschroder, M., Pfeiffer, F., Schulze, S., & Halim, M. F. A. (2018). Archaeal cell surface
biogenesis. *FEMS Microbiology Reviews*. https://doi.org/10.1093/femsre/fuy027

Poweleit, N., Ge, P., Nguyen, H. H., Loo, R. R. O., Gunsalus, R. P., & Zhou, Z. H. (2016).
CryoEM structure of the Methanospirillum hungatei archaellum reveals structural
features distinct from the bacterial flagellum and type IV pili. *Nature Microbiology*, *2*,
16222. https://doi.org/10.1038/nmicrobiol.2016.222

Pum, D, Messner, P., & Sleytr, U. B. (1991). Role of the S layer in morphogenesis and cell
division of the archaebacterium Methanocorpusculum sinense. *Journal of*
Bacteriology, *173*(21), 6865–6873. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/1938891

Pum, Dietmar, & Sleytr, U. B. (2014). Reassembly of S-layer proteins. *Nanotechnology*.
https://doi.org/10.1088/0957-4484/25/31/312001

Quax, T.E.F., Daum, B. (2018) Structure and assembly mechanism of virus-associated
pyramids. Biophysical Reviews 2018 Apr;10(2):551-557. doi: 10.1007/s12551-017-
0357-4. Epub 2017 Dec 4. https://doi.org/10.1007/s12551-017-0357-4

Quemin, E.R.J., Lucas, S., Daum, B., Quax, T.E.F, Kühlbrandt, W., Forterre, P., Albers,
S.V., Prangishvili, D., Krupovic, M. (2013) First insights into the entry process of
hyperthermophilic archaeal viruses. Journal of Virology 87(24):13379-85.
https://doi.org/ 10.1128/JVI.02742-13

Rodrigues-Oliveira, T., Belmok, A., Vasconcellos, D., Schuster, B., & Kyaw, C. M. (2017).
567   Archaeal S-layers: Overview and current state of the art. *Frontiers in Microbiology*.
568   https://doi.org/10.3389/fmicb.2017.02597
569 She, Q., Zhang, C., Deng, L., Peng, N., Chen, Z., & Liang, Y. X. (2009). Genetic analyses
570   in the hyperthermophilic archaeon Sulfolobus islandicus . *Biochemical Society*
571   *Transactions*. https://doi.org/10.1042/bst0370092
572 Taylor, K. A., Deatherage, J. F., & Amos, L. A. (1982). Structure of the S-layer of
573   Sulfolobus acidocaldarius. *Nature*. https://doi.org/10.1038/299840a0
574 Tschitschko, B., Williams, T.J., Allen, M.A., Páez-Espino, D., Kyrpides, N., Zhong, L.,
575   Raftery, M.J., Cavicchioli, R. (2015). Antarctic archaea-virus interactions:
576   metaproteome-led analysis on invasion, evasion and adaption. ISME J., 9(9):2094-
577   2107. https://doi.org/10.1038/ismej.2015.110
578 Veith, A., Klingl, A., Zolghadr, B., Lauber, K., Mentele, R., Lottspeich, F., … Kletzin, A.
579   (2009). *Acidianus* , *Sulfolobus* and *Metallosphaera* surface layers: structure,
580   composition and gene expression. *Molecular Microbiology*, *73*(1), 58–72.
581   https://doi.org/10.1111/j.1365-2958.2009.06746.x
582 Zhang, C., Wipfler, R. L., Li, Y., Wang, Z., Hallett, E. N., & Whitaker, R. J. (2018).
583   Revealing S-layer Functions in the Hyperthermophilic Crenarchaeon Sulfolobus
584   islandicus. *BioRxiv*. https://doi.org/10.1101/444406

Figures

Figure 1

Figure 1 – Cryo electron tomography of *Sulfolobus* cells. A, tomographic slice through a *Sisl* cell. **B**, cross section through the cell membrane and S-layer. **C**, segmented surface representation of the *Sisl* S-layer. Scale bar (A), 500 nm. **D-G**, Sub-tomogram averages of *Saci* (D,E) and Sisl (F,G) S-layers. Maps are coloured by proximity to the membrane plane. Yellow, proximal; red, distal. Scale bars (D, F), 20 nm.

Figure 2

Figure 2 - Structure of isolated S-layers from *S. acidocaldarius* and *S. solfataricus*. A, B, S-layer maps of *Saci* (A) and *Ssol* (B) at 21 Å or 16 Å resolution, respectively. **C, D,** S-layer unit of *Saci* (C) and *Ssol* (D), including three stalks and a hexameric pore. Maps are coloured by proximity to the membrane plane. Yellow, proximal; red, distal; purple, conical protrusions. Scale bars, 20 nm.

Figure 3

Figure 3 - Structural dissection of the *S. solfataricus* S-layer. A, SDS PAGE. M, marker; 1, washed once; 2, washed twice; 3, watched three times in detergent. **B**, sub-tomogram average of fully-assembled S-layer. **C**, sub-tomogram average of SlaB-depleted S-layer. **D**, difference map (pink) overlayed with the complete S-layer visualises location of SlaB.

Figure 4

Figure 4 - S-layers recrystallise with and without SlaB. A-F, reassembly of *Sulfolobus* S-layers from SlaA and SlaB. G-L, reassembly of S-layers from SlaA only. (A,D,G,J, tomographic slices; B,E,H,K, 3D surface representation; C,F,I,L, sub-tomogram averages). White circle highlights presence of SlaB (C&F) and red circles absence of SlaB (I&L). Scale bar: 20 nm

Figure 5

Figure 5 - Assembly models for the *S. acidocaldarius* (A-C) and *S. solfataricus* (D-F) S-layers. A, **D** outward facing surfaces (grey mesh) segmented into SlaA dimers (each dimer in a different colour) and SlaB trimers (white). **B**, **E** inward-facing surface. **C**, **F** side view, perpendicular to the membrane plane. Scale bar: 20 nm

Figure 6

Figure 6 – SlaA structure determines S-layer topology and unit dimensions. Comparison of the structures of SlaA from *Saci* (**A**) and *Ssol* (**B**) show differences in length, height and angular shape of the dimer. **C, D**, location of one SlaA dimer within the S-layer of *Saci* (**C**) and *Ssol* (**D**). Red circles / arrowheads indicate apical domains that determine shape and topology of the hexameric pores.

Figure 7

Figure 7 – Model of the *Sulfolobus* S-layer compared to the single-subunit S-layer of haloarchaea. The Sulfolobus S-layer (upper panel) consists of the two protein subunits: SlaA dimers (red, orange, yellow) form the outer S-layer canopy. Each SlaA protein is predicted to be rich in **β** strands. The SlaA dimer has a boomerang-like shape, the angle of which determines the S-layer unit cell size. SlaB trimers (grey) form the membrane anchors of the S-layer. Each SlaB is predicted to consist of an N-terminal transmembrane domain (TMD), a coiled-coil domain (CC) and 2-3 C-terminal **β**-sandwich domains (**β**). SlaA and SlaB proteins are highly glycosylated (green). In contrast, the S-layers of many halophile archaea (such as that of Haloferax volcanii; bottom panel) consist of multiple copies of only one S-layer glycoprotein (SLG) subunit and are inserted into the membrane by a posttranslationally added lipid anchor (LA).

Figure 1S1

A

B

Figure 1 S1 – Sequence alignment comparing SlaA (A) and SlaB (B) from *SisI* and *Ssol*.
The sequence identity for SlaA and SlaB is 87.4% and 87.7%, respectively.

Figure 1S2

A

B

C

D

Figure 1 S2 – Sequence alignment comparing SlaA and SlaB from *SisI, SsoI* and *Ssaci*. The sequence identity: *SisI*_SlaA / S*aci*_SlaA, 24% (A); *SisI*_SlaB / S*aci*_SlaB, 25% (B); *Ssol*_SlaA / S*aci*_SlaA, 25% (C); *SisI*_SlaB / S*aci*_SlaB, 26% (D).

Figure 1S3

Saci   *Sisl*

Figure 1 S3 – Tomographic slices through S-layer planes in whole-cell tomograms. Left panel, tomographic slice through the S-layer of Sulfolobus acidocaldarius; right panel, tomographic slice through the S-layer of Sulfolobus islandicus. Insets, power spectra reveal 2D lattices with hexagonal symmetry.

Figure 1S4

Saci (in situ)
membrane-distal

membrane-proximal

Sisl (in situ)
membrane-distal

membrane-proximal

Figure 1 S4 – Consecutive slices through sub-tomogram averages of S-layers. Top panel, in situ map calculated from whole-cell tomogram of Sulfolobus acidocaldarius, bottom panel, map calculated from whole-cell tomogram of Sulfolobus islandicus.

Figure 1S5

In situ

S. acidocaldarius

S. islandicus

Isolated

S. acidocaldarius

S. solfataricus

Figure 1 S5 – Resolution of S-layer maps determined by power spectra. Top panel, power spectra of in situ maps calculated from whole-cell tomograms, bottom panel, power spectra of maps calculated from isolated S-layers. Ny, Nyquist frequency