Running Head:

White spotting in the American Paint Horse

Title:

Impact of white spotting alleles, including *W20*, on phenotype in the American

Paint Horse

Samantha A. Brooks[*], Katelyn M. Palermo[*], Alisha Kahn[*], and Jessica Hein[#]

[*]University of Florida Department of Animal Sciences, UF Genetics Institute,

Gainesville FL, 32611-0910

[#]American Paint Horse Association, Fort Worth TX, 76161-0023

Acknowledgments:

The authors would like to thank the many APHA staff members for their efforts in

submitting and collating the data analyzed in this study. Thanks to the UF

undergraduate researchers who generously volunteered for data-entry work on this

project: Hannah Hillard, Kalisse Horne, Rachel Kullman, Erica Riano, Matt Winter,

Courtney McCreary, Rachel Shepherd, Anna Moskovitz, and Kaycie Miller. Our

gratitude to Dr. Ernie Bailey for proofreading the manuscript.

Conflicts of Interest:

The Authors have no conflicts to declare.

Summary:

The American Paint Horse Association (APHA) records pedigree and performance

information for their breed, a stock-type horse valued as a working farm or ranch horse

and as pleasure horses. As the name implies, the breed is also valued for attractive

white spotting patterns on the coat. The APHA utilizes visual inspections of

photographs to determine if coat spotting exceeds threshold anatomical landmarks

considered characteristic of desirable patterns. Horses with sufficient white patterning

enter the "Regular" registry, rather than the "Solid Paint-bred" division, providing a

threshold modeled phenotype. Genetic studies previously defined sequence variants

corresponding to 35 alleles for white spotting in the horse. Here, we calculate the allele

frequency for nine common white spotting alleles in the American Paint horse using a

sample of 1,054 registered animals. The APHA spotting phenotype is altered by

additive interactions among spotting loci, and epistatically by the *MC1R* and *ASIP*

genes controlling pigment production. The *W20* allele within the *KIT* gene, independent

of other known spotting alleles, was strongly associated with the APHA-defined white

spotting phenotype (p = 1.86 x10^{-18}), refuting reports that *W20* acts only as a modifier of

other underlying white spotting patterns. The parentage of an individual horse, either

American Paint or American Quarter Horse, did not alter the likelihood of entering the

APHA Regular registry. An empirical definition of the action of these genetic loci on the

APHA-defined white spotting phenotype will allow more accurate application of genome-

assisted selection for improving color production and marketability of APHA horses.

Key words:

KIT

coat color

ASIP

MC1R

W20

American Paint Horse

Introduction:

Founded in 1962, the American Paint Horse Association's mission includes preserving the lineage of colorful, versatile stock horses through registration with the association (Hood 1987). Since its inception, registration of the Paint Horse has been based primarily on a combination of pedigree and phenotype in order to select for the white-patterned, stock-type horse valued by APHA's founders—this includes acceptance of "cropout" Quarter Horses, who were refused registration with the American Quarter Horse Association until 2004 (AQHA 2019). As of January 1, 1980, the association requires all fully-registered APHA horses to have a sire and dam registered with either the American Paint Horse Association, the American Quarter Horse Association or The Jockey Club, a standard that remains today (APHA 2018). With more than 1.09 million horses registered with its organization as of May 2019, the APHA ranks as the second-largest equine breed association in the world. In the US alone, the equine industry contributes a total of approximately $122 billion to the U.S. economy and 1.7 million jobs (American Horse Council Foundation 2019). While the American Paint Horse is valued for conformation and performance traits, the white spotting pattern on the coat exerts an overwhelming influence on the economic value of individual horses within this large breed registry (Brooks et al. 2007).

During the APHA registration process, horses are designated into one of two sub-registries based on the presence or absence of qualifying natural white markings on their coats: the Regular registry or the Solid Paint-bred registry. This is done through phenotypic observation of photographs submitted of the horse, analyzed by trained APHA staff members against a set of thresholds (reference lines along the hocks, knees and head) determined by the association's board of member-elected directors, not empirical scientific data. Regular registry status is granted for horses that meet white spotting requirements as outlined in the *APHA Rule Book* which has contained a diagram describing these thresholds since 1973 (APHA 2018). The patterns desired by the APHA extend beyond what is typically observed as an average white marking on the face or legs (Haase et al. 2013) and exclude markings created by the *Leopard* locus (Bellone et al. 2013).

In brief, unpigmented areas characterizing APHA-accepted white spotting

patterns must include both white hair and skin extending beyond reference lines

determined by the association's elected governing body; those reference lines are not

based on empirical thresholds of known white spotting patterns' distribution zones. On

the face, qualifying depigmented areas must extend beyond a reference line from the

base of the horse's ear to the outside corner of the eye to the corner of the mouth and

under the chin to the opposite corner of the mouth (Figure 1). On the body, qualifying

depigmented areas must occur above a perpendicular line around the leg located at the

center of the knee or hock (Figure 1). These spotting patterns must include a minimum

of two inches of solid white hair with some underlying unpigmented skin in the qualifying

area. The Solid Paint-Bred registry is used for horses who meet the parentage

requirement but lack the minimum requirement for white spotted skin and coats as

defined in the rulebook (APHA 2018).

When registering a Paint Horse today, the applicant provides identifying

information pertaining to the foal, including its date of birth, color, pattern, sex,

parentage and ownership. The APHA requires parentage verification via genetic testing

when the horse is the product of breeding with transported or frozen semen, embryo

transfer or vitrified embryo transfer, or other special situations, such as when registering

a horse over age 10, and for all Quarter Horses and Thoroughbreds applying for

registration as Paint Horses (APHA 2018).

The applicant also provides a minimum of four full-color photographs of the

horse, showing the entire animal from the left side, right side, front and rear. Additional

photographs of spotted areas of the coat might be required by APHA staff to verify that

the pattern meets registration guidelines in terms of the location, size and requirement

for underlying skin to also be unpigmented (APHA 2018). The photographs become part

of the horse's permanent record at APHA and are used to identify the animal throughout

its life. Trained APHA staff members determine the registration category (Solid Paint-

Bred Registry or Regular Registry for each horse by visual inspection to confirm that a

horse's natural white spotting patterns meets APHA's standard. The quality of the

submitted photographs is variable, adding additional challenges to the interpretation of

photographic evidence of the pattern relative to the APHA standards.

In the horse, at least 35 known white spotting polymorphisms underly white

spotting patterns on the skin and coat: for a comprehensive and current listing of

spotting alleles in the horse see OMIA.org (OMIA 2019). Of these known alleles, nine

are common to and routinely tested for in the APHA population (Table 1). Others, like

MITF- Splash White 4 and many of the *KIT*W* alleles, are breed specific and not

present in the Paint Horse, while some like *MITF- Splash White 5* were not yet

commercially available for testing at the time the study began and therefore were not

examined (Henkel et al. 2019).

Among the APHA community, significant controversy surrounds the action of

KIT*W20. The first published observation of the *KIT*W20* allele came from a study

searching for causative variants in horse with white coats, phenotypes that left the

majority of the skin surface totally unpigmented, and therefore did not attempt to

document relatively less extensive white spotting phenotypes among their sampled

population (Haase et al. 2007). Subsequently, the authors examined the *KIT*W20*

variant in a larger set of horses (n= 52) and noted a compound heterozygote effect

creating extensive white patterning in horses with the *KIT*W5/W20* genotype, and

observing a significant effect of the *KIT*W20* allele generating a white spotting

phenotype more expansive than the typical white markings on the face and legs

(Hauswirth et al. 2013). These authors state the circumstances in their publication: "We

previously discovered a missense variant in exon 14 of the equine *KIT* gene

(c.2045G>A; p.Arg682His) but did not immediately recognise [sic] its functional role

(Haase et al. 2007)." However, in the eyes of the lay audience, the latter publication was

eclipsed by the 2007 paper containing limited observations, and the lack of data

specifically addressing the phenotype produced by the *KIT*W20* allele led to public

claims that there is no, or only a "subtle" impact from the *KIT*W20* allele on white

spotting (https://www.vgl.ucdavis.edu/services/horse/dominantwhite.php).

As a whole, the population frequencies of all white-spotting alleles lack

investigation, and despite the easily applied Mendelian inheritance patterns for most of

these alleles, genetic testing for selection of breeding stock remains underutilized in the

industry. Only recently has the APHA industry expressed an interest in using readily

available genetic tools to improve their color production. Previously, selective breeding

for spotting patterns in the American Paint Horse was commonly based on hearsay,

personal experience, and limited observations. Starting in 2017, the APHA has

developed rules that help designate a horse to the Regular Registry based on presence

of white-spotting alleles. These new rules give breeders and horse owners additional

tools to make responsible breeding decisions in an effort to produce the healthiest, most

marketable horse possible. This more inclusive approach encourages objective

registration of foals with phenotypes not well characterized by the APHA threshold

method alone, adding significant economic value to these horses, which in turn can

ultimately improve their long-term welfare (Hein 2017).Recent work highlighting

"exceptions" to proposed qualitative phenotypes for these loci emphasizes the need for

application of impartial and quantitative assessments capturing the full breadth of

phenotypes resulting from these variants (Druml et al. 2018). Toward this ideal, the

goals of this study were to: establish allele frequencies for known white spotting alleles

in the American Paint Horse, investigate association of these loci with a threshold

phenotype uniformly defined by the APHA, identify interactions between loci, any effect

due to the sex of the horse and to test association of the *W20* allele with the APHA

photo-based threshold phenotype.

Materials and Methods:

Retrospective Registration Records

The American Paint Horse Association provided data for a sampling of 1,054

horses born and registered between 1992 and 2018 including: registered name and

number, registration type ("Regular" registry *i.e.* possessing a spotted coat or "Solid

Paint-Bred" phenotype), age, and sex for each horse, as well as the registry and

registration type of the sire and dam for that horse. The original photos submitted by the

applicant for evaluation of the white spotting pattern on the horse at the time of

registration were also provided by the APHA and used to confirm predicted phenotypes

by visual inspection from a single experienced observer blinded to the identity and

genotype of the horse (SAB). In 2012, the APHA began partnering with commercial

providers of genetic testing to record genotypes on their registered horses. These

1,054 horses were voluntarily tested for color alleles by their owners for their own

educational use or interests. Results were already on file with APHA and although they

are not a randomly extracted subset of all APHA registrations, the large sample size

does offer the opportunity to investigate important questions regarding the action of

these alleles.

Genotypes

Each horse was genotyped for nine spotting pattern loci using a commercial

service (Veterinary Genetics Laboratory, University of California, Davis) and the data

shared with the APHA through a collaborative agreement. Alleles at the following loci

were examined (Table 1): *KIT- TO, SB1, W5, W10, W20* or "*N*" (Brooks & Bailey 2005;

Brooks *et al.* 2007; Haase et al. 2009; Hauswirth *et al.* 2013), *EDNRB- O* or "*N*"

(Metallinos et al. 1998; Santschi et al. 1998; Yang et al. 1998)*, MITF- SW1, SW3* or *N*

and *PAX3- SW2* or "*N*" (Hauswirth et al. 2012). The testing service provider designates

the wild type allele as "*N*". Additional genetic variants associated with white spotting

patterns in the horse are documented in the scientific literature, but were either not

available commercially at the time of testing and/or do not commonly segregate in the

APHA and are therefore not relevant to this population (for a comprehensive catalog of

spotting alleles in the horse see OMIA.org). Additionally, two known pigmentation loci

were genotyped for investigation of epistasis: *MC1R- E* or *e* (Marklund et al. 1996)*,*

ASIP- A or *a* (Rieder et al. 2001)*.* For the purpose of calculating allele frequencies we

refer to the allele carried by the reference genome assembly (EquCab3.0) as the wild

type (Kalbfleisch et al. 2018).

 Statistical analyses

Analyses were conducted in the JMP Pro v14.1.0 package (SAS Institute Inc.).
The additive effect of multiple white spotting loci was tested by a logistic regression of
the number of alleles against the two possible registry categories. Linkage, assessed as
LOD scores between coat color loci on ECA3, was assessed using HaploView v4.2
(Barrett 2009). To measure the impact of the *MC1R-ASIP* signaling system on white
spotting, a subset of 364 horses that possessed only one spotting allele (controlling for
any effect due to interactions between multiple spotting loci) was utilized, excluding all
horses possessing a *Tobiano* allele (in order to avoid the impact of linkage between
MC1R alleles and *TO).* Given the well documented interaction between alleles at *ASIP*
and its antagonistic target, the *MC1R* receptor, a logistic regression model was
constructed comparing the APHA registry phenotype (Regular or Solid Paint-Bred) with
the linear ranking of the genotypes by the predicted signaling activity and base color
phenotype. Thus, the *E/- a/a* genotype, likely resulting in a constitutively active *MC1R*
receptor and the black base color, was scored "0"; the wild-type genotype *E/- A/-*,
predicted to have normal signaling activity and a bay base color, was given a "1"; and
the *e/e* genotype, resulting in a loss of *MC1R* signaling regardless of the *ASIP* genotype
and a chestnut base coat, received a score of "2" (Barrett et al. 2005). Association
between genotypes for the *W20* allele and the APHA threshold phenotype was
assessed in a subset 529 horses bearing no other white spotting alleles at any of the
other eight sites tested and using a Chi-square test under a dominant model (merging
the *W20/N* and *W20/W20* genotypes).

Results and Discussion:

Phenotype distribution varies for each white spotting genotype

The 1,054 horses registered with the APHA in the examined time-period included
406 stallions, 90 geldings and 558 mares submitted for registration at a mean age of
1.07 years between 1992 and 2018. 90% of the sample comprised horses registered
post-2004. Among these 1,054 horses, 773 were designated to the Regular registry,

while 281 entered the Solid Paint-Bred category. This sampling did not represent an

unbiased observation of all foals produced from APHA-registered breedings. Not all

foals are submitted for registration (especially if the foal is unlikely to achieve Regular

registry status), and because DNA genetic testing for white spotting alleles is not

required for registration, the available data for this retrospective study is limited to

horses registered with APHA that also had DNA testing results available for

investigation. Records are not retained by the APHA on submitted horses ultimately

rejected for registration. Nonetheless, this sample is the largest collection of genotype

data and white spotting phenotypes in the horse.

Genotypic and allele frequencies for each of the nine variants examined are

presented in Table 2. Presence of just a single alternate allele resulted in APHA

designation to the Regular registry for *KIT*TO*, *MITF*SW3,* and *KIT*SB1,* although the

latter two genotypes were only rarely observed (n = 3 and 8 respectively, Table 3). As

expected, we did not observe a homozygous *EDNRB*O* genotype as this causes the

well-documented Lethal White Overo Syndrome (Metallinos *et al.* 1998; Santschi *et al.*

1998; Yang *et al.* 1998). However, a single horse homozygous for the *SW2* allele was

identified, which is a state previously hypothesized to be lethal based on comparisons to

similar variants in the *PAX3* gene of other species (Hauswirth *et al.* 2012). The

SW2/SW2 genotype was confirmed in a second DNA sample submitted independently

and anonymously. Veterinary examinations of this horse were unavailable at the time of

writing, but his owner reports that aside from apparent deafness she is healthy and

strong (Figure 3.).

109 horses possessed at least one white spotting allele but were APHA

registered Solid Paint-bred based on visual examination of photographs. Visual

inspection of the photographs by a single blinded observer revealed that the extent and

distribution of white markings in these horses often fell just below the guidelines of the

APHA Rule Book (2018) but were more extensive than the common white markings

described as "socks" on the limbs or a "blaze" on the face in previous studies (Haase *et*

al. 2013). The phenotype of these horses likely represents the tail of the natural

distribution of total depigmented area characteristic of the unique reduction in

melanocyte migration for each of these alleles. Previous work observed production of

foals with phenotypes very typical of the *TO* allele from parents with a *TO* allele but a

very minimal white phenotype (Stachurska & Jansen 2015). Thus, these horses likely

possess genetic value as producers of future spotted foals, although on the surface they

do not fit the visual description of a Regular registry horse.

One horse categorized as Solid Paint-Bred possessed a total of five white

spotting alleles and an entirely white coat. Although this animal has a high breeding

value for white spotting, it was registered in the less-valued Solid Paint-Bred category

because the horse did not exhibit at least two inches of contrasting colored hair in his

coat, per APHA registration guidelines. In these cases, phenotyping by photograph and

use of the *APHA Rule Book* description for Regular registry white spotting patterns

misses horses with superior genetic value for production of white spotting patterns.

For 97 horses recorded by APHA in the Regular registry based on their

photographic phenotype, genetic testing results did not identify possession of any of the

nine white spotting alleles evaluated. In some cases, white spotting patterns in these

horses may be due to rare genetic markers not tested in this study, or patterns for which

genetic markers/causative variations are not yet known. Visual inspection of the

registration photographs for these 97 horses revealed that 31 possessed patterns

consistent with white spotting loci. These 31 included many phenotypes resembling

those generated by alleles known but not genotyped in this cohort, including *MITF*SW4*

(Hauswirth *et al.* 2013), *MITF*SW5* (Henkel *et al.* 2019)*, and more than 27 other alleles

at the *KIT* locus. We also observed among these 31 horses some well-recognized

patterns with yet unknown genetic etiology (*i.e.* the "Rabicano" pattern) and two horses

with previously undocumented and distinct white spotting phenotypes. Phenotypes for

the remaining 66 (68%) horses fell within the described range for common white

markings of the face and legs (Haase *et al.* 2013), or typified the roan and grey coat

colors. Roan and grey do introduce white hair into the coat, but in a more interspersed

manner than desired by the APHA (Marklund et al. 1999; Sundstrom et al. 2012).

In total, the predicted phenotype from the genotypes at the nine spotting variants

investigated here agreed with the Regular registry (spotted) classification in 95.4% of

horses, demonstrating that these nine alleles account for an overwhelming majority of

spotting phenotypes present in the APHA. However, the photo-based designation to the

Regular or Solid Paint-Bred categories disagreed with the genotyped presence or

absence of a white spotting allele in 17% of horses (206) submitted for registration. This

discordance demonstrates the need for separate selection tools and interpretations for

each of the functions of the registry: quantification of desirable visual phenotypes versus

assessment of breeding value for production of white spotted foals.

Cumulative interactions among spotting loci, and between spotting loci and alleles for

base coat color, additively increase white spotting phenotypes above the APHA

threshold

The frequency of the Regular registry threshold phenotype varied by allele and

by allele combination (Table 3), however sample sizes within each possible combination

were still too small to statistically examine specific interactions between loci. However,

excluding the one horse possessing five alleles and an entirely white coat (considered a

Solid Paint-Bred under current rules,) horses with an increasing number of spotting

alleles are more likely to exhibit a phenotype that exceeds the APHA threshold for the

Regular registry (Chi-square = 226.11, p < 0.0001, Table 4). Thus, the nine spotting

alleles examined here act collectively in an additive fashion, increasing the proportion of

white on the horse above the APHA Regular registry phenotype threshold.

Although the *MC1R* signaling pathway primarily contributes to pigment switching

between eumelanin and pheomelanin, loss-of-function alleles at the *MC1R* locus can

also reduce migration of melanocyte stem cells (Chou et al. 2013). Assessing the effect

of MC1R signaling on depigmentation phenotypes resulting from the *KIT* locus is not

straightforward as the physical proximity of these two genes on ECA3 could produce an

association due to linkage. Indeed, suppression of recombination by the *Tobiano* allele,

a 36 Mb paracentric inversion encompassing a 41 Mb span between the *KIT* and *MC1R*

genes is well documented (Trommershausen-Smith 1978; Brooks *et al.* 2007). In this

dataset, we observed strong linkage between *MC1R*E* and the *TO* inversion allele (n =

1054, LOD= 40.96), but not between either *MC1R* allele and *KIT*W20* (LOD= 0.05).

Before assessing modification of spotting phenotype by alleles at *MC1R* and

ASIP, we selected a subset of horses that possessed just one spotting allele (avoiding

any coincidental interaction across multiple spotting alleles) and excluding any horse

with a *TO* allele (likely to produce association to *MC1R*E* due to recombination

suppression generated by the *TO* inversion). Among these 368 horses we observed a

significant effect of the *MC1R-ASIP* signaling system on the APHA-defined Regular

registry white spotting phenotype (Table 4, p = 0.0171), with the *e/e* genotype producing

the highest proportion of Regular registry horses.

The sex of the individual horse did not significantly impact the registry

designation when examined within 461 horses possessing just one white spotting allele

(Chi-square test, p = 0.209). Across the entire sample population of 1,054 horses,

stallions were more likely to enter the Regular registry than mares or geldings (p<

0.0001), but this may reflect a tendency for applicants to submit registrations on colts

with some prospect as future stallions, rather than an influence of sex or castration on

white spotting phenotype.

The KIT*W20 allele results in white spotting, independent of other known alleles

We investigated association between *KIT*W20,* a common allele in the APHA

population, and the APHA Regular registry phenotype using a subset of the sample

comprised of 529 horses possessing only the *N/N* (n= 270)*, W20/N* (n= 225) and

W20/W20 (n= 34) genotypes (no other spotting alleles at any of the 8 remaining loci

tested, to control for potential interactions with other variants.) *KIT*W20* is indeed

significantly associated with white spotting patterns on the coat, as defined by the

Regular registry APHA threshold (Chi-square test under a dominant model, p= 1.86 x10^{-18}

, Table 3). Clearly, *KIT*W20* strongly correlates with the white spotting phenotype as

defined by the *APHA Rule Book.* Novel white spotting patterns likely appear in 2.9% of

the population (discussed above), and thus would not be sufficient on their own to

coincidentally explain the effect of *KIT*W20* on the Regular registry spotting phenotype,

as has been previously suggested by breeders of APHA horses. The significantly

increased likelihood of achieving Regular registry status among horses carrying only the

KIT*W20 allele conclusively demonstrates the value of this trait in genomic selection

schemes for breeders seeking to improve their production of white spotted, Regular

registry eligible foals.

No evidence for unknown white spotting phenotype "modifying" alleles

Given the historical selective pressure to eliminate white spotting phenotypes in

the American Quarter Horse breed (AQHA 2019) and the rarity of white spotting

phenotypes within the Thoroughbred breed (often used in the production of American

Quarter Horses), APHA breeders are concerned that undiscovered "modifying" alleles

capable of reducing the white spotting phenotypes may be present in the American

Quarter Horse population. As APHA rules permit cross-breeding with American Quarter

Horses or Thoroughbred horses, as well as registration of horses from these breeds

that exhibit approved white spotting phenotypes, in-flow of "modifying" alleles could

negatively impact the expression of white spotting phenotypes on APHA horses in

future generations. 335 registrations were examined where the sire of the submitted

horse belonged to the APHA Regular registry and the dam was registered as either

Solid Paint-Bred (n= 189) versus those with dams from the American Quarter Horse

Association or The Jockey Club (Thoroughbred) registries (n= 146). No significant

difference was observed in the Regular vs. Solid Paint-Bred registry status of foals from

dams belonging to the Solid Paint-Bred registry compared to those with dams registered

with the AQHA or The Jockey Club (Chi-square test p= 0.704). Finally, at the genome-

wide scale, the genetic background of horses from the APHA and AQHA registries is

very similar (Petersen et al. 2013), and written history documents many AQHA-

registered foundation animals at the inception of the APHA.

Conclusions

Given the large population size of the APHA, estimated at over a million

animals, our sampling of just 1,054 animals may not be large enough to accurately

reflect allele frequencies across the breed as a whole. However, this is the largest

sampling of this breed accomplished to date and provides a first glimpse into the

potential value of genotype assisted selection for spotting pattern in the APH. Here we

document that most of the genetic markers responsible for the iconic color patterns

valued by American Paint Horse Association members, breeders and owners. These

loci are valuable tools for prediction of the genetic value of breeding stock, thereby

optimizing future production of spotted foals. Identification of phenotype-altering

interactions between the nine spotting loci tested, as well as with the *MC1R-ASIP*

signaling system, will improve genotype-based phenotype predictions. Genetic tools

could rapidly improve the accuracy of selection for white spotting in the horse and may

provide significant economic savings compared to the time-consuming photo-analysis

approach currently used to register APHA horses.

Literature Cited:

1. American Horse Council Foundation (2019) 2017 Economic Impact Study of the U.S. Horse Industry.
2. APHA (2018) *2018 APHA Rule Book*. American Paint Horse Association, Fort Worth, Texas.
3. AQHA (2019) Flashy Paint Coat Color. URL https://www.aqha.com/-/flashy-paint-coat-color.
4. Barrett J.C. (2009) Haploview: Visualization and analysis of SNP genotype data. Cold Spring Harb Protoc 2009, pdb ip71.
5. Barrett J.C., Fry B., Maller J. & Daly M.J. (2005) Haploview: analysis and visualization of LD and haplotype maps. Bioinformatics 21, 263-5.
6. Bellone R.R., Holl H., Setaluri V., Devi S., Maddodi N., Archer S., Sandmeyer L., Ludwig A., Foerster D., Pruvost M., Reissmann M., Bortfeldt R., Adelson D.L., Lim S.L., Nelson J., Haase B., Engensteiner M., Leeb T., Forsyth G., Mienaltowski M.J., Mahadevan P., Hofreiter M., Paijmans J.L.A., Gonzalez-Fortes G., Grahn B. & Brooks S.A. (2013) Evidence for a Retroviral Insertion in TRPM1 as the Cause of Congenital Stationary Night Blindness and Leopard Complex Spotting in the Horse. Plos One 8.
7. Brooks S.A. & Bailey E. (2005) Exon skipping in the KIT gene causes a Sabino spotting pattern in horses. Mamm Genome 16, 893-902.
8. Brooks S.A., Lear T.L., Adelson D.L. & Bailey E. (2007) A chromosome inversion near the KIT gene and the Tobiano spotting pattern in horses. Cytogenetic and genome research 119, 225-30.
9. Chou W.C., Takeo M., Rabbani P., Hu H., Lee W., Chung Y.R., Carucci J., Overbeek P. & Ito M. (2013) Direct migration of follicular melanocyte stem cells to the epidermis after wounding or UVB irradiation is dependent on Mc1r signaling. Nat Med 19, 924-9.
10. Druml T., Grilz-Seger G., Neuditschko M., Horna M., Ricard A., Pausch H. & Brem G. (2018) Novel insights into Sabino1 and splashed white coat color patterns in horses. Anim Genet 49, 249-53.
11. Haase B., Brooks S.A., Schlumbaum A., Azor P.J., Bailey E., Alaeddine F., Mevissen M., Burger D., Poncet P.A., Rieder S. & Leeb T. (2007) Allelic heterogeneity at the equine KIT locus in dominant white (W) horses. Plos Genetics 3, 2101-8.
12. Haase B., Brooks S.A., Tozaki T., Burger D., Poncet P.A., Rieder S., Hasegawa T., Penedo C. & Leeb T. (2009) Seven novel KIT mutations in horses with white coat colour phenotypes. Animal Genetics 40, 623-9.
13. Haase B., Signer-Hasler H., Binns M., Obexer-Ruff G., Hauswirth R., Bellone R., Burger D., Rieder S., Wade C. & Leeb T. (2013) Accumulating Mutations in Series of Haplotypes at the KIT and MITF Loci Are Major Determinants of White Markings in Franches-Montagnes Horses. Plos One 8.
14. Hauswirth R., Haase B., Blatter M., Brooks S., Burger D., Drogemuller C., Gerber V., Henke D., Janda J., Jude R., Magdesian K., Matthews J., Poncet P., Svansson V., Tozaki T., Wilkinson-White L., Penedo M., Rieder S. & Leeb T. (2012) Mutations in MITF and PAX3 Cause "Splashed White" and Other White Spotting Phenotypes in Horses. Plos Genetics 8, 404-12.
15. Hauswirth R., Jude R., Haase B., Bellone R., Archer S., Holl H., Brooks S., Tozaki T., Penedo M., Rieder S. & Leeb T. (2013) Novel variants in the KIT and PAX3 genes in horses with white-spotted coat colour phenotypes. Animal Genetics 44, 763-5.
16. Hein J. (2017) Color Me Paint. pp. 69-72, Fort Worth, TX.
17. Henkel J., Lafayette C., Brooks S.A., Martin K., Patterson-Rosa L., Cook D., Jagannathan V. & Leeb T. (2019) Whole-genome sequencing reveals a large deletion in the MITF gene in horses with white spotted coat colour and increased risk of deafness. Anim Genet 50, 172-4.
18. Hood N. (1987) Luncheon Honors Association Founders. In: *Paint Horse Journal*, pp. 30-2.

432   19.   Kalbfleisch T.S., Rice E.S., DePriest M.S., Walenz B.P., Hestand M.S., Vermeesch J.R., O Connell
433   B.L., Fiddes I.T., Vershinina A.O., Saremi N.F., Petersen J.L., Finno C.J., Bellone R.R., McCue M.E.,
434   Brooks S.A., Bailey E., Orlando L., Green R.E., Miller D.C., Antczak D.F. & MacLeod J.N. (2018)
435   Improved reference genome for the domestic horse increases assembly contiguity and
436   composition. Commun Biol 1, 197.
437   20.   Marklund L., Moller M.J., Sandberg K. & Andersson L. (1996) A missense mutation in the gene
438   for melanocyte-stimulating hormone receptor (MC1R) is associated with the chestnut coat color
439   in horses. Mamm Genome 7, 895-9.
440   21.   Marklund S., Moller M., Sandberg K. & Andersson L. (1999) Close association between sequence
441   polymorphism in the KIT gene and the roan coat color in horses. Mamm Genome 10, 283-8.
442   22.   Metallinos D.L., Bowling A.T. & Rine J. (1998) A missense mutation in the endothelin-B receptor
443   gene is associated with Lethal White Foal Syndrome: an equine version of Hirschsprung disease.
444   Mammalian genome : official journal of the International Mammalian Genome Society 9, 426-
445   31.
446   23.   OMIA (2019) Online Mendelian Inheritance in Animals. URL http://omia.org/.
447   24.   Petersen J., Mickelson J., Cothran E., Andersson L., Axelsson J., Bailey E., Bannasch D., Binns M.,
448   Borges A., Brama P., Machado A., Distl O., Felicetti M., Fox-Clipsham L., Graves K., Guerin G.,
449   Haase B., Hasegawa T., Hemmann K., Hill E., Leeb T., Lindgren G., Lohi H., Lopes M., McGivney
450   B., Mikko S., Orr N., Penedo M., Piercy R., Raekallio M., Rieder S., Roed K., Silvestrelli M.,
451   Swinburne J., Tozaki T., Vaudin M., Wade C. & McCue M. (2013) Genetic Diversity in the Modern
452   Horse Illustrated from Genome-Wide SNP Data. Plos One 8.
453   25.   Rieder S., Taourit S., Mariat D., Langlois B. & Guerin G. (2001) Mutations in the agouti (ASIP), the
454   extension (MC1R), and the brown (TYRP1) loci and their association to coat color phenotypes in
455   horses (Equus caballus). Mammalian genome : official journal of the International Mammalian
456   Genome Society 12, 450-5.
457   26.   Santschi E.M., Purdy A.K., Valberg S.J., Vrotsos P.D., Kaese H. & Mickelson J.R. (1998) Endothelin
458   receptor B polymorphism associated with lethal white foal syndrome in horses. Mammalian
459   genome : official journal of the International Mammalian Genome Society 9, 306-9.
460   27.   Stachurska A. & Jansen P. (2015) Crypto-tobiano horses in Hucul breed. Czech Journal of Animal
461   Science 60, 1-9.
462   28.   Sundstrom E., Imsland F., Mikko S., Wade C., Sigurdsson S., Pielberg G.R., Golovko A., Curik I.,
463   Seltenhammer M.H., Solkner J., Lindblad-Toh K. & Andersson L. (2012) Copy number expansion
464   of the STX17 duplication in melanoma tissue from Grey horses. BMC Genomics 13, 365.
465   29.   Trommershausen-Smith A. (1978) Linkage of Tobiano Coat Spotting and Albumin Markers in a
466   Pony Family. Journal of Heredity 69, 214-6.
467   30.   Yang G.C., Croaker D., Zhang A.L., Manglick P., Cartmill T. & Cass D. (1998) A dinucleotide
468   mutation in the endothelin-B receptor gene is associated with lethal white foal syndrome
469   (LWFS); a horse variant of Hirschsprung disease. Human molecular genetics 7, 1047-52.

472 **Tables and Figures:**

474 Table 1: Details of the nine polymorphisms genotyped for this analysis. The
475 "Reference" allele refers to the EquCab3.0 assembly and also happens to be the
476 presumed ancestral or wild type for each of these nine variants.

Locus	Reference Allele	Alternate Allele	Phenotype	Inheritance Pattern for Alternate Allele	Reference
EDNRB	*N*[*]	*O*	Overo spotting/OLWS	Dominant, homozygous lethal	Metallinos *et al.* 1998; Santschi *et al.* 1998; Yang *et al.* 1998
ECA3	*N*[*]	*TO*	Tobiano spotting	Dominant	Brooks *et al.* 2007
KIT	*N*[*]	*SB1*	Sabino spotting	Additive	Brooks & Bailey 2005
KIT	*N*[*]	*W5*	Variable white spotting	Dominant, presumed homozygous lethal	Haase *et al.* 2009
KIT	*N*[*]	*W10*	Variable white spotting	Dominant, presumed homozygous lethal	Haase *et al.* 2009
KIT	*N*[*]	*W20*	Variable white spotting	Dominant	Hauswirth *et al.* 2013
MITF	*N*[*]	*SW1*	Splashed white spotting	Dominant	Hauswirth *et al.* 2012
MITF	*N*[*]	*SW3*	Splashed white spotting	Dominant, presumed homozygous lethal	Hauswirth *et al.* 2012
PAX3	*N*[*]	*SW2*	Splashed white spotting	Dominant	Hauswirth *et al.* 2012
ASIP	*A*	*a*	Black/Bay pigment distribution	Recessive	Rieder *et al.* 2001
MC1R	*E*	*e*	Black/Red pigmentation	Recessive	Marklund *et al.* 1996

[*]For these loci the "*N*" designation is used by the UC Davis commercial testing service to communicate a wild type allele.

Table 2. Genotype counts and allele frequencies for nine white spotting variants and
two base coat color loci genotyped among 1054 APHA registered horses, within each of
the two registration categories. Data are presented as genotype counts for the
homozygous reference (Hom Ref), heterozygous (Het) and homozygous alternate (Hom
Alt) states, as well as the frequency of the reference (fRef) and alternate (fAlt) alleles.

	Regular Registry (n= 773)					Solid Paint-bred Registry (n= 281)					Total (n= 1054)				
	Hom Ref	Het	Hom Alt	fRef	fAlt	Hom Ref	Het	Hom Alt	fRef	fAlt	Hom Ref	Het	Hom Alt	fRef	fAlt
EDNRB*O (Frame Overo)	570	203	0	0.87	0.13	263	18	0	0.97	0.06	833	221	0	0.88	0.12
TO (Tobiano)	572	142	59	0.83	0.17	281	0	0	1.00	0.00	853	142	59	0.88	0.12
KIT*SB1 (Sabino1)	748	24	1	0.98	0.02	281	0	0	1.00	0.00	1029	24	1	0.99	0.01
KIT*W05 (Variable White)	773	0	0	1.00	0.00	281	0	0	1.00	0.00	1054	0	0	1.00	0.00
KIT*W10 (Variable White)	771	2	0	1.00	0.00	281	0	0	1.00	0.00	1052	2	0	1.00	0.00
KIT*W20 (Variable White)	404	319	50	0.73	0.27	196	74	11	0.83	0.17	600	393	61	0.76	0.24
PAX3*SW2 (Splashed White)	757	15	1	0.99	0.01	272	9	0	0.98	0.03	1029	24	1	0.99	0.01
MITF*SW1 (Splashed White)	680	90	3	0.94	0.06	263	17	1	0.96	0.07	943	107	4	0.95	0.05
MITF*SW3 (Splashed White)	769	4	0	0.99	0.01	281	0	0	1.00	0.00	1050	4	0	1.00	0.00
ASIP*A (Bay/Black)	316	323	134	0.62	0.38	115	118	48	0.62	0.38	431	441	182	0.62	0.38
MC1R*E (Black/Red)	78	234	461	0.25	0.75	9	82	190	0.16	0.82	87	316	651	0.23	0.77

489 Table 3: Horse designation to the Regular or Solid Paint-bred registry divisions by
490 genotype and genotype combinations (for simplicity, the table reports only
491 genotypes/combinations observed in at least three horses.)

	Genotype	Registry Category		% Regular
		Regular	Solid Paint-bred	
	N/N	97	172	36%
Single Allele Genotypes	N/SW2	7	3	70%
	W20/W20	24	10	71%
	N/W20	166	58	74%
	N/SW1	34	8	81%
	N/O	64	10	86%
	N/SW3	3	0	100%
	N/TO	97	0	100%
	TO/TO	57	0	100%
	N/SB1	8	0	100%
Multi-allele Genotypes	N/SW2, N/W20	1	3	25%
	N/SW1, N/SW2	5	2	71%
	N/SW1, N/W20	16	6	73%
	N/O, N/SW	7	1	88%
	N/O, N/W20	82	7	92%
	N/TO, N/SW1	5	0	100%
	N/TO, N/O	5	0	100%
	N/TO, N/W20	25	0	100%
	N/O, N/SB1	4	0	100%
	N/SB1, N/W20	5	0	100%
	N/O, W20/W20	19	0	100%
	N/SW1, W20/W20	4	0	100%
	N/TO, N/O, N/W20	7	0	100%
	N/O, N/SW1, N/W20	9	0	100%
	N/O, N/SW1, W20/W20	3	0	100%

512 Table 4: Horse counts for the Regular and Solid Paint-bred phenotypes compared to
513 the total number of white spotting pattern alleles at the nine loci tested, and genotypes
514 at the *MC1R* and *ASIP* coat color loci in a subset of 368 horses.

			Registry Category		
			Regular	Solid Paint-bred	% Regular
Total # of White Spotting Alleles (n= 1054)		*0*	97	172	36%
		1	382	79	83%
		2	240	29	89%
		3	51	0	100%
		4	3	0	100%
		5	1	0	100%
MC1R -ASIP Geno. (n=368)		*E/-, a/a*	13	7	65%
		E/-, A/-	57	23	71%
		e/e, -/-	219	49	82%

517
518

521 **Figure 1:** Anatomical landmarks on the face and limbs of the APHA horse are used to
522 determine the extent of white spotting patterns, and to designate each registered horse
523 to either the Regular or Solid Paint-bred sub-registries. To enter the Regular registry, a
524 proposed horse must possess at least 2" of white hair, as well as underlying
525 unpigmented skin, on the body surface beyond these reference lines.

528
529 **Figure 2:** Diverse white spotting phenotypes are captured by the American Paint Horse
530 Registry. Horse A is categorized in the Solid registry though he possesses the same
531 *EDNRB*O* white spotting genotype as horse E. Horses B-F are all classified as Regular
532 registry by photographic assessment (photo credits: J. Hein)

534
535 **Figure 3:** Previously thought to be lethal in the homozygous state, this carrier of the
536 *PAX3 SW2/SW2* homozygous genotype has extensive depigmentation of the skin and
537 coat (photo credit: Paint Horse Journal)

538