Pax2a regulates angiogenesis to facilitate mmp2-dependent basement membrane remodeling of the optic fissure.

Megan L. Weaver[1], Piedade. W. P[1], Meshram, N.N[1] and J.K. Famulski[1].

1. Department of Biology, University of Kentucky

ABSTRACT:

Vertebrate retinal development requires timely and precise fusion of the optic fissure (OF). Recent studies have suggested hyaloid vasculature to be involved in optic fissure fusion. In order to examine this link, we analyzed OF fusion and hyaloid vasculogenesis in the zebrafish pax2a^{noi} mutant line. We determined that OF basement membrane (BM) remodeling, normally preceded by F-actin accumulation is mis-regulated in pax2a$^{-/-}$ embryos. Comparing transcriptomic profiles of pax2a$^{-/-}$ and wildtype eyes we discovered a novel connection between regulation of angiogenesis and fusion. Pax2a$^{-/-}$ eyes exhibited a significant reduction of *talin1* expression, a regulator of hyaloid vasculature, in addition to increased expression of an anti-angiogenic protease, *adamts1*. Using 3D and live imaging we observed reduced OF hyaloid vascularization in pax2a$^{-/-}$ embryos. Additionally, pharmacological inhibition of VEGF signaling or *adamts1* mRNA overexpression phenocopied the pax2a$^{-/-}$ vasculature, F-actin and BM remodeling phenotypes. Finally, we show that hyaloid vasculature expresses *mmp2* which is necessary for remodeling the fissure BM. Taken together we propose a pax2a driven mechanism that restricts anti-angiogenic activity of adamts1 enabling hyaloid vasculature invasion of the OF and delivery of the BM remodeler mmp2.

INTRODUCTION:

Ocular development is a highly conserved process amongst vertebrate species. Assembly of the hemispherical, retinal structure from an initially flat sheet of cells requires many complex morphogenetic movements. One such morphogenetic movement involves the invagination of the optic vesicle which results in a fissure forming at the ventral region of the developing retina. This fissure, known as the choroid or optic fissure (OF), enables hyaloid vasculature cell migration into the developing retina and subsequent establishment of the hyaloid vasculature. Hyaloid vasculature is a temporary circulatory system required for ocular development, and in most cases will degenerate once mature blood vessels begin to grow (1-4). As soon as the hyaloid vasculature has been established, the two opposing retinal epithelial sheets of the optic fissure will undergo fusion. Thereby, they encase the ganglion cell axons localized in the optic stalk and complete retinal morphogenesis. Failure of optic fissure fusion leads to a congenital blinding disorder known as coloboma (5-7). Coloboma is a prevalent cause of pediatric blindness, accounting for approximately 10% of cases worldwide (6, 8). This makes it one of the leading causes of pediatric blindness. Coloboma is a spectrum disorder presenting unilaterally or bilaterally and ranging in severity from minor visual impairment, to complete blindness in the affected eye (9). This spectrum of severity is associated with the location and degree to which the choroid fissure was able to fuse and the severity of subsequent loss of ganglion cell axons (7).

Coloboma has been studied for many decades in many different species. Work over this time has led to a general outline of the signaling and morphogenetic pathways required for proper optic fissure formation and fusion (recently reviewed in (10)). In particular, opposing action of bone morphogenetic protein (BMP) and sonic hedgehog (Shh) signaling establishing the dorsal-ventral pattern of the optic vesicle and ensuring proper expression of optic stalk and OF regulators pax2, vax1 and vax2 (11, 12). However, the actual molecular mechanisms driving optic fissure fusion remains largely unknown. The process of epithelial tissue fusion is not unique to the eye and occurs throughout development, encompassing neural tube closure, palatal shelf formation and eyelid development (13). Epithelial fusion has been studied for over a century, and is

known to involve transcriptional regulation, cell signaling pathways and morphogen

gradients (14, 15). The actin cytoskeleton is thought to be a crucial component of the

machinery driving epithelial fusion in many tissues (16). The importance of the actin

cytoskeleton during epithelial fusion involves lamellipodial and filopodial projections

between the two opposing epithelia. These help to "zipper" the cells together to form a

single continuous sheet. When lamellipodial and filopodial projections are precluded,

epithelial fusion often fails (17). Lamellipodia and filopodia have been observed during

OF fusion almost 3 decades ago (18-20). However, the functional and regulatory

mechanisms behind these projections remain unknown. Another cellular mechanism

known to be directly involved in epithelial fusion is the remodeling of the basement

membrane (BM). During epithelial fusion, the BM acts as a physical barrier restricting

the establishment of cell-cell contacts, which must be removed in order to complete

fusion. Recent work from several labs, working on different species, has characterized

progressive removal of the BM during OF fusion (21-24). However, the molecular

mechanisms facilitating this process, in particular BM remodeling, also remain largely

unknown. It was recently suggested that hyaloid vasculature cells migrating through the

OF could potentially signal or facilitate BM remodeling (22). Hyaloid vasculature

nourishes the developing retina and lens while connecting to the choroid vasculature for

proper blood flow (3). Hyaloid vasculogenesis takes advantage of an open optic fissure

so that vasculature cells can migrate into the developing optic cup. Once optic fissure

fusion is completed, hyaloid vasculature is fully established. James *et al.* 2016, showed

that mutations in zebrafish *talin1*, an actin cytoskeleton scaffolding protein known to be

required for endothelial cell migration (25), result in OF fusion defects. Theirs, and

previous studies also indicated that *cloche* mutants, which lack all early hyaloid

vasculature, have delayed basement membrane breakdown in the region of the choroid

fissure (22, 26). Since the hyaloid vasculature requires an open fissure to complete

establishment of its network, it has been proposed that migrating hyaloid vasculature

cells may regulate the timing of fissure fusion. This mechanism could potentially involve

vasculature-mediated activation of the fusion machinery within the retinal rim cells, or

direct supply of molecular factors, such as matrix proteases (27). In additional support

of this hypothesis, recent optic cup transplantation experiments in zebrafish embryos

confirm that in the absence of hyaloid vasculature, ectopic retinal OFs fail to initiate

fusion (28). Hence, there is a clear link between OF fusion and hyaloid vasculature.

In our current study, we have undertaken a detailed analysis of zebrafish OF

fusion comparing WT and the pax2a$^{-/-}$ coloboma model (29). In particular, we have

characterized the timing of BM remodeling, cytoskeletal responses, morphological

apposition and hyaloid vascularization. Using transcriptomic analysis to compare WT

and pax2a$^{-/-}$ eyes we discovered mis-regulation in angiogenic signaling, resulting in

decreased OF hyaloid vascularization and OF fusion failure. In particular we found that

pax2a$^{-/-}$ embryos exhibit a decrease in *talin1* expression coupled with increased

expression of an anti-angiogenic protease, *a disintegrin and metalloproteinase with*

thrombospondin motifs 1 (adamts1), (30, 31). Modulation of vascularization via

pharmacological inhibition of VEGF signaling or overexpression of adamts1 both

phenocopied the pax2a$^{-/-}$ hyaloid vasculature and coloboma phenotypes.

Mechanistically, we also show that hyaloid vasculature is a source of mmp2, mmp14a

and mmp14b and that mmp2 activity is necessary for OF BM remodeling. Taken

together, we propose a novel pathway for the regulation of OF fusion where pax2a

mediates proper expression of *adamts1* in order to facilitate hyaloid vasculature

recruitment to the OF and subsequent vasculature supplied mmp2-dependent BM

remodeling.

RESULTS:

Optic fissure basement membrane remodeling is preceded by F-actin

accumulation.

Several recent studies have undertaken a detailed time course to map out the

exact timing of optic fissure fusion in numerous species, including zebrafish (21-24).

Overall, in zebrafish, the data point to ~32-36hpf as the time of OF fusion initiation, as

observed by BM remodeling. To decipher the molecular mechanisms regulating OF

fusion we also performed a detailed time course analysis of OF BM remodeling using

laminin immunohistochemistry (IHC) and additionally analyzed F-actin. Our goal was to

determine whether changes in F-actin levels in the OF were correlative with fusion.

James *et al*, 2016 had recently suggested that infiltrating vasculature endothelial cells

migrating through the fissure could be a source of signal for fissure fusion. In fact, they

have shown that interactions between vasculature and the fissure resulted in an

increase of F-actin (22). As such, we performed whole mount IHC for laminin deposition

in embryos starting at 24hpf. We sampled every 4 hours up to 48hpf and then at 56 and

72hpf (Fig 1A). To analyze the progression of the fusion process, we quantified the

laminin signal within the OF from 3D confocal scans. Regions of the BM outside the OF

and juxtaposed to the developing lens were used to normalize the laminin signal (Fig

S1A). We quantified distal, medial and proximal regions of the fissure (Fig S1B,C).

When comparing results from the three regions we did not observe significant

differences in the distribution of laminin or F-actin. Going forward we focused on

quantifying laminin signal in the central proximal region (medial) of the fissure (Fig S1B).

This region has been previously shown to represent the site of fusion initiation and is

easily identified in 24-72hpf embryos. In agreement with recent studies in zebrafish, we

found that OF fusion initiates at ~32-36hpf and does so in a central region of the fissure,

subsequently proceeding proximally and distally (Fig 1B). By 48hpf, we observed most

of the laminin signal was removed from the OF, and by 72hpf, little to no laminin

persists in the region (Fig 1A, B).

To track F-actin levels during fusion we stained the laminin labeled embryos with

phalloidin and quantified the signal within the fissure. To normalize phalloidin staining

intensity we used an interior region of the retina (Fig S1A). Our analysis revealed an

increase in OF associated F-actin signal between 24-32hpf, preceding the time that BM

remodeling is initiated (Fig 1A,B). We also noticed a second increase in F-actin around

44hpf, a time of active fusion (Fig 1C). The observed increase of F-actin signal

correlates with a) the timing of hyaloid vasculature migration (24-32hpf) into the fissure

and b) the expected timing of actin protrusions involved in epithelial fusion (36-48hpf).

When measuring distance between the fusing lobes of the retina we observed

apposition to complete around 36-40hpf (Fig S2). Taken together, we conclude that

both apposition and BM remodeling are preceded by an increase of F-actin within the

OF.

Optic fissure fusion mechanics are disrupted upon loss of pax2a function.

In order to test our theory that an increase in F-actin is involved in the initiation of OF fusion we compared our findings to an established model of coloboma, the pax2a[noi] line (29). Pax2 is required for OF fusion in several model systems and has been documented in human coloboma cases (32-34). The noi mutation is predicted to result in a loss of pax2a function due to a premature stop codon at position 198 (35). Originally characterized for their no-isthmus phenotype, pax2a[noi/noi] embryos (referred to as pax2a$^{-/-}$ from this point on) elicit a fully penetrant unfused OF. Unfortunately, homozygous mutants are not viable and do not enable the study of coloboma at juvenile or adult stages. We hypothesized that examining the molecular events leading to OF fusion in the pax2a[noi] system would inform us whether these events are functionally important. Similar to our WT study, we examined apposition, laminin and F-actin levels throughout the time of OF fusion (Fig 2A, Fig S1C). In stark contrast to WT, pax2a$^{-/-}$ embryos did not exhibit a significant decrease in laminin signal between 32-48hpf. In fact, as expected, laminin appears to be largely retained in the OF of pax2a$^{-/-}$ embryos up to 72hpf (Fig 2B). Previous studies in pax2a null mice had indicated failure of BM remodeling and retention of laminin (34, 36, 37). While there is a moderate decrease in laminin signal in pax2a$^{-/-}$ embryos from 32-36hpf, this result is significantly different from WT. IHC images clearly show that laminin persists in the fissure up to and including 72hpf (Fig 2A). When examining OF F-actin levels in pax2a$^{-/-}$ embryos we did not detect the expected increase between 24 and 32hpf (Fig 2C). We observed F-actin levels fluctuating between 36 and 44hpf and finally dropping to WT amounts by 48hpf. Similar results were also observed when quantifying distal and proximal regions of the fissure (Fig S1D). Our data therefore suggest that the absence of OF F-actin accumulation between 24 and 32hpf, correlates with failure to initiate fusion. Lastly, when examining retinal lobe distance in pax2a$^{-/-}$ embryos we did not observe any significant defects in the timing of apposition (Fig S2). Taken together, we conclude that in the absence of pax2a F-actin fails to accumulate in the OF and this correlates with the failure to remodel the BM. Based on our analysis of apposition, we also conclude that pax2a does not regulate morphogenetic movements of the retinal lobes. In summary, we

propose that accumulation of F-actin is necessary for the initiation of OF BM

remodeling. We therefore next sought to investigate the mechanism of OF F-actin

accumulation.

Pax2a retinal transcriptomic profile during optic fissure fusion.

Although pax2 has been studied in many systems, there are currently no obvious

transcriptional targets of pax2 that would directly point to mechanistic regulation of OF

fusion. In an effort to understand the absence of OF BM remodeling and mis-regulation

of F-actin observed in pax2a$^{-/-}$ embryos we compared whole eye transcriptomic profiles

between WT and pax2a$^{-/-}$ embryos. Eyes from 48hpf WT and pax2a$^{-/-}$ embryos were

isolated and total RNA was subsequently purified and sequenced using Illumina

sequencing. 48hpf was chosen for our experimental time point as pax2a$^{-/-}$ embryos are

easily phenotyped at this age thanks to observable heart malformations. When

comparing three replicates for WT and pax2a$^{-/-}$ embryos using RSEM software, we

detected 1215 transcripts significantly upregulated (>95% confidence interval) and 1202

transcripts significantly downregulated (Fig 3A and 3D). Gene ontology analysis

indicated a wide spread of biological function being affected, including cytoskeletal

signaling, adhesion and developmental processes (Fig 3B, E). In fact, the distribution of

biological function between up and downregulated genes was highly similar (Fig 3B, D).

A list of the top 20 up and down regulated genes is outlined in Figure 3C and F. Results

of all the statistically significant up and downregulated genes identified in our assay are

presented as supplementary data (Table 1 and Table 2).

When examining our list of downregulated targets, one, *talin1* (tln1), stood out in

particular. Not only was *tln1* one of the most downregulated targets (~200 fold

decrease) it was also recently shown to be necessary for the migration of hyaloid

vasculature cells into the OF and subsequent fusion (22). As such, we next sought to

investigate the relationship between pax2a, tln1, vasculogenesis and optic fissure fusion

mechanics.

Loss of pax2a function leads to reduced hyaloid vasculature within the optic

fissure.

To validate our RNAseq data, we performed wholemount *in situ* hybridization

(WISH) for *tln1* comparing WT siblings to pax2a$^{-/-}$ mutant embryos. WT expression of

tln1 was observed in the optic fissure between 28 and 48hpf coinciding with pax2a

expression (Fig 4A, Fig S3). In pax2a$^{-/-}$ embryos, OF *tln1* expression appears

significantly reduced compared to WT while retaining similar expression in periocular

regions (Fig 4A). *Tln1* expression is also reduced in the mid brain-hind brain boundary,

another region of strong *pax2a* expression (Fig 4A, S3A). Our WISH data was further

supported by qPCR results showing a significant decrease in *tln1* expression at 32hpf

(Fig 4B). To determine whether the reduction of *tln1* expression in pax2a$^{-/-}$ embryos had

functional consequences on hyaloid vasculogenesis we examined pax2a$^{-/-}$

Tg[kdrl:mCherry] embryos. Using 3D *in vivo* time-lapse confocal microscopy we

recorded migration of mCherry expressing cells through the OF from 24 to 30hpf (Fig

4B, Movie 1). At 24hpf, both WT and pax2a$^{-/-}$ embryos contain mCherry expressing cells

within the fissure. However, over the next six hours of imaging it is apparent that pax2a^{-}

$^{/-}$ embryos have significantly fewer mCherry expressing cells pass through the OF (Fig

4C, Movie 1). To quantify this effect, we fixed WT and pax2a$^{-/-}$ Tg[kdrl:mCherry]

embryos at 24, 32, 36 and 48hpf, collected 3D confocal stacks and counted the number

of mCherry positive cells found within the OF (Fig 4D). The data indicate that in pax2a$^{-/-}$

embryos there is a significant reduction in the number of mCherry expressing cells at

both 32 and 36hpf (Fig 4E). Furthermore, using 3D rendering, we noted that in 48hpf

pax2a$^{-/-}$ embryos the hyaloid vasculature established in the back of the lens is reduced

in size and lacks proper connections to the newly forming choroidal and superficial

vasculature systems (Movie 2-3). Overall, our data indicate that pax2a$^{-/-}$ embryos exhibit

decreased expression of *tln1* and impaired hyaloid vascularization of the optic fissure.

<u>Inhibition of VEGF signaling impairs optic fissure fusion mechanics.</u>

Based on our discovery of impaired OF hyaloid vasculature in pax2a$^{-/-}$ embryos,

we next examined whether this phenomenon is associated with failure of optic fissure

fusion. Hence, we turned our attention to vascular endothelial growth factor (VEGF)

signaling. VEGF, the ligand for vascular endothelial growth factor receptor (VEGFR), is

a prime candidate for regulating the migration and proliferation of hyaloid vasculature

cells. In support of this notion, when examining expression patterns for VEGF ligands,

vegfaa and ab, ba, c and d, we found *vegfaa, ab* and *c* to be expressed in the retina and

periocular regions (Fig S4A). To inhibit VEGF signaling, we took advantage of the

dorsomorphin derivative DMH4, which has been shown in zebrafish to selectively inhibit

VEGF signaling independent from BMP (38). Based on published working

concentrations, we conducted a dose response to examine DHM4 effects on hyaloid

vasculature using Tg[kdrl:mCherry] as a readout (38). Treatment of embryos from 12-

24hpf, ranging from 1-100μM, resulted in a dose dependent reduction of mCherry signal

in the developing retina (Fig S4B). We decided to use the 100μM concentration for

subsequent experiments as this concentration was able to completely inhibit

vascularization of the retina up to 72hpf without any significant impact on overall embryo

health (Fig 5A, S4C). Embryos were treated starting at 12hpf and examined for fissure

fusion status via whole mount laminin IHC at 24, 32, 36, 48, 56 and 72hpf (Fig 5B). 3D

confocal imaging revealed a persistence of laminin signal within the fissure up and

including 72hpf in all embryos examined (Fig 5B). Measurement of the laminin signal

generated very similar results to what we observed in pax2a$^{-/-}$ embryos (Fig 5C). This is

highlighted by retention of laminin signal at 48hpf and persistence of laminin in 100%

(n=49) of embryos treated up to 72hpf (Fig 5B-C). In addition to laminin, we also imaged

and quantified OF F-actin levels during the treatment (Fig 5B). Similar to pax2a$^{-/-}$

embryos DMH4 treatment prevented the accumulation of OF F-actin between 24-32hpf

(Fig 5D). This was in contrast to DMSO treated embryos which exhibited the

accumulation of F-actin in the fissure between 24-32hpf like wildtype embryos. DMH4

treatment did not affect the size of the retina nor the apposition of retinal lobes (Fig S2).

Lastly, we also observed a significant decrease of *tln1,* but not *pax2a,* expression in

DMH4 treated embryos (Fig S4E,F). This suggests that *tln1* is likely expressed by the

vasculature cells within the OF. Based on these findings, we conclude that inhibition of

VEGF signaling results in failure of F-actin accumulation and basement membrane

breakdown, presumably due to the absence of hayloid vasculature cells in the OF.

Anti-angiogenic protease *adamts1* is upregulated in OF of pax2a mutant embryos.

Having established a connection between pax2a function and recruitment of
hyaloid vasculature cells into the optic fissure, we next sought to examine the
mechanism behind it. We therefore re-examined our transcriptomic comparison of WT
and pax2a$^{-/-}$ eyes, with a focus on angiogenic regulation. This new examination
revealed misexpression of *a disintegrin and metalloproteinase with thrombospondin*
motifs 1 (adamts1), a secreted protease known to be involved in regulating VEGF
signaling (39). Being a member of the ADAMTS family, adamts1 contains three
thrombospondin motifs enabling it to directly bind and sequester VEGF (31).
Furthermore, the protease activity of adamts1 has bene shown to target
thrombospondin 1 and 2 for cleavage, ultimately liberating their active forms which bind
to and block VEGF from binding the VEGF receptor (30). Our RNAseq data indicated
that *adamts1* expression was upregulated almost two-fold in pax2a$^{-/-}$ eyes (table 1). This
would suggest that the expression of *adamts1*, and therefore its anti-angiogenic activity,
is normally kept in check by pax2a. Pax2a has not been reported to harbor
transcriptionally repressive function, so this may involve an indirect mechanism. To
confirm our RNAseq results we performed WISH, examining *adamts1* expression at 28
and 36hpf. In agreement with our transcriptomic analysis, we observed an upregulation
in *adamts1* in the ventral region of pax2a$^{-/-}$ eyes, specifically surrounding the optic
fissure (Fig 6A). A statistically significant increase in *adamts1* expression was also
confirmed using qPCR (Fig 6B). To test whether upregulation of *adamts1* expression
would have direct effects on hyaloid vascularization and optic fissure fusion we injected
Tg[*kdrl*:mCherry] embryos with *adamts1* mRNA. We subsequently examined hyaloid
vasculature using 3D confocal imaging between 24-48hpf. Injection of *adamts1* mRNA
resulted in a significant decrease in the number of *kdrl*:mCherry cells found within the
optic fissure (Fig 6B, C, Movie 5). The observed decrease mirrored the results from
pax2a$^{-/-}$ embryos (Fig 6C). Our findings therefore suggest that *adamts1* expression
needs to be tightly regulated in order to ensure proper vascularization of the retina,
including the OF. Furthermore, the observed up-regulation of *adamts1* expression in
pax2a$^{-/-}$ embryos may explain the resulting OF hyaloid vasculature deficiency.
335
Overexpression of *adamts1* inhibits optic fissure fusion mechanics.

Having mimicked the vasculature phenotype of pax2a mutants by over

expressing *adamts1*, we next determined whether it had any effects on OF fusion. To

do so, we performed whole mount laminin and F-actin staining in *adamts1* mRNA

injected embryos at 24, 32, 36, and 48hpf (Fig 7A). Confocal imaging of the fissure

indicated that *adamts1* mRNA injection also phenocopied pax2a$^{-/-}$ associated

persistence of laminin and therefore failure of optic fissure fusion. Quantified levels of

laminin persisting at 48hpf was similar to pax2a$^{-/-}$ embryos (Fig 7B). Similar to pax2a$^{-/-}$

and DMH4 treated embryos, we again noted a lack F-actin accumulation between 24-

32hpf in *adamts1* overexpressing embryos (Fig 7C). As was observed with pax2a$^{-/-}$,

timing for apposition of the lobes was not affected in adamts1 overexpressing embryos

(Fig S2). Taken together, our data suggest that maintaining proper levels of *adamts1*

expression is necessary to enable VEGF signaling and subsequent recruitment of

hyaloid vasculature required for initiation of OF BM remodeling.

<u>Hyaloid vasculature is the source of mmp2 necessary for OF BM remodeling.</u>

The above data confirm and support a model where hyaloid vasculature drives or

initiates the OF fusion process. However, a missing key to this model is the mechanism

by which vasculature cells induce fissure fusion. James *et al*, along with others, have

suggested vasculature cells may be a source of BM remodeling enzymes, such as

matrix metaloproteases (mmp). To investigate this further, we used WISH to examine

mmp expression within the OF between 24-48hpf. Our data indicated that *mmp2, 14a*

and *14b* were expressed within the fissure between 28-36hpf (Fig S5 and Famulski lab

unpublished data). Mmp2 had recently been associated with OF fusion in the mouse

while evidence from an mmp2 metalloproteinase activity probe in zebrafish indicated

mmp2 activity is present in the developing eye and likely OF (40, 41). mmp14 is an

activator of mmp2 and its co-expression with mmp2 within the fissure suggests mmp2 is

in fact active (42, 43). To test whether mmp2 fits within our model we assayed *mmp2,*

mmp14a and mmp14b expression in pax2a$^{-/-}$ and DMH4 treated embryos. In both

cases, OF expression of *mmp2, mmp14a and mmp14b* was reduced as observed by

WISH and confirmed by qPCR (Fig 8A, B). This suggested that all three are expressed

by the hyaloid vasculature cells. To confirm this, we performed 2 color WISH for *mmp2*

and *kdrl*. Confocal imaging verified that *mmp2* expression is co-localized with that of

kdrl within the OF (Fig 8C). Conversely, when performing 2 color WISH for *mmp2* and

rorB, a retina specific probe, we did not detect co-localization of the signals. We

therefore conclude that the source of mmp2, 14a and 14b enzymes in the OF are

hyaloid vasculature cells. Finally, to determine whether mmp2 activity is necessary for

OF fusion we treated embryos with ARP101, an mmp2 specific inhibitor previously

shown to be effective in zebrafish (44). Treating embryos from 24-48hpf inhibited OF

BM remodeling in a dose dependent manner (Fig 8D,E, S6). When compared to DMSO,

embryos treated with 20μM ARP101 either completely or partially retain their OF BM up

to 48hpf. Mmp2 activity is therefore necessary for BM remodeling in the OF. To

determine when mmp2 activity is required we began the ARP101 treatment between

24-32hpf and assayed BM status at 48hpf (Fig 8F). Treatments started later than 30hpf

had little to no effect on OF BM remodeling, indicating that mmp2 activity is required

between ~24-30hpf (Fig 8F). This finding also correlates with the observed timing of

mmp2 expression in the OF (Fig S5). However, because we cannot determine exactly

how quickly ARP101 inhibits enzymatic activity in our assay, active mmp2 may persist

in the OF longer than 30hpf.

DISCUSSION:

Studies of optic fissure fusion dating back several decades have been suggesting a

direct connection between the fusion process and hyaloid vasculature found within the

fissure. In fact, this hypothesis has been recently strengthened by data showing that a

reduction of hyaloid vasculature in the fissure, or removal of optic vesicles from sources

of vasculature inhibits or significantly delays fusion (22, 28). In our study, we have

characterized a pax2a driven mechanism that ensures proper vascularization of the OF

by restricting anti-angiogenic activity of adamts1 to enable hyaloid vasculature invasion

of the OF and delivery of BM remodelers mmp2, 14a and 14b. In conclusion, our

findings further validate the notion that hyaloid vasculature is an active component of

the machinery driving OF fusion.

Several recent reports, including this one, have comprehensively characterized optic

fissure fusion timing (21-24). Work from zebrafish, mice and chick all point to an orderly

progression involving: **1)** retinal growth and cellular rearrangement leading to nasal and

temporal retinal lobe apposition, **2)** invasion of the fissure by endothelial and neural

crest cells forming the hyaloid vasculature system, **3)** cellular signaling, either between

retinal rim cells or between rim cells and the migrating vasculature cells, **4)** remodeling

or removal of the basement membrane to enable physical connection of the rim cells

and subsequent formation of a continuous retinal epithelial sheet via re-polarization and

cell-cell adhesion. Step 1 has been nicely characterized in a few recent publications

outlining the flow of retinal cells and morphological formation of the fissure (45-47). For

step 2, several reports have carefully characterized the formation of the hyaloid

vasculature system, including migration of hyaloid vasculature precursor cells into the

fissure as soon as it forms (1-3). Importantly, perturbation of this process, or removal of

the developing eye from its source, has been shown to lead to fissure fusion failure (22,

28). To date, steps 3 and 4 are the least understood. Recent work from our lab has

characterized the composition of core BM components within the fissure (23).

Additionally, work from other labs has identified several molecular components

associated with cell-cell adhesion and epithelial sheet fusion to function within the

fissure, including α-catenin, n-cadherin, and netrin (22, 24, 48-51). However, the timing

and the molecular mechanism organizing and regulating these components remains

uncharacterized. Finally, the elephant in the room has always been the identity of the

BM remodeling mechanism. To date, only adamts16 has been functionally examined in

context of fissure fusion, mmp2 activity was detected during mouse OF fusion and

mmp23bb was implicated in the fusion process from transcriptomic data (41, 50, 52).

While attempting to address the mechanistic aspects of steps 3 and 4 using the

pax2a[noi] model we first uncovered a relationship between F-actin and BM remodeling.

Work by *James et al. 2016* suggested that F-actin accumulation may be indicative of

hyaloid cell interaction with retinal rim cells. During our detailed time course analysis,

we discovered that BM remodeling is in fact preceded by an increase in F-actin signal

within the OF (Fig 1). When assayed in pax2a[-/-] embryos the F-actin accumulation is

absent while the hyaloid vasculature in the OF is also diminished. The timing of F-actin

accumulation coincides with the active migration of hyaloid vasculature cells through the

fissure (Fig 4). Furthermore, we show pharmacologically, via VEGF inhibition, that

hyaloid vasculature is necessary for the accumulation of F-actin and OF BM

remodeling. Transcriptionally, we show that the decrease in hayloid vasculature in

pax2a$^{-/-}$ embryos coincides with a significant decrease in *tln1* expression. Tln1 is a key

regulator of endothelial cell migration, recently shown to be directly involved in OF

fusion (22). In fact, loss of tln1 function has been previously shown to decrease in

fissure associated vasculature and inhibit OF BM remodeling (22). We also show a

significant decrease in *tln1* expression upon VEGF inhibition (DMH4 treatment),

suggesting that *tln1* expression within the OF is associated with hayloid vasculature.

Talin1 is known to be a direct link between the actin cytoskeleton and the BM via

integrin (53, 54), and has been associated with the formation of adhesion junctions (55,

56). One could therefore hypothesize several different models for its role in optic fissure

fusion. We predict that tln1 is regulating the ability of hayloid vasculature cells to migrate

properly through the OF, however we cannot rule out the possibility that tln1 directly

participates in the OF fusion process. The absence of tln1 expression is most likely a

correlation to the decrease of hayloid vasculature cells within the fissure of pax2a$^{-/-}$

embryos. Our transcriptomic analysis was performed using whole eye tissue, which

would have included the hayloid vasculature. As such, the decrease in *tln1* expression

is likely indicative of decreased numbers of hayloid vasculature cells which require *tln1*

expression for proper migration into the fissure and or proliferation.

Our study strongly supports the notion that vasculature plays an integral part in OF

fusion. As such, modulation of angiogenesis represents a potential mechanism to

ensure proper vascularization of the OF. In our transcriptomic comparison of WT and

pax2a mutants we did in fact discover a connection to angiogenesis. In the absence of

pax2a function we observed an upregulation of *adamts1*. Examining *adamts1*

expression in the eye further confirmed an upregulation of expression in the ventral

regions pax2a$^{-/-}$ retinas. Encoding 3 thrombospondin (thsb) motifs able to directly bind

and sequester VEGF, in addition to targeting thsb1 and 2 for cleavage into their active

VEGF inhibiting form, adamts1 is well known for it's anti-angiogenic function in cancer,

aortic and renal biology (57). In our study, we show that upregulating adamts1

expression reduces the number of hyaloid vasculature cells in the OF. Up regulation of

adamts1 also prevents the accumulation of F-actin and subsequent BM remodeling.

Taken together we propose that the reduction in hayloid vasculature observed in

pax2a$^{-/-}$ embryos likely stems from an upregulation of adamts1 leading to suppressed

VEGF signaling and therefore limited vascularization of the OF.

The last aspect of our proposed model pertains to the mechanism of OF BM

remodeling. It had been suggested over the years that vasculature may be the source of

BM remodeling activity during OF fusion. To that end, we have discovered that *mmp2*,

mmp14a and *mmp14b* are all expressed within the hayloid vasculature during OF

fusion. In fact, timing of their expression matches precisely with the initiation of BM

remodeling. When we examine pax2a mutants or inhibit hayloid vasculature completely

we no longer detect expression of either of these mmps. Furthermore, using ARP101, a

specific mmp2 inhibitor, we show that mmp2 activity is necessary for OF BM remodeling

and this activity is specifically required between 26-32hpf. Mmp2 has been implicated

in OF fusion previously. Mouse studies have shown mmp2 expression within

macrophages residing in the OF. Furthermore, recent examination of mmp2 activity,

using a reporter construct, indicates that mmp2 is active in the eye and likely the OF

(40). Mmp14 is a known to activate mmp2 activity by cleaving the inhibitory pro-peptide

of mmp2. As such, co-expression of mmp2 and 14 is further indicative of mmp2 playing

a functional role in OF BM remodeling.

To date, the only BM remodeling enzyme to be associated with OF fusion is

adamts16. Loss of adamts16 function in zebrafish, via morpholino, led to a coloboma

like phenotype which the authors credited to the inability to degrade laminin in the

fissure (52). We have not been able to reproduce adamts16 expression within the

fissure. Recent studies in mice, zebrafish and chick have compared expression in OF

cells pre, during and post fusion (24, 50, 58, 59). Surprisingly, none of these studies

identified any obvious candidates for carrying out BM remodeling, including mmp2 and

mmp14. However, it remains possible that the OF tissue examined lacked the hayloid

vasculature and therefore prevented the identification of mmp2 and mmp14 as

candidates. Lastly, expression of *mmp2, 14a* and *14b* is transient in the fissure which
may have contributed to their OF expression being missed by other groups. Our data
indicate mmp2 activity is necessary for BM remodeling of the OF, but does not rule out
a role for additional proteases. There may be several proteases involved, such as
adamts16, mmp2 and others yet inhibition of just one could trigger failure of the
remodeling process and lead to fissure fusion failure. Continuing studies into matrix
protease activity will be needed to fully characterize the remodeling mechanisms during
OF fusion.
499
In conclusion, we present a new model for the mechanism of OF BM remodeling
where pax2a restricts expression of *adamts1* in the ventral retina to enable hayloid
vasculature invasion of the OF. Once in the fissure, vasculature cells express *mmp2,*
14a and *14b* to initiate BM remodeling. As soon as fusion begins, vasculature becomes
restricted from the fissure and ultimately the two retinal lobes fuse. Our model aligns
with the molecular events observed in the fissure, including accumulation of F-actin at
the time of vasculature migration through the fissure (24-28hpf), expression of mmps
(26-36hpf) and subsequent BM remodeling (32-48hpf). For future studies it will be
important to understand how pax2a regulates *adamts1* expression and to examine
whether mutations in *adamts1*, *mmp2* or *mmp14* are potential markers for coloboma.
510

 FIGURE LEGENDS:

 Figure 1: **An increase in F-actin dynamics precedes laminin remodeling during**
 optic fissure fusion.
 A) Whole mount immunohistochemistry was used to simultaneously visualize F-actin
 (red) and laminin (green) during optic fissure fusion, 24-72hpf. Central-proximal sections
 obtained using confocal imaging were collected and quantified. Scale bar = 50um. **B)**
 Quantification of laminin signal intensity within the optic fissure, normalized to regions of
 laminin staining juxtaposed to the lens. Relative pixel intensities are displayed. The
 green dotted line depicts the trend in laminin intensity over time. ANOVA $p < 0.0001$. **C)**
 Quantification of F-actin intensity (phalloidin staining) within the optic fissure, normalized
 to regions of F-actin signal within the lobe of the retina. Relative pixel intensities are
 displayed. The blue dotted line represents the trend in F-actin intensity over time.
 ANOVA $p < 0.0001$.

 Figure 2: **Pax2 mutant embryos lack F-actin activity increase and laminin**
 remodeling within the optic fissure.
 A) Whole mount Immunohistochemistry was used to simultaneously visualize F-actin
 (red) and laminin (green) during optic fissure fusion, 24-72hpf, in pax2$^{-/-}$ embryos.
 Central-proximal regions of the optic fissure are displayed. Scale bar = 50um **B)**
 Quantification of laminin signal intensity within the central-proximal region of the optic
 fissure, normalized to regions of laminin staining juxtaposed to the lens. Relative pixel
 intensities are displayed. ANOVA $p < 0.0001$. **C)** Quantification of F-actin intensity
 (phalloidin staining) within the central-proximal region of the optic fissure, normalized to
 regions of F-actin signal within the lobe of the retina. Relative pixel intensities are
 displayed. The blue dotted line represents the trend of F-actin intensity observed in WT
 embryos. ANOVA $p < 0.0001$.

 Figure 3: **Retinal transcriptomic comparison between WT and pax2a$^{-/-}$.**
 A) Heat map representing transcripts found to be significantly down regulated in pax2a$^{-/-}$
 embryos. **B)** Gene ontology for pax2a$^{-/-}$ downregulated transcripts. **C)** Top 20

downregulated transcripts are outlined in order of fold change. **D)** Heat map

representing transcripts found to be significantly upregulated in pax2a$^{-/-}$ embryos. **E)**

Gene ontology for pax2a$^{-/-}$ upregulated transcripts. **F)** Top 20 upregulated transcripts

are outlined in order of fold change.

<u>Figure 4:</u> **Pax2a is necessary for *tln1* expression and recruitment of vasculature**

into the optic fissure.

A) Wholemount in situ hybridization comparing *tln1* expression between WT and

pax2a$^{-/-}$ embryos at 28 and 32hpf. *tln1* signal within the optic fissure (yellow arrowhead)

appears reduced in Pax2a$^{-/-}$ embryos (red *). **B)** qPCR analysis of *tln1* expression in WT

and pax2a-/- embryos at 32hpf confirms a reduction in *tln1* expression. **C)** *In vivo* 4D

confocal imaging of Tg[kdrl:mCherry] WT or pax2a$^{-/-}$ embryos. Time lapse series

depicting the region of the optic fissure (dotted white lines) and mCherry positive

vasculature endothelial cells migrating through the fissure from 24-29hpf. Scale bar =

10um. **D)** Comparison of WT and pax2a$^{-/-}$ vascularization during optic fissure fusion, 24-

48hpf. 3D reconstructions of whole mount Tg[kdrl:mCherry] (red) WT or pax2a$^{-/-}$

embryos with DAPI (blue) stained DNA. Scale bar = 50μm. **E)** Quantification of the

number of mCherry positive cells from 3D confocal stacks within the region of the optic

fissure. Individual embryo results are depicted. ANOVA p<0.0001.

<u>Figure 5:</u> **Inhibiting angiogenesis prevents optic fissure fusion.**

A) 3D confocal imaging of Tg[kdrl:mCherry] (red) embryos treated with DMSO or 100μM

DMH4 between 24-72hpf. DNA was stained with DAPI (blue). Scale bar = 50μm. DMH4

treatment results in the absence of hyaloid vasculature the fissure and retina. **B)** Whole

mount Immunohistochemistry was used to simultaneously visualize F-actin (red) and

laminin (green) in DMH4 treated embryos from 24-72hpf. Central-proximal regions of

the optic fissure are displayed. Treatment with DMH4 results in retention of laminin and

optic fissure fusion failure. Scale bar = 50μm. **C)** Quantification of laminin signal

intensity within the central-proximal region of the optic fissure, normalized to regions of

laminin staining juxtaposed to the lens. Relative pixel intensities are displayed. ANOVA

p<0.0001. **D)** Quantification of F-actin signal intensity (phalloidin staining) within the

central-proximal region of the optic fissure, normalized to regions of F-actin signal within

the lobe of the retina. Relative pixel intensities are displayed. A blue dotted line

represents the WT trend in F-actin intensity. ANOVA p<0.0001.

Figure 6: **Adamts1 inhibits retinal vascularization.**

A) Wholemount in situ hybridization comparing *adamts1* expression between WT and

pax2a$^{-/-}$ embryos at 28 and 32hpf. *adamts1* signal within the optic fissure (yellow

arrowhead) is appears increased in Pax2a$^{-/-}$ embryos (red *). **B)** qPCR analysis

confirms an increase in *adamts1* expression in pax2a-/- embryos. **C)** Comparison of WT

and adamts1 mRNA injected embryo vascularization during optic fissure fusion, 24-

48hpf. 3D reconstructions of whole mount Tg[kdrl:mCherry] (red) WT or adamts1 mRNA

injected embryos with DNA stained by DAPI (blue). Scale bar = 50μm. **D)** Quantification

of mCherry positive cells from 3D confocal stacks within the region of the optic fissure.

Individual embryo results are depicted. * p<0.001, ANOVA p<0.0001.

Figure 7: *Adamts1* **misregulation leads to failure of optic fissure fusion.**

A) Whole mount Immunohistochemistry was used to simultaneously visualize F-actin

(red) and laminin (green) during optic fissure fusion, 24-72hpf in adamts1 mRNA

injected embryos. Central-proximal regions of the optic fissure are displayed. Scale bar

= 50μm. **B)** Quantification of laminin signal intensity within the central-proximal regions

of the optic fissure, normalized to regions of laminin staining juxtaposed to the lens.

Relative pixel intensities are displayed. ANOVA p<0.0001. **C)** Quantification of F-actin

signal intensity (phalloidin staining) within the central-proximal regions of the optic

fissure, normalized to regions of F-actin signal within the lobe of the retina. Relative

pixel intensities are displayed. A blue dotted line represents the WT trend in F-actin

intensity over time. ANOVA p<0.0001.

Figure 8: **Hyaloid vasculature is the source of mmp2 activity required for optic**

fissure BM remodeling.

A) Wholemount in situ hybridization comparing *mmp2, mmp14a* and *mmp14b*

expression in WT vs pax2a$^{-/-}$ and DMSO vs DMH4 treated embryos at 32hpf. *Mmp2,*

14a and 14b signal within the optic fissure (yellow arrowhead) appears decreased in

pax2a$^{-/-}$ and DMH4 treated embryos. **B)** qPCR analysis confirms a decrease in

expression of *mmp2, 14a* and *14b* in both pax2a$^{-/-}$ and DMH4 treated embryos. **C)** Two

color wholemount in situ hybridization simultaneously examining *mmp2* and *kdrl*, or

mmp2 and *rorB* expression at 32hpf. DNA was stained with DAPI. Scale bar = 50μm.

Clear overlap of signal is observed for *mmp2* and *kdrl*, but not *mmp2* and *rorB*. **D)**

Whole mount Immunohistochemistry was used to visualize laminin (red) in ARP101

treated Tg[*rx3*:GFP] embryos at 48hpf. Central-proximal regions of the optic fissure are

displayed. **E)** Quantification of 48hpf ARP101 treated embryos for fissure fusion

(absence of laminin signal), partial fusion (partial retention of laminin) or failure to fuse

(retention of laminin throughout the fissure). n= 41 (10mM), 38 (15mM), 42 (20mM)

ANOVA p<0.0001. **F)** Quantification of 48hpf ARP101 treated embryos for fissure

fusion, partial fusion or failure to fuse after various treatment initiation times from 24-

32hpf. n= 42 (24-48hpf), 29 (26-48hpf), 29 (28-48hpf), 23 (30-48hpf), 37 (32-48hpf).

ANOVA p<0.0001.

Supplemental Material

Figure S1: **Optic Fissure quantification**

A) Graphical representation of the region of OF analyzed using confocal microscopy. **B**)

Sample image depicting regions of the OF (1), laminin normalization (2) and F-actin

normalization (3) used for signal intensity quantification**. C)** Quantification of laminin and

F-actin signal intensity in wild type embryos within the distal, medial and proximal

regions of the optic fissure, normalized to regions of laminin staining juxtaposed to the

lens and F-actin signal within the lobe of the retina. Relative pixel intensities are

displayed. ANOVA p<0.0001. **D)** Quantification of laminin and F-actin signal intensity in

pax2a$^{-/-}$ embryos within the distal, medial and proximal regions of the optic fissure,

normalized to regions of laminin staining juxtaposed to the lens and F-actin signal within

the lobe of the retina. Relative pixel intensities are displayed. ANOVA p<0.0001.

Figure S2: **Optic Fissure apposition measurements.**

Measurements of the distance between retinal lobes (apposition) in WT, pax2a$^{-/-}$, DMH4

treated or ADAMTS1 mRNA injected embryos at 48hpf. Measurements were made

using laminin staining as reference for edges of retinal lobes. Distance was measured

as pixels.

Figure S3: *pax2a* **and** *tln1* **expression during development**

A) Whole mount in situ hybridization of *pax2a* probe at 24, 48, 54 and 72hpf. Lateral

(top) and ventral (bottom) images depicting optic fissure expression are shown. Pax2a

expression persists in the fissure up to 54hpf. **B**) Whole mount in situ hybridization of

tln1 probe at 24, 48, 54 and 72hpf. Lateral (top) and ventral (bottom) images depicting

optic fissure expression are shown. *tln1* expression is detected in the optic fissure from

24-48hpf (yellow arrowheads).

Figure S4: **DMH4 dose response**

A) Whole mount in situ hybridization of *vegfaa, ab, ba, c* and *d* probes at 32hpf. **B)** 3D

reconstructions of 48hpf whole mount Tg[kdrl:mCherry] (red) embryos treated with 1, 5,

50 or 100uM DMH4. DNA stained with DAPI (blue). Increasing concentration of DMH4

eliminates mCherry expressing cells from the optic fissure and retina. Scale bar =

50μm. **C)** Brightfield images of DMSO or DMH4 treated embryos at 24, 48 and 72hpf.

D) Whole mount in situ hybridization of *pax2a* or *tln1* probe at 32hpf in DMSO or DMH4

treated embryos. OF expression is indicated with a yellow arrowhead. DMH4 treatment

does not appear to alter *pax2a* expression but eliminates *tln1* expression from the OF.

E) qPCR results for *tln1* expression in DMH4 treated embryos.

Figure S5: **Time course of *mmp2, mmp14a* and *mmp14b* ocular expression**

Whole mount in situ hybridization of *mmp2, mmp14a, mmp14b* probes at 24, 26, 30 32,

34, 36 and 38hpf. *Mmp2* expression is present in the optic fissure from 26-34hpf (yellow

arrowheads). *Mmp14a* expression is present in the optic fissure from 32-36hpf(yellow

arrowheads). *Mmp14b* expression is present in the optic fissure from 32-36hpf (yellow

arrowheads).

Figure S6: **ARP101 treatment dose response**

A) Brightfield images of DMSO and ARP treated embryos at 48hpf. Concentrations of

ARP101 exceeding 20uM result in toxic effects. **B)** Whole mount Immunohistochemistry

was used to visualize laminin (red) in ARP101 treated Tg[*rx3*:GFP] embryos at 48hpf.

Central-proximal and distal regions of the optic fissure are displayed.

Movie 1: WT Tg[kdrl:mCherry] migration within the optic fissure, 24-30hpf

Movie 2: pax2a$^{-/-}$ Tg[kdrl:mCherry] migration within the optic fissure, 24-30hpf

Movie 3: 3D rotation of WT Tg[kdrl:mCherry] embryo at 48hpf

Movie 4: 3D rotation of pax2a$^{-/-}$ Tg[kdrl:mCherry] embryo at 48hpf

Movie 5: 3D rotation of 48hpf Tg[kdrl:mCherry] embryo injected with ADAMTS1 mRNA

Movie 6: 3D rotation of 48hpf Tg[kdrl:mCherry] embryo injected with ADAMTS1

morpholino

Table 1: 95th percentile RNA sequencing comparison

Table 2: Raw data RNA sequencing comparison

Table 3: Primer sequences

MATERIALS AND METHODS:

Zebrafish and embryo maintenance

Zebrafish were maintained using husbandry procedures approved by University of

Kentucky IACUC committee. Embryos were kept at 28.5°C in E3 embryo media. AB and

TL strains were used as wild-type, Tg[*kdrl*:mCherry] transgenic line was used to

visualize retinal vascularization (60), Tg[*rx3*:GFP] was used to visualize retinal cells

(61).

Pax2^{noi} embryos were a gift from Dr. Gregory-Evans. Genotyping analysis was

conducted by amplifying the region of gDNA with the noi mutation using the forward

primer: 5'- CTCGCTCTGCCTCCATGATTG3-'and the reverse: 5'-

GGCACTGAAAGAGCACAGG -3'. The resultant 460bp amplicon was digested with

TaqI (NEB) which would recognize and digest the WT allele sequence but not the

mutant allele.

Immunohistochemistry (IHC)

Dechorionated embryos were fixed with 4% PFA in PBS at room temperature for 3h

and washed with PBST 4 times for 5 minutes. Embryos were then permeabilized with

Proteinase K, 30μg/mL 10 minutes for 24-28hpf, 50μg/mL 15-20 minutes for 32-48hpf

and 75μg/mL 20 minutes for 56-72hpf, washed 2 times in PBST for 5 minutes and

blocked overnight at 4°C with 10% sheep serum, 0.8% Triton X-100 and 1% BSA in

PBS. Primary mouse anti-laminin antibody (ThermoFisher – 1:100) in blocking buffer

(1% sheep serum, 1% BSA and 0.8% Triton X-100 in PBS) were incubated overnight at

4°C and washed 5 times in PBST for 15 minutes. Secondary antibody, goat anti-rabbit

(Alexa Fluor® 488 – Abcam – 1:1000), DAPI 1:1000, and phalloidin (Alexa Fluor® 555 -

1:50) were incubated overnight at 4°C in the dark. Tg[*kdrl*:mCherry] embryos were

treated with proteinase K as described above and stained with DAPI 1:1000 overnight.

The embryos were washed 2 times in PBST for 15 minutes and visualized using a

Nikon C2+ confocal microscope equipped with a 40X (1.15NA) water immersion

objective. Embryos were embedded in 1.2% low melting point agarose on glass bottom

35mm dishes (Fluorodish, World Precision Instruments). Images were captured in steps

of 3.5 microns for a total of 31.5 microns using Nikon Elements software. Image

adjustment, such as cropping and brightness/contrast was performed using Adobe

Photoshop.

<u>Analysis of Fluorescence signal</u>

Fiji software (https://fiji.sc) was used to measure the fluorescence intensity of laminin

and phalloidin signal from raw image data. In order to account for variability in staining,

normalization values were measured for laminin and actin pixel intensity where an area

directly outside of the choroid fissure was measured and a ratio was generated between

the two values (Fig S1B). In cases where the fissure edges were farther apart then the

size of the box used for analysis, the box was divided into two and used to measure

fluorescence intensity on each side of the fissure. Measured values from both boxes

was added and then normalized. For Tg[krdl:mCherry], 3D reconstructions of the optic

fissure were generated and individual cells were counted (from the opening of the OF

through the back of the lens).

<u>Stats</u>

Student's t-test was used to compare individual time points. One-way ANOVA was

used to analyze across treatments. Graphs are displayed as mean +/- standard

deviation. Analysis was performed using Prism8 graphing software (GraphPad).

<u>Total RNA Sequencing</u>

WT and Pax2$^{-/-}$ embryos were dissected and collected at 48hpf and RNA was

extracted from dissected eyes using trizol. Pax2a$^{-/-}$ embryos were phenotyped by

distinct heart defects only observed in pax2a[-/-] embryos, as verified by previous

genotyping experiments. Dissected eyes were suspended in 1 mL of trizol and sheared

with a 22-gauge needle. Samples were incubated at room temperature for 5 minutes,

then 200 μL of chloroform was added and vortexed for 1 minute. Samples were then

centrifuged at 12,000 g for 15 minutes at 4℃. Aqueous phase was removed and put into

fresh RNAse free tube. 3 μL of Glycoblue was added to samples and vortexed for 5

seconds. 500 μL of 100% Isopropanol was then added and vortexed for an additional 10

seconds. Samples then were incubated at room temperature for 10 minutes.

Subsequently, samples were centrifuged at 12,000 g for 10 minutes at 4℃. Supernatant

was removed and 1 mL of 75% EtOH was added and vortexed for 10 seconds. Samples

were then centrifuged at 7,500 g for 5 minutes at 4℃. Supernatant was removed and

samples then underwent a pulse centrifugation and any remaining supernatant was

removed. Samples were air-dried for 7 minutes at room temperature under a fume

hood. 20 μL of RNAse free ddH$_2$O was added to the samples and mixed until pellet

dissolved. Finally, samples were incubated at 60℃ for 12 minutes and stored at -80℃.

The RNA then underwent a DNase treatment using the DNA-free Kit. 2 μL of 10X

DNaseI Buffer, and 1 μL of rDNaseI was added to the entire 20 μL sample. Samples

were incubated at 37℃ for 20 minutes, and then 2 μL of DNase Inactivation Reagent

was added. Samples were then incubated at room temperature for 2 minutes, mixing

samples 3 times. Samples were centrifuged at 10,000 g for 1.5 minutes. Supernatant

was then removed and placed into fresh RNase free tube and stored at -80℃.

Purified total RNA was sent to Applied Biological Materials Inc. for Illumina

sequencing (PE150bp, 30-40 million reads/sample). Bioinformatic analyses were

completed using RSEM. Those transcripts which had a greater than 0.95 posterior

probability of being differentially expressed were considered significant. Panther

(http://pantherdb.org) was used for gene ontology and pie chart generation.

<u>Whole-mount in situ hybridization (WISH)</u>

Whole-mount in situ hybridization was performed as previously described (62). RNA

probes were generated using PCR with T7 promoter sequence linkers and

subsequently transcribed [DIG or FITC labeled] using T7 polymerase (Roche). Primer

sequences are all found in Table 3. Images were captured using a Nikon Digital sight

DS-Fi2 camera mounted on a Nikon SZM800 stereo scope using Elements software.

Dissected eyes from 24, to 72hpf embryos were mounted in 70% glycerol and imaged

under DIC using a Nikon TiE compound microscope equipped with a 20X (0.7NA)

objective and Elements software. Image adjustment, such as cropping and

brightness/contrast was performed using Adobe Photoshop.

qPCR analysis

32hpf embryos were anesthetized with tricane, tail tips were collected for genotyping

and the heads, dissected just posterior of the eyes, were fixed in RNAlater. After

genotyping, embryos corresponding to WT and pax2a-/- were pooled, 5-10 embryos,

and total RNA isolated using a RNAaqueous kit (Ambion). DMSO and DMH4 treated

embryos were harvested in the same fashion absent any genotyping. qPCR was

performed as previously described (63).

2 color fluorescent in situ hybridization

Fluorescent whole mount in situ hybridization was performed as previously described

(64). Images were collected using a Nikon C2+ confocal microscope with a 20X

(0.95NA) objective and images were adjusted for brightness and contrast using Adobe

Photoshop.

Live imaging analysis

Live imaging of Tg[kdrl:mCherry] embryos was conducted using a Nikon C2+ confocal

microscope equipped with at 20X (0.95NA) water immersion objective. 22hpf embryos

were imbedded in 1.1% low gelling agarose in 1-inch glass bottomed Flourodish cell

culture dishes (World Precision Instruments) and covered in embryo media, 3-amino

benzoic acidethylester (tricaine) to anaesthetize the embryos and 1-phenyl 2-thiourea

(PTU) to inhibit pigmentation. Z-stacks 75μm thick with a step size of 2.5μm were

captured over the course of 6 hours at 10 minute intervals. The time lapse data were

reconstructed in 3D using Elements software. Image adjustment, such as cropping and

brightness/contrast was performed using Adobe Photoshop. After imaging, embryos

were removed and genotyped.

<u>Inhibitor treatments</u>

Embryos were incubated in embryo media with 5, 25, 50, or 100 μM of DMH-4 (Sigma

Life Science) in DMSO starting at 12hpf. Fresh DHM4 containing media was added

every 12hpf for timepoints past 24hpf. For ARP101 (Tocris) treatments embryos were

dechorionated and treated with 10, 15, 20, 25 or 30μM ARP101 in fish water from 24 to

48hpf, unless stated otherwise.

<u>Cloning and mRNA synthesis.</u>

Full coding domain sequences for *adamts1* (NCBI reference sequence

XM_021475923.1), were amplified from 24hpf zebrafish cDNA using primers 5'-

CTCGAGTCAACAGGGAGTCAGATTGC-3' for *adamts1*, cloned into pGEMT

(Promega), digested with AsiSI/XhoI, BamHI/XhoI respectively and subsequently cloned

into pCS2+. All constructs were verified by sanger sequencing (Eurofinsgenomics).

mRNA was synthesized from linearized pCS2 constructs using SP6 mMessage

mMachine kit (Ambion) and purified using YM-50 Microcon columns (Amicon, Millipore).

<u>Ethics Statement</u>

The use of zebrafish in this study was approved by the University of Kentucky IACUC

committee, Institutional PHS Assurance #D16-00217 (A3336-01) with a protocol

number: 2015-1370.

ACKNOWLEDGEMENTS:

This work for supported by faculty start-up funds provided to JKF from the department

of Biology, University of Kentucky. MLW was supported by the Lyman T Johnson

scholarship from the University of Kentucky. WPP was supported by Brazilian National

Council for Scientific and Technological Development (CNPq) under grant number

202970/2014-0. We thank Dr. Jeramiah Smith for assistance with bioinformatic analysis

of our RNAseq data and members of the Famulski lab for helpful discussions.

REFERENCES:

1.   Hartsock A, Lee C, Arnold V, Gross JM. In vivo analysis of hyaloid vasculature morphogenesis in zebrafish: A role for the lens in maturation and maintenance of the hyaloid. Dev Biol. 2014;394(2):327-39.

2.   Hashiura T, Kimura E, Fujisawa S, Oikawa S, Nonaka S, Kurosaka D, et al. Live imaging of primary ocular vasculature formation in zebrafish. PLoS One. 2017;12(4):e0176456.

3.   Kaufman R, Weiss O, Sebbagh M, Ravid R, Gibbs-Bar L, Yaniv K, et al. Development and origins of zebrafish ocular vasculature. BMC Dev Biol. 2015;15:18.

4.   Yoshikawa Y, Yamada T, Tai-Nagara I, Okabe K, Kitagawa Y, Ema M, et al. Developmental regression of hyaloid vasculature is triggered by neurons. J Exp Med. 2016;213(7):1175-83.

5.   AS AL, Gregory-Evans CY, Gregory-Evans K. An update on the genetics of ocular coloboma. Hum Genet. 2019.

6.   Chang L, Blain D, Bertuzzi S, Brooks BP. Uveal coloboma: clinical and basic science update. Curr Opin Ophthalmol. 2006;17(5):447-70.

7.   Gregory-Evans CY, Williams MJ, Halford S, Gregory-Evans K. Ocular coloboma: a reassessment in the age of molecular neuroscience. J Med Genet. 2004;41(12):881-91.

8.   Stoll C, Alembik Y, Dott B, Roth MP. Congenital eye malformations in 212,479 consecutive births. Ann Genet. 1997;40(2):122-8.

9.   Huynh N, Brooks BP. Uveal coloboma. JAMA Ophthalmol. 2013;131(10):1274.

10.   Patel A, Sowden JC. Genes and pathways in optic fissure closure. Semin Cell Dev Biol. 2017.

11.   Take-uchi M, Clarke JD, Wilson SW. Hedgehog signalling maintains the optic stalk-retinal interface through the regulation of Vax gene activity. Development. 2003;130(5):955-68.

12.   Macdonald R, Wilson SW. Pax proteins and eye development. Curr Opin Neurobiol. 1996;6(1):49-56.

13.   Ray HJ, Niswander L. Mechanisms of tissue fusion during development. Development. 2012;139(10):1701-11.

14.   Hayes P, Solon J. Drosophila dorsal closure: An orchestra of forces to zip shut the embryo. Mech Dev. 2017;144(Pt A):2-10.

15.   Jacinto A, Martinez-Arias A, Martin P. Mechanisms of epithelial fusion and repair. Nat Cell Biol. 2001;3(5):E117-23.

16.   Vasioukhin V, Bauer C, Yin M, Fuchs E. Directed actin polymerization is the driving force for epithelial cell-cell adhesion. Cell. 2000;100(2):209-19.

17.   Gumbiner BM. Cell adhesion: the molecular basis of tissue architecture and morphogenesis. Cell. 1996;84(3):345-57.

18.   Hero I. The optic fissure in the normal and microphthalmic mouse. Exp Eye Res. 1989;49(2):229-39.

19.   Hero I. Optic fissure closure in the normal cinnamon mouse. An ultrastructural study. Invest Ophthalmol Vis Sci. 1990;31(1):197-216.

20.   Geeraets R. An electron microscopic study of the closure of the optic fissure in the golden hamster. Am J Anat. 1976;145(4):411-31.

21.   Bernstein CS, Anderson MT, Gohel C, Slater K, Gross JM, Agarwala S. The cellular bases of choroid fissure formation and closure. Dev Biol. 2018;440(2):137-51.

22.   James A, Lee C, Williams AM, Angileri K, Lathrop KL, Gross JM. The hyaloid vasculature
facilitates basement membrane breakdown during choroid fissure closure in the zebrafish eye.
Dev Biol. 2016;419(2):262-72.
23.   Carrara N, Weaver M, Piedade WP, Vocking O, Famulski JK. Temporal characterization of
optic fissure basement membrane composition suggests nidogen may be an initial target of
remodeling. Dev Biol. 2019.
24.   Hardy H, Prendergast JG, Patel A, Dutta S, Trejo-Reveles V, Kroeger H, et al. Detailed
analysis of chick optic fissure closure reveals Netrin-1 as an essential mediator of epithelial
fusion. Elife. 2019;8.
25.   Monkley SJ, Kostourou V, Spence L, Petrich B, Coleman S, Ginsberg MH, et al.
Endothelial cell talin1 is essential for embryonic angiogenesis. Dev Biol. 2011;349(2):494-502.
26.   Dhakal S, Stevens CB, Sebbagh M, Weiss O, Frey RA, Adamson S, et al. Abnormal retinal
development in Cloche mutant zebrafish. Dev Dyn. 2015;244(11):1439-55.
27.   Heissig B, Hattori K, Friedrich M, Rafii S, Werb Z. Angiogenesis: vascular remodeling of
the extracellular matrix involves metalloproteinases. Curr Opin Hematol. 2003;10(2):136-41.
28.   Gestri G, Bazin-Lopez N, Scholes C, Wilson SW. Cell Behaviors during Closure of the
Choroid Fissure in the Developing Eye. Front Cell Neurosci. 2018;12:42.
29.   Macdonald R, Scholes J, Strahle U, Brennan C, Holder N, Brand M, et al. The Pax protein
Noi is required for commissural axon pathway formation in the rostral forebrain. Development.
1997;124(12):2397-408.
30.   Lee NV, Sato M, Annis DS, Loo JA, Wu L, Mosher DF, et al. ADAMTS1 mediates the
release of antiangiogenic polypeptides from TSP1 and 2. EMBO J. 2006;25(22):5270-83.
31.   Luque A, Carpizo DR, Iruela-Arispe ML. ADAMTS1/METH1 inhibits endothelial cell
proliferation by direct binding and sequestration of VEGF165. J Biol Chem. 2003;278(26):23656-
65.
32.   Sanyanusin P, McNoe LA, Sullivan MJ, Weaver RG, Eccles MR. Mutation of PAX2 in two
siblings with renal-coloboma syndrome. Hum Mol Genet. 1995;4(11):2183-4.
33.   Sanyanusin P, Schimmenti LA, McNoe LA, Ward TA, Pierpont ME, Sullivan MJ, et al.
Mutation of the PAX2 gene in a family with optic nerve colobomas, renal anomalies and
vesicoureteral reflux. Nat Genet. 1995;9(4):358-64.
34.   Torres M, Gomez-Pardo E, Gruss P. Pax2 contributes to inner ear patterning and optic
nerve trajectory. Development. 1996;122(11):3381-91.
35.   Brand M, Heisenberg CP, Jiang YJ, Beuchle D, Lun K, Furutani-Seiki M, et al. Mutations in
zebrafish genes affecting the formation of the boundary between midbrain and hindbrain.
Development. 1996;123:179-90.
36.   Favor J, Sandulache R, Neuhauser-Klaus A, Pretsch W, Chatterjee B, Senft E, et al. The
mouse Pax2(1Neu) mutation is identical to a human PAX2 mutation in a family with renal-
coloboma syndrome and results in developmental defects of the brain, ear, eye, and kidney.
Proc Natl Acad Sci U S A. 1996;93(24):13870-5.
37.   Otteson DC, Shelden E, Jones JM, Kameoka J, Hitchcock PF. Pax2 expression and retinal
morphogenesis in the normal and Krd mouse. Dev Biol. 1998;193(2):209-24.
38.   Hao J, Ho JN, Lewis JA, Karim KA, Daniels RN, Gentry PR, et al. In vivo structure-activity
relationship study of dorsomorphin analogues identifies selective VEGF and BMP inhibitors. ACS
Chem Biol. 2010;5(2):245-53.

39. Tan Ide A, Ricciardelli C, Russell DL. The metalloproteinase ADAMTS1: a comprehensive
review of its role in tumorigenic and metastatic pathways. Int J Cancer. 2013;133(10):2263-76.
40. Jeffrey EJ, Crawford BD. The epitope-mediated MMP activation assay: detection and
quantification of the activation of Mmp2 in vivo in the zebrafish embryo. Histochem Cell Biol.
2018;149(3):277-86.
41. Tsuji N, Matsuura T, Narama I, Yoshiki A, Ozaki K. Macrophage-Associated Gelatinase
Degrades Basement Membrane at the Optic Fissure Margins During Normal Ocular
Development in Mice. Invest Ophthalmol Vis Sci. 2018;59(3):1368-73.
42. Nishida Y, Miyamori H, Thompson EW, Takino T, Endo Y, Sato H. Activation of matrix
metalloproteinase-2 (MMP-2) by membrane type 1 matrix metalloproteinase through an
artificial receptor for proMMP-2 generates active MMP-2. Cancer Res. 2008;68(21):9096-104.
43. Strongin AY, Collier I, Bannikov G, Marmer BL, Grant GA, Goldberg GI. Mechanism of cell
surface activation of 72-kDa type IV collagenase. Isolation of the activated form of the
membrane metalloprotease. J Biol Chem. 1995;270(10):5331-8.
44. Theodore LN, Hagedorn EJ, Cortes M, Natsuhara K, Liu SY, Perlin JR, et al. Distinct Roles
for Matrix Metalloproteinases 2 and 9 in Embryonic Hematopoietic Stem Cell Emergence,
Migration, and Niche Colonization. Stem Cell Reports. 2017;8(5):1226-41.
45. Eckert P, Knickmeyer MD, Schutz L, Wittbrodt J, Heermann S. Morphogenesis and axis
specification occur in parallel during optic cup and optic fissure formation, differentially
modulated by BMP and Wnt. Open Biol. 2019;9(2):180179.
46. Heermann S, Schutz L, Lemke S, Krieglstein K, Wittbrodt J. Eye morphogenesis driven by
epithelial flow into the optic cup facilitated by modulation of bone morphogenetic protein.
Elife. 2015;4.
47. Knickmeyer MD, Mateo JL, Eckert P, Roussa E, Rahhal B, Zuniga A, et al. TGFbeta-
facilitated optic fissure fusion and the role of bone morphogenetic protein antagonism. Open
Biol. 2018;8(3).
48. Chen S, Lewis B, Moran A, Xie T. Cadherin-mediated cell adhesion is critical for the
closing of the mouse optic fissure. PLoS One. 2012;7(12):e51705.
49. Lahrouchi N, George A, Ratbi I, Schneider R, Elalaoui SC, Moosa S, et al. Homozygous
frameshift mutations in FAT1 cause a syndrome characterized by colobomatous-
microphthalmia, ptosis, nephropathy and syndactyly. Nat Commun. 2019;10(1):1180.
50. Richardson R, Owen N, Toms M, Young RM, Tracey-White D, Moosajee M.
Transcriptome profiling of zebrafish optic fissure fusion. Sci Rep. 2019;9(1):1541.
51. Smith R, Huang YT, Tian T, Vojtasova D, Mesalles-Naranjo O, Pollard SM, et al. The
Transcription Factor Foxg1 Promotes Optic Fissure Closure in the Mouse by Suppressing Wnt8b
in the Nasal Optic Stalk. J Neurosci. 2017;37(33):7975-93.
52. Cao M, Ouyang J, Guo J, Lin S, Chen S. Metalloproteinase Adamts16 Is Required for
Proper Closure of the Optic Fissure. Invest Ophthalmol Vis Sci. 2018;59(3):1167-77.
53. Anthis NJ, Wegener KL, Ye F, Kim C, Goult BT, Lowe ED, et al. The structure of an
integrin/talin complex reveals the basis of inside-out signal transduction. EMBO J.
2009;28(22):3623-32.
54. Wegener KL, Partridge AW, Han J, Pickford AR, Liddington RC, Ginsberg MH, et al.
Structural basis of integrin activation by talin. Cell. 2007;128(1):171-82.

55.   Calderwood DA, Campbell ID, Critchley DR. Talins and kindlins: partners in integrin-
mediated adhesion. Nat Rev Mol Cell Biol. 2013;14(8):503-17.
56.   Prout M, Damania Z, Soong J, Fristrom D, Fristrom JW. Autosomal mutations affecting
adhesion between wing surfaces in Drosophila melanogaster. Genetics. 1997;146(1):275-85.
57.   Boerboom D, Lafond JF, Zheng X, Lapointe E, Mittaz L, Boyer A, et al. Partially redundant
functions of Adamts1 and Adamts4 in the perinatal development of the renal medulla. Dev Dyn.
2011;240(7):1806-14.
58.   Brown JD, Dutta S, Bharti K, Bonner RF, Munson PJ, Dawid IB, et al. Expression profiling
during ocular development identifies 2 Nlz genes with a critical role in optic fissure closure. Proc
Natl Acad Sci U S A. 2009;106(5):1462-7.
59.   Cao M, Ouyang J, Liang H, Guo J, Lin S, Yang S, et al. Regional Gene Expression Profile
Comparison Reveals the Unique Transcriptome of the Optic Fissure. Invest Ophthalmol Vis Sci.
2018;59(15):5773-84.
60.   Wang Y, Kaiser MS, Larson JD, Nasevicius A, Clark KJ, Wadman SA, et al. Moesin1 and
Ve-cadherin are required in endothelial cells during in vivo tubulogenesis. Development.
2010;137(18):3119-28.
61.   Hocking JC, Famulski JK, Yoon KH, Widen SA, Bernstein CS, Koch S, et al. Morphogenetic
defects underlie Superior Coloboma, a newly identified closure disorder of the dorsal eye. PLoS
Genet. 2018;14(3):e1007246.
62.   Holly VL, Widen SA, Famulski JK, Waskiewicz AJ. Sfrp1a and Sfrp5 function as positive
regulators of Wnt and BMP signaling during early retinal development. Dev Biol.
2014;388(2):192-204.
63.   Pereira Piedade W, Veith S, Famulski JK. Ubiquitin-mediated proteasome degradation
regulates optic fissure fusion. Biol Open. 2019;8(6).
64.   Lauter G, Soll I, Hauptmann G. Two-color fluorescent in situ hybridization in the
embryonic zebrafish brain using differential detection systems. BMC Dev Biol. 2011;11:43.
1004

Figure 1

Figure 2

A

B

Figure 3

A

Wild-Type   pax2a -/-

B

- biological adhesion
- biological regulation
- cellular component of organization or biogenesis
- cellular process
- developmental process
- immune system process
- localization
- locomotion
- metabolic process
- multicellular organismal process
- reproduction
- reponse to stimulus

C

Gene ID	Gene Name	GO Term
aifm1	Apoptosis-inducing factor, mitochondrion-associated 1	oxidoreductase activity
sumo3a	small ubiquitin-like modifier 3a	protein sumoylation
lim2.1	Lens intrinsic membrane protein 2.1	Membrane
tfb2m	transcription factor B2, mitochondrial	transferase activity
tln1	Talin 1	actin binding
mgst2	microsomal glutathione S-transferase 2	enzyme activator activity
si:ch211-57b15.1	si:ch211-57b15.1	NA
ptprna	protein tyrosine phosphatase, receptor type, Na	protein dephosphorylation
dnajb11	DnaJ (Hsp40) homolog, subfamily B, member 11	protein folding
krt96	Keratin 96	structural molecule activity
sept9a	septin 9a	GTP binding
si:rp71-36a1.5	si:rp71-36a1.5	membrane
si:dkey-9c18.3	si:dkey-9c18.3	NA
arf5	ADP Ribosylation Factor 5	GTP binding
dock1	Dedicator Of Cytokinesis 1	small GTPase mediated signal transduction
hbbe1.1	Embryonic 1 beta-globin	Hemoglobin Complex
zgc:152951	Zgc:152951	electron transfer activity
pcbp3	Poly(rC)-binding protein 3	RNA binding
gpsm1b	G protein-signaling modulator 1b	GTPase regulator activity
abracl	Costars family protein ABRACL	regulation of actin filament-based process

D

Wild-Type   pax2a -/-

E

- biological adhesion
- biological regulation
- cellular component of organization or biogenesis
- cellular process
- developmental process
- immune system process
- localization
- locomotion
- metabolic process
- multicellular organismal process
- reproduction
- reponse to stimulus
- rhythmic process

F

Gene ID	Gene Name	GO Term
gmnn	geminin, DNA replication inhibitor	regulation of DNA replication
col11a1a	collagen, type XI, alpha 1a	extracellular matrix structural constituent
fubp3	far upstream element (FUSE) binding protein 3	RNA binding
si:dkey-26g8.4	Cathepsin Lb	proteolysis
ublcp1	ubiquitin-like domain containing CTD phosphatase 1	protein dephosphorylation
smarcal1	SWI/SNF related, matrix, actin dependent regulator of chromatin, a-like 1	DNA repair
he1b	hatching enzyme 1b	metalloendopeptidase activity
anxa3a	annexin A3a	regulation of NMDA receptor activity
si:dkey-269i1.4	cathepsin L1-like	proteolysis
zgc:174855	zgc:174855	
gsta.1	glutathione S-transferase, alpha tandem duplicate 1	glutathione metabolic process
tbc1d7	TBC1 domain family, member 7	activation of GTPase activity
srek1	splicing regulatory glutamine/lysine-rich protein 1	regulation of alternative mRNA splicing
scg5	secretogranin V	neuropeptide signaling pathway
ggact.3	gamma-glutamylamine cyclotransferase, tandem duplicate 3	lyase activity
si:dkey-26g8.4	Cathepsin L1-like	proteolysis
si:dkey-239j18.3	si:dkey-239j18.3	proteolysis
ahsa1a	AHA1, activator of heat shock protein ATPase homolog 1a	ATPase activator activity
he1a	hatching enzyme 1a	metalloendopeptidase activity
dpf3	D4, zinc and double PHD fingers, family 3	zinc ion binding

Figure 4

Figure 5

Figure 6

A

	Wild Type	pax2a -/-
28 hpf		
36 hpf		

adamts1

B

adamts1

Relative expression

WT   pax2a-/-

C

Wild Type · adamts1 mRNA · Wild Type · adamts1 mRNA · Wild Type · adamts1 mRNA · Wild Type · adamts1 mRNA

kdrl:mCherry

kdrl:mCherry DAPI

24 hpf   32 hpf   36 hpf   48 hpf

D

Vascularization of the Optic Fissure

Number of cells

<0.0001
****

ns

0.0002
***

ns

WT · adamts1 mRNA · pax2a-/-   WT · adamts1 mRNA · pax2a-/-

32hpf   36hpf

Figure 7

A

B

C

Figure 8

Supplementary Figure 1

A

distal

nasal ↔ temporal

proximal

distal
medial
proximal

B

1: Optic Fissure Signal

2: Laminin signal normalization

3: Phalloidin (F-actin) signal normalization

C

Wild Type Optic Fissure Laminin Intensity

Relative pixel Intensity

Distal   Medial   Proximal

24hpf
32hpf
36hpf
44hpf
48hpf

Wild Type Optic Fissure F-actin Intensity

Relative Pixel Intensity

Distal   Medial   Proximal

24hpf
32hpf
36hpf
44hpf
48hpf

D

pax2a-/- Optic Fissure Laminin Intensity

Relative pixel Intensity

Distal   Medial   Proximal

24hpf
32hpf
36hpf
44hpf
48hpf

pax2a-/- Optic Fissure F-actin Intensity

Relative pixel Intensity

Distal   Medial   Proximal

24hpf
32hpf
36hpf
44hpf
48hpf

Supplementary figure 2

Optic Fissure Apposition

Supplementary Figure 3

A

Supplementary Figure 4

Supplementary Figure 5

A

Supplementary Figure 6

A

B

10 uM ARP101   15 uM ARP101   20 uM ARP101   25 uM ARP101   30 uM ARP101