1
2
3

Pax2a regulates angiogenesis to facilitate fusion of the optic fissure

Megan L. Weaver[1], Piedade. W. P[1], Meshram, N.N[1] and J.K. Famulski[1].

1. Department of Biology, University of Kentucky

ABSTRACT:

Formation of the vertebrate retina involves timely and precise fusion of the optic fissure, failure of which results in the congenital blinding disorder coloboma. The optic fissure forms in consequence of optic cup morphogenesis and has a functional role as a conduit for migration of endothelial vasculature during hyaloid vasculogenesis. Recent studies have linked hyaloid vasculature precursor cells to be potential triggers, initiating optic fissure fusion. In order to examine this link, we analyzed the molecular events associated with optic fissure fusion and hyaloid vasculogenesis in the zebrafish pax2a mutant coloboma model. By means of wholemount immunohistochemistry, we determined that optic fissure fusion initiation, as indicated by laminin remodeling, is preceded by a spike in F-actin signaling. In contrast, the spike in F-actin activity and the disassembly of the basement membrane (BM) were misregulated in $pax2a^{-/-}$ embryos. To examine how the fusion process is disrupted in the absence of pax2a function, we compared transcriptomic profiles between $pax2a^{-/-}$ and wildtype (WT) embryos. This analysis uncovered a novel connection between regulation of angiogenesis and fusion. Loss of pax2a function resulted in significant reduction of talin1 expression, a known regulator of endothelial vasculature migration associated with fissure fusion. In addition, loss of pax2a also resulted in increased expression of an anti-angiogenic protease, ADAMTS1. 3D confocal and live imaging of retinal hyaloid vascularization in Tg[kdrl:mCherry] embryos indicated a significant deficit in $pax2a^{-/-}$ embryos while pharmacological inhibition of VEGF using DMH4 or ADAMTS1 mRNA overexpression phenocopied the pax2a mutant phenotype. Taken together, we propose that Pax2a positively regulates tln1 while negatively regulating ADAMTS1 expression. This enables timely hyaloid vascularization of the retina which in turn directly signals to initiate fissure fusion via cytoskeletal rearrangements and subsequent BM remodelling.

66 **AUTHOR SUMMARY**
67
Formation of a fully functioning human eye requires precise and coordinated cellular

mechanisms during very early development. One such mechanism is responsible for

the formation of the eyeball structure from a flat sheet of cells to a hemispherical organ.

In order to facilitate the biophysical properties of this shape change, an opening on the

ventral side of the eye forms and must be closed in order to properly form the eyeball.

Failure of this event leads to pediatric blindness in the form of a disease termed

Coloboma. While it has been studied by researchers for almost a century, we still lack a

significant understanding of how and why some children are born with this disease. In

this study, we aimed to better understand how the closure of this opening is triggered by

the developing tissues. What we found is, that non-eye cells which are found within this

opening and are making vasculature networks to nourish the eye, are likely triggers for

the fusion event. We found that regulating their migration and proliferation, also known

as angiogenesis, is required for closure of this opening and therefore proper

development of the eye.

INTRODUCTION:

Ocular development is a highly conserved process amongst vertebrate species. Assembly of the hemispherical retinal structure from an initially flat sheet of cells requires many complex morphogenetic movements. One such morphogenetic movement involves the invagination of the optic vesicle which results in a fissure forming at the ventral region of the developing retina. This fissure, known as the choroid or optic fissure (OF), enables hyaloid vasculature endothelial cell migration into the developing retina and subsequent establishment of the hyaloid vasculature. Hyaloid vasculature is a temporary circulatory system required for ocular development, and in most cases will degenerate once mature blood vessels begin to grow (1-4). As soon as the hyaloid vasculature has been established, the two opposing retinal epithelial sheets of the optic fissure will undergo fusion. Thereby, they encase the ganglion cell axons localized in the optic stalk and complete retinal morphogenesis. Failure of OF fusion leads to a congenital blinding disorder known as coloboma (5-7). Coloboma is a prevalent cause of pediatric blindness, accounting for approximately 10% of cases worldwide (6, 8). This makes it one of the leading causes of pediatric blindness. Coloboma is a spectrum disorder presenting unilaterally or bilaterally and ranging in severity from minor visual impairment, to complete blindness in the affected eye (9). This spectrum of severity is associated with the location and degree to which the choroid fissure was able to fuse and the severity of subsequent loss of ganglion cell axons (7). The process of epithelial tissue fusion is not unique to the eye and occurs throughout development, encompassing neural tube closure, palatal shelf formation and eyelid development to name a few. Similar to the failure of optic fissure fusion, congenital disorders have been associated with various organs where epithelial fusion fails (10). Epithelial fusion has been studied for over a century, and is known to involve transcriptional regulation, cell signaling pathways, and morphogen gradients (11, 12). However, in regard to the optic fissure we are still lacking a basic understanding of how these molecular mechanisms coordinate and directly facilitate fusion. Recent work has begun to implicate some of the cellular processes involved in optic fissure fusion. For example, the actin cytoskeleton is known to be a crucial component of the machinery driving fusion of the optic fissure as well as other tissues (13). The importance of the

actin cytoskeleton during epithelial fusion is thought to involve lamellipodia and filopodia

projections between the two opposing epithelia. These help to "zipper" the cells together

to form a single continuous sheet. When lamellipodial and filopodial projections are

precluded, fusion often fails (14). Interestingly, these projections were observed during

OF fusion almost 3 decades ago (15-17). However, while it has been observed that

actin protrusions likely play a functional role during OF fusion, it still remains to be

determined what signals rearrangement of the actin cytoskeleton. The second cellular

mechanism known to be directly involved in epithelial fusion is the remodeling of the

basement membrane (BM). During epithelial fusion, the BM acts as a physical barrier

restricting the establishment of cell to cell contacts, which must be removed in order for

fusion to complete. Recent work, including our own, across several species has

characterized progressive removal of the BM during choroid fissure fusion (18, 19).

However, the molecular mechanisms facilitating this process, in particular BM

remodeling, remain largely unknown. It was recently shown that migrating hyaloid

vasculature precursors found in the OF could potentially signal BM remodeling in

anticipation of fusion (19). James *et al.* 2016, showed that mutations in zebrafish talin1,

an actin cytoskeleton scaffolding protein known to be required for endothelial cell

migration (20), result in choroid fissure fusion defects. Theirs, and previous studies also

indicated that *cloche* mutants, which lack all early hyaloid vasculature, have delayed

basement membrane breakdown in the region of the choroid fissure (19, 21). Hyaloid

vasculature has long been a hypothesized to be an initiating factor of optic fissure

fusion. It's role during retinal development is to establish a vasculature supply to the

developing retina and lens while connecting to the choroid vasculature for proper blood

flow (3). Hyaloid vasculogenesis takes advantage of an open optic fissure so that

vasculature cells can migrate into the developing optic cup. Once optic fissure fusion is

completed, hyaloid vasculature is fully established. It has therefore been proposed that

because hyaloid vasculature requires an open fissure to complete establishment of its

network, migrating hyaloid vasculature cells could directly signal to the opposing retinal

lobes and regulate the timing of fissure fusion. This mechanism could potentially involve

vasculature-mediated activation of the fusion machinery within the retinal rim cells, or

direct supply of molecular factors, such as matrix proteases (22). In additional support

of this hypothesis, recent optic cup transplantation experiments in zebrafish embryos

confirm that in the absence of hyaloid vasculature, ectopic retinal OFs fail to initiate

fusion (23). There is therefore a clear link between OF fusion initiation and hyaloid

vasculature migration through the fissure.

Coloboma has been studied for many decades in many different species. This

has led to a general outline of the signaling and morphogenetic pathways required for

proper OF formation and fusion (recently reviewed in (24)). In particular, opposing

action of bone morphogenetic protein (BMP) and sonic hedgehog (Shh) signaling

establishes the dorso-ventral pattern of the optic vesicle and ensures proper expression

of optic stalk and OF regulators pax2, vax1 and vax2 (25, 26). However, as outlined

above, the actual molecular mechanisms driving the fusion process remain largely

unknown. Recent work from our lab has begun to decipher the composition and timing

of remodeling of the OF BM (27), however, the field still lacks a detailed understanding

of the cellular processes during and leading up to OF fusion. Hence, we have

undertaken a detailed analysis of zebrafish OF fusion comparing WT and the pax2a$^{-/-}$

coloboma model (28). In particular we have characterized the timing of BM remodeling,

F-actin dynamics, morphological apposition and hyaloid vascularization. Furthermore,

using transcriptomic analysis to compare WT and pax2a$^{-/-}$ eyes we discovered

misregulation in angiogenic signaling, resulting in decreased vasculogenesis of the

early retina and OF fusion failure. In particular we found that pax2a$^{-/-}$ embryos exhibit a

decrease in talin1 expression coupled with an increase in *A disintegrin and*

*metalloproteinase with thrombospondin motifs 1 (*ADAMTS1) expression, an anti-

angiogenic protease (29, 30). Modulation of vascularization via pharmacological

inhibition of VEGF signaling or overexpression of ADAMTS1 both phenocopied the

pax2a$^{-/-}$ hyaloid vasculature and coloboma phenotypes. This has led us to propose a

novel pathway for the regulation of optic fissure fusion where pax2a mediates proper

regulation of anti-angiogenic factors in order to facilitate hyaloid vasculature migration

through the OF. Timely migration of hyaloid vasculature cells in turn signals to the retina

rim cells, via F-actin, and initiates the fissure fusion machinery, in particular remodeling

of the BM.

RESULTS:

Optic Fissure basement membrane remodeling is preceded by F-actin signaling.

Several recent studies have undertaken a detailed time course to map out the exact timing of optic fissure fusion in numerous species, including zebrafish (18, 19). Overall, in zebrafish, the data point to ~32-36hpf as the time of OF fusion initiation, observed by BM remodeling. To better understand the molecular mechanisms regulating OF fusion, we also performed a detailed time course analysis of OF BM remodeling, using laminin immunohistochemistry (IHC) as a read out of this process. We performed whole mount IHC for laminin deposition in embryos starting at 24hpf and sampling every 4 hours up to 48hpf and then at 56 and 72hpf (Fig 1A). To analyze the progression of the fusion process, we quantified the laminin signal within the OF from 3D z-stacks of confocal scans. Regions of the BM outside the OF and juxtaposed to the developing lens were used to normalize the laminin signal (Fig S1A). In particular we quantified laminin signal in the central proximal region of the fissure (Fig S1B). This region has been shown to represent the site of fusion initiation. In agreement with recent studies in zebrafish, we found that OF fusion initiates at ~32-36hpf and does so in a central region of the fissure, subsequently proceeding proximally and distally (Fig 1B). By 48hpf, we observed most of the laminin signal is removed from the OF, and by 72hpf, little to no laminin persists in the region (Fig 1A, B). In addition to examining the timing of OF BM remodeling, we also determined whether changes in F-actin dynamics correlate to the timing of fissure fusion. James *et al*, 2016 had recently suggested that infiltrating vasculature endothelial cells migrating through the fissure could be a source of signal for fissure fusion. In fact, they showed that vasculature cell interaction with the fissure results in an increase of cytoskeletal, F-actin, rearrangements (19). To track F-actin signaling during fusion we stained the laminin labeled embryos using phalloidin and quantified the signal within the fissure. To normalize staining intensity we used an interior region of the retina (Fig S1A). Our analysis revealed a spike in OF associated F-actin signal between 24-32hpf, preceding the time that BM remodeling is initiated (Fig 1A,B). We also noticed a second spike of F-actin activity around 44hpf, a time of active

fusion (Fig 1C). Interestingly, the observed spikes in F-actin activity correlate with the

timing of hyaloid vasculature migration (24-32hpf) through the fissure and the expected

timing of actin protrusions facilitating epithelial fusion (36-48hpf). When measuring

distance between the fusing lobes of the retina we observed apposition to complete

around 36-40hpf (Fig S2). Taken together, we concluded that both apposition and BM

remodeling are preceded by a spike in F-actin within the OF driving initiation of BM

remodeling and subsequently fusion.

Optic fissure fusion mechanics are altered and disrupted upon loss of Pax2a

function.

In order to test our theory that an F-actin signaling wave precedes and drives

subsequent OF fusion we compared our findings to an established model of coloboma,

the pax2a[noi] line (28). Pax2 is a critical regulator of OF fusion in several model systems

and has been documented in human coloboma cases (31-33). The noi mutation is

predicted to result in a loss of Pax2a function due to a premature stop codon at position

198 (34). Originally characterized for their no-isthmus phenotype, pax2a[noi/noi] embryos

(referred to as pax2a[-/-] from this point on) elicit a fully penetrant unfused OF in addition

to severe heart defects. Unfortunately, homozygotes are not viable and do not enable

the study of coloboma at juvenile or adult stages. We hypothesized that examining the

molecular events leading to OF fusion in the pax2a[noi] system would inform us whether

these events are functionally important. Similar to our WT study, we examined lobe

distance, in addition to laminin and F-actin levels throughout the time OF fusion (Fig 2A,

Fig S1C). In stark contrast to WT, pax2a[-/-] embryos did not exhibit a significant decrease

in laminin signal between 32-48hpf, in fact, laminin appears to be largely retained in the

OF of pax2a[-/-] embryos (Fig 2B). While there is moderate decrease in laminin signal in

pax2a[-/-] embryos from 32-36hpf, this is well above the levels observed in WT. IHC

images clearly show that laminin persists in the fissure up to and including 72hpf (Fig

2A). A time when WT OF fusion is already completed and clearly absent of laminin

signal (Fig 1A). When examining F-actin levels in pax2a[-/-] embryos we did not detect a

spike in activity between 24 and 32hpf. Also, F-actin levels fluctuate between 36 and

44hpf and finally drop to WT levels by 48hpf. As such, it appears the absence of F-actin

signaling at early timepoints, 24-32hpf, correlates with failure to initiate fusion. Lastly,

when examining retinal lobe distance in pax2a$^{-/-}$ embryos we do not observe any

significant defects in apposition (Fig S2). Taken together, we conclude that in the

absence of pax2a the molecular mechanism regulating OF fusion, likely involving F-

actin signaling, fails to initiate. This is apparent by the persistence of OF BMs. While

neither F-actin signaling or BM remodeling is not initiated in pax2a$^{-/-}$ embryos,

apposition of the lobes appears largely unaffected indicating that pax2a does not

regulate morphogenetic movements of the retinal lobes. Based on these data we

propose that the F-actin signaling spike we observe just prior to initiation of OF BM

remodeling, in conjunction with pax2a activity, is critical for the completion of OF fusion.

Pax2a retinal transcriptomic profile during optic fissure fusion.

Interestingly, although pax2a has been the subject of intensive research, there

are currently no obvious transcriptional targets of pax2a that would directly point to

mechanistic regulation of OF fusion. In an effort to understand the absence of OF BM

remodeling and misregulation of F-actin activity observed in pax2a$^{-/-}$ embryos we

compared retinal transcriptomic profiles between WT and pax2a$^{-/-}$ embryos. Whole

retinas from 48hpf WT and pax2a$^{-/-}$ embryos were isolated and total RNA was

subsequently purified and sequenced using Illumina sequencing. 48hpf was chosen for

our experimental time point as pax2a$^{-/-}$ embryos are easily phenotyped at this age due

to severe heart malformation. When comparing three replicates for WT and pax2a$^{-/-}$

embryos using RSEM software, we detected 1215 transcripts significantly upregulated

(>95% confidence interval) and 1202 transcripts significantly downregulated (Fig 3A and

3D). Gene ontology analysis indicated a wide spread of biological function being

affected, including cytoskeletal signaling, adhesion and developmental processes for

both down and upregulated genes (Fig 3B, E). In fact, the distribution of biological

function between up and downregulated genes was highly similar (Fig 3B, D). A list of

top 20 up and down regulated genes is outlined in Figure 3C and F. Results of all the

statistically significant up and downregulated genes identified in our assay are

presented as supplementary data (Table 1).

From our analysis of downregulated targets, one, talin1 (tln1), stood out in

particular. As outlined in the introduction, tln1 was recently shown to be involved in the

migration of vasculature endothelial cells and subsequently fusion of the OF (19). As

such, we next sought to investigate the relationship between pax2a, tln1,

vasculogenesis and optic fissure fusion mechanics.

Loss of pax2a function leads to a reduction of hyaloid vasculature migration into

the optic fissure.

Tln1 has recently been shown to play a role in optic fissure fusion and our

discovery of reduced tln1 expression in pax2a$^{-/-}$ retinas engaged us to evaluate this

connection further. To validate our RNAseq data, we performed wholemount *in situ*

hybridization (WISH) for tln1 comparing WT siblings to pax2a$^{-/-}$ mutant embryos. WT

expression of tln1 was observed in the optic fissure between 28 and 48hpf coinciding

with pax2a expression (Fig 4A, Fig S3). In pax2a$^{-/-}$ embryos, OF tln1 expression

appears significantly reduced compared to WT while retaining similar expression in

periocular regions (Fig 4A). Tln1 expression is also reduced in the mid brain-hind brain

boundary, another region of strong pax2a expression (Fig 4A, S3A). To determine

whether the reduction of tln1 expression in pax2a$^{-/-}$ embryos had functional

consequences on optic fissure fusion and hyaloid vasculogenesis we examined pax2a$^{-/-}$

embryos expressing mCherry in vasculature cell, Tg[kdrl:mCherry]. Using 3D *in vivo*

time-lapse confocal microscopy we recorded migration of mCherry expressing cells

through the OF from 24 to 30hpf (Fig 4B, Movie 1). Both WT and pax2a$^{-/-}$ embryos

contain vasculature endothelial cells within the fissure at 24hpf, however, over the next

6 hours of imaging it becomes clear that a higher number of mCherry expressing cells

migrate through the OF in WT than in pax2a$^{-/-}$ embryos (Fig 4B, Movie 1). To quantify

this effect, we processed WT and pax2a$^{-/-}$ Tg[kdrl:mCherry] embryos at 24, 32, 36 and

48hpf, collected 3D confocal stacks and counted the number of mCherry positive cells

found within the fissure (Fig 4C). The data clearly indicate that there is a significant

reduction in the number of kdrl:mCherry cells found within the pax2a$^{-/-}$ optic fissure at

both 32 and 36hpf (Fig 4D). Furthermore, using 3D rendering rotation, we observed that

in 48hpf pax2a$^{-/-}$ embryos the hyaloid vasculature established in the back of the lens is

reduced in size and lacks the proper connections to the newly forming choroidal and

superficial vasculature systems (Movie 2-3). Overall, our data indicate that loss of pax2a

function leads to a decreased and impaired migration of hyaloid vasculature endothelial

cells into the optic fissure. Ultimately leading to a disorganized and reduced hyaloid

vasculature.

Inhibition of VEGF signaling impair optic fissure fusion.

Based on our discovery of reduced hyaloid vasculature in $pax2a^{-/-}$ embryos, we

next sought to examine whether this phenomenon is associated with failure of optic

fissure fusion. To do so, we turned our attention to vascular endothelial growth factor

(VEGF) signaling. VEGF, the ligand for vascular endothelial growth factor receptor

(VEGFR), is an angiogenic factor and the prime candidate for signaling the migration

and proliferation of hyaloid vasculature endothelial cells in the optic fissure. To inhibit

VEGF signaling, we took advantage of the dorsomorphin derivative DMH4, which has

been shown to selectively inhibit VEGF signaling independent from BMP (35). Based on

previously published working concentrations, we conducted a dose response to

examine DHM4 effects on hyaloid vasculature formation using Tg[kdrl:mCherry] as a

readout. Treatment of embryos from 12-24hpf, ranging from 1-100μM, resulted in a

dose response of reduction of mCherry signal in the developing retina (Fig S4). We

decided to use the 100μM concentration for subsequent experiments as this

concentration was able to completely inhibit vascularization of the retina up to 72hpf

(Fig 5A). Embryos were treated starting at 12hpf and examined for fissure fusion status

via whole mount laminin IHC at 24, 32, 36, 48, 56 and 72hpf (Fig 5B). 3D confocal

imaging revealed a persistence of laminin signal within the fissure up and including

72hpf (Fig 5B) upon DMH4 treatment. Measurement of the laminin signal over our time

course generated very similar results to what we observed in $pax2a^{-/-}$ embryos, with

laminin persisting within the OF at timepoints where WT embryos have completed

remodeling (Fig 5C). In addition to laminin, we also quantified F-actin activity during the

treatment (Fig 5B). Again, similar to $pax2a^{-/-}$ embryos we did not observe a spike in F-

actin activity between 24-32hpf as observed in WT embryos (Fig 5D). Similar to $pax2a^{-/-}$

embryos, DMH4 treatment also did not affect the size of the retina nor the apposition of

retinal lobes (Fig S2). Based on these findings, we conclude that inhibition of VEGF

signaling results in failure of basement membrane breakdown, presumably due to the

absence of vasculature endothelial cell migration through the optic fissure. This is

further supported by a lack of increased F-actin signaling during the time of vasculature

migration through the OF. As such, our results give direct evidence for the necessity of

vasculature endothelial cell migration through the OF in order to initiate BM remodeling

and ultimately achieve fusion of the fissure.

Pax2a negatively regulates ADAMTS1 expression to inhibit anti-angiogenesis.

Having established a clear connection between pax2a function and proper

recruitment of hyaloid vasculature endothelial cells into the optic fissure, we sought to

examine the mechanisms behind this phenotype. Our transcriptomic comparison of WT

and pax2a$^{-/-}$ eyes, with a focus on angiogenesis regulation, revealed misexpression of *a*

disintegrin and metalloproteinase with thrombospondin motifs 1 (ADAMTS1), a secreted

protease known to be involved in regulating VEGF signaling (36). Being a member of

the ADAMTS family, ADAMTS1 encodes three thrombospondin motifs enabling it to

directly bind and sequester VEGF (30). Furthermore, the protease activity of ADAMTS1

has bene shown to target *thrombospondin* (thsb) 1 and 2 for cleavage, ultimately

liberating their active forms of which bind to and block VEGF from binding the VEGF

receptor (29). In our transcriptomic comparison we found that ADAMTS1 expression

was upregulated almost two-fold in pax2a$^{-/-}$ retinas. This would suggest that the

expression of ADAMTS1, and therefore its anti-angiogenic activity, is normally

somehow kept in check by pax2a. Pax2a has not been reported to harbor

transcriptionally repressive function, so this may involve an indirect mechanism. To

confirm our RNAseq results we performed WISH, examining ADAMTS1 expression at

28 and 36 hpf. In agreement with our transcriptomic analysis, we observed an

upregulation in ADAMTS1 in eyes of pax2a$^{-/-}$ embryos, specifically in the region of the

optic fissure (Fig 6A). To test whether upregulation of ADAMTS1 expression would have

direct effects on hyaloid vasculature migration and optic fissure fusion we injected

Tg[kdrl:mCherry] expressing embryos with 100pg ADAMTS1 mRNA. We subsequently

examined hyaloid vasculature using 3D confocal imaging between 24-48hpf. Injection of

ADAMTS1 mRNA resulted in a significant decrease in the number of kdrl:mCherry cells

found within the optic fissure (Fig 6B, Movie 5). Quantification of mCherry positive cells

within the optic fissure confirmed this observation (Fig 6C). The reduction in the

numbers of mCherry positive cells within the fissure was similar to what we observed in

pax2a$^{-/-}$ embryos (Fig 4, Movies 2, 5). This result indicated that the anti-angiogenic

function of ADAMTS1 applies to regulating hyaloid vasculogenesis. In contrast to

ADAMTS1 mRNA overexpression, we also examined the consequences of ADAMTS1

loss of function using morpholino-mediated transient knockdown of ADAMTS1.

Tg[kdrl:mCherry] expressing embryos injected with ADAMTS1 morpholinos had a

contrasting phenotype to that of pax2a$^{-/-}$ or ADAMTS1 mRNA injected embryos.

Morphant ocular vasculature appeared expanded and significantly disorganized, in

particular within the dorsal regions (Fig 6D, Movie 6). Morphants did not display any

distinct phenotypes in hyaloid vasculature of the optic fissure at 48hpf, however, we did

note improper connections being formed as well as a disorganization of the dorsal

vasculature network as compared to WT (Fig 6D and Movies 4,6). These findings

suggest that while ADAMTS1 activity needs to be suppressed in the optic fissure, other

regions of the retinal vasculature may rely on proper levels of ADAMTS1 to enact

proper connections within the hyaloid/choroidal network.

<u>Misregulation of ADAMTS1 expression leads to optic fissure fusion failure</u>

Having mimicked the vasculature phenotype of pax2a mutants, we next

examined whether overexpression of ADAMTS1 would also inhibit optic fissure fusion.

To do so, we performed whole mount laminin and F-actin IHC in ADAMTS1 mRNA

injected embryos at 24, 32, 36, 48 and 72hpf (Fig 7A). Confocal imaging of the fissure

indicated that like the vasculature phenotype, ADAMTS1 mRNA injection also

phenocopied pax2a$^{-/-}$ associated persistence of laminin and therefore failure of optic

fissure fusion (Fig 7B). Levels of laminin persisting at 48hpf was similar to pax2a$^{-/-}$

embryos. As was observed with pax2a$^{-/-}$, timing for apposition of the lobes was not

affected in ADAMTS1 overexpressing embryos. Furthermore, as documented for

pax2a$^{-/-}$ and DMH4 treated embryos, we again noted a lack of the F-actin signaling

spike between 24-32hpf, and no significant difference to WT levels by 48hpf (Fig 7C).

Maintaining proper levels of ADAMTS1 expression is therefore necessary to ensure

timely fusion of the fissure. Taken together, we envision that pax2a functions to restrict

expression of ADAMTS1 within the optic fissure to ensure proper recruitment and

migration of hyaloid vasculature endothelial cells. One established, the hyaloid

vasculature in turn signals to the retinal rim cells to initiate BM remodeling and leads to

subsequent OF fusion.

DISCUSSION

Pax2 is the best recognized regulator of optic fissure fusion with a direct

connection to pediatric cases of coloboma. Interestingly, while the association of pax2

and coloboma is undoubted, little molecular detail exists of how pax2 actually confers

this important function. At the same time, studies of optic fissure fusion dating back

several decades have been suggesting a direct connection between the fusion process

and migrating vasculature endothelial, or periocular mesenchyme, cells found within the

fissure prior to and during fusion. A notion strongly supported by recent findings where

reduction of hyaloid vasculature in the fissure, or removal of optic vesicles from sources

of vasculature inhibits or significantly delays fusion (19, 23). In our study, we outline a

novel connection between pax2a function and hyaloid vasculature-mediated initiation of

optic fissure fusion, further strengthening the hypothesis that vasculature directly signals

initiation of fissure fusion.

Several recent reports, including this one, have comprehensively characterized

optic fissure fusion timing. Work from zebrafish, mice and chick all point to an orderly

progression involving: **1)** retinal growth and cellular rearrangement leading to nasal and

temporal retinal lobe apposition, **2)** invasion of the fissure by endothelial and neural

crest cells forming the hyaloid vasculature system, **3)** cellular signaling, either between

retinal rim cells or between rim cells and the migrating vasculature cells, likely in the

form of cytoskeletal activation (f-actin), **4)** remodeling or removal of the basement

membrane to enable physical connection of the rim cells and subsequent formation of a

continuous retinal epithelial sheet via re-polarization and cell-cell adhesion. Step 1 has

been nicely characterized in a few recent publications outlining the flow of retinal cells

and morphological formation of the fissure (37-39). For step 2, several reports have

carefully characterized the formation of the hyaloid vasculature system, including

migration of hyaloid vasculature precursor cells into the fissure as soon as it forms (1-3).

Importantly, perturbation of this process, or removal of the developing eye from its

source, has been shown to lead to fissure fusion failure (19, 23). To date, steps 3 and 4

are the least well understood. Several molecular components associated with cell-cell

adhesion and epithelial sheet fusion have been associated with the fissure, including α-

catenin, n-cadherin, and netrin (19, 40-43). However, the timing or the molecular

mechanism organizing and regulating these components remain uncharacterized.

Furthermore, very little information exists pertaining to the identity of protease enzymes

responsible for BM remodeling. Recent work from our lab has characterized the

composition of core BM components within the fissure, but to date, only ADAM16 has

been functionally examined in context of fissure fusion, while MMP2 was detected

during fusion in mice and MMP23bb implicated from transcriptomic comparison of pre

and post fusion fissure expression (42, 44, 45). Identifying proteases active in OF fusion

continues to be an area of needed attention. In our current study, we attempted to

address the questions of step 3 by concurrently analyzing f-actin and BM remodeling. In

doing so we discovered that BM remodeling is preceded by an uptick in F-actin

signaling. The timing of this coincides with the active migration of hyaloid vasculature

precursors through the fissure. An event shown previously to involve in F-actin signaling

within in the fissure (19). Furthermore, we show both genetically and pharmacologically

that hyaloid vasculature precursors are necessary for OF fusion, and ultimately outline a

mechanism where suppression of anti-angiogenic activity within the fissure is critical for

OF fusion. These findings not only confirm a long-held hypothesis of hyaloid vasculature

involvement in OF fusion, but also uncover previously unexplored mechanisms of

angiogenesis playing a role in the OF fusion process.

Based on our IHC time course analysis of optic fissure fusion dynamics, in

particular F-actin and laminin status, we propose that BM remodeling is preceded by an

increase of F-actin activity. This likely stemming from the interaction of retinal and

vasculature cells within the fissure (Fig 1). Interestingly, when examining this process in

the pax2a[noi] mutant line we did not detect an increase of Factin signaling at times

preceding fusion. This coincided with a decrease in the number of vasculature cells

found within the fissure and ultimately lead to failure of optic fissure basement

membrane remodeling and lack of fusion. In an attempt to understand the molecular

consequence driving these differences we uncovered misregulation of tln1 expression in

the absence of pax2a function. Tln1 is a key regulator of endothelial cell migration

recently shown to be directly involved in OF fusion (19). In fact, loss of tln1 function

leads to a decrease in fissure associated vasculature and persistence of the OF BM.

Our finding that pax2a regulates expression of tln1 has several compelling implications.

Talin1 is known to be a direct link between the actin cytoskeleton and the BM via

integrin (46, 47). Furthermore, tln1 has been associated with the formation of adhesion

junctions and other cell-cell interaction processes (48, 49). One could therefore

hypothesize several different models for its role in optic fissure fusion. Our, and

previous analyses of tln1 expression do not directly indicate which cells within the

fissure are expressing tln1. It could be the retinal lobe rim cells, the migrating

vasculature cells or both. As such, we can envision that tln1 could function through

differing mechanisms. If expressed solely in the retinal cells, then tln1 may function to

act as a sensor for the apposition of retinal lobes, or for the presence, or absence, of

vasculature endothelial cells. James et al. 2016 showed that vasculature cells within the

fissure trigger F-actin accumulation and presumably cytoskeletal signaling, potentially

facilitated by tln1. A feature that is missing in pax2a mutant fissures. Furthermore, tln1

may also act as a direct mechanism required for the formation of adhesions between

the fusing lobes. However, as observed in both tln1 and pax2 mutants, the BM persists

at times fusion should be occurring, suggesting that tln1 first plays a functional role in

the initiation of BM remodeling, and it remains unclear whether its directly involved in

cell-cell adhesion. Future work will focus on answering these questions.

Our study clearly supports the notion that vasculature plays an integral part in OF

fusion. As such, regulation of angiogenesis may also be part of the mechanism. Using a

transcriptomic comparison of WT and pax2a mutants we did in fact discover a

connection between angiogenesis and pax2a. In the absence of pax2a function we

observe an upregulation of ADAMTS1. Encoding 3 thrombospondin (thsb) motifs able to

directly bind and sequester VEGF, in addition to targeting thsb1 and 2 for cleavage into

their VEGF inhibitory active forms, ADAMTS1 has been extensively studied anti-

angiogenic factor in cancer, aortic and renal biology (50). In fact, we showed that

upregulating ADAMTS1 expression partially inhibits hyaloid vasculature migration into

the optic fissure, an outcome similarly observed in pax2a[noi] embryos. Up regulation of

ADAMTS1 also prevented the F-actin spike and OF BM remodeling while having no

effect on eye size or retinal lobe apposition. Along those same lines we also showed

that optic fissure fusion and preceding F-actin signaling are abolished when

pharmacologically inhibiting VEGF using DHM4. Interestingly, knockdown of ADAMTS1

using morpholino resulted in hyaloid vasculature disorganization, suggesting anti-

angiogenesis may function to guide and/or shape the developing vasculature network

during retinal development. Taken together, our study confirms an undeniable link

between hyaloid vasculogenesis and optic fissure fusion. It also uncovers a molecular

mechanism involving angiogenic regulation to ensure proper and timely establishment

of hyaloid vasculature and subsequent OF fusion.

MATERIALS AND METHODS

Zebrafish and embryo maintenance

Zebrafish were maintained using husbandry procedures approved by University of Kentucky IACUC committee. Embryos were kept at 28.5°C in E3 embryo media. AB and TL strains were used as wild-type, Tg[*kdrl*:mCherry] transgenic line was used to visualize retinal vascularization (51).

Pax2^{noi} embryos were a gift from Dr. Gregory-Evans. Genotyping analysis was conducted by amplifying the region of gDNA with the noi mutation using the forward primer: 5'- CTCGCTCTGCCTCCATGATTG3-'and the reverse: 5'- GGCACTGAAAGAGCACAGG -3'. The resultant 460bp amplicon was digested with TaqI (NEB) which would recognize and digest the WT allele sequence but not the mutant allele.

Immunohistochemistry (IHC)

Dechorionated embryos were fixed with 4% PFA in PBS at room temperature for 3h and washed with PBST 4 times for 5 minutes. Embryos were then permeabilized with Proteinase K, 30μg/mL 10 minutes for 24-28 hpf, 50μg/mL 15-20 minutes for 32-48 hpf and 75μg/mL 20 minutes for 56-72 hpf, washed 2 times in PBST for 5 minutes and blocked overnight at 4°C with 10% sheep serum, 0.8% Triton X-100 and 1% BSA in PBS. Primary mouse anti-laminin antibody (ThermoFisher – 1:100) in blocking buffer (1% sheep serum, 1% BSA and 0.8% Triton X-100 in PBS) were incubated overnight at 4°C and washed 5 times in PBST for 15 minutes. Secondary antibody, goat anti-rabbit (Alexa Fluor® 488 – Abcam – 1:1000), DAPI 1:1000, and phalloidin (Alexa Fluor® 555 - 1:50) were incubated overnight at 4°C in the dark. Tg[*kdrl*:mCherry] embryos were treated with proteinase K as described above and stained with DAPI 1:1000 overnight. The embryos were washed 2 times in PBST for 15 minutes and visualized using a Nikon C2+ confocal microscope equipped with a 40X (1.15NA) water immersion objective. Embryos were embedded in 1.2% low melting point agarose on glass bottom 35mm dishes (Fluorodish, World Precision Instruments). Images were captured in steps of 3.5 microns for a total of 31.5 microns using Nikon Elements software. Image

adjustment, such as cropping and brightness/contrast was performed using Adobe

Photoshop.

<u>Analysis of Fluorescence signal</u>

Fiji software (https://fiji.sc) was used to measure the fluorescence intensity of laminin

and phalloidin signal from raw image data. In order to account for variability in staining,

normalization values were measured for laminin and actin pixel intensity where an area

directly outside of the choroid fissure was measured and a ratio was generated between

the two values (Fig S1B). For Tg[krdl:mCherry], 3D reconstructions of the optic fissure

were generated and individual cells were counted (from the opening of the OF through

the back of the lens).

<u>Stats</u>

Student's t-test was used to compare individual time points. One-way ANOVA was

used to analyze across treatments. Graphs are displayed as mean +/- standard

deviation. Analysis was performed using Prism8 graphing software (GraphPad).

<u>Total RNA Sequencing</u>

WT and Pax2$^{-/-}$ embryos were dissected and collected at 48 hpf and RNA was

extracted from dissected eyes using trizol. Pax2a$^{-/-}$ embryos were phenotyped by

distinct heart defects only observed in pax2a$^{-/-}$ embryos, as verified by previous

genotyping experiments. Dissected eyes were suspended in 1 mL of trizol and sheared

with a 22-gauge needle. Samples were incubated at room temperature for 5 minutes,

then 200 μL of chloroform was added and vortexed for 1 minute. Samples were then

centrifuged at 12,000 g for 15 minutes at 4℃. Aqueous phase was removed and put into

fresh RNAse free tube. 3 μL of Glycoblue was added to samples and vortexed for 5

seconds. 500 μL of 100% Isopropanol was then added and vortexed for an additional 10

seconds. Samples then were incubated at room temperature for 10 minutes. Samples

were then centrifuged at 12,000 g for 10 minutes at 4℃. Supernatant was removed and

1 mL of 75% EtOH was added and vortexed for 10 seconds. Samples were then

centrifuged at 7,500 g for 5 minutes at 4℃. Supernatant was removed and samples then

underwent a pulse centrifugation and any remaining supernatant was removed.

Samples were then air-dried for 7 minutes at room temperature under a fume hood. 20

μL of RNAse free ddH$_2$O was added to the samples and mixed until pellet dissolved.

Samples were finally incubated at 60℃ for 12 minutes and then stored at -80℃.

The RNA then underwent a DNase treatment using the DNA-free Kit. 2 μL of 10X

DNaseI Buffer, and 1 μL of rDNaseI was added to the entire 20 μL sample. Samples

were incubated at 37℃ for 20 minutes, and then 2 μL of DNase Inactivation Reagent

was added. Samples were then incubated at room temperature for 2 minutes, mixing

samples 3 times. Samples were centrifuged at 10,000 g for 1.5 minutes. Supernatant

was then removed and placed into fresh RNase free tube and stored at -80℃.

Purified total RNA was sent to Applied Biological Materials Inc. for Illumina

sequencing. Bioinformatic analyses were completed using RSEM. Those transcripts

which had a greater than 0.95 posterior probability of being differentially expressed

were considered significant. Panther (http://pantherdb.org) was used for gene ontology

and pie chart generation.

Whole-mount in situ hybridization (WISH)

Whole-mount in situ hybridization was performed as previously described (52). RNA

probes were generated using PCR with T7 promoter sequence and subsequently

transcribed [DIG labeled] using T7 polymerase (Roche). Primer sequences were 5'-

GAGGGCAAGAGAATTCAGTAC-3' and 5'-

TAATACGACTCACTATAGGGCAGATTGCACAAGTTCAGAAG-3' for ADAMTS1, 5'-

ATGGATATTCACTGCAAAGCAG-3' and 5'-

TAATACGACTCACTATAGGGCTAGTGGCGGTCATAGGCAGTG-3' for pax2a, 5'-

GCCACAGCTAAAGCAGTGGCTG-3' and 5'-

TAATACGACTCACTATAGGGGTACTGCTGCTGTCGGATCAT-3' for talin 1. Images

were captured using a Nikon Digital sight DS-Fi2 camera mounted on a Nikon SZM800

stereo scope using Elements software. Dissected eyes from 24, to 72hpf embryos were

mounted in 70% glycerol and imaged under DIC using a Nikon TiE compound

microscope equipped with a 20X (0.7NA) objective and Elements software. Image

adjustment, such as cropping and brightness/contrast was performed using Adobe

Photoshop.

Morpholino Injections

Translation blocking Morpholino was obtained from Gene Tools, LLC and used to

knock down the expression of ADAMTS1. The morpholino used in this study was:

Control MO: 5'-CCTCTTACCTCAGTTACAATTTATA-3', Adamts1: 5'-

AGACATTATATTCCTGTCAAGTTCT-3'. 6ng of of ADAMTS1 or control morpholino

was injected into 1-2 cell stage embryos.

Live imaging analysis

Live imaging of Tg[kdrl:mCherry] embryos was conducted using a Nikon C2+ confocal

microscope equipped with at 20X (0.95NA) water immersion objective. 22 hpf embryos

were imbedded in 1.1% low gelling agarose in 1-inch glass bottomed Flourodish cell

culture dishes (World Precision Instruments) and covered in embryo media, 3-amino

benzoic acidethylester (tricaine) to anaesthetize the embryos and 1-phenyl 2-thiourea

(PTU) to inhibit pigmentation. Z-stacks 75μm thick with a step size of 2.5μm were

captured over the course of 6 hours at 10 minute intervals. The time lapse data were

reconstructed in 3D using Elements software. Image adjustment, such as cropping and

brightness/contrast was performed using Adobe Photoshop. After imaging, embryos

were removed and genotyped.

DMH-4 treatment

Embryos were incubated in embryo media with 5, 25, 50, or 100 μM of DMH-4 (Sigma

Life Science) in DMSO starting at 12hpf. Fresh DHM4 containing media was added

every 12hpf for timepoints past 24hpf.

Cloning and mRNA synthesis.

Full coding domain sequences for ADAMTS1 (NCBI reference sequence

XM_021475923.1), Pax2a (Ensembl transcript ID: ENSDARG00000028148) were

amplified from 24hpf zebrafish cDNA using primers 5'-

CATGGATCCATGGATATTCACTGCAAAGCA-3' and 5'-

ATGCTCGAGCTAGTGGCGGTCATAGGCAGTG-3' for pax2a, 5'-

GCGATCGCATGTCTTTTTTGCGTGTGGTG-3' and 5'-

CTCGAGTCAACAGGGAGTCAGATTGC-3' for ADAMTS1, cloned into pGEMT

(Promega), digested with AsiSI/XhoI, BamHI/XhoI respectively and subsequently cloned

into pCS2+. All constructs were verified by sanger sequencing (Eurofinsgenomics).

mRNA was synthesized from linearized pCS2 constructs using SP6 mMessage

mMachine kit (Ambion) and purified using YM-50 Microcon columns (Amicon, Millipore).

<u>Ethics Statement</u>

The use of zebrafish in this study was approved by the University of Kentucky IACUC

committee, Institutional PHS Assurance #D16-00217 (A3336-01) with a protocol

number: 2015-1370.

638
639
ACKNOWLEDGEMENTS

This work for supported by faculty start-up funds provided to JKF from the department

of Biology, University of Kentucky. MLW was supported by the Lyman T Johnson

scholarship from the University of Kentucky. PPW was supported by Brazilian National

Council for Scientific and Technological Development (CNPq) under grant number

202970/2014-0. We thank Dr. Jeramiah Smith for assistance with bioinformatic analysis

of our RNAseq data and members of the Famulski lab for helpful discussions.

REFERENCES:
1.   Hartsock A, Lee C, Arnold V, Gross JM. In vivo analysis of hyaloid vasculature morphogenesis in zebrafish: A role for the lens in maturation and maintenance of the hyaloid. Dev Biol. 2014;394(2):327-39.
2.   Hashiura T, Kimura E, Fujisawa S, Oikawa S, Nonaka S, Kurosaka D, et al. Live imaging of primary ocular vasculature formation in zebrafish. PLoS One. 2017;12(4):e0176456.
3.   Kaufman R, Weiss O, Sebbagh M, Ravid R, Gibbs-Bar L, Yaniv K, et al. Development and origins of zebrafish ocular vasculature. BMC Dev Biol. 2015;15:18.
4.   Yoshikawa Y, Yamada T, Tai-Nagara I, Okabe K, Kitagawa Y, Ema M, et al. Developmental regression of hyaloid vasculature is triggered by neurons. J Exp Med. 2016;213(7):1175-83.
5.   AS AL, Gregory-Evans CY, Gregory-Evans K. An update on the genetics of ocular coloboma. Hum Genet. 2019.
6.   Chang L, Blain D, Bertuzzi S, Brooks BP. Uveal coloboma: clinical and basic science update. Curr Opin Ophthalmol. 2006;17(5):447-70.
7.   Gregory-Evans CY, Williams MJ, Halford S, Gregory-Evans K. Ocular coloboma: a reassessment in the age of molecular neuroscience. J Med Genet. 2004;41(12):881-91.
8.   Stoll C, Alembik Y, Dott B, Roth MP. Congenital eye malformations in 212,479 consecutive births. Ann Genet. 1997;40(2):122-8.
9.   Huynh N, Brooks BP. Uveal coloboma. JAMA Ophthalmol. 2013;131(10):1274.
10.   Ray HJ, Niswander L. Mechanisms of tissue fusion during development. Development. 2012;139(10):1701-11.
11.   Hayes P, Solon J. Drosophila dorsal closure: An orchestra of forces to zip shut the embryo. Mech Dev. 2017;144(Pt A):2-10.
12.   Jacinto A, Martinez-Arias A, Martin P. Mechanisms of epithelial fusion and repair. Nat Cell Biol. 2001;3(5):E117-23.
13.   Vasioukhin V, Bauer C, Yin M, Fuchs E. Directed actin polymerization is the driving force for epithelial cell-cell adhesion. Cell. 2000;100(2):209-19.
14.   Gumbiner BM. Cell adhesion: the molecular basis of tissue architecture and morphogenesis. Cell. 1996;84(3):345-57.
15.   Hero I. The optic fissure in the normal and microphthalmic mouse. Exp Eye Res. 1989;49(2):229-39.
16.   Hero I. Optic fissure closure in the normal cinnamon mouse. An ultrastructural study. Invest Ophthalmol Vis Sci. 1990;31(1):197-216.
17.   Geeraets R. An electron microscopic study of the closure of the optic fissure in the golden hamster. Am J Anat. 1976;145(4):411-31.
18.   Bernstein CS, Anderson MT, Gohel C, Slater K, Gross JM, Agarwala S. The cellular bases of choroid fissure formation and closure. Dev Biol. 2018;440(2):137-51.
19.   James A, Lee C, Williams AM, Angileri K, Lathrop KL, Gross JM. The hyaloid vasculature facilitates basement membrane breakdown during choroid fissure closure in the zebrafish eye. Dev Biol. 2016;419(2):262-72.
20.   Monkley SJ, Kostourou V, Spence L, Petrich B, Coleman S, Ginsberg MH, et al. Endothelial cell talin1 is essential for embryonic angiogenesis. Dev Biol. 2011;349(2):494-502.

21.   Dhakal S, Stevens CB, Sebbagh M, Weiss O, Frey RA, Adamson S, et al.
Abnormal retinal development in Cloche mutant zebrafish. Dev Dyn.
2015;244(11):1439-55.
22.   Heissig B, Hattori K, Friedrich M, Rafii S, Werb Z. Angiogenesis: vascular
remodeling of the extracellular matrix involves metalloproteinases. Curr Opin Hematol.
2003;10(2):136-41.
23.   Gestri G, Bazin-Lopez N, Scholes C, Wilson SW. Cell Behaviors during Closure
of the Choroid Fissure in the Developing Eye. Front Cell Neurosci. 2018;12:42.
24.   Patel A, Sowden JC. Genes and pathways in optic fissure closure. Semin Cell
Dev Biol. 2017.
25.   Take-uchi M, Clarke JD, Wilson SW. Hedgehog signalling maintains the optic
stalk-retinal interface through the regulation of Vax gene activity. Development.
2003;130(5):955-68.
26.   Macdonald R, Wilson SW. Pax proteins and eye development. Curr Opin
Neurobiol. 1996;6(1):49-56.
27.   Carrara N, Weaver M, Piedade WP, Vocking O, Famulski JK. Temporal
characterization of optic fissure basement membrane composition suggests nidogen
may be an initial target of remodeling. Dev Biol. 2019.
28.   Macdonald R, Scholes J, Strahle U, Brennan C, Holder N, Brand M, et al. The
Pax protein Noi is required for commissural axon pathway formation in the rostral
forebrain. Development. 1997;124(12):2397-408.
29.   Lee NV, Sato M, Annis DS, Loo JA, Wu L, Mosher DF, et al. ADAMTS1 mediates
the release of antiangiogenic polypeptides from TSP1 and 2. EMBO J.
2006;25(22):5270-83.
30.   Luque A, Carpizo DR, Iruela-Arispe ML. ADAMTS1/METH1 inhibits endothelial
cell proliferation by direct binding and sequestration of VEGF165. J Biol Chem.
2003;278(26):23656-65.
31.   Sanyanusin P, McNoe LA, Sullivan MJ, Weaver RG, Eccles MR. Mutation of
PAX2 in two siblings with renal-coloboma syndrome. Hum Mol Genet. 1995;4(11):2183-
4.
32.   Sanyanusin P, Schimmenti LA, McNoe LA, Ward TA, Pierpont ME, Sullivan MJ,
et al. Mutation of the PAX2 gene in a family with optic nerve colobomas, renal
anomalies and vesicoureteral reflux. Nat Genet. 1995;9(4):358-64.
33.   Torres M, Gomez-Pardo E, Gruss P. Pax2 contributes to inner ear patterning and
optic nerve trajectory. Development. 1996;122(11):3381-91.
34.   Brand M, Heisenberg CP, Jiang YJ, Beuchle D, Lun K, Furutani-Seiki M, et al.
Mutations in zebrafish genes affecting the formation of the boundary between midbrain
and hindbrain. Development. 1996;123:179-90.
35.   Hao J, Ho JN, Lewis JA, Karim KA, Daniels RN, Gentry PR, et al. In vivo
structure-activity relationship study of dorsomorphin analogues identifies selective
VEGF and BMP inhibitors. ACS Chem Biol. 2010;5(2):245-53.
36.   Tan Ide A, Ricciardelli C, Russell DL. The metalloproteinase ADAMTS1: a
comprehensive review of its role in tumorigenic and metastatic pathways. Int J Cancer.
2013;133(10):2263-76.

37.   Eckert P, Knickmeyer MD, Schutz L, Wittbrodt J, Heermann S. Morphogenesis
and axis specification occur in parallel during optic cup and optic fissure formation,
differentially modulated by BMP and Wnt. Open Biol. 2019;9(2):180179.
38.   Heermann S, Schutz L, Lemke S, Krieglstein K, Wittbrodt J. Eye morphogenesis
driven by epithelial flow into the optic cup facilitated by modulation of bone
morphogenetic protein. Elife. 2015;4.
39.   Knickmeyer MD, Mateo JL, Eckert P, Roussa E, Rahhal B, Zuniga A, et al.
TGFbeta-facilitated optic fissure fusion and the role of bone morphogenetic protein
antagonism. Open Biol. 2018;8(3).
40.   Chen S, Lewis B, Moran A, Xie T. Cadherin-mediated cell adhesion is critical for
the closing of the mouse optic fissure. PLoS One. 2012;7(12):e51705.
41.   Lahrouchi N, George A, Ratbi I, Schneider R, Elalaoui SC, Moosa S, et al.
Homozygous frameshift mutations in FAT1 cause a syndrome characterized by
colobomatous-microphthalmia, ptosis, nephropathy and syndactyly. Nat Commun.
2019;10(1):1180.
42.   Richardson R, Owen N, Toms M, Young RM, Tracey-White D, Moosajee M.
Transcriptome profiling of zebrafish optic fissure fusion. Sci Rep. 2019;9(1):1541.
43.   Smith R, Huang YT, Tian T, Vojtasova D, Mesalles-Naranjo O, Pollard SM, et al.
The Transcription Factor Foxg1 Promotes Optic Fissure Closure in the Mouse by
Suppressing Wnt8b in the Nasal Optic Stalk. J Neurosci. 2017;37(33):7975-93.
44.   Cao M, Ouyang J, Guo J, Lin S, Chen S. Metalloproteinase Adamts16 Is
Required for Proper Closure of the Optic Fissure. Invest Ophthalmol Vis Sci.
2018;59(3):1167-77.
45.   Tsuji N, Matsuura T, Narama I, Yoshiki A, Ozaki K. Macrophage-Associated
Gelatinase Degrades Basement Membrane at the Optic Fissure Margins During Normal
Ocular Development in Mice. Invest Ophthalmol Vis Sci. 2018;59(3):1368-73.
46.   Anthis NJ, Wegener KL, Ye F, Kim C, Goult BT, Lowe ED, et al. The structure of
an integrin/talin complex reveals the basis of inside-out signal transduction. EMBO J.
2009;28(22):3623-32.
47.   Wegener KL, Partridge AW, Han J, Pickford AR, Liddington RC, Ginsberg MH, et
al. Structural basis of integrin activation by talin. Cell. 2007;128(1):171-82.
48.   Calderwood DA, Campbell ID, Critchley DR. Talins and kindlins: partners in
integrin-mediated adhesion. Nat Rev Mol Cell Biol. 2013;14(8):503-17.
49.   Prout M, Damania Z, Soong J, Fristrom D, Fristrom JW. Autosomal mutations
affecting adhesion between wing surfaces in Drosophila melanogaster. Genetics.
1997;146(1):275-85.
50.   Boerboom D, Lafond JF, Zheng X, Lapointe E, Mittaz L, Boyer A, et al. Partially
redundant functions of Adamts1 and Adamts4 in the perinatal development of the renal
medulla. Dev Dyn. 2011;240(7):1806-14.
51.   Wang Y, Kaiser MS, Larson JD, Nasevicius A, Clark KJ, Wadman SA, et al.
Moesin1 and Ve-cadherin are required in endothelial cells during in vivo tubulogenesis.
Development. 2010;137(18):3119-28.
52.   Holly VL, Widen SA, Famulski JK, Waskiewicz AJ. Sfrp1a and Sfrp5 function as
positive regulators of Wnt and BMP signaling during early retinal development. Dev Biol.
2014;388(2):192-204.
785

FIGURE LEGENDS

Figure 1: **An increase in F-actin dynamics precedes laminin remodeling during optic fissure fusion.**

A) Whole mount Immunohistochemistry was used to simultaneously visualize F-actin (red) and laminin (green) during optic fissure fusion, 24-72hpf. Central-proximal sections obtained using confocal imaging were collected and quantified. Scale bar = 50um. **B)** Quantification of laminin signal intensity within the optic fissure, normalized to regions of laminin staining juxtaposed to the lens. Relative pixel intensities are displayed. A green dotted line depicts the trend in laminin intensity over time. ANOVA p<0.0001. **C)** Quantification of F-actin signal intensity (phalloidin staining) within the optic fissure, normalized to regions of F-actin signal within the lobe of the retina. Relative pixel intensities are displayed. A red dotted line represents the trend in F-actin intensity over time. ANOVA p<0.0001.

Figure 2: **Pax2 mutant embryos lack F-actin activity increase and laminin remodeling within the optic fissure.**

A) Whole mount Immunohistochemistry was used to simultaneously visualize F-actin (red) and laminin (green) during optic fissure fusion, 24-72hpf, in pax2$^{-/-}$ embryos. Central-proximal sections obtained using confocal imaging were collected and quantified. Scale bar = 50um **B)** Quantification of laminin signal intensity within the optic fissure, normalized to regions of laminin staining juxtaposed to the lens. Relative pixel intensities are displayed. A green dotted line depicts the trend in laminin intensity over time. Blue dotted line represents the trend observed in wildtype embryos. ANOVA p<0.0001. **C)** Quantification of F-actin signal intensity (phalloidin staining) within the optic fissure, normalized to regions of F-actin signal within the lobe of the retina. Relative pixel intensities are displayed. A red dotted line represents the trend in F-actin intensity over time. Blue dotted line represents the trend observed in wildtype embryos. ANOVA p<0.0001.

<u>Figure 3</u>: **Retinal transcriptomic comparison between WT and pax2a⁻/⁻.**

A) Heat map representing transcripts found to be significantly down regulated in pax2a⁻/⁻

embryos. **B)** Gene ontology for pax2a⁻/⁻ downregulated transcripts. **C)** Top 20

downregulated transcripts are outlined in order of fold change. **D)** Heat map

representing transcripts found to be significantly upregulated in pax2a⁻/⁻ embryos. **E)**

Gene ontology for pax2a⁻/⁻ upregulated transcripts. **F)** Top 20 upregulated transcripts

are outlined in order of fold change.

<u>Figure 4</u>: **Pax2a is necessary for tln1 expression and recruitment of vasculature**

into the optic fissure.

A) Wholemount in situ hybridization comparing tln1 expression between WT and

pax2a⁻/⁻ embryos at 28 and 32hpf. Tln1 signal within the optic fissure (yellow arrowhead)

is clearly reduced in Pax2a⁻/⁻ embryos (red *). **B)** *In vivo* 4D confocal imaging of

Tg[kdrl:mCherry] WT or pax2a⁻/⁻ embryos. Time lapse series depicting the region of the

optic fissure (dotted white lines) and mCherry positive vasculature endothelial cells

migrating through the fissure. Compared to WT, pax2a⁻/⁻ embryos display a reduction in

the number of vasculature cells migrating in the fissure from 24-29hpf. Scale bar =

10um. **C)** Comparison of WT and pax2a⁻/⁻ vascularization during optic fissure fusion, 24-

48hpf. 3D reconstructions of whole mount Tg[kdrl:mCherry] (red) WT or pax2a⁻/⁻

embryos with DNA stained by DAPI (blue). Scale bar = 50um. **D)** Quantification of

mCherry positive cells from 3D confocal stacks within the region of the optic fissure.

Individual embryo results are depicted. ANOVA p<0.0001.

<u>Figure 5</u>: **Inhibiting angiogenesis prevents optic fissure fusion.**

A) 3D confocal imaging of Tg[kdrl:mCherry] (red) embryos treated with DMH4 at 24-

72hpf. DNA was stained with DAPI (blue). Scale bar = 50um. DMH4 treatment restricts

the migration of vasculature endothelial cells into the fissure and retina. **B)** Whole mount

Immunohistochemistry was used to simultaneously visualize F-actin (red) and laminin

(green) in DMH4 treated embryos during the time of optic fissure fusion, 24-72hpf.

Central-proximal sections obtained using confocal imaging were collected and

quantified. Treatment with DMH4 inhibits laminin removal and results in optic fissure

fusion failure. Scale bar = 50um. **C)** Quantification of laminin signal intensity within the

optic fissure, normalized to regions of laminin staining juxtaposed to the lens. Relative

pixel intensities are displayed. A green dotted line depicts the trend in laminin intensity

over time, a blue dotted line represents the WT trend. ANOVA p<0.0001. **D)**

Quantification of F-actin signal intensity (phalloidin staining) within the optic fissure,

normalized to regions of F-actin signal within the lobe of the retina. Relative pixel

intensities are displayed. A red dotted line represents the trend in F-actin intensity over

time, a blue dotted line represents the WT trend. ANOVA p<0.0001.

Figure 6: **ADAMTS1 inhibits retinal vascularization.**

A) Wholemount in situ hybridization comparing ADAMTS1 expression between WT and

pax2a$^{-/-}$ embryos at 28 and 32hpf. ADAMTS1 signal within the optic fissure (yellow

arrowhead) is clearly increased in Pax2a$^{-/-}$ embryos (red *). **B)** Comparison of WT and

ADAMTS1 mRNA injected embryo vascularization during optic fissure fusion, 24-48hpf.

3D reconstructions of whole mount Tg[kdrl:mCherry] (red) WT or ADAMTS1 mRNA

injected embryos with DNA stained by DAPI (blue). Scale bar = 50um. **C)** Quantification

of mCherry positive cells from 3D confocal stacks within the region of the optic fissure.

Individual embryo results are depicted. * p<0.001, ANOVA p<0.0001. **D)** 3D

reconstruction of Tg[kdrl:mCherry] (red) WT or ADAMTS1 MO injected embryos at

48hpf. DNA is stained with DAPI (blue). Inhibiting ADAMTS1 function leads to

significant disorganization of the dorsal vasculature (yellow *) as well as misconnections

within the vascular network (yellow arrowhead). Scale bar = 50um.

Figure 7: **ADAMTS1 misregulation leads to failure of optic fissure fusion.**

A) Whole mount Immunohistochemistry was used to simultaneously visualize F-actin

(red) and laminin (green) during optic fissure fusion, 24-72hpf in ADAMTS1 mRNA

injected embryos. Central-proximal sections obtained using confocal imaging were

collected and quantified. Scale bar = 50um. **B)** Quantification of laminin signal intensity

within the optic fissure, normalized to regions of laminin staining juxtaposed to the lens.

Relative pixel intensities are displayed. A green dotted line depicts the trend in laminin

intensity over time, blue line depicts WT trends. ANOVA p<0.0001. **C)** Quantification of

879 F-actin signal intensity (phalloidin staining) within the optic fissure, normalized to

880 regions of F-actin signal within the lobe of the retina. Relative pixel intensities are

881 displayed. A red dotted line represents the trend in F-actin intensity over time, blue line

882 represents WT trends. ANOVA $p < 0.0001$.

Supplemental Material

Figure S1: **Optic Fissure quantification strategy**

A) Graphical representation of the region of OF analyzed using confocal microscopy. **B**) Sample image depicting regions of the OF (1), laminin normalization (2) and f-actin normalization (3) used for signal intensity quantification.

Figure S2: **Optic Fissure apposition measurements.**

Measurements of the distance between retinal lobes (apposition) in WT, pax2a$^{-/-}$, DMH4 treated or ADAMTS1 mRNA injected embryos at 48hpf. Measurements were made using laminin staining as reference for edges of retinal lobes. Distance was measured as pixels.

Figure S3: *pax2a* and *tln1* expression during development

A) Whole mount in situ hybridization of pax2a probe at 24, 48, 54 and 72hpf. Lateral (top) and ventral (bottom) images depicting optic fissure expression are shown. Pax2a expression persists in the fissure up to 54hpf. **B**) Whole mount in situ hybridization of tln1 probe at 24, 48, 54 and 72hpf. Lateral (top) and ventral (bottom) images depicting optic fissure expression are shown. Tln1 expression is detected in the optic fissure from 24-48hpf (yellow arrowheads).

Figure S4: **DMH4 dose response**

3D reconstructions of 48hpf whole mount Tg[kdrl:mCherry] (red) embryos treated with 1, 5, 50 or 100uM DMH4. DNA stained with DAPI (blue). Increasing concentration of DMH4 eliminates mCherry expressing cells from the optic fissure and retina. Scale bar = 50um.

Movie 1: WT Tg[kdrl:mCherry] migration within the optic fissure, 24-30hpf

Movie 2: pax2a$^{-/-}$ Tg[kdrl:mCherry] migration within the optic fissure, 24-30hpf

<u>Movie 3:</u> 3D rotation of WT Tg[kdrl:mCherry] embryo at 48hpf

<u>Movie 4:</u> 3D rotation of pax2a$^{-/-}$ Tg[kdrl:mCherry] embryo at 48hpf

<u>Movie 5:</u> 3D rotation of 48hpf Tg[kdrl:mCherry] embryo injected with ADAMTS1 mRNA

<u>Movie 6:</u> 3D rotation of 48hpf Tg[kdrl:mCherry] embryo injected with ADAMTS1

morpholino

<u>Table 1:</u> pax2$^{-/-}$ 48hpf retinal expression changes.

Figure 1

A

| 24 hpf | 28 hpf | 32 hpf | 36 hpf | 40 hpf | 44 hpf | 48 hpf | 56 hpf | 72 hpf |

laminin

F-actin

merge

B

Optic Fissure Laminin Intensity

Relative pixel intensity

ns   0.0011 **   0.0001 ***   0.0001 ***   ns

24hpf 32hpf 36hpf 40hpf 44hpf 48hpf

C

Optic Fissure F-actin Intensity

Relative pixel intensity

0.0001 ***   0.0001 ***   ns   0.018 *   0.0003 ***

24hpf 32hpf 36hpf 40hpf 44hpf 48hpf

Figure 2

A

B

Pax2a⁻/⁻ Optic Fissure Laminin Intensity

C

Pax2a⁻/⁻ Optic Fissure F-actin Intensity

Figure 3

A

Wild-Type   Pax2 -/-

B

- ■ biological adhesion
- ■ biological regulation
- ■ cellular component of organization or biogenesis
- ■ cellular process
- ■ developmental process
- ■ immune system process
- ■ localization
- ■ locomotion
- ■ metabolic process
- ■ multicellular organismal process
- ■ reproduction
- ■ reponse to stimulus

C

Gene ID	Gene Name	GO Term
aifm1	Apoptosis-inducing factor, mitochondrion-associated 1	oxidoreductase activity
sumo3a	small ubiquitin-like modifier 3a	protein sumoylation
lim2.1	Lens intrinsic membrane protein 2.1	Membrane
tfb2m	transcription factor B2, mitochondrial	transferase activity
tln1	Talin 1	actin binding
mgst2	microsomal glutathione S-transferase 2	enzyme activator activity
si:ch211-57b15.1	si:ch211-57b15.1	NA
ptprna	protein tyrosine phosphatase, receptor type, Na	protein dephosphorylation
dnajb11	DnaJ (Hsp40) homolog, subfamily B, member 11	protein folding
krt96	Keratin 96	structural molecule activity
sept9a	septin 9a	GTP binding
si:rp71-36a1.5	si:rp71-36a1.5	membrane
si:dkey-9c18.3	si:dkey-9c18.3	NA
arf5	ADP Ribosylation Factor 5	GTP binding
dock1	Dedicator Of Cytokinesis 1	small GTPase mediated signal transduction
hbbe1.1	Embryonic 1 beta-globin	Hemoglobin Complex
zgc:152951	Zgc:152951	electron transfer activity
pcbp3	Poly(rC)-binding protein 3	RNA binding
gpsm1b	G protein-signaling modulator 1b	GTPase regulator activity
abracl	Costars family protein ABRACL	regulation of actin filament-based process

D

Wild-Type   Pax2 -/-

E

- ■ biological adhesion
- ■ biological regulation
- ■ cellular component of organization or biogenesis
- ■ cellular process
- ■ developmental process
- ■ immune system process
- ■ localization
- ■ locomotion
- ■ metabolic process
- ■ multicellular organismal process
- ■ reproduction
- ■ reponse to stimulus
- ■ rhythmic process

F

Gene ID	Gene Name	GO Term
gmnn	geminin, DNA replication inhibitor	regulation of DNA replication
col11a1a	collagen, type XI, alpha 1a	extracellular matrix structural constituent
fubp3	far upstream element (FUSE) binding protein 3	RNA binding
si:dkey-26g8.4	Cathepsin Lb	proteolysis
ublcp1	ubiquitin-like domain containing CTD phosphatase 1	protein dephosphorylation
smarcal1	SWI/SNF related, matrix, actin dependent regulator of chromatin, a-like 1	DNA repair
he1b	hatching enzyme 1b	metalloendopeptidase activity
anxa3a	annexin A3a	regulation of NMDA receptor activity
si:dkey-269i1.4	cathepsin L1-like	proteolysis
zgc:174855	zgc:174855	
gsta.1	glutathione S-transferase, alpha tandem duplicate 1	glutathione metabolic process
tbc1d7	TBC1 domain family, member 7	activation of GTPase activity
srek1	splicing regulatory glutamine/lysine-rich protein 1	regulation of alternative mRNA splicing
scg5	secretogranin V	neuropeptide signaling pathway
ggact.3	gamma-glutamylamine cyclotransferase, tandem duplicate 3	lyase activity
si:dkey-26g8.4	Cathepsin L1-like	proteolysis
si:dkey-239j18.3	si:dkey-239j18.3	proteolysis
ahsa1	AHA1, activator of heat shock protein ATPase homolog 1a	ATPase activator activity
he1a	hatching enzyme 1a	metalloendopeptidase activity
dpf3	D4, zinc and double PHD fingers, family 3	zinc ion binding

Figure 4

A

B

C

D

Vascularization of the Optic Fissure

Figure 5

A

DMH4 100μM

kdrl:mCherry DAPI

24 hpf   32 hpf   36 hpf   48hpf   56 hpf   72 hpf

B

DMH4 100μM

laminin

F-actin

merge

24 hpf   32 hpf   36 hpf   48hpf   56 hpf   72 hpf

C

DMH4 treatment Laminin intensity

Relative pixel intensity

<0.0001
****

32hpf   36hpf   48hpf   WT 48hpf

D

DMH4 treatment F-actin intensity

Relative pixel intensity

ns   0.0076   0.0002
***   ****

24hpf   32hpf   36hpf   48hpf

Figure 6

A

ADAMTS1

Wildtype	Pax2-/-	Wildtype	Pax2-/-
28 hpf		36 hpf	

B

kdrl:mCherry

kdrl:mCherry DAPI

Wild Type	ADAMTS1 mRNA	Wild Type	ADAMTS1 mRNA	Wild Type	ADAMTS1 mRNA	Wild Type	ADAMTS1 mRNA
24 hpf		32 hpf		36 hpf		48 hpf	

C

Vascularization of the Optic Fissure

D

kdrl:mCherry

kdrl:mCherry DAPI

Control	ADAMTS1 MO
48 hpf	

Figure 7

A

ADAMTS1 mRNA   Control

laminin

F-actin

merge

24 hpf   32 hpf   36 hpf   48 hpf   48 hpf

B   C

ADAMTS1 overexpression Laminin Intensity

ADAMTS1 overexpression F-actin Intensity

Supplementary Figure 1

A

distal

nasal ← → temporal

proximal

central

B

1: Optic Fissure Signal

2: Laminin signal normalization

3: Phalloidin (F-actin) signal normalization

Supplementary figure 2

Optic Fissure Apposition

Supplementary Figure 3

Supplementary Figure 4

DMH-4

kdrl:mCherry

1 uM   5 uM   50 uM   100 uM

kdrl:mCherry DAPI

48hpf