**Generation of Recombinant Rotavirus Expressing NSP3-UnaG Fusion Protein by a**

**Simplified Reverse Genetics System**

Asha A. Philip[a], Jacob L. Perry[b], Heather E. Eaton[c], Maya Shmulevitz[c], Joseph M. Hyser[b], and

John T. Patton[a#]

[a]Department of Biology, Indiana University, Bloomington, Indiana, USA

[b]Department of Molecular Virology and Microbiology

[c]Department of Medical Microbiology and Immunology, University of Alberta, Edmonton,

Alberta, Canada

**#Corresponding address:** John T. Patton, Department of Biology, Indiana University, 212 S.

Hawthorne Drive, Simon Hall 011, Bloomington, IN 47405 USA

**Running title:** Rotavirus Encoding NSP3-Fluorescent Protein

**Email addresses:**

Asha A. Philip, aaphilip@iu.edu; Heather E. Heaton, heeaton@ualberta.ca; Maya Shmulevitz,

shmulevi@ualberta.ca; John T. Patton. jtpatton@iu.edu

revised_version1

**Abstract**

Rotavirus is a segmented double-stranded (ds)RNA virus that causes severe gastroenteritis in

young children. We have established an efficient simplified rotavirus reverse genetics (RG)

system that uses eleven T7 plasmids, each expressing a unique simian SA11 (+)RNA, and a

CMV support plasmid for the African swine fever virus NP868R capping enzyme. With the

NP868R-based system, we generated recombinant rotavirus (rSA11/NSP3-FL-UnaG) with a

genetically modified 1.5-kB segment 7 dsRNA that encodes full-length NSP3 fused to UnaG, a

139-aa green fluorescent protein (FP). Analysis of rSA11/NSP3-FL-UnaG showed that the virus

replicated efficiently and was genetically stable over 10 rounds of serial passage. The NSP3-

UnaG fusion product was well expressed in rSA11/NSP3-FL-UnaG-infected cells, reaching

levels similar to NSP3 in wildtype rSA11-infected cells. Moreover, the NSP3-UnaG protein, like

functional wildtype NSP3, formed dimers *in vivo*. Notably, NSP3-UnaG protein was readily

detected in infected cells via live cell imaging, with intensity levels ~3-fold greater than that of

the NSP1-UnaG fusion product of rSA11/NSP1-FL-UnaG.  Our results indicate that FP-

expressing recombinant rotaviruses can be made through manipulation of the segment 7 dsRNA

without deleting or interrupting any of the twelve open reading frames of the virus. Because

NSP3 is expressed at levels higher than NSP1 in infected cells, rotaviruses expressing NSP3-

based FPs may be a more sensitive tool for studying rotavirus biology than rotaviruses

expressing NSP1-based FPs. This is the first report of a recombinant rotavirus containing a

genetically engineered segment 7 dsRNA.

**Importance**

Previous studies have generated recombinant rotaviruses that express fluorescent proteins (FPs)

by inserting reporter genes into the NSP1 open reading frame (ORF) of genome segment 5.

Unfortunately, NSP1 is expressed at low levels in infected cells, making viruses expressing FP-

fused NSP1 less than ideal probes of rotavirus biology. Moreover, FPs were inserted into

segment 5 in such a way as to compromise NSP1, an interferon antagonist affecting viral growth

and pathogenesis. We have identified an alternative approach for generating rotaviruses

expressing FPs, one relying on fusing the reporter gene to the NSP3 ORF of genome segment 7.

This was accomplished without interrupting any of the viral ORFs, yielding recombinant viruses

likely expressing the complete set of functional viral proteins. Given that NSP3 is made at

moderate levels in infected cells, rotavirus encoding NSP3-based FPs should be more sensitive

probes of viral infection than rotaviruses encoding NSP1-based FPs.

## Introduction

Group A rotavirus (RVA) is a primary cause of acute gastroenteritis in infants and
children under 5 years of age (Crawford et al., 2017). The RVA genome consists of eleven
segments of double-stranded (ds)RNA, with a total size of 18.5 kbp, and is contained within a
non-enveloped icosahedral capsid composed of three concentric protein layers (Settembre et al.,
2011). During replication, the genome segments are transcribed, producing eleven 5'-capped, but
non-polyadenylated (+)RNAs (Imai et al., 1983; Trask et al., 2012). The (+)RNAs are generally
monocistronic, each with a single open reading frame (ORF) that specifies one of the six
structural (VP1-VP4, VP6-VP7) or six nonstructural (NSP1-NSP6) proteins of the virus. The
(+)RNAs also serve as templates for the synthesis of dsRNA genome segments (Gugliemli et al.,
2010).

Insights into RVA biology have been severely hampered by the lack of a reverse genetics
(RG) system to unravel details into the replication and pathogenesis mechanisms of the virus.
This limitation was recently overcome by Kanai et al. (2017), who described the development of
a fully plasmid-based RG system that allowed genetic engineering of the prototypic RVA simian
SA11 strain. Key to the RG system was co-transfection of baby hamster kidney cells expressing
T7 RNA polymerase (BHK-T7 cells) with T7 plasmids directing synthesis of the eleven SA11
(+)RNAs, two CMV plasmids encoding the vaccinia virus D1L-D12R capping enzyme complex
(Kyrieleis et al., 2014), and another CMV plasmid encoding the avian reovirus p10FAST fusion
protein (Salsman et al., 2005). Subsequent publications described changes to the Kanai RG
system designed to reduce its complexity and/or enhance the recovery of recombinant virus.
Notably, Komoto et al (2018) showed that recombinant virus could be produced simply by

transfecting BHK-T7 cells with eleven SA11 T7 plasmids, with the caveat that plasmids for the

viroplasm building blocks (NSP2 and NSP5) (Fabbretti et al., 1999; Eichwald et al., 2004) be

added at levels three-fold greater than the other plasmids. Of possible significance, the Komoto

RG system used a set of SA11 T7 plasmids with vector backbones that differed in size and

sequence from the SA11 T7 plasmids described by Kanai et al (2017).

Plasmid-based RG systems have been used to modify several SA11 genome segments,

with the focus mostly on segment 5, which encodes NSP1 (Kanai et al., 2017, 2018; Komoto et

al., 2018). Through insertion of reporter genes into the segment 5 dsRNA, recombinant RVAs

have been produced that express fluorescent reporter proteins (FPs) (e.g., mCherry, eGFP)

(Kanai et al., 2017, 2018; Komoto et al., 2018); these FP-RVAs are important tools for studying

virus replication and pathogenesis via live cell imaging and other fluorescence-based approaches.

Unfortunately, the segment 5 product NSP1 is expressed at low levels in infected cells and is

subject to proteasomal degradation making FP-fused NSP1 proteins less than ideal probes of

RVA biology (Martinez-Alvarez et al., 2013). Moreover, FP genes were inserted into the

segment 5 dsRNA in such a way as to alter the NSP1 open reading frame (ORF), likely

compromising the protein's function as an IFN-antagonist (Barro and Patton, 2005; Davis and

Patton, 2017) and, thus, impacting the virus's biological properties.

In this study, we explored an alternative approach for making FP-RVAs, one relying on

modification of the genome segment that expresses NSP3 (segment 7), a viral translation

enhancer expressed at moderate levels in the infected cell that may not be required for virus

replication (Montero et al., 2006; Gratia et al., 2015). In generating recombinant RVAs, we

employed a simplified RG system requiring only a single support plasmid: a CMV expression

vector for the African swine fever virus (ASFV) NP868R capping enzyme (Dixon et al., 2013).

Using the NP868R-based RG system, we produced a novel SA11 strain with modified segment 7

dsRNA that expressed NSP3 fused to the green fluorescent protein (FP) UnaG. This is the first

RVA strain engineered to produce a FP that did not involve deleting or interrupting any of one of

the 12 viral ORFs, thus yielding a recombinant virus likely expressing a complete set of

functional viral proteins.

**Materials and methods**

*Cell culture.* Embryonic monkey kidney cells (MA104) were grown in M199 complete

medium [Medium 199 (Lonza) and 1% penicillin-streptomycin (P/S) (Corning)] containing 5%

fetal bovine serum (FBS) (Gibco) (Arnold et al., 2009). BHK-T7 cells were a kind gift from Drs.

Ulla Buchholz and Peter Collins, Laboratory of Infectious Diseases, NIAID, NIH.  BHK-T7 cells

were grown in Glasgow complete medium [Glasgow minimum essential medium (G-MEM)

(Lonza), 10% tryptone-peptide broth (Gibco), 1% P/S, 2% non-essential amino acids (NEAA)

(Gibco), and 1% glutamine (Gibco)] containing 5% heat-inactivated FBS. Medium used to

cultivate BHK-T7 cells was supplemented with 2% G418 geneticin (Invitrogen) every other

passage.

*Plasmid construction.* RVA (simian SA11 strain) plasmids pT7/VP1SA11,

pT7/VP2SA11, pT7/VP3SA11, pT7/VP4SA11, pT7/VP6SA11, pT7/VP7SA11, pT7/NSP1SA11,

pT7/NSP2SA11, pT7/NSP3SA11, pT7/NSP4SA11, and pT7/NSP5SA11 were kindly provided

by Dr. Takeshi Kobayashi (Kanai et al., 2017) through the Addgene plasmid repository

{https://www.addgene.org/Takeshi_Kobayashi/}. To generate the pCMV/NP868R plasmid, a

DNA representing the ASFV NP868R ORF (Genbank NP_042794), bound by upstream *Xba*I

and downstream *Bam*HI sites, was synthesized by Genscript and inserted into the *Eco*RV site of

the pUC57 plasmid. A DNA fragment containing the NP868R ORF was recovered from the

plasmid by digestion with *Not*I and *Pvu*I and ligated into the plasmid pCMV-Script (Agilent

Technologies), cut with the same two restriction enzymes. pT7/NSP3-FL-UnaG was constructed

by fusing a DNA fragment containing the ORF for 3xFL-UnaG to the 3'-end of the NSP3 ORF in

pT7/NSP3SA11 using a Takara In-fusion HD cloning kit. The 3xFL-UnaG fragment was

amplified from the reovirus S1 3xFL-UnaG plasmid (Eaton et al, 2017) using the primer pair 5'-

tcatttggttgcgaagactacaaagaccatgacggtgattataaaga-3'

and 5'-catgtatcaaaatggtcattctgtggcccttctgtagct-3' and the pT7/NSP3SA11 plasmid was amplified

using the NSP3 primer pair 5'-ccattttgatacatgttgaacaatcaaatacagtgt-3' and 5'-

ttcgcaaccaaatgaatattgataattacatctctgtattaat-3'. pT7/NSP1-FL-UnaG was constructed by inserting

the 3xFL-UnaG fragment into the NSP1 ORF of the segment 5 cDNA (position 1199) in

pT7/NSP1SA11. For cloning, the pT7/NSP1SA11 plasmid was amplified using the NSP1 primer

pair 5'- tgaagaagtgtttaatcacatgtcgcc-3' and 5'- tttgatccatgtgattagtaaacaaactccaaa-3 and the 3xFL-

UnaG insert was amplified from pT7/NSP3-R2A-FL-UnaG using the primer pair 5'-

tcacatggatcaaacctacaaagaccatgacggtgattataaagatcat-3' and 5'-

ttaaacacttcttcatcattctgtggcccttctgtagc-3'. Transfection quality plasmids were prepared

commercially (www.plasmid.com) or using a QIAprep spin miniprep kit. Primers and

sequencing services were provided by Eurofin Genomics. Sequences of recombinant RVAs were

determined from cDNAs prepared from viral RNAs using a Superscript III reverse transcriptase

kit (Invitrogen).

*Optimized RVA RG protocol.* On Day 0, freshly confluent monolayers of BHK-T7 cells

were disrupted using trypsin-versene and resuspended in G418-free Glasgow complete medium

containing 5% FBS. Cell numbers were determined with a Nexcelom Cellometer AutoT4

counter. The cells were seeded in the same medium into 12-well plates (2-4 x $10^5$ cells/well). On

Day 1, plasmid mixtures were prepared that contained 0.8 μg each of the 11 RVA pT7 plasmids,

except pT7/NSP2SA11 and pT7/NSP5SA11, which were 2.4 μg each. Included in plasmid

mixtures was 0.8 μg of pCMV/NP868R. The plasmid mixtures were added to 100 μl of pre-

warmed (37°C) Opti-MEM (Gibco) and mixed by gently pipetting up and down. Afterwards, 25

μl of TransIT-LTI transfection reagent (Mirus) was added, and the transfection mixtures gently

vortexed and incubated at room temperature for 20 min. During the incubation period, BHK-T7

cells in 12-well plates were washed once with Glasgow complete medium, and then 1 ml of

SMEM complete medium [MEM Eagle Joklik (Lonza), 10% TBP, 2% NEAA, 1% P/S, and 1%

glutamine)] was placed in each well. The transfection mixture was added drop-by-drop to the

medium in the wells and the plates returned to a 37°C incubator. On Day 3, 2 x$10^5$ MA104 cells

in 250 μl of M199 complete medium were added to wells, along with trypsin to a final

concentration of 0.5 μg/ml (porcine pancreatic type IX, Sigma Aldrich). On Day 5, cells in plates

were freeze-thawed 3-times and the lysates placed in 1.5 ml microfuge tubes. After

centrifugation at 500 x g for 10 min (4°C), 300 μl of the supernatant was transferred onto

MA104 monolayers in 6-well plates containing 2 ml of M199 complete medium and 0.5 μg/ml

trypsin. The plates were incubated at 37°C for 7 days or until complete cytopathic effects (CPE)

were observed. Typically, complete CPE was noted at 4-6 days p.i. for wells containing

replicating RVA.

*Analysis of recombinant viruses*. RVAs were propagated in MA104 cells in M199

complete medium containing 0.5 μg/ml trypsin. Viruses were isolated by plaque purification and

titered by plaque assay or fluorescence focus assay on MA104 cells (Arnold et al., 2009). To

determine peak viral titers, virus isolates were grown in parallel in MA104 cell monolayers until

cytopathic effects reached 100%. Plaque titers in lysates were calculated from six independent

plaque assays. Viral RNAs were recovered from clarified infected-cell lysates by Trizol

extraction, resolved by electrophoresis on Novex 8% polyacrylamide gels (Invitrogen), and

detected by staining with ethidium bromide.

For immunoblot assays, proteins in lysates prepared at 8 h p.i. from MA104 cells infected

with RVA at an MOI of 5 were resolved by electrophoresis on Novex linear 8-16%

polyacrylamide gels and transferred to nitrocellulose membranes. After blocking with phosphate-

buffered saline containing 5% nonfat dry milk, blots were probed with guinea pig polyclonal

anti-NSP3 (Lot 55068, 1:2000) or anti-VP6 (Lot 53963, 1:2000) antisera (Arnold et al., 2012), or

mouse monoclonal Flag M2 (Sigma F1804, 1:2000) or rabbit monoclonal PCNA (13110S, Cell

Signaling Technology (CST), 1:1000) antibody. Primary antibodies were detected using

1:10,000 dilutions of horseradish peroxidase (HRP)-conjugated secondary antibodies: horse anti-

mouse IgG (CST), anti-guinea pig IgG (KPL), or goat anti-rabbit IgG (CST). Signals were

developed using Clarity Western ECL Substrate (Bio-Rad) and detected using a Bio-Rad

ChemiDoc imaging system. Image J analysis was used to determine the intensity of bands on

immunoblots {https://imagej.net/ImageJ}.

*Assessment of genetic stability*. MA104 cell monolayers in 6-well plates were infected

with recombinant RVA at an MOI of ~0.1. When cytopathic effects were complete (4-5 days

p.i.), the cells were freeze-thawed twice in their medium, and lysates were centrifuged at low

speed to remove debris. Virus in clarified lysates was serially passaged 10-times, infecting

MA104 cells with 2 µl of lysate combined with 2 ml of fresh M199 complete medium. To

analyze viral dsRNA content, 0.6 ml of clarified lysates were incubated with 1 µl of RNase T1

(Fermentas, 1000 U/ml) for 30 min at 37°C, extracted with Trizol, and pelleted by ethanol

precipitation. Purified dsRNA was resolved by electrophoresis on 8% polyacrylamide gels and

detected by staining with ethidium bromide.  Plaque purification was used to recover six virus

isolates from the passage 10 pool of rSA11/NSP3-FL-UnaG. The segment 6 (VP6) and 7 (NSP3)

RNAs of three isolates were sequenced by Eurofins Genomics.

*Live-cell imaging of UnaG fluorescence in infected cells.* MA104 cells were seeded at a

density of 25,000 cells per well in a 96-well optical bottom plate (Greiner Bio-One) and grown

for 2 days at $37°C/5\%$ $CO_2$ before being infected with rSA11/NSP1-FL-UnaG or rSA11/NSP3-

FL-NSP3-UnaG at ~4 fluorescent focus units per cell. After 1 h the inoculum was removed and

200 μl FluoroBrite (Thermo Fisher Scientific) was added to each well for maintenance media.

The plate was mounted into an Oko Labs stage-top environmental chamber equilibrated to

$37°C/5\%$ $CO_2$. Wells were imaged with a widefield epifluorescence Nikon TiE inverted

microscope using a SPECTRAX LED light source (Lumencor) and a 20x Plan Fluor (NA 0.45)

phase contrast objective. Fluorescence light images were recorded using an ORCA-Flash 4.0

sCMOS camera (Hamamatsu), and Nikon Elements Advanced Research v4.5 software was used

for multipoint position selection, data acquisition, and image analysis. Single cells were selected

as Regions of Interest (ROI) and fluorescence intensity measured for the experiment.

Fluorescence intensity values were exported to Microsoft Excel and normalized to the initial

baseline fluorescence value. The baseline and maximum fluorescence values for each ROI

selected were calculated and from this the change in fluorescence ($\Delta F$) was calculated by

subtracting the maximum and baseline values. The $\Delta F$ was plotted as percent change and

biostatistical analyses was performed using GraphPad Prism (version 8.1) with results presented

as mean ± standard deviation. Comparisons used a nonparametric Mann-Whitney test and

differences were considered statistically significant for $p < 0.05$.

*Nucleotide sequence accession numbers.* pCMV/NP868R, MH212166; pT7/NSP3-FL-

UnaG, MK868472; and, pT7/NSP1-FL-UnaG, MH197081.

**Results and Discussion**

*NP868R-based RG system*. A previous study showed that addition of a CMV plasmid

expressing the ASFV NP868R capping enzyme to the mammalian reovirus RG system

significantly increased recovery of recombinant virus (Eaton et al., 2017). Based on this finding,

we constructed a similar CMV plasmid for NP868R (pCMV/NP868R) and evaluated whether its

co-transfection with SA11 pT7 plasmids into BHK-T7 cells was sufficient to allow recovery of

recombinant RVAs. In our RG experiments, two different types of pT7 vectors for segment 7

(NSP3) were used, one (pT7/NSP3SA11) designed to introduce a wild type segment 7 dsRNA

into recombinant RVA and the second (pT7/NSP3-FL-UnaG) designed to test the possibility of

introducing a modified segment 7 RNA that expressed NSP3 with a fluorescent tag. To construct

pT7/NSP3-FL-UnaG, the C-terminus of the NSP3 ORF in pT7/NSP3SA11 was fused to the ORF

for UnaG, a 139-amino acid (aa) green fluorescent protein of the Unagi eel that utilizes bilirubin

as a fluorophore (Kumagai et al., 2013) (Fig. 1). To further ease detection of the protein product

of the modified pT7 plasmid, a 3x Flag sequence was inserted between the NSP3 ORF and UnaG

ORF. As a result of the addition of Flag and UnaG sequences, the pT7/NSP3-FL-UnaG vector

expresses a 1.6-kb RNA that encodes a 477-aa protein instead of the 1.1-kb RNA and 315-aa

protein of pT7/NSP3SA11 (Fig. 1). The NSP3-FL-UnaG fusion protein retained the same RNA-

binding domain, coiled-coil dimerization domain, and eIF4G-binding domain present in wildtype

NSP3 (Deo et al., 2002; Groft et al., 2002 ).

Rotavirus RG experiments were performed using stocks of BHK-T7 cells maintained in

Glasgow medium enriched with FBS, tryptone-peptide broth, and non-essential amino acids.

Mixtures of RG plasmids were transfected into BHK-T7 cells that were ~90% confluent and had

been seeded into 12-well plates the day before. In our optimized protocol, plasmid mixtures

included pCMV/NP868R, 3x levels of pT7/NSP2SA11 and pT7/NSP5SA11, either

pT7/NSP3SA11 or pT7/NSP3-FL-UnaG, and the remaining SA11 pT7 plasmids. Transfected

BHK-T7 cells were over seeded with MA104 cells to promote amplification of recombinant

viruses. At 5 days post transfection, the cells were freeze-thawed thrice and large debris removed

by low speed centrifugation. Recombinant viruses in cell lysates were amplified by passage in

MA104 cells, plaque isolated, and amplified again in MA104 cells.

*Recombinant SA11 expressing fused NSP3-UnaG.* Analysis of the cell lysates showed

that transfection of BHK-T7 cells with RG plasmid mixtures using the optimized protocol

supported the generation of recombinant RVAs, including SA11 isolates with a wildtype

segment 7 (NSP3) dsRNA (rSA11/wt) or a modified segment 7 RNA (rSA11/NSP3-FL-UnaG)

(Fig. 2). The identity of the modified segment 7 dsRNA in rSA11/NSP3-FL-UnaG was verified

by gel electrophoresis (Fig. 2A), which revealed that the wildtype 1-kB segment 7 dsRNA had

been replaced with a segment co-migrating with the 1.5-kB segment 5 (NSP1) dsRNA. The

authenticity of the segment 7 dsRNA in rSA11/NSP3-FL-UnaG was confirmed by RT-PCR and

sequencing (data not shown). Immunoblot analysis with anti-NSP3 and anti-FLAG antibodies of

infected cell lysates indicated that segment 7 of rSA11/NSP3-FL-UnaG expressed a protein of

the expected size (55 kD) for NSP3-FL-UnaG and did not express the wildtype 37-kD NSP3

(Fig. 2B). As a probe of the properties of NSP3-FL-UnaG, we examined whether the protein was

able to form dimers in infected cells, as previously reported for wildtype NSP3 (Arnold et al,

2012). Indeed, electrophoretic analysis of rSA11/NSP3-UnaG-infected cell lysates treated with

denaturing sample buffer at 25°C showed that the NSP3-FL-UnaG migrated as a dimer (Fig. 3),

suggesting that the NSP3 coiled-coil dimerization domain retains its function in the fusion

product. Under these same electrophoretic conditions, the VP6 inner capsid protein of both

rSA11/NSP3-FL-UnaG and rSA11/wt formed trimers that were stable at 25°C (Fig. 3) (Clapp

and Patton, 1991). Quantitation of bands appearing on immunoblots probed with anti-NSP3 and

anti-VP6 antibodies indicated that the steady-state level of NSP3-FL-UnaG (normalized to VP6)

in rSA11/NSP3-FL-UnaG-infected cells approximated that of the steady-state level of NSP3 in

rSA11/wt infected cells (Fig. 3, lanes 4 versus 6). Thus, the segment 7 (+)RNAs of rSA11/NSP3-

FL-UnaG and rSA11/wt appear to be translated with near equal efficiency.

Plaque analysis showed that rSA11/wt grew to a peak titer in MA104 cells that was ~2-3-

fold greater than rSA11/NSP3-FL-UnaG (1.6 - 4.8 x $10_7$ and 0.6 - 2.0 x $10_7$, respectively) and

generated plaques that were ~2-fold larger (Fig. 4). Ten rounds of serial passage of rSA11/NSP3-

FL-UnaG at low MOI revealed no difference in the dsRNA profile of the starting virus and the

passage 10 virus, suggesting that the recombinant RVA was genetically stable (Fig. 5). To

further evaluate this issue, the segment 6 (VP6) and 7 (NSP3) RNAs of three plaque isolates

recovered from the passage 10 pool of rSA11/NSP3-FL-UnaG were sequenced. The sequences

of segment 6 and 7 RNAs were the same as those present in the pT7/VP6SA11 and pT7/NSP3-

FL-UnaG plasmids used to make the recombinant viruses. Thus, based on serial passage and

sequencing of P10 viruses, rSA11/NSP3-FL-UnaG appears to stably retain its foreign sequence.

*Fluorescence signal of rSA11/NSP3-FL-UnaG*. Examination of MA104 cells infected

with rSA11/NSP3-FL-UnaG by live-cell imaging from 0 to 16 h p.i. confirmed that UnaG was

functional, emitting fluorescent light in a range overlapping green fluorescent protein (GFP)

(Fig. 6A, Movie S1) (Rodriguez et al., 2017). At early times of infection (0 to 8 h p.i.), the signal

localized predominantly to the cytoplasm. At later times, UnaG fluorescent signal was also

readily detected in the nucleus (Move S1). To contrast the intensity of the fluorescent signal

produced by recombinant viruses expressing UnaG fused to NSP3 versus NSP1, we generated

rSA11/NSP1-FL-UnaG (Fig. 7) using the optimized NP868R-based RG protocol. To produce

this virus, a FL-UnaG ORF terminating with a stop codon was inserted into NSP1 ORF of the

segment 5 cDNA of pT7/NSP1SA11. As illustrated in Fig. 7, pT7/NSP1-FL-UnaG produces a

2.1-kb RNA that encodes a 575-aa protein instead of the 1.6-kb RNA and 520-aa protein of

pT7/NSP1SA11. The segment-5 protein product of pT7/NSP1-FL-UnaG ends with the same

Flag-UnaG cassette as the segment-7 protein product of pT7/NSP3-FL-UnaG.

As determined by gel electrophoresis (Fig. 7) and sequencing (not shown), the genome of

rSA11/NSP1-FL-UnaG included the expected large 2.1-kB segment 5 dsRNA. Immunoblot

analysis of rSA11/NSP1-FL-UnaG-infected cell lysates using anti-Flag antibody indicated that

the virus also encoded the NSP1-FL-UnaG product. Based on plaque assay, rSA11/NSP1-FL-

UnaG grew to a peak titer (0.8 - 1.0 x $10_7$) that was ~3-fold less than rSA11/wt and produced

plaques on MA104 cells that were ~2-fold smaller than rSA11/wt (Fig. 4) The small plaque

phenotype of rSA11/NSP1-FL-NSP1 is similar to small plaque phenotype described before for

RVAs encoding truncated or altered NSP1 proteins (Patton et al., 2001).

Quantification of UnaG fluorescence signals generated by MA104 cells infected with

rSA11/NSP3-FL-UnaG were approximately 3-times greater than those generated by cells

infected with rSA11/NSP1-FL-UnaG (Fig. 6B). This result suggests that RVAs expressing fused

NSP3-FPs may be more sensitive probes of viral infection that RVAs expressing fused NSP1-

FPs.

*Stem-loop structure in the segment 7 3'-UTR*. rSA11/NSP3-FL-UnaG was generated by

placing a nonviral 500-bp insert into the segment 7 dsRNA at the junction of the NSP3 ORF and

3'-UTR. Similar recombinant RVAs have been made by inserting nonviral sequences between

the NSP2 ORF and 3'-UTR of the segment 8 dsRNA (Navarro et al., 2013). Thus, the junction

between the viral ORF and 3'-UTR may represent a site well suited for the introduction of long

foreign sequences into RVA genome segments. Interestingly, for RVAs with naturally occurring

genome rearrangements, this is the same site in segment 5, 6, 7, 10 and 11 dsRNAs in which

viral sequence duplications have been noted to initiate (Ballard et al., 1992; Shen et al., 1994;

Gault et al., 2001; Patton et al., 2001; Arnold et al., 2012). The 3'-UTR contains multiple *cis-*

acting signals important to rotavirus replication, including sequences that are recognized by the

RVA RNA polymerase VP1 and the translation enhancer NSP3 (Deo et al., 2002; Tortorici et al.,

2003). The fact that well-growing genetically-stable recombinant RVAs have been recovered in

which a viral ORF has been separated from its 3'-UTR indicates that *cis*-acting signals in the 3'-

UTR continue to function even though displaced linearly a long distance from the remaining

viral sequence of the RNA.

In a previous study (Navarro et al., 2013), an *in silico* RNA folding analysis

{*http://rna.tbi.univie.ac.at/cgi-bin/RNAfold.cgi*} was performed to probe how the insertion of

sequence duplications and foreign sequences effected the predicted secondary structure of the

mutant segment 8 (+)RNAs used in making recombinant RVAs. The results showed that despite

extensive differences in the overall folding predictions of the mutant RNAs, in all cases their 5'-

and 3'-UTRs interacted to form stable 5'-3' panhandles. In addition, the predictions all revealed

an identical stem-loop structure projecting from the 5' side of the 5'-3 panhandle, formed by

residues that are highly conserved among RVA segment 8 RNAs. The conservation of the

structure and its sequence suggested that the stem-loop may function as a segment specific

packaging signal (Navarro et al., 2013). We performed a similar *in silico* RNA folding analysis,

contrasting the secondary structures predicted for the segment 7 RNAs of rSA11/wt and

rSA11/NSP3-FL-UnaG. The results showed that the overall secondary structures predicted for

the RNAs differed considerably, with the notable exception that extending from the 3'-UTR of

both RNAs was a long (~70 base) stable stem-loop structure formed by sequences that are highly

conserved in RVA segment 7 RNAs (Fig. 8). The stability and location of the stem-loop suggests

that this structure may function as a segment specific packaging signal, in a manner previously

proposed for the conserved stem-loop detected in the segment 8 RNA.

*Summary*. rSA11/NSP3-FL-UnaG is the first recombinant RVA to be described with a

modified segment 7 dsRNA. Segment 7 joins segments 4 (VP4) (Johne et al., 2015; Mohanty et

al., 2017), 5 (NSP1) (Kanai et al., 2019), 8 (NSP2) (Trask et al., 2010; Navarro et al., 2013), and

11 (NSP5/NSP6) (Komoto et al., 2017) as targets altered by RG and represents only the second

RVA segment to be used as a vector for FP expression. Our analysis of rSA11/NSP3-FL-UnaG

indicates that it is possible to generate recombinant RVAs that express FPs through their fusion

to the C-terminus of NSP3. Given that NSP3 is expressed at moderate levels in infected cells,

RVAs expressing NSP3-based FPs may be more effective indicators of viral replication in live

cell imagining experiments and other fluorescence-based assay systems than RVAs expressing

NSP1-based FPs, since NSP1 is expressed at low levels *in vivo* (Martinez-Alvarez et al., 2013).

Although several recombinant RVAs that express FPs have been described, rSA11/NSP3-FL-

UnaG is unique among them in that none of its ORFs have been deleted or interrupted. Instead,

the only impact on rSA11/NSP3-FL-UnaG was to fuse its NSP3 ORF to a FL-UnaG ORF.

Importantly, although the NSP3 ORF in RVA strains is not naturally extended and does not

encode NSP3 fused to a downstream protein, the NSP3 ORF of group C rotaviruses (RVCs) is

extended, encoding an NSP3 protein that is fused to a 2A stop-start translational element

(Donnelly et al., 2001) and double-stranded RNA binding protein (dsRBP) (James et al., 1999;

Langland et al., 1994). Given that RVC segment 6 encodes an NSP3 fusion protein, it seems

likely that the NSP3 fusion protein of rSA11/NSP3-FL-UnaG remains functional, even when

fused to a downstream protein. Interestingly, despite repeated attempts, we were unsuccessful in

generating recombinant RVAs using mutated pT7/NSP3SA11 plasmids in which the NSP3 ORF

was interrupted through insertion of stop codons (data not shown). This result implies that NSP3

is essential for RVA replication or is required to generate recombinant viruses using the RG

system.

Our results suggest that the RVA segment 7 RNA can be re-engineered to function as an

expression vector of foreign proteins, without compromising the function of any of the viral

ORFs. It remains unclear how much foreign sequence can be inserted into the segment 7 RNA,

but our analysis so-far indicates that it is possible to generate well replicating viruses carrying

>500 bp of extra sequence. Given the remarkable flexibility so far noted in the ability of the

rotavirus to accommodate changes in the size and sequences of its RNA, the virus may turn out

to be particularly empowering tool for unraveling the shared mechanisms used by the *Reoviridae*

to package and replicate their genomes.

**Acknowledgments.** We are grateful to all the members of the Patton and Danthi laboratory for

their support and encouragement on this project. Our thanks also go to Ulla Buckholtz and Peter

Collins, NIAID, NIH for their gift of BHK-T7 cells. This work was supported by National

Institutes of Health grants R03 AI131072 and R21 AI144881, Indiana University Start-Up

Funding, and the Lawrence M. Blatt Endowment.

**References**

**References**

1. Arnold, M., Patton, J.T., McDonald, S.M. 2009. Culturing, storage, and quantification of
rotaviruses. Curr Protoc Microbiol Chapter 15:Unit 15C.3.

2. Arnold, M.M., Brownback, C.S., Taraporewala, Z.F., Patton, J.T. 2012. Rotavirus variant
replicates efficiently although encoding an aberrant NSP3 that fails to induce nuclear
localization of poly(A)-binding protein. J. Gen. Virol. 93:1483-1494.

3. Ballard, A., McCrae, M.A., Desselberger, U. 1992. Nucleotide sequences of normal and
rearranged RNA segments 10 of human rotaviruses. J. Gen. Virol. 73:633-638.

4. Barro, M., Patton, J.T. 2005. Rotavirus nonstructural protein 1 subverts innate immune
response by inducing degradation of IFN regulatory factor 3. Proc. Natl. Acad. Sci. USA
102:4114-4119.

5. Clapp, L.L., Patton, J.T. 1991. Rotavirus morphogenesis: domains in the major inner capsid
protein essential for binding to single-shelled particles and for trimerization. Virology
180:697-708.

6. Crawford, S.E., Ramani, S., Tate, J.E., Parashar, U.D., Svensson, L., Hagbom, M., Franco,
M.A., Greenberg, H.B., O'Ryan, M., Kang, G., Desselberger, U., Estes, M.K. 2017.
Rotavirus infection. Nat. Rev. Dis. Primers 3:17083.

7. Davis, K.A., Patton, J.T. 2017. Shutdown of interferon signaling by a viral-hijacked E3
ubiquitin ligase. Microb. Cell. 4:387-389.

8. Deo, R.C., Groft, C.M., Rajashankar, K.R., Burley, S.K. 2002. Recognition of the rotavirus
mRNA 3' consensus by an asymmetric NSP3 homodimer. Cell 108:71-81.

9. Dixon, L.K., Chapman, D.A., Netherton, C.L., Upton, C. 2013. African swine fever virus
replication and genomics. Virus Res. 173:3-14.

10. Donnelly, M.L., Hughes, L.E., Luke, G., Mendoza, H., ten Dam, E., Gani, D., Ryan, M.D.

2001. The 'cleavage' activities of foot-and-mouth disease virus 2A site-directed mutants and

naturally occurring '2A-like' sequences. J. Gen. Virol. 82:1027-1041.

11. Eaton, H.E., Kobayashi, T., Dermody, T.S., Johnston, R.N., Jais, P.H., Shmulevitz, M. 2017.

African swine fever virus NP868R capping enzyme promotes reovirus rescue during reverse

genetics by promoting reovirus protein expression, virion assembly, and RNA incorporation

into infectious virions. J. Virol. 91:e02416-16.

12. Eichwald, C., Rodriguez, J.F., Burrone, O.R. 2004. Characterization of rotavirus NSP2/NSP5

interactions and the dynamics of viroplasm formation. J. Gen. Virol. 85:625-634.

13. Fabbretti, E., Afrikanova, I., Vascotto, F., Burrone, O.R.. 1999. Two non-structural rotavirus

proteins, NSP2 and NSP5, form viroplasm-like structures in vivo. J. Gen. Virol. 80:333-339.

14. Gault, E., Schnepf, N., Poncet, D., Servant, A., Teran, S., Garbarg-Chenon, A. 2001. A

human rotavirus with rearranged genes 7 and 11 encodes a modified NSP3 protein and

suggests an additional mechanism for gene rearrangement. J. Virol. 75:7305-7314.

15. Gratia, M., Sarot, E., Vende, P., Charpilienne, A., Baron, C.H., Duarte, M., Pyronnet, S.,

Poncet, D. 2015. Rotavirus NSP3 is a translational surrogate of the poly(A)-binding protein-

poly(A) complex. J. Virol. 89:8773-8782.

16. Groft, C.M., Burley, S.K. 2002. Recognition of eIF4G by rotavirus NSP3 reveals a basis for

mRNA circularization. Mol. Cell. 9:1273-1283.

17. Guglielmi, K.M., McDonald, S.M., Patton, J.T. 2010. Mechanism of intraparticle synthesis

of the rotavirus double-stranded RNA genome. J. Biol. Chem. 285:18123-18128.

18. Hofacker, I.L. Vienna RNA secondary structure server. 2003. Nucleic Acids Res. 31:3429-

3431.

19. Hofacker, I.L., Fontana, W., Stadler, P.F., Bonhoeffer, S., Tacker, M., Schuster, P. 1994. Fast

folding and comparison of RNA secondary structures. Monatsh. Chemie 125:167–188.

20. Imai, M., Akatani, K., Ikegami, N., Furuichi, Y. 1983. Capped and conserved terminal

structures in human rotavirus genome double-stranded RNA segments. J. Virol. 47:125–136.

21. James, V.L., Lambden, P.R., Deng, Y., Caul, E.O., Clarke, I.N. 1999. Molecular

characterization of human group C rotavirus genes 6, 7 and 9. J. Gen. Virol. 80:3181-3187.

22. Johne, R., Reetz, J., Kaufer, B.B., Trojnar, E. 2015. Generation of an avian-mammalian

rotavirus reassortant by using a helper virus-dependent reverse genetics system. J. Virol.

90:1439-1443.

23. Kanai, Y., Komoto, S., Kawagishi, T., Nouda, R., Nagasawa, N., Onishi, M., Matsuura, Y.,

Taniguchi, K., Kobayashi, T. 2017. Entirely plasmid-based reverse genetics system for

rotaviruses. Proc. Natl. Acad. Sci. USA 114:2349-2354.

24. Kanai, Y., Kawagishi, T., Nouda, R., Onishi, M., Pannacha, P., Nurdin, J.A., Nomura, K.,

Matsuura, Y., Kobayashi, T. 2018. Development of stable rotavirus reporter expression

systems. J. Virol. 93: e01774-18.

25. Komoto, S., Kanai, Y., Fukuda, S., Kugita, M., Kawagishi, T., Ito, N., Sugiyama, M.,

Matsuura, Y., Kobayashi, T., Taniguchi, K. 2017. Reverse genetics system demonstrates that

rotavirus nonstructural protein NSP6 is not essential for viral replication in cell culture. J.

Virol. 91 pii: e00695-17.

26. Komoto, S., Fukuda, S., Ide, T., Ito, N., Sugiyama, M., Yoshikawa, T., Murata, T.,

Taniguchi, K. 2018. Generation of recombinant rotaviruses expressing fluorescent proteins

by using an optimized reverse genetics system. J. Virol. 92:e00588-18.

27. Kumagai, A., Ando, R., Miyatake, H., Greimel, P., Kobayashi, T., Hirabayashi, Y.,

Shimogori, T., Miyawaki, A. 2013. A bilirubin-inducible fluorescent protein from eel

muscle. Cell 153:1602-1611.

28. Kyrieleis, O.J., Chang, J., de la Peña, M., Shuman, S., Cusack, S. 2014. Crystal structure of

vaccinia virus mRNA capping enzyme provides insights into the mechanism and evolution of

the capping apparatus. Structure 22:452-465.

29. Langland, J.O., Pettiford, S., Jiang, B., Jacobs, B.L. 1994. Products of the porcine group C

rotavirus NSP3 gene bind specifically to double-stranded RNA and inhibit activation of the

interferon induced protein kinase PKR. J. Virol. 68:3821-3829.

30. Martínez-Álvarez, L., Piña-Vázquez, C., Zarco, W., Padilla-Noriega, L. 2013. The shift from

low to high non-structural protein 1 expression in rotavirus-infected MA-104 cells. Mem.

Inst. Oswaldo Cruz. 108:421-428.

31. Mohanty, S.K., Donnelly, B., Dupree, P., Lobeck, I., Mowery, S., Meller, J., McNeal, M.,

Tiao, G. 2017. A point mutation in the rhesus rotavirus VP4 protein generated through a

rotavirus reverse genetics system attenuates biliary atresia in the murine model. J. Virol. 91

pii: e00510-17.

32. Montero, H., Arias, C.F., Lopez, S. 2006. Rotavirus nonstructural protein NSP3 is not

required for viral protein synthesis. J. Virol. 80:9031-9038.

33. Navarro, A., Trask, S.D., Patton, J.T. 2013. Generation of genetically stable recombinant

rotaviruses containing novel genome rearrangements and heterologous sequences by reverse

genetics. J. Virol. 87:6211-6220.

34. Patton, J.T., Taraporewala, Z., Chen, D., Chizhikov, V., Jones, M., Elhelu, A., Collins, M.,

Kearney, K., Wagner, M., Hoshino, Y., Gouvea, V. 2001. Effect of intragenic rearrangement

and changes in the 3' consensus sequence on NSP1 expression and rotavirus replication. J.

Virol. 75:2076-2086.

35. Rodriguez, E.A., Campbell, R.E., Lin, J.Y., Lin, M.Z., Miyawaki, A., Palmer, A.E., Shu, X.,

Zhang, J., Tsien, R.Y. 2017. The growing and glowing toolbox of fluorescent and

photoactive proteins. Trends Biochem Sci. 42:111-129.

36. Salsman, J., Top, D., Boutilier, J., Duncan, R. 2005. Extensive syncytium formation

mediated by the reovirus FAST proteins triggers apoptosis-induced membrane instability. J.

Virol. 79:8090-8100.

37. Settembre, E.C., Chen, J.Z., Dormitzer, P.R., Grigorieff, N., Harrison, S.C. 2011. Atomic

model of an infectious rotavirus particle. EMBO J. 30:408–416.

38. Shen, S., Burke, B., Desselberger, U. 1994. Rearrangement of the VP6 gene of a group A

rotavirus in combination with a point mutation affecting trimer stability. J. Virol. 68:1682-

1688.

39. Tortorici, M.A., Broering, T.J., Nibert, M.L., Patton, J.T. 2003. Template recognition and

formation of initiation complexes by the replicase of a segmented double-stranded RNA

virus. J. Biol. Chem. 278:32673-32682.

40. Trask, S.D, Taraporewala, Z.F., Boehme, K.W., Dermody, T.S., Patton, J.T. 2010. Dual

selection mechanisms drive efficient single-gene reverse genetics for rotavirus. Proc. Natl.

Acad. Sci. USA 107:18652-18657.

41. Trask, S.D., McDonald, S.M., Patton, J.T. 2012. Structural insights into the coupling of

virion assembly and rotavirus replication. Nat. Rev. Microbiol. 10:165–177

**Figure 1.** Wildtype and modified NSP3 proteins encoded by pT7 plasmids and rotaviruses. (A)

Organization of pT7 plasmids expressing wild type NSP3 and NSP3-FL/UnaG (+)RNAs,

indicating locations of T7 promoter (prm) and Hepatitis delta virus (HDV) self-cleaving

ribozyme (Rz). Nucleotide positions are labeled. (B) Products of recombinant RVAs expressing

wild type NSP3 and NSP3-FL-UnaG and RVC (Bristol strain) expressing NSP3-2A-dsRBP,

including approximate locations of functional domains in NSP3 (Deo et al., 2002; Groft and

Burley, 2002). Red arrow indicates the position of the stop-restart cleavage site in the 2A-like

element in the RVC NSP3-2A-dsRBP ORF. Amino acid positions are labeled. FL represents a 3x

Flag tag.

**Figure 2.** Recovery of recombinant RVA with a modified segment 7 dsRNA that expresses the

fused NSP3-FL-UnaG protein. (A) Profiles of the eleven genomic dsRNAs recovered from

rSA11 viruses resolved by PAGE. Red arrow notes the position of the segment 7 dsRNA. (B)

Western blot analysis of proteins present at 8 h p.i. in MA104 cells infected with recombinant

viruses.

**Figure 3**. Dimerization of NSP3-FL-UnaG. MA104 cells were mock infected, or infected with

rSA11/wt or rSA11/NSP3-FL-UnaG and incubated until 8 h p.i., when cells were harvested. Cell

lysates were mixed with sample buffer containing sodium dodecyl sulfate and b-

mercaptoethanol, incubated for 10 min either at 25 or 95$_\circ$C, resolved by electrophoresis on a

Novex 8-16% polyacrylamide gel, and the blotted onto a nitrocellulose membrane. Blots were

probed with guinea pig polyclonal anti-NSP3 or anti-VP6 antibodies, or with a mouse anti-

PCNA monoclonal antibody. Primary antibodies were detected using horseradish peroxidase-

conjugated secondary antibodies. Sizes (kD) of protein markers (M) are indicated. NSP3 band

intensities were determined by Image J analysis and were normalized to VP6 band intensities.

**Figure 4.** Comparison of plaques formed by wildtype rSA11 and mutant rSA11 strains

containing NSP3-FL-UnaG or NSP1-FL-UnaG sequences. (A) Plaques were generated on

MA104 monolayers and detected at 7 days p.i. by crystal violet staining (Arnold et al., 2009).

(B) Sizes of 40 randomly selected plaques from six independent plaque assays were measured

and the means determined. Mean values and 95% confidence intervals are plotted (black lines).

Significance values were calculated using an unpaired Student's t-test (GraphPad Prism, version

8). **** , $P < 0.0001$

**Figure 5.** Genetic stability of rSA11/NSP3-FL-UnaG. Virus was serially passaged 10-times in

MA104 cells using 1:1000 dilutions of infected cell lysate as inoculum. RNA was recovered

from infected cell lysates using Trizol, treated 30 min with RNase T1 at room temperature, and

resolved by electrophoresis on a 8% polyacrylamide gel. RNA gel was detected by staining gel

with ethidium bromide. Red arrow denotes the position of the segment 7 (NSP3) RNA.

**Figure 6.** Comparison of UnaG expression by rSA11/NSP3-FL-UnaG versus rSA11/NSP1-FL-

UnaG. (A) Images taken at 7.5 h p.i. with an epifluorescence Nikon TiE inverted microscope

with a 20x Plan Apo (NA 0.75) differential interference contrast (DIC) objective. (B)

Quantification of percent change in UnaG fluorescence (ΔF) in MA104 cells infected with

rSA11/ NSP1-FL-UnaG or rSA11/NSP3-FL-UnaG. Single cell analysis (N=270 cells each) was

performed to determine the percent ΔF from baseline. Data shown as mean ± standard deviation.

****p <0.0001

**Figure 7.** pT7 plasmids and rotaviruses expressing NSP1. (A) Organization of pT7 plasmids

expressing wild type NSP1 and NSP1-FL-UnaG (+)RNAs, indicating locations of T7 promoter

and Hepatitis delta virus self-cleaving ribozymes. The red arrow notes the position of the stop

codon in the NSP1-FL-UnaG ORF. (B) Products of recombinant RVAs expressing wild type

NSP1 and NSP1-FL-UnaG. (C) Profiles of the eleven genomic dsRNAs recovered from rSA11

viruses resolved by PAGE. Red arrow notes the position of the segment 5 dsRNA. (D) Western

blot analysis of proteins present at 8 h p.i. in MA104 cells infected with recombinant viruses.

**Figure 8.** Conservation of a predicted stable stem-loop structure formed by the 3'-UTR sequence

of rSA11/wt and rSA11/NSP3-FL-UnaG. Secondary structures associated with minimum free

energy were calculated for segment 7 (+)RNAs using *RNAfold* {http://rna.tbi.univie.ac.at/} and

color coded to indicate base-pairing probability (Hokacker, 2003; Hofacker et al., 1994).

Portions of the secondary structures are shown that include the 5' and 3' ends of the (+)RNAs

(labeled) and the conserved 3' stem-loops (3'SL) (boxed). Also labeled are the start and stop

codons (green and red arrowheads, respectively) of both the NSP3 and NSP3-FL-UnaG ORFs.

**Movie S1.** Expression of fluorescent signal by rSA11/NSP3-FL-UnaG and rSA11/NSP1-FL-

UnaG over the course of infection. Images were collected at 6 min intervals from 0 to 16 h p.i.

using an epifluorescence Nikon TiE inverted microscope with a 20x Plan Apo (NA 0.75)

differential interference contrast (DIC) objective.

**Figure 1.** Wildtype and modified NSP3 proteins encoded by pT7 plasmids and rotaviruses. (A) Organization of pT7 plasmids expressing wild type NSP3 and NSP3-FL/UnaG (+)RNAs, indicating locations of T7 promoter (prm) and Hepatitis delta virus (HDV) self-cleaving ribozyme (Rz). Nucleotide positions are labeled. (B) Products of recombinant RVAs expressing wild type NSP3 and NSP3-FL-UnaG and RVC (Bristol strain) expressing NSP3-2A-dsRBP, including approximate locations of functional domains in NSP3 (Deo et al., 2002; Groft and Burley, 2002). Red arrow indicates the position of the stop-restart cleavage site in the 2A-like element in the RVC NSP3-2A-dsRBP ORF. Amino acid positions are labeled. FL represents a 3x Flag tag.

**Figure 2.** Recovery of recombinant RVA with a modified segment 7 dsRNA that expresses the fused NSP3-FL-UnaG protein. (A) Profiles of the eleven genomic dsRNAs recovered from rSA11 viruses resolved by PAGE. Red arrow notes the position of the segment 7 dsRNA. (B) Western blot analysis of proteins present at 8 h p.i. in MA104 cells infected with recombinant viruses.

**Figure 3**. Dimerization of NSP3-FL-UnaG. MA104 cells were mock infected, or infected with rSA11/wt or rSA11/NSP3-FL-UnaG and incubated until 8 h p.i., when cells were harvested. Cell lysates were mixed with sample buffer containing sodium dodecyl sulfate and β-mercaptoethanol, incubated for 10 min either at 25 or 95°C, resolved by electrophoresis on a Novex 8-16% polyacrylamide gel, and the blotted onto a nitrocellulose membrane. Blots were probed with guinea pig polyclonal anti-NSP3 or anti-VP6 antibodies, or with a mouse anti-PCNA monoclonal antibody. Primary antibodies were detected using horseradish peroxidase-conjugated secondary antibodies. Sizes (kD) of protein markers (M) are indicated. NSP3 band intensities were determined by Image J analysis and were normalized to VP6 band intensities.

**Figure 4.** Comparison of plaques formed by wildtype rSA11 and mutant rSA11 strains containing NSP3-FL-UnaG or NSP1-FL-UnaG sequences. (A) Plaques were generated on MA104 monolayers and detected at 6 days p.i. by crystal violet staining (Arnold et al., 2009). (B) Sizes of 40 randomly selected plaques from six independent plaque assays were measured and the means determined. Mean values and 95% confidence intervals are plotted (black lines). Significance values were calculated using an unpaired Student's t-test (GraphPad Prism, version 8). **** , $P < 0.0001$

**Figure 5**. Genetic stability of rSA11/NSP3-FL-UnaG. Virus was serially passaged 10-times in MA104 cells using 1:1000 dilutions of infected cell lysate as inoculum. RNA was recovered from infected cell lysates using Trizol, treated 30 min with RNase T1 at room temperature, and resolved by electrophoresis on a 8% polyacrylamide gel. RNA gel was detected by staining gel with ethidium bromide. Red arrow denotes the position of the segment 7 (NSP3) RNA.

A

B

**Figure 6**. Comparison of UnaG expression by rSA11/NSP3-FL-UnaG versus
rSA11/NSP1-FL-UnaG. (A) Images taken at 7.5 h p.i. with an epifluorescence Nikon
TiE inverted microscope with a 20x Plan Apo (NA 0.75) differential interference
contrast (DIC) objective. (B) Quantification of percent change in UnaG fluorescence
($\Delta$F) in MA104 cells infected with rSA11/ NSP1-FL-UnaG or rSA11/NSP3-FL-UnaG.
Single cell analysis (N=270 cells each) was performed to determine the percent $\Delta$F
from baseline. Data shown as mean ± standard deviation. ****p <0.0001

**Figure 7.** pT7 plasmids and rotaviruses expressing NSP1. (A) Organization of pT7 plasmids expressing wild type NSP1 and NSP1-FL-UnaG (+)RNAs, indicating locations of T7 promoter and Hepatitis delta virus self-cleaving ribozymes. The red arrow notes the position of the stop codon in the NSP1-FL-UnaG ORF. (B) Products of recombinant RVAs expressing wild type NSP1 and NSP1-FL-UnaG. (C) Profiles of the eleven genomic dsRNAs recovered from rSA11 viruses resolved by PAGE. Red arrow notes the position of the segment 5 dsRNA. (D) Western blot analysis of proteins present at 8 h p.i. in MA104 cells infected with recombinant viruses.

**Figure 8**. Conservation of a predicted stable stem-loop structure formed by the 3'-UTR sequence of rSA11/wt and rSA11/NSP3-FL-UnaG. Secondary structures associated with minimum free energy were calculated for segment 7 (+)RNAs using *RNAfold* {http://rna.tbi.univie.ac.at/} and color coded to indicate base-pairing probability (Hokacker, 2003; Hofacker et al., 1994). Portions of the secondary structures are shown that include the 5' and 3' ends of the (+)RNAs (labeled) and the conserved 3' stem-loops (3'SL) (boxed). Also labeled are the start and stop codons (green and red arrowheads, respectively) of both the NSP3 and NSP3-FL-UnaG ORFs.