1 **The evolution of biotic and abiotic realized niches within freshwater**
2 ***Synechococcus***
3
4

Authors: Nicolas Tromas[1#], Mathieu Castelli[2#], Zofia E. Taranu[3], Juliana S. M. Pimentel[4], Daniel

A. Pereira[4], Romane Marcoz[1], Alessandra Giani[4] and B. Jesse Shapiro[1*]

1- Département de sciences biologiques, Université de Montréal, 90 Vincent-d'Indy, Montréal,

QC, Canada, Montréal, QC H2V 2S9, Canada

2- mathieu.castelli@gmail.com

3- Department of Biology, University of Ottawa, Gendron Hall, 30 Marie Curie, Ottawa

4- Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil
13
14
15 *Corresponding authors: B. Jesse Shapiro. Phone: 514-343-6033. E-mail:
16 jesse.shapiro@umontreal.ca; Nicolas Tromas. Phone 514-343-3188. E-mail:
17 nicolas.tromas@umontreal.ca.
18
19 # These authors contributed equally to this work
20
21

22 **Originality-Significance Statement**
23

We address a fundamental question in ecology and evolution: how do niche preferences change

over evolutionary time? Using time-series analysis of 16S rRNA gene amplicon sequencing data,

we develop a new approach to highlight the importance of biotic factors in defining realized

niches, and show how niche preferences change "clock-like" within the genus *Synechococcus.*

Ours is also one of few studies on the ecology of freshwater *Synechococcus,* adding significantly

to our knowledge about this abundant and widespread lineage of Cyanobacteria.

Summary

Understanding how ecological traits have changed over evolutionary time is a fundamental question in biology. When closely-related organisms share similar environmental preferences, "habitat filtering" is expected to determine how communities are assembled. Yet in practice, it is challenging to assess the impact of habitat filtering, due to our inability to measure all relevant abiotic variables and distinguish the impact of biotic versus abiotic factors. Here we explored the co-occurrence patterns of freshwater cyanobacteria at the sub-genus level to investigate whether closely-related taxa share similar niches, and to what extent these niches were defined by abiotic or biotic variables. We used deep 16S rRNA gene amplicon sequencing and measured several environmental parameters in water samples collected over time and space in Furnas Reservoir, Brazil. We found that closely-related *Synechococcus* did not have similar preferences for abiotic niche dimensions. However, closely-related *Synechococcus* did tend to co-occur with one another, and also with similar surrounding microbial communities. These results suggest that biotic factors may be stronger niche determinants than abiotic factors. Alternatively, cryptic abiotic drivers may determine niche and community structure, but biotic factors provide the most informative measure of niche similarity.

Introduction

A bacterial community is a group of potentially interacting organisms that coexist at a particular place and time (Magurran, 2003). Environmental selective pressures are a strong force shaping microbial community assembly (Martiny et al, 2015). We know, for example, that certain abiotic factors explain a large portion of the variation in microbial community composition (e.g. the effect of pH on soil bacterial communities; Fierer and Jackson, 2006). Therefore, associations between microbial traits - generally defined as a phenotypic response to a specific environmental condition - and abiotic niches could help explain community assembly rules (Green et al., 2008; Burke et al., 2011). However, trait-based approaches to understand communities may be challenging, as important abiotic variables may go unmeasured. Even less is known about biotic interactions, despite their importance in determining community composition, diversity, and dynamics (Needham et al., 2016).

Abiotic factors are generally thought to determine an organism's fundamental niche (where it is theoretically capable of living), whereas biotic factors determine its realized niche (where it actually lives in nature; Hutchinson, 1957). Species (or taxonomic) co-occurrence networks are often used to infer niche similarity among organisms that tend to co-occur in nature over space and time, and microbial co-occurrence networks can easily be constructed from 16S rRNA gene amplicon sequencing surveys (Friedman and Alm, 2012; Röttjers and Faust, 2018; Tromas et al., 2018). However, co-occurrence can be driven by both abiotic and biotic factors, which are hard to disentangle in practice (Kraft et al., 2015). Regardless, organisms sharing a similar niche are expected to be associated with similar surrounding communities (Cohan and Koeppel, 2008; Faust et al.2012; Pascual-García et al., 2014).

A fundamental question spanning ecology and evolution is how ecological traits change over evolutionary time. For example, some traits (such as bacteriophage host range) evolve rapidly at the tips of a phylogenetic tree, whereas other traits (such as salinity preference) are deeply conserved (Martiny *et al.* 2015). When closely related organisms share similar ecological preferences, so-called "habitat filtering" or "environmental" filtering is expected to result in phylogenetic clustering, meaning that a community tends to contain more closely related organisms than expected by a random draw from the phylogeny (Webb et al., 2002; Horner-Devine & Bohannan, 2006; Martiny et al., 2015). In contrast, if close relatives evolve different traits to avoid competitive exclusion, this will result in phylogenetic overdispersion (*i.e.* a community composed of more distant relatives than expected by chance). However, the relative importance of these two processes in shaping microbial communities is still widely debated, and difficult to distinguish (Koeppel and Wu, 2014; Cadotte and Tucker, 2017). We have previously shown that within the cyanobacterial genus *Dolichospermum* (Tromas et al., 2018), the relationships between phylogenetic distance and ecological similarity varies by trait, suggesting that it might be necessary to analyze each niche dimension or trait separately (Martiny *et al*., 2015).

In this study, we explored the co-occurrence patterns of freshwater cyanobacteria at the sub-genus level, to investigate if closely related taxa have more similar niches, and to what extent these niches can be quantified by abiotic or biotic variables. We focused on *Synechococcus*, the most abundant cyanobacterial genus in Furnas Reservoir (Brazil) at the time of sampling (2006-2008). *Synechococcus* is among the most abundant organisms living in oceans and lakes (Stockner et al. 2000; Scanlan, 2003). The phylogenetic coherence of the genus has been

questioned by a recent study showing it to be polyphyletic (Coutinho et al 2016). *Synechococcus*

is physiologically highly plastic, ubiquitous, and able to acclimate to different environmental

conditions (Callieri 1996; Vörös et al. 1998; Callieri et al., 2011). Previous studies have shown

that different *Synechococcus* strains could co-exist in the same site but respond differently to

environmental changes, suggesting niche partitioning (Ferris et al., 2003; Allewalt et al., 2006;

Becker et al., 2007; Becraft et al., 2011; Callieri et al. 2012). Recently, Zheng et al. (2018),

observed a geographical pattern of heterotrophic bacteria associated with different marine

Synechococcus strains, indicating that strains living in the same area tend to be associated with

similar communities. It remains, however, unclear whether the *Synechococcus* genotype or the

environment are the main drivers of *Synechococcus* interactions with the surrounding microbial

community.

Using deep 16S rRNA gene amplicon sequencing of 86 water samples collected in time series

across nine locations in the Furnas Reservoir, we tracked genetic diversity within the

Synechococcus genus, along with the surrounding microbial community, and measured several

abiotic variables. We found that that closely-related *Synechococcus* tended to co-occur with one

another and also with similar surrounding microbial communities. Such phylogenetic clustering

indicates that overall realized niche similarity tends to evolve "clock-like" in the *Synechococcus*

lineage. However, closely-related *Synechococcus* did not have similar abiotic niche preferences

(with the exception of total phosphorus). These results suggest that biotic factors may be stronger

niche determinants than abiotic factors. Alternatively, cryptic abiotic drivers may determine niche

and community structure, but biotic factors provide the most informative measure of niche

similarity.

Materials and methods

Sampling, environmental data measurements

A total of 90 water samples were collected from September 2006 to April 2008 at nine stations in Furnas Reservoir (Minas Gerais, Brazil). Furnas is a large reservoir (1440 km^2), located in southeastern Brazil (20o40'S; 46o19'W) and formed by the damming of two main rivers (Rio Grande and Rio Sapucaí), which divide the reservoir in two separated branches (Figure S1). Sampling stations S1, S4, S6, S9 are from a relatively pristine branch of the reservoir, whereas S12, S14, S18 and S20 are impacted by human activities. Temperature profiles were measured in the water column by aid of a multi-parameter probe (556 YSI, USA). Water samples were collected from the euphotic zone (determined by Secchi disc depth) by a Van Dorn sampler. Samples were stored in bottles that had been acid-washed and rinsed with deionized water. A portion of each water sample was immediately filtered through glass fiber filters (Whatmann GF/F, 0.7 μm pore size). The exact filtered volumes were recorded and filters were kept frozen until further analyses (chlorophyll, DNA and microcystin). For dissolved nutrient analyses, 200 mL of filtered water samples were stored at −20 °C. For total phosphorus, samples were frozen with no previous filtration. Nutrient analyses were performed using spectrophotometric methods according to APHA (2005). All nutrient analyses (Nitrate, Nitrite and total phosphorus (TP) were performed on three replicates. For phytoplankton analyses, samples were Lugol-preserved for subsequent cyanobacteria identification and quantification.

DNA extraction, purification and sequencing

DNA was extracted from frozen filters according to Kurmayer et al. (2003), with few modifications. Briefly, filters were treated with a sucrose buffer (25% w/v sucrose, 50 Mm Tris-HCl, 100 Mm EDTA, pH 8) on ice for 2 h and with addition of lysozyme (5mg/mL, 1h, 37°C).

Proteinase K (100 μg/mL) in sodium dodecyl sulfate (2% v/v) was added and filters were

incubated overnight at 55°C. A phenol:chloroform:isoamyl alcohol solution (25:24:1, v/v) was

used for protein precipitation and DNA isolation. The DNA was cleaned in 100% ethanol and

pellets rinsed with 70% ethanol. The DNA was resuspended in TE (10 mM Tris- HCl, pH 8, and

1 mM EDTA). The DNA extract was quantified by a spectrophotometer, at 260 nm and 280 nm,

and its quality checked in 1% (w/v) agarose gel, stained with ethidium bromide.

Sequence analysis

We followed the same protocol described in Tromas et al., (2017) to generate a library of

V4 region amplicons. We performed one sequencing run using MiSeq reagent Kit V2 (Illumina,

San Diego, CA, USA) on a MiSeq instrument (Illumina). A total of 4,476,747 sequences of the

16S rRNA gene V4 region were obtained from 90 lake samples, two negative controls, and two

mock community samples. We obtained a median of 37,682 sequences per sample. Using a

similar approach as described in Tromas *et al*., (2018), we processed the sequences with

SmileTrain (https://github.com/almlab/SmileTrain/wiki; Preheim et al., 2013) for read quality

filtering, primer removal, chimera filtering, and merging using USEARCH (version 7.0.1090,

http://www.drive5.com/usearch/, default parameter) (Edgar, 2010), Mothur (version 1.33.3)

(Schloss *et al*., 2009), and Biopython (version 2.7). Minimum Entropy Decomposition (MED)

was then applied to the filtered and merged reads to partition sequence reads into MED nodes

(Eren *et al*., 2015). MED was performed using the following parameters: −M noise filter set to

1000, resulting in ~17.5% of reads filtered and 466 MED nodes representing the whole bacterial

community. Samples with less than 1000 reads were removed, yielding a final dataset of 86

reservoir samples. Finally, we assigned taxonomy to MED nodes using the assign_taxonomy.py

QIIME script (default parameters), and a combination of GreenGenes and a freshwater-specific

database (Freshwater database 2016 August 18 release; Newton *et al.*, 2011), using TaxAss

(https://github.com/McMahonLab/TaxAss, installation date: September 13[th] 2016; Rohwer *et al.*,

2017).

Spatio-temporal analysis

We used multivariate regression tree analyses (Breiman *et al.* 1984; De'ath 2002) to

investigate if spatio-temporal variables could explain genetic variation within *Synechococcus*. We

used two different temporal predictors: year and month and one spatial predictor: station. The

analysis was performed using the function *mvpart*() and *rpart.pca*() from the R mvpart package

(Therneau and Atkinson, 1997; De'ath, 2007). Prior to analysis, the *Synechococcus* MED nodes

table was Hellinger transformed to downweight the effect of double-zeros (Rao, 1995).

Genetic distance between Synechococcus nodes

We measured the genetic distance between *Synechococcus* MED nodes using the software

MEGA (Kumar et al., 2016; version 7.0.18) with the p-distance (the proportion of nucleotide

sites at which two sequences differ), calculated by dividing the number of sites with nucleotide

differences by the total number of sites compared (excluding sites with gaps).

Co-response to abiotic factors

As described in Tromas *et al.* (2018), we used a Latent Variable Model (LVM)

framework (boral package in R; Hui 2015, Warton *et al.* 2015) to explore how *Synechococcus*

nodes co-responded to abiotic gradients and used these co-responses as indicators of niche

similarity. That is, for each abiotic factor, we ran separate LVMs, regressing the bacterial

community as a function (both linear and non-linear) of the given factor. This component of the

LVM thus defined the taxon-specific environmental responses. To then identify remaining

patterns of co-occurrence after accounting for all measured environmental variables, we fit a

global LVM which included all abiotic factors and two latent variables (sensu Letten *et al.* 2015

and Warton *et al.* 2015). To visualize patterns of co-occurrence arising from the different

environmental factors, we calculated two types of correlation matrices. The first, a co-response

correlation matrix, was constructed by calculating, for any two nodes, the correlation between

their fitted values. This correlation matrix thus represented the correlation between nodes that can

be attributed to a shared or diverging environmental responses. A significant positive correlations

between the fitted response of any two taxa to an environmental variable represented a co-

response (i.e., as one taxa increases in response to an environmental variable, the other likewise

increases), whereas a significant negative correlations between the fitted response of two taxa

represented some degree of niche separation (i.e., as one taxa increases in response to an

environmental variable, the other decreases).

To account for the correlation between nodes that may be attributable to biotic processes or

missing environmental covariates, a second type of correlation matrix, a residual correlation

matrix, was calculated using the latent variable coefficients of the global LVM. Since the latent

variables are the output of an ordination of the residuals, we expected weak residual correlations

induced by the latent variables if the environmental variables were the dominant force structuring

patterns of species co-occurrence (*i.e. t*he model explained most of the variance, with little left

unexplained in the residuals). Conversely, if any unmeasured environmental factors and/or biotic

processes are equally, or more, important than measured environmental factors, we expected

strong correlations based on the latent variables (*i.e. t*he model was poor and much of the

observed variability remained in the residuals).

To test how *Synechococcus* niche similarity varied with genetic distance, we examined the

relationship between the co-response of *Synechococcus* taxa to environmental parameters and

their genetic distance (i.e., plotting the correlation coefficient of the LVM co-responses vs.

genetic distances) (R_script1). However, given the large number of environmental factors driving

phytoplankton community dynamics (Hutchinson, 1961), we expected some degree of co-

limitation whereby the response of a taxon to an environmental variable (and consequently the

degree of co-response among any two taxa) would be limited by other environmental variables.

In such cases, the taxon's response and rate of change would have an upper limit (set by all

measured environmental factors) but may not reach this limit if other, unmeasured factors are co-

limiting (Cade and Noon 2003). As an increasing number of unmeasured factors become limiting

at some sample location, or time point, the relationship between the response and the measured

factor becomes increasingly heterogeneous or wedge shaped. When such heterogeneous

variances are observed (wedge shape biplot), it suggests that there is not a single slope coefficient

that characterizes the relationship, and that focusing solely the coefficient fit by ordinary least

squares regression (mean response) may underestimate the true rate of change. Thus, to

accurately examine co-limitations among measured and unmeasured factors, we applied quantile

regressions using 'quantreg' R package (Koenker, 2015) (R_script1), which, instead of fitting a

regression to the mean response (ordinary least squares regression), fits regression curves to other

quantiles of the response variable (Cade and Noon 2003).

240
Co-occurrence with biotic factors

A caveat of the above LVM-quantile regressions is that although it identifies whether

some relationships are limited by unmeasured abiotic and/or biotic factors, it could not tease apart

the relative importance of each type of factor (everything is instead lumped as an unknown). In

order to determine the relative contribution of biotic processes, we thus quantified node co-

occurrence patterns among *Synechococcus* and the remaining community. In particular, we

explored the relationship between *Synechococcus* nodes and the surrounding community by

measuring: (1) co-occurrences between *Synechococcus* and other taxa, and (2) paired differences

in co-occurrence among *Synechococcus* nodes and a specific taxon. For the former, we calculated

co-occurrences among taxa using SparCC (Friedman and Alm, 2012), including 20 iterations to

estimate the median correlation of each pair of MED nodes, 500 bootstraps to assess the

statistical significance and centered log ratio (CLR) transform to correct for compositionality.

Correlations were then filtered using a false discovery rate threshold ($Q < 0.05$). For the latter, we

further calculated the absolute difference of Sparcc correlation (*r*) between a *Synechococcus* node

(X_i) and a specific taxon T, such that $|\Delta r| = |\text{Corr}(X_1, T) - \text{Corr}(X_2, T)|$ where Corr is defined

here as the SparCC correlation between X_i and T.

For each non-*Synechococcus* taxon, we then estimated the relationship between $|\Delta r|$ and

the distance between the given *Synechococcus* nodes (X_1 and X_2). A positive correlation would

indicate that closely related *Synechococcus* nodes have a lower $|\Delta r|$ than more distant nodes, *i.e.*

closely related nodes would have more similar correlations with potentially interacting

community members. To reduce potential bias or noise due to a small sample size of

Synechococcus pairs, we selected all non-*Synechococcus* nodes that were significantly co-

occurrent with at least seven different *Synechococcus*. This cutoff was chosen to ensure at least

20 unique pairs of *Synechococcus* for each non-*Synechococcus* taxon. A total of 373 non-

Synechococcus taxa were selected using this cutoff, and the correlation between $|\Delta r|$ and genetic

distance was non-significant (Spearman correlation, P<0.05) for 120 of them. Finally, we

performed a permutation test to quantify any bias in the method due to data structure. For each of

the significant non-*Synechococcus* taxa, we estimated the rate of false positive correlations by

sampling values of *Synechococcus* genetic distance with replacement, while randomizing the

association between genetic distance and |Δ*r*|. We performed 1000 permutations for each

significant non-*Synechococcus* taxon using the script R_script2. We then recorded the proportion

(*p*) of permutations yielding a larger correlation than observed, after adding a pseudocount of 1 to

both the numerator (the number of permutations yielding a larger correlation than observed) and

the denominator (the total number of permutations).

275
276   ***Phylogenetic analysis***
277
We first verified that *Synechococcus* group was monophyletic by building a phylogenetic

tree with FastTree (version 2.1.8, Price et al., 2009), using all MED node sequences aligned with

MAFFT (v7.154b; Katoh and Standley, 2013). We then specifically aligned *Synechococcus* MED

node sequences using muscle (in Mega, version 7.0.26; Kumar et al., 2016*)* and built

phylogenetic trees using PhyML (version 3.0; Guindon et al., 2010). We used the ALDEx2 R

package (version: 1.5.0; Fernandes *et al*., 2014) and the aldex() function to identify MED nodes

associated with the pristine or human-impacted branches of Furnas Reservoir. To do so, we used

Welsh's t-test and 128 Monte Carlo samples. ALDEx2 uses the CLR transformation to avoid

compositional effects. A Q-value below 0.05 after Benjamini-Hochberg correction was taken as

evidence for association. We also used a second association method, DESeq2 implemented in the

"MicrobiomeAnalyst" web-based tool (Dhariwal et al., 2017).

289
290   ***Statistical analysis***
291
292   All statistical tests were performed in R.

293
294

Results

Abiotic factors shape genetic variation within Synechococcus

Furnas Reservoir is divided into two branches: one, which is heavily human-impacted (Figure S1,

sampling stations 12-20) and the other, less impacted (stations 1-9). *Synechococcus* dominated

the bacterial community in both branches during the sampling period (2006-2008) (Figure S2)

and formed a monophyletic group (Figure S3, S4). We first asked whether particular

Synechococcus strains (which we used interchangeably with MED nodes) were associated with

different branches of the reservoir, different time periods, or different abiotic factors. We applied

a multivariate regression tree analysis (Breiman et al. 1984; De'ath 2002) to test whether the

composition of *Synechococcus* strains changed over time and space. We found that the year and

month of sampling did not explain much variance (R^2=0.02 and R^2=0.03, respectively), nor did

the station (R^2=0.06) or branch of the reservoir (R^2=0.02). In contrast, Chl *a*, Nitrite, TP, Depth,

TC, PTC, and water temperature all explained a significant proportion of the variance (Figure S5;

40% of variance explained), and depth, temperature and nitrite alone explained most of the

variability (Figure S6; 16% unique, and an additional 6% co-explained with Chl *a*).

Closely related Synechoccus do not share similar preferences for measured abiotic niches

Having established that abiotic factors, but not temporal or spatial variation, explain a significant

amount of variation in the relative abundances of *Synechococcus* strains, we asked whether

closely related strains tended to share similar abiotic niches. Here we defined abiotic niche

similarity by estimating the co-responses of pairs of *Synechococcus* strains to each abiotic factor

using LVM-quantile regression (Methods). For most abiotic factors, we did not observe any

significant relationship between co-responses and genetic distance, with the exception of total

phosphorus for which the relationship was negative (Figure 1, S7). Similarly, the degree of

abiotic niche separation (*i.e.* significant negative correlations between the fitted response of any

two *Synechococcus* to an abiotic factor) was largely independent of genetic distance (Figure S8,

Figure S9).

Figure 1. Relationship between *Synechococcus* niche similarity and genetic distance. Niche

similarity is defined as a positive LVM co-response (y-axis). Shown are the scatterplots of LVM

co-responses versus each variables superimposed with the lines for 5th, 10th, 15th, 25th, 35th, 45th,

55th, 65th, 75th, 85th and 90th quantile regression fits (grey lines, for each co-response quantile),

the median fit (50th quantile; blue line), and the least squares estimate or mean fit (dashed red

line).

To take into account the correlation between nodes that may be attributable to biotic processes or

missing environmental covariates, we used the latent variables of the global LVM to examine

how the correlation among node residuals varied with genetic distance. For the positive

correlations (Figure 2A, B), the correlation coefficients tended to decrease with genetic distance.

This result suggests a negative relationship between the residuals (*i.e.* unmeasured abiotic

variables and/or biotic processes) and phylogenetic distance. For the negative residual

correlations (Figure 2C, D), there were no apparent relationships with the genetic distance.

Therefore, there remain significant co-responses that were not explained by the abiotic factors

measured in this study. Overall, most of the measured abiotic environmental variables were not

related to *Synechococcus* genetic distance. However, preferences for unmeasured biotic or abiotic

factors do tend to change with genetic distance, such that more closely-related *Synechococcus*

have similar realized niche preferences.

Figure 2. Relationship between *Synechococcus* co-responses (LVM residuals) and genetic distance. Positive co-responses are shown in panels **A** and **B**; negative co-responses in panels **C** and **D.** *(*A, C*)* Scatterplots of LVM co-responses versus LVM residuals superimposed with the lines for 5th, 10th, 15th, 25th, 35th, 45th, 55th, 65th, 75th, 85th and 90th quantile regression fits (grey lines), the median fit (50th quantile; blue line), and the least squares estimate or mean fit (dashed red line). (B, D) Panels show the quantile slope estimates (dark grey line) and corresponding confidence intervals (light grey bands) of the response models across all quantiles (arrows, with the median shown in blue). Significant quantile slopes occur when confidence intervals do not overlap zero.

Closely related Synechococcus share similar biotic interactions

The observed tendency for closely-related *Synechococcus* to share similar realized niches (Figure

2A) could be explained by shared preferences for either unmeasured abiotic factors or biotic

interactions. We used co-occurrence network analysis to determine the role of biotic interactions.

First, we asked if closely related *Synechococcus* tend to co-occur across samples. We observed a

negative relationship between node co-occurrence and pairwise genetic distance (Figure 3),

indicating that genetically similar *Synechococcus* nodes are indeed more likely to co-occur, and

thus to have similar realized niches. This pattern was significant [linear regression, $F_{(1,488)}$ =

108.6, $P < 0.001$, adjusted R^2 =0.18].

Figure 3. **Negative relationship between pairwise SparCC correlation coefficients and**

pairwise genetic distance between *Synechococcus* nodes. Only significant SparCC correlations

(Q < 0.05) were included. Pairwise genetic distance was computed as percent nucleotide identity

(p-distance). Blue diamonds represent the mean SparCC correlation for each distance. Boxplots

show the median (horizontal line), the 25th and 75th percentile (enclosed in box) and 95%

confidence intervals (whiskers).

We further investigated whether more closely-related *Synechococcus* nodes have more similar

associations with surrounding community members. To do so, we performed a pairwise analysis

to examine the relationship between *Synechococcus* genetic distance and $|\Delta r|$, a measure of the

similarity of *Synechococcus* co-occurrence with non-*Synechococcus* taxa (Methods). A total of

373 non-*Synechococcus* taxa were analysed and the correlation between $|\Delta r|$ and genetic distance

was non-significant for 120 of them. The remaining 253 significant non-*Synechococcus* have

consistently positive relationships between co-occurrence and genetic distance, *i.e.* closely related

Synechococcus nodes have a higher chance of co-occurring with similar non-*Synechococcus* taxa

(Figure 4). This result is robust to data structure, as determined by permuting the associations

between genetic distance and $|\Delta r|$ (Figure S10; Methods).

Figure 4. Closely related *Synechococcus* nodes co-occur with similar surrounding community. The histogram shows the distribution of Spearman correlations between the genetic distance of *Synechococcus* nodes and their association with similar surrounding communities. Spearman correlations were calculated between the genetic distance of *Synechococcus* pairs and the absolute difference of Sparcc correlation (r) of a *Synechococcus* node with a specific taxon T, such that $|\Delta r| = |\mathrm{Corr}(X_1, T) - \mathrm{Corr}(X_2, T)|$ where Corr is defined here as the SparCC correlation score of X_i (*Synechococcus* node) and T (non-*Synechococcus* node). We then estimated the correlation of $|\Delta r|$ and the genetic distance between nodes X_1 and X_2. The mean Spearman correlation estimate was 0.373 (red line) and 99.6% of correlations are positive.

Finally, we selected the non-*Synechococcus* taxa with the highest correlation scores between |Δr|

and *Synechococcus* distance genetic. Five taxa had correlation estimates higher than 0.6,

including three Proteobacteria, one Verrucomicrobia, and one Planctomycetes (Table S1).

Previous studies have shown that members of the *Comamonadaceae* family and *Spirobacillales*

order (members of Proteobacteria; Table S1) were associated with *Synechococcus* (Guedes et al.,

2018) or more broadly with algae (Fullbright et al., 2019). Moreover, the *Gallionella* genus was

previously observed colonizing filamentous algae (Mori et al., 2015).

Overall, these results show that the co-occurring surrounding community is, on average, more

similar for closely related *Synechococcus*. Members of this surrounding community could

interact directly with *Synechococcus,* or share similarity in their abiotic niche.

Discussion

In this study, we investigated *Synechococcus* niche preferences at relatively fine (sub-

genus) taxonomic resolution using a 16S gene sequence variants approach previously used to

study other bacterial lineages (Koeppel and Wu, 2014; Tromas et al., 2018). We found that

abiotic factors, such as reservoir depth and nitrite concentration, greatly affected the relative

abundance of *Synechococcus* strains (Figure S5, S6). However, the measured abiotic factors were

not related to genetic distance, indicating that these traits are probably loosely conserved and may

evolve on shorter time scales, being thus consistent with phylogenetic overdispersion. An

exception was total phosphorus preference, an abiotic niche dimension that was correlated with

genetic distance, consistent with phylogenetic clustering. Phosphorus use may therefore evolve

'"clock-like" along the *Synechococcus* phylogeny. This contrasts with the idea that organic

phosphate niches are fast-evolving and relatively uncorrelated with phylogeny, perhaps due to

horizontal transfer of relevant genes (Coleman & Chisholm 2010; Martiny et al., 2015).

Therefore, how phosphorus-related traits evolve along phylogenies likely depend on the lineage

considered (*e.g. Synechococcus* vs. *Prochlorococcus* or *Pelagibacter*) and evolutionary time

scales. We note that only 77 of the 780 *Synechococcus* strain pairs showed a significant co-

response to phosphorus (Figure 1), indicating that the effect is driven by a small number of

closely-related strains with similar co-responses, and cannot be generalized to all *Synechococcus*.

This result might be explained if *Synechococcus* represents a polyphyletic group (Coutinho et al.,

2016), composed of different clades with different phosphorus niches. However, in our sample,

Synechococcus is monophyletic (Figure S3, S4), excluding this explanation. Overall, our results

demonstrate how abiotic factors can shape community composition within a genus, but measured

abiotic niches generally do not evolve along the genus-level phylogeny.

Habitat filtering was originally defined to describe how abiotic factors select for genetically

similar (closely related) organisms (Cadotte and Tucker 2017). Yet in practice, it is difficult to

disentangle the contributions of biotic and abiotic factors, and thus to satisfactorily define habitat

filtering (Kraft et al., 2015). In this study, we therefore estimated the influence of both

unmeasured abiotic variables and biotic factors, and found a negative association with genetic

distance (Figure 2). This suggests that closely related *Synechococcus* tend to share similar

realized niches. As a result, closely related *Synechococcus* tend to co-occur (Figure 3), as

observed in studies using genomic (rather than 16S) similarity (Kamneva, 2017). This result is

consistent with habitat filtering, broadly defined to include both biotic and abiotic factors.

We suggest that biotic niches are the major drivers of habitat filtering and the prime determinants

of realized niches in *Synechococcus*. This is because closely-related *Synechococcus* tend to co-

occur with more similar surrounding communities (Figure 4), whereas they do not tend to share

abiotic niche preferences (Figure 1). Of course, it is possible that unmeasured abiotic factors are

driving the observed biotic effects. However, we measured abiotic factors such as temperature

and nutrients, which are important in structuring microbial communities, and none of these

factors are sufficient to account for realized niches or their distribution across the *Synechococcus*

phylogeny. Thus, as previously suggested (Cohan and Koeppel, 2008), biotic factors (quantified

here with 16S amplicon sequencing) provide more information about niches than abiotic factors,

which are difficult to measure comprehensively. For example, the presence of certain taxa could

indicate the importance of an unmeasured abiotic factor (*e.g* Acidophiles or Alkaliphiles as

indicators of pH). In this case, considering biotic factors allowed us to uncover a signal of habitat

filtering within *Synechococcus* that would have been obscured by only considering commonly

measured abiotic factors.

References

Allewalt, J.P., Bateson, M.M., Revsbech, N.P., Slack, K., and Ward, D.M. (2006) Effect of
Temperature and Light on Growth of and Photosynthesis by Synechococcus Isolates
Typical of Those Predominating in the Octopus Spring Microbial Mat Community of
Yellowstone National Park. *Appl Environ Microbiol* **72**: 544–550.

Becker, S., Richl, P., and Ernst, A. (2007) Seasonal and habitat-related distribution pattern of
Synechococcus genotypes in Lake Constance. *FEMS Microbiol Ecol* **62**: 64–77.

Becraft, E.D., Cohan, F.M., Kühl, M., Jensen, S.I., and Ward, D.M. (2011) Fine-Scale
Distribution Patterns of Synechococcus Ecological Diversity in Microbial Mats of
Mushroom Spring, Yellowstone National Park. *Appl Environ Microbiol* **77**: 7689–7697.

Breiman L, Friedman JH, Olshen RA, Stone CJ. (1984) Classification and Regression Trees.
Wadsworth International Group, Belmont, CA, USA.

Burke, C., Steinberg, P., Rusch, D., Kjelleberg, S., and Thomas, T. (2011) Bacterial community
assembly based on functional genes rather than species. *Proc Natl Acad Sci USA* **108**:
14288–14293.

Cade, B.S. and Noon, B.R. (2003) A Gentle Introduction to Quantile Regression for Ecologists.
Frontiers in Ecology and the Environment **1**: 412–420.

Cadotte, M.W. and Tucker, C.M. (2017) Should Environmental Filtering be Abandoned? *Trends
in Ecology & Evolution* **32**: 429–437.

Callieri, C. (1996) Extinction coefficient of red, green and blue light and its influence on
picocyanobacterial types in lakes at different trophic levels. *Memorie dell'Istituto Italiano
di Idrobiologia* 54: 135-142.

Callieri, C., Caravati, E., Corno, G., and Bertoni, R. (2012) Picocyanobacterial community
structure and space-time dynamics in the subalpine Lake Maggiore (N. Italy). *Journal of
Limnology* **71**: e9–e9.

Callieri, C., Lami, A., and Bertoni, R. (2011) Microcolony Formation by Single-Cell
Synechococcus Strains as a Fast Response to UV Radiation. *Appl Environ Microbiol* **77**:
7533–7540.

Cohan, F.M. and Koeppel, A.F. (2008) The origins of ecological diversity in prokaryotes. *Curr
Biol* **18**: R1024-1034.

Coleman, M. L., and Chisholm, S. W. (2010). Ecosystem-specific selection pressures revealed
through comparative population genomics. *Proc. Natl. Acad. Sci. U. S. A.* **107** : 18634–
18639.

Coutinho, F., Tschoeke, D.A., Thompson, F., and Thompson, C. (2016) Comparative genomics

506   of Synechococcus and proposal of the new genus Parasynechococcus. *PeerJ* **4**: e1522.
507
508   De'ath G. (2007) mvpart: Multivariate partitioning. R package version 1.6-2.
509
510   Dhariwal, A., Chong, J., Habib, S., King, I.L., Agellon, L.B., and Xia, J. (2017)
511   MicrobiomeAnalyst: a web-based tool for comprehensive statistical, visual and meta-
512   analysis of microbiome data. *Nucleic Acids Res* **45**: W180–W188.
513
514   Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
515   **26**: 2460–2461.
516
517   Eren, A.M., Morrison, H.G., Lescault, P.J., Reveillaud, J., Vineis, J.H., and Sogin, M.L. (2015)
518   Minimum entropy decomposition: unsupervised oligotyping for sensitive partitioning of
519   high-throughput marker gene sequences. *ISME J* **9**: 968–979.
520
521   Faust, K., Sathirapongsasuti, J.F., Izard, J., Segata, N., Gevers, D., Raes, J., and Huttenhower, C.
522   (2012) Microbial Co-occurrence Relationships in the Human Microbiome. *PLOS*
523   *Computational Biology* **8**: e1002606.
524
525   Fernandes, A.D., Reid, J.N., Macklaim, J.M., McMurrough, T.A., Edgell, D.R., and Gloor, G.B.
526   (2014) Unifying the analysis of high-throughput sequencing datasets: characterizing
527   RNA-seq, 16S rRNA gene sequencing and selective growth experiments by
528   compositional data analysis. *Microbiome* **2**: 15.
529
530   Ferris, M.J., Kühl, M., Wieland, A., and Ward, D.M. (2003) Cyanobacterial Ecotypes in
531   Different Optical Microenvironments of a 68°C Hot Spring Mat Community Revealed by
532   16S-23S rRNA Internal Transcribed Spacer Region Variation. *Appl Environ Microbiol*
533   **69**: 2893–2898.
534
535   Fierer, N. and Jackson, R.B. (2006) The diversity and biogeography of soil bacterial
536   communities. *Proc Natl Acad Sci USA* **103**: 626–631.
537
538   Friedman, J. and Alm, E.J. (2012) Inferring Correlation Networks from Genomic Survey Data.
539   *PLOS Computational Biology* **8**: e1002687.
540
541   Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010)
542   New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
543   performance of PhyML 3.0. *Syst Biol* **59**: 307–321.
544
545   Green, J.L., Bohannan, B.J.M., and Whitaker, R.J. (2008) Microbial Biogeography: From
546   Taxonomy to Traits. *Science* **320**: 1039–1043.
547
548   Horner-Devine, M.C. and Bohannan, B.J.M. (2006) Phylogenetic Clustering and Overdispersion
549   in Bacterial Communities. *Ecology* **87**: S100–S108.
550
Hui, F.K.C., Taskinen, S., Pledger, S., Foster, S.D., and Warton, D.I. (2015) Model-based
approaches to unconstrained ordination. *Methods in Ecology and Evolution* **6**: 399–411.

Hutchinson, G.E. (1957) Concluding Remarks. *Cold Spring Harb Symp Quant Biol* **22**: 415–427.

Hutchinson, G.E. (1961) The Paradox of the Plankton. *The American Naturalist* **95**: 137–145.

Kamneva, O.K. (2017) Genome composition and phylogeny of microbes predict their co-
occurrence in the environment. *PLOS Computational Biology* **13**: e1005366.

Katoh, K. and Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol **30**: 772–780.

Koeppel, A.F. and Wu, M. (2014) Species matter: the role of competition in the assembly of
congeneric bacteria. *ISME J* **8**: 531–540.

Koenker, R. (2015). Package 'quantreg'

Kraft, N. J. B., Adler, P. B., Godoy, O., James, E. C., Fuller, S., and Levine, J. M. (2015).
Community assembly, coexistence and the environmental filtering metaphor. Funct. Ecol.
29, 592–599.

Kumar, S., Stecher, G., and Tamura, K. (2016) MEGA7: Molecular Evolutionary Genetics
Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* **33**: 1870–1874.

Kurmayer, R., Christiansen, G., and Chorus, I. (2003) The Abundance of Microcystin-Producing
Genotypes Correlates Positively with Colony Size in Microcystis sp. and Determines Its
Microcystin Net Production in Lake Wannsee. *Appl Environ Microbiol* **69**: 787–795.

Letten, A.D., Keith, D.A., Tozer, M.G., and Hui, F.K.C. (2015) Fine-scale hydrological niche
differentiation through the lens of multi-species co-occurrence models. *Journal of*
Ecology 103: 1264–1275.

Magurran, A.E. and Henderson, P.A. (2003) Explaining the excess of rare species in natural
species abundance distributions. *Nature* **422**: 714.

Martiny, J.B.H., Jones, S.E., Lennon, J.T., and Martiny, A.C. (2015) Microbiomes in light of
traits: A phylogenetic perspective. *Science* **350**: aac9323.

Needham, D. M., Fuhrman, J. A. (2016) Pronounced daily succession of phytoplankton, archaea
and bacteria following a spring bloom. *Nature Microbiology*, **1**: 16005.

Newton, R.J., Jones, S.E., Eiler, A., McMahon, K.D., and Bertilsson, S. (2011) A guide to the
natural history of freshwater lake bacteria. *Microbiol Mol Biol Rev* **75**: 14–49.

Pascual-García, A., Tamames, J., and Bastolla, U. (2014) Bacteria dialog with Santa Rosalia: Are
aggregations of cosmopolitan bacteria mainly explained by habitat filtering or by
ecological interactions? *BMC Microbiol* **14**: 284.

Preheim, S.P., Perrotta, A.R., Martin-Platero, A.M., Gupta, A., and Alm, E.J. (2013)
Distribution-based clustering: using ecology to refine the operational taxonomic unit.
Appl Environ Microbiol **79**: 6593–6603.

Price MN, Dehal PS, Arkin AP. (2009). FastTree: computing large minimum evolution trees with
profiles instead of a distance matrix. *Mol Biol Evol* **26**:1641–1650.

Rohwer, R.R., Hamilton, J.J., Newton, R.J., and McMahon, K.D. (2018) TaxAss: Leveraging a
Custom Freshwater Database Achieves Fine-Scale Taxonomic Resolution. *mSphere* **3**:
e00327-18.

Röttjers, L. and Faust, K. (2018) From hairballs to hypotheses–biological insights from microbial
networks. *FEMS Microbiol Rev* **42**: 761–780.

Scanlan, D.J. (2003) Physiological diversity and niche adaptation in marine Synechococcus. In,
Advances in Microbial Physiology. Academic Press, pp. 1–64.

Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., et al. (2009)
Introducing mothur: Open-Source, Platform-Independent, Community-Supported
Software for Describing and Comparing Microbial Communities. *Appl Environ Microbiol*
75: 7537–7541.

Stockner, J., Callieri, C., and Cronberg, G. (2002) Picoplankton and Other Non-Bloom-Forming
Cyanobacteria in Lakes. In, Whitton,B.A. and Potts,M. (eds), *The Ecology of*
Cyanobacteria: Their Diversity in Time and Space. Dordrecht: Springer Netherlands, pp.
195–231.

Tromas, N., Taranu, Z.E., Martin, B.D., Willis, A., Fortin, N., Greer, C.W., and Shapiro, B.J.
(2018) Niche Separation Increases With Genetic Distance Among Bloom-Forming
Cyanobacteria. *Front Microbiol* **9** : 438

Vörös, L., Callieri, C., Balogh, K.V., and Bertoni, R. (1998) Freshwater picocyanobacteria along
a trophic gradient and light quality range. *Hydrobiologia* **369**: 117–125.

Warton, D.I., Blanchet, F.G., O'Hara, R.B., Ovaskainen, O., Taskinen, S., Walker, S.C., and Hui,
F.K.C. (2015) So Many Variables: Joint Modeling in Community Ecology. *Trends in*
Ecology & Evolution **30**: 766–779.

Webb, C.O., Ackerly, D.D., McPeek, M.A., and Donoghue, M.J. (2002) Phylogenies and
Community Ecology. *Annual Review of Ecology and Systematics* **33**: 475–505.

Zheng, Q., Wang, Y., Xie, R., Lang, A.S., Liu, Y., Lu, J., et al. (2018) Dynamics of
Heterotrophic Bacterial Assemblages within Synechococcus Cultures. *Appl Environ*
Microbiol **84**: e01517-17.

Data availability

Raw sequence data have been deposited NCBI GenBank under BioProject number

PRJNA544938.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

NT is funded by a project from the European Union's Horizon 2020 research and innovation

program under the Marie Sklodowska-Curie grant agreement No 656647. BJS was supported by

a Canada Research Chair and NSERC Discovery Grant. Sampling and samples processing were

supported by a grant from Furnas Centrais Hidroelétrica S.A. (Brazil) to AG. JSMP and DAP

received a scholarship from CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível

Superior). A sabbatical visit of AG to the Université de Montréal AG was supported by a CAPES

Senior Fellowship.