The evolution of realized niches within freshwater *Synechococcus*

Authors: Nicolas Tromas[1#]*, Zofia E. Taranu[2#], Mathieu Castelli[3#], Juliana S. M. Pimentel[4],

Daniel A. Pereira[4], Romane Marcoz[1], B. Jesse Shapiro[1]* and Alessandra Giani[4]

1- Département de sciences biologiques, Université de Montréal, 90 Vincent-d'Indy, Montréal,

QC, Canada, Montréal, QC H2V 2S9, Canada

2- Environnement et Changement Climatique Canada,105 Rue McGill, Montréal, QC H2Y 2E7

3- mathieu.castelli@gmail.com

4- Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil

*Corresponding authors: B. Jesse Shapiro. Phone: 514-343-6033. E-mail: jesse.shapiro@umontreal.ca; Nicolas Tromas. Phone 514-343-3188. E-mail: nicolas.tromas@umontreal.ca

# These authors contributed equally to this work

Originality-Significance Statement

We address a fundamental question in ecology and evolution: how do niche preferences change

over evolutionary time? Using time-series analysis of 16S rRNA gene amplicon sequencing data,

we develop an approach to highlight the importance of biotic factors in defining realized niches,

and show how niche preferences change proportionally with the 16S gene molecular clock within

the genus *Synechococcus.* Ours is also one of few studies on the ecology of freshwater

Synechococcus, adding significantly to our knowledge about this abundant and widespread

lineage of Cyanobacteria.

Summary

Understanding how ecological traits have changed over evolutionary time is a fundamental question in biology. Specifically, the extent to which more closely-related organisms share similar ecological preferences due to phylogenetic conservation – or if they are forced apart by competition – is still debated. Here we explored the co-occurrence patterns of freshwater cyanobacteria at the sub-genus level to investigate whether more closely-related taxa share more similar niches, and to what extent these niches were defined by abiotic or biotic variables. We used deep 16S rRNA gene amplicon sequencing and measured several abiotic environmental parameters (nutrients, temperature, etc.) in water samples collected over time and space in Furnas Reservoir, Brazil. We found that relatively more closely-related *Synechococcus* (in the continuous range of 93-100% nucleotide identity in 16S) had an increased tendency to co-occur with one another (*i.e.* had similar realized niches). This tendency could not be easily explained by shared preferences for measured abiotic niche dimensions. Thus, commonly measured abiotic parameters might not be sufficient to characterize, nor to predict community assembly or dynamics. Rather, co-occurrence between *Synechococcus* and the surrounding community (whether or not they represent true biological interactions) may be a more sensitive measure of realized niches. Overall, our results suggest that realized niches are phylogenetically conserved, at least at the sub-genus level and at the resolution of the 16S marker. Determining how these results generalize to other genera and at finer genetic resolution merits further investigation.

Introduction

A bacterial community is a group of potentially interacting organisms that coexist at a particular place and time (Magurran, 2003). Environmental selective pressures are a strong force shaping microbial community assembly (Martiny et al, 2015). We know, for example, that certain abiotic factors explain a large portion of the variation in microbial community composition (e.g. the effect of pH on soil bacterial communities; Fierer and Jackson, 2006). Therefore, associations between microbial traits - generally defined as a phenotypic response to a specific environmental condition - and abiotic niches could help explain community assembly rules (Green et al., 2008; Burke et al., 2011). However, using trait-based approaches to understand communities may be challenging, as important abiotic variables may go unmeasured. Even less is known about biotic interactions, despite their importance in determining community composition, diversity, and dynamics (Needham et al., 2016).

Abiotic factors are generally thought to determine an organism's fundamental niche (where it is theoretically capable of living), whereas biotic factors determine its realized niche (where it actually lives in nature; Hutchinson, 1957). Species (or taxonomic) co-occurrence networks are often used to infer niche similarity among organisms that tend to co-occur in nature over space and time, and microbial co-occurrence networks can easily be constructed from 16S rRNA gene amplicon sequencing surveys (Friedman and Alm, 2012; Röttjers and Faust, 2018; Tromas et al., 2018). However, co-occurrence can be driven by both abiotic and biotic factors, which are hard to disentangle in practice (Kraft et al., 2015). Regardless, organisms sharing a similar niche are expected to be associated with similar surrounding communities (Cohan and Koeppel, 2008; Faust et al.2012; Pascual-García et al., 2014).

A fundamental question spanning ecology and evolution is how ecological traits change over

evolutionary time. For example, some traits (such as bacteriophage host range or phosphorus

utilization) evolve rapidly at the tips of a phylogenetic tree, whereas other traits (such as salinity

preference) are deeply conserved (Martiny *et al.* 2015). When more closely related organisms

share similar ecological preferences, so-called "habitat filtering" or "environmental" filtering is

expected to result in phylogenetic clustering, meaning that a community tends to contain more

closely related organisms than expected by a random draw from the phylogeny (Webb et al.,

2002; Horner-Devine & Bohannan, 2006; Martiny et al., 2015). In contrast, if close relatives

evolve different traits to avoid competitive exclusion, this will result in phylogenetic

overdispersion (*i.e.* a community composed of more distant relatives than expected by chance).

However, the relative importance of these two processes in shaping microbial communities is still

widely debated, and difficult to distinguish (Koeppel and Wu, 2014; Cadotte and Tucker, 2017).

We have previously shown that within the cyanobacterial genus *Dolichospermum* (Tromas et al.,

2018), the relationships between phylogenetic distance and ecological similarity varies by trait,

suggesting that it might be necessary to analyze each niche dimension or trait separately (Martiny

et al., 2015).

In this study, we explored the co-occurrence patterns of freshwater cyanobacteria at the sub-

genus level, to investigate if more closely related taxa have more similar niches, and to what

extent these niches can be quantified by abiotic or biotic variables. We use the term "closely

related" as a relative term, describing a continuous range of genetic similarity (93-100%

nucleotide identity) in 16S rRNA gene sequences. We focused on *Synechococcus*, the most

abundant cyanobacterial genus in Furnas Reservoir (Brazil) at the time of sampling (2006-2008).

Synechococcus is among the most abundant organisms living in oceans and lakes (Stockner et al.

2000; Scanlan, 2003). The phylogenetic coherence of the genus has been questioned by a recent

study showing it to be polyphyletic (Coutinho et al 2016). *Synechococcus* is physiologically

highly plastic, ubiquitous, and able to acclimate to different environmental conditions (Callieri

1996; Vörös et al. 1998; Callieri et al., 2011). Previous studies have shown that different

Synechococcus strains could co-exist in the same site but respond differently to environmental

changes, suggesting niche partitioning (Ferris et al., 2003; Allewalt et al., 2006; Becker et al.,

2007; Becraft et al., 2011; Callieri et al. 2012). Recently, Zheng et al. (2018), observed a

geographical pattern of heterotrophic bacteria associated with different marine *Synechococcus*

strains, indicating that strains living in the same area tend to be associated with similar

communities. It remains, however, unclear whether the *Synechococcus* genotype or the

environment are the main drivers of *Synechococcus* interactions with the surrounding microbial

community.

Using deep 16S rRNA gene amplicon sequencing of 86 water samples collected in time series

across nine locations in the Furnas Reservoir, we tracked genetic diversity within the

Synechococcus genus, along with the surrounding microbial community, and measured several

abiotic variables. We found that that more closely-related *Synechococcus* tended to co-occur with

one another and also with similar surrounding microbial communities. Such phylogenetic

clustering indicates that realized niche similarity tends to evolve proportionally with substitutions

in the 16S gene in the *Synechococcus* lineage (*i.e.* niches evolve according to the 16S gene

molecular clock). However, more closely-related *Synechococcus* did not have similar abiotic

niche preferences (with the exception of total phosphorus). These results suggest that biotic

factors may be strong niche determinants, which could be particularly useful when relevant

abiotic factors are unknown or cannot be readily measured. Alternatively, cryptic abiotic drivers

may determine niche and community structure, but biotic factors provide the most informative

measure of niche similarity.

Materials and methods

Sampling, environmental data measurements

A total of 90 water samples were collected from September 2006 to April 2008 at nine stations in

Furnas Reservoir (Minas Gerais, Brazil). Furnas is a large reservoir (1440 km^2), located in

southeastern Brazil (20^{o}40'S; 46^{o}19'W) and formed by the damming of two main rivers (Rio

Grande and Rio Sapucaí), which divide the reservoir in two separated branches (Figure S1).

Sampling stations F1, F4, F6, F9 are from a relatively pristine branch of the reservoir, whereas

F12, F14, F18 and F20 are impacted by human activities. Temperature profiles were measured in

the water column by aid of a multi-parameter probe (556 YSI, USA). Water samples were

collected from the euphotic zone (determined by Secchi disc depth) by a Van Dorn vertical water

sampler. Samples were stored in bottles that had been acid-washed and rinsed with deionized

water. A portion of each water sample was immediately filtered through glass fiber filters

(Whatmann GF/F, 0.7 μm pore size). The exact filtered volumes were recorded and filters were

kept frozen until further analyses (chlorophyll a, DNA and microcystin). For dissolved nutrient

analyses, 200 mL of filtered water samples were stored at −20 °C. For total phosphorus, samples

were frozen with no previous filtration. Nutrient analyses were performed using

spectrophotometric methods according to APHA (2005). All nutrient analyses (Nitrate, Nitrite

and total phosphorus (TP) were performed on three replicates.

DNA extraction, purification and sequencing

DNA was extracted from frozen filters according to Kurmayer et al. (2003), with few

modifications. Briefly, filters were treated with a sucrose buffer (25% w/v sucrose, 50 mM Tris-

HCl, 100 Mm EDTA, pH 8) on ice for 2 h and with addition of lysozyme (5mg/mL, 1h, 37°C).

Proteinase K (100 μg/mL) in sodium dodecyl sulfate (2% v/v) was added and filters were

incubated overnight at 55°C. A phenol:chloroform:isoamyl alcohol solution (25:24:1, v/v) was

used for protein precipitation and DNA isolation. The DNA was cleaned in 100% ethanol and

pellets rinsed with 70% ethanol. The DNA was resuspended in TE (10 mM Tris- HCl, pH 8, and

1 mM EDTA). The DNA extract was quantified by a spectrophotometer (Perkin Elmer, Lambda

25), at 260 nm and 280 nm, and its quality checked in 1% (w/v) agarose gel, stained with

ethidium bromide. DNA samples were stored at -20ºC.

Sequence analysis

We followed the same protocol described in Tromas et al., (2017) to generate a library of

V4 region amplicons. Libraries for 2x250bp paired-end Illumina sequencing were prepared using

a two-step 16S rRNA gene amplicon PCR as described in Preheim et al. (2013). We performed

one sequencing run using MiSeq reagent Kit V2 (Illumina, San Diego, CA, USA) on a MiSeq

instrument (Illumina). A total of 4,476,747 sequences of the 16S rRNA gene V4 region were

obtained from 90 lake samples, two negative controls, and two mock community samples. We

obtained a median of 37,682 sequences per sample. Using a similar approach as described in

Tromas *et al*., (2018), we processed the sequences with SmileTrain

(https://github.com/almlab/SmileTrain/wiki; Preheim et al., 2013) for read quality filtering,

primer removal, chimera filtering, and merging using USEARCH (version 7.0.1090,

http://www.drive5.com/usearch/, default parameter) (Edgar, 2010), Mothur (version 1.33.3)

(Schloss *et al*., 2009), and Biopython (version 2.7). Minimum Entropy Decomposition (MED)

177 was then applied to the filtered and merged reads to partition sequence reads into MED nodes

178 (Eren *et al*., 2015). MED was performed using the following parameters: –M noise filter set to

179 1000, resulting in ~17.5% of reads filtered and 466 MED nodes representing the whole bacterial

180 community. Samples with less than 1000 reads were removed, yielding a final dataset of 86

181 reservoir samples. Finally, we assigned taxonomy to MED nodes using the assign_taxonomy.py

182 QIIME script (default parameters), and a combination of GreenGenes and a freshwater-specific

183 database (Freshwater database 2016 August 18 release; Newton *et al*., 2011), using TaxAss

184 (https://github.com/McMahonLab/TaxAss, installation date: September 13[th] 2016; Rohwer *et al*.,

185 2017). We found a total of 40 *Synechococcus* nodes, which we refer to as strains below.

187 ***Spatio-temporal analysis***

188  We used multivariate regression tree analyses (Breiman *et al*. 1984; De'ath 2002) to

189 investigate if spatio-temporal variables could explain genetic variation within *Synechococcus*. We

190 used two different temporal predictors: year and month and one spatial predictor: station. The

191 analysis was performed using the function *mvpart*() and *rpart.pca*() from the R mvpart package

192 (Therneau and Atkinson, 1997; De'ath, 2007). Prior to analysis, the *Synechococcus* MED node

193 table was Hellinger transformed to downweight the effect of double-zeros (Rao, 1995).

195 ***Genetic distance between Synechococcus nodes***

196  We measured the genetic distance between *Synechococcus* MED nodes using the software

197 MEGA (Kumar et al., 2016; version 7.0.18) with the p-distance (the proportion of nucleotide

198 sites at which two sequences differ), calculated by dividing the number of sites with nucleotide

199 differences by the total number of sites compared (excluding sites with gaps).

Co-response to abiotic factors

As described in Tromas *et al.* (2018), we used a Latent Variable Model (LVM) framework (boral package in R; Hui 2015, Warton *et al.* 2015) to explore how *Synechococcus* nodes co-responded to abiotic gradients and used these co-responses as indicators of niche similarity. That is, for each abiotic factor, we ran separate LVMs, regressing the bacterial community as a function (both linear and non-linear) of the given factor. This component of the LVM thus defined the taxon-specific environmental responses. To then identify remaining patterns of co-occurrence after accounting for all measured environmental variables, we fit a global LVM which included all abiotic factors and two latent variables (*sensu* Letten *et al.* 2015 and Warton *et al.* 2015). To visualize patterns of co-occurrence arising from the different environmental factors, we calculated two types of correlation matrices. The first, a co-response correlation matrix, was constructed by calculating, for any two nodes, the correlation between their fitted values. This correlation matrix thus represented the correlation between nodes that can be attributed to a shared or diverging environmental responses. A significant positive correlation between the fitted response of any two taxa to an environmental variable represented a co-response (i.e., as one taxa increases in response to an environmental variable, the other likewise increases), whereas a significant negative correlations between the fitted response of two taxa represented some degree of niche separation (i.e., as one taxa increases in response to an environmental variable, the other decreases).

To account for the correlation between nodes that may be attributable to biotic processes or missing environmental covariates, a second type of correlation matrix, a residual correlation matrix, was calculated using the latent variable coefficients of the global LVM. Since the latent variables are the output of an ordination of the residuals, we expected weak residual correlations

induced by the latent variables if the environmental variables were the dominant force structuring

patterns of species co-occurrence (*i.e.* the model explained most of the variance, with little left

unexplained in the residuals). Conversely, if any unmeasured environmental factors and/or biotic

processes are equally, or more, important than measured environmental factors, we expected

strong correlations based on the latent variables (*i.e.* the model was poor and much of the

observed variability remained in the residuals).

To test how *Synechococcus* niche similarity varied with genetic distance, we examined the

relationship between the co-response of *Synechococcus* taxa to environmental parameters and

their genetic distance (i.e., plotting the correlation coefficient of the LVM co-responses vs.

genetic distances) (R_script1). However, given the large number of environmental factors driving

phytoplankton community dynamics (Hutchinson, 1961), we expected some degree of co-

limitation whereby the response of a taxon to an environmental variable (and consequently the

degree of co-response among any two taxa) would be limited by other environmental variables.

In such cases, the taxon's response and rate of change would have an upper limit (set by all

measured environmental factors) but may not reach this limit if other, unmeasured factors are co-

limiting (Cade and Noon 2003). As an increasing number of unmeasured factors become limiting

at some sample location, or time point, the relationship between the response and the measured

factor becomes increasingly heterogeneous or wedge shaped. When such heterogeneous

variances are observed (wedge shape biplot), it suggests that there is not a single slope coefficient

that characterizes the relationship, and that focusing solely the coefficient fit by ordinary least

squares regression (mean response) may underestimate the true rate of change. Thus, to

accurately examine co-limitations among measured and unmeasured factors, we applied quantile

regressions using 'quantreg' R package (Koenker, 2015) (R_script1), which, instead of fitting a

regression to the mean response (ordinary least squares regression), fits regression curves to other

quantiles of the response variable (Cade and Noon 2003). Finally, to provide an effect size for the

linear quantile relationships of abiotic co-responses vs. genetic distance, we calculated the

goodness of fit measure for quantile regression for a suite of quantiles (where the goodness of fit

is estimated as 1 minus the ratio between the sum of absolute deviations of the parameterized

model and the sum of absolute deviations of the unparameterized, null model). Note that this

goodness of fit measure (referred hereafter as R1) is not comparable to the OLS coefficient of

determination (R2); it is based on the variance of absolute deviations as opposed to the variance

of squared deviations, and consequently, R1 is always smaller than R2 (Koenker and Machado,

1999).

258
Co-occurrence with biotic factors

A caveat of the above LVM-quantile regressions is that although it identifies whether

some relationships are limited by unmeasured abiotic and/or biotic factors, it could not tease apart

the relative importance of each type of factor (everything is instead lumped as an unknown). In

order to determine the relative contribution of biotic processes, we thus quantified node co-

occurrence patterns among *Synechococcus* and the remaining community. In particular, we

explored the relationship between *Synechococcus* nodes and the surrounding community by

measuring: (1) co-occurrences between *Synechococcus* and other taxa, and (2) paired differences

in co-occurrence among *Synechococcus* nodes and a specific taxon. For the former, we calculated

co-occurrences among taxa using SparCC (Friedman and Alm, 2012), including 20 iterations to

estimate the median correlation of each pair of MED nodes, 500 bootstraps to assess the

statistical significance and centered log ratio (CLR) transform to correct for compositionality.

Correlations were then filtered using a false discovery rate threshold ($Q < 0.05$). For the latter, we

further calculated the absolute difference of SparCC correlation (*r*) between a *Synechococcus*

node (X_i) and a specific taxon T, such that $|\Delta r| = |\ \text{Corr}(X_1, T) - \text{Corr}(X_2, T)\ |$ where Corr is

defined here as the SparCC correlation between X_i and T.

For each non-*Synechococcus* taxon, we then estimated the relationship between $|\Delta r|$ and

the distance between the given *Synechococcus* nodes (X_1 and X_2). A positive correlation would

indicate that more closely related *Synechococcus* nodes have a lower $|\Delta r|$ than more distant

nodes, *i.e.* more closely related nodes would have more similar correlations with potentially

interacting community members. To reduce potential bias or noise due to a small sample size of

Synechococcus pairs, we selected all non-*Synechococcus* nodes that were significantly co-

occurrent with at least seven different *Synechococcus*. This cutoff was chosen to ensure at least

20 unique pairs of *Synechococcus* for each non-*Synechococcus* taxon. A total of 373 non-

Synechococcus taxa were selected using this cutoff, and the correlation between $|\Delta r|$ and genetic

distance was non-significant (Spearman correlation, $P<0.05$) for 120 of them. Finally, we

performed a permutation test to quantify any bias in the method due to data structure. For each of

the significant non-*Synechococcus* taxa, we estimated the rate of false positive correlations by

sampling values of *Synechococcus* genetic distance with replacement, while randomizing the

association between genetic distance and $|\Delta r|$. We performed 1000 permutations for each

significant non-*Synechococcus* taxon using the script R_script2. We then recorded the proportion

(*p*) of permutations yielding a larger correlation than observed, after adding a pseudocount of 1 to

both the numerator (the number of permutations yielding a larger correlation than observed) and

the denominator (the total number of permutations).

293
294   ***Phylogenetic analysis***
295
We first verified that our *Synechococcus* sequences were monophyletic by building a

phylogenetic tree with FastTree (version 2.1.8, Price et al., 2009), using all MED node sequences

aligned with MAFFT (v7.154b; Katoh and Standley, 2013). We then specifically aligned

Synechococcus MED node sequences using muscle (in Mega, version 7.0.26; Kumar et al., 2016*)*

and built phylogenetic trees using PhyML with the general time-reversible (GTR) model of

nucleotide substitution (version 3.0; Guindon et al., 2010). We used the ALDEx2 R package

(version: 1.5.0; Fernandes *et al*., 2014) and the aldex() function to identify MED nodes associated

with the pristine or human-impacted branches of Furnas Reservoir. To do so, we used Welsh's t-

test and 128 Monte Carlo samples. ALDEx2 uses the CLR transformation to avoid compositional

effects. A Q-value below 0.05 after Benjamini-Hochberg correction was taken as evidence for

association. We also used a second association method, DESeq2 implemented in the

"MicrobiomeAnalyst" web-based tool (Dhariwal et al., 2017).

Statistical analysis
310

311   All statistical tests were performed in R.

Results

Abiotic factors shape Synechococcus community structure

Furnas Reservoir is divided into two branches: one, which is heavily human-impacted (Figure S1, sampling stations 12-20) and the other, less impacted (stations 1-9). *Synechococcus* dominated the bacterial community in both branches during the sampling period (2006-2008) (Figure S2) and formed a monophyletic group (Figure S3, S4). Furnas Reservoir is meso-eutrophic based on phosphorus concentrations. Total phosphorus (TP) did not vary significantly across stations but did vary significantly within each station over time (Figure S5). Temperature did not vary widely across sampling stations or time. We first asked whether particular *Synechococcus* strains (which we used interchangeably with MED nodes) were associated with different branches of the reservoir, different time periods, or different abiotic factors. We applied a multivariate regression tree analysis (Breiman et al. 1984; De'ath 2002) to test whether the composition of *Synechococcus* strains changed over time and space. We found that the year and month of sampling did not explain much variance ($R^2=0.02$ and $R^2=0.03$, respectively), nor did the station ($R^2=0.06$) or branch of the reservoir ($R^2=0.02$). In contrast, Nitrite, TP, depth, and water temperature all explained a significant proportion of the variance (Figure S6; 22.5% of variance explained). Nitrite is the single most explanatory variable (13% of variance, but also colinear with temperature and depth; Figure S6), followed by TP (5.5% of variance; Figure S7).

More closely related Synechoccus do not share similar preferences for most measured abiotic niches

Having established that abiotic factors, but not temporal or spatial variation, explain a significant amount of variation in the relative abundances of *Synechococcus* strains, we asked whether more

closely related strains tended to share similar abiotic niches. We examined *Synechococcus* 16S

sequences across a range of ~93-100% nucleotide identity, which is a rather large range

considering that niche partitioning can occur at relatively high identity, >97% (Martiny *et al*.

2015 ; Larkin and Martiny, 2017). Here we defined abiotic niche similarity by estimating the co-

responses of pairs of *Synechococcus* strains to each abiotic factor using LVM-quantile regression

(Methods). For most abiotic factors, we did not observe any significant relationship between co-

responses and genetic distance, with the exception of total phosphorus for which the relationship

was negative (Median p-value across quantiles = 0.021; median R1 across quantiles = 0.12)

(Figure 1 & S8, Table S2). Similarly, the degree of abiotic niche separation (*i.e.* significant

negative correlations between the fitted response of any two *Synechococcus* strains to an abiotic

factor) was generally independent of genetic distance (Figure S9, S10), except for depth and

Nitrite which were significant at some quantiles but with quite a poor fit (median R1 across

quantiles ≤ 0.02)(Table S2C).

We also evaluated non-linear quantile relationships among co-responses and genetic

distances, but we never found a significant non-linear relationship over the total range of genetic

distances (Table S2A-C). We repeated the quantile regression analysis in a restricted range of

100-97% identity, to see if any linear trend is discernible in a range roughly corresponding with

species or genera (Figure S11). The negative linear trend observed between co-responses and

genetic distance for TP in the full range of genetic identify (Figure 1) was still observed in the

restricted range (Figure S11) although not significant at all quantiles (Table S2D). There was also

some evidence for a non-linear relationship between genetic distance and both positive and

negative co-responses to TP (Table S2 D, E), suggesting possible niche separation among closely

related (~99% identical) strains. However, this trend is driven by just three pairs of very closely

related strains with quite different TP preferences (Figure S12). Overall, these results suggest that

TP-related niches show some evidence for phylogenetic conservation within freshwater *Synechoccus*, while the evidence for other niches is weak or inconclusive.

Figure 1. Relationship between *Synechococcus* niche similarity and genetic distance. Niche similarity is defined as a positive LVM co-response (y-axis). Shown are the scatterplots of LVM co-responses versus each variables superimposed with the lines for 5th, 10th, 15th, 25th, 35th, 45th, 55th, 65th, 75th, 85th and 90th quantile regression fits (grey lines, for each co-response quantile), the median fit (50th quantile; blue line), and the least squares estimate or mean fit (dashed red line).

Biotic processes or unmeasured environmental variables could define *Synechococcus* realized niches, and are captured in the residual variation of the global LVM. We next asked whether more closely-related *Synechococcus* strains had more similar co-responses to this residual variation. For the positive co-responses (Figure 2A, B), we observed a decrease with genetic

distance. This result suggests a negative relationship between the residuals (*i.e.* unmeasured

abiotic variables and/or biotic niche preferences) and phylogenetic distance. For the negative

residual correlations (Figure 2C, D), there was no apparent relationship with genetic distance.

Therefore, there remain significant co-responses that were not explained by the abiotic factors

measured in this study. Overall, most of the measured abiotic environmental variables were not

related to *Synechococcus* genetic distance. However, preferences for unmeasured biotic or abiotic

factors do tend to change with genetic distance, such that more closely-related *Synechococcus*

have similar realized niche preferences.

Figure 2. Relationship between *Synechococcus* co-responses (LVM residuals) and genetic distance. Positive co-responses are shown in panels **A** and **B**; negative co-responses in panels **C** and **D.** *(*A, C*)* Scatterplots of LVM co-responses versus LVM residuals superimposed with the lines for 5th, 10th, 15th, 25th, 35th, 45th, 55th, 65th, 75th, 85th and 90th quantile regression fits (grey lines), the median fit (50th quantile; blue line), and the least squares estimate or mean fit (dashed red line). (B, D) Panels show the quantile slope estimates (dark grey line) and corresponding confidence intervals (light grey bands) of the response models across all quantiles (arrows, with the median shown in blue). Significant quantile slopes occur when confidence intervals do not overlap zero.

More closely related Synechococcus share similar biotic interactions

The observed tendency for more closely-related *Synechococcus* to share similar realized niches

(Figure 2A) could be explained by shared preferences for either unmeasured abiotic factors or

biotic interactions. We used co-occurrence network analysis to explore the role of potential biotic

interactions. First, we confirmed that more closely related *Synechococcus* tend to co-occur across

samples. We observed a negative relationship between node co-occurrence and pairwise genetic

distance (Figure 3), indicating that genetically similar *Synechococcus* nodes are indeed more

likely to co-occur, and thus to have similar realized niches. This pattern was significant [linear

regression, $F(1,488) = 108.6$, $P < 0.001$, adjusted $R^2 = 0.18$].

Figure 3. **Negative relationship between pairwise SparCC correlation coefficients and**

pairwise genetic distance between *Synechococcus* nodes. Only significant SparCC correlations

(Q < 0.05) were included. Pairwise genetic distance was computed as percent nucleotide identity

(p-distance). Blue diamonds represent the mean SparCC correlation for each distance. Boxplots

show the median (horizontal line), the 25th and 75th percentile (enclosed in box) and 95%

confidence intervals (whiskers).

If more closely-related *Synechococcus* tend to co-occur, it follows that they should also co-occur

with more similar surrounding microbial communities. The composition of the surrounding

microbial community could provide information about the nature of the realized niche, while also

suggesting possible biotic interactions. To confirm the expectation that more closely-related

Synechococcus should also share similar surrounding communities, we performed a pairwise

analysis to examine the relationship between *Synechococcus* genetic distance and $|\Delta r|$, a measure

of the similarity of *Synechococcus* co-occurrence with non-*Synechococcus* taxa (Methods). A

total of 373 non-*Synechococcus* taxa were analysed and the correlation between $|\Delta r|$ and genetic

distance was non-significant for 120 of them. The remaining 253 significant non-*Synechococcus*

have consistently positive relationships between co-occurrence and genetic distance, *i.e.* more

closely related *Synechococcus* nodes have a higher chance of co-occurring with similar non-

Synechococcus taxa (Figure 4). This result is robust to data structure, as determined by permuting

the associations between genetic distance and $|\Delta r|$ (Figure S13; Methods).

Figure 4. More closely related *Synechococcus* nodes co-occur with similar surrounding community. The histogram shows the distribution of Spearman correlations between the genetic distance of *Synechococcus* nodes and their association with similar surrounding communities. Spearman correlations were calculated between the genetic distance of *Synechococcus* pairs and the absolute difference of SparCC correlation (r) of a *Synechococcus* node with a specific taxon T, such that $|\Delta r| = |\ Corr(X_1, T) - Corr(X_2, T)\ |$ where Corr is defined here as the SparCC correlation score of X_i (*Synechococcus* node) and T (non-*Synechococcus* node). We then estimated the correlation of $|\Delta r|$ and the genetic distance between nodes X_1 and X_2. The mean Spearman correlation estimate was 0.373 (red line) and 99.6% of correlations are positive.

Finally, we selected the non-*Synechococcus* taxa with the highest correlation scores between $|\Delta r|$ and *Synechococcus* distance genetic. Five taxa had correlation estimates higher than 0.6,

including three Proteobacteria, one Verrucomicrobia, and one Planctomycetes (Table S1).

Previous studies have shown that members of the *Comamonadaceae* family and *Spirobacillales*

order (members of Proteobacteria; Table S1) were associated with *Synechococcus* (Guedes et al.,

2018) or more broadly with algae (Fullbright et al., 2018). Moreover, the *Gallionella* genus was

previously observed colonizing filamentous algae (Mori et al., 2015).

Overall, these results show that the co-occurring surrounding community is, on average, more

similar for more closely related *Synechococcus*. Members of this surrounding community could

interact directly with *Synechococcus,* or share similarity in their abiotic niches.

Discussion

In this study, we investigated *Synechococcus* niche preferences at relatively fine (sub-

genus) taxonomic resolution using a 16S sequencing approach previously applied to other

bacterial lineages (Koeppel and Wu, 2014; Tromas et al., 2018). However, we acknowledge that

even at the finest possible taxonomic resolution, the 16S gene may be too slow-evolving to be a

good marker for fast-evolving traits, such as toxin production (Berry et al., 2017) or

bacteriophage resistance (Martiny et al., 2015). We found that water temperature, reservoir depth,

and Nitrite concentration greatly affected the relative abundance of *Synechococcus* strains (Figure

S6, S7). However, we did not find significant linear or non-linear relationships between genetic

distance and preference for any of these abiotic niches. This was true when considering the whole

range of genetic distance (>93% identity) or a restricted range of more closely related strains

(>97%), suggesting that these traits are phylogenetically unconserved (at least within our dataset

and at the resolution of the 16S marker). Such a lack of conservation would be expected for a trait

such as Nitrite or Nitrate utilization in an organism capable of fixing nitrogen. Although some

marine *Synechococcus* can fix nitrogen (Zehr et al., 2001), further investigation will be needed to

determine if this is also the case in our freshwater community.

In contrast, total phosphorus (TP) preference was more phylogenetically constrained, even

though it explained less variation in community structure than Nitrite (Figure S6, S7).

Phosphorus use may therefore evolve "clock-like" along the *Synechococcus* phylogeny, following

an approximately linear model in the range of 96-100% nucleotide identity (Figure 1). This

contrasts with the observation in another cyanobacterium (*Prochlorococcus*) that organic

phosphate niches are fast-evolving and thus correlated with the phylogeny only near the tips of

the phylogenetic tree (Coleman & Chisholm 2010; Martiny et al., 2015). Consistent with this

idea, we identified significant niche separation among very closely related (~99% identical in

16S) *Synechococcus*, although the effect was driven by only three pairs of strains (Figure S12).

We also note that TP is a fairly crude measurement, encompassing both organic and inorganic

forms of phosphorus, and adaptation to these finer-grained niche dimensions may evolve

differently in *Synechoccus*. TP may not even be the driving selective pressure and may simply be

a marker of another important factor (*e.g.* runoff, overall biomass, etc.). Furnas Reservoir is

meso-eutrophic and phosphorus is probably not limiting (concentration varies from 8 to 53µg/L,

with a mean ~25µg/L). This might explain why only 77 of the 780 *Synechococcus* strain pairs

showed a significant co-response to phosphorus (Figure 1), indicating that the effect is driven by

a small number of closely-related strains with similar co-responses, and cannot be generalized to

all *Synechococcus*. This result might also be explained if *Synechococcus* represents a

polyphyletic group (Coutinho et al., 2016), composed of different clades with different

phosphorus niches. However, in our sample, *Synechococcus* is monophyletic (Figure S3, S4),

excluding this explanation. Overall, our results demonstrate how abiotic factors can shape

community composition within a genus but measured abiotic niches generally do not evolve

along the genus-level phylogeny.

Habitat filtering was originally defined to describe how abiotic factors select for genetically

similar (more closely related) organisms (Cadotte and Tucker 2017). Yet in practice, it is difficult

to disentangle the contributions of biotic and abiotic factors, and thus to satisfactorily define

habitat filtering (Kraft et al., 2015). In this study, we therefore estimated the influence of both

unmeasured abiotic variables and biotic factors, and found a negative association with genetic

distance (Figure 2). This suggests that more closely related *Synechococcus* tend to share similar

realized niches. As a result, we confirmed that genetically similar *Synechococcus* nodes tend to

co-occur, (Figure 3), as observed in studies using genomic (rather than 16S) similarity (Kamneva,

2017). This result could be unintuitive especially if *Synechococcus* nodes co-occur, but do not

share abiotic dimensions. For example, two taxa could be present in the same samples, but at very

different relative abundances. Therefore, they would co-occur but respond quantitatively to

different environmental variables. Overall, this result is consistent with habitat filtering, broadly

defined to include both biotic and abiotic factors.

Here we found that commonly-measured abiotic environmental factors (*e.g.* nutrients,

temperature) may be insufficient to capture relevant niche dimensions, and biotic co-occurrence

networks could be more informative about the evolution of niche preferences. In this study, we

measured abiotic factors such as temperature and nutrients, which are important in structuring

microbial communities, and yet these factors are largely insufficient to account for realized

niches or their distribution across the *Synechococcus* phylogeny. However, more closely-related

Synechococcus tend to co-occur with more similar surrounding microbial communities. As

previously suggested, the surrounding microbial community could provide hints as to the nature

of a realized niche (Cohan and Koeppel, 2008; Tromas et al., 2018). For example, the presence of

certain taxa could indicate the importance of an unmeasured abiotic factor (*e.g* Acidophiles or

Alkaliphiles as indicators of pH). In some cases, co-occurrence may indicate real biological

interactions (*e.g.* physical association, predation, cross-feeding, etc.). Some of the bacteria whose

co-occurrence with *Synechococcus* was strongly phylogenetically conserved have previously

been found in physical association with algae and cyanobacteria (Table S1; Mori et al. 2015;

Fullbright et al., 2018) and future work could experimentally test the nature of possible

interactions with *Synechococcus* strains. Alternatively, co-occurring microbes could simply share

similar unmeasured niches (including unmeasured abiotic or biotic factors such as phage or

protozoan grazers), and mining their genome sequences or metabolic capabilities could provide

insight into the nature of these niches. In this case, considering biotic factors allowed us to

uncover a signal of habitat filtering within *Synechococcus* that would have been obscured by only

considering commonly measured abiotic factors.

References

Allewalt, J.P., Bateson, M.M., Revsbech, N.P., Slack, K., and Ward, D.M. (2006) Effect of
530   Temperature and Light on Growth of and Photosynthesis by Synechococcus Isolates
531   Typical of Those Predominating in the Octopus Spring Microbial Mat Community of
532   Yellowstone National Park. *Appl Environ Microbiol* **72**: 544–550.
533
534   Becker, S., Richl, P., and Ernst, A. (2007) Seasonal and habitat-related distribution pattern of
535   Synechococcus genotypes in Lake Constance. *FEMS Microbiol Ecol* **62**: 64–77.
536
537   Becraft, E.D., Cohan, F.M., Kühl, M., Jensen, S.I., and Ward, D.M. (2011) Fine-Scale
538   Distribution Patterns of Synechococcus Ecological Diversity in Microbial Mats of
539   Mushroom Spring, Yellowstone National Park. *Appl Environ Microbiol* **77**: 7689–7697.
540
541   Berry, M. A., White, J. D., Davis, T. W., Jain, S., Johengen, T. H., Dick, G. J., et al. (2017). Are
542   oligotypes meaningful ecological and phylogenetic units? A case study of Microcystis in
543   freshwater lakes. *Front Microbiol* **8** : 365.
544
545   Breiman L, Friedman JH, Olshen RA, Stone CJ. (1984) Classification and Regression Trees.
546   Wadsworth International Group, Belmont, CA, USA.
547
548   Burke, C., Steinberg, P., Rusch, D., Kjelleberg, S., and Thomas, T. (2011) Bacterial community
549   assembly based on functional genes rather than species. *Proc Natl Acad Sci USA* **108**:
550   14288–14293.
551
552   Cade, B.S. and Noon, B.R. (2003) A Gentle Introduction to Quantile Regression for Ecologists.
553   *Frontiers in Ecology and the Environment* **1**: 412–420.
554
555   Cadotte, M.W. and Tucker, C.M. (2017) Should Environmental Filtering be Abandoned? *Trends
556   in Ecology & Evolution* **32**: 429–437.
557
558   Callieri, C. (1996) Extinction coefficient of red, green and blue light and its influence on
559   picocyanobacterial types in lakes at different trophic levels. *Memorie dell'Istituto Italiano
560   di Idrobiologia* 54: 135-142.
561
562   Callieri, C., Caravati, E., Corno, G., and Bertoni, R. (2012) Picocyanobacterial community
563   structure and space-time dynamics in the subalpine Lake Maggiore (N. Italy). *Journal of
564   Limnology* **71**: e9–e9.
565
566   Callieri, C., Lami, A., and Bertoni, R. (2011) Microcolony Formation by Single-Cell
567   Synechococcus Strains as a Fast Response to UV Radiation. *Appl Environ Microbiol* **77**:
568   7533–7540.
569
570   Cohan, F.M. and Koeppel, A.F. (2008) The origins of ecological diversity in prokaryotes. *Curr
571   Biol* **18**: R1024-1034.
572
573   Coleman, M. L., and Chisholm, S. W. (2010). Ecosystem-specific selection pressures revealed

574   through comparative population genomics. *Proc. Natl. Acad. Sci. U. S. A.* **107** : 18634–
575   18639.
576
577 Coutinho, F., Tschoeke, D.A., Thompson, F., and Thompson, C. (2016) Comparative genomics
578   of Synechococcus and proposal of the new genus Parasynechococcus. *PeerJ* **4**: e1522.
579
580 De'ath G. (2007) mvpart: Multivariate partitioning. R package version 1.6-2.
581
582 Dhariwal, A., Chong, J., Habib, S., King, I.L., Agellon, L.B., and Xia, J. (2017)
583   MicrobiomeAnalyst: a web-based tool for comprehensive statistical, visual and meta-
584   analysis of microbiome data. *Nucleic Acids Res* **45**: W180–W188.
585
586 Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
587   **26**: 2460–2461.
588
589 Eren, A.M., Morrison, H.G., Lescault, P.J., Reveillaud, J., Vineis, J.H., and Sogin, M.L. (2015)
590   Minimum entropy decomposition: unsupervised oligotyping for sensitive partitioning of
591   high-throughput marker gene sequences. *ISME J* **9**: 968–979.
592
593 Faust, K., Sathirapongsasuti, J.F., Izard, J., Segata, N., Gevers, D., Raes, J., and Huttenhower, C.
594   (2012) Microbial Co-occurrence Relationships in the Human Microbiome. *PLOS*
595   *Computational Biology* **8**: e1002606.
596
597 Fernandes, A.D., Reid, J.N., Macklaim, J.M., McMurrough, T.A., Edgell, D.R., and Gloor, G.B.
598   (2014) Unifying the analysis of high-throughput sequencing datasets: characterizing
599   RNA-seq, 16S rRNA gene sequencing and selective growth experiments by
600   compositional data analysis. *Microbiome* **2**: 15.
601
602 Ferris, M.J., Kühl, M., Wieland, A., and Ward, D.M. (2003) Cyanobacterial Ecotypes in
603   Different Optical Microenvironments of a 68°C Hot Spring Mat Community Revealed by
604   16S-23S rRNA Internal Transcribed Spacer Region Variation. *Appl Environ Microbiol*
605   **69**: 2893–2898.
606
607 Fierer, N. and Jackson, R.B. (2006) The diversity and biogeography of soil bacterial
608   communities. *Proc Natl Acad Sci USA* **103**: 626–631.
609
610 Friedman, J. and Alm, E.J. (2012) Inferring Correlation Networks from Genomic Survey Data.
611   *PLOS Computational Biology* **8**: e1002687.
612
613 Fulbright, S.P., Robbins-Pianka, A., Berg-Lyons, D., Knight, R., Reardon, K.F., Chisholm, S.T.
614   (2018) Bacterial community changes in an industrial algae production system. *Algal Res*
615   **31**:147–156.
616
617 Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010)
618   New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
619   performance of PhyML 3.0. *Syst Biol* **59**: 307–321.
620

621 Green, J.L., Bohannan, B.J.M., and Whitaker, R.J. (2008) Microbial Biogeography: From
622   Taxonomy to Traits. *Science* **320**: 1039–1043.
623

624 Guedes, I.A., Rachid, C.T.C.C., Rangel, L.M., Silva, L.H.S., Bisch, P.M., Azevedo, S.M.F.O.,
625   and Pacheco, A.B.F. (2018) Close Link Between Harmful Cyanobacterial Dominance and
626   Associated Bacterioplankton in a Tropical Eutrophic Reservoir. *Front Microbiol* **9**: 424.
627

628 Horner-Devine, M.C. and Bohannan, B.J.M. (2006) Phylogenetic Clustering and Overdispersion
629   in Bacterial Communities. *Ecology* **87**: S100–S108.
630

631 Hui, F.K.C., Taskinen, S., Pledger, S., Foster, S.D., and Warton, D.I. (2015) Model-based
632   approaches to unconstrained ordination. *Methods in Ecology and Evolution* **6**: 399–411.
633

634 Hutchinson, G.E. (1957) Concluding Remarks. *Cold Spring Harb Symp Quant Biol* **22**: 415–427.
635

636 Hutchinson, G.E. (1961) The Paradox of the Plankton. *The American Naturalist* **95**: 137–145.
637

638 Kamneva, O.K. (2017) Genome composition and phylogeny of microbes predict their co-
639   occurrence in the environment. *PLOS Computational Biology* **13**: e1005366.
640

641 Katoh, K. and Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7:
642 improvements in performance and usability. Mol Biol Evol **30**: 772–780.
643

644 Koeppel, A.F. and Wu, M. (2014) Species matter: the role of competition in the assembly of
645   congeneric bacteria. *ISME J* **8**: 531–540.
646

647 Koenker, R. (2015). Package 'quantreg'
648

649 Koenker, R., Machado, J.A.F. (1999). Goodness of fit and related inference processes for quantile
650   regression. Journal of the American Statistical Association. **94**, 1296-1310.
651

652 Kraft, N. J. B., Adler, P. B., Godoy, O., James, E. C., Fuller, S., and Levine, J. M. (2015).
653   Community assembly, coexistence and the environmental filtering metaphor. Funct. Ecol.
654   **29**, 592–599.
655

656 Kumar, S., Stecher, G., and Tamura, K. (2016) MEGA7: Molecular Evolutionary Genetics
657   Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* **33**: 1870–1874.
658

659 Kurmayer, R., Christiansen, G., and Chorus, I. (2003) The Abundance of Microcystin-Producing
660   Genotypes Correlates Positively with Colony Size in Microcystis sp. and Determines Its
661   Microcystin Net Production in Lake Wannsee. *Appl Environ Microbiol* **69**: 787–795.
662

663 Larkin, A.A and Martiny, A.C. (2017) Microdiversity shapes the traits, niche space, and
664   biogeography of microbial taxa. Environmental Microbiology Reports **9**:55-70.
665

666 Letten, A.D., Keith, D.A., Tozer, M.G., and Hui, F.K.C. (2015) Fine-scale hydrological niche
667   differentiation through the lens of multi-species co-occurrence models. *Journal of*

668   *Ecology 103*: 1264–1275.
669

670   Magurran, A.E. and Henderson, P.A. (2003) Explaining the excess of rare species in natural
671   species abundance distributions. *Nature* **422**: 714.
672

673   Martiny, J.B.H., Jones, S.E., Lennon, J.T., and Martiny, A.C. (2015) Microbiomes in light of
674   traits: A phylogenetic perspective. *Science* **350**: aac9323.
675

676   Mori, J.F., Neu, T.R., Lu, S., Händel, M., Totsche, K.U., and Küsel, K. (2015) Iron encrustations
677   on filamentous algae colonized by Gallionella-related bacteria in a metal-polluted
678   freshwater stream. *Biogeosciences* **12**: 5277–5289.
679

680   Needham, D. M., Fuhrman, J. A. (2016) Pronounced daily succession of phytoplankton, archaea
681   and bacteria following a spring bloom. *Nature Microbiology*, **1**: 16005.
682

683   Newton, R.J., Jones, S.E., Eiler, A., McMahon, K.D., and Bertilsson, S. (2011) A guide to the
684   natural history of freshwater lake bacteria. *Microbiol Mol Biol Rev* **75**: 14–49.
685

686   Nusch, E.A. (1980) Comparison of different methods for chlorophyll and phaeopigment
687   determination. Arch. Hydrobiol. Beih. **14** : 14–36."
688

689   Pascual-García, A., Tamames, J., and Bastolla, U. (2014) Bacteria dialog with Santa Rosalia: Are
690   aggregations of cosmopolitan bacteria mainly explained by habitat filtering or by
691   ecological interactions? *BMC Microbiol* **14**: 284.
692

693   Preheim, S.P., Perrotta, A.R., Martin-Platero, A.M., Gupta, A., and Alm, E.J. (2013)
694   Distribution-based clustering: using ecology to refine the operational taxonomic unit.
695   *Appl Environ Microbiol* **79**: 6593–6603.
696

697   Price MN, Dehal PS, Arkin AP. (2009). FastTree: computing large minimum evolution trees with
698   profiles instead of a distance matrix. *Mol Biol Evol* **26**:1641–1650.
699

700   Rohwer, R.R., Hamilton, J.J., Newton, R.J., and McMahon, K.D. (2018) TaxAss: Leveraging a
701   Custom Freshwater Database Achieves Fine-Scale Taxonomic Resolution. *mSphere* **3**:
702   e00327-18.
703

704   Röttjers, L. and Faust, K. (2018) From hairballs to hypotheses–biological insights from microbial
705   networks. *FEMS Microbiol Rev* **42**: 761–780.
706

707   Scanlan, D.J. (2003) Physiological diversity and niche adaptation in marine Synechococcus. In,
708   *Advances in Microbial Physiology*. Academic Press, pp. 1–64.
709

710   Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., et al. (2009)
711   Introducing mothur: Open-Source, Platform-Independent, Community-Supported
712   Software for Describing and Comparing Microbial Communities. *Appl Environ Microbiol*
713   **75**: 7537–7541.
714

715 Steunou, A.-S., Bhaya, D., Bateson, M.M., Melendrez, M.C., Ward, D.M., Brecht, E., et al.
716   (2006) In situ analysis of nitrogen fixation and metabolic switching in unicellular
717   thermophilic cyanobacteria inhabiting hot spring microbial mats. Proc Natl Acad Sci USA
718   **103**: 2398–2403.
719

720 Stockner, J., Callieri, C., and Cronberg, G. (2002) Picoplankton and Other Non-Bloom-Forming
721   Cyanobacteria in Lakes. In, Whitton,B.A. and Potts,M. (eds), *The Ecology of*
722   *Cyanobacteria: Their Diversity in Time and Space*. Dordrecht: Springer Netherlands, pp.
723   195–231.
724

725 Tromas, N., Taranu, Z.E., Martin, B.D., Willis, A., Fortin, N., Greer, C.W., and Shapiro, B.J.
726   (2018) Niche Separation Increases With Genetic Distance Among Bloom-Forming
727   Cyanobacteria. *Front Microbiol* **9** : 438
728

729 Tromas, N., Fortin, N., Bedrani, L., Terrat, Y., Cardoso, P., Bird, D., et al. (2017) Characterising
730   and predicting cyanobacterial blooms in an 8-year amplicon sequencing time course. The
731   *ISME Journal* **11**: 1746–1763.
732

733 Vörös, L., Callieri, C., Balogh, K.V., and Bertoni, R. (1998) Freshwater picocyanobacteria along
734   a trophic gradient and light quality range. *Hydrobiologia* **369**: 117–125.
735

736 Warton, D.I., Blanchet, F.G., O'Hara, R.B., Ovaskainen, O., Taskinen, S., Walker, S.C., and Hui,
737   F.K.C. (2015) So Many Variables: Joint Modeling in Community Ecology. *Trends in*
738   *Ecology & Evolution* **30**: 766–779.
739

740 Webb, C.O., Ackerly, D.D., McPeek, M.A., and Donoghue, M.J. (2002) Phylogenies and
741   Community Ecology. *Annual Review of Ecology and Systematics* **33**: 475–505.
742

743 Zheng, Q., Wang, Y., Xie, R., Lang, A.S., Liu, Y., Lu, J., et al. (2018) Dynamics of
744   Heterotrophic Bacterial Assemblages within Synechococcus Cultures. *Appl Environ*
745   *Microbiol* **84**: e01517-17.
746
747

748 **Data availability**

749 Raw sequence data have been deposited NCBI GenBank under BioProject number

750 PRJNA544938.

752 **Conflict of Interest**

753 The authors declare no conflict of interest.

Acknowledgments

NT is funded by a project from the European Union's Horizon 2020 research and innovation

program under the Marie Sklodowska-Curie grant agreement No 656647. BJS was supported by

a Canada Research Chair and NSERC Discovery Grant. Sampling and samples processing were

supported by a grant from Furnas Centrais Hidroelétrica S.A. (Brazil) to AG. JSMP and DAP

received a scholarship from CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível

Superior). A sabbatical visit of AG to the Université de Montréal AG was supported by a CAPES

Senior Fellowship.