Inter-animal variability, experience-dependence and continuous dorsoventral tuning of integrative properties of stellate cells in the medial entorhinal cortex

Hugh Pastoll[1], Derek Garden[1], Ioannis Papastathopoulos[2,3], Gülşen Sürmeli[1], Matthew F. Nolan[1]

Affiliations: [1]Centre for Discovery Brain Sciences, University of Edinburgh, Edinburgh, United Kingdom. EH8 9XD. [2]School of Mathematics, Maxwell Institute and Centre for Statistics, University of Edinburgh, Edinburgh, United Kingdom. EH3 3FD, [3]The Alan Turing Institute, 96 Euston Road, London, NW1 2DB.

Abstract

Distinctions between cell types underpin organisational principles for nervous system function, but it is also clear that functional variation exists between neurons of the same type. This is exemplified by correspondence between grid cell spatial scales and synaptic integrative properties of stellate cells (SCs) in the medial entorhinal cortex. However, we know little about how this variability is structured either within or between individuals. To address this, we used ex-vivo patch-clamp recordings to determine integrative properties from many (up to 55) SCs per mouse. We find that the expression and topographic tuning of SC integrative properties varies between mice, is in part dependent on experience and, in contrast to modularity of grid cell spatial scales, has a continuous dorsoventral organisation. Our results constrain mechanisms for modular grid firing and provide evidence for basal and experience-dependent inter-animal phenotypic variability among neurons of the same type.

Introduction

The concept of cell types provides a general organising principle for understanding biological structures including the brain (Regev et al., 2017; Zeng and Sanes, 2017). The simplest conceptualisation of a neuronal cell type, as a population of phenotypically similar neurons with features that cluster around a single point, is extended by observations of variability in cell type features, suggesting that some neuronal cell types may be conceived as clustering along a line rather than a point in a feature space (Cembrowski and Menon, 2018; O'Donnell and Nolan, 2011)(Figure 1A). Correlations between the functional organisation of sensory, motor and cognitive circuits and electrophysiological properties of individual neuronal cell types suggest that this feature variability is important for nervous system computations (e.g. (Adamson et al.,

2002; Angelo et al., 2012; Fletcher and Williams, 2018; Garden et al., 2008; Giocomo et al., 2007; Kuba et al., 2005; O'Donnell and Nolan, 2011)). This within cell type variability has typically been deduced by combining electrophysiological data obtained from multiple animals. In contrast, the structure of variation within individual animals has received little attention. For example, intra-animal variability in electrophysiological phenotypes could result from modular or continuous distributions of cell properties (Figure 1B). In addition, investigations of invertebrate nervous systems identify inter-animal differences as a further source of variation between neurons of the same type (Goaillard et al., 2009), but it is not clear whether mammalian neurons exhibit similar phenotypic diversity (Figure 1B). Distinguishing these possibilities requires many more electrophysiological observations per animal than are obtained in typical studies.

Stellate cells in layer 2 (SCs) of the medial entorhinal cortex (MEC) provide a striking example of correspondence between functional organisation of neural circuits and variability of electrophysiological features within a single cell type. The MEC contains neurons that encode an animal's location through grid-like firing fields (Fyhn et al., 2004). The spatial scale of grid fields follows a dorsoventral organisation (Hafting et al., 2005), which is mirrored by a dorsoventral organization in key electrophysiological features of SCs (Boehlen et al., 2010; Dodson et al., 2011; Garden et al., 2008; Giocomo and Hasselmo, 2008a; Giocomo et al., 2007; Pastoll et al., 2012a). However, recordings of many grid cells within a single animal indicate that they are organised in discrete modules (Stensola et al., 2012), with each cell in a module having a similar grid scale and orientation (Barry et al., 2007; Gu et al., 2018; Stensola et al., 2012), and progressively more ventral modules composed of cells with wider grid spacing (Stensola et al., 2012). Studies that demonstrate dorsoventral organisation of integrative properties of SCs have so far relied on pooling of relatively few measurements per animal. As a result, it is unclear whether organisation of these cellular features is modular, as one might expect if these properties directly set the scale of the firing fields of individual grid cells (Giocomo et al., 2007), or if instead it follows a continuous dorsoventral organisation. The possibility of basal or experience-dependent population level differences between individual animals has also not previously been considered.

Assessment of variability between animals and the structure of variability within an animal requires statistical approaches not typically used in single-cell electrophysiological investigations. Given appropriate assumptions, inter-animal differences can be evaluated using mixed effect models that are well established in other fields (Baayen et al., 2008; Geiler-

Samerotte et al., 2013). Detection of data that arise from modular as opposed to continuous distributions has received less general attention. To facilitate detection of modularity using relatively few observations, we introduce a modification of the gap statistic algorithm (Tibshirani et al., 2001) that estimates the number of modes in a dataset while controlling for observations expected by chance (see Methods and Supplemental Figures 1 - 5). This algorithm performs well compared with other measures (Giocomo et al., 2014; Stensola et al., 2012), which we find are prone to high false positive rates (Supplemental Figure 4A). Using the modified gap statistic algorithm, we find that recordings from approximately 30 SCs per animal should be sufficient to detect modularity given the experimentally observed separation between grid modules (see Methods and Supplemental Figures 2-3). Although methods for high quality recording from SCs in ex-vivo brain slices are well established (Pastoll et al., 2012b), in previous studies typically fewer than 5 recordings per animal have been made, which is much less than our estimate of the minimum number of observations required to test for modularity.

To investigate intra- and inter-animal variation in electrophysiological features of SCs we therefore carried out experiments that maximise the number of SCs recorded per animal. Using these data we establish the relative dependence of 12 electrophysiologically measured integrative properties on dorsoventral position, show that dorsoventral tuning is much stronger in SCs compared to nearby pyramidal cells, and demonstrate substantial variability between individual animals. To test whether electrophysiological features are affected by experience we compared mice housed in standard home cages with mice housed in large environments. We find that the size of housing is associated with differences in specific electrophysiological features, but that substantial inter-animal differences remain after accounting for housing and other experimental factors. Evaluation of modularity indicates that the integrative properties of SCs follow a continuous rather than a modular dorsoventral organisation. By dissociating the organisation of electrophysiological features of SCs from that of grid cell firing fields our results impose new constraints on possible mechanisms for generation of spatial codes. Our evidence for inter-animal variability suggests that tuning of electrophysiological phenotypes in part depends on experience and in part involves set points at the level of neuronal populations.

Results

Sampling integrative properties from many neurons per animal

Before addressing intra- and inter-animal variability we first describe the data set used for the analyses that follow. We established procedures to facilitate recording of integrative properties of many SCs from a single animal (see Methods). With these procedures, we measured and analysed electrophysiological features of 836 SCs (n/mouse: range 11-55; median = 35) from 27 mice (median age = 37.4 days, age range = 18 - 57 days). The mice were housed either in a standard home cage (N = 18 mice, n = 583 neurons) or from postnatal day 16 in a 2.4 x 1.2 m cage, which provided a large environment that could be freely explored (N = 9, n = 253, median age = 38 days)(Supplemental Figure 6). For each neuron we assessed six measures of its sub-threshold integrative properties (Figure 2A-B) and six measures of its supra-threshold integrative properties (Figure 2C). Until indicated otherwise we report analysis of datasets that combine the groups of mice housed in standard and large home cages and that span the full range of ages.

Because SCs are found intermingled with pyramidal cells in layer 2 (L2PCs), and as misclassification of L2PCs as SCs would likely confound investigation of intra-animal variation, we validated our criteria for distinguishing each cell type. To establish characteristic electrophysiological properties of L2PCs we recorded from neurons in layer 2 identified by Cre-dependent marker expression in a Wfs1^{Cre} mouse line (Sürmeli et al., 2015). Expression of Cre in this line, and a similar line (Kitamura et al., 2014), labels pyramidal cells in layer 2 (L2PCs) that project to the CA1 region of the hippocampus, but does not label SCs (Kitamura et al., 2014; Sürmeli et al., 2015). We identified two populations of neurons in layer 2 of MEC that were labelled in Wfs1^{Cre} mice (Figure 3A-C). The more numerous population had properties consistent with L2PCs (Figure 3A, G) and could be separated from the unidentified population on the basis of a lower rheobase (Figure 3C). The unidentified population had firing properties typical of layer 2 interneurons (cf. (Gonzalez-Sulser et al., 2014)). A principal component analysis (PCA)(Figure 3D-F) clearly separated the L2PC population from the SC population, but did not identify subpopulations of SCs. The less numerous population has properties similar to those of inhibitory interneurons and also clearly distinct from SCs (Figure 3A, C). These data demonstrate that the SC population used for our analyses is distinct from excitatory pyramidal cells also found in layer 2 of the MEC.

To further validate the large SC dataset we assessed the location-dependence of individual features, several of which have previously been found to depend on the dorso-ventral location of the recorded neuron (Boehlen et al., 2010; Booth et al., 2016; Garden et al., 2008; Giocomo

et al., 2007; Pastoll et al., 2012a). We initially fit the dependence of each feature on dorsoventral position using a standard linear regression model. We found substantial (adjusted $R^2 > 0.1$) dorsoventral gradients in input resistance, sag, membrane time constant, resonant frequency, rheobase and the current-frequency (I-F) relationship (Table 1, Figure 3G). In contrast to SCs, we did not find evidence for dorsoventral organisation of these features in L2PCs (Figure 3G). Thus, our large dataset replicates the previously observed dependence of integrative properties of SCs on their dorsoventral position, and shows that this location dependence further distinguishes SCs from L2PCs.

Inter-animal differences in intrinsic properties of stellate cells

To what extent does variability between the integrative properties of SCs at a given dorsoventral location (Figure 3G) arise from differences between animals? Comparing specific features from individual animals suggested that their distributions could be almost completely non-overlapping despite consistent and strong dorsoventral tuning (Figure 4A). If this apparent inter-animal variability results from random sampling of a distribution determined by a common underlying set point, then fitting the complete data set with a mixed model in which animal identity is included as a random effect should reconcile the apparent differences between animals (Figure 4B). In this scenario, the conditional R^2 estimated from the mixed effect model, in other words the estimate of variance explained by animal identity and location, should be similar to the marginal R^2 value, which indicates the variance explained by location only. In contrast, if differences between animals contribute to experimental variability, the mixed model should predict different fitting parameters for each animal, and the estimated conditional R^2 should be greater than the corresponding marginal R^2 (Figure 4C).

Fitting the experimental measures for each feature with mixed models suggests that differences between animals contribute substantially to the variability in properties of SCs. In contrast to simulated data in which inter-animal differences are absent (Figure 4B), differences in fits between animals remained after fitting with the mixed model (Figure 4D). This corresponds well with expectations from fits to simulated data containing inter-animal variability (cf. Figures 4C). To visualise inter-animal variability for all measured features we plot for each animal the intercept of the model fit (I), the predicted value at a location 1 mm ventral from the intercept (I+S) and the slope (lines)(Figure 4E). Strikingly, even for features such as rheobase and input resistance (IR) that are highly tuned to a neurons' dorsoventral position, the extent of variability

between animals is similar to the degree to which the property changes between dorsal and mid-levels of the MEC.

If set points that determine integrative properties of SCs do indeed differ between animals, then mixed models should provide a better account of the data than linear models generated by pooling data across all animals. Consistent with this, we found that mixed models for all electrophysiological features gave a substantially better fit to the data than linear models that considered all neurons as independent (adjusted $p < 2$ x 10^{-17} for all models, χ^2 test, Table 1). Furthermore, even for properties with substantial (R^2 value > 0.1) dorsoventral tuning, the conditional R^2 value for the mixed effect model was substantially larger than the marginal R^2 value (Figure 4D and Table 1). Together, these analyses demonstrate substantial inter-animal variability in all of the measured electrophysiological features of SCs suggesting that set points that establish integrative properties of SCs differ between animals.

Electrophysiological features of stellate cells are modified by experience

Because experience-dependent modification of neuronal integrative properties can be important for learning (Zhang and Linden, 2003), we asked if experience influences the electrophysiological features of SCs. Since entorhinal neurons represent location, we reasoned that modifying the space through which animals can navigate may drive experience-dependent plasticity. In particular, as standard mouse housing has dimensions less than the distance between firing fields of more ventrally located grid cells (Brun et al., 2008; Hafting et al., 2005), we were interested to know whether electrophysiological features of SCs that follow a dorsoventral organisation differ between mice housed in larger environments (28800 cm^2) compared with standard home cages (530 cm^2).

We compared the mixed models described above to models in which housing was also included as a fixed effect. To minimise the effects of age on SCs (Boehlen et al., 2010; Burton et al., 2008)(Supplemental Table 2), we focussed these and subsequent analyses on mice between P33 and P44 (N = 25, n = 779). We found that larger housing was associated with a smaller sag coefficient indicating an increased sag response, a lower resonant frequency and a larger spike half-width (adjusted $p < 0.05$; Figure 4E, Supplemental Table 3, Supplemental analyses). These differences were primarily from changes to the magnitude rather than the location-dependence of each feature. Other electrophysiological features appeared unaffected by housing.

Given the effect of housing on several electrophysiological features of SCs, we asked if inter-animal differences in these or other features remain after accounting for housing. To address this we compared mixed models that include dorsoventral location and housing as fixed effects with equivalent linear models in which individual animals are not accounted for. Mixed models incorporating animal identity continued to provide a better account of the data, both for features that were dependent on housing (adjusted $p < 2.8 \times 10^{-21}$) and for features that were not (adjusted $p < 1.4 \times 10^{-7}$)(Supplemental Table 4).

Together, these data suggest that experience of large environments modifies specific electrophysiological features of SCs. After accounting for environment significant inter-animal variation remains, suggesting that additional mechanisms acting at the level of animals rather than individual neurons also determine differences between SCs.

Inter-animal differences remain after accounting for additional experimental parameters
To evaluate whether factors other than housing can account for inter-animal differences between SCs, we assessed potential effects of additional experimental variables. We found that the measured electrophysiological features of SCs were largely independent of brain hemisphere (Supplemental Table 5), but that properties primarily related to the action potential waveform depended on mediolateral position of the recorded neuron (Supplemental Table 6)(cf. (Canto and Witter, 2012)). We also found that the identity of the experimenter (Supplemental Table 7) and the time since slice preparation (Supplemental Tables 8 and 9) introduce a degree of systematic variability into measurements of electrophysiological features of SCs, but overall these features and their dorsoventral organisation are robust to these potential sources of technical variation. Moreover, our data collection minimised possible effects of these parameters when evaluating the influence of housing, for example by counterbalancing the dorsoventral direction along which recordings were made (see Methods). Nevertheless, to address the possibility that these, or additional factors, account for inter-animal variation, we carried out several further tests.

First, we evaluated models that incorporate housing, mediolateral position, experimenter identity and the direction in which sequential recordings were obtained as fixed effects. When we compared versions of these models that contained animal identity as a random effect with otherwise equivalent linear models, we again found highly significant differences (adjusted p <

10^{-8}) indicating that the additional factors are unable to fully account for inter-animal differences (Supplemental Table 10).

Second, we evaluated minimal datasets from each experimenter in which housing, mediolateral position and the recording direction were identical. Because of the smaller size of these datasets the statistical power to detect inter-animal variation is reduced. Nevertheless, in these analyses the conditional R^2 of the mixed model fit was again substantially higher than the marginal R^2, and most (9/12) features were better fit by a mixed model compared to a corresponding linear model in both datasets (Supplemental Table 11).

We also considered whether variation in tissue quality as a contributor to inter-animal variability. To minimise this possibility we standardised our procedures for tissue preparation (see Methods), such that slices were of consistent high quality as assessed by low numbers of unhealthy cells and by visualisation of soma and dendrites of neurons in the slice. We further reasoned that if the condition of the slices differed between animals, then in better quality slices it would be easier to record from more neurons, in which case any features that depend on tissue quality would correlate with the number of recorded neurons. We found that the majority (10/12) of electrophysiological features were not significantly ($p > 0.2$) associated with the number of recorded neurons (Supplemental Table 12). Moreover, when models were fit to data from animals from which > 35 recordings were made (N = 11, n = 459), comparison of marginal and conditional R^2 values continued to indicate substantial inter-animal variance, and fits obtained with mixed models remained significantly different to fits that did not account for animal identity ($p < 4.4 \times 10^{-5}$)(Supplemental Table 13).

Together, these analyses indicate that differences between animals remain after accounting for experimental and technical factors that might contribute to variation in measured features of SCs. We also note that conditional R^2 values of electrophysiological features of L2PCs are much lower than for SCs (cf. Table 1 and Supplemental Table 1) suggesting that inter-animal variation may be specific to SCs, and further arguing that inter-animal differences result from biological rather than technical variability.

The distribution of intrinsic properties is consistent with a continuous rather than a modular organisation

To evaluate modularity of intrinsic properties we used datasets with n ≥ 34 (N = 15, median age = 37 days, age range = 18 - 43 days). We focus initially on rheobase, which is the property with strongest correlation to dorsoventral location, and resonant frequency, which is related to the oscillatory dynamics underlying dorsoventral tuning in some models of grid firing. For n ≥ 34 we expect that if properties are modular then this would be detected by the modified gap statistic in at least 50% of animals (Supplemental Figures 2C, 3). In contrast, we find that for datasets from the majority of animals the modified gap statistic identifies only a single mode in the distribution of rheobase values (Figure 5A and Figure 6)(N = 13/15) and of resonant frequencies (Figure 5B and Figure 6) (N = 14/15), indicating that these properties have a continuous rather than a modular distribution. Consistent with this, smoothed distributions did not show clearly separated peaks for either property (Figure 5). The mean and 95% confidence interval for p_{detect}, the probability of evaluating a dataset as clustered, was 0.133 and 0.02–0.4 for rheobase and 0.067 and 0.002–0.32 for resonant frequency. These values of p_{detect} were not significantly different from the proportions expected from the complete absence of clustering (p = 0.28 and 0.66, binomial test). Thus, rheobase and resonant frequency of SCs, while depending strongly on a neuron's dorsoventral position, do not have a detectable modular organisation.

When we investigated the other measured integrative properties we also failed to find evidence for modularity. Across all properties, for any given property at most 3 out of 15 mice were evaluated as exhibiting clustered organisation using the modified gap statistic (Figure 6). This does not differ significantly from the proportion expected by chance when no modularity is present (p > 0.05, binomial test). Consistent with this the average proportion of datasets evaluated as modular across all measured features was 0.072 ± 0.02 (± SEM), which is similar to the expected false positive rate. In contrast, for the 7 grid datasets with n ≥ 20 considered in Supplemental Figure 5, the mean for p_{detect} is 0.86, with 95 % confidence intervals of 0.42 to 0.996. We note here that previously published discontinuity algorithms did indicate significant modularity in the majority of the intrinsic properties measured in our dataset (N = 13/15 and N = 12/15 respectively), but this was attributable to false positives resulting from the relatively even sampling of recording locations (see Supplemental Figure 4A). Therefore, we conclude that it is unlikely that within individual animals any of the intrinsic properties we examined have organisation resembling the modular organisation of grid cells in behaving animals.

Multiple sources of variance contribute to diversity in stellate cell intrinsic properties

Finally, we considered potential relationships between the measured electrophysiological features. Estimation of conditional independence for measurements at the level of individual neurons revealed strong partial correlations between many of the features (Figure 7A). At the level of individual animals, which we assessed using the intercept of fits of mixed models to the data, we found a smaller set of strong partial correlations (Figure 7B). The conditional independence structure indicated by the partial correlation matrices is consistent with the expectation that particular membrane ion channels will influence multiple electrophysiologically measured features. For example, conditionally on all remaining measured features, rheobase was negatively correlated with input resistance and membrane potential, and in SCs all three features are strongly influenced by HCN and leak potassium channels (Garden et al., 2008).

Given the strong partial correlations between electrophysiological features, we wanted to know whether dorsoventral variation found for many of these features can be reduced to a smaller number of dimensions in feature space, as we would expect if dorsoventral variation results from common core mechanisms. To address this we performed a principal components analysis of all measured electrophysiological features. To remove variance caused by animal age and the experimenter identity, we used a reduced dataset obtained by one experimenter and restricted to animals between 32 and 45 days old (N = 25, n = 572). The first principal component accounted for almost 30% of the variance in the measured electrophysiological features, and the first 5 components accounted for almost 80% of the variance (Figure 7C). Examination of the rotations used to generate the principal components suggested relationships between individual features that are consistent with our evaluation of the conditional independence structure of the measured features (cf. Figure 7D and 7A). When we fit the principal components using mixed models with location as a fixed effect and animal identity as a random effect, we found that the first two components depended significantly on dorsoventral location (Figure 7E and Supplemental Table 14)(marginal R^2 = 0.50 and 0.09 and adjusted p = 1.09 x 10^{-15} and 1.05 x 10^{-4} respectively). These observations suggest that that the dependence of multiple electrophysiological features on dorsoventral position may be reducible to two core mechanisms that together account for much of the variability between SCs in their intrinsic electrophysiology.

We conclude by asking whether experience-dependence and inter-animal variation are present in PCA dimensions that account for dorsoventral variation, and whether they extend to additional dimensions in feature space. We found that the intercept, but not the slope, of the

dependence of the first two principal components on dorsoventral position depended on housing (adjusted p = 0.039 and 0.027)(Figure 7E, F and Supplemental Table 15). After accounting for housing, the first two principal components were still better fit by models that include animal identity as a random effect (adjusted p = 3.3 x 10^{-9} and 4.1 x 10^{-86}), indicating remaining inter-animal differences in these components (Supplemental Table 16). Nine of the next ten higher order principal components did not depend significantly on housing (adjusted p > 0.1)(Supplemental Table 15), while 8 differed significantly between animals (adjusted p < 0.05)(Supplemental Table 16).

Together, these analyses indicate that dorsoventral organisation of multiple electrophysiological features of SCs reflect two main sources of variance, both of which are dependent on experience. Significant inter-animal variation in the major sources of variance remains after accounting for experience and experimental parameters.

Discussion

Phenotypic variation is well established across many areas of biology (Geiler-Samerotte et al., 2013), but has received relatively little attention in investigations of mammalian nervous systems. Our analysis indicates that within animal variation in functional properties of SCs is continuous, and suggests substantial and unexpected inter-animal variability in SC properties that in part depends on experience. These results identify new constraints on models for grid cell firing, indicate the importance of accounting for animal identity in experimental design and suggest that set points for electrophysiological features are influenced by circuit level mechanisms whose activity differs between animals.

Continuous dorsoventral organisation of stellate cell integrative properties
Dorsoventral gradients in electrophysiological features of SCs have stimulated cellular models for the organisation of grid firing (Burgess, 2008; Giocomo and Hasselmo, 2008b; Grossberg and Pilly, 2012; O'Donnell and Nolan, 2011; Widloski and Fiete, 2014). Increases in spatial scale following deletion of HCN1 channels (Giocomo et al., 2011), which in part determine the dorsoventral organisation of SC integrative properties (Garden et al., 2008; Giocomo and Hasselmo, 2009), support a relationship between electrophysiological properties of SCs and grid cell spatial scales. Our data argue against models that explain this relationship through

single cell computations (Burgess, 2008; Giocomo et al., 2007), as in this case modularity of integrative properties of SCs is required to generate modularity of grid firing. A continuous dorsoventral organisation of electrophysiological properties of SCs could support modular grid firing generated by self-organising maps (Grossberg and Pilly, 2012), or by synaptic learning mechanisms (Kropff and Treves, 2008; Urdapilleta et al., 2017). It is less clear how a continuous gradient would affect predictions of grid firing from continuous attractor network models (Shipston-Sharman et al., 2016; Widloski and Fiete, 2014), which can account for modularity by limiting synaptic interactions between modules. In continuous attractor models modularity can instead be achieved by restricting synaptic connectivity between modules.

The continuous variation of electrophysiological features of SCs suggested by our analysis is consistent with continuous gradients in gene expression along layer 2 of the MEC (Ramsden et al., 2015). It is also consistent with single cell RNA sequencing analysis of other brain areas, which indicates that while expression profiles for some cell types cluster around a point in feature space, others lie along a continuum (Cembrowski and Menon, 2018). It will be interesting in future to determine whether gene expression continua establish corresponding continua of electrophysiological features (cf. (Liss et al., 2001)).

Inter-animal variation and experience-dependence of neuronal membrane properties
In invertebrate nervous systems, for which individual neurons are identifiable, striking inter-animal variability in key electrophysiological properties has been demonstrated (Goaillard et al., 2009). Mammalian nervous systems contain many more neurons, which are not individually identifiable, but that can be grouped into neuronal cell types. From a developmental perspective in which cell identity is considered as a trough in a state-landscape through which each cell moves (Wang et al., 2011b), variation in the population of neurons of the same type could be conceived as noisy deviations from the trough (Figure 8A). From this perspective, our finding that variability among neurons of the same type manifests between as well as within animals, could be explained by differences between animals in the position of the trough in the developmental landscape (Figure 8B).

What factors contribute to variability between animals? We find that a subset of electrophysiological properties of SCs are sensitive to the size of the environment in which an animal is housed. The reduction in the sag coefficient in larger environments could be explained by an increase in HCN channel mediated hyperpolarization-activated currents (I_h) in SCs (cf.

(Nolan et al., 2007)). This is consistent with activity-dependent increases in I_h in other neuron types (Fan et al., 2005; Narayanan and Johnston, 2007). Since HCN1 channels affect spatial memory and grid codes (Giocomo et al., 2011; Nolan et al., 2004), these changes could mediate a behavioural adaptation to the larger environment. However, substantial inter-animal differences in electrophysiological features remain after accounting for environment and for multiple biological and technical sources of variation (Supplemental Tables 4-16). An important future goal will be to distinguish intrinsic from further possible extrinsic causes of variation between animals. For example, social interactions may affect brain circuitry (Wang et al., 2011a, 2014), although these effects appear focussed on frontal cortical structures rather than circuits for spatial computations (Wang et al., 2014). Alternatively, stochastic mechanisms operating at the population level may drive emergence of inter-animal differences during development of SCs (Donato et al., 2017; Ray and Brecht, 2016).

What are the functional roles of inter-animal variability? In the stomatogastric ganglion, inter-animal variation correlates with circuit performance (Goaillard et al., 2009). Accordingly, variation in intrinsic properties of SCs might correlate with differences in grid firing (Domnisoru et al., 2013; Gu et al., 2018; Rowland et al., 2018; Schmidt-Hieber and Häusser, 2013) or behaviours that rely on SCs (Kitamura et al., 2014; Qin et al., 2018; Tennant et al., 2018). Modification of grid field scaling following deletion of HCN1 channels is consistent with this possibility (Giocomo et al., 2011; Mallory et al., 2018). Alternatively, inter-animal differences may reflect multiple ways to achieve a common higher order phenotype. According to this view, coding of spatial location by SCs would not differ between animals despite lower level variation in intrinsic electrophysiological features. This is related to the idea of degeneracy at the level of single cell electrophysiological properties (Marder and Goaillard, 2006; Mittal and Narayanan, 2018; O'Leary et al., 2014; Swensen and Bean, 2005), except that here the electrophysiological features differ between animals but higher order circuit computations may nevertheless be similar.

In conclusion, our results suggest that variation operating at the level of neural circuits determines inter-animal differences in neuronal integrative properties. This has implications for experimental design and model building as the distribution of replicates from the same animal will differ from those obtained from different animals (Marder and Taylor, 2011). It also suggests new questions about circuit development; for example, what is the origin of variability acting at the level of the circuit and how does this interact with cell autonomous variation arising from

stochasticity of gene expression and with experience-dependent mechanisms? Addressing these questions may turn out to be critical to understanding how cellular biophysics determines circuit level computations in cognitive circuits.

Acknowledgements

We thank Vanessa Stempel for comments on the manuscript. This work was supported by grants to MN from the Wellcome Trust (200855/Z/16/Z) and the BBSRC (BB/L010496/1, BB/1022147/1 and BB/H020284/1).

Author contributions

HP and MFN conceived of the study. HP and DG performed experiments. HP and MFN analysed the data. IP contributed to statistical analyses. GS assisted with experiments and contributed reagents. MFN obtained funding, supervised the study and wrote the paper. All authors contributed to review and editing of the manuscript.

Declaration of interests

The authors declare no competing interests.

Figure 1. Classification and variability of neuronal cell types

(A) Neuronal cell types are identifiable by features clustering around a distinct point (blue, green and yellow) or a line (red) in feature space.

(B) Phenotypic variability of a single neuronal type may only be apparent by measuring features from multiple neurons from multiple individual animals. Examples of variability not detectable with few recorded neurons per animal include modular organisation of a feature, which would be masked by inter-animal variability (upper), variability between animals in features that may be orthogonal to features varying within an animal (middle) and inter-animal differences in the offset of the range of a variable feature (lower).

Figure 2. Dorsoventral organization of intrinsic properties of stellate cells from a single animal

(A-C) Waveforms (grey traces) and example responses (black traces) from a single mouse for step (A), ZAP (B) and ramp (C) stimuli (left). Properties derived from each protocol are shown plotted against recording location (each data point is a black cross)(right). KSDs with arbitrary bandwidth are displayed to the right of each data plot to facilitate visualisation of the property's distribution when location information is disregarded (light grey pdfs). A. Injection of a series of current steps enables measurement of the resting membrane potential (Vrest) (i), the input resistance (IR) (ii), the sag coefficient (sag) (iii) and the membrane time constant (Tm) (iv).

(B) Injection of ZAP current waveform enables calculation of an impedance amplitude profile, which was used to estimate the resonance resonant frequency (Res. F) (i) and magnitude (Res. mag) (ii).

(C) Injection of a slow current injection ramp enabled measurement of the rheobase (i), the slope of the current-frequency relationship (I-F slope) (ii), and using only the first 5 spikes in each response (enlarged zoom, lower left) the AHP minimum value (AHP$_{min}$) (iii), the spike maximum (Spk. max) (iv) , the spike width at half height (Spk. HW) (v) and the spike threshold (Spk. thr.) (vi).

Figure 3. Distinct and dorsoventrally organised properties of layer 2 stellate cells

(A) Representative action potential afterhyperpolarization waveforms from a SC (left), a pyramidal cell (middle) and an unidentified cell (right). The pyramidal and unidentified cells were both positively labelled in Wfs1^{Cre} mice.

(B) Plot of the first versus the second principal component from PCA of the properties of labelled neurons in Wfs1^{Cre} mice reveals two populations of neurons.

(C) Histogram showing the distribution of rheobase values of cells positively labelled in Wfs1^{Cre} mice. The two groups identified in B can be distinguished by their rheobase.

(D) Plot of the first two principal components from PCA of the properties of the L2PC (n = 44, green) and SC populations (n = 840, black). Putative pyramidal cells (x) and SCs (+) are colored according to their dorsoventral location (inset shows the scale).

(E) Proportion of total variance explained by the first five principal components for the analysis in (D).

(F) Histograms of locations of recorded SCs (upper) and L2PCs (lower).

(G) All values of measured features from all mice are plotted as a function of the dorsoventral location of the recorded cells. Lines indicate fits of a linear model to the complete datasets for SCs (black) and L2PCs (green). Putative pyramidal cells (x, green) and SCs (+, black). Adjusted R^2 values use the same colour scheme.

Figure 4. Inter-animal variability and dependence on environment of intrinsic properties of stellate cells

(A) Examples of rheobase as a function of dorsoventral position for two mice.

(B-C) Each line is the fit of simulated data from a different subject for datasets in which there is no inter-subject variability (B) or in which intersubject variability is present (C). Fitting data from each subject independently with linear regression models suggests intersubject variation in both datasets (left). In contrast, after fitting mixed effect models (right) intersubject variation is no longer suggested for dataset in which it is absent (B) but remains for the dataset in which it is present (C).

(D) Each line is the fit of rheobase as a function of dorsoventral location for a single mouse. The fits were carried out independently for each mouse (left) or using a mixed effect model with mouse identity as a random effect (right).

(E) The intercept (I), sum of the intercept and slope (I + S), and slopes (right) for each mouse obtained by fitting mixed effect models for each property as a function of dorsoventral position.

Figure 5. Input resistance and resonant frequency do not have a detectable modular organisation

(A-B) Input resistance (A) and resonant frequency (B) are plotted as a function of dorsoventral position separately for each every animal. Marker colour and types indicate the age and housing

conditions of the animal ('+'s standard housing, 'x's large housing). KSDs (arbitrary bandwidth, same for all animals) are also shown. The number of clusters in the data (k_{est}) is indicated for each animal.

Figure 6. Significant modularity is not detectable for any measured property

(A-B) Histograms showing the k_{est} counts across all mice for each different measured sub-threshold (A) and supra-threshold (B) intrinsic properties. The maximum k evaluated was 8. P values from binomial tests (with N = 15) indicate that for every measured property, the number datasets evaluated as clustered is significantly less than expected if the underlying distribution of that property is always clustered with all separation between modes ≥ 5 std (pink text). Similarly, p-values from binomial tests show that no property has a k_{est} count greater than that expected by chance if the underlying distribution of the property is uniform (purple text).

Figure 7. Feature relationships and inter-animal variability after reducing dimensionality of the data

(A-B) Partial correlations between the electrophysiological features investigated at the level of individual neurons (A) and at the level of animals (B). Partial correlations outside the 95% confidence intervals generated from a bootstrapped dataset are colour coded. Non-significant correlations are coloured white.

(C) Proportion of variance explained by each principal component.

(D) Loading plot for the first two principal components.

(E) The first five principal components plotted as a function of position.

(F) Intercept (I), intercept plus the slope (I + S) and slopes aligned to the same intercept, for fits for each animal of the first five principal components to a mixed model with location as a fixed effect and animal as a random effect.

Figure 8. Models for intra- and inter-animal variation
(A) The configuration of a cell type can be conceived of as a trough in a developmental landscape (solid line). In this scheme, the developmental landscape is established at the level of individual animals and differences between cells (filled circles) reflect noisy variation away from the trough.
(B) In this framework differences between animals (red and blue) can be conceptualised as different troughs in the developmental landscape.

Feature	Slope	p (slope)	Marginal R2	Conditional R2	Slope (min)	Slope (max)	p (vs linear)
Vm (mV)	-0.8842	3.66e-05	0.0458	0.3478	-1.9646	0.1495	8.73e-35
IR (MΩ)	11.7939	8.40e-17	0.3833	0.5321	9.6294	14.2622	4.33e-40
Sag	0.0327	6.06e-10	0.1211	0.3470	0.0163	0.0432	1.91e-38
Tm (ms)	2.6458	3.70e-12	0.1925	0.3430	1.8095	3.9786	1.20e-29
Res. frequency (Hz)	-1.3341	4.13e-09	0.1223	0.5525	-2.2988	-0.3424	6.37e-65
Res. magnitude	-0.1142	6.35e-08	0.0636	0.1985	-0.1381	-0.0872	9.13e-20
Spike thresold (mV)	0.0823	8.20e-01	0.0001	0.5097	-2.4677	2.3797	2.03e-17
Spike maximum (mV)	1.8997	1.85e-05	0.0642	0.4365	-1.2875	3.2971	1.14e-50
Spike width (ms)	0.0167	1.93e-02	0.0097	0.6427	-0.0206	0.0553	7.04e-139
Rheobase (pA)	-119.8845	9.08e-15	0.3818	0.6522	-153.8736	-76.1257	6.55e-43
Spike AHP (mV)	-0.6448	1.93e-02	0.0115	0.2571	-1.8280	0.4084	1.82e-17
I-F slope (Hz/pA	0.0364	6.06e-10	0.2282	0.5609	0.0188	0.0872	6.82e-34

Table 1. Dependence of electrophysiological features of SCs on dorsoventral position
Estimates of the population slope and corresponding p value, marginal and conditional R^2, and the minimum and maximum slopes across all mice, obtained by fitting the dependence of each feature on dorsoventral position using mixed effect models. The data are from 27 mice. Estimates of significance of the slope were obtained using a χ^2 test to compare models that include location as a fixed effect with models in which location was removed. Estimates of significance for comparison of the mixed with a corresponding linear model without random effects were obtained using a χ^2 test. Significance estimates were adjusted for multiple comparisons using the method of Benjamini and Hochberg (Benjamini and Hochberg, 1995). Units for the slope measurements are units for the property mm^{-1}.

Methods

Mouse strains. All experimental procedures were performed under a United Kingdom Home Office license and with approval of the University of Edinburgh's animal welfare committee. Recordings of many SCs per animal used male C57/Bl6J mice (Charles River). Recordings targeting calbindin cells used a Wfs1[Cre] line (Wfs1-Tg3-CreERT2) obtained from Jackson Labs (Strain name: B6;C3-Tg(Wfs1-cre/ERT2)3Aibs/J; stock number:009103). Mice were group housed on a 12 h light/dark cycle (light on 07.30–19.30 h). Mice were usually housed in standard breeding cages, with a subset of mice instead housed from pre-weaning ages in a larger 2.4 x 1.2 m cage that was enriched with bright plastic objects (Supplemental Figure 6). At weaning the mother was removed from the large cage.

Viral injections. To label Cre expressing neurons in Wfs1[Cre] heterozygous mice we stereotaxically injected AAV-FLEX-GFP, generated with a chimeric 1/2 serotype, which expresses GFP from a CBA promoter (Murray et al., 2011)(titer: 1.5×1012 cp/ml) as described in Surmeli et al., 2015.

Slice preparation. Methods for preparation of parasagittal brain slices containing medial entorhinal cortex were based on procedures described previously (Pastoll et al., 2012b). Briefly, mice were sacrificed by cervical dislocation and their brains carefully removed and placed in cold (2-4 °C) modified ACSF, with composition (in mM): NaCl 86, NaH_2PO_4 1.2, KCl 2.5, $NaHCO_3$ 25, Glucose 25, Sucrose 75, $CaCl_2$ 0.5, $MgCl_2$ 7. Brains were then hemisected and sectioned, also in modified ACSF at 4-8 °C, using a vibratome (Leica VT1200S). After cutting brains were placed at 36 °C for 30 minutes in standard ACSF, with composition (in mM): NaCl 124, NaH_2PO4 1.2, KCl 2.5, $NaHCO_3$ 25, Glucose 20, $CaCl_2$ 2, $MgCl_2$ 1. They were then allowed to cool passively to room temperature. All slices were prepared by the same experimenter (HP) following the same procedure on each day.

Recording methods. Whole-cell patch-clamp recordings were made between 1 to 10 hours after slice preparation using procedures described previously (Pastoll et al., 2012a, 2012b, 2013; Sürmeli et al., 2015). Patch electrodes were filled with the following intracellular solution (in mM): K Gluconate 130; KCl 10, HEPES 10, $MgCl_2$ 2, EGTA 0.1, Na_2ATP 2, Na_2GTP 0.3 NaPhosphocreatine 10. The open tip resistance was 4-5 MΩ, all seal resistances were > 2 GΩ and series resistance were < 30 MΩ. Recordings were made in current clamp mode using

Multiclamp 700B amplifiers (Molecular Devices, Sunnyvale, CA, USA) connected to PCs via Instrutech ITC-18 interfaces (HEKA Elektronik, Lambrecht, Germany) and using Axograph X acquisition software (http://axographx.com/). Pipette capacitance and series resistances were compensated using the capacitance neutralisation and bridge-balance amplifier controls. An experimentally measured liquid junction potential of 12.9 mV was not corrected for. Stellate cells were identified by their large sag response and the characteristic waveform of their action potential afterhyperpolarization (see (Alonso and Klink, 1993; Gonzalez-Sulser et al., 2014; Nolan et al., 2007; Pastoll et al., 2012a)).

To maximize the number of cells recorded per animal we adopted two strategies. First, to minimize the time required to obtain data from each recorded cell, we measured electrophysiological features using a series of three short protocols following initiation of stable whole-cell recordings. We used responses to sub-threshold current steps to estimate passive membrane properties (Figure 2A), a frequency modulated sinusoidal current waveform (ZAP waveform) to estimate impedance amplitude profiles (Figure 2B) and a linear current ramp to estimate the action potential threshold and firing properties (Figure 2C). From analysis of data obtained with these protocols we obtained 12 quantitative measures that describe sub- and supra-threshold integrative properties of each recorded cell (Figure 2A-C). Second, for the majority of mice two experimenters recorded in parallel from neurons in two sagittal brain sections from the same hemisphere. The median dorsal-ventral extent of the recorded SCs was 1644 µm (range 0 - 2464 µm). Each experimenter aimed to evenly sample recording locations across the dorsoventral extent of the MEC, and for most animals each experimenter recorded sequentially from opposite extremes of the dorsoventral axis. Each experimenter varied the starting location for recording between animals. For several features the direction of recording affected their measured dependence on dorsoventral location, but the overall dependence of these features on dorsoventral location was robust to this effect (Supplemental Table 9).

Measurement of electrophysiological features. Electrophysiological recordings were analysed in Matlab (Mathworks) using a custom written semi-automated pipeline. We defined each feature as follows (see also (Nolan et al., 2007; Pastoll et al., 2012a)): resting membrane potential was the mean of the membrane potential during 1 s prior to injection of current steps used to assess subthreshold properties; input resistance was the steady-state voltage response to the negative current steps divided by their amplitude; membrane time constant was the time constant of an exponential function fit to the initial phase of membrane potential responses to

the negative current steps; the sag coefficient was the steady-state divided by the peak membrane potential response to the negative current steps; resonance frequency was the frequency at which the peak membrane potential impedance was found to occur; resonance magnitude was the ratio between the peak impedance and the impedance at a frequency of 1 Hz; action potential threshold was calculated from responses to positive current ramps as the membrane potential at which the first derivative of the membrane potential crossed 20 mv ms^{-1} averaged across the first five spikes, or fewer if less spikes were triggered; rheobase was the ramp current at the point when the threshold was crossed on the first spike; spike maximum was the mean peak amplitude of the action potentials triggered by the positive current ramp; spike width was the duration of the action potentials measured at the voltage corresponding to the midpoint between the spike threshold and spike maximum; the AHP minimum was the negative peak membrane potential of the afterhyperpolarization following the first action potential when a second action potential also occurred; the F-I slope was the linear slope of the relationship between the spike rate and injected ramp current over a 500 ms window.

Analysis of location-dependence, experience-dependence and inter-animal variability

Analyses of location-dependence and inter-animal differences used R 3.4.3 (R Core Team, Vienna, Austria) and R Studio 1.1.383 (R Studio Inc., Boston, MA).

To fit linear mixed effect models we used the lme4 package (Bates et al., 2014). Features of interest were included as fixed effects and animal identity was included as a random effect. All reported analyses are for models with the maximal random effect structure, in other words the random effect was specified with uncorrelated slope and intercept. We also evaluated models in which only the intercept, or correlated intercept and slope were specified as the random effect. We compared the models used their Akaike Information Criterion (AIC) scores. In general, models with either random slope and intercept, or correlated random slope and intercept, had lower AIC scores than random intercept only models, indicating a better fit to the data. We used the former set of models for all analyses of all properties for consistency and because a maximal effect structure may be preferable on theoretical grounds (Barr et al., 2013). We evaluated assumptions including linearity, normality, homoscedasticity and influential data points. For some features we found modest deviations from these assumptions that could be corrected using copula transformation of the data. Model fits were similar following transformation of the data. However, because transformation loses scale information we focus here on analyses of the untransformed data.

To evaluate location-dependence of a given feature, p-values were calculated using a χ^2 test comparing models to null models with no location information but identical random effect specification. To calculate marginal and conditional R^2 of mixed effect models we used the MuMin package (Bartoń, 2014). To evaluate additional fixed effects we used Type II Wald χ^2 test tests provided by the car package (Fox and Weisberg, 2018). To compare mixed effect with equivalent linear models we used a χ^2 test to compare the calculated deviance for each model. For clarity we have reported key statistics in the main text and provide full test statistics in the Supplemental Tables. In addition the code will be provided as an R markdown document in which the analyses can be fully reproduced.

To evaluate partial correlations between features we used the function cor2pcor from the R package corpcor (Schafer et al., 2017). Principal components analysis used core R functions.

Implementation of tests for modularity. To establish statistical tests to distinguish 'modular' from 'continuous' distributions given relatively few observations, we classified datasets as continuous or modular by modifying the gap statistic algorithm (Tibshirani et al., 2001). The gap statistic estimates the number of clusters that best account for the data (k_{est}) in any given dataset (Supplementary Figure 1A-C). However, this estimate may be prone to false positives, particularly where the numbers of observations are low. We therefore introduced a thresholding mechanism for tuning the sensitivity of the algorithm so that the false positive rate, which is the rate of misclassifying datasets drawn from continuous (uniform) distributions as 'modular', is low, constant across different numbers of cluster modes and insensitive to dataset size (Supplementary Figure 1D-G). With this approach we are able to estimate whether a dataset is best described as lacking modularity ($k_{est} = 1$), or having a given number of modes ($k_{est} > 1$). We describe below tests carried out to validate the approach.

To illustrate the sensitivity and specificity of the modified gap statistic algorithm we applied it to simulated datasets drawn either from a uniform distribution (k = 1, n = 40) or a bimodal distribution with separation between the modes of 5 standard deviations (k = 2, n = 40, sigma = 5)(Supplemental Figure 2A). We set the thresholding mechanism so that k_{est} for each distinct k (where $k_{est} \geq 2$) has a false positive rate of 0.01. In line with this, testing for $2 \leq k_{est} \leq 8$ (the maximum k expected to occur in grid spacing in the MEC), across multiple (N = 1000) simulated

datasets drawn from the uniform distribution, produced a low false positive rate ($P(k_{est}) \geq 2 =$ ~0.07), whereas when the data were drawn from the bimodal distribution the ability to detect modular organisation (p_{detect}) was good ($P(k_{est}) \geq 2 = ~0.8$)(Supplemental Figure 2B). The performance of the statistic improved with larger separation between clusters and with greater numbers of data points per dataset (Supplemental Figure 2C) and is relatively insensitive to the numbers of clusters (Supplemental Figure 2D). The algorithm maintains high rates of p_{detect} when modes have varying densities and when sigma between modes varies in a manner similar to grid spacing data (Supplemental Figure 3).

Recently described algorithms (Giocomo et al., 2014; Stensola et al., 2012) address the problem of identifying modularity when data are sampled from multiple locations and data values vary as a function of location, as is the case for the mean spacing of grid fields for cells at different dorsoventral locations (Hafting et al., 2005). They generate log normalised discontinuity (which we refer to here as lnDS) or discreteness scores, which are the log of the ratio of discontinuity or discreteness scores for the data points of interest and for the sampling locations, with positive values interpreted as evidence for clustering (Giocomo et al., 2014; Stensola et al., 2012). However, in simulations of datasets generated from a uniform distribution with evenly spaced recording locations, we find that the lnDS is always greater than zero (Supplemental Figure 4A). This is because evenly spaced locations result in a discontinuity score that approaches zero and therefore the log ratio of the discontinuity of the data to this score will be positive. Thus, for evenly spaced data the lnDS is guaranteed to produce false positive results. When locations are instead sampled from a uniform distribution, approximately a half of simulated datasets have a log discontinuity ratio greater than 0 (Supplemental Figure 4A), which in previous studies would be interpreted as evidence for modularity (Giocomo et al., 2014). Similar discrepancies arise for the discreteness measure (Stensola et al., 2012). To address these issues we introduced a log discontinuity ratio threshold, so that the discontinuity method could be matched to produce a similar false positive rate to the adapted gap statistic algorithm used in the example above. After including this modification, we found that for a given false positive rate the adapted gap statistic is more sensitive at detecting modularity in the simulated datasets (Supplementary Figure 4B).

To establish whether the modified gap statistic detects clustering in experimental data we applied it to previously published grid cell data (Stensola et al., 2012). We find that the modified gap statistic identified clustered grid spacing for 6 of 7 animals previously identified as having

grid modules and with n ≥ 20. For these animals the number of modules was similar (but not always identical) to the number of previously identified modules (Supplementary Figure 5). In contrast, the modified gap statistic does not identify clustering in 5 of 6 sets of recording locations, confirming that the grid clustering is likely not a result of uneven sampling of locations (we could not test the 7th as location data were not available). The thresholded discontinuity score also detects clustering in the same 5 of the 6 tested sets of grid data. From the 6 grid datasets detected as clustered with the modified gap statistic we estimated the separation between clusters by fitting the data with a mixture of Gaussians, with the number of modes set by the value k obtained with the modified gap statistic. This analysis suggested that the largest spacing between contiguous modules in each mouse is always > 5.6 standard deviations (mean = 20.5 ± 5.0 standard deviations). Thus, the modified gap statistic detects modularity found in the grid system and indicates that previous descriptions of grid modularity are in general robust to the possibility of false positives associated with the discreteness and discontinuity methods.

Data and code availability

Data will be made available through the University of Edinburgh Datashare resource (https://datashare.is.ed.ac.uk/). All analyses will be made available at https://github.com/MattNolanLab/.

Supplemental Figures

Supplemental Figure 1. A quantitative adaption of the Gap Statistic clustering algorithm
(A-C) Logic of the Gap Statistic. (A) Simulated clustered dataset with three modes (k = 3)(open grey circles) (upper) and corresponding simulated reference dataset drawn from a uniform distribution with lower and upper limits set by the minimum and maximum values from the corresponding clustered dataset (open grey diamonds). Data were allocated to clusters by running a K-means algorithm 20 times and selecting the partition with the lowest average intracluster dispersion. Red, green blue and yellow dashed ovals show data subsets allocated to

each cluster when k_{Eval} = 1, 2, 3 and 4 modes are considered respectively. (B) The average value of the log intracluster dispersion for the clustered (open circles) and reference (open diamonds) datasets for each value of k tested in (A). (C) Gap values resulting from the difference between the clustered and reference values for each k in (B). Many ($\geq$ 500 in this paper) reference distributions are generated and their mean intracluster dispersion values are subtracted from those arising from the clustered distribution to maximise the reliability of Gap values.

(D) A procedure for selecting the optimal k depending on the associated gap values. Quantitative procedure for selecting optimal k (k_{est}). ΔGap values (open circles) are obtained by subtracting from the Gap value of a given k the Gap value for the previous k (ΔGap_k = Gap_k - Gap_{k-1}). For each ΔGap_k, if its ΔGap value greater than a threshold (filled triangles) associated with that ΔGap_k, that ΔGap_k will be k_{est}, if no ΔGap exceeds its threshold, k_{est} = 1. In the figure, for ΔGap_k = 2, 3, 4 (brown, pink and cyan), the ΔGap value exceeds its threshold only when ΔGap_k = 3. Therefore k_{est} = 3. (E-G) Determination of ΔGap_k thresholds. (E) Histograms of ΔGap values calculated from 20000 simulated datasets drawn from uniform distributions for each ΔGap_k (brown, pink and cyan respectively for ΔGap_k = 2, 3, 4) for a single dataset size (n = 40). ΔGap thresholds (filled triangles) are the values beyond which 1% of the ΔGap values fall and vary with ΔGap_k. (F) Histograms of ΔGap values for a range of dataset sizes (n = 20, 40, 100) and their associated thresholds. (G) Plot of the ΔGap thresholds as a function of dataset size and ΔGap_k. For separate ΔGap_k, ΔGap threshold values are well fit by a hyperbolic function of dataset size. These fits provide a practical method of finding the appropriate ΔGap threshold for an arbitrary dataset size.

Supplemental Figure 2. Discrimination of continuous from modular organisations using the adapted Gap Statistic algorithm

(A) Simulated datasets (each n = 40) drawn from continuous (uniform, k = 1 mode) (upper) and modular (multimodal mixture of Gaussians with k = 2 modes)(lower) distributions, and plotted

against simulated dorsoventral locations. Also shown are probability density functions (pdf) used to generate each dataset (light blue) and the densities estimated post-hoc from the generated data as kernel smoothed densities (light grey pdfs).

(B) Histograms showing the distribution of k_{est} from 1000 simulated datasets drawn from each pdf in (A). k_{est} is determined for each dataset by a modified gap statistic algorithm (see Supplemental Figure 1 above). When $k_{est} = 1$, the dataset is considered continuous (unclustered), when $k_{est} \geq 2$ the dataset is considered modular (clustered). The algorithm operates only on the feature values and does not use location information.

(C) Illustration of a set of clustered (k = 2) pdfs with the distance (in standard deviations) between clusters ranging from 2 to 6 (upper). Systematic evaluation of the ability of the modified gap statistic algorithm to detect clustered organisation ($k_{est} \geq 2$) in simulated datasets of different size (n = 20 to 100) drawn from the clustered (filled blue) and continuous (open blue) pdfs (lower). The proportion of datasets drawn from the continuous distribution that have $k_{est} \geq 2$ is the false positive (FP) rate (pFP = ~0.07). The light grey filled circle shows the smallest dataset size (n = 40) with SD = 5 where the proportion of datasets detected as clustered (p_{detect}) is ~ 0.8.

(D). Plot showing how p_{detect} at n = 40, SD = 5 changes when datasets are drawn from pdfs with different numbers of clusters (n modes from 2 to 8). Further evaluation of analysis of additional clusters is in the following figure.

Supplemental Figure 3. Additional evaluation of the adapted Gap Statistic algorithm

(A – C) Plots showing the results showing how p_{detect} (the ability of the modified gap statistic algorithm to detect clustered organisation) depends on dataset size and separation between cluster modes in simulated datasets drawn from clustered pdfs with different numbers of modes. The grey markers indicate n = 40, SD = 5 (as shown in Figure 1E). In each plot, p_{detect} is shown as a function of simulated dataset size and separation between modes when k = 3 (A), k = 5 (B) and k = 8 (C), which was the maximum number of clusters evaluated.

(D – F) Histograms showing the counts of k_{est} from the 1000 simulated n = 40, SD = 5 datasets (grey filled circles) illustrated in (A) – (C) respectively.

(G) p_{detect} as a function of dataset size and mode separation with k = 5 but when cluster modes are unevenly sampled. Sample sizes from clusters vary randomly with each dataset. A single mode can contribute from all to none of the points in any simulated dataset.

(H) p_{detect} as a function of dataset size and mode separation with k = 5 but when the distance between mode centres increases by a factor of sqrt(2) between sequential cluster pairs. Data is shown for different initial separations (the distance between the first two cluster centres) ranging from 1 to 4 (with corresponding separations between the final cluster pair ranging from 4 to 16).

Supplemental Figure 4. Comparing delta gap with discontinuity measures for discreteness

(A) Counts of log discontinuity ratio scores generated from a simulated uniform data distribution. The data distribution was ordered and then sampled either at positions drawn at random from a

uniform distribution (dark blue), or at positions with a fixed increment (light blue). For the data sampled at random positions approximately half of the scores are > 0 and for even sampling all scores are > 0. Therefore, a threshold score > 0 does not distinguish discrete from continuous distributions.

(B) Comparison of p_{detect} as a function of dataset size for the adapted Gap Statistic Algorithm, the discontinuity (upper) and the discreteness algorithm (lower). Each algorithm is adjusted to yield a 7% false positive rate. Each column shows simulations of data with different numbers of modes (k).

Supplemental Figure 5. Evaluation of modularity of grid firing using a modified gap statistic algorithm

Examples of grid spacing for individual neurons (crosses) from different mice. Kernel smoothed densities (KSDs) were generated with either a wide (solid grey) or narrow (dashed lines) kernel. The number of modes estimated using the modified gap statistic algorithm ranges is ≥ 2 for all but one animal (animal 4) with n ≥ 20 (animals 3 and 7 have < 20 recorded cells). We did not have location information for animal 2.

Supplemental Figure 6. Large environment for housing
(A-B) The large cage environment viewed from above (A) and from inside (B).

Supplemental Tables

Property	Slope	p (slope)	Marginal R2	Conditional R2	Slope (min)	Slope (max)	p (vs linear)
Vm	2.0086	0.1558	0.1104	0.1104	2.0086	2.0086	1
IR	0.1655	0.9652	0.0001	0.0181	0.1655	0.1655	1
Sag	-0.0137	0.5928	0.0142	0.0142	-0.0137	-0.0137	1
Tm	3.0716	0.2531	0.0861	0.1681	1.7960	4.2054	1
Res. frequency	0.2200	0.8135	0.0042	0.0042	0.2200	0.2200	1
Res. magnitude	0.0940	0.3552	0.0673	0.1764	0.0623	0.1553	1
Spike thresold	2.5595	0.1558	0.0986	0.0986	2.5595	2.5595	1
Spike maximum	1.1065	0.4628	0.0280	0.1533	1.1065	1.1065	1
Spike width	0.0495	0.0298	0.1922	0.1922	0.0495	0.0495	1
Rheobase	-15.4864	0.5928	0.0165	0.0165	-15.4864	-15.4864	1
Spike AHP	1.6079	0.3742	0.0470	0.0982	1.0252	2.2523	1
I-F slope	-0.0037	0.8625	0.0019	0.0019	-0.0037	-0.0037	1

Supplemental Table 1. Dependence of calbindin cell properties on dorsoventral position
Analyses are as described for Table 1. Data are from GFP positive putative pyramidal neurons
(n = 42, N = 3).

	All ages					P32 < age					P32 < age < P45				
property	dvloc	age	dvloc:age	N	n	dvloc	age	dvloc:age	N	n	dvloc	age	dvloc:age	N	n
vm	3.32e-07	0.557584	0.0833	27	836	1.65e-07	0.869	0.00263	26	795	4.44e-06	0.691	0.732	25	779
ir	5.07e-75	0.000804	0.6174	27	836	3.57e-81	0.468	0.42610	26	795	2.89e-73	0.985	0.732	25	779
sag	4.38e-19	0.004268	0.0833	27	836	1.06e-20	0.468	0.02552	26	795	7.84e-18	0.691	0.732	25	779
tau	5.19e-28	0.091904	0.9044	27	836	1.45e-28	0.869	0.96715	26	795	4.16e-28	0.844	0.732	25	779
resf	1.37e-17	0.006574	0.1811	27	836	7.11e-21	0.468	0.00242	26	795	1.54e-17	0.985	0.732	25	779
resmag	5.78e-12	0.006574	0.2673	27	836	4.42e-13	0.869	0.16209	26	795	7.84e-12	0.691	0.732	25	779
spkthr	8.40e-01	0.648357	0.6174	27	836	8.47e-01	0.869	0.42923	26	795	6.71e-01	0.985	0.806	25	779
spkmax	3.97e-08	0.784797	0.6678	27	836	1.96e-07	0.869	0.67149	26	795	3.98e-07	0.985	0.732	25	779
spkhlf	1.31e-02	0.053866	0.9182	27	836	2.26e-02	0.869	0.43513	26	795	4.21e-02	0.985	0.732	25	779
rheo	4.20e-79	0.247460	0.6174	27	836	2.65e-73	0.869	0.96715	26	795	2.25e-71	0.691	0.732	25	779
ahp	1.14e-02	0.648357	0.9182	27	836	1.67e-02	0.579	0.97670	26	795	2.31e-02	0.691	0.891	25	779
fi	6.48e-21	0.257819	0.3016	27	692	5.65e-21	0.468	0.42923	26	666	7.27e-19	0.691	0.732	25	656

Supplemental Table 2. Dependence of SC properties on age
The distinguishing electrophysiological features of SCs and their dorsoventral organisation were
apparent at all ages, with some features depending significantly on age (left columns),
consistent with the idea that SCs continue to mature beyond P18 (Boehlen et al., 2010; Burton
et al., 2008). When we considered only animals between P33 and P44 we did not find any

significant effect of age (right columns). Significance estimates for the effects of dorsoventral position (dvloc), age (age) and interactions between dorsoventral position and age (dvloc:age) were estimated using type II ANOVA and Wald χ^2 test from fits to mixed models containing age and location as fixed effects and animal identity as random effects. Significance estimates were adjusted for multiple comparisons using the Benjamini and Hochberg method.

property	N	n	Fixed effects				raw p			adjusted p		
			Int	dvloc	housing	dv:housing	dvloc	housing	dv:housing	dvloc_adj	housing_adj	dv:housing_adj
vm	25	779	-63.657	-0.836	0.276	0.0524	6.1e-06	0.5330	0.8873	8.2e-06	0.581	0.972
ir	25	779	20.015	11.288	-2.306	0.3679	7.2e-74	0.0942	0.7807	4.3e-73	0.188	0.972
sag	25	779	0.531	0.032	0.023	0.0009	3.0e-19	0.0031	0.9042	8.9e-19	0.031	0.972
tau	25	779	8.408	2.301	-1.110	0.4297	9.1e-28	0.0606	0.3824	3.7e-27	0.145	0.765
resf	25	779	9.187	-1.024	1.189	-0.2756	3.9e-17	0.0113	0.3543	7.8e-17	0.045	0.765
resmag	25	779	1.804	-0.091	0.048	-0.0339	8.7e-12	0.7457	0.3255	1.5e-11	0.746	0.765
spkthr	25	779	-39.222	0.140	0.715	0.0275	6.8e-01	0.4332	0.9725	6.8e-01	0.537	0.972
spkmax	25	779	43.696	2.343	1.073	-0.6996	3.3e-07	0.4478	0.3664	5.0e-07	0.537	0.765
spkhlf	25	779	0.501	0.022	-0.055	-0.0136	4.0e-02	0.0052	0.3060	4.4e-02	0.031	0.765
rheo	25	779	398.932	-97.192	65.598	-35.5348	7.3e-96	0.0275	0.0031	8.8e-95	0.083	0.038
ahp	25	779	-56.474	-0.470	0.980	-0.2299	2.1e-02	0.2537	0.6794	2.5e-02	0.380	0.972
fi	25	656	0.034	0.039	0.011	-0.0032	5.0e-19	0.1874	0.7107	1.2e-18	0.321	0.972

Supplemental Table 3. Dependence of SC properties on housing
Significance estimates for the effects of dorsoventral position (dvloc), housing (housing) and interactions between dorsoventral position and housing (dvloc:housing) estimated using type II ANOVA and Wald χ^2 test from fits to mixed models containing age and location as fixed effects and animal identity as random effects. Initial significance estimates (raw p) were adjusted for multiple comparisons (adjusted p) using the Benjamini and Hochberg method.

property	deviance (mixed)	deviance (linear)	df (mixed)	df (linear)	p	p_adj
vm	3216.17	3368.84	7	5	7.05e-34	2.12e-33
ir	5212.59	5309.84	7	5	7.65e-22	1.31e-21
sag	-2740.77	-2645.37	7	5	1.92e-21	2.88e-21
tau	3704.02	3786.67	7	5	1.13e-18	1.50e-18
resf	2801.35	2955.20	7	5	3.91e-34	1.56e-33
resmag	-81.94	-50.32	7	5	1.37e-07	1.37e-07
spkthr	3814.26	3861.51	7	5	5.49e-11	5.99e-11
spkmax	4071.15	4274.50	7	5	6.98e-45	4.19e-44
spkhlf	-2237.45	-1943.78	7	5	1.70e-64	2.04e-63
rheo	8742.69	8885.94	7	5	7.84e-32	1.57e-31
ahp	3909.56	3977.71	7	5	1.59e-15	1.91e-15
fi	-2766.11	-2622.51	7	5	6.55e-32	1.57e-31

Supplemental Table 4. Inter-animal differences in electrophysiological features remain after accounting for housing
Results from comparison of mixed effect model incorporating dorsoventral location and housing, with an equivalent linear model. The significance estimate (p) is calculated using a chi squared test and adjusted for multiple comparisons (p_adj) using the Benjamini and Hochberg method.

property	N	n	Fixed effects				raw p			adjusted p		
			Int	dvloc	hemi	dv:hemi	dvloc	hemi	dv:hemi	dvloc_adj	hemi_adj	dv:hemi_adj
vm	15	488	-63.032	-1.109	-0.651	0.3327	1.5e-05	0.518	0.45215	2.0e-05	0.95	0.6782
ir	15	488	18.039	12.787	-0.250	-0.7602	4.8e-55	0.630	0.66028	5.8e-54	0.95	0.8804
sag	15	488	0.528	0.050	0.021	-0.0269	3.9e-15	0.798	0.00197	7.8e-15	0.95	0.0118
tau	15	488	6.850	3.035	1.449	-0.5172	9.1e-23	0.041	0.38699	3.6e-22	0.45	0.6782
resf	15	488	10.472	-1.829	-0.994	0.7301	7.0e-16	0.134	0.04116	1.7e-15	0.54	0.1646
resmag	15	488	1.954	-0.210	-0.176	0.1422	4.0e-10	0.884	0.00022	6.9e-10	0.95	0.0026
spkthr	15	488	-38.212	-0.494	-0.239	0.0677	3.3e-01	0.871	0.94670	3.3e-01	0.95	0.9775
spkmax	15	488	43.582	3.043	0.754	-1.2048	5.5e-07	0.850	0.21065	8.2e-07	0.95	0.6319
spkhlf	15	488	0.467	0.024	0.016	-0.0017	4.6e-03	0.413	0.92148	5.0e-03	0.95	0.9775
rheo	15	488	436.558	-118.477	1.454	0.5655	1.2e-38	0.954	0.97750	7.1e-38	0.95	0.9775
ahp	15	488	-55.426	-1.217	-0.664	0.4963	3.5e-03	0.774	0.44470	4.2e-03	0.95	0.6782
fi	15	399	0.046	0.031	-0.016	0.0114	3.9e-16	0.075	0.28022	1.2e-15	0.45	0.6725

Supplemental Table 5. Dependence of SC properties on hemisphere
We did not find significant effects of brain hemisphere on any features except for the relationship between dorsoventral location and sag. Significance estimates for the effects of

dorsoventral position (dvloc), brain hemisphere (hemi) and interactions between dorsoventral position and hemisphere (dvloc:hemi) were estimated using type II ANOVA and Wald χ^2 test from fits to mixed models containing age and location as fixed effects and animal identity as random effects. Initial significance estimates (raw p) were adjusted for multiple comparisons (adjusted p) using the Benjamini and Hochberg method.

property	N	n	Fixed effects				raw p			adjusted p		
			Int	dvloc	ml	dv:ml	dvloc	ml	dv:ml	dvloc_adj	ml_adj	dv:ml_adj
vm	18	650	-63.390	-1.069	-0.4528	0.8628	1.1e-03	4.9e-03	0.00531	1.5e-03	5.3e-03	0.0159
ir	18	650	16.951	12.533	3.0417	-1.1815	1.6e-63	1.9e-03	0.28948	1.9e-62	2.9e-03	0.3860
sag	18	650	0.547	0.036	-0.0052	-0.0092	1.2e-14	3.1e-05	0.16974	3.7e-14	5.3e-05	0.2546
tau	18	650	7.659	3.027	0.8499	-1.4678	5.6e-16	2.4e-03	0.00033	2.2e-15	3.2e-03	0.0020
resf	18	650	9.552	-1.307	-0.0139	0.3652	1.0e-12	3.1e-03	0.12308	2.1e-12	3.7e-03	0.2110
resmag	18	650	1.847	-0.154	-0.0318	0.1077	1.8e-09	1.0e-05	0.00138	2.6e-09	2.0e-05	0.0055
spkthr	18	650	-40.060	0.300	2.2349	-0.1210	5.9e-01	1.2e-21	0.78976	5.9e-01	3.6e-21	0.8482
spkmax	18	650	46.435	2.376	-3.6126	-1.1149	5.4e-10	1.0e-107	0.00910	9.2e-10	1.2e-106	0.0218
spkhlf	18	650	0.497	0.012	-0.0107	0.0019	9.0e-02	8.6e-02	0.84819	9.8e-02	8.6e-02	0.8482
rheo	18	650	456.554	-121.469	-28.2336	3.1109	6.1e-35	4.7e-06	0.78003	3.7e-34	1.1e-05	0.8482
ahp	18	650	-56.987	-1.126	1.8540	1.5587	8.0e-02	3.2e-57	0.00017	9.5e-02	1.9e-56	0.0020
fi	18	548	0.027	0.033	0.0176	0.0118	2.6e-13	1.9e-33	0.01812	6.2e-13	7.4e-33	0.0362

Supplemental Table 6. Dependence of SC properties on mediolateral position
Mediolateral as well as dorsoventral position has been reported to determine the sub-threshold electrophysiological features of SCs (Canto and Witter, 2012). We found significant effects of mediolateral position on all measured electrophysiological features. However, the sizes of the effects of mediolateral position on subthreshold features (vm, ir, sag, tau, resf, resmag, rheo) were much smaller than for dorsoventral position. In contrast, supra-threshold features (spkthr, spkmax, ahp) were more greatly affected by mediolateral position, with more medial neurons having a higher spike threshold, and lower amplitudes of the spike peak and after-hyperpolarization. Fixed effects are the intercept and slope coefficients for mixed models containing dorsoventral and mediolateral location as fixed effects and animal identity as random effects Significance estimates for the effects of dorsoventral position (dvloc), mediolateral position (ml) and interactions between dorsoventral position and mediolateral position (dvloc:ml) are estimated using type II ANOVA and Wald χ^2 test from the fits of the mixed models. Initial significance estimates (raw p) were adjusted for multiple comparisons (adjusted p) using the Benjamini and Hochberg method.

property	N	n	Fixed effects				raw p			adjusted p		
			Int	dvloc	exp	dv:exp	dvloc	exp	dv:exp	dvloc_adj	exp_adj	dv:exp_adj
vm	25	779	-63.157	-0.844	-0.4325	0.07527	1.0e-05	2.1e-02	0.82154	1.6e-05	2.6e-02	0.8962
ir	25	779	23.677	9.057	-7.2059	3.39094	7.2e-78	6.0e-12	0.00382	8.7e-77	1.0e-11	0.0229
sag	25	779	0.554	0.024	-0.0075	0.01003	6.1e-18	4.1e-01	0.17241	1.5e-17	4.1e-01	0.3448
tau	25	779	8.070	2.941	-0.7249	-0.33590	5.1e-30	3.3e-07	0.45881	2.0e-29	4.4e-07	0.6882
resf	25	779	8.805	-0.966	1.6859	-0.39172	1.2e-20	1.0e-31	0.09830	3.5e-20	2.5e-31	0.2359
resmag	25	779	1.751	-0.052	0.1123	-0.07864	1.7e-12	1.0e-01	0.04924	2.9e-12	1.1e-01	0.1969
spkthr	25	779	-36.296	-0.361	-3.3736	0.82167	4.1e-01	1.1e-34	0.06640	4.1e-01	3.4e-34	0.1992
spkmax	25	779	42.418	1.564	2.9575	0.15435	1.6e-05	7.2e-38	0.76938	2.1e-05	2.9e-37	0.8962
spkhlf	25	779	0.476	0.042	-0.0237	-0.03258	7.9e-03	6.4e-44	0.00027	9.5e-03	3.8e-43	0.0033
rheo	25	779	417.475	-115.526	35.8258	-6.35544	5.6e-63	3.1e-08	0.57950	3.4e-62	4.7e-08	0.7727
ahp	25	779	-53.306	0.044	-3.7310	-0.47087	1.8e-01	2.5e-111	0.24805	1.9e-01	3.0e-110	0.4252
fi	25	656	0.058	0.039	-0.0253	-0.00025	1.4e-16	2.1e-26	0.96190	2.8e-16	4.2e-26	0.9619

Supplemental Table 7. Dependence of SC properties on experimenter
We found that for many electrophysiological features the identity of the experimenter affected the intercept, but not the slope, of their relationship with dorsoventral position. All features except for spike threshold nevertheless followed a dorsoventral organisation after accounting for the experimenter. Significance estimates for the effects of dorsoventral position (dvloc), experimenter (exp) and interactions between dorsoventral position and experimenter (dvloc:exp) were estimated using type II ANOVA and Wald χ^2 test from fits to mixed models containing age and location as fixed effects and animal identity as random effects. Initial significance estimates (raw p) were adjusted for multiple comparisons (adjusted p) using the Benjamini and Hochberg method.

property	N	n	Fixed effects				raw p			adjusted p		
			Int	dvloc	rect	dv:rect	dvloc	rect	dv:rect	dvloc_adj	rect_adj	dv:rect_adj
vm	25	778	-70.3538	0.425	1.0e+01	-1.972	5.0e-07	2.2e-19	1.7e-01	6.8e-07	1.3e-18	4.0e-01
ir	25	778	14.8133	15.190	5.4e+00	-5.535	5.6e-72	8.5e-01	3.0e-01	6.7e-71	9.3e-01	5.0e-01
sag	25	778	0.5482	0.023	-9.3e-04	0.013	1.2e-18	4.9e-01	6.8e-01	2.8e-18	5.9e-01	6.8e-01
tau	25	778	8.2726	3.356	-9.7e-01	-1.142	2.7e-27	8.4e-02	5.8e-01	1.1e-26	1.3e-01	6.8e-01
resf	25	778	7.2369	-0.871	4.2e+00	-0.569	2.8e-16	1.5e-06	6.2e-01	5.7e-16	3.6e-06	6.8e-01
resmag	25	778	1.5901	-0.005	3.7e-01	-0.167	7.4e-12	9.0e-02	3.4e-01	1.3e-11	1.3e-01	5.0e-01
spkthr	25	778	-55.9007	6.033	2.6e+01	-9.539	2.7e-01	5.8e-36	1.3e-06	2.7e-01	7.0e-35	1.5e-05
spkmax	25	778	40.7004	-0.699	5.6e+00	3.866	5.1e-07	3.5e-09	1.3e-01	6.8e-07	1.0e-08	3.9e-01
spkhlf	25	778	0.5921	0.045	-2.0e-01	-0.045	2.4e-02	4.9e-17	3.1e-01	2.6e-02	1.9e-16	5.0e-01
rheo	25	778	451.9554	-98.901	-1.3e+01	-30.573	1.3e-66	1.7e-01	5.6e-01	7.7e-66	2.3e-01	6.8e-01
ahp	25	778	-60.7014	3.712	7.5e+00	-6.633	2.0e-02	9.7e-01	4.3e-03	2.4e-02	9.7e-01	1.7e-02
fi	25	655	0.0092	0.113	5.0e-02	-0.115	1.1e-22	4.5e-06	2.6e-06	3.4e-22	9.1e-06	1.6e-05

Supplemental Table 8. Dependence of SC properties on time since slice preparation

We anticipated that the interval between slice preparation and recording may influence measured electrophysiological features. Consistent with our expectation, analyses of the data were consistent with changes to some electrophysiological features of SCs with time since slice preparation, but dorsoventral gradients could not be explained by these changes (see also Supplemental Table 8 below). Significance estimates for the effects of dorsoventral position (dvloc), time since slice preparation (rect) and interactions between dorsoventral position and experimenter (dvloc:rect) estimated using type II ANOVA and Wald χ^2 test from fits to mixed models containing age and location as fixed effects and animal identity as random effects. Initial significance estimates (raw p) were adjusted for multiple comparisons (adjusted p) using the Benjamini and Hochberg method.

property	N	n	Fixed effects Int	dvloc	dir	dv:dir	raw p dvloc	dir	dv:dir	adjusted p dvloc_adj	dir_adj	dv:dir_adj
vm	18	650	-64.127	-0.063	1.351	-1.3535	2.8e-04	0.509	2.4e-05	3.7e-04	0.61	9.4e-05
ir	18	650	18.337	11.758	0.427	0.1689	8.7e-64	0.374	8.8e-01	1.0e-62	0.57	9.2e-01
sag	18	650	0.534	0.040	0.021	-0.0164	1.3e-16	0.411	1.9e-02	3.1e-16	0.57	3.8e-02
tau	18	650	7.975	2.269	0.091	0.2595	2.5e-20	0.144	5.4e-01	1.0e-19	0.49	7.2e-01
resf	18	650	9.372	-0.905	0.443	-0.5214	1.9e-12	0.429	3.7e-02	3.8e-12	0.57	6.3e-02
resmag	18	650	1.846	-0.104	-0.030	-0.0035	1.0e-08	0.122	9.2e-01	1.7e-08	0.49	9.2e-01
spkthr	18	650	-40.893	1.947	4.997	-4.2461	5.2e-01	0.219	1.0e-23	5.2e-01	0.49	1.2e-22
spkmax	18	650	43.529	3.044	2.134	-1.7908	5.0e-08	0.556	2.4e-03	7.4e-08	0.61	5.9e-03
spkhlf	18	650	0.511	-0.012	-0.042	0.0491	1.2e-01	0.059	2.8e-06	1.3e-01	0.49	1.7e-05
rheo	18	650	439.066	-119.612	10.708	-1.5603	2.0e-43	0.168	9.0e-01	1.2e-42	0.49	9.2e-01
ahp	18	650	-56.018	-0.474	0.408	-0.4158	1.1e-02	0.872	4.0e-01	1.3e-02	0.87	6.0e-01
fi	18	548	0.047	0.028	-0.023	0.0189	7.5e-18	0.245	6.5e-04	2.2e-17	0.49	1.9e-03

Supplemental Table 9. Dependence of SC properties on direction in which sequential recordings are made

In anticipation of effects of the time since slice preparation on electrophysiological features of SCS, we varied between experimenters and experimental days the direction along the dorsoventral axis from which consecutive recordings were made (see Methods). Consistent with effects of time on electrophysiological features (see Supplemental Table 7 above), we found that the direction in which sequential recordings were made influence the slope, but not the intercept of several electrophysiological features. Significance estimates for the effects of dorsoventral position (dvloc), direction in which sequential recordings were made (dir) and interactions between dorsoventral position and recording direction (dvloc:dir) estimated using type II ANOVA and Wald χ^2 test from fits to mixed models containing age and location as fixed

effects and animal identity as random effects. Initial significance estimates (raw p) were adjusted for multiple comparisons (adjusted p) using the Benjamini and Hochberg method.

property	deviance (mixed)	deviance (linear)	df (mixed)	df (linear)	p	p_adj
vm	2636.81	2789.34	13	11	7.56e-34	3.02e-33
ir	4315.73	4407.44	13	11	1.22e-20	1.62e-20
sag	-2293.74	-2212.49	13	11	2.27e-18	2.72e-18
tau	2997.16	3122.25	13	11	6.86e-28	1.37e-27
resf	2122.53	2266.61	13	11	5.16e-32	1.55e-31
resmag	-115.70	-80.54	13	11	2.31e-08	2.31e-08
spkthr	2936.17	2974.72	13	11	4.25e-09	4.63e-09
spkmax	3008.01	3239.16	13	11	6.40e-51	7.67e-50
spkhlf	-2067.19	-1911.08	13	11	1.26e-34	7.55e-34
rheo	7247.31	7388.78	13	11	1.91e-31	4.58e-31
ahp	2832.78	2943.86	13	11	7.58e-25	1.14e-24
fi	-2473.53	-2349.35	13	11	1.08e-27	1.85e-27

Supplemental Table 10. Interanimal differences in extended models
Results from comparison of mixed effect model incorporating dorsoventral location, housing, mediolateral position, experimenter identity and direction in which recordings were obtained with an equivalent linear model. Data are from animals between 32 and 45 days old. The significance estimate (p) is calculated using a chi squared test and adjusted for multiple comparisons (p_adj) using the Benjamini and Hochberg method.

Property	Nobs	Ngrps	marginal.R2	Conditional.R2	Dev(mixed)	Dev(linear)	Dev_DF (mixed)	Dev_DF (linear)	p	p(adj)
vm	155	5	0.0206282	0.2653056	618	639	5	3	3.54e-05	6.07e-05
ir	155	5	0.4338500	0.5813127	962	991	5	3	6.41e-07	1.54e-06
sag	155	5	0.1800715	0.4129883	-576	-543	5	3	5.57e-08	1.67e-07
tau	155	5	0.0931872	0.0931872	696	696	5	3	1.00e+00	1.00e+00
resf	155	5	0.0564489	0.2271287	531	551	5	3	6.27e-05	9.41e-05
resmag	155	5	0.0904314	0.2491198	-39.5	-24.1	5	3	4.59e-04	5.01e-04
spkthr	155	5	0.2037580	0.3875554	667	695	5	3	1.12e-06	2.24e-06
spkmax	155	5	0.2940325	0.4130673	639	655	5	3	4.12e-04	4.94e-04
spkhlf	155	5	0.0298569	0.5313239	-546	-458	5	3	7.54e-20	9.05e-19
rheo	155	5	0.2125328	0.7326873	1736	1788	5	3	7.33e-12	4.40e-11
ahp	155	5	0.0051765	0.1661218	549	566	5	3	2.12e-04	2.82e-04
fi	138	5	0.3032774	0.6856170	-778	-741	5	3	9.29e-09	3.72e-08

Property	Nobs	Ngrps	marginal.R2	Conditional.R2	Dev(mixed)	Dev(linear)	Dev_DF (mixed)	Dev_DF (linear)	p	p(adj)
vm	70	6	0.0938959	0.4186968	314	322	5	3	2.05e-02	2.73e-02
ir	70	6	0.1473580	0.5692115	475	502	5	3	2.05e-06	5.41e-06
sag	70	6	0.0086794	0.2642029	-223	-212	5	3	5.73e-03	9.83e-03
tau	70	6	0.0068137	0.0507558	327	327	5	3	7.95e-01	8.67e-01
resf	70	6	0.0008116	0.4514622	203	229	5	3	2.25e-06	5.41e-06
resmag	70	6	0.0238201	0.3629621	14.5	28.8	5	3	7.62e-04	1.52e-03
spkthr	70	6	0.1837547	0.4722489	342	352	5	3	8.09e-03	1.21e-02
spkmax	70	6	0.0519267	0.0745947	356	356	5	3	9.62e-01	9.62e-01
spkhlf	70	6	0.1474714	0.7119071	-268	-230	5	3	6.49e-09	3.89e-08
rheo	70	6	0.3428390	0.8415186	787	834	5	3	7.09e-11	8.51e-10
ahp	70	6	0.0207732	0.1760733	332	336	5	3	1.35e-01	1.62e-01
fi	60	6	0.2925095	0.7238371	-285	-248	5	3	1.00e-08	4.01e-08

Supplemental Table 11. Interanimal differences in minimal models

Results from comparison of mixed effect models with dorsoventral location as a fixed effect and animal identity as a random effect using a minimal datasets obtained by either HP (upper) or DG (lower). Data are from animals between 32 and 45 days old.

			Fixed effects				raw p			adjusted p		
property	N	n	Int	dvloc	counts	dv:counts	dvloc	counts	dv:counts	dvloc_adj	counts_adj	dv:counts_adj
vm	25	779	-63.262	-1.334	-0.00470	1.5e-02	7.2e-06	0.72309	0.384	9.7e-06	0.9641	0.58
ir	25	779	20.165	8.240	-0.04768	9.6e-02	1.2e-75	0.61560	0.113	1.4e-74	0.9234	0.23
sag	25	779	0.568	0.010	-0.00055	6.2e-04	8.3e-20	0.98892	0.080	2.5e-19	0.9889	0.23
tau	25	779	5.964	3.987	0.04659	-3.9e-02	8.6e-28	0.34149	0.092	3.4e-27	0.9234	0.23
resf	25	779	12.067	-1.880	-0.06141	1.8e-02	1.2e-18	0.00231	0.194	2.8e-18	0.0139	0.33
resmag	25	779	1.777	-0.066	0.00166	-1.3e-03	5.3e-12	0.83262	0.433	9.1e-12	0.9889	0.58
spkthr	25	779	-38.810	0.278	0.00217	-3.3e-03	6.7e-01	0.98673	0.925	6.7e-01	0.9889	0.97
spkmax	25	779	44.731	0.051	-0.00543	5.4e-02	1.5e-07	0.46922	0.117	2.2e-07	0.9234	0.23
spkhlf	25	779	0.327	0.064	0.00408	-1.4e-03	1.4e-02	0.00022	0.019	1.6e-02	0.0027	0.22
rheo	25	779	388.527	-73.875	1.56603	-1.3e+00	8.3e-74	0.45667	0.043	5.0e-73	0.9234	0.23
ahp	25	779	-56.673	-0.185	0.02539	-1.2e-02	2.7e-02	0.59590	0.659	2.9e-02	0.9234	0.79
fi	25	656	0.052	0.036	-0.00032	1.3e-05	5.5e-18	0.32090	0.973	1.1e-17	0.9234	0.97

Supplemental Table 12. Electrophysiological features and the number of recorded neurons

Significance estimates for the effects of dorsoventral position (dvloc), number of recorded neurons (counts) and interactions between dorsoventral position and number of recorded neurons (dvloc:counts) estimated using type II ANOVA and Wald χ^2 test from fits to mixed models containing age and location as fixed effects and animal identity as random effects. Initial

significance estimates (raw p) were adjusted for multiple comparisons (adjusted p) using the Benjamini and Hochberg method.

ngrps	nobs	marginal.R2	conditional.R2	mm_vslinear_pdiff_adj
11	459	0.0568	0.373	2.77e-29
11	459	0.5551	0.634	5.68e-13
11	459	0.1629	0.345	1.48e-13
11	459	0.2357	0.405	1.99e-16
11	459	0.2809	0.496	2.42e-23
11	459	0.1392	0.239	1.08e-07
11	459	0.3139	0.416	4.43e-05
11	459	0.4810	0.703	4.63e-31
11	459	0.3246	0.537	1.29e-20
11	459	0.5444	0.681	4.46e-26
11	459	0.4594	0.568	7.63e-16
11	390	0.5159	0.742	1.33e-12

Supplemental Table 13. Interanimal differences for experiments with > 35 recorded neurons
Analyses are as for Supplemental Table 1, but are restricted to experiments in which > 35 neurons were recorded from.

Feature	Slope	p (slope)	Marginal R2	Conditional R2	Slope (min)	Slope (max)	p (vs linear)
PC1	-2.43035	1.09e-15	0.49545	0.73250	-3.16510	-2.07369	3.86e-37
PC2	0.95272	1.05e-04	0.09299	0.80751	-1.07716	2.11870	2.20e-97
PC3	0.25632	1.73e-01	0.01187	0.51214	-0.43529	1.11057	2.96e-38
PC4	-0.26627	2.53e-01	0.00941	0.79049	-1.47230	1.06857	4.45e-63
PC5	-0.11523	4.01e-01	0.00390	0.43605	-0.49011	0.44712	3.10e-24
PC6	-0.01593	9.03e-01	0.00006	0.64068	-1.12576	1.08528	7.98e-31
PC7	-0.06545	4.90e-01	0.00250	0.05729	-0.18349	0.07602	2.94e-02
PC8	-0.01013	9.03e-01	0.00007	0.23751	-0.31346	0.24927	3.49e-10
PC9	-0.13547	3.95e-02	0.01617	0.18557	-0.22543	0.08319	4.98e-12
PC10	-0.01408	9.03e-01	0.00031	0.04810	-0.02747	-0.00033	8.19e-02
PC11	-0.00545	9.03e-01	0.00007	0.02613	-0.00794	-0.00383	2.74e-01
PC12	-0.07795	1.50e-02	0.02402	0.08892	-0.19333	0.01590	7.02e-04

Supplemental Table 14. Dependence of principal components on dorsoventral position and animal identity

Analyses are as described for Table 1, but are applied to principal components of the electrophysiological features of SCs.

property	N	n		Fixed effects				raw p			adjusted p	
			Int	dvloc	housing	dv:housing	dvloc	housing	dv:housing	dvloc_adj	housing_adj	dv:housing_adj
PC1	25	487	2.055	-2.373	0.9582	-0.0754	2.7e-86	0.0098	0.77164	3.3e-85	0.039	0.8890
PC2	25	487	-1.859	1.067	1.2929	-0.1590	2.1e-07	0.0045	0.68640	1.2e-06	0.027	0.8890
PC3	25	487	-0.157	0.301	-0.2277	-0.0681	6.3e-02	0.4571	0.81489	1.5e-01	0.609	0.8890
PC4	25	487	0.754	-0.369	-0.6951	0.1501	1.1e-01	0.1441	0.67220	2.2e-01	0.247	0.8890
PC5	25	487	0.284	-0.114	-0.1393	-0.0016	2.3e-01	0.6192	0.99353	4.0e-01	0.681	0.9935
PC6	25	487	0.436	-0.238	-0.4509	0.3286	8.8e-01	0.2440	0.21869	9.4e-01	0.366	0.6561
PC7	25	487	0.398	-0.279	-0.4526	0.3226	3.1e-01	0.0991	0.02447	4.6e-01	0.238	0.0979
PC8	25	487	0.277	-0.057	-0.3795	0.0706	8.7e-01	0.0037	0.59369	9.4e-01	0.027	0.8890
PC9	25	487	0.134	-0.177	-0.0042	0.0658	1.3e-02	0.6239	0.56105	3.8e-02	0.681	0.8890
PC10	25	487	0.097	-0.138	-0.1354	0.2118	9.4e-01	0.0430	0.00576	9.4e-01	0.129	0.0345
PC11	25	487	0.045	-0.039	-0.0537	0.0535	8.8e-01	0.8677	0.41405	9.4e-01	0.868	0.8890
PC12	25	487	-0.076	0.046	0.2241	-0.1870	3.7e-03	0.1242	0.00056	1.5e-02	0.247	0.0067

Supplemental Table 15. Dependence of principal components of SC properties on housing

Analyses are as described for Supplemental Table 3, but are applied to principal components of the electrophysiological features of SCs.

component	deviance (mixed)	deviance (linear)	df (mixed)	df (linear)	p	p_adj
PC1	1463.86	1596.78	7	5	1.37e-29	3.28e-29
PC2	1260.20	1658.40	7	5	3.40e-87	4.08e-86
PC3	1394.81	1567.87	7	5	2.63e-38	1.05e-37
PC4	1180.62	1441.66	7	5	2.07e-57	1.24e-56
PC5	1217.87	1319.67	7	5	7.83e-23	1.57e-22
PC6	1104.30	1237.28	7	5	1.33e-29	3.28e-29
PC7	1070.53	1078.31	7	5	2.05e-02	2.46e-02
PC8	904.79	929.29	7	5	4.78e-06	7.17e-06
PC9	827.72	881.34	7	5	2.27e-12	3.89e-12
PC10	590.98	592.69	7	5	4.25e-01	4.25e-01
PC11	432.20	434.54	7	5	3.12e-01	3.40e-01
PC12	116.24	128.36	7	5	2.34e-03	3.12e-03

Supplemental Table 16. Dependence of principal components on animal identity in models that account for housing

Analyses are as for Supplemental Table 10, but are applied to principal components of the electrophysiological features of SCs.

References

Adamson, C.L., Reid, M.A., Mo, Z.-L., Bowne-English, J., and Davis, R.L. (2002). Firing features and potassium channel content of murine spiral ganglion neurons vary with cochlear location. J. Comp. Neurol. *447*, 331–350.

Alonso, A., and Klink, R. (1993). Differential electroresponsiveness of stellate and pyramidal-like cells of medial entorhinal cortex layer II. J. Neurophysiol. *70*, 128–143.

Angelo, K., Rancz, E.A., Pimentel, D., Hundahl, C., Hannibal, J., Fleischmann, A., Pichler, B., and Margrie, T.W. (2012). A biophysical signature of network affiliation and sensory processing in mitral cells. Nature *488*, 375–378.

Baayen, R.H., Davidson, D.J., and Bates, D.M. (2008). Mixed-effects modeling with crossed random effects for subjects and items. J. Mem. Lang. *59*, 390–412.

Barr, D.J., Levy, R., Scheepers, C., and Tily, H.J. (2013). Random effects structure for confirmatory hypothesis testing: Keep it maximal. J. Mem. Lang. *68*.

Barry, C., Hayman, R., Burgess, N., and Jeffery, K.J. (2007). Experience-dependent rescaling of entorhinal grids. Nat. Neurosci. *10*, 682–684.

Bartoń, K. (2014). MuMIn: multi-model inference. R package version 1.10. 0. See http://CRAN. R-Project. Org/package= MuMIn.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2014). Fitting Linear Mixed-Effects Models using lme4.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statisitical Society, Series B *57*, 289–300.

Boehlen, A., Heinemann, U., and Erchova, I. (2010). The range of intrinsic frequencies represented by medial entorhinal cortex stellate cells extends with age. J. Neurosci. *30*, 4585–4589.

Booth, C.A., Ridler, T., Murray, T.K., Ward, M.A., de Groot, E., Goodfellow, M., Phillips, K.G., Randall, A.D., and Brown, J.T. (2016). Electrical and Network Neuronal Properties Are Preferentially Disrupted in Dorsal, But Not Ventral, Medial Entorhinal Cortex in a Mouse Model of Tauopathy. J. Neurosci. *36*, 312–324.

Burgess, N. (2008). Grid cells and theta as oscillatory interference: theory and predictions. Hippocampus *18*, 1157–1174.

Burton, B.G., Economo, M.N., Lee, G.J., and White, J.A. (2008). Development of theta rhythmicity in entorhinal stellate cells of the juvenile rat. J. Neurophysiol. *100*, 3144–3157.

Canto, C.B., and Witter, M.P. (2012). Cellular properties of principal neurons in the rat entorhinal cortex. II. The medial entorhinal cortex. Hippocampus *22*, 1277–1299.

Cembrowski, M.S., and Menon, V. (2018). Continuous Variation within Cell Types of the Nervous System. Trends Neurosci.

Dodson, P.D., Pastoll, H., and Nolan, M.F. (2011). Dorsal-ventral organization of theta-like activity intrinsic to entorhinal stellate neurons is mediated by differences in stochastic current fluctuations. J. Physiol. *589*, 2993–3008.

Domnisoru, C., Kinkhabwala, A.A., and Tank, D.W. (2013). Membrane potential dynamics of grid cells. Nature *495*, 199–204.

Donato, F., Jacobsen, R.I., Moser, M.-B., and Moser, E.I. (2017). Stellate cells drive maturation of the entorhinal-hippocampal circuit. Science *355*.

Fan, Y., Fricker, D., Brager, D.H., Chen, X., Lu, H.-C., Chitwood, R.A., and Johnston, D. (2005). Activity-dependent decrease of excitability in rat hippocampal neurons through increases in I(h). Nat. Neurosci. *8*, 1542–1551.

Fletcher, L.N., and Williams, S.R. (2018). Neocortical Topology Governs the Dendritic Integrative Capacity of Layer 5 Pyramidal Neurons. Neuron *0*.

Fox, J., and Weisberg, S. (2018). An R Companion to Applied Regression (SAGE Publications).

Fyhn, M., Molden, S., Witter, M.P., Moser, E.I., and Moser, M.-B. (2004). Spatial representation in the entorhinal cortex. Science *305*, 1258–1264.

Garden, D.L.F., Dodson, P.D., O'Donnell, C., White, M.D., and Nolan, M.F. (2008). Tuning of synaptic integration in the medial entorhinal cortex to the organization of grid cell firing fields. Neuron *60*, 875–889.

Geiler-Samerotte, K.A., Bauer, C.R., Li, S., Ziv, N., Gresham, D., and Siegal, M.L. (2013). The details in the distributions: why and how to study phenotypic variability. Curr. Opin. Biotechnol. *24*, 752–759.

Giocomo, L.M., and Hasselmo, M.E. (2008a). Time constants of h current in layer ii stellate cells differ along the dorsal to ventral axis of medial entorhinal cortex. J. Neurosci. *28*, 9414–9425.

Giocomo, L.M., and Hasselmo, M.E. (2008b). Computation by oscillations: implications of experimental data for theoretical models of grid cells. Hippocampus *18*, 1186–1199.

Giocomo, L.M., and Hasselmo, M.E. (2009). Knock-out of HCN1 subunit flattens dorsal-ventral frequency gradient of medial entorhinal neurons in adult mice. J. Neurosci. *29*, 7625–7630.

Giocomo, L.M., Zilli, E.A., Fransén, E., and Hasselmo, M.E. (2007). Temporal frequency of subthreshold oscillations scales with entorhinal grid cell field spacing. Science *315*, 1719–1722.

Giocomo, L.M., Hussaini, S.A., Zheng, F., Kandel, E.R., Moser, M.-B., and Moser, E.I. (2011). Grid cells use HCN1 channels for spatial scaling. Cell *147*, 1159–1170.

Giocomo, L.M., Stensola, T., Bonnevie, T., Van Cauter, T., Moser, M.-B., and Moser, E.I. (2014). Topography of head direction cells in medial entorhinal cortex. Curr. Biol. *24*, 252–262.

Goaillard, J.-M., Taylor, A.L., Schulz, D.J., and Marder, E. (2009). Functional consequences of animal-to-animal variation in circuit parameters. Nat. Neurosci. *12*, 1424–1430.

Gonzalez-Sulser, A., Parthier, D., Candela, A., McClure, C., Pastoll, H., Garden, D., Sürmeli, G., and Nolan, M.F. (2014). GABAergic projections from the medial septum selectively inhibit

interneurons in the medial entorhinal cortex. J. Neurosci. *34*, 16739–16743.

Grossberg, S., and Pilly, P.K. (2012). How entorhinal grid cells may learn multiple spatial scales from a dorsoventral gradient of cell response rates in a self-organizing map. PLoS Comput. Biol. *8*, e1002648.

Gu, Y., Lewallen, S., Kinkhabwala, A.A., Domnisoru, C., Yoon, K., Gauthier, J.L., Fiete, I.R., and Tank, D.W. (2018). A Map-like Micro-Organization of Grid Cells in the Medial Entorhinal Cortex. Cell.

Hafting, T., Fyhn, M., Molden, S., Moser, M.-B., and Moser, E.I. (2005). Microstructure of a spatial map in the entorhinal cortex. Nature *436*, 801–806.

Kitamura, T., Pignatelli, M., Suh, J., Kohara, K., Yoshiki, A., Abe, K., and Tonegawa, S. (2014). Island cells control temporal association memory. Science *343*, 896–901.

Kropff, E., and Treves, A. (2008). The emergence of grid cells: Intelligent design or just adaptation? Hippocampus *18*, 1256–1269.

Kuba, H., Yamada, R., Fukui, I., and Ohmori, H. (2005). Tonotopic specialization of auditory coincidence detection in nucleus laminaris of the chick. J. Neurosci. *25*, 1924–1934.

Liss, B., Franz, O., Sewing, S., Bruns, R., Neuhoff, H., and Roeper, J. (2001). Tuning pacemaker frequency of individual dopaminergic neurons by Kv4.3L and KChip3.1 transcription. EMBO J. *20*, 5715–5724.

Mallory, C.S., Hardcastle, K., Bant, J.S., and Giocomo, L.M. (2018). Grid scale drives the scale and long-term stability of place maps. Nat. Neurosci. 1–1.

Marder, E., and Goaillard, J.-M. (2006). Variability, compensation and homeostasis in neuron and network function. Nat. Rev. Neurosci. *7*, 563–574.

Marder, E., and Taylor, A.L. (2011). Multiple models to capture the variability in biological neurons and networks. Nat. Neurosci. *14*, 133–138.

Mittal, D., and Narayanan, R. (2018). Degeneracy in the robust expression of spectral selectivity, subthreshold oscillations, and intrinsic excitability of entorhinal stellate cells. J. Neurophysiol. *120*, 576–600.

Murray, A.J., Sauer, J.-F., Riedel, G., McClure, C., Ansel, L., Cheyne, L., Bartos, M., Wisden, W., and Wulff, P. (2011). Parvalbumin-positive CA1 interneurons are required for spatial working but not for reference memory. Nat. Neurosci. *14*, 297–299.

Narayanan, R., and Johnston, D. (2007). Long-term potentiation in rat hippocampal neurons is accompanied by spatially widespread changes in intrinsic oscillatory dynamics and excitability. Neuron *56*, 1061–1075.

Nolan, M.F., Malleret, G., Dudman, J.T., Buhl, D.L., Santoro, B., Gibbs, E., Vronskaya, S., Buzsáki, G., Siegelbaum, S.A., Kandel, E.R., et al. (2004). A behavioral role for dendritic integration: HCN1 channels constrain spatial memory and plasticity at inputs to distal dendrites of CA1 pyramidal neurons. Cell *119*, 719–732.

Nolan, M.F., Dudman, J.T., Dodson, P.D., and Santoro, B. (2007). HCN1 channels control

resting and active integrative properties of stellate cells from layer II of the entorhinal cortex. J. Neurosci. *27*, 12440–12451.

O'Donnell, C., and Nolan, M.F. (2011). Tuning of synaptic responses: an organizing principle for optimization of neural circuits. Trends Neurosci. *34*, 51–60.

O'Leary, T., Williams, A.H., Franci, A., and Marder, E. (2014). Cell types, network homeostasis, and pathological compensation from a biologically plausible ion channel expression model. Neuron *82*, 809–821.

Pastoll, H., Ramsden, H.L., and Nolan, M.F. (2012a). Intrinsic electrophysiological properties of entorhinal cortex stellate cells and their contribution to grid cell firing fields. Front. Neural Circuits *6*, 17.

Pastoll, H., White, M., and Nolan, M. (2012b). Preparation of parasagittal slices for the investigation of dorsal-ventral organization of the rodent medial entorhinal cortex. J. Vis. Exp.

Pastoll, H., Solanka, L., van Rossum, M.C.W., and Nolan, M.F. (2013). Feedback Inhibition Enables Theta-Nested Gamma Oscillations and Grid Firing Fields. Neuron *77*, 141–154.

Qin, H., Fu, L., Hu, B., Liao, X., Lu, J., He, W., Liang, S., Zhang, K., Li, R., Yao, J., et al. (2018). A Visual-Cue-Dependent Memory Circuit for Place Navigation. Neuron 1–9.

Ramsden, H.L., Sürmeli, G., McDonagh, S.G., and Nolan, M.F. (2015). Laminar and dorsoventral molecular organization of the medial entorhinal cortex revealed by large-scale anatomical analysis of gene expression. PLoS Comput. Biol. *11*, e1004032.

Ray, S., and Brecht, M. (2016). Structural development and dorsoventral maturation of the medial entorhinal cortex. Elife *5*, e13343–e13343.

Regev, A., Teichmann, S.A., Lander, E.S., Amit, I., Benoist, C., Birney, E., Bodenmiller, B., Campbell, P., Carninci, P., Clatworthy, M., et al. (2017). Science forum: the human cell atlas. Elife *6*, e27041.

Rowland, D.C., Obenhaus, H.A., Skytøen, E.R., Zhang, Q., Kentros, C.G., Moser, E.I., and Moser, M.-B. (2018). Functional properties of stellate cells in medial entorhinal cortex layer II. Elife *7*.

Schafer, J., Opgen-Rhein, R., Zuber, V., Ahdesmaki, M., Silva, A.P.D., and Strimmer, K. (2017). corpcor: Efficient estimation of covariance and (partial) correlation (R package version 1.6. 9).

Schmidt-Hieber, C., and Häusser, M. (2013). Cellular mechanisms of spatial navigation in the medial entorhinal cortex. Nat. Neurosci. *16*, 325–331.

Shipston-Sharman, O., Solanka, L., and Nolan, M.F. (2016). Continuous attractor network models of grid cell firing based on excitatory-inhibitory interactions. J. Physiol. *594*, 6547–6557.

Stensola, H., Stensola, T., Solstad, T., Frøland, K., Moser, M.-B., and Moser, E.I. (2012). The entorhinal grid map is discretized. Nature *492*, 72–78.

Sürmeli, G., Marcu, D.C., McClure, C., Garden, D.L.F., Pastoll, H., and Nolan, M.F. (2015). Molecularly Defined Circuitry Reveals Input-Output Segregation in Deep Layers of the Medial Entorhinal Cortex. Neuron *88*, 1040–1053.

Swensen, A.M., and Bean, B.P. (2005). Robustness of burst firing in dissociated purkinje neurons with acute or long-term reductions in sodium conductance. J. Neurosci. *25*, 3509–3520.

Tennant, S.A., Fischer, L., Garden, D.L.F., Gerlei, K.Z., Martinez-Gonzalez, C., McClure, C., Wood, E.R., and Nolan, M.F. (2018). Stellate Cells in the Medial Entorhinal Cortex Are Required for Spatial Learning. Cell Rep. *22*, 1313–1324.

Tibshirani, R., Walther, G., and Hastie, T. (2001). Estimating the number of clusters in a data set via the gap statistic. J. R. Stat. Soc. Series B Stat. Methodol. *63*, 411–423.

Urdapilleta, E., Si, B., and Treves, A. (2017). Selforganization of modular activity of grid cells. Hippocampus *27*, 1204–1213.

Wang, F., Zhu, J., Zhu, H., Zhang, Q., Lin, Z., and Hu, H. (2011a). Bidirectional control of social hierarchy by synaptic efficacy in medial prefrontal cortex. Science *334*, 693–697.

Wang, F., Kessels, H.W., and Hu, H. (2014). The mouse that roared: neural mechanisms of social hierarchy. Trends Neurosci. *37*, 674–682.

Wang, J., Zhang, K., Xu, L., and Wang, E. (2011b). Quantifying the Waddington landscape and biological paths for development and differentiation. Proc. Natl. Acad. Sci. U. S. A. *108*, 8257–8262.

Widloski, J., and Fiete, I.R. (2014). A model of grid cell development through spatial exploration and spike time-dependent plasticity. Neuron *83*, 481–495.

Zeng, H., and Sanes, J.R. (2017). Neuronal cell-type classification: challenges, opportunities and the path forward. Nat. Rev. Neurosci. *18*, 530–546.