On the expression of co-operative feeding behaviour in 3rd instar *Drosophila melanogaster* larvae!

Lucas Khodaei[1], Tara Newman[2], Samantha Lum[2], Henry Ngo[2], Matthew Maoloni[1] & Tristan A.F. Long[1]*

[1] Department of Biology, Wilfrid Laurier University, 75 University Ave. Waterloo, Ontario, N2L 3C5.
[2] Department of Health Science, Wilfrid Laurier University, 75 University Ave. Waterloo, Ontario, Canada, N2L 3C5.

*Corresponding author: tlong@wlu.ca

Abstract: Under poor nutritional conditions, 3rd instar *Drosophila melanogaster* larvae will work collaboratively in feeding clusters to obtain resources that cannot be reached individually. To better understand the conditions that influence the expression of this behaviour we examined the frequencies, the size and the membership in vials of flies that were initially seeded with either 100 or 200 eggs each using flies from both a large, outbred population and a replicate population that was homozygous for the *bw* allele. Overall, more feeding clusters, containing more larval participants were observed in the higher density vials compared to the lower density vials, consistent with the idea that this social behaviour is a response to dwindling resources in the environment. The presence of the *bw* allele did not result in greater egg-to-adult mortality, nor did it result in lower participation in feeding clusters.

Keywords: social behaviour, group feeding, foraging, density, survivorship, *Drosophila melanogaster*, larvae,

1. Introduction:

The fruit fly, *Drosophila melanogaster*, has a rich and celebrated history as a model organism in scientific research, and whose study has contributed to some of the greatest biological discoveries of all time (Weiner 1999, Brookes 2001). In recent years, it has also become a popular model species in which the ultimate and proximate causes of social behaviour can be examined (*e.g.* Wu *et al.* 2003, Sokolowski 2010, Durisko & Dukas 2013, Durisko *et al.* 2014, Camiletti & Thompson 2016).

Of the many interesting behaviours exhibited by *D. melanogaster*, our study focuses on the expression of social feeding among 3rd instar larvae (*see* Dombrovski *et al.* 2017). In this species, females lay eggs in fermenting vegetation (Reaume & Sokolowski 2006), where their offspring are often confined until their eclosion as adults (Sokolowski 1985). The structural and nutritional environment experienced by larval *D. melanogaster* often deteriorates over time, through the combined action of their burrowing and feeding, the buildup of their waste products, the effects of secreted digestive enzymes, and the activity of the microbial community. While desirable resources may potentially be found at lower stratums, the liquidification of the upper layers of the larvaes' habitat makes it difficult for them to access them, as they risk suffocation (Dombrovski *et al.* 2017). In these situations, individuals aggregate to form "feeding clusters" by synchronizing their digging movements to dig a 'well' (*sensu* Louis & de Polavieja 2017) that allows them to gain access to deeply buried resources, while still allowing respiration (using their posterior breathing spiracles) via a common air cavity (Wu *et al.* 2003, Justice *et al.* 2012, Dombrovski *et al.* 2017).

This study was undertaken to better understand some of the factors that may potentially influence the expression of this group feeding behaviour, as part of a larger investigation of this social phenomenon (Khodaei & Long 2019). Specifically, we set out to examine whether this phenomenon was influenced by larval density, and/or the genetic background of the larvae.

First, we explored the extent to which the formation and appearance of feeding clusters is associated with the local density of larvae. The nutritional environment experienced by larvae is often influenced by the abundance of conspecifics, with higher densities associated with greater competition for resources, slower development, higher mortality, and worsened individual condition (*reviewed in* Ashburner *et al.* 2005, Tennant *et al.* 2014). Since the local density of larvae is likely to be strongly associated with the rate/amount of resource depletion and/or environmental degradation, it is possible that larvae will be more likely to engage in group foraging at higher densities in order to

access needed resources. However, as *D. melanogaster* larval cannibalism is
expressed in times of nutritional stress and can involve individuals consuming
other larvae (Vijendravarma *et al.* 2013), adult carcasses (Yang 2018) and/or eggs
(Ahmad *et al.* 2015; *but see* Narasimha *et al.* 2019), at higher densities one
might alternatively see less social behaviour being expressed, if individuals
become more adverse to being in close contact with hungry conspecifics.
In their study of group feeding, Dombrovski *et al.* (2017) found that both
mechanosensory cues and vision are critical for the efficient co-ordination of
feeding behaviours between adjacent larvae. The importance of sight was
demonstrated in one of their assays as larvae from three different vision-
impaired mutant lines, norpA^{P41}, GMR-hid1, and GMR-hid2, formed fewer, and
smaller feeding clusters than flies from a wild-type population. In Khodaei &
Long (2019), our focal larvae were obtained from a *wild-type* population of *D.*
melanogaster, but competed in some of our assays with flies from a replicate
population where individuals homozygous for the *bw* allele (where adults have
brown coloured eyes). While adult *bw* homozygotes may have have
compromised vision that renders them selectively inferior to *wild-type*
individuals (*e.g.* MacLellan *et al.* 2009, but see Burnet *et al.* 1968), whether there
are any phenotypic differences with respect to larval group foraging is unknown.
92
2. Materials and methods
For our assays we used *Drosophila melanogaster* that we obtained from
two populations: *Ives* (hereafter 'IV'), and IV-*bw*. The IV population is an outbred,
wild-type population which originated from a sample of 200 mated females
collected in Amherst MA, USA in 1975 (Rose 1984). The IV-*bw* population was
created through the introgression of the recessive, brown-eyed allele, *bw* into
the IV genetic background via repeated back-crossing. Both of these
populations are kept at large sizes (~3500 adults per generation) in non-
overlapping generations, and are maintained under standardized conditions
(incubated at 25°C, 60% RH on a 12hr-L:12hr-D diurnal light cycle) (Martin &
Long 2015). Flies are cultured in narrow *Drosophila* vials containing 10mL of a
banana, killed-yeast and agar media, set at an initial density of approximately of
100 eggs/vial. Populations are cultured *en masse* every 14 days (Rose 1984,
Martin & Long 2015), by mixing all eclosed adult flies together under light CO_2
anesthetic, distributing them into vials containing fresh media and a small
amount of live yeast for 2-3 hours before being removed. The eggs laid in these
vials are then culled by hand to the appropriate density, and these vials are used
to found the next generation of culture.

In our experiment we set out to examine survivorship and co-operative
behaviour of both the IV and IV-*bw* populations at two initial egg density
treatments: 100 eggs/vial (the typical culture density), as well as 200 eggs/vial.
Under these more crowded conditions, IV flies may experience more resource
limitation (Tennant *et al.* 2014). We established 26 replicate vials for each of
these four experimental treatments. Eggs were obtained by placing adult IV and
IV-*bw* flies into oviposition chambers that contained grape juice/agar medium
(Sullivan *et al.,* 2000) for ~18h (overnight). From these surfaces we carefully
counted and transferred sets of 100 or 200 eggs into new vials containing 10ml
of standard media before being placed in the incubator.
Starting 24h after the experimental vials' creation, we began surveying
them for signs of larval cooperative feeding. Surveys were done on 3
consecutive days in 3 sessions/day, spaced 3h apart by a team of well-trained
observers who were blind to the treatment identity of the vials. These observers
counted the total number of feeding clusters present in each vial, where a
cluster was defined as a grouping of 3 or more larvae which were all feeding in a
downward direction creating a depression in the food media. Feeding clusters
were defined as being separate from one another if there was evidence that
larvae in the two groups were not working together (*i.e.* oriented in opposite
directions) and were separated from each other by at least 0.5mm. For those
clusters that had formed along the side of the vial, we measured the width (the
distance from one side of the foraging group to the other at the food surface
level) and depth (the distance from the food surface level to the lowest point in
the foraging group) of each of the clusters using a caliper to the nearest 0.5 mm.
We estimated the size of each of the cluster's depression observed as the area
of a triangle using our width and depth measurements, and then calculated the
mean cluster depression areas measured across all sessions. The total number of
larvae along the contour of each feeding cluster that were visible through the
transparent wall was also counted.
Fourteen days after the experimental vials' creation, all eclosed adult flies
were carefully removed from the vial using light CO_2 anesthesia and were
counted.
All statistical analyses were performed using R version 3. 3.2 (R Core
Team 2016) with vials as the unit of replication. We compared the egg-to-adult
survivorship rates using a generalized linear model (GLM), with a quasibinomial
error distribution, where the response variable was the number of flies (out of
the initial number of eggs in the vial) that survived to adulthood, and the
independent variables were the population source, the initial egg density and
their interaction. We determined the statistical significance of independent

variables using a likelihood-ratio Chi-square test using the *Anova* function in the
car package (Fox & Weisberg 2011).
We compared the frequency of cluster formation using a GLM with
quasipoisson error distribution, where the independent variables were the
population source, the initial egg density and their interaction, and the
dependent variable was the number of clusters in each vial summed across all
observation sessions. As above, we used a likelihood-ratio Chi-square test to
determine the statistical significance of independent variables.
As both the data on mean cluster size and mean number of larvae present
met parametric assumptions, we examined the effects of population origin,
initial egg density and their interaction using the Two-Way Analysis of Variance
(ANOVA) method, which was followed by a *post-hoc* Tukey HSD test to
compare the means of the treatments, if necessary.
163
3. Results
When comparing the rates of egg-to-adult survivorship across our
experimental treatments (Figure 2), we found a significant effect of density (LR
χ^2=172.57, df=1, p<0.0001), but no difference between the IV and IV-
populations (LR χ^2= 1.29, df=1, p=0.25) and no significant interaction between
population and density (LR χ^2=0.67, df=1, p=0.42).
The total number of feeding clusters (Figure 3) observed was also greater
in vials that had been seeded with 200 eggs compared to those from the 100
egg/vial treatment (LR χ^2=11.09, df=1, p=0.0009). We found no significant
difference between the vials that contained IV larvae and those which contained
IV-*bw* larvae (LR χ^2= 0.06, df=1, p=0.8), and no significant interaction between
the population origin and the egg density treatments (LR χ^2=0.21, df=1,
p=0.65).
When examining variation in the size of feeding clusters (Figure 4), we
found no significant effect of either population origin (F=1.73, df=1,99, p=0.19),
egg density (F=0.01, df=1,99, p=0.91) or their interaction (F=2.70, df=1,99,
p=0.10).
Finally, when we examined the number of larvae found in each cluster
(Figure 5) the main effect of population origin was not significant (F=0.70,
df=1,99, p=0.40). There were, however, more larvae on average in clusters in
those vials that had been seeded with 200 eggs compared to those from the
100 egg/vial treatment (F=14.85, df=1,99, p=0.0002) and there was a significant
interaction between the two main factors (F=11.15, df=1,99, p=0.001).
Specifically, we observed significantly more larvae, on average, in clusters from

the IV-*bw* 200 treatment vials, compared to the three other treatment
combinations ($\bar{x}$ IV-*bw* 100 = 11.18, $\bar{x}$ IV-*bw* 200 = 14.29, $\bar{x}$ IV 100 = 12.26, $\bar{x}$ IV 200 = 12.47).
190

4. Discussion

In this study we set out to examine whether the manifestation of
collaborative feeding clusters in 3rd instar *D. melanogaster* larvae was influenced
by the local density of conspecifics in their environment and/or whether there
were differences between larvae obtained from two populations that differed at
the *bw* locus. Overall, we saw that in the more resource-depleted high-density
vials, that co-operative feeding clusters were more commonly observed.
Furthermore, we saw no evidence that larvae homozygous for *bw* allele were
less likely to participate in group feeding behaviours.
Increasing the density of eggs in the vials clearly led to an environment
that was more challenging to the development of our *D. melanogaster* larvae,
which was reflected in our lower egg-to-adult survival rates in those vials that
had been initially seeded with 200 eggs compared to those that had initially
contained 100 eggs apiece (Figure 2). This is consistent with many other studies
that have documented the adverse effects associated with the more inhospitable
environment that results from increased conspecific densities (*reviewed in*
Ashburner *et al.* 2005, Tennant *et al.* 2014). Successful larval development in *D.*
melanogaster requires individuals to ingest between 3 to 5 times their body
weight in yeasts and decaying matter each day (Chiang & Hodson 1950), and
consequently resources will become depleted more quickly as the number of
developing larvae within a vial increases. Thus, seeing more feeding clusters
within the high-density vials (Figure 3) is indicative that the cooperative feeding
behaviour is dependent on the quality of the larvae's environment. The
burrowing action of larvae causes the upper layers of their media to become
both nutrient-poor, and increasingly liquefied (Dombrovski *et al.* 2017), which
would only be exacerbated by larger numbers of conspecifics.
Despite observing more clusters in higher density vials, we did not see
any statistically significant difference in the size of the feeding clusters compared
to the lower density vials (Figure 4). This somewhat surprising negative result
suggests several possibilities, which may be worth pursuing in future studies.
First, there may be an optimal size for feeding clusters, where individuals obtain
the greatest resources in return for their efforts, and that if clusters grow beyond
that point, individuals may move to smaller clusters with greater net benefits.
Secondly, the upper layers of media in the higher density vials may be more
fluid than those in the lower density vials, and whose structural instability may
prevent larvae in the 200 egg vials from successfully excavating deeper wells.

Finally, the 3rd instar larvae in the higher density vials may be of worse
physiological condition (due to more limited nutritional resources) and may be
less capable of the sustained, co-ordination of their movements needed to
create a larger air cavity.
When comparing the number of larvae within each cluster (Figure 5) we
observed that, on average, feeding clusters in the higher density vials had ~1.67
more members than those from the lower density vials ($\bar{x}_{100}$ = 11.71, $\bar{x}_{200}$ =
13.38; 2 sample t-test, t = -3.66, df = 101, p = 0.0004), which could reflect the
greater motivation to work together in the more challenging environment and/or
the greater absolute number of larvae present in the higher density vials.
However, the difference between 100 and 200 egg vials was much larger in
those vials that contained IV-*bw* larvae than those that contained IV larvae,
which resulted in a significant population or origin x density treatment
interaction. It is unclear what may have contributed to this observed difference.
Apart from this one situation, we did not see any other evidence that larvae from
the IV-*bw* population differed in their larval feeding behaviour, or response to
differences in larval density from those larvae obtained from the IV population. If
being homozygous at the *bw* allele impaired the larvae's' ability to work
collectively, we would have expected to see fewer clusters, smaller clusters
and/or fewer larvae in clusters when larvae originated from the IV-*bw*
population. None of these predicted differences were observed. While the
phenotype associated with *bw* homozygosity can be deleterious to adults
(MacLellan *et al.* 2009, Colpitts *et al.* 2017), in our study, it does not appear to
interfere with their ability to feed collectively as larvae (Figures 3, 4, & 5), nor
have a deleterious effect on egg-to-adult survivorship (Figure 1).
This study contributes to the growing body of research examining the
proximate and ultimate causes of group feeding behaviour in *Drosophila*
melanogaster (Wu *et al.* 2003, Justice *et al.* 2012, Dombrovski *et al.* 2017, Louis
& de Polavieja 2017), and provides insight into the conditions that favour the
evolution of social behaviours.
257
258

259   **Data Accessibility Statement**: Please see the Supplemental Cluster Data and
260   Supplemental Survivorship Data files associated with this manuscript.
261
262   **Competing Interests Statement**: The authors report no competing interests.
263

264   **Author Contributions:** The project was conceived by TAFL and LK. Counting of
265   eggs, clusters, and/or adult flies was performed by LK, TN, SL, DH, MM and/or
266   TAFL. TAFL wrote the manuscript.
267

268   **Acknowledgements**: TAFL was funded with a Natural Sciences and Engineering
269   Research Council (NSERC) Discovery grant. We would like to thank the
270   anonymous referee who was the inspiration for these experiments. This work was
271   conducted at Wilfrid Laurier University, which exists on the traditional territory of
272   the Neutral, Anishnawbe, and Haudenosaunee peoples.
273

275
276
Figure 1: Photo depicting a *Drosophila melanogaster* larval feeding cluster with
lines illustrating how width and depth of each cluster were measured. The width
of each cluster was defined as the horizontal distance between the outermost
cluster group members, while the depth of each cluster was defined as the
vertical distance from the media surface level to the lowest point in the foraging
group. We also counted the number of larvae along the contour of the feeding
cluster that were visible through the transparent wall of the vial as an index of
how many larvae were co-operating.
285

286
Figure 2: Boxplots illustrating egg-to-adult survivorship rates in vials of
Drosophila melanogaster obtained from the IV or the IV-*bw* that were initiated
with either 100 or 200 eggs. Each box encloses the middle 50% of data (Inter-
Quartile Range, IQR), with the location of the median represented by a
horizontal line. Values >±1.5 x the IQR outside the box are considered outliers
and depicted as open circles. Whiskers extend to the largest and smallest values
that are not outliers.
294
295

296
297 **Figure 3:** Boxplots illustrating the total number of larval feeding clusters
298 observed in vials of *Drosophila melanogaster* obtained from the IV or the IV-*bw*
299 population that were initiated with either 100 or 200 eggs. Boxplot components
300 as described as in Figure 2.
301

302
303 **Figure 4.** Boxplots illustrating the mean size of all larval feeding clusters
304 observed in vials of *Drosophila melanogaster* obtained from the IV and IV-*bw*
305 populations that were initiated with either 100 or 200 eggs. Boxplot
306 components as described as in Figure 2.
307
308

309
310 **Figure 5.** Boxplots illustrating the mean number of larvae present in all larval
311 feeding clusters observed in vials of *Drosophila melanogaster* obtained from the
312 IV and IV-*bw* populations that were initiated with either 100 or 200 eggs.
313 Boxplot components as described in Figure 2.

References:

Ahmad, M., Chaudhary, S., Afzal, A., Tariq, M. (2015) Starvation-induced dietary behaviour in *Drosophila melanogaster* larvae and adults. Scientific Reports. 5(1). doi:10.1038/srep14285

Ashburner M., Golic K.G, & Hawley R.S. (2005). *Drosophila*: a laboratory handbook (2nd ed). Cold Spring Harbor, NY:Cold Spring Harbor Laboratory Press.

Brookes, M. (2001). Fly: the unsung hero of twentieth-century science. New York: Ecco.

Burnet, B., Connolly, K. and Beck, J., 1968. Phenogenetic studies on visual acuity in *Drosophila melanogaster*. Journal of Insect Physiology, 14(6), pp.855-860.

Camiletti, A., & Thompson, G. (2016). Drosophila as a genetically tractable model for social insect behavior. Frontiers in Ecology and Evolution, 4. doi: 10.3389/fevo.2016.00040

Chiang, H, & Hodson, A. (1950). An analytical study of population growth in *Drosophila melanogaster*. Ecological Monographs, 20(3), 173-206. doi: 10.2307/1948580

Colpitts, J., Williscroft, D., Sekhon, H.S. and Rundle, H.D., (2017). The purging of deleterious mutations in simple and complex mating environments. Biology Letters, 13(10), p.20170518.

Crespi, BJ., Yanega, D. 1995. The definition of eusociality. Behavioral Ecology. 6(1):109-15.

Louis, M., & de Polavieja, G. (2017). Collective behavior: Social digging in *Drosophila* larvae. Current Biology, 27(18), R1010-R1012. doi: 10.1016/j.cub.2017.08.023

Dombrovski, M., Poussard, L., Moalem, K., Kmecova, L., Hogan, N., Schott, E., Vaccari, A., Acton, S., & Condron B. (2017). Cooperative behavior emerges among *Drosophila* larvae. Current Biology, 27, 2821-2826. doi: 10.1016/j.cub.2017.07.054

Durisko, Z., & Dukas, R. (2013). Attraction to and learning from social cures in

fruitfly larvae. Proceedings of the Royal Society B, 280, 20131398. doi:

10.198/rspb.2013.1398

Durisko, Z., Kemp, R., Mubasher, R., & Dukas, R. (2014). Dynamics of social

behavior in fruit fly larvae. PLoS ONE, 9(4), e95495. doi:

10.1371/journal.pone.0095495

Fox, J., & Weisberg, S. (2011). An {R} Companion to Applied Regression, Second

Edition. Thousand Oaks CA: Sage. URL:

http://socserv.socsci.mcmaster.ca/jfox/Books/Companion

Justice, E.D, Macedonia, N.J., Hamilton, C., & Condron, B. (2012). The simple fly

larval visual system can process complex images. Nature Communications, 23

(3),1156.

Khodaei, L., & Long, T.A.F. (2019) Kin recognition and co-operative foraging in

Drosophila melanogaster larvae. Journal of Evolutionary Biology, 32(12): 1352-

1361. doi: /10.1111/jeb.13531

Martin, E., & Long, T.A.F. (2015). Are flies kind to kin? The role of intra- and

inter-sexual relatedness in mediating reproductive conflict. Proceedings of the

Royal Society B, 282(1821), 20151991. doi: 10.1098/rspb.2015.1991

MacLellan, K., Whitlock, M.C., & Rundle, H.D. (2009) Sexual selection against

deleterious mutations via variable male search success. Biology Letters 5 (6):795-

797.

R Core Team (2016). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL: https://www.R-

project.org/.

Narasimha, S., Nagornov, K., Menin, L., Mucciolo, A., Rohwedder, A., Humbel,

B., Stevens, M., Thum, A.S., Tsybin, Y.O., & Vijendravarma, R.K. (2019)

Drosophila melanogaster cloak their eggs with pheromones, which prevents

cannibalism. PLoS Biology 17(1): e2006012. doi: 10.1371/journal.pbio.2006012

Reaume, C.J., & Sokolowski, M.B. (2006). The nature of *Drosophila*
melanogaster. Current Biology, 16(16), R623-628.
394

395   Rose, M. (1984). Laboratory evolution of postponed senescence in *Drosophila*
396   *melanogaster*. Evolution, 38(5), 1004-1010. doi: 10.1111/j.1558-
397   5646.1984.tb00370.x
398

399   Sokolowski, M.B. (2010). Social interactions in "simple" model systems. Neuron,
400   65, 780–794.
401

402   Tennant M.E., Sonser, E., & Long, T.A.F. (2014). When the going gets tough, do
403   the tough get choosy? Condition-dependency of female choosiness in
404   *Drosophila melanogaster*. In Causes and consequences of female mate choice in
405   *Drosophila melanogaster*: A hemiclonal analysis. Tennant M.E, MSc Integrative
406   Biology Thesis 1645. https://scholars.wlu.ca/etd/1645
407

408   Vijendravarma, R.K., Narasimha, S., & Kawecki, T.J. (2013) Predatory cannibalism
409   in *Drosophila melanogaster* larvae. Nature Communications 4: 1-8.
410   https://doi.org/10.1038/ncomms2744
411

412   Weiner, J., (2000). Time, love, memory: a great biologist and his quest for the
413   origins of behavior. New York: Vintage Books
414

415   Wu, Q., Wen, T., Lee, G., Park, J., Cai, H., & Shen, P. (2003). Developmental
416   control of foraging and social behavior by the *Drosophila* neuropeptide Y-like
417   system. Neuron, 39(1), 147-161. doi: 10.1016/s0896-6273(03)00396-9
418

419   Yang, D. (2018) Carnivory in the larvae of *Drosophila melanogaster* and other
420   *Drosophila* species. Scientific Reports, 8(1). doi:10.1038/s41598-018-33906-w
421