Big data analysis of all bacterial genomes establishes a triple whammy of carbapenemases, ICEs and multiple clinically-relevant bacteria

João Botelho[1#*], Joana Mourão[2,3], Adam P. Roberts[4,5], Luísa Peixe[1*]

[1]UCIBIO/REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal.

[2]Center for Neuroscience and Cell Biology, University of Coimbra, Coimbra, Portugal.

[3]Institute for Interdisciplinary Research, University of Coimbra, Coimbra, Portugal.

[4]Department of Tropical Disease Biology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom.

[5]Centre for Drugs and Diagnostics, Liverpool School of Tropical Medicine, Liverpool, United Kingdom.

[#]New affiliation: Antibiotic Resistance Evolution Group, Max-Planck-Institute for Evolutionary Biology, 24306 Plön, Germany; Department of Evolutionary Ecology and Genetics, Zoological Institute, Christian-Albrechts-Universität zu Kiel, 24118 Kiel, Germany.

*Addresses for correspondence: João Botelho, Antibiotic Resistance Evolution Group, Max-Planck-Institute for Evolutionary Biology, 24306 Plön, Germany; Department of Evolutionary Ecology and Genetics, Zoological Institute, Christian-Albrechts-Universität zu Kiel, 24118 Kiel, Germany.; e-mail: botelho@evolbio.mpg.de; Luísa Peixe, Laboratório de Microbiologia. Faculdade de Farmácia da Universidade do Porto, Rua Jorge Viterbo Ferreira nº 228, 4050-313 Porto, Portugal; phone: +351 220428500; e-mail: lpeixe@ff.up.pt.

Abstract

Carbapenemases inactivate most β-lactam antibiotics, including carbapenems and have

been frequently reported among *Enterobacteriaceae*, *Acinetobacter* spp. and

Pseudomonas spp. Traditionally, the horizontal gene transfer of carbapenemase-

encoding genes (CEGs) has been linked to plasmids. However, given that integrative and

conjugative elements (ICEs) are possibly the most abundant conjugative elements

among prokaryotes, we conducted an *in silico* analysis to ascertain the likely role of ICEs

in the spread of CEGs among all bacterial genomes. We detected 7457 CEGs, of which

224 were located within putative ICEs among several clinically-relevant bacterial species

(including *Klebsiella pneumoniae*, *Escherichia coli, Citrobacter freundii, Enterobacter*

hormaechei and *Pseudomonas aeruginosa*). We also identified a *Bacillus subtilis* strain

with an NDM-1-encoding ICE. Most CEGs detected within ICEs belong to the KPC, IMP

and NDM families and different mechanisms were likely responsible for acquisition of

these genes. The majority of CEG-bearing ICEs belong to the MPF_T, MPF_F, MPF_G and MPF_I

classes and often encode additional adaptive traits such as resistance to other

antibiotics (e.g. aminoglycosides and fluoroquinolones), metal and biocide tolerance

and bacteriocin production. This study provides a snapshot of the different CEGs

associated with ICEs among all bacterial genomes and sheds light on the

underappreciated contribution of ICEs to the spread of carbapenem resistance globally.

Introduction

Due to the importance of carbapenems for the treatment of serious infections in humans, the WHO has stated these antibiotics should be reserved for infections caused by multidrug resistant Gram-negative bacteria in humans [1]. Recently, the same agency presented a list of bacterial pathogens for which research and development of new antibiotics are urgently required, and the top priority pathogens were the carbapenem-resistant strains of *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacteriaceae* [2].

The evolution of carbapenem-resistance in bacteria is often driven by the horizontal gene transfer (HGT) of carbapenemase-encoding genes (CEGs) [3]. Carbapenemases are beta-lactamases able to hydrolyze carbapenems as well as most of other beta-lactam antibiotics. The CEGs are often located on integrons or transposons that themselves target mobile genetic elements (MGEs) such as plasmids [3, 4], which makes the dissemination of these genes unpredictably complex within bacterial communities. Recently, we demonstrated that another type of MGE, the integrative and conjugative elements (ICEs), play a significant role as vehicles for the dissemination of CEGs among *P. aeruginosa* [5]. ICEs are self-transmissible MGEs that can integrate into and excise from the genome (as transposons and phages do) and can exist as circular, sometimes replicable, extrachromosomal elements and be transferred by conjugation (as plasmids do) [6–9]. ICEs appear to have a bipartite lifestyle that shifts between vertical and horizontal transmission [7, 10]. Eight mating pair formation (MPF) classes were proposed (B, C, F, FA, FATA, G, I and T), based on the phylogeny of VirB4, the only ubiquitous protein in all known type-IV secretion systems (T4SS) involved in conjugation [11]. The MPF_T is widely distributed both in conjugative plasmids and ICEs, while MPF_F is more prevalent in plasmids and MPF_G on ICEs [12].

Given that ICEs were identified in most bacterial clades and were proposed to be more prevalent than conjugative plasmids [6], we conducted an *in silico* analysis to explore the distribution of CEG-bearing ICEs among all sequenced bacterial genomes. Our results demonstrate that CEG-bearing ICEs mainly belong to three MPF families and are primarily located in several bacterial pathogens. Our analysis highlights the importance

of thoroughly investigating these elements as important vehicles for the spread of

antibiotic resistance (AR), particularly to carbapenems.

Material and methods

Bacterial genomes and carbapenemases search

All RefSeq bacterial genomes available in NCBI (accessed on 02/03/2018), including

complete and draft genome sequences were downloaded and blasted against a local

database of 688 carbapenemases (**Table S1**) adapted from AMRfinder

(https://www.ncbi.nlm.nih.gov/pathogens/antimicrobial-resistance/AMRFinder/) [13],

using diamond v0.9.21 (https://github.com/bbuchfink/diamond) [14] with the following

parameters: 'diamond blastx –d DB.dmnd –o hits.txt --id 100 --subject-cover 100 -f 6 --

sensitive'.

Tracing ICEs among the bacterial genomes

The CEG-bearing genomes were annotated with prokka v1.13

(https://github.com/tseemann/prokka) [15], and the GenBank files were used as input

to mine ICEs on the standalone version of ICEfinder

(http://202.120.12.136:7913/ICEfinder/ICEfinder.html). Putative MGEs were annotated

as ICEs if complete conjugation and relaxase modules, plus an integrase or DDE

transposase superfamily were identified within the boundaries of the element, and as

integrative and mobilizable elements (IMEs) if the elements contained all except genes

involved in conjugation. The translated coding sequences of extracted MGEs were

analyzed on the CONJscan module of MacSyFinder (https://github.com/gem-

pasteur/macsyfinder) [16, 17]. A MGE was classified as an ICE when a positive prediction

was verified by ICEfinder and/or the CONJscan module. To predict the multi-locus

sequence type of each CEG-bearing MGEs, we used mlst v2.16.1

(https://github.com/tseemann/mlst), which scans the genomes against PubMLST typing

schemes (https://pubmlst.org/) [18].

Characterization of the CEG-bearing ICEs

Screening of AR genes among ICEs was performed using ResFinder v3.1

(https://cge.cbs.dtu.dk/services/ResFinder/) [19]. Metal and biocides tolerance genes

were blasted against a protein database downloaded from BacMet

(http://bacmet.biomedicine.gu.se) [20] using the aforementioned diamond parameters.

Associated MGEs were annotated using Galileo™ AMR

(https://galileoamr.arcbio.com/mara/) (Arc Bio, Cambridge, MA) [21]. Bagel4

(http://bagel4.molgenrug.nl/) was used to mine ICEs for bacteriocins and ribosomally

synthesized and posttranslationally modified peptides [22]. We ran our extracted ICEs

against REBASE (http://rebase.neb.com/rebase/rebase.html) to look for restriction-

modification systems [23]. We also used CRISPRCasFinder (https://crisprcas.i2bc.paris-

saclay.fr/CrisprCasFinder/Index) to look for CRISPR (clustered regularly interspaced

short palindromic repeats) arrays and their associated (Cas) proteins within ICE

sequences [24].

Results

Carbapenemase-encoding genes are mainly found in Gram-negative bacteria

We retrieved a total of 107,887 bacterial genomes from NCBI and we identified 7457

CEGs belonging to 33 beta-lactamase families among 7343 genomes (**Table S2**). Most

CEG-bearing genomes were detected in the USA, China and Europe (**Figure 1A** and **Table**

S2). Also, the majority of the bacterial genomes were sequenced in these countries. A

total of 43 genera were detected, although dominated by a limited number of bacterial

genus. Our results show that CEGs are mostly present on Gram-negative bacteria.

Exceptions included the report of OXA-72 in *Staphylococcus aureus* and NDM-1 in

Bacillus subtilis. As suspected, we observed that the ESKAPEEc pathogens *K.*

pneumoniae, *A. baumannii*, *P. aeruginosa, Enterobacter spp.* and *Escherichia coli* carry a

wide diversity of CEG families including the most promiscuous OXA, KPC, NDM, IMP, VIM

and GES-types (**Figure 1B**). In addition, we observed that several CEGs (such as bla_{SPM-1})

have a narrow host range and were only identified within strains belonging to the same

genus (**Figure 1C**). Even though carbapenemases of the DIM family were not identified

on bacterial genomes other than the *Pseudomonas* genus, a previous study has reported

the presence of bla_{DIM-1} in *Enterobacteriaceae* [25].

A

B

C

Figure 1. Dispersion of carbapenemase-encoding bacterial genomes. **A.** Worldwide
distribution of carbapenemase families. The size of the circles reflects the number of
hits. **B.** Distribution of carbapenemases observed in more than one genus. The legend
for the most frequently identified carbapenemases and bacterial genus is present on the
left and right axis, respectively. **C.** Representation of carbapenemases identified in only
one bacterial genus. Only carbapenemases that were identified more than 4 times were
included on the map and on the graphs. For a complete list of all carbapenemases,
please refer to **Table S2**.

A large proportion of CEG-bearing ICEs belong to three MPF families and target
clinically-relevant gram-negative bacteria

Our analysis with ICEfinder resulted in the identification of 251 CEGs within 248 putative
MGEs (**Table S3 and Table S4**). Of these, 224 putative ICEs were predicted. The
remaining 24 MGEs were considered as IMEs or putative conjugative regions. The later
designation is attributed to regions where an integrase and a qualified T4SS but no
relaxase or a type-IV coupling protein were identified. Using the CONJscan module of

MacSyFinder, we noted that the majority of the ICEfinder hits belong to three MPF

families: MPF_T, MPF_F and MPF_G (**Figure 2** and **Table S3**). ICEs of the MPF_T class were

particularly promiscuous, being responsible for the spread of several CEGs such as $bla_{KPC-2/3}$,

bla_{NDM-1} and bla_{IMP-1} among clinically-relevant pathogens as *P. aeruginosa*, *E. coli* and

species belonging to the *Klebsiella* genus and the *Enterobacter cloaceae* complex. The

bla$_{SPM-1}$ gene was exclusively identified in ICEs of the MPF_T class.

The bacterial hosts housing these elements belong to more than 70 sequence types

(STs), of which the *K. pneumoniae* ST258 is the most frequently represented (**Figure 3**

and **Table S3**). The worldwide spread of carbapenemase-producing *K. pneumoniae* has

been linked to some high-risk clones, including ST258 and close relatives that were also

identified in this study, such as ST11 and ST512 [26, 27]. We also reported several strains

belonging to ST78 from the *Enterobacter hormaechei*, recognized as a global driver of

CEGs [28, 29].

Figure 2. Sankey diagram showing the contribution of different MPF families for the spread of CEGs among several bacteria. *Aci*sp stands for

Acinetobacter sp., *A*sp for *Achromobacter* sp., *Bs* for *Bacillus subtilis*, *Cf* for *Citrobacter freundii*, *C*sp for *Citrobacter* sp., *Ea* for *Elizabethkingia*

anophelis, *Ec* for *E. coli*, *Ecc* for *E. cloaceae* complex, *Ka* for *Klebsiella aerogenes*, *Km* for *Klebisella michiganensis*, *Ko* for *Klebsiella oxytoca, Kp* for

K. pneumoniae, *Kq* for *Klebsiella quasipneumoniae*, *Kv* for *Klebsiella variicola*, *Mm* for *Morganella morganii*, *Pa* for *P. aeruginosa* and Pm for

Proteus mirabilis.

Figure 3. Sequence types (STs) of the ICEfinder hits, including ICEs, IMEs and putative

conjugative regions. Only STs that were identified more than once were included on the

graph. For a complete list of all STs, please refer to **Table S3**.

A variable repertoire of CEG-bearing integrons and transposons target ICEs

We identified 20 CEG variants among the 251 hits, dominated by bla_{KPC-2}, bla_{KPC-3}, bla_{IMP-1} and bla_{NDM-1}, and rare descriptions of bla_{KPC-4}, $bla_{IMP-10/13}$, $bla_{VIM-1/34}$, bla_{DIM-1}, bla_{SPM-1} and $bla_{OXA-23/48/181}$ (**Figure 4** and **Table S3**). KPC-encoding ICEs although mostly observed in *Klebsiella* spp. were also identified in *E. coli*, *Enterobacter* spp., *Citrobacter* spp. and *Aeromonas* spp. IMP-encoding ICEs were restricted to *Enterobacter* spp. and *Pseudomonas* spp.

The $bla_{KPC-2/3}$ genes were spread mainly by MPF_T and MPF_G classes (**Figure 2**). We noted that composite transposons (e.g. ISCR4 and ISCR24) were frequently linked to the acquisition of bla_{SPM-1}, bla_{NDM-1} and bla_{OXA-48} [30, 31], while bla_{IMP} and bla_{VIM} were found on class I integrons [3]. In addition, we observed that bla_{KPC} genes were frequently flanked by the insertion sequences (ISs) IS*Kpn7* and IS*Kpn6* within the Tn*4401* transposon, which is capable of conferring a high frequency of transposition [32]. Interestingly, we found a bla_{NDM-1} gene associated with a MPF_T class ICE on a *B. subtilis* strain (**Figure 2** and **4** and **Table S3**). This is the only situation where we found a CEG-bearing ICE on Gram positive bacteria. Besides CEGs, the ICEs also harbor genes conferring resistance to other antibiotics, such as aminoglycosides, fluoroquinolones, macrolides, sulfonamides and trimethoprim (**Table S5**), widening the spectrum of transmissible AR genes selectable by carbapenems due to linkage.

Figure 4. Distribution of carbapenemases among ICEfinder hits, including ICEs, IMEs and putative conjugative regions. Different colour links are related to the number of associations between a carbapenemase and a given bacterial species. The size of circles is proportional to the number of times each carbapenemase or bacterial strain was identified. *Acib* stands for *A. baumannii*, *Acip* for *Acinetobacter pittii*, *Aci*sp for *Acinetobacter* sp., *Aer*sp for *Aeromonas* sp., *Ah* for *Aeromonas hydrophila*, *A*sp for *Achromobacter* sp., *Bs* for *Bacillus subtilis*, *Cf* for *Citrobacter freundii*, *C*sp for *Citrobacter* sp., *Ea* for *Elizabethkingia anophelis*, *Ec* for *E. coli*, *Ecc* for *E. cloaceae* complex, *Ka* for *Klebsiella aerogenes*, *Km* for *Klebisella michiganensis*, *Ko* for *Klebsiella oxytoca, Kp* for *K. pneumoniae*, *Kq* for *Klebsiella quasipneumoniae*, *Kv* for *Klebsiella variicola*, *Mm* for *Morganella morganii*, *Pa* for *P. aeruginosa*, Pm for *Proteus mirabilis* and *P*sp for *Pseudomonas* sp.

Acquisition of additional traits by ICEs include competitive weapons as bacteriocins

Besides genes conferring antibiotic resistance, some of the CEG-bearing ICEs harbor other cargo genes that may confer a selective advantage to the ICE host, as bacteriocins and restriction-modification systems. We identified class III bacteriocins among 6 *K. pneumoniae* ST258 strains and associated with MPF$_F$ and MPF$_T$ classes (**Figure S1** and **Table S6**). Additionally, our results show that some representatives of this ST and other *Enterobacteriaceae* (e.g. *E. coli*, *Enterobacter* sp.) strains carry type I and II restriction-modification systems or "orphan" methyltransferases (**Table S7**). Ammonium quaternary compounds and metal tolerance genes, such as the *qac* (ammonium quaternary), *mer* (mercury), *ars* (arsenic) and *cus/czc* (copper) operons were also found in several genera, including *Enterobacter*, *Klebsiella* and *Pseudomonas* (**Table S8**). No CRISPR-Cas systems were identified within these elements.

Discussion

We have set out to comprehensively identify the CEGs among all bacterial genomes deposited in NCBI and the CEG-bearing ICE sequences. Our study considerably expands the knowledge of the repertoire of CEGs-bearing ICEs. We uncovered more than 200 putative MGEs that may be involved in HGT of CEGs among all bacterial genomes. To expand our predictions, we also used the CONJscan module of Macsyfinder to trace the MPF families likely involved in spreading these genes.

Even though CEGs might rapidly spread worldwide, local selection is likely required for them to reach fixation, as can be seen for the clonal expansion of *K. pneumoniae* ST258 harboring *bla*$_{KPC-2/3}$ [26, 27]. The scenario observed for the acquisition of the most important CEGs by ICEs (Tn*4401* for *bla*$_{KPC-2/3}$, composite transposons for *bla*$_{NDM-1}$ and class I integrons for *bla*$_{IMP}$ and *bla*$_{VIM}$) resembles that of plasmids [3] and provide additional support to the notion that the line separating these elements is blurred [8, 9, 33]. We now show that besides plasmids, this promiscuous repertoire of integrons and transposons frequently targets ICEs of different MPF families. Surprisingly, we noted that only *bla*$_{SPM-1}$ was unable to spread beyond the same clonal lineage [31]. Indeed, all hits were identified in *P. aeruginosa* ST277 strains from Brazil and within a MPF$_T$ ICE

family, indicating that likely these ICEs are only transferring vertically. Understanding

the limitations on this family of ICE's horizontal gene transfer may be translatable to

other, more transferable, ICEs and could underpin a control strategy to prevent the

spread of these elements in the future.

CEGs are rarely identified in Gram-positive bacteria. Local selection and enriched

metagenome might have also contributed to the ICE encoding NDM-1 observed in a *B.*

subtilis from a highly polluted marine environment [34]. Despite the works on ICEs in

Gram positive bacteria [35, 36], this first report of a CEG-bearing ICE in ubiquitous and

resilient bacterial groups is of remarkable epidemiological interest.

In addition to AR, ICEs can be involved in other traits such as heavy metal resistance,

carbon-source utilization, symbiosis, restriction-modification and bacteriocin synthesis

[9]. Since bacteria commonly inhabit highly competitive environments, the production

of toxins (such as bacteriocins) confers a selective advantage to the host [37]. Class III

bacteriocins – bacteriolysins - consists of high-molecular-weight and heat-labile toxins

[38]. We speculate that the presence of these toxins within the ICEs of six *K. pneumoniae*

ST258 strains can promote ICE stability by preferentially selecting for cells harboring the

ICE. Restriction-modification systems protect a bacterial host against the invasion of

foreign DNA through the action of a restriction enzyme that cuts DNA lacking the

epigenetically methylated DNA modified by the methyltransferase [39]. The presence of

these systems within ICEs may also allow their stable maintenance through

postsegregational killing. This has already been demonstrated for plasmids [40].

Additionally, these systems may also prevent further infection of the bacterial host by

another ICE or MGE lacking the methylation.

One caveat of our studies is that these results do not yet expose the complete set of

CEG-bearing ICEs present in all bacteria. There is an inherent bias in the number of times

we detect a particular CEG in certain pathogen genomes as some are over represented

in the database compared to others. It is possible that certain observations will flatten

out as more genomes are analyzed. This analysis also underestimates the extent of host

range because we only used assembled genomes. Less than 10% of the bacterial

genomes currently present on NCBI are complete. This is a major drawback since

ICEfinder can only detect ICEs within complete genomes or contigs with or without

directed repeats as the tRNA-distal boundaries, meaning that if the ICE was integrated
into the 3' or 5' ends of protein-encoding genes and/or split into several contigs, this
tool may miss the element. Even though ICEfinder uses the profile Hidden Markov
Models (HMM) for signature proteins coded by ICEs, we identified some plasmid hits
among our data using PlasmidFinder (**Table S9**) [41]. Also, ICEfinder may not provide the
accurate boundaries of ICEs. Indeed, the relaxase and the T4SS encoded by ICEs
resemble those of plasmids [6, 8, 11]. Plus, it is possible that ICEs and plasmids have
swapped conjugation modules along their evolutionary history [6]. We believe that a
more thorough exploration of this issue, especially regarding the precise delimitation of
ICEs, will be an important further step toward an improved understanding of the
contribution of these elements to bacterial adaptation and evolution of AR.

While we have chosen to focus on CEG-bearing genomes, our computational approach
can be applied to trace other relevant AR genes and other cargo genes that may confer
a selective advantage to the ICE host. Leveraging knowledge linking the accurate
prediction of ICE sequences to the carriage of AR genes, will not only improve our
understanding of HGT, but may also uncover potential approaches to tackle the spread
of AR.

Acknowledgments

This work was supported by the Applied Molecular Biosciences Unit- UCIBIO which is
financed by national funds from FCT/MCTES (UID/Multi/04378/2019).

References

1. EFSA Panel on Biological Hazards (BIOHAZ). Scientific Opinion on Carbapenem
resistance in food animal ecosystems. *EFSA J* 2013; **11**.

2. World Health Organization. GLOBAL PRIORITY LIST OF ANTIBIOTIC-RESISTANT
BACTERIA TO GUIDE RESEARCH, DISCOVERY, AND DEVELOPMENT OF NEW
ANTIBIOTICS. http://www.who.int/news-room/detail/27-02-2017-who-

publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed.

Accessed 11 Aug 2018.

3.   Partridge SR, Kwong SM, Firth N, Jensen SO. Mobile Genetic Elements

Associated with Antimicrobial Resistance. *Clin Microbiol Rev* 2018; **31**: e00088-

17.

4.   Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B,

Guerra B, et al. Plasmids carrying antimicrobial resistance genes in

Enterobacteriaceae. *J Antimicrob Chemother* 2018; **73**: 1121–1137.

5.   Botelho J, Roberts AP, León-Sampedro R, Grosso F, Peixe L. Carbapenemases on

the move: it's good to be on ICEs. *Mob DNA* 2018; **9**: 37.

6.   Guglielmini J, Quintais L, Garcillán-Barcia MP, de la Cruz F, Rocha EPC. The

Repertoire of ICE in Prokaryotes Underscores the Unity, Diversity, and Ubiquity

of Conjugation. *PLoS Genet* 2011; **7**: e1002222.

7.   Carraro N, Poulin D, Burrus V. Replication and Active Partition of Integrative and

Conjugative Elements (ICEs) of the SXT/R391 Family: The Line between ICEs and

Conjugative Plasmids Is Getting Thinner. *PLOS Genet* 2015; **11**: e1005298.

8.   Cury J, Touchon M, Rocha EPC. Integrative and conjugative elements and their

hosts: composition, distribution and organization. *Nucleic Acids Res* 2017; **45**:

8943–8956.

9.   Johnson CM, Grossman AD. Integrative and Conjugative Elements (ICEs): What

They Do and How They Work. *Annu Rev Genet* 2015; **49**: 577–601.

10.   Delavat F, Miyazaki R, Carraro N, Pradervand N, van der Meer JR. The hidden life

of integrative and conjugative elements. *FEMS Microbiol Rev* 2017; **41**: 512–537.

11.   Guglielmini J, de la Cruz F, Rocha EPC. Evolution of Conjugation and Type IV

Secretion Systems. *Mol Biol Evol* 2013; **30**: 315–331.

12.   Guglielmini J, Néron B, Abby SS, Garcillán-Barcia MP, de la Cruz F, Rocha EPC.

Key components of the eight classes of type IV secretion systems involved in

bacterial conjugation or protein secretion. *Nucleic Acids Res* 2014; **42**: 5715–27.

13. Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I, et al. Using the
NCBI AMRFinder Tool to Determine Antimicrobial Resistance Genotype-
Phenotype Correlations Within a Collection of NARMS Isolates. *bioRxiv* 2019;
550707.

14. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using
DIAMOND. *Nat Methods* 2015; **12**: 59–60.

15. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;
30: 2068–2069.

16. Abby SS, Cury J, Guglielmini J, Néron B, Touchon M, Rocha EPC. Identification of
protein secretion systems in bacterial genomes. *Sci Rep* 2016; **6**: 23080.

17. Abby SS, Néron B, Ménager H, Touchon M, Rocha EPC. MacSyFinder: A Program
to Mine Genomes for Molecular Systems with an Application to CRISPR-Cas
Systems. *PLoS One* 2014; **9**: e110726.

18. Jolley KA, Maiden MCJ. BIGSdb: Scalable analysis of bacterial genome variation
at the population level. *BMC Bioinformatics* 2010; **11**: 595.

19. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al.
Identification of acquired antimicrobial resistance genes. *J Antimicrob
Chemother* 2012; **67**: 2640–2644.

20. Pal C, Bengtsson-Palme J, Rensing C, Kristiansson E, Larsson DGJ. BacMet:
antibacterial biocide and metal resistance genes database. *Nucleic Acids Res*
2014; **42**: D737–D743.

21. Partridge SR, Tsafnat G. Automated annotation of mobile antibiotic resistance in
Gram-negative bacteria: the Multiple Antibiotic Resistance Annotator (MARA)
and database. *J Antimicrob Chemother* 2018; **73**: 883–890.

22. van Heel AJ, de Jong A, Song C, Viel JH, Kok J, Kuipers OP. BAGEL4: a user-
friendly web server to thoroughly mine RiPPs and bacteriocins. *Nucleic Acids Res*
2018; **46**: W278–W281.

23. Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE—a database for DNA restriction

and modification: enzymes, genes and genomes. *Nucleic Acids Res* 2015; **43**:

D298–D299.

24.   Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, et al.

CRISPRCasFinder, an update of CRISRFinder, includes a portable version,

enhanced performance and integrates search for Cas proteins. *Nucleic Acids Res*

2018; **46**: W246–W251.

25.   Leski TA, Bangura U, Jimmy DH, Ansumana R, Lizewski SE, Li RW, et al.

Identification of blaOXA-51-like, blaOXA-58, blaDIM-1, and blaVIM

Carbapenemase Genes in Hospital Enterobacteriaceae Isolates from Sierra

Leone. *J Clin Microbiol* 2013; **51**: 2435–2438.

26.   Chen L, Mathema B, Pitout JDD, DeLeo FR, Kreiswirth BN. Epidemic Klebsiella

pneumoniae ST258 Is a Hybrid Strain. *MBio* 2014; **5**: e01355-14.

27.   Bowers JR, Kitchel B, Driebe EM, MacCannell DR, Roe C, Lemmer D, et al.

Genomic Analysis of the Emergence and Rapid Global Dissemination of the

Clonal Group 258 Klebsiella pneumoniae Pandemic. *PLoS One* 2015; **10**:

e0133727.

28.   Peirano G, Matsumura Y, Adams MD, Bradford P, Motyl M, Chen L, et al.

Genomic Epidemiology of Global Carbapenemase-Producing *Enterobacter* spp.,

2008–2014. *Emerg Infect Dis* 2018; **24**: 1010–1019.

29.   Gomez-Simmonds A, Annavajhala MK, Wang Z, Macesic N, Hu Y, Giddins MJ, et

al. Genomic and Geographic Context for the Evolution of High-Risk Carbapenem-

Resistant *Enterobacter cloacae* Complex Clones ST171 and ST78. *MBio* 2018; **9**.

30.   Ding Y, Teo JWP, Drautz-Moses DI, Schuster SC, Givskov M, Yang L. Acquisition of

resistance to carbapenem and macrolide-mediated quorum sensing inhibition

by Pseudomonas aeruginosa via ICETn43716385. *Commun Biol* 2018; **1**: 57.

31.   Nascimento APB, Ortiz MF, Martins WMBS, Morais GL, Fehlberg LCC, Almeida

LGP, et al. Intraclonal Genome Stability of the Metallo-β-lactamase SPM-1-

producing Pseudomonas aeruginosa ST277, an Endemic Clone Disseminated in

Brazilian Hospitals. *Front Microbiol* 2016; **7**: 1946.

32.   Cuzon G, Naas T, Nordmann P. Functional characterization of Tn4401, a Tn3-
419   based transposon involved in blaKPC gene mobilization. *Antimicrob Agents*
420   *Chemother* 2011; **55**: 5370–3.

421   33.   Carraro N, Burrus V. The dualistic nature of integrative and conjugative
422   elements. *Mob Genet Elements* 2015; **5**: 98–102.

423   34.   Zeng W, Chen G, Tang Z, Wu H, Shu L, Liang Z. Draft Genome Sequence of
424   Bacillus subtilis GXA-28, a Thermophilic Strain with High Productivity of Poly-γ-
425   Glutamic Acid. *Genome Announc* 2014; **2**.

426   35.   Brophy JAN, Triassi AJ, Adams BL, Renberg RL, Stratis-Cullum DN, Grossman AD,
427   et al. Engineered integrative and conjugative elements for efficient and
428   inducible DNA transfer to undomesticated bacteria. *Nat Microbiol* 2018; **3**:
429   1043–1053.

430   36.   Kohler V, Keller W, Grohmann E. Regulation of Gram-Positive Conjugation. *Front*
431   *Microbiol* 2019; **10**: 1134.

432   37.   Inglis RF, Bayramoglu B, Gillor O, Ackermann M. The role of bacteriocins as
433   selfish genetic elements. *Biol Lett* 2013; **9**: 20121173.

434   38.   Hols P, Ledesma-García L, Gabant P, Mignolet J. Mobilization of Microbiota
435   Commensals and Their Bacteriocins for Therapeutics. *Trends Microbiol* 2019; **0**.

436   39.   Takahashi N, Naito Y, Handa N, Kobayashi I. A DNA methyltransferase can
437   protect the genome from postdisturbance attack by a restriction-modification
438   gene complex. *J Bacteriol* 2002; **184**: 6100–8.

439   40.   Naito T, Kusano K, Kobayashi I. Selfish behavior of restriction-modification
440   systems. *Science* 1995; **267**: 897–9.

441   41.   Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, et
442   al. In silico detection and typing of plasmids using PlasmidFinder and plasmid
443   multilocus sequence typing. *Antimicrob Agents Chemother* 2014; **58**: 3895–903.

445