Comprehensive genome data analysis establishes a triple whammy of carbapenemases, ICEs and multiple clinically-relevant bacteria

João Botelho[1#*], Joana Mourão[2,3,4], Adam P. Roberts[5,6], Luísa Peixe[1*]

[1]UCIBIO/REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal.

[2]University of Coimbra, Center for Innovative Biomedicine and Biotechnology, Coimbra, Portugal.

[3]University of Coimbra, Center for Neuroscience and Cell Biology, Coimbra, Portugal.

[4]University of Coimbra, Institute for Interdisciplinary Research, Coimbra, Portugal.

[5]Department of Tropical Disease Biology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom.

[6]Centre for Drugs and Diagnostics, Liverpool School of Tropical Medicine, Liverpool, United Kingdom.

[#]New affiliation: Antibiotic Resistance Evolution Group, Max-Planck-Institute for Evolutionary Biology, 24306 Plön, Germany; Department of Evolutionary Ecology and Genetics, Zoological Institute, Christian-Albrechts-Universität zu Kiel, 24118 Kiel, Germany.

*Addresses for correspondence: João Botelho, Antibiotic Resistance Evolution Group, Max-Planck-Institute for Evolutionary Biology, 24306 Plön, Germany; Department of Evolutionary Ecology and Genetics, Zoological Institute, Christian-Albrechts-Universität zu Kiel, 24118 Kiel, Germany.; e-mail: botelho@evolbio.mpg.de; Luísa Peixe, Laboratório de Microbiologia. Faculdade de Farmácia da Universidade do Porto, Rua Jorge Viterbo Ferreira nº 228, 4050-313 Porto, Portugal; phone: +351 220428500; e-mail: lpeixe@ff.up.pt.

Abstract

Carbapenemases inactivate most β-lactam antibiotics, including carbapenems and have been frequently reported among *Enterobacteriaceae*, *Acinetobacter* spp. and *Pseudomonas* spp. Traditionally, the horizontal gene transfer of carbapenemase-encoding genes (CEGs) has been linked to plasmids. However, given that integrative and conjugative elements (ICEs) are possibly the most abundant conjugative elements among prokaryotes, we conducted an *in-silico* analysis to ascertain the likely role of ICEs in the spread of CEGs among all bacterial genomes (n=182,663). We detected 17,520 CEGs, of which 66 were located within putative ICEs among several bacterial species (including clinically-relevant bacteria as *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* and *Escherichia coli*). Most CEGs detected within ICEs belong to the metallo-beta-lactamases IMP, NDM and SPM families and different mechanisms were likely responsible for acquisition of these genes. The majority of CEG-bearing ICEs belong to the MPF_G, MPF_T and MPF_F classes and often encode resistance to other antibiotics (e.g., aminoglycosides and fluoroquinolones). This study provides a snapshot of the different CEGs associated with ICEs among available bacterial genomes and sheds light on the underappreciated contribution of ICEs to the spread of carbapenem resistance globally.

Keywords

Integrative and conjugative elements, carbapenemases, antibiotic resistance.

Author Notes

53 All supporting data has been provided within the article or through supplementary data

54 files. Supplementary material is available with the online version of this article.

Abbreviations

57 AR, antibiotic resistance; CEG, carbapenemase-encoding gene; HGT, horizontal gene

58 transfer; ICE, integrative and conjugative element; MGE, mobile genetic element; MPF,

59 mating pair formation; NCBI, National Center for Biotechnology Information; T4SS, Type-

60 IV Secretion System; WHO, World Health Organization.

Impact Statement

Carbapenems are commonly used to treat severe infections in humans. Resistance is

often mediated by carbapenemases. These enzymes degrade carbapenems and are

frequently present in plasmids. Here, we demonstrate that carbapenemase-encoding

genes (CEGs) can also be located within integrative and conjugative elements (ICEs).

CEG-bearing ICEs belong to three mating-pair formation families. These mobile elements

may be particularly important in bacteria where plasmids do not seem to play a

significant role in the spread of antibiotic resistance genes, as *Pseudomonas* spp. This

study considerably expands the knowledge of the repertoire of CEGs-bearing ICEs

among clinically-relevant bacterial pathogens, such as *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* and *Escherichia coli*.

Data Summary

All the bacterial genomes scanned in this study have been deposited previously in the National Center for Biotechnology Information genome database and are listed on the supplementary tables. The extracted 66 ICEs in fasta format and the outputs for the profile HMMs scanned on the 386 putative MGEs identified in this study are deposited on figshare at https://figshare.com/projects/_Comprehensive_genome_data_analysis_establishes_a _triple_whammy_of_carbapenemases_ICEs_and_multiple_clinically-relevant_bacteria/78369.

Introduction

Due to the importance of carbapenems for the treatment of severe infections in humans, the WHO stated that these antibiotics should be reserved for infections caused by multidrug resistant Gram-negative bacteria in humans [1]. Recently, the same agency presented a list of bacterial pathogens for which research and development of new antibiotics are urgently required, and the top priority pathogens were the carbapenem-resistant strains of *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacteriaceae* [2].

The evolution of carbapenem-resistance in bacteria is often driven by the horizontal

gene transfer (HGT) of carbapenemase-encoding genes (CEGs) [3, 4]. Carbapenemases

are beta-lactamases able to hydrolyse carbapenems as well as most of other beta-

lactam antibiotics. These enzymes are members of the serine beta-lactamases classes A

and D, and the class B metallo-beta-lactamases [5]. The CEGs are often located on

integrons or transposons that themselves target mobile genetic elements (MGEs) such

as plasmids [3, 4], which makes the dissemination of these genes unpredictably complex

within bacterial communities. Recently, we demonstrated that another type of MGE,

the integrative and conjugative elements (ICEs), are likely to play a significant role as

vehicles for the dissemination of CEGs among *P. aeruginosa* [6]. Besides genes

conferring antibiotic resistance, ICEs may harbour additional cargo genes that provide

an adaptive advantage over other elements. Some of these examples include the

presence of the siderophore yersiniabactin encoded within the ICEKp in hypervirulent

clonal-group CG23 *Klebsiella pneumoniae* [7]; the Tn*5252*-related ICEs carrying

bacteriocin clusters in *Streptococcus suis* [8]; the type I-C CRISPR-Cas systems identified

within pKLC102-like ICEs in *P. aeruginosa* [9] and the type III restriction-modification

systems from SXT/R391-related ICEs in *Shewanella* spp. [10].

ICEs are self-transmissible MGEs that can integrate into and excise from the genome (as

transposons and phages do) and can exist as circular, sometimes replicable,

extrachromosomal elements and be transferred by conjugation (as some plasmids do)

[11–14]. ICEs appear to have a bipartite lifestyle that shifts between vertical and

horizontal transmission [15, 16]. HGT by conjugation requires three main components:

a relaxase, a mating pair formation (MPF) system, and a type-IV coupling protein, with

the last two forming a spanning-membrane multi-protein complex named type-IV

secretion system (T4SS) [17]. Until now, eight MPF classes were proposed (B, C, F, FA,

FATA, G, I and T), based on the phylogeny of VirB4, the only ubiquitous protein among

the T4SS. The MPF$_T$ is widely distributed both in conjugative plasmids and ICEs, while

MPF$_F$ is more prevalent in plasmids and MPF$_G$ on ICEs [11].

Given that ICEs were identified in most bacterial clades and were proposed to be more

prevalent than conjugative plasmids [11], we conducted an *in silico* analysis to explore

the distribution of CEG-bearing ICEs among all sequenced bacterial genomes. Our results

demonstrate that CEG-bearing ICEs belong to three MPF families and are primarily

located in several clinically-relevant bacterial pathogens. Our analysis highlights the

importance of thoroughly investigating these elements as important vehicles for the

spread of antibiotic resistance (AR), particularly to carbapenems.

Material and methods

Bacterial genomes and carbapenemases search

In **Figure 1,** we present the workflow used in this study, from the acquisition of bacterial

genomes to the identification and characterization of putative ICEs. We retrieved all

bacterial genomes available in NCBI Reference Sequence Database (RefSeq, accessed on

21/03/2020), including complete and draft genome sequences, using ncbi-genome-

download v0.2.12 (https://github.com/kblin/ncbi-genome-download). We downloaded

over 6,000 curated AR protein sequences from AMRfinder database

(https://ftp.ncbi.nlm.nih.gov/pathogen/Antimicrobial_resistance/AMRFinderPlus/data

base/3.6/2020-01-22.1/) [18] and we built an in-house database only including the

proteins that code for a carbapenemase (n=1014, **Table S1**). We then blasted the

genomes against the extracted carbapenemases using diamond v0.9.29.130

(http://www.diamondsearch.org/index.php) [19], using minimum 100% identity and

subject cover and with the sensitive mode enabled.

Bacterial genomes from RefSeq NCBI

Step 1
Diamond blastx against database of carbapenemases

Hits with CEGs

Step 2
Extract protein annotations from CEG–bearing hits

Proteomes

Step 3
Identification of profile HMMs for relaxases/integrases and delimitation of CEGs–bearing ICEs

Putative ICEs

ICE characterization:
– CONJscan (MPF system)
– AMRFinder (AR genes)
– REBASE (RM system)
– CRISPRCasFinder
– antismash (metabolites)
– eggNOG–mapper

Abbreviations: AR, Antibiotic Resistance; DB, Database; CEGs, carbapenemase–encoding genes; HMMs, Hidden Markov Models; ICEs, Integrative and Conjugative Elements; MLST, Multilocus Sequence Typing; MPF, Mating–Pair Formation; RM, Restriction–Modification

Figure 1 – Overview of the workflow followed in this study. All assemblies available in

NCBI RefSeq were downloaded and blasted against an in-house database of

carbapenemases using diamond blastx (Step 1). NCBI annotated proteins from CGE-

bearing genomes were then extracted (Step 2) and used for the identification of relaxase

and serine or tyrosine recombinase (Step 3). Search of directed repeats and delimitation

of putative ICEs was also performed. CONJscan was used to identify the MPF family of

each element. We then looked for AR genes, restriction-modification systems, CRISPR

arrays and their associated (Cas) proteins as well as secondary metabolites within

extracted ICEs. We also characterized the functional annotations of their proteomes and

the MLST of the genomes carrying a CEG-bearing ICE.

Tracing ICEs among the bacterial genomes

The RefSeq protein files from the CEG-bearing genomes identified by diamond were

extracted. We used the hmmsearch function of the HMMER3 software package v3.3

(http://hmmer.org/) [20] to search the proteomes against the standalone version of

MOBfamDB, a curated hidden Markov models (HMM) relaxase database

(https://castillo.dicom.unican.es/mobscan_about/) [21]. We also used this function to

search the pfam v33.0 database for serine or tyrosine recombinase accessions numbers

(Pfam IDs PF00589 and PF07508). The hmmsearch command was used with default

parameters and an E-value threshold of 0.01. The CEG-bearing genomes with relaxase

and integrase hits were further analysed. We manually inspected the hits to look for

direct repeats and to delimit CEG-harbouring ICEs. To predict the MPF families, the

translated coding sequences of delimited ICEs were analysed on the standalone

CONJscan module of MacSyFinder v1.0.5 (https://github.com/gem-

pasteur/macsyfinder) [22, 23]. To identify the multi-loci sequence type of the genomes

containing CEG-bearing ICEs, we used mlst v2.16.1

(https://github.com/tseemann/mlst), which scans the genomes against PubMLST typing

schemes (https://pubmlst.org/) [24].

Characterization of the CEG-bearing ICEs

Screening of AR genes among ICEs was performed using amrfinder v3.6.10

(https://github.com/ncbi/amr) [18]. The genetic platforms involved in the acquisition of

CEGs by ICEs were annotated using Galileo^{TM} AMR

(https://galileoamr.arcbio.com/mara/) (Arc Bio, Cambridge, MA) [25]. We ran our

extracted ICEs against REBASE (http://rebase.neb.com/rebase/rebase.html) to look for

restriction-modification systems [26]. We also used CRISPRCasFinder

(https://crisprcas.i2bc.paris-saclay.fr/CrisprCasFinder/Index) to look for CRISPR

(clustered regularly interspaced short palindromic repeats) arrays and their associated

(Cas) proteins within ICE sequences [27]. Secondary metabolite biosynthetic gene

clusters were traced using antismash v5.1.2

(https://antismash.secondarymetabolites.org/#!/start) [28]. We used eggNOG-mapper

v2 (http://eggnog-mapper.embl.de/) for functional annotation based on orthology

assignments of the ICE proteomes [29].

Results

Carbapenemase-encoding genes are mainly found in Proteobacteria

We retrieved a total of 182,663 bacterial genomes from NCBI (16,798 complete

genomes and 165,865 genomes assembled at the chromosome, scaffold or contig level).

We identified a total of 17,520 CEGs, with 1,422 CEGs on 1,236 complete genomes

(including 512 chromosomes and 724 plasmids) and 16,098 CEGs on 16,038 draft

genomes (**Table S2**). We identified a total of 377 carbapenemase variants among the

17,520 hits. Our results show that CEGs are mostly located on Proteobacteria and

dominated by clinically-relevant pathogens as *A. baumannii*, *K. pneumoniae*, *Escherichia*

coli and *P. aeruginosa* (**Table S2**). These genomes encode a wide diversity of

carbapenemases, including OXA-23 (15.6%, n=2,739/17,520), KPC-2 (13.2%), KPC-3

(8.9%) and NDM-1 (6.7%). As we are tracing MGEs integrated in the chromosome, the

724 plasmid hits in complete genomes were excluded from the analysis. Among the hits

on the 16,038 draft genomes, we filtered out sequences with the word 'plasmid' present

on the fasta-header (n=131, **Table S3**). To maximise the chances of detecting entire ICEs,

we filtered out sequences shorter than 40kb (n=10,050). All the excluded ones are

available for analysis in **Table S4**. The remaining sequences from draft genomes

(n=5,857) were inspected for the presence of CEG-harbouring ICEs.

A large proportion of CEG-bearing ICEs belong to three families and target

clinically-relevant gram-negative bacteria

We identified a total of 66 ICEs, including 42 newly characterized elements associated

with 17 different CEGs (**Table 1**). We could predict the boundaries from 55 of these

elements. The terminal region of the remaining 11 putative ICEs could not be

determined due to a fragmented contig or assembly gaps within the sequence. Nearly

half of the putative ICEs (48.5%, n=32/66) were integrated at the 3' end of a tRNA^{Gly}

gene. The bacterial hosts housing these elements belong to 23 sequence types (STs)

(**Table S5**).

Using the CONJscan module of MacSyFinder we identified the MPF family for 62 hits (no

family could be determined for the remaining 4 hits) and we noted that the majority of

the putative ICEs belong to three families: MPF$_G$ (69%, n=43/62), MPF$_T$ (26%) and MPF$_F$

(5%) (**Figure 2**, **Table 1** and **Table S6**). ICEs of the MPF$_G$ and MPF$_T$ classes were

particularly promiscuous, being responsible for the spread of several CEGs of the

metallo-beta-lactamase family such as *bla*$_{NDM-1}$, *bla*$_{SPM-1}$ and *bla*$_{IMP}$ variants among

clinically-relevant pathogens as *P. aeruginosa*, *K. pneumoniae* and *E. coli*. ICEs of the

MPF$_F$ class carrying blaIMP$_{-6}$ or *bla*$_{KPC-2}$ were restricted to *E. coli* and *K. pneumoniae*. The

bla$_{SPM-1}$ gene was exclusively identified in *P. aeruginosa* and in ICEs of the MPF$_T$ class

(**Table 1**).

Additionally, we also identified 386 hits encoding an integrase and a relaxase in the

vicinity of CEGs (**Table S7**). For these hits, however, we could not predict if the CEG is

located on a plasmid or an ICE, since the contig is fragmented and tracing the boundaries

of the element is not possible, or the sequences have assembly gaps that make this

prediction challenging. Also, some plasmids may encode a tyrosine or serine

recombinase, and some ICEs may encode replicases and partition systems that are

typical of plasmids [30], which can hinder the accurate prediction of genetic platforms

when the sequence has poor quality or is highly fragmented due to short-read

sequencing approaches.

Table 1 – Diversity and characterization of carbapenemases-encoding genes in integrative and conjugative elements-associated genomes.

MPF family	CEG	Bacterial Species	ICE genetic platforms	GC content (%)	Sequence length (kb)[1]	References
MPF$_G$ (n=43)	AFM-1 (n=1)	*Bordetella trematum*	Flanked by IS*91* family ISs	51.9	130	This study
	DIM-1 (n=1)	*Pseudomonas aeruginosa*	Class I In flanked by IS*6100*	62.8	89	This study, [6]
	GES-5 (n=5)	*Pseudomonas aeruginosa*	Class I In within Tn*3*-like Tn	60.5-62.1	93-116	This study, [6]
	GES-24 (n=1)	*Pseudomonas aeruginosa*	Class I In within Tn*3*-like Tn	63.0	63	This study
	IMP-1 (n=4)	*Pseudomonas aeruginosa*	Class I In within Tn*3*-like Tn	61.3-62.3	76-109	This study, [6]
	IMP-13 (n=9)	*Pseudomonas aeruginosa*	Class I In within Tn*3*-like Tn	61.5-62.0	65-89	This study, [6]

IMP-14 (n=2)	*Achromobacter xylosoxidans*	Class I In within Tn*3*-like Tn	61.7-62.1	106-122	This study
IMP-16 (n=1)	*Pseudomonas monteilii*	Class I In within Tn*3*-like Tn	62.6	86	This study
IMP-54 (n=1)	*Pseudomonas aeruginosa*	Class I In within Tn*3*-like Tn	61.7	91	This study
KPC-2 (n=1)	*Pseudomonas aeruginosa*	Tn*4401*	64.2	115	This study, [31]
NDM-1 (n=12)	*Pseudomonas aeruginosa, Pseudomonas asiatica, Morganella morganii*	Next to (or flanked by) IS*91* family ISs	58.7-64.5	97-167	This study, [32, 33]
VIM-2 (n=1)	*Pseudomonas aeruginosa*	Class I In within Tn*3*-like Tn	63.4	65	This study
VIM-4 (n=4)	*Pseudomonas aeruginosa,*	Class I In within Tn*3*-like Tn	61.0-62.3	88-102	This study

		Klebsiella pneumoniae, Alcaligenes faecalis				
MPF$_T$ (n=16)	AFM-1 (n=1)	*Stenotrophomonas maltophilia*	Flanked by IS*91* family ISs	63.7	77	This study
	KPC-2 (n=1)	*Pseudomonas* sp.	Next to IS*Kpn6* and IS*26*	59.2	62	This study, [6]
	NDM-1 (n=3)	*Pseudomonas aeruginosa, Pseudomonas asiatica*	Flanked by IS*91* family ISs	63.6-64.6	73-74	This study, [34]
	SPM-1 (n=11)	*Pseudomonas aeruginosa*	Flanked by IS*CR3*-like elements	64.9-65.5	44-58	This study, [35]
MPF$_F$ (n=3)	IMP-6 (n=1)	*Escherichia coli*	Class I In within Tn*3*-like Tn	51.9	118	This study
	KPC-2 (n=2)	*Klebsiella pneumoniae*	Tn*4401*	54.4	75	This study

	IMP-8 (n=1)	*Enterobacter cloacae*	Class I In within Tn*3*-like Tn	53.2	124	This study, [36]
Not assigned (n=4)	KPC-2 (n=1)	*Achromobacter* sp.	Next to IS*Kpn6* and IS*26*	60.4	43	This study
	NDM-1 (n=1)	*Pseudomonas aeruginosa*	Flanked by IS*CR3*-like elements	63.6	98	This study
	NDM-5 (n=1)	*Escherichia coli*	Flanked by IS*CR1* and IS*26*	52.4	54	This study, [37]

Abbreviations: CEG, carbapenemase-encoding gene; ICE, Integrative and Conjugative Elements; In, Integron; MPF, Mating Pair Formation; ST, Sequences Type.

[1]Some sequences are not complete. For more details, please refer to **Table S5**.

Abbreviations: Ax, Achromobacter xylosoxidans; Af, Alcaligenes faecalis; Bt, Bordetella trematum; Ec, Escherichia coli, Kp, Klebsiella pneumoniae; Mm, Morganella morganii; Pas, Pseudomonas asiatica; Pm, Pseudomonas monteilii; Psp, Pseudomonas sp.; Sm, Stenotrophomonas maltophilia.

241   **Figure 2 –** Sankey diagram showing the contribution of different MPF families for the spread of CEGs among several bacterial genomes. The left,

242   centre and right axis represents the association between the identified carbapenemases, MPF type and the bacterial genus, respectively, while

243   the width of each connection is proportional to the number of positive hits.

A variable repertoire of CEG-bearing integrons and transposons target ICEs

We identified 17 CEG variants among the 66 putative ICEs, dominated by bla_{NDM-1} and

bla_{SPM-1} (**Table 1**). Insertion sequences (ISs, e.g. ISCR3-like elements) were frequently

linked to the acquisition of bla_{SPM-1} and bla_{NDM-1} [34, 35], while bla_{IMP}, bla_{VIM}, and bla_{GES}

were found on class I integrons frequently integrated into Tn3-family transposons [6].

The bla_{KPC-2} gene was typically found within Tn4401-like transposons, which are capable

of conferring a high frequency of transposition [38]. Curiously, the recently identified

AFM-1 metallo-beta-lactamase (Genbank accession number MK143105.1) was here

identified in two ICEs inserted in *Bordetella trematum* and *Stenotrophomonas*

maltophilia genomes (**Table 1**). We also found bla_{NDM-1} genes in ICEs integrated into the

genomes of a recently proposed *Pseudomonas asiatica* species, which is spreading in

hospital settings in Myanmar [39]. Besides CEGs, the ICEs identified in this study also

harbour genes conferring resistance to other antibiotics, such as aminoglycosides,

fluoroquinolones, macrolides, and tetracyclines (**Table S8**), widening the spectrum of

transmissible AR genes selectable by carbapenems due to linkage.

Acquisition of additional traits by ICEs include competitive weapons as bacteriocins and siderophores

Besides genes conferring antibiotic resistance, the CEG-bearing ICEs here identified

harbour other cargo genes that may confer a selective advantage to the ICE host. We

found DUF692 domains typical of bacteriocin production genes among the 6 MPF$_G$ ICEs

from *P. asiatica* strains and the *P. aeruginosa* N15-01092, 1334/14 and ST773 strains

(**Table S5**). All these ICEs carry a *bla*$_{NDM-1}$ gene and a similar copy of the bacteriocin-

encoding gene. Curiously, the bacteriocin production gene was not identified in the

MPF$_T$ ICE from *P. asiatica* strain MY569. Additionally, we found the siderophore

aerobactin operon within an ICE in *E. coli* strain E302 (**Table S5**). This operon is usually

found in enterobacterial plasmids, and was also identified in a pathogenicity island in

uropathogenic *E. coli* strain CFT073 [40]. We identified no CRISPR-Cas systems nor

restriction-modification systems within the 66 putative ICEs.

The majority of the proteins encoded within the 66 ICEs refer to replication,

recombination, transcription and intracellular trafficking functions (**Figure 3**). Several

proteins, however, encoded for unknown functions (34.0%, n=1,615/4,754, **Table S9**),

highlighting the lack of knowledge on the ICE proteome.

Figure 3 – Grouped bar chart representing the incidence of each eggNOG function broken down by category (represented by different colours)

in the CEG-bearing ICE sequences.

Discussion

We have set out to comprehensively identify the CEGs among all bacterial genomes deposited in NCBI and the CEG-bearing ICE sequences. Our study considerably expands the knowledge of the repertoire of CEGs-bearing ICEs. We uncovered 66 putative ICEs that may be involved in HGT of CEGs amongst bacterial genomes. To expand our predictions, we also used the CONJscan module of Macsyfinder to trace the MPF families likely involved in HGT.

The scenario observed for the acquisition of the most important CEGs by ICEs (ISs for bla_{NDM-1} and bla_{SPM-1}, class I integrons for bla_{IMP} and bla_{VIM} and Tn4401 for bla_{KPC-2}) resembles that of plasmids [3] and provide additional support to the notion that the line separating these elements is blurred [13, 14]. We now show that besides plasmids, this promiscuous repertoire of integrons and transposons frequently targets ICEs of different MPF families. Even though CEGs might rapidly spread worldwide, local selection is likely required for them to reach fixation, as can be seen for the clonal expansion of *P. aeruginosa* ST277 harbouring bla_{SPM-1} [35]. Surprisingly, we noted that this gene was unable to be detected beyond the same clonal lineage. Indeed, all hits were identified in *P. aeruginosa* ST277 strains from Brazil and within a MPF_T ICE family, indicating that these ICEs are only transferring vertically and/or horizontally within this ST. Understanding the limitations on HGT of this family of ICE may be translatable to other, more transferable, ICEs and could underpin a control strategy to prevent the spread of these elements in the future.

Although bla_{KPC} and bla_{OXA} were the most frequently identified CEGs within the analysed genomes (**Table S2**), we only found 5 ICEs carrying bla_{KPC-2} and 2 ICEs with bla_{OXA} genes

(**Table S8**). The bla_{KPC} genes are mostly located in Tn*4401* transposons that target *K.*

pneumoniae plasmids, while the bla_{OXA} genes are frequently associated with ISs that

tend to target the *Acinetobacter* spp. chromosomes and plasmids [3]. These genes may

take advantage of the copy number and the higher genetic plasticity of plasmids [30].

This plasticity may increase the rate of appearance of novel mutations and the high copy

number may amplify the effect due to the increased gene dosage [41].

In addition to AR, ICEs can be involved in other adaptive traits such as carbon-source

utilization, symbiosis, restriction-modification, siderophore and bacteriocin synthesis

[14]. Since bacteria commonly inhabit highly competitive environments, the production

of specific secondary metabolites (such as bacteriocins and siderophores) may confer a

selective advantage to the host [42]. We speculate that the presence of these

metabolites within the ICEs here characterized may promote their stability by

preferentially selecting for cells harbouring the ICE.

One caveat of our studies is that these results do not yet expose the complete set of

CEG-bearing ICEs present in all bacteria. There is an inherent bias in the number of times

we detect a particular CEG in certain pathogen genomes as some are over represented

in the database compared to others. It is possible that certain observations will flatten

out as more genomes are analysed. Less than 10% of the bacterial genomes currently

present on NCBI are complete. This is a major drawback since putative ICEs present in

draft genomes tend to be fragmented due to the presence of repetitive regions that are

not resolved using short-read sequencing. Also, the relaxase and the T4SS encoded by

ICEs resemble those of plasmids [11, 13, 43]. Plus, it is possible that ICEs and plasmids

have swapped conjugation modules throughout their evolutionary history [11]. We

believe that a more thorough exploration of this issue, especially regarding the precise

delimitation of ICEs, will be an important further step toward an improved

understanding of the contribution of these elements to bacterial adaptation and

evolution of AR.

While we have chosen to focus on CEG-bearing genomes, our computational approach

can be applied to trace other relevant AR genes and other cargo genes that may confer

a selective advantage to the ICE host. Leveraging knowledge linking the accurate

prediction of ICE sequences to the carriage of AR genes, will not only improve our

understanding of HGT, but may also uncover potential approaches to tackle the spread

of AR.

Data bibliography

The DNA sequences used in this study are available in NCBI, under the accession

numbers provided in supplementary tables S1-S9.

Funding information

This work was supported by the Applied Molecular Biosciences Unit - UCIBIO which is

financed by national funds from FCT (UIDB/04378/2020).

Author contributions

J. B. conceptualized the project. J. B. ran the analyses. J.B. and J. M. analysed the data.

J.B. wrote the manuscript. J. M., A. P. R. and L. P. edited and revised the manuscript. All

authors read, commented on and approved the final manuscript.

Conflicts of interest

353 The authors declare that there are no conflicts of interest.

References

1. EFSA Panel on Biological Hazards (BIOHAZ). Scientific Opinion on Carbapenem resistance

in food animal ecosystems. *EFSA J* 2013; **11**.

2. World Health Organization. GLOBAL PRIORITY LIST OF ANTIBIOTIC-RESISTANT BACTERIA

TO GUIDE RESEARCH, DISCOVERY, AND DEVELOPMENT OF NEW ANTIBIOTICS.

http://www.who.int/news-room/detail/27-02-2017-who-publishes-list-of-bacteria-for-

which-new-antibiotics-are-urgently-needed. Accessed 11 Aug 2018.

3. Partridge SR, Kwong SM, Firth N, Jensen SO. Mobile Genetic Elements Associated with

Antimicrobial Resistance. *Clin Microbiol Rev* 2018; **31**: e00088-17.

4. Botelho J, Grosso F, Peixe L. Antibiotic resistance in Pseudomonas aeruginosa –

Mechanisms, epidemiology and evolution. *Drug Resist Updat* 2019; **44**: 100640.

5. Queenan AM, Bush K. Carbapenemases: the versatile beta-lactamases. *Clin Microbiol Rev*

2007; **20**: 440–58, table of contents.

6. Botelho J, Roberts AP, León-Sampedro R, Grosso F, Peixe L. Carbapenemases on the

move: it's good to be on ICEs. *Mob DNA* 2018; **9**: 37.

7. Lam MMC, Wyres KL, Duchêne S, Wick RR, Judd LM, Gan YH, et al. Population genomics

of hypervirulent Klebsiella pneumoniae clonal-group 23 reveals early emergence and

rapid global dissemination. *Nat Commun* 2018; **9**.

8. Libante V, Nombre Y, Coluzzi C, Staub J, Guédon G, Gottschalk M, et al. Chromosomal

Conjugative and Mobilizable Elements in Streptococcus suis: Major Actors in the

Spreading of Antimicrobial Resistance and Bacteriocin Synthesis Genes. *Pathog (Basel,*

Switzerland) 2019; **9**.

9. van Belkum A, Soriaga LB, LaFave MC, Akella S, Veyrieras J-B, Barbu EM, et al.

Phylogenetic   Distribution   of   CRISPR-Cas   Systems   in   Antibiotic-Resistant

Pseudomonas aeruginosa. *MBio* 2015; **6**: e01796-15.

10. Fang Y, Wang Y, Li Z, Liu Z, Li X, Diao B, et al. Distribution and Genetic Characteristics of

SXT/R391 Integrative Conjugative Elements in Shewanella spp. From China. *Front*

Microbiol 2018; **9**: 920.

11. Guglielmini J, Quintais L, Garcillán-Barcia MP, de la Cruz F, Rocha EPC. The Repertoire of

ICE in Prokaryotes Underscores the Unity, Diversity, and Ubiquity of Conjugation. *PLoS*

Genet 2011; **7**: e1002222.

12. Carraro N, Burrus V. The dualistic nature of integrative and conjugative elements. *Mob*

Genet Elements 2015; **5**: 98–102.

13. Cury J, Touchon M, Rocha EPC. Integrative and conjugative elements and their hosts:

composition, distribution and organization. *Nucleic Acids Res* 2017; **45**: 8943–8956.

14. Johnson CM, Grossman AD. Integrative and Conjugative Elements (ICEs): What They Do

and How They Work. *Annu Rev Genet* 2015; **49**: 577–601.

15. Carraro N, Poulin D, Burrus V. Replication and Active Partition of Integrative and

Conjugative Elements (ICEs) of the SXT/R391 Family: The Line between ICEs and

Conjugative Plasmids Is Getting Thinner. *PLOS Genet* 2015; **11**: e1005298.

16.   Delavat F, Miyazaki R, Carraro N, Pradervand N, van der Meer JR. The hidden life of
integrative and conjugative elements. *FEMS Microbiol Rev* 2017; **41**: 512–537.

17.   Smillie C, Garcillán-Barcia MP, Francia MV, Rocha EPC, de la Cruz F. Mobility of plasmids.
Microbiol Mol Biol Rev 2010; **74**: 434–52.

18.   Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I, et al. Validating the
AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance
Genotype-Phenotype Correlations in a Collection of Isolates. *Antimicrob Agents*
Chemother 2019; **63**.

19.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. *Nat*
Methods 2015; **12**: 59–60.

20.   Eddy SR. Accelerated profile HMM searches. *PLoS Comput Biol* 2011; **7**: e1002195.

21.   Garcillán-Barcia MP, Redondo-Salvo S, Vielva L, de la Cruz F. MOBscan: Automated
Annotation of MOB Relaxases. *Methods in Molecular Biology*. 2020. Humana Press Inc.,
pp 295–308.

22.   Abby SS, Néron B, Ménager H, Touchon M, Rocha EPC. MacSyFinder: A Program to Mine
Genomes for Molecular Systems with an Application to CRISPR-Cas Systems. *PLoS One*
2014; **9**: e110726.

23.   Abby SS, Cury J, Guglielmini J, Néron B, Touchon M, Rocha EPC. Identification of protein
secretion systems in bacterial genomes. *Sci Rep* 2016; **6**: 23080.

24.   Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb
software, the PubMLST.org website and their applications. *Wellcome Open Res* 2018; **3**:
124.

25. Partridge SR, Tsafnat G. Automated annotation of mobile antibiotic resistance in Gram-

negative bacteria: the Multiple Antibiotic Resistance Annotator (MARA) and database. *J*

Antimicrob Chemother 2018; **73**: 883–890.

26. Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE—a database for DNA restriction and

modification: enzymes, genes and genomes. *Nucleic Acids Res* 2015; **43**: D298–D299.

27. Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, et al.

CRISPRCasFinder, an update of CRISRFinder, includes a portable version, enhanced

performance and integrates search for Cas proteins. *Nucleic Acids Res* 2018; **46**: W246–

W251.

28. antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline | Nucleic

Acids   Research | Oxford   Academic.

https://academic.oup.com/nar/article/47/W1/W81/5481154. Accessed 21 Feb 2020.

29. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology

resource based on 5090 organisms and 2502 viruses | Nucleic Acids Research | Oxford

Academic. https://academic.oup.com/nar/article/47/D1/D309/5173662. Accessed 24

Feb 2020.

30. Cury J, Oliveira PH, de la Cruz F, Rocha EPC. Host Range and Genetic Plasticity Explain the

Coexistence of Integrative and Extrachromosomal Mobile Genetic Elements. *Mol Biol*

Evol 2018; **35**: 2230–2239.

31. Abril D, Marquez-Ortiz RA, Castro-Cardozo B, Moncayo-Ortiz JI, Olarte Escobar NM,

Corredor Rozo ZL, et al. Genome plasticity favours double chromosomal Tn4401b-bla

KPC-2 transposon insertion in the Pseudomonas aeruginosa ST235 clone. *BMC Microbiol*

2019; **19**.

32. Urbanowicz P, Izdebski R, Baraniak A, Żabicka D, Ziółkowski G, Hryniewicz W, et al.

Pseudomonas aeruginosa with NDM-1, DIM-1 and PME-1 β-lactamases, and RmtD3 16S

rRNA methylase, encoded by new genomic islands. *J Antimicrob Chemother* 2019.

33.   Khan A, Shropshire WC, Hanson B, Dinh AQ, Wanger A, Ostrosky-Zeichner L, et al.

Simultaneous infection with enterobacteriaceae and pseudomonas aeruginosa

harboring multiple carbapenemases in a returning traveler colonized with candida auris.

Antimicrob Agents Chemother 2020; **64**.

34.   Ding Y, Teo JWP, Drautz-Moses DI, Schuster SC, Givskov M, Yang L. Acquisition of

resistance to carbapenem and macrolide-mediated quorum sensing inhibition by

Pseudomonas aeruginosa via ICETn43716385. *Commun Biol* 2018; **1**: 57.

35.   Nascimento APB, Ortiz MF, Martins WMBS, Morais GL, Fehlberg LCC, Almeida LGP, et al.

Intraclonal Genome Stability of the Metallo-β-lactamase SPM-1-producing Pseudomonas

aeruginosa ST277, an Endemic Clone Disseminated in Brazilian Hospitals. *Front Microbiol*

2016; **7**: 1946.

36.   Zhan Z, Hu L, Jiang X, Zeng L, Feng J, Wu W, et al. Plasmid and chromosomal integration

of four novel blaIMP-carrying transposons from Pseudomonas aeruginosa, Klebsiella

pneumoniae and an Enterobacter sp. *J Antimicrob Chemother* 2018; **73**: 3005–3015.

37.   McCracken MG, Adam HJ, Blondeau JM, Walkty AJ, Karlowsky JA, Hoban DJ, et al.

Characterization of carbapenem-resistant and XDR Pseudomonas aeruginosa in Canada:

results of the CANWARD 2007–16 study. *J Antimicrob Chemother* 2019; **74**: iv32–iv38.

38.   Cuzon G, Naas T, Nordmann P. Functional characterization of Tn4401, a Tn3-based

transposon involved in blaKPC gene mobilization. *Antimicrob Agents Chemother* 2011;

55: 5370–3.

39.   Tohya M, Tada T, Watanabe S, Kuwahara-Arai K, Zin KN, Zaw NN, et al. Emergence of

carbapenem-resistant pseudomonas asiatica producing NDM-1 and VIM-2 Metallo-

Lactamases in Myanmar. *Antimicrob Agents Chemother* 2019; **63**.

40.   Banerjee R, Weisenhorn E, Schwartz KJ, Myers KS, Glasner JD, Perna NT, et al. Tailoring a

Global Iron Regulon to a Uropathogen. *MBio* 2020; **11**.

41.   San Millan A, Escudero JA, Gifford DR, Mazel D, MacLean RC. Multicopy plasmids

potentiate the evolution of antibiotic resistance in bacteria. *Nat Ecol Evol* 2016; **1**: 0010.

42.   Inglis RF, Bayramoglu B, Gillor O, Ackermann M. The role of bacteriocins as selfish genetic

elements. *Biol Lett* 2013; **9**: 20121173.

43.   Guglielmini J, Néron B, Abby SS, Garcillán-Barcia MP, de la Cruz F, Rocha EPC. Key

components of the eight classes of type IV secretion systems involved in bacterial

conjugation or protein secretion. *Nucleic Acids Res* 2014; **42**: 5715–27.

475