Structure-based drug repositioning explains ibrutinib as VEGFR2 inhibitor

Melissa F. Adasme[1,◑], Daniele Parisi[2,◑], Kristien Van Belle[3], Sebastian Salentin[1], V. Joachim Haupt[1,7], Gary S. Jennings[1], Jörg-Christian Heinrich[1], Jean Herman[3,4,5], Ben Sprangers[3,4,6], Thierry Louat[3], Yves Moreau[2], Michael Schroeder[1,*]

[1] Biotechnology Center (BIOTEC), Technische Universität Dresden, 01307 Dresden, Germany
[2] ESAT-STADIUS, KU Leuven, B-3001 Heverlee, Belgium
[3] Interface Valorisation Platform (IVAP), KU Leuven, 3000 Leuven, Belgium
[4] Laboratory of Molecular Immunology (Rega institute), KU Leuven, Belgium
[5] Department of Pediatric Nephrology and Solid Organ Transplantation, University Hospitals Leuven, 3000 Leuven, Belgium
[6] Department of Nephrology, University Hospitals Leuven, 3000 Leuven, Belgium
[7] PharmAI GmbH, 01307 Dresden, Germany

◑These authors contributed equally to this work.
* correspondingauthor@institute.edu

Abstract

Many drugs are promiscuous and bind to multiple targets. On the one hand, these targets may be linked to unwanted side effects, but on the other, they may achieve a combined desired effect (polypharmacology) or represent multiple diseases (drug repositioning). With the growth of 3D structures of drug-target complexes, it is today possible to study drug promiscuity at the structural level and to screen vast amounts of drug-target interactions to predict side effects, polypharmacological potential, and repositioning opportunities. Here, we pursue such an approach to identify drugs inactivating B-cells, whose dysregulation can function as a driver of autoimmune diseases. Screening over 500 kinases, we identified 22, whose knock out impeded the activation of B-cells. Among these 22 is the gene KDR, whose gene product VEGFR2 is a prominent cancer target with anti-VEGFR2 drugs on the market for over a decade. The main result of this paper is that structure-based drug repositioning for the identified kinase targets identified the cancer drug ibrutinib as micromolar VEGFR2 inhibitor with a very high therapeutic index in B-cell inactivation. These findings prove that ibrutinib is not only acting on the Bruton's tyrosine kinase BTK, against which it was designed. Instead, it may be a polypharmacological drug, which additionally targets angiogenesis via inhibition of VEGFR2. Therefore ibrutinib carries potential to treat other VEGFR2 associated disease. Structure-based drug repositioning explains ibrutinib's anti VEGFR2 action through a specific pattern of interactions shared between BTK and VEGFR2. Overall, structure-based drug repositioning was able to predict these findings at a fraction of the time and cost of a conventional screen.

Introduction

The number of ways a drug can interact with its target is limited. Alex et al. estimate that there are 10000 interfaces [2] and Gao et al. reduce this number to 1000 distinct

interface types [10]. While the exact number may be debatable, a limit itself has important consequences: there is redundancy and re-use in drug-target interaction, which makes it difficult to design a target-specific or one target-only drug. Hence, drug promiscuity, a drug hitting multiple targets, is rather the normality than the exception [16, 19, 27].

While drug promiscuity can be a source of a drug's side effects, it can also be advantageous: In polypharmacology, drugs are designed to hit multiple targets associated to multiple desired effects in a disease. This ensures that the drug is more robust [3] to changes in the targets such as expression levels or mutation. Polypharmacology is successfully applied in practice. As an example, consider the kidney cancer drugs sunitinib, sorafenib, and pazopanib [23, 28], which are approved since 2006 and 2010, respectively. These drugs simultaneously act on proliferation and angiogenesis by binding (among others) both the platelet-derived growth factor receptors PDGFR and the vascular endothelial growth factor receptor VEGFR2. Targets such as PDGFR and VEGFR2 are particularly suitable to polypharmacology as they are kinases, the largest family of targets that bind to a common substrate adenosine triphosphate (ATP) [23].

Overington et al. argue that by nature of their close evolutionary relation and similar function, several members of the same family may be affected by a drug, which is thus promiscuous. In [19, 29], the authors quantify such polypharmacology by constructing a network of targets, which are related by shared drugs. The network is particularly dense for kinases, underlying the potential for side effects and polypharmacology. It also covers other classes of drug targets, such as e.g. G protein-coupled receptors (GPCRs), for which structural studies of selectivity and promiscuity [30, 34] pave the way for novel avenues in drug design through polypharmacology.

Besides polypharmacology, drug promiscuity can also support drug repositioning. Drug repositioning aims to identify novel indication for an existing drug. Often this is achieved by exploring diseases closely related to the original indication or by linking the known main drug target to a new disease. That latter was the case for Viagra with its repositioning from heart disease to erectyle disfunction. For both, old and new indication, the main target is the phosphodiesterase PDE5. However, repositioning is also possible if a drug hits two targets, one linked to the existing indication and the other to a new one. For example, the anti-herpes drug BVDU binds a viral thymidine kinase, but also the heat shock protein Hsp27, an anti-cancer target. Thus, BVDU's promiscuity achieves two very different effects supporting its repositioning efforts [18] from herpes to cancer.

If drug promiscuity can be beneficial in polypharmacology and in drug repositioning, it becomes important to understand its source. As argued above, one reason for a drug to bind multiple targets is the limited space of binding interfaces [2, 16]. As a consequence, there are binding sites and interfaces which are very similar across unrelated targets and drugs. These similarities can be studied and uncovered algorithmically. In [24] the authors review target- and drug-centric approaches to polypharmacology. The former comprise alignments of binding sites and the latter fingerprints [7, 8], which characterize a drug and its interaction to a target. They can comprise chemical interactions such as hydrogen bonds, pi-stacking, or hydrophobic interactions together with geometric information on distances and angles. These interactions are encoded as standardized binary vectors, which become comparable. As a consequence, it is possible to search large repositories of drug-target complexes. Here, we will present and pursue such a structural approach to polypharmacology and drug repositioning with the goal of identifying candidate drugs for B-cell inactivation (see Fig 1).

B-cells produce antibodies and establish effective inter cellular communication for a

Fig 1. Overview of the study. Background. B- and T-cells play an important role in autoimmune diseases. Current treatments focus on inhibition of T-cell activation. Novel or improved treatments may emerge by inhibiting B-cells. **Application.** Through an RNAi screen, we identified kinases, that play a role in B-cell activation and are therefore suitable autoimmune targets. There are two routes to validate inhibitors against these kinases. First, known inhibitors may exist and second, structural drug repositioning may yield novel inhibitors after *in vitro* validation. Finally, all validated compounds are checked *in vitro* for selectivity and efficacy of B-cell inactivation.

controlled immune response through expression of cell surface markers and secretion of signaling molecules. B-cells play an important role in many blood cancers and in autoimmune diseases such as systemic lupus erythematosus, rheumatoid arthritis and psoriasis. Interestingly, there is a connection between cancer and psoriasis as angiogenesis, one of the cancer hall marks [14], and high expression of vascular endothelial growth factors (VEGFs) are present in psoriasis [13]. Subsequently, the polypharmacological cancer drugs pazopanib, sunitinib, and sorafenib lead to improvements in psoriasis [39]. This is particularly remarkable, as their initial indication is kidney cancer and not a blood cancer suggesting that blood cancer may be a

repositioning opportunity. In fact, there are links between blood cancer and autoimmunity: Ibrutinib is used to treat leukemias and lymphomas as well as chronic graft vs. host disease. It was designed against Bruton's tyrosine kinase (BTK), which are highly expressed in B-cells. Despite the designed specific inhibition of BTKs, there is recent evidence that ibrutinib is anti-angiogenic [31] suggesting that there are other targets.

Rituximab and bortezomib are two other cancer and autoimmune drugs, which act on B-cells. Rituximab is a monoclonal antibody, which binds to a surface marker of B-cells and then triggers cell-death [32]. Bortezomib is a cytotoxic small molecule widely used in cancer chemotherapy. Recently, it was tested for its autoimmune potential in lupus [1]. Both drugs kill B-cells rather than modulating their activity.

While there are links from cancer to auto-immune diseases for these drugs, they are nonetheless cytotoxic. Therefore, it is interesting to complement these drugs with novel small molecules, which inactivate B-cells and which have low toxicity. To this end, we will employ an assay to identify suitable targets driving B-cell inactivation and we will identify known and novel binders using a structural polypharmacology and drug repositioning approach. For target identification and lead validation, we will use a recently introduced assay [38], in which B-cell inactivation is read out via two cell surface markers. In [38], the authors compared six cell lines, thirteen stimuli, and eight read outs concluding that human Burkitt lymphoma cells (Namalwa) stimulated by the TLR9 ligand ODN2006 using expression-levels of the cell surface markers CD70 and CD80 is best suited to study B-cell inactivation. Cell surface markers such as CD70 and CD80 act as co-stimulatory molecules for the activation and differentiation of CD4+ T-cells. This is necessary for the development of an effective immune response. Furthermore, In vitro studies with murine B cells [37] and in vivo studies with murine disease models and transgenic mice [4] have indicated that markers CD70 and CD80 are also key molecules in the signalling for the regulation of B cell's effector functions such as Ig production. Naive B-cells from patients with common variable immunodeficiency are markedly impaired in upregulating the costimulatory molecules CD80 and CD70 upon BCR cross-linking and the expression remained reduced even in presence of autologous helper CD4+ T cells. The insufficient upregulation of these two crucial costimulatory molecules could explain the poor class switching and, hence, reduced Ig serum levels, except for IgM [36].

Materials and methods

The general methodology applied in this study is represented in Fig 1.Application and below there is a detailed description of all the step-by-step protocols performed in this work:

RNAi screening and target selection. The approach developed, uses human Burkitt lymphoma Namalwa cells [26] stimulated with ODN2006. For RNAi screening, the MISSION LentiExpress TM shRNA library (Sigma-Aldrich, Diegem, Belgium) was used, which has 3 - 10 shRNA clones per kinase target. After transduction, cells were stimulated by ODN2006 for 24 hours. Cells were incubated with fluorescent antibodies for CD70 and CD80. After FACS sorting, expression was measured by amount of fluorescence. Then, the y-value was calculated as follows:

$$y = \frac{SC - SS}{SC - NSC}$$

where SC is the expression of stimulated control, SS the stimulated sample, and NSC the non-stimulated control. Based on a close observation of the data, since generally, as

observed, silencing of "hit" kinases gave a more potent inhibition of marker CD70 compared to marker CD80, clones with $y > 0.8$ for CD70 and $y > 0.5$ for CD80 were selected and resulted in the 22 kinase targets.

Known inhbitors. Known inhibitors were retrieved from BindingDB in April 2018. A drug's target in BindingDB had 100% sequence similarity to one of the 22 kinase targets and the Ki, Kd, or IC50 had to be less than 50μM.

Structural drug-repositioning. All ligand-target complexes in Protein Data Bank (PDB) [15] were represented as interaction fingerprints. Interaction fingerprints are binary vectors, where each feature represents a combination of interactions within an angle and distance range (see Fig 2). Interactions are obtained from PLIP [33] and comprise e.g. hydrogen bonds, hydrophobic contacts, pi stacking. Two interaction fingerprints designed by PharmAI company were compared by computing their Tanimoto score, i.e. the number of shared features divided by the number of overall features present in either vector. The score ranges from 0 (no features in common) to 1 (all features in common). For a set of fingerprints, the significance of a score is judged by p-value. When screening a template ligand-target complex against the PDB, the interaction fingerprint scores are sorted by p-value and only those with p-value less than 10^{-3} are retained. To apply structural drug-repositioning, we identified suitable templates in PDB. We found structures for 14 of the 22 kinase targets. This reduced to 10 after removing structures without interacting compounds or binding ions or biologically non-relevant small molecules according to BioLiP (zhanglab.ccmb.med.umich.edu/BioLiP). For these 10 kinases, there were a total of 141 drug-target templates.

Kinase activity test. Compounds predicted from structural drug repositioning were further filtered for FDA approval using DrugBank. Nutraceuticals, already known binders, and 6 compounds not available from Sigma-Aldrich were discarded. Dasatanib, a strong binder to all targets, was added as control. The 111 compounds were bought from Sigma-Aldrich and Selleckchem. Kinase inhibition was tested for them in duplicate at 1μM and 10μM concentrations resulting in 888 assays carried out by ReactionBiology. Data were normalized to percentage Enzyme Activity relative to DMSO controls. Compounds inhibiting more than 50 percent of the enzyme activity have been defined as strong inhibitors and inhibition between 30 and 50 percent as weak inhibition.

Chemical structures and similarity. Chemical similarity was computed with the PubChem Score Matrix Service [22] with standard settings. Chemical structures were downloaded in SDF format. Heatmaps were plotted with the Heatplus package v2.16.0 in R 3.2.5 using hierarchical clustering on average distances.

Links between targets and disease. To establish links between the 22 targets and disease we manually search PubMed (April-July 2018) and Open Targets [5] (December 2018). The search was done using the target name together with disease keywords cancer, cancer of the immune system, and autoimmune disease. We refined the latter with specific autoimmune disease such as lupus, multiple sclerosis, psoriasis, type 1 diabetes, Grave's syndrome, inflammatory bowel disease, Hashimoto's thyroiditis, Sjögren syndrome, celiac disease, rheumatoid arthritis. Cancer of the immune system was expanded with keywords myeloma, leukemia, lymphoma and cancer with tumour, adenoma, sarcoma, carcinoma, blastoma.

A PLIP Interaction Data (1VSN:NFT:A:283)

B 3D Pattern Pairings

Angle and distance binning

$1.5\ \text{Å} < d_2 \leq 2.5\ \text{Å}$
$0 < a_1 \leq 30\ °$

HYB : HAL : d_2 : a_1

+1

0 0 0 0 **1** 0 0 0 0 0 0 — bin

C Structure-Invariant Fingerprint

Fig 2. Structural drug-repositioning with PLIP interaction fingerprints.
A. Schematic for the PLIP interaction fingerprints. Raw interaction data from PLIP are
calculated for a complex. **B.** For each pair of interaction vectors, their distances and
angles are calculated. **C.** Finally, the geometric measures are binned and combined with
the types of the pairing into a string representation. Each such combination corresponds
to a feature in the fingerprint, which is incremented for each observation of the encoded
pairing. The resulting fingerprint is structure-invariant since it only considers the
interaction vectors.

Links between drugs and disease. PubChem was used (May 2018) to link drugs   156
to diseases, which were grouped into the five high-level groups cancer, cancer of the   157
immune system, autoimmune diseases, infection, and others.   158

Drug-target-disease network. A drug-target-disease network was constructed by   159
adding links from known as well as predicted and validated binders to their respective   160
targets and adding the above target-disease and drug-disease links. The network was   161
visualized with Cytoscape v3.6.0.   162

Drugs' known targets. The number of targets in Fig 7 was retrieved from   163
BindingDB [12]. A target qualified if it is binding with a Ki, IC50 or Kd less than   164
50μM. Targets are grouped by unique Uniprot Id.   165

Kinase binding assay. Dose response curves for ibrutinib and suramin against   166
VEGFR2 were determined with the KINOMEscan TM by DisvcoverX, a competition   167
binding assay that quantitatively measures the ability of a compound to compete with   168
an immobilized, active-site directed ligand. It is measured via quantitative PCR of the   169
DNA tag linked to the kinase. An 11-point 3-fold serial dilution of each test compound   170
was prepared in 100% DMSO at 100x of the final test concentration and subsequently   171

diluted to 1x in the assay (final DMSO concentration = 1%). Most Kds were ₁₇₂ determined using a compound top concentration of 30000 nM. If the initial Kd ₁₇₃ determined was lower than 0.5 nM (the lowest concentration tested), the measurement ₁₇₄ was repeated with a serial dilution starting at a lower top concentration. A Kd value ₁₇₅ reported as 40000 nM indicates that the Kd was determined to be higher than 30000 ₁₇₆ nM. Binding constants (Kds) were calculated with a standard dose-response curve using ₁₇₇ the Hill equation: ₁₇₈

$$Response = Background + \frac{Signal - Background}{1 + (KdHillSlope/DoseHillSlope)}$$

The Hill Slope was set to -1. Curves were fitted using a non-linear least square fit with ₁₇₉ the Levenberg-Marquardt algorithm. ₁₈₀

Interactions of Ibrutinib and suramin.   Interactions in Fig 3.B were obtained ₁₈₁ from PLIP v1.4.0 [33] applied to PDB IDs 3cjg, 4ifg, 3c7q, 3gan. ₁₈₂

Therapeutic and selectivity index.   In order to test the efficacy and selectivity of ₁₈₃ the hit compounds on B-cells, 4 assays were performed: B-cell assay, Mixed ₁₈₄ Lymphocyte Reaction (MLR), WST-1 assay on RPMI1788 cells (B-cell line) and ₁₈₅ cytotoxicity WST-1 assay on Jurkat cells (T-cell line). Compounds were prepared as 10 ₁₈₆ mM stock solutions in dimethylsulfoxide (DMSO)and were tested at 50-10-1-0.1-0.01 ₁₈₇ and 0.001 μM in the 4 assays. ₁₈₈

Blood samples of healthy volunteers were collected at the Red Cross of Mechelen, ₁₈₉ Belgium. Each donor consents to the use of his blood for research purposes. Human ₁₉₀ peripheral blood mononuclear cells (PBMCs) were obtained by density gradient ₁₉₁ centrifugation of the heparinized venous blood over Lymphoprep™ (Axis Shield PoC AS; ₁₉₂ density 1.077 ± 0.001 g/mL). The human B cell line RPMI1788 (American Type ₁₉₃ Culture Collection, USA) and the human T cell line Jurkat (European Collection of Cell ₁₉₄ Cultures, ECACC, England) were maintained in RPMI1640 culture medium ₁₉₅ (BioWhittaker®, Lonza, Verviers, Belgium) containing 10% foetal calf serum (FCS, ₁₉₆ HyClone® Thermo Scientific, United Kingdom) and 5 μg/mL gentamicin sulphate ₁₉₇ (BioWhittaker®, Lonza, Verviers, Belgium) at 37°C and 5% CO2. ₁₉₈

Highly purified naive peripheral human B cells were separated from fresh human ₁₉₉ PBMCs on magnetic columns by positive selection using CD19 magnetic beads ₂₀₀ according to the manufacturer's instructions (MACS Miltenyi Biotech, Leiden, The ₂₀₁ Netherlands). The purity of the isolated naive B cells was $\geq 95\%$ as analysed by flow ₂₀₂ cytometry. Freshly isolated human CD19+ B cells were then plated at 25 000 cells per ₂₀₃ well in a 384-well plate (Perkin Elmer, Zaventem, Belgium) in 55 μL in DMEM medium ₂₀₄ containing 10% FCS and 5 μg/mL gentamicin sulphate. After 7 days of stimulation ₂₀₅ with 0.1 μM ODN2006 (InvivoGen, Toulouse, France), supernatant was taken for ₂₀₆ analysis of IgG using the AlphaLISA human IgG kit according to the manufacturer's ₂₀₇ instructions (Perkin Elmer, Zaventem, Belgium). Analysis was performed with the ₂₀₈ EnVisionTM 2103 Multilabel Reader (Perkin Elmer, Zaventem, Belgium). ₂₀₉

MLR assay (T cell assay): Freshly isolated human PBMCs and (responder cells) ₂₁₀ were resuspended in RPMI1640 medium containing 10% FCS and 5nμg/mL gentamicin ₂₁₁ sulphate. RPMI1788 cells (stimulator cells) were inactivated by treatment with ₂₁₂ mitomycin C (Kyowa®, Takeda Belgium, Brussels, Belgium) for 20 min at 37 °C, ₂₁₃ washed and finally suspended in culture medium. An amount of 100 μL of each cell ₂₁₄ suspension was mixed with 20 μL of diluted compound. The mixed cells were cultured ₂₁₅ at 37 °C for 6 days in 5% CO2. DNA synthesis was assayed by the addition of 10 μCi ₂₁₆ (methyl-3H) thymidine (Perkin Elmer, Zaventem, Belgium) per well during the last 18 h ₂₁₇ in culture. Thereafter, the cells were harvested on glass filter paper and the counts per ₂₁₈

A Dose response curve for ibrutinib and suramin

B Interaction similarities of ibrutinib and suramin to their targets

Fig 3. Ibrutinib and suramin are inhibitors of VEGFR2. A. Dose response curves for ibrutinib (KD value of 2μM) and suramin (KD value of 25μM) from the competition binding assay determined with the KINOMEscan TM by DisvcoverX prove that ibrutinib and suramin are μM inhibitors of VEGFR2. **B.** Structural repositioning based in interaction of inhibitors (orange) to their targets (blue), predicts ibrutinib as VEGFR2 inhibitor since ibrutinib's interaction to CDPK1 is similar to pazopanib's interaction to VEGFR2. Both have (1) a double hydrogen bond (blue lines) and (2)(3)(4)(5)(6) hydrophobic interactions (gray dashed lines) in common. Similarly, suramin is predicted as VEGFR2 inhibitor. They have (1) a double hydrogen bond, (2) a simple hydrogen bond, (3)(4)(5) hydrophobic interactions and (6) a salt bridge (yellow dashed lines) in common. Subsequent in vitro validation proves both predictions correct.

minute determined in a liquid scintillation counter (TopCount, Perkin Elmer, Zaventem, Belgium).

WST-1 assays on RPMI-1788 cells and Jurkat cells were performed as follows: 20 μL of test compound dilution were added to each well containing exponential growing cells and the plates were incubated for 48 h at 37 °C, 5% CO2. Untreated cells and positive control (1% triton X-100, for the last 15 min) served as reference for maximum and

minimum viability. At the end of incubation 100 μL of supernatant were removed and replaced by 10 μL of WST-1 solution (Cell Proliferation Reagent WST-1, Roche Applied Science). After 3 to 5 h incubation at 37°C, 5% CO2, optical density was measured at 450nm with the EnVisionTM 2103 Multilabel Reader.

B cell Therapeutic Index (B cell TI) was calculated as the ratio IC50 WST-1 assay RPMI1788/IC50 B cell assay; T cell Therapeutic Index (T cell TI) referred to the ratio IC50 WST-1 assay Jurkat/IC50 B CELL assay; while the selectivity index was calculated as the ratio IC50 MLR assay/IC50 B cell assay.

Results

Kinase targets for B-cell inactivation. As a first step towards novel B-cell modulators, we identified suitable kinase targets (see Fig 1.Application). To this end, human kinases were knocked out one by one and the knock out's effect on cell survival and inactivation was measured using the assay introduced in [38]. Cell survival is important, as our goal is modulation of B-cells and not their depletion. As shown in Fig 4.A nearly all human kinases (501) were present in the RNAi library. After knock out, 400 proved to be non-lethal and 22 inactivate B-cells, i.e. the cell surface markers CD70 and CD80 were downregulated in expression (see Table 1).

Fig 4. Selection of targets and inhibitors. A. From 538 human kinases to 22 involved in B-cell activation in Namalwa cells. Nearly all human kinases (501) are present in the RNAi library and for most of them (400) knock out is not lethal. Of those 400, 22 inhibit expression of both B-cell activation markers CD70 and CD80. **B.**From 2538 FDA approved drugs to 1μM B-cell inactivator. Out of the 875 drug with available structures, the *in silico* screening yielded 157 positive binders. From the 110 drugs available for *in vitro* testing, 20 were validated as target inhibitors, 2 validated in kinase binding assay as positive micromolar inhibitors and finally one validated drug with positive effects in B-cells inactivation.

The read out of CD70 and CD80 expression was crucial for this initial target identification. To quantify it, three steps must be considered: without stimulation, CD70 and CD80 expression in B-cells is low; upon stimulation, the markers are upregulated and finally, upon knock out of a kinase essential in B-cell activation, expression of the markers decreases. These two phases, the increase and the decrease, are compared to each other, i.e. it is not sufficient to compare expression levels of non-activated controls to activated knock outs. Instead activated controls have to be factored in. This is captured in the y-value (see Methods), which compares the markers' upregulation after stimulation to their subsequent downregulation (or not) after knock

Table 1. Targets identified in RNAi B-cell activation screen sorted by availability of structural data. The top ten targets are suitable for structure-based drug repositioning screening. For 10 targets (number 11 to 20) no structural were available and thus these target could not be used for structure -based drug repositioning. Silencing of all listed targets leads to strong the downregulation of CD70 and CD80 expressions in the B cell Namalwa cell line activated by the TLR9 agonist ODN2006 without causing cell death (column cell survival). A literature study links nearly all targets to cancer (C) and some to autoimmune diseases (AD) or cancer of the immune system (CIS). Overall, there are many novel autoimmune targets and KDR (VEGFR2) emerges as very promising target with 36 protein structures, potent decrease in expression of CD70 and CD80 following knock out of target in activated Namalwa cells without affecting cell viability and links to both cancer and auto-immunity.

	Gene name	Protein name	PDB structures	CD70 down regulation	CD80 down regulation	% Cell survival	C	CIS	AD
1	DAPK1	Death-associated protein kinase 1	39	1.21	0.87	100	x	x	x
2	**KDR**	**Vascular endothelial growth factor receptor 2 (VEGFR2)**	**36**	**1.27**	**0.98**	**75.6**	**x**	**x**	**x**
3	NEK2	Serine/threonine-protein kinase Nek2	24	1.37	0.79	100	x	x	
4	CSNK1D	Casein kinase I isoform delta (CK1d)	17	0.97	1.28	100	x		
5	EPHB4	Ephrin type-B receptor 4	14	1.95	0.74	100	x		
6	CDC7	Cell division cycle 7-related protein kinase	4	1.88	0.83	100	x		
7	ILK	Integrin-linked protein kinase	2	1.28	0.95	100	x		
8	MST-4	Serine/threonine-protein kinase 26	2	2.63	0.73	100	x		x
9	MAP2K2	Dual specificity mitogen-activated protein kinase kinase 2	2	1.32	0.67	82.9	x		
10	MLK4	Mitogen-activated protein kinase kinase kinase 21	1	1.81	0.54	99.7	x		
11	PXK	PX domain-containing protein kinase-like protein	0	1.16	0.54	94.4			x
12	PTK7	Inactive tyrosine-protein kinase 7	0	1.07	0.71	86.2	x	x	
13	PAK3	Serine/threonine-protein kinase PAK 3	0	2.83	0.98	86.8	x		
14	BRSK1	Serine/threonine-protein kinase	0	1.76	0.83	100	x		
15	FER	Tyrosine-protein kinase Fer	0	1.60	0.95	100	x		
16	IRAK-3	Interleukin-1 receptor-associated kinase 3	0	1.70	1.02	93.3	x		x
17	MAP3K10	Mitogen-activated protein kinase kinase kinase 10	0	1.65	0.93	87.0	x		
18	PRKAG2	5'-AMP-activated protein kinase subunit gamma-2	0	1.57	0.74	100	x		
19	SCYL1	N-terminal kinase-like protein	0	1.18	0.68	84.6	x		
20	STK33	Serine/threonine-protein kinase 33	0	1.43	0.92	100	x		
21	TXK	Tyrosine-protein kinase TXK	0	1.49	1.06	79.9			
22	WNK2	Serine/threonine-protein kinase WNK2	0	1.47	0.86	100	x		

out. Concretely, it is the fraction of expression level of activated control minus knock out and expression level of activated control minus non-activated control. A y-value close to zero indicates that the knock out does not influence B-cell's activation, a value higher than one means that the knock out potentially inhibits activation of B-cells. Table 1 shows these fractions for the best clone of each of the 22 kinases. Besides the definition of the y-value, quality of knock out is important to consider. Per kinase to be silenced there are multiple RNA constructs, hitting different regions of the target gene, which may suffer from different off-targets. Due to this variability, there were between one and ten clones across the 22 kinases with varying degree of effectiveness in silencing. Subsequently, kinases were considered as suitable targets for B-cell inactivation/modulation if at least one clone shows y-value greater 0.5 for CD80 and greater 0.8 for CD70 (for more details check S1 File and S1 Fig).

The 22 kinases, as shown in Table 1, were the starting point for the drug screen. As hinted in the introduction, drugs targeting B-cells are among others used in anti-cancer therapy and against autoimmune diseases. Therefore, we characterized the disease relation of the 22 identified targets focusing in particular on cancer, cancer of the immune system, and autoimmunity. A literature review revealed that nearly all targets are linked to cancer, four to cancer of the immune system, and five to autoimmunity (see Table 1). Among those five is VEGFR2, which is a key cancer target with approved drugs. Clinical trials are ongoing with exploring the inhibition of VEGFR2 as treatment for psoriasis (pazopanib, phase 1) and systemic scleroderma (nintedanib, phase 3). The other four targets with a link to autoimmunity are DAPK1 (inflammatory bowel

disease), MST4 (Grave's disease), PXK (systemic lupus erythematosus), and IRAK3 ²⁷³
(rheumatoid arthritis) (for literature evidence check S1 Table). ²⁷⁴

To assess the 22 kinases' potential as drug targets we consulted the Open Targets ²⁷⁵
Platform [5], which covers over 20.000 targets with associated information on drugs and ²⁷⁶
clinical trials. VEGFR2, EPHB4, CDC7, MAP2K2, and MAP3K10 are listed with ²⁷⁷
VEGFR2 as most established target with a number of drugs already approved. ²⁷⁸
Furthermore, we checked BindingDB [12], the largest database for drug-target binding ²⁷⁹
affinities and found 22 drugs, each binding one or more of the 22 targets (see S3 Fig). ²⁸⁰
14 of the 22 kinases have at least one inhibitor in micromolar range and all but six ²⁸¹
drugs are polypharmacological drugs hitting multiple targets. In fact, 15 FDA approved ²⁸²
drugs inhibit the same 13 kinases including VEGFR2, which documents the ²⁸³
polypharmacological potential of these drugs. Most of these drugs are chemically closely ²⁸⁴
related. Consider for example, erlontinib, vandetanib, and gefitinib (see Fig 5.A for ²⁸⁵
their chemical structure), which belong to the same chemical class of quinazolines. To ²⁸⁶
find a different compound with a better profile of toxicity and specificity we aimed to ²⁸⁷
expand the scaffold space by running a structure-based drug repositioning screen. ²⁸⁸

Fig 5. Chemical structures of known (A) and predicted (B) inhibitors. Drugs with drug name and compound ID (CID) are grouped by therapeutic indication: Autoimmune diseases(green), Cancer immune system (blue), Cancer (red), Infections (yellow), and others (gray).

Structure-based screening to expand scaffold space. The starting point of a ²⁸⁹
structure-based drug repositioning screen is a template drug-target complex, from which ²⁹⁰
the chemical space beyond the drug's scaffold and the given target's fold is explored. A ²⁹¹
structure-based screening ignores the template's global structure and instead focuses ²⁹²
specifically on two aspects only: the binding site and the drug's interaction to it (for ²⁹³
more details see S2 File). These two features were extracted from the template and ²⁹⁴
were compared to other drug-target complexes by aligning binding sites and scoring the ²⁹⁵
similarity between the interactions (see Fig 2). ²⁹⁶

A prerequisite for structure-based screening is the availability of structural templates. Table 1 shows that 10 out of the 22 kinase targets have structural data available. For these 10 kinases targets there were a total of 141 templates. They were compared to over 300.000 ligand-target complexes in the Protein Databank PDB and ranked by p-value. The table also shows that there is huge variation from only one structure to over 30 for VEGFR2. Regarding candidate drugs, which may be predicted, PDB covers suitable structural data for 875 out of some 2500 FDA approved drugs. Based on these data, the structural drug repositioning screen predicts 157 approved drugs (see Fig 4.B). The 157 predictions are not equally distributed across the ten targets, but they correlate with the amount of structural data available for the target. e.g. for VEGFR2 with its 36 structures, there are 53 predictions, while MLK4 with one structure has three predictions only (for more details see S3 File).

Subsequently these 110 drugs were tested at 10 μM and 1 μM on the panel of the 10 selected kinases (for more details see S4 File). We considered a 50% inhibition of the kinase activity as strong and 30 to 50% inhibition as weak. Table 2 summarizes the result. Only one compound, Ibrutinib, displays strong inhibition at 10 and 1 μM on one of the kinase (VEGFR2); while 9 other compounds show strong inhibition at 10 μM only. Ten additional compounds show weak inhibition at 10 μM.

Table 2. List of new drug-target interactions predicted by interaction profile similarity and confirmed by experimental kinase activity assay.
Enzyme activity was measured at a compound concentration of 1μM and 10μM. Drugs able to lower the enzyme activity to less than 50% (above dashed line) at 10μM were defined as strong inhibitors while drugs that lowered kinase activity at 10μM between 50% and 70% (below dashed line) were classified as weak inhibitors. Discarding the disinfectant Hexachlorophene because of its low medical interest, two compounds have been highlighted for showing inhibition at both 1μM and 10μM, ibrutinib and suramin .

Compound	Target	% Enzyme activity at 1μM	% Enzyme activity at 10μM
Hexachlorophene	NEK2	84.92	00.02
Ibrutinib	**VEGFR2**	**19.52**	**01.50**
Hexachlorophene	VEGFR2	109.5	05.14
Suramin	**VEGFR2**	**72.67**	**06.48**
Toremifene	DAPK1	90.42	14.77
Ibrutinib	DAPK1	95.59	23.72
Saquinavir	DAPK1	87.09	27.81
Tolcapone	CK1d	91.69	35.58
Ritonavir	DAPK1	92.58	38.46
Etravirine	VEGFR2	108.4	41.08
Thyroxine	NEK2	90.16	48.11
Lopinavir	DAPK1	90.43	49.89
Adenosine	CK1d	87.72	50.22
Diethyldtilbestrol	NEK2	97.92	50.73
Proflavine	NEK2	104.1	51.57
Triiodothyronine	NEK2	111.9	55.67
Adenosine	NEK2	96.13	58.99
Triclosan	NEK2	99.17	64.62
Triclosan	CDC7	70.45	65.43
Propofol	EPHB4	66.96	68.44
Prochlorperazine	NEK2	70.10	68.73
Estradiol	DAPK1	110.6	68.98
Amodiaquine	CK1d	107.2	69.07
Niacinamide	EPHB4	69.30	69.74

Thus, taking all analyses together, we have identified a total of 42 drugs (Fig 5), 22 known and 20 predicted and validated, which target at least one of the 20 target kinases and which thus carry the potential to inactivate B-cells. Before testing this potential,

we summarize the chemical and disease space captured through these 42 drugs.   318

Chemical and disease space of identified drugs. Fig 6 shows a   319
drug-target-disease network for the 42 drugs. The network depicts nicely that the   320
majority of known drugs are used in cancer and are promiscuously targeting multiple   321
targets. This is complemented by tofacitinib, an autoimmunity drug targeting many   322
shared targets as well. In contrast, there is a sparse region in the network, where   323
anti-infectives hit some of the targets including in particular VEGFR2. Ibrutinib is the   324
exception to cancer drugs targeting only two targets.   325

Fig 6. Drug-target-disease network. Inner circle (14 targets), mid circle (known
and predicted inhibitors), outer circle (diseases). There is a tight network of cancer
drugs and targets while there is only one known autoimmune compound, which binds,
however, many of the cancer targets. VEGR2, the promising autoimmune target, we
identified, is inhibited by cancer and anti-infection drugs.

The dense region of cancer drugs in the network may be due to their chemical   326
similarity and resulting similar target profiles. Conversely, it can be expected that   327
anti-infectives will have very different scaffolds. To understand this relation in detail, we   328
compared the chemical structure of all 42 drugs against each other. Consider Fig 5 with   329
the drugs' structure. Visual inspection already reveals some similar scaffolds. As   330
mentioned above, the anti-cancer drugs erlotinib, vandetanib, gefitinib, and afatinib are   331
all quinazoline derivatives. To systematically assess these similarities we clustered the   332
drugs by their chemical similarity according to PubChem's CACTVS fingerprints [22].   333
CACTVS fingerprints represent chemicals as binary vectors of 880 dimensions. Each   334
dimension represents a specific chemical feature. The chemical similarity of two drugs   335
can then be defined as number of shared features divided by the overall number of   336

features present in the two drugs, i.e. the Tanimoto score of the two fingerprints. The ₃₃₇
similarity can range between 0 (dissimilar) and 1 (similar). ₃₃₈

Fig 7 depicts these chemical similarities for all pairs of the 42 drugs. Dark blue ₃₃₉
indicates high similarity. The dark blue group of four compounds in the top left ₃₄₀
quadrant represents the highly similar erlotinib, vandetanib, gefitinib, and afatinib ₃₄₁
mentioned above. This high-level view of the chemical compound space breaks down ₃₄₂
into the top left quadrant of similar compounds and the bottom right quadrant, which ₃₄₃
is completely dissimilar from the top left quadrant. To understand this grouping, we ₃₄₄
enriched the figure with additional information. On the bottom horizontal axis we ₃₄₅
labelled known binders in orange and novel binders in green. Broadly, the top left ₃₄₆
quadrant contains the known compounds, whereas the novel ones are the bottom right ₃₄₇
quadrant. This supports that the structural screen identified novel scaffolds not present ₃₄₈
in the known binders. Furthermore, we added to the vertical axis on the left the disease ₃₄₉
information for the drugs and on the right the number of targets. These bars show that ₃₅₀
the known binders in the top left quadrant have many targets and are anti-cancer, ₃₅₁
whereas the novel compounds in the bottom right cover various diseases and have few ₃₅₂
targets. At the border between known and novel quadrant is ibrutinib, an anti-cancer ₃₅₃
drug structurally close to many of the known drugs but not a known binder to any of ₃₅₄
the targets according to BindingDB. Furthermore, suramin, an anti-infective shows ₃₅₅
some weak structural similarity to the cancer drug mitoxantrone. The former has only ₃₅₆
very few targets, while the latter is highly promiscuous. ₃₅₇

Ibrutinib and suramin are micromolar VEGFR2 inhibitors. Ibrutinib and ₃₅₈
suramin emerged as promising leads from the kinase activity assay. The activity of ₃₅₉
ibrutinib on VEGFR2 is also confirmed by other experimental results in the LINCS ₃₆₀
kinomescan database [21]. To further sustain these findings, we carried out a dose ₃₆₁
response study. The curve in Fig 3.A plots drug concentration in nM against amount of ₃₆₂
bound kinase. Both experiments were carried out in duplicates. Ibrutinib binds to ₃₆₃
VEGFR2 at a Kd of 2μM and suramin at a Kd of 25μM. ₃₆₄

To better understand how ibrutinib and suramin achieve this high affinity against ₃₆₅
VEGFR2, we checked their interaction profiles. The prediction of ibrutinib was based ₃₆₆
among others on ibrutinib's interaction with CDPK1 (PDB 4ifg), which is very similar ₃₆₇
to pazopanib's interaction with VEGFR2 (PDB 3cjg). This is a remarkable similarity, ₃₆₈
as CDPK1 is from the parasite Toxoplasma gondii and VEGFR2 from human. Despite ₃₆₉
two targets from completely different species, Fig 3.B shows that both interactions ₃₇₀
comprise a double hydrogen bond (1) and five hydrophobic interactions in similar ₃₇₁
position (2), (3), (4), (5), and (6). In a similar way, suramin's binding to RDM1 (PDB ₃₇₂
3gan) is similar to nintenadib's interaction with VEGFR2 (PDB 3c7q). Again, this ₃₇₃
similarity is very remarkable, as RDM1 is a regulator of DNA methylation in ₃₇₄
Arabidopsis thaliana and VEGFR2 is from human. Nonetheless, both comprise in their ₃₇₅
ligand interaction a double hydrogen bond (1), a simple hydrogen bond (2), three ₃₇₆
hydrophobic interactions (3), (4), (5) and a salt bridge (6). ₃₇₇

These examples not only document scaffold hopping from pazopanib to ibrutinib and ₃₇₈
from nintenadib to suramin, but also target hopping from parasitic CDPK1 to human ₃₇₉
VEGFR2 and from plant RDM1 to human VEGFR2. These jumps in target space are ₃₈₀
large. They cross family and species boundaries. As such they illustrate the limited ₃₈₁
space of binding sites, which was eluded to in the introduction. ₃₈₂

Ibrutinib inactivates B-cells. In a final validation step, we tested the drugs' ability ₃₈₃
to impede B-cell's activation (for more details see S5 File). We wanted inhibition of ₃₈₄
activation without cell death and specific to B-cells. We introduce the therapeutic index, ₃₈₅
which relates toxicity to inactivation. We calculate it for B- and T-cells. From this we ₃₈₆

Fig 7. Pairwise chemical similarity of known and predicted inhibitors.
Known inhibitors labeled with orange tag on bottom line and predicted ones with a
green tag on bottom line. Dark blue boxes represent high chemical similarity of
inhibitors (see Methods). Bars on the left are therapeutic indication (red=cancer,
blue=cancer of the immune system, green=autoimmune disease, yellow=infection, and
gray=other) and gray bars on the right represent the number of known targets (size of
bar). The top left quadrant are mostly known inhibitors, which are chemically similar
to each other and are mostly linked to cancer. The bottom right quadrant are new,
predicted inhibitors, which have different scaffolds and are hence not derived from the
known inhibitors. Consequently, these inhibitors have other therapeutic indications.

compute the specificity index as ratio of B-cell to T-cell therapeutic index. Our goal is ₃₈₇
to uncover compounds with high B-cell therapeutic index and high specificity index. ₃₈₈
Indices are computed from duplicates. For details on cell lines used see Methods section. ₃₈₉

As a main result, ibrutinib has a very high B-cell therapeutic index of over 40000 ₃₉₀
confirming ibrutinib's ability to inactivate B-cells at low toxicity. A selectivity index of ₃₉₁
over 5000 also means that ibrutinib interferes only little with T-cells. Table 3 ₃₉₂
summarize these indices including 15 other compounds. For suramin, the therapeutic ₃₉₃
and specificity indices are close to one and hence suramin is not a B-cell modulator, ₃₉₄
although it is a micromolar VEGFR2 inhibitor. Tofacitinib, which can be seen as a ₃₉₅
control, due to its approval for autoimmune indications, has a good B-cell therapeutic ₃₉₆
index but lacks specificity. ₃₉₇

Table 3. Potency on B-cell, B-cell therapeutic index (TI), T-cell therapeutic index, and specificity index (SI). The B cell IC50 is given in μM and is the mean of 4 independent experiments. The therapeutic index relates toxicity to activity. B cell Therapeutic index (B cell TI) is the ratio between the toxic concentration and active concentration on B-cell and was calculated as IC50 WST-1 assay RPMI1788/IC50 B-cell assay. T cell Therapeutic index (T cell TI) is the ratio between toxic and active concentrations on T-cells and was calculated as IC50 WST-1 assay Jurkat/IC50 B-cell assay. The selectivity index is the ration between active concentrations on T-cells and B-cells and was calculated as IC50 MLR assay/IC50 B-cell assay. A high value of Therapeutic Index corresponds to high activity at low toxicity and is hence desirable. A high value of Selectivity Index indicates high specificity to B-cells rather than T-cells. Ibrutinib has high B-cell TI and SI.

Drug	B-cell IC50 (μM)	B-cell TI	T-cell TI	T/B-cell SI
Ibrutinib	**0.0002**	**3856**	**81090**	**5530**
Dasatinib	0.046	0,71	160,0	0,07
Tofacitinib	**0.432**	**>115,6**	**>115,6**	**0,61**
Proflavine	0.59	5,14	8,55	7,50
Amodiaquin	0.66	42,33	39,43	14,73
Hexachlorophene	5.34	0,79	1,10	0,27
Etravirine	5.90	1,35	5,15	1,23
Prochlorperazine	6.12	1,67	3,52	0,76
Triclosan	20.88	1.32	1,25	1.64
Diethylstilbestrol	31.07	1,01	0,22	0,59
Estradiol	33.91	1,06	0,58	0,62
Thyroxine	34.10	>1,47	>1,47	>1,47
Suramin	**35.20**	**>1,42**	**>1,42**	**0,84**
Thyronine	36.82	>1,36	>1,36	1,20
Adenosine	>50	1,00	1,00	1,00
Nicotinamide	>50	1,00	1,00	1,00

Discussion

Drug promiscuity is a common phenomenon and we have exploited it here in a structure-based drug repositioning screen to identify ibrutinib, a novel VEGFR2 inhibitor and a B-cell inactivator.

VEGFR2 is a receptor-type tyrosine kinase (RTK) with a key role in the regulation of angiogenesis and lymphangiogenesis, which makes it an attractive target for antiangiogenic and proangiogenic therapies [35]. A lot of VEGFR2 neutralizing agents have been developed for cancer treatment, such as the human antibody bevacizumab (Avastin) [35], and the kinase inhibitors sunitinib, sorafenib, axitinib and vandetanib. This kinase showed already an interesting connection with the autoimmune disease psoriasis [17], and two of its inhibitors have been repositioned from the anti cancer indication to immunomodulation and are currently in clinical trials for the treatment of psoriasis (pazopanib) and systemic scleroderma (Nintedanib).

While this discovery of a confirmed binder of a new scaffold proves the power of the in silico approach, it raises the question of the approach's general applicability. In drug repositioning, one can take a target- or a drug-centric approach. In a target-centric approach, the target of the repositioned drug is the key. It was associated with one disease and is now found to be relevant in another disease. Hence, the drug may be repositioned from the first to the second disease. In a drug-centric approach to drug repositioning, the prediction of a novel target, which is linked to a disease, forms the

base for repositioning. In this work, we pursued both approaches. ₄₁₈

The RNAi screen on the human B-cell line Namalwa identified novel B-cell targets ₄₁₉ by using CD70 and CD80 expression as read-out for B-cell activity. We must clarify ₄₂₀ that the role of CD70 and CD80 is not exclusive of B-cells activity, and therefore their ₄₂₁ inhibition might also be important in contrasting T-lymphocytes over-activation with ₄₂₂ an advantageous pleiotropic effect in immunosuppressive therapy. Among the various ₄₂₃ stimuli investigated for B-cell activation, in vitro stimulation with TLR9 agonist ₄₂₄ ODN2006 resulted in the broadest phenotypic change profile in human polyclonal B-cells. ₄₂₅ Compared with an assay of human B-cell IgG, which checks the level of IgG production ₄₂₆ by primary B-cells after 7 days incubation with the same stimulation agent ODN2006, ₄₂₇ the Namalwa-ODN2006 protocol takes only 24 hours, making it ideal for a fast first line ₄₂₈ screening. Furthermore, the heterogeneity between the different human blood donors, ₄₂₉ the heterogeneity after isolation and purification from peripheral blood, the limited ₄₃₀ yield of the purification and the short longevity make primary B-cells less optimal for ₄₃₁ repeatable assays. So a monoclonal population of dividing cells such as the human ₄₃₂ B-cell line Namalwa stabilizes the assay by overcoming the aforementioned limitations. ₄₃₃

Our analysis of known binders to these targets is target-based drug repositioning. It ₄₃₄ was fruitful as many of the targets were established drug targets with associated ₄₃₅ approved drugs. In fact, our analysis showed that these drugs are highly promiscuous ₄₃₆ and hit many of the targets, which is possibly an intended polypharmacological design. ₄₃₇ Their chemical analysis showed highly similar compounds of the same scaffolds. ₄₃₈ Therefore, we applied structure-based screening to implement scaffold hopping away ₄₃₉ from these known binders into novel, uncharted chemical space. This second approach ₄₄₀ was drug-centric, as the structure-based screen across the Protein Databank PDB led to ₄₄₁ novel drug-target predictions. The most prominent ones were suramin and ibrutinib, ₄₄₂ both binding VEGFR2. If we consider this repositioning at a higher level, then suramin ₄₄₃ may be moved from infectious disease (sleeping sickness and river blindness) to cancer ₄₄₄ and ibrutinib from cancer to autoimmunity. While both drugs suffer from side effects, ₄₄₅ they are nonetheless serious drug repositioning candidates: suramin is tested in a phase ₄₄₆ I/II study against autism (see clinicaltrials.gov) and the FDA approved ibrutinib in ₄₄₇ 2017 for host-vs-graft disease. ₄₄₈

While the above results focus on the most established target, VEGFR2, the RNAi ₄₄₉ screen generated another 21 novel B-cell targets overall. For some there is a link to ₄₅₀ autoimmunity and for a few there are drugs known to inhibit them. However, future ₄₅₁ work should fully explore all candidates regarding their relation to disease. This is ₄₅₂ difficult for any target and any drug: Even if a drug and a target can be linked to a ₄₅₃ disease, it still does not necessarily mean that the drug acts through the target. As an ₄₅₄ example, the RNAi screen suggests that silencing of VEGFR2 leads to B-cell ₄₅₅ inactivation, the inhibition of VEGFR2 with ibrutinib confirmed this effect, whereas ₄₅₆ inhibition with suramin did not. Generally, besides a methodological difference in ₄₅₇ silencing and inhibition, ibrutinib may have a more favorable polypharmacological ₄₅₈ profile for B-cell inactivation than suramin has. In a nutshell, both target- and ₄₅₉ drug-centric drug repositioning can be successful. The latter has the advantage of ₄₆₀ moving to novel scaffolds. And last, but not least, the structure-based drug ₄₆₁ repositioning does not only comprise the prediction, but also a transparent ₄₆₂ physio-chemical explanation for the prediction. Thus, giving confidence and a starting ₄₆₃ point for subsequent optimization steps. ₄₆₄

A prerequisite for these successes is the availability of structures available in the ₄₆₅ Protein Data Bank (PDB). While there are far less structures than sequences in public ₄₆₆ databases, the growth of structural data is none the less impressive: PDB has more ₄₆₇ than doubled in size over the last seven years. Today, it contains 3D structures of over ₄₆₈ 1200 different drug targets and more than 60% of all PDB structures contain proteins in ₄₆₉

complexe with biologically relevant ligands. For the present screen, PDB contained suitable structures for about half of the targets. S2 Fig shows how these structures became available over time. It reveals that 15 years ago, structure-based drug repositioning would not have been possible for the identified 22 targets. Assuming a continuing linear growth, the figure suggests that in another 15 years the complete set of targets can be surveyed. Thus, the potential of structure-based drug repositioning is visible today and will further increase over time. Technological advance is also likely to speed up progress, since improvements in structure prediction and modelling may soon reach the maturity to close the structural gap faster.

The structural drug repositioning screen as carried out here considered two aspects: the ligand-target interaction and superposition of binding site and ligand. Generally, many approaches in the past have focused either on the ligand (see e.g. [6]) or on the target (geometrical and/or chemical similarities of the target proteins [20]). Our approach combined both views. The quality of such an integrated approach can be measured by the hit rates achieved. Here, we considered 110 compounds after prediction and filtering steps. For 15 of these prediction there was experimental evidence of high affinity binding in BindingDB and for other 20 unknown binders our kinase inhibition assay provided confirmation. This corresponds to a very high hit rate of 32%.

Such hit rates are unprecedented in traditional high-throughput screening, but they come at a cost. In classical high-throughput screening, there is a large compound library, which may be already focused on a specific type of compounds. In contrast, the structural screen is limited by the compounds in PDB. Overall, there are over 30000, which is substantial. But in comparison to the million of compounds available in databases such as PubChem it is small. This gap in size is reduced if one adds additional selection criteria. In drug repositioning, this can be e.g. the approval of the compounds. There are some 2500 approved drugs and for the structural screening some 800 are available in PDB. So, the orders of magnitude in size difference reduce to a factor of three. A possible disadvantage of a smaller library may be the inability to cover a large chemical space. However, this problem does not arise in the structural screen as carried out here, since the focus on drug-target interaction supported scaffold hopping. It ensured that huge steps in chemical space were possible and were indeed observed in the resulting data. Finally, *in silico* screening implies low cost and time. Instead of 8750 validation experiments (875 approved drugs with structure times 10 targets), there were only some 100, a cost reduction by nearly two orders of magnitude. In summary, the combination of a small in silico library with the algorithmic approach ultimately leads to a very high hit rate over a large chemical space at a low screening cost.

Other computational approaches also pursue *in silico* drug-target prediction by using text mining, machine learning or chemical similarities, among other methods. When comparing our method to the three most cited approaches [9, 11, 25], in order to check whether they are able to predict the ibrutinib-VEGFR2 connection or not, we came out with the following: SwissTargetPrediction (129 citations) is based on similarity measures of chemical structure and molecular shape. Performing a target screening for Ibrutinib, it predicted a long list of TYR kinases, including EGFR, but no specific link to VEGFR2. On the other hand, SuperPred (98 citations) is based on structural fingerprint similarities and text mining for target prediction. It identifies no known targets for Ibrutinib and while it is able to predict 5 TYR kinases with good E values, non of them is a direct link to VEGFR2. Finally, Hitpick (45 citations) combines two methods (1-Nearest-Neighbour and machine learning) in order to calculate 2D similarity. For ibrutinib, it predicted only 2 TYR kinases as targets, but non of them VEGFR2.

Although they all successfully predicted targets from the Receptor protein tyrosine kinase class, which is the same of VEGFR2, none of them was able to actually make the direct link as our method did. Furthermore, all of the mentioned approaches are based

in 2D/3D similarity measures of the compounds and in available assay data coming ₅₂₂
from databases. Therefore, their screening results often in similar scaffolds and protein ₅₂₃
classes, being unable to make such specific prediction for distant biochemical entities. ₅₂₄

Conclusion ₅₂₅

B-cells play an important role in diseases ranging from auto-immunity to cancer. There ₅₂₆
is an unmet need for small molecules inactivating B-cells. Here, we present ibrutinib as ₅₂₇
such a candidate immuno-modulator. In a first step, we ran an RNAi screen with a ₅₂₈
recently introduced drug screening assay for B-cell inactivation. We found 22 kinases, ₅₂₉
whose silencing inactivates B-cells. Many of these targets are already drug targets with ₅₃₀
approved drugs, which will be beneficial for future drug repositioning efforts. We ₅₃₁
focused specifically on VEGFR2, as it is already an established cancer drug target and ₅₃₂
there is evidence for a role in autoimmunity, too. For lead identification, we pursued ₅₃₃
two approaches. We evaluated existing drugs affecting the identified targets. As we ₅₃₄
found mostly chemically similar cancer drugs with a polypharmacological design hitting ₅₃₅
many of our targets, we wanted to expand the scaffold space. We used structural drug ₅₃₆
repositioning to predict novel inhibitors, which we then validated in an enzyme activity ₅₃₇
and binding study. We identified the cancer drug ibrutinib and the anti-infective ₅₃₈
suramin as micromolar VEGFR2 inhibitors. In a final step, we tested B-cell inactivation ₅₃₉
and found ibrutinib as a very effective inhibitor with very high therapeutic index and ₅₄₀
good specificity to B-cells rather than T-cells. The discovery is particularly remarkable ₅₄₁
as the underlying scaffold hopping is tied to target hopping. At the source of the ₅₄₂
prediction are similarities in the binding sites of human VEGFR2 to a parasitic and to a ₅₄₃
plant protein. These similarities are remarkable as they are unlikely of an evolutionary ₅₄₄
origin since not only family boundaries are crossed, but also species boundaries. Instead, ₅₄₅
they document the large, but limited number of binding sites, as postulated in [2, 10]. ₅₄₆
Overall, the in silico structural screen produced results at a very high hit rate, which ₅₄₇
translates into low cost and time. It was able to predict chemically distant compounds ₅₄₈
thus covering a large chemical space. ₅₄₉

Supporting information ₅₅₀

S1 File.   RNAi screening. Results of the shRNA transfection of human Burkitt ₅₅₁
lymphoma cells (Namalwa) stimulated by the TLR9 ligand ODN2006. Per each clone ₅₅₂
the name of the silenced gene ('Gene' column), the number of Namlwa cells aquired ₅₅₃
by the FACS ('Cell number'), the percentage of inhibition of over-expression of surface ₅₅₄
proteins CD70 ('% inh CD70') and CD80 ('% inh CD70') are reported. ₅₅₅

S2 File.   Compounds screening. Dataset of 157 compounds resulting from the ₅₅₆
computational screening based in interaction similarities. It contains the compound ₅₅₇
name, compound id (CID), inchikey, drug category if possible, binding evidence coming ₅₅₈
from Binding DB, related diseases coming from PubChem, best p-value and finally ₅₅₉
individual P-values for each compound-targets prediction. ₅₆₀

S3 File.   Protein-drug predictions. List of 1828 predictions between the 157 ₅₆₁
compounds and the 22 targets. It contains the compound id (CID), the protein target, ₅₆₂
the complex id (pdbid:hetid:chain:position) of the template, the complex id of the hit, ₅₆₃
the method making the prediction and the p-value of the prediction. ₅₆₄

S4 File.   Activity assay. Results of the kinase activity assay performed for the 111
purchased compounds. It contains the compound name, the target name and the
enzyme activity value of 2 data points at 1 and 10 μM.

S5 File.   Bcell assay. Results of the efficacy and cytotoxicity tests of 16 of our
kinase inhibitors ('Compound' column) on B-cell and T-cell lines. The four type of
assays are reported: 'B-cell efficacy', 'T-cell efficacy', 'Cytotoxicity/ WST1 RPMI 1788'
and 'Cytotoxicity/ WST-1 Jurkat'. For each assay the mean of 4 different independent
measures and the standard deviation (SD) are reported. Three different indexes have
been calculated: 'Therapeutic Index (TI)/ WST1 RPMI 1788/B-cell' (B-cell/B-cell),
'Therapeutic Index (TI)/ WST1 Jurkat/B-cell' (T-cell/B-cell), and 'Selectivity Index
(SI)/ MLR/B-cell' (T-cell/B-cell). Mycophenolate Mofetyl and Cyclosporine A have
been used as positive controls.

S1 Fig.   RNAi screening for target identification. The level of inhibition of
upregulation (y) , of CD70 (dark blue dots) and CD80 (light blue dots) for each clone of
the selected genes (x axe) have been plotted plotted. The clones laying in the bottom
part of the graph with y ¡ 0 (red part), showed an expression of the surface receptor
(Ssample or Ss on the side bar) higher than the stimulated control cells (Sctrl or Sc); the
clones (Ss) with 0 ¿ y ¿1 (yellow part) had a level of CD70/CD80 expression lower than
the stimulated controls (Sc) but higher than the non-stimulated control cells (NSctrl or
NSc); the clones (Ss) placed in the area with y ¿1 (green part) showed an exprssion of
the activation markers lower also than the non-stimulated controls NSc . Those genes
have been selected for showing the y of at least one clone above both the threshold of
0.8 for CD70 (dark blue dotted line) and 0.5 for CD80 (light blue dotted line).

S2 Fig.   Available structures for 22 targets over time. Over the last 15 years,
structures for half of the identified target kinases were deposited in PDB, so that today
there is sufficient data for structure-based drug repositioning available. Before the year
2002, this type of screening would not have been possible. In the future, it will further
improve.

S3 Fig.   Known drugs bind unspecifically. Known drugs binding 22 identified
kinases with Ki, Kd, or IC50 ¡50μM. Drugs are colored by indication: red for cancer,
blue for cancer of immune system, green for immunomodulation, yellow for infection
and gray for other indications. Overall, the majority of drugs are anti-cancer and bind
unspecifically.

S1 Table.   Literature evidence for a target's association to disease. Links in
litterature between each gene target ('Target' column) and some pathological conditions
('Disease' column) such as Cancer, Tumors of Immune System (Lymphoma, Leukemia
and Multiple Myeloma), and Autoimmune Diseases (Inflammatory Bowel Disease,
Psoriasis, Lupus Erythematosus, Graves' Disease and Rheumatoid Arthritis) are
reported ('PMID' column).

Acknowledgments

Thanks to Florian Kaiser and Arnout Voet for valuable feedback.

Author contributions statement

MA, DP, MS conceived the study, generated and analysed data, and wrote the manuscript. SS, VJH, MA implemented structural screening. GJ, JCH, JH, KvB, BS carried out experiments and analysed data.

Competing interests

The authors declare no competing financial interests.

References

1. Tobias Alexander, Ramona Sarfert, Jens Klotsche, Anja A Kühl, Andrea Rubbert-Roth, Hannes-Martin Lorenz, Jürgen Rech, Bimba F Hoyer, Qingyu Cheng, Aderajew Waka, et al. The proteasome inhibitior bortezomib depletes plasma cells and ameliorates clinical manifestations of refractory systemic lupus erythematosus. *Annals of the rheumatic diseases*, pages annrheumdis–2014, 2015.

2. Patrick Aloy and Robert B Russell. Ten thousand interactions for the molecular biologist. *Nature biotechnology*, 22(10):1317, 2004.

3. Andrew Anighoro, Jurgen Bajorath, and Giulio Rastelli. Polypharmacology: challenges and opportunities in drug discovery: miniperspective. *Journal of medicinal chemistry*, 57(19):7874–7887, 2014.

4. Ramon Arens, Kiki Tesselaar, Paul A Baars, Gijs M.W van Schijndel, Jenny Hendriks, Steven T Pals, Paul Krimpenfort, Jannie Borst, Marinus H.J van Oers, and René A.W van Lier. Constitutive cd27/cd70 interaction induces expansion of effector-type t cells and results in ifn-mediated b cell depletion. *Immunity*, 15(5):801 – 812, 2001.

5. Denise Carvalho-Silva, Andrea Pierleoni, Miguel Pignatelli, ChuangKee Ong, Luca Fumis, Nikiforos Karamanis, Miguel Carmona, Adam Faulconbridge, Andrew Hercules, Elaine McAuley, et al. Open targets platform: new developments and updates two years on. *Nucleic acids research*, 47(D1):D1056–D1065, 2018.

6. Zhan Deng, Claudio Chuaqui, and Juswinder Singh. Structural interaction fingerprint (sift): a novel method for analyzing three-dimensional protein- ligand binding interactions. *Journal of medicinal chemistry*, 47(2):337–344, 2004.

7. Jérémy Desaphy, Guillaume Bret, Didier Rognan, and Esther Kellenberger. sc-pdb: a 3d-database of ligandable binding sites—10 years on. *Nucleic acids research*, 43(D1):D399–D404, 2014.

8. Je´re´my Desaphy, Eric Raimbaud, Pierre Ducrot, and Didier Rognan. Encoding protein–ligand interaction patterns in fingerprints and graphs. *Journal of chemical information and modeling*, 53(3):623–637, 2013.

9. Mathias Dunkel, Stefan Günther, Jessica Ahmed, Burghardt Wittig, and Robert Preissner. Superpred: drug classification and target prediction. *Nucleic acids research*, 36(suppl_2):W55–W59, 2008.

10. Mu Gao and Jeffrey Skolnick. Structural space of protein–protein interfaces is degenerate, close to complete, and highly connected. *Proceedings of the National Academy of Sciences*, 107(52):22517–22522, 2010.

11. David Gfeller, Aurélien Grosdidier, Matthias Wirth, Antoine Daina, Olivier Michielin, and Vincent Zoete. Swisstargetprediction: a web server for target prediction of bioactive small molecules. *Nucleic acids research*, 42(W1):W32–W38, 2014.

12. Michael K Gilson, Tiqing Liu, Michael Baitaluk, George Nicola, Linda Hwang, and Jenny Chong. Bindingdb in 2015: a public database for medicinal chemistry, computational chemistry and systems pharmacology. *Nucleic acids research*, 44(D1):D1045–D1053, 2015.

13. Christopher EM Griffiths and Jonathan NWN Barker. Pathogenesis and clinical features of psoriasis. *The Lancet*, 370(9583):263–271, 2007.

14. Douglas Hanahan and Robert A Weinberg. Hallmarks of cancer: the next generation. *Cell*, 144(5):646–674, 2011.

15. Helen M. Berman, John Westbrook, Zukang Feng, et al. The Protein Data Bank *Nucleic Acid Research.* 235–242, 2000.

16. V Joachim Haupt, Simone Daminelli, and Michael Schroeder. Drug promiscuity in pdb: protein binding site similarity is key. *PLoS one*, 8(6):e65894, 2013.

17. Regina Heidenreich, Martin Röcken, and Kamran Ghoreschi. Angiogenesis drives psoriasis pathogenesis. *International Journal of Experimental Pathology*, 90(3):232–248, 2009.

18. Jörg C Heinrich, Sainitin Donakonda, V Joachim Haupt, Petra Lennig, Yixin Zhang, and Michael Schroeder. New hsp27 inhibitors efficiently suppress drug resistance development in cancer cells. *Oncotarget*, 7(42):68156, 2016.

19. Andrew L Hopkins. Network pharmacology: the next paradigm in drug discovery. *Nature chemical biology*, 4(11):682, 2008.

20. Howook Hwang, Fabian Dey, Donald Petrey, and Barry Honig. Structure-based prediction of ligand–protein interactions on a genome-wide scale. *Proceedings of the National Academy of Sciences*, page 201705381, 2017.

21. Alexandra B Keenan, Sherry L Jenkins, Kathleen M Jagodnik, Simon Koplev, Edward He, Denis Torre, Zichen Wang, Anders B Dohlman, Moshe C Silverstein, Alexander Lachmann, et al. The library of integrated network-based cellular signatures nih program: system-level cataloging of human cells response to perturbations. *Cell systems*, 6(1):13–24, 2018.

22. Sunghwan Kim, Jie Chen, Tiejun Cheng, Asta Gindulyte, Jia He, Siqian He, Qingliang Li, Benjamin A Shoemaker, Paul A Thiessen, Bo Yu, et al. Pubchem 2019 update: improved access to chemical data. *Nucleic acids research*, 47(D1):D1102–D1109, 2018.

23. Zachary A Knight, Henry Lin, and Kevan M Shokat. Targeting the cancer kinome through polypharmacology. *Nature Reviews Cancer*, 10(2):130, 2010.

24. Antonio Lavecchia and Carmen Cerchia. In silico methods to address polypharmacology: current status, applications and future perspectives. *Drug Discovery Today*, 21(2):288–298, 2016.

25. Xueping Liu, Ingo Vogt, Tanzeem Haque, and Mónica Campillos. Hitpick: a web server for hit identification and target prediction of chemical screenings. *Bioinformatics*, 29(15):1910–1912, 2013.

26. Henault M, Lee LN, Evans GF, and Zuckerman SH. The human burkitt lymphoma cell line namalwa represents a homogenous cell system characterized by high levels of toll-like receptor 9 and activation by cpg oligonucleotides. *Journal of Immunological Methods*, 300, 2005.

27. Jordi Mestres, Elisabet Gregori-Puigjane, Sergi Valverde, and Ricard V Sole. Data completeness—the achilles heel of drug-target networks. *Nature biotechnology*, 26(9):983, 2008.

28. John P Overington, Bissan Al-Lazikani, and Andrew L Hopkins. How many drug targets are there? *Nature reviews Drug discovery*, 5(12):993, 2006.

29. Gaia V Paolini, Richard HB Shapland, Willem P van Hoorn, Jonathan S Mason, and Andrew L Hopkins. Global mapping of pharmacological space. *Nature biotechnology*, 24(7):805, 2006.

30. Yao Peng, John D McCorvy, Kasper Harpsøe, Katherine Lansu, Shuguang Yuan, Petr Popov, Lu Qu, Mengchen Pu, Tao Che, Louise F Nikolajsen, et al. 5-ht2c receptor structures reveal the structural basis of gpcr polypharmacology. *Cell*, 172(4):719–730, 2018.

31. Lingyan Ping, Ning Ding, Yunfei Shi, Lixia Feng, Jiao Li, Yalu Liu, Yufu Lin, Cunzhen Shi, Xing Wang, Zhengying Pan, et al. The bruton's tyrosine kinase inhibitor ibrutinib exerts immunomodulatory effects through regulation of tumor-infiltrating macrophages. *Oncotarget*, 8(24):39218, 2017.

32. Greg L. Plosker and David P. Figgitt. Rituximab. *Drugs*, 63(8):803–843, Apr 2003.

33. Sebastian Salentin, Sven Schreiber, V Joachim Haupt, Melissa F Adasme, and Michael Schroeder. Plip: fully automated protein–ligand interaction profiler. *Nucleic acids research*, 43(W1):W443–W447, 2015.

34. Patrick M Sexton and Arthur Christopoulos. To bind or not to bind: Unravelling gpcr polypharmacology. *Cell*, 172(4):636–638, 2018.

35. Masabumi Shibuya. Vascular endothelial growth factor and its receptor system: physiological functions in angiogenesis and pathological roles in various diseases. *The Journal of Biochemistry*, 153(1):13–19, 11 2012.

36. G. P. Spickett. Current perspectives on common variable immunodeficiency (cvid). *Clinical & Experimental Allergy*, 31(4):536–542, 2001.

37. S Suvas, V Singh, S Sahdev, H Vohra, and J N. Agrewala. Distinct role of cd80 and cd86 in the regulation of the activation of b cell and b cell lymphoma. *J. Biol. Chem.*, 277(10):7766–7775, 2002.

38. Kristien Van Belle, Jean Herman, Louis Boon, Mark Waer, Ben Sprangers, and Thierry Louat. Comparative in vitro immune stimulation analysis of primary human b cells and b cell lines. *Journal of immunology research*, 2016, 2016.

39. Anja K Weidemann, Ania A Crawshaw, Emily Byrne, and Helen S Young. Vascular endothelial growth factor inhibitors: investigational therapies for the treatment of psoriasis. *Clinical, cosmetic and investigational dermatology*, 6:233, 2013.