Title: Reproducible microbial community dynamics of two drinking water systems treating

similar source waters.

Sarah C Potgieter[a], Zihan Dai[b], Minette Havenga[a], Solize Vosloo[c] Makhosazana Sigudu[d] and

Ameet J Pinto[c,] Stefanus N Venter[a]*

Author Affiliations

[a] Rand Water Chair in Water Microbiology, Department of Biochemistry, Genetics and

Microbiology, University of Pretoria, South Africa

[b] School of Engineering, University of Glasgow, United Kingdom.

[c] Department of Civil and Environmental Engineering, Northeastern University, Boston, USA

[d] Scientific Services, Rand Water, Vereeniging, South Africa

Corresponding author:

Prof. Stephanus N Venter,

Department of Biochemistry, Genetics and Microbiology,

University of Pretoria,

Pretoria,

South Africa, 0002

Email: fanus.venter@up.ac.za

22   Graphical abstract

Abstract

Understanding whether the spatial-temporal dynamics of the drinking water microbiome are reproducible in full-scale drinking water systems is an important step towards devising engineering strategies to manipulate it. Yet, direct comparisons across full-scale drinking water systems are challenging because multiple factors, from source water to treatment process choice and configuration, can be unique to each system. This study compared the spatial-temporal dynamics of the drinking water microbiome in two drinking water treatment plants (DWTPs) with identical sequence of treatment strategies treating source waters from the same river system and with treated drinking water distributed in same large-scale (but independent) distribution system (DWDS) with similar disinfectant residual regiment. Dissimilarities in source water communities were tempered by the pre-disinfection treatments, resulting in highly similar post-filtration microbial communities between the two systems. However, high community turnover due to disinfection resulted in highly dissimilar microbial communities in the finished water between the two systems. Interestingly however, the microbial communities in the two systems increased in similarity during transit through the DWDS despite presence of a disinfectant residual. Overall our study finds that the drinking water microbiome demonstrated reproducible spatial and temporal dynamics within both independent but nearly identical DWTPs and their corresponding DWDSs.

Keywords: drinking water treatment; drinking water distribution; disinfection; microbial community dynamics; amplicon sequencing

1. Introduction

Drinking water treatment operations are designed to reduce microbial concentrations and limit microbial growth in drinking water distribution systems (DWDS) in order to deliver safe water to the consumer[1-3]. The choice of treatment strategies implemented to achieve this is highly site specific and is based on a range of considerations including source water type, local regulations, etc[3,4]. Drinking water treatment itself can be considered as a series of ecological disturbances implemented sequentially on the microbial community as it transits from the source water to the consumers taps[5]. These disturbances shape the microbial community composition (i.e., who is there)[6], as well as the population and community size (i.e., cell concentrations). Previous studies have shown that the drinking water microbiome is shaped by the choice of treatment strategy and conditions in the DWDS[3,5,7-10]. For instance, the microbial concentrations in finished drinking water can vary from 10^3 and 10^5 cells/mL[11-15], depending on the presence, absence, and concentrations of the disinfectant residual. Disinfectant residuals not only impact community size, but also impact community structure[3,10,16-18] and functional potential[18,19]. Similarly, filtration processes impact downstream microbial communities both by filtration-mediated seeding (Pinto *et al*., 2012) and through the biologically mediated removal of nutrients making them unavailable for microbial growth downstream[11,12,20,21].

Factors influencing the drinking water microbiome are site specific due to the unique type and combination of treatment technologies across DWTPs and DWDS configurations, source water types, and operational practices[21-23]. Thus, while the impact of individual treatment process or environmental conditions on the drinking water microbiome may be compared across DWTP's and DWDSs, it is often impossible to perform paired comparison between DWSs due to the inability to control for the type and sequential combination of treatment and distribution practices. For instance, studies have investigated drinking water

microbiome dynamics in two DWTPs that use similar treatment processes but utilize different

source waters or utilize the same source water but different treatment approaches[24-27]. Further,

often such comparisons are limited to either the treatment system or the distribution system,

but rarely encompass a comprehensive source-to-tap analysis. To our knowledge, no study has

yet investigated the impact of identical drinking water treatment regimens and similar

distribution practices on the drinking water microbiome in full-scale systems utilizing similar

source waters. This is a significant knowledge gap in the field, because it has direct

implications for our understanding of reproducibility of the drinking water microbiome

dynamics and the impact of engineering interventions.

This study presents unique insights from systematic comparisons between two

DWTPs, using the same treatment strategies to treat similar source waters originating from the

same river system and their corresponding distribution systems. The specific objectives of this

study were to investigate: (i) whether identical sequence of treatment technologies within two

DWTP's treating similar source waters results in similar changes in the microbial community,

(ii) the extent to which similarities in temporal dynamics between DWTP's are conserved (or

not) between their respective distribution systems, and (iii) and to what extent the impact of

physicochemical parameters on microbial community structure is conserved between the two

DWSs. To our knowledge this mirrored study has not been previously performed and is critical

towards improving our understanding of the spatial-temporal dynamics of the drinking water

microbiome.

2. Materials and methods

2.1. Site description. This is study involves two drinking water systems (System R and S) that

are operated by same drinking water utility (Fig. 1A). These two systems supply on an average

3653 million liters per day to approximately 12 million people within large metropolitan

region and local municipalities as well as for industrial use through a network of large diameter

pipelines stretching over 3056 km. The source water is drawn primarily from the Vaal river

and dam system into two drinking water treatment plants (R_DWTP and S_DWTP), which

abstract, treat, and distribute 98% (approximately 4320 ML/d) of the total water supplied by

the utility. The R_DWTP (river intake pumping site) treats source water from the river

downstream of the dam and the S_DWTP treats source water from a canal directly from the

dam. Treatment of the source waters in both DWTPs consists of the identical treatment steps

(Fig. 1B). Briefly, source water in both DWTPs is dosed with polyelectrolyte coagulants with

low lime for coagulation and flocculation, with no need for pH correction after sedimentation.

Although in some months in System R, a combination of polyelectrolyte and silica lime are

used for coagulation and flocculation (Table S1). In these instances, following sedimentation,

the water pH is adjusted to near neutral by bubbling CO_2 gas followed by filtration through

rapid gravity sand filters. Finally, the filter effluent is dosed with chlorine gas via bubbling as

the primary disinfection step. The total chlorine at sites following chlorination varies between

1.0 mg/L and 2.5 mg/L after 20 min contact time. Chlorinated water leaving both DWTPs is

again dosed with chloramine (approximately 2 mg/L) at secondary disinfection boosting

stations. For the purpose of this study chlorinated water originating from the R_DWTP was

monitored to the booster station, which supplies approximately 1,100 ML/d of chloraminated

water, serving predominately the northwest area of the distribution system (R_DWDS).

Chlorinated water originating from the S_DWTP was also monitored to another booster

station, supplying approximately 700 ML/d of chloraminated water to the eastern parts of the

distribution system (S_DWDS) (Fig. 1A). Samples were also collected after the booster

stations in the chloraminated sections of each DWDS, with monochloramine residuals varying

on average between 0.8 mg/L in the autumn and 1.5 mg/L in the spring. Further details on

range of physical-chemical parameters for both systems were obtained from the utility (Table

S1 and S2).

2.2. Sample collection and processing. Samples were collected for 8 months (February 2016

– September 2016) on a monthly basis from the two DWTPs (R_DWTP and S_DWTP) and

their associated DWDS (R_DWDS and S_DWDS) (Fig. 1A). At each DWTP, samples

collected included source water (SW), filter influent (FI, i.e. water entering the rapid sand filter

following coagulation, flocculation, sedimentation and carbonation), filter bed media (FB),

filter effluent (FE) and chlorinated water leaving the treatment plant (CHLA). In both DWDSs,

chlorinated water samples were collected immediately before the booster station (CHLB),

chloraminated water leaving the booster station (CHM) and chloraminated bulk water samples

at two points with the DWDSs (DS1 and DS2, respectively) (Fig. 1A and 1B). Within the two

DWTPs, 1 L of source water, 4 L of filter influent, and 8 L of filter effluent were collected.

Typically, 8 – 16 L of bulk water was collected for samples collected post disinfection and in

the DWDS. Collected water samples were filtered to harvest microbial cells followed by

phenol:chloroform DNA extraction as described by Potgieter *et al*., 2018. To obtain microbial

biomass from the filter bed media samples, 10 g of filter media was mixed with 50 ml

extraction buffer (i.e., 0.4 g/L EDTA, 1.2 g/L TRIS, 1 g/L peptone and 0.4 g/L N-dodecyl-N,

N dimethyl-3-amminio-1-propanesulfonate) followed by sonication for 1 min to remove the

microbial biomass attached to sand particles[28]. After sonication, the aqueous phase was filtered

through a Sterivex™-GP 0.22 μm polycarbonate membrane filter unit (Merck Milipore, South

Africa) followed by phenol:chloroform DNA extraction, as with bulk water samples.

2.3. Sequencing and data processing. Extracted DNA from samples were sent to the

Department of Microbiology and Immunology, University of Michigan Medical School (Ann

Arbor, USA) for the 2x250 bp sequencing of the V4 hypervariable region of 16S rRNA gene

using the Illumina MiSeq platform[29]. All raw sequence data have been deposited with links to

BioProject accession number PRJNA529765 on NCBI. The resultant 16S rRNA gene

amplicon sequences were processed using the Divisive Amplicon Denoising Algorithm,

DADA2 v1.14[30] workflow including sequence filtering, dereplication, inferring sample

composition, chimera identification and removal, merging of paired-end reads and

construction on a sequence table. Initial trimming and filtering of reads followed standard

filtering parameters described for Illumina MiSeq 2x250 V4 region of the 16S rRNA gene.

Specifically, reads with ambiguous bases were removed (maxN=0) were removed, the

maximum number of "expected errors" was defined (maxEE=2) and reads were truncated at

the first instance of a quality score less than or equal to truncQ (truncQ=2). Dereplication was

performed where identical sequences are combined into "unique sequences" while

maintaining the corresponding abundance of the number of reads for that unique sequence.

The core sample inference algorithm was applied to dereplicated data and forward and reverse

reads were merged together to obtain fully denoised sequences[30]. Merged reads were then used

to construct an amplicon sequence variant (ASV) table[31], chimeras were identified and

removed and taxonomic assignments were called using the SILVA reference database

(https://www.arb-silva.de) through the DADA2 chimera removal and taxonomy assignment

script.

2.4. Microbial community analysis. Resulting ASV table was imported into the mothur (v

1.35.1)[32] and the shared sequences between sample locations from the two DWTPs and

corresponding DWDSs as well as the unique sequences within each sample location were

calculated using the venn function in mothur. Alpha diversity measures (i.e., richness,

Shannon Diversity Index and Pielou's evenness) were calculated using the summary.single

function in mothur with the parameters, subsampling=1263 (sample with the least amount of

sequences) and iters=1000 (1000 subsampling of the entire dataset). Due to subsampling, 10

samples were excluded from the analyses and Good's coverage estimates were calculated to

assess whether sufficient number of sequences were retained for each sample after

subsampling. This indicated that subsampling at a library size of 1263 retained the majority of

the richness for all samples (i.e., average Good's coverage = 95.84 ± 0.02%). One-way

analysis of variance (ANOVA)[33] and post-hoc Tukey Honest Significant Differences (HSD)

test were performed in R (http://www.R-project.org) using the stats package[34] to determine

the statistical significance between spatial and temporal groupings within the alpha diversity.

Bray-Curtis and weighted UniFrac were used to determine pair-wise dissimilarity in

community structure between samples, whereas Jaccard and unweighted UniFrac were used

to infer dissimilarity in community membership. Bray-Curtis and Jaccard distances were

calculated using the dist.shared function in mothur with the parameters, subsampling=1263

and iters=1000. Weighted and unweighted UniFrac distances were calculated through the

construction of a phylogenetic tree with representative sequences using the clearcut command

in mothur also with the parameters subsampling=1263 and iters=1000[35,36]. Pairwise Analysis

of Molecular Variance (AMOVA) was performed using the amova function in mothur on all

beta diversity matrices, to determine the effect of sample groupings based on DWDS sample

location, DWDS section, and season[37,38]. Beta diversity metrics and metadata files containing

sample location, sample type, disinfection type and season were imported into R

(http://www.R-project.org) for statistical analysis and visualization. Principal-coordinate

analyses (PCoA) using Bray-Curtis and Jaccard distances was performed using the phyloseq

package[39]. All plots were constructed using the ggplot2 package[40]. To identify the contribution

of environmental parameters and their combinations towards microbial community structure,

distance-based redundancy analysis (dbRDA) was used. The function dbrda() from R package

"vegan"[41] was applied on Bray-Curtis distances estimated between samples using ASV counts

to investigate relationships between the scaled environmental parameters and microbial

community structure. In addition, the faction of variation explained by the environmental

parameters identified as significantly associated with ASV count-based Bray-Curtis distance

matrices was determined using the function varpart() in the "vegan" package.

3. Results and discussion

A total of 172 samples were sequenced for this study (Table S3) which resulted in 4,921,399

sequences post quality control and a total of 10,012 ASVs. Taxonomic classification of these

ASV's revealed that bacteria dominated the microbial community (mean relative abundance,

MRA: $98.74 \pm 0.02\%$ across all samples) followed by archaea (MRA: $1.04 \pm 0.01\%$). The

bacterial community was primarily composed of *Gammaproteobacteria* (including

Betaproteobacteriales), *Actinobacteria*, and *Planctomycetes* (Fig. S1 and Table S4).

3.1. Treatment processes increased similarities between microbial communities across

the two drinking water systems. Despite being drawn from the same surface water system,

the two-source water microbial communities were significantly different (AMOVA, $F_{ST} \leq$

3.04, $p < 0.001$, depending on the beta diversity measure) and only shared 22.5% of the

detected ASV's (Fig. S2). These differences could likely be a result of varying hydrological

conditions at the two locations (i.e., river drawn R_SW vs S_SW from the dam), which may

translate to the variable occurrence of low abundance and transient taxa[42]. Despite harboring

more ASV's over the period of the study (Fig. S2), R_SW was less diverse than S_SW

(ANOVA; $p < 0.05$) per timepoint (Fig. S3; Table S5). This was because R_SW was likely

much more influenced by strong hydrological conditions such as runoff and increased flow

rates during heavy rainfall events[43,44] as compared to S_SW. This is supported by the fact that

R_SW exhibited higher temporal variability as compared to S_SW based on pairwise beta

diversity measures (Fig. S4).

227   The pre-chlorination treatment processes played an important role in tempering the

differences in microbial communities across the two DWTPs. The microbial community

composition of both DWTPs was highly diverse and in addition to *Proteobacteria* included

other dominant phyla (i.e. MRA greater than 1%) such as *Acidobacteria, Cyanobacteria,*

Verrucomicrobia and *Planctomycetes* (Fig. S1; Table S4). The 33 most abundant ASV's i.e.,

ASV's with an MRA of > 0.5%, showed similar trends between the two DWTPs in terms of

increase/decrease in MRA with each sequential treatment step (Fig. 2; Table S6). These

dominant ASV's within both DWTP included members of *Actinobacteria, Acidobacteria,*

Gammaproteobacteria, Bacteroidetes and *Thaumarcheota.* This reproducible effect of

sequential treatment steps was also evident at the community level. Specifically, the treatment

process resulted in similar changes in alpha diversity measures in both DWTPs (Fig. S3).

Based on ANOVA these changes were found to be significant (richness: $F_{ST} = 19.67, p < 0.05,$

Shannon Diversity Index: $F_{ST} = 9.78$, p < 0.05, Inverse Simpson Diversity Index: $F_{ST} = 15.64,$

p < 0.05 and Pielou's evenness: $F_{ST} = 4.79$, p < 0.05). Richness and diversity consistently

decreased along treatment processes (excluding filter bed samples (FB)), with the most

significant decrease immediately following chlorination (Fig. S3; Table S5). This shows that

the decrease in microbial abundance and diversity, typical during treatment processes[1,5,45-47],

was reproducible in both systems.

245   Beta-diversity analyses indicated that microbial communities became increasingly

similar from source water through treatment and filtration where the microbial communities

between the filter bed and the filter effluent were approximately 40% similar in community

structure (Bray-Curtis: 0.57 ± 0.15) and 30% similar in community membership (Jaccard: 0.71

± 0.11); this was comparable to the similarity of the filter bed and filter influent communities

(community structure: Bray-Curtis: 0.55 ± 0.17, community membership: Jaccard: 0.7 ± 0.13)

(Fig. 3A). Similar trends in community structure and membership were observed in weighted

and unweighted Unifrac analyses (Fig. S5). Following similar prefiltration treatments of the

source water, filter influent samples between the two systems showed the greatest similarity

(Fig. 3B). Although, filter bed microbial communities (R_FB and S_FB) showed higher

dissimilarity in community structure and membership (Fig. 3B; Table S7) between the two

DWTPs, they were also significantly different from both the filter inflow (AMOVA: $F_{ST} \le$

5.07, p < 0.001) and filter effluent (AMOVA: $F_{ST} \le 6.51$, p < 0.001). The greater dissimilarity

between the FB from the two DWTP's could also be attributed to inherent high heterogeneity

of attached growth microbial communities in the filter media[7,23]. Yet, microbial communities

were more similar between filter bed samples of the two DWTPs, than the source waters that

feed them. Further, an increase in the number of shared ASV's between the two filter effluents

(36.04%), indicates that conditions in the filter beds were sufficiently similar to have the same

effect on the resulting effluent and the selection of dominant taxa in both systems.

3.2. Rapid microbial community turnover due to disinfection increases dissimilarity

across the two drinking water systems. In contrast to the DWTP (i.e., pre-chlorination)

where treatment processes enhance similarity in microbial community between the two

DWTPs, the rapid change in microbial community due to chlorination resulted in significant

increase in dissimilarity between locations within the DWTP/DWDS and across the two

systems. Following chlorination, the microbial communities between filter effluent (FE) and

bulk water (CHLA) were only $8 - 11\%$ similar in community structure (Bray-Curtis: $0.89 \pm$

0.2) and membership (Jaccard: 0.92 ± 0.15) (AMOVA: $F_{ST} \le 18.22$, p < 0.001 depending on

the beta diversity measure) (Fig. 3A). This significant decrease in similarity was also observed

in weighted and unweighted Unifrac analyses (Fig. S5). In addition, this significant change

following disinfection was also observed as shift in the relative abundance of dominant ASVs

(Fig. 2).

Further, chlorinated samples CHLA and CHLB increased slightly in similarity in

community structure and membership (Bray-Curtis: 0.77 ± 0.16 and Jaccard: 0.86 ± 0.11) but

remained highly dissimilar, though not significantly different. However, following secondary

disinfection, the microbial communities between chlorinated (CHLB) and chloraminated

water (CHM) again increased in dissimilarity in community structure (Bray-Curtis: 0.84 ±

0.12) and membership (Jaccard: 0.91 ± 0.08) (AMOVA: $F_{ST} \leq 4.09$, p < 0.001 depending on

the beta diversity measure) in both systems (Fig. 3A) (same trends observed for Unifrac

analyses, Fig. S5). This was consistent for both, the R and S systems. Beta diversity

comparisons between paired samples from System R and S immediately after chlorination

(R_CHLA and S_CHLA) showed an increase in dissimilarity in both community structure and

membership (Fig. 3B; Table S7). This was also observed between R_CHLB and S_CHLB

samples. Samples within the DWDS (CHM, DS1 and DS2) showed consistent temporal trends

where samples from both systems increased in dissimilarity 6 months apart, consistent with

the observed changes in temperature (Fig. S4; Fig. S6). However, temporal variability

remained high within DS1 and DS2 samples where pair-wise comparisons between

consecutive months within each sample location were dissimilar in community membership

(i.e., Jaccard: 0.80 ± 0.06 and unweighted UniFrac: 0.67 ± 0.06) and structure (i.e., Bray-

Curtis: 0.69 ± 0.13 and weighted UniFrac: 0.52 ± 0.13) (Fig. 3B).

Disinfection significantly reduces bacterial cell concentrations and has a substantial

influence on community composition and structure [3,10,24,45,48,49]. While microbial communities

prior to chlorination were similar between the two DWTPs, the microbial community

composition and structure in both systems were highly dissimilar post-chlorination. This

dissimilarity was also observed on a temporal scale, where samples post disinfection showed

greater temporal variability as compared to the pre-chlorinated samples. This variability could

potentially arise from a few different factors. For instance, chlorine and chloramine are both

strong oxidants and are likely to inactivate microorganisms indiscriminately; this could be one

potential reason for higher temporal variability post-disinfection and greater dissimilarity

between paired samples between the two drinking water systems. Alternatively, those

surviving disinfection may be associated with biofilms. The presence of disinfectant residual

has been reported to have limited impact in preventing biofilm development and in some cases

may even promote biofilm formation as a stress response[50,51]. Another explanation could be

reduction in microbial abundance and diversity due to disinfection may lead to the detection

of low abundance or rare taxa that may not have well defined niche within the drinking water

community and thus are variably present and/or detected.

Ultimately, the application of disinfection can be viewed as an ecological disturbance,

where the sequential and controlled addition of disinfectants, disturb the drinking water

microbial continuum[5]. The diversity of microbial communities generally decreases in response

to environmental stress and disturbances, ultimately shifting the ecological balance of

microbial populations within the community [6,52,53]. Following a disturbance, the surviving

populations are considered to have specific properties (such as biofilm formation or oxidative

stress response mechanisms), allowing them to persist in the disrupted environment.

3.3. Chloramine residuals and distribution conditions promote microbial community

stability. Considering the aforementioned similar impacts of treatment process and

distribution system on the microbial communities between the two systems, it is not surprising

that microbial communities from both systems clustered together depending on sampling

location (i.e., DWTP or DWDS), rather than system (R or S) (Fig. 4). For instance, the DWTP

locations clustered independently from the DWDS across both systems (Fig. 4A). Even within

the DWTP, microbial communities clustered depending on sampling location despite the

temporal variability (Fig. 4B). The same was consistent for the DWDS, where the chlorinated

and chloraminated sections of the DWDS clustered together (Fig. 4C). Further, the measured

water quality parameters were highly similar at each location between the two DWTPs and

DWDS (Fig. S7; Table S2). To assess the extent to which measured water quality parameters

impacted the microbial community, we performed dbRDA analysis by focusing primarily on

the post-chlorination samples (i.e., chlorinated and chloraminated DWDS). This was done

because each section of the DWDS across the two DWS's has multiple samples per month

over the eight-month sampling period resulting in significant sample size for both dbRDA and

subsequent variance partitioning tests. Consistent with PCoA analyses, dbRDA revealed that

samples clustered based on disinfectant residual type (chlorinated or chloraminated) and not

based on the system they originated from. Based on dbRDA analysis, ammonium, water

temperature, monochloramine and free Cl_2 emerged as the best indicators among water quality

parameters that explained the variability in microbial community structure in both systems

(ANOVA: $p < 0.001$) (Fig. 5; Table S8). Increased concentrations of free chlorine (1.66 ± 0.55

mg/L) was identified as a significant variable in chlorinated samples ($p = 0.0021$).

Alternatively, among other water quality parameters, ammonium (0.36 ± 0.13 mg/L as

N) and monochloramine concentrations (1.60 ± 0.38 mg/L) as well as temperature ($18.28 \pm$

4.47 °C) were identified as significant variables in chloraminated samples ($p < 0.001$).

Variance partitioning analyses were used to determine the proportion of variance that could

be attributed to variables (individual or combined) identified as most significant in the dbRDA

analysis (Fig. 5). This resulted in ammonium, free Cl_2, monochloramine and temperature each

explaining 2.5%, 1.9%, 1.6% and 1.3% of the variance, respectively. While the contribution

of these parameters towards explaining the variance in community structure was significant,

it was small. In fact, 90% of the variance could not be explained by these four parameters (Fig.

S8; Table S9). Here, ammonium and monochloramine concentrations and temperature best

explained the variability in chloraminated samples from both distribution systems as

disinfectant residual concentration and water temperatures did not differ greatly between the

two lines. This is well supported as these variables are long considered to be important factors

in shaping the drinking water microbiome[3]. Therefore, the observed dissimilarity between the

two distribution lines may be accredited to the initial differential response of the microbial

community to chlorine residuals.

Interestingly, the changes in the microbial community in the chlorinated sections were

driven predominately by chlorine residuals (specifically free Cl_2) as an initial significant

ecological disturbance. However, chlorine residuals are typically short-lived and therefore

chloramine residuals are added as a more stable alternative to ensure residuals are maintained

over longer distances[54,55]. Nevertheless, the transport of bulk water over longer distances leads

to increased water age and decreased disinfectant residual due to disinfectant decay, ultimately

contributing to microbial regrowth[50]. In addition to disinfectant residuals, the microbial

community is also exposed to various other distribution conditions and water quality

parameters including (but not limited to) pipe material, hydraulic conditions, residence time,

water temperature, concentrations of nitrogen species, organic carbon, etc. All these factors

have a considerable impact on the drinking water microbiome[3].

An increase in similarity was observed in locations towards the end of the DWDS

(DS2 samples). Through distribution, water is continually seeded by similar microbial

communities over time thereby selecting for the same dominant taxa through similarities in

pipe material, residence times, hydraulic conditions and operation practices contributing to

site specific taxa and biofilms. Biofilms have also been shown to seed bulk water communities

through biofilm erosion, detachment and sloughing[56]. In this study, the established (historical)

biofilms in both systems have been long exposed to similar distribution conditions thereby

potentially contributing to the development of similar biofilm microbial communities. This

increase in similarity in community membership and structure with increasing residence time

in the DWDS was more pronounced in samples from summer and autumn. Here, elevated

water temperatures in summer months may affect the bacterial community composition and

structure by positively influencing the growth kinetics and competition processes of specific

bacterial species in each section of the DWDS[3]. Alternatively, seasonal increase in water

demand could increase shear stress resulting in increased detachment of similar established

biofilms[56].

In addition, where chlorine may be non-specific in its action in reducing bacterial cell

concentrations, chloramine (with the addition of ammonia) may be more selective towards

microbial taxa capable of using ammonia as an energy source. This may explain the observed

increase in *Nitrosomonas* sp. (ASV_6) in chloraminated distribution system samples from

both systems. Here, the addition of chloramine as a secondary disinfectant has been shown to

support the growth of nitrifying bacteria in DWDS[57]. The long residence time and associated

lower disinfectant residual concentrations, together with the release of ammonia through

disinfection decay results in increased abundance of nitrifiers and therefore potential

nitrification[49]. Thus, decreasing monochloramine residual concentrations associated with

increased water age at nearing the end of both DWDSs may also contribute to the growth of

similar bacterial assemblages at these sites.

This study provided a unique opportunity to compare the effects of the same treatment

strategies (disturbances) on similar source waters as well as the distribution of treated water

on the drinking water microbiome in a large-scale system. The spatial-temporal dynamics of

two independent DWTP and their corresponding DWDSs with in the same large-scale

distribution system were shown to be reproducible. Similarities in design and operational

parameters of the two DWTPs resulted in the development of similar pre-disinfection

microbial communities. However, the differential impact of chlorination was highlighted in post-disinfection samples from the two systems, resulting in dissimilar microbial communities between the two systems. Lastly, distribution was observed to select for certain dominant taxa, thereby increasing similarity between microbial communities due to the impact of inherent biofilms seeding the distribution system. However, dissimilarities in microbial community throughout distribution may arise from initial differences in the source waters and the differential response to chlorination ultimately leading to the presence of site specific rare/low abundant taxa. This study confirms the role of treatment and distribution in shaping the microbial community and may provide utilities with the reassurance that under that same conditions, drinking water quality will be reproducible throughout the varying stages of the drinking water system.

Supporting Information

Table S1: Water quality parameters of the source water from both systems

Table S2: Water quality parameters for DWDS sites

Table S3: The number of samples collected and processed

Table S4: Mean relative abundance of dominant bacterial phyla

Table S5: Mean and standard deviations of alpha diversity indexes

Table S6: Mean percentage relative abundance of the top 33 abundant sequence variants

Table S7: Pair-wise beta diversity comparisons

Table S8: Permutation test for distance-based redundancy analysis (dbRDA)

Table S9: Variance Partition analyses

Fig. S1: Phylum-level and proteobacterial classes of bacterial sequences detected

Fig. S2: Venn diagram of shared amplicon sequence variants (ASVs) between source waters.

Fig. S3: Spatial changes in alpha diversity indexes

Acknowledgements

This research was funded and supported by Rand Water, Gauteng, South Africa through the

Rand water Chair in Water Microbiology at the University of Pretoria. Sarah Potgieter would

also like to acknowledge the National Research Foundation (NRF) for additional funding.

Ameet J. Pinto was supported by NSF Award 1749530. Furthermore, the authors would like

to acknowledge the Centre for Microbial Systems Molecular Biology Lab, University of

Michigan, USA for their services in Illumina MiSeq sequencing.

References

1. Hammes, F., Berney, M., Wang, Y., Vital, M., Koster, O. and Egli, T. (2008). Flow-

cytometric total bacterial cell counts as a descriptive microbiological parameter for drinking

water treatment processes. *Water Research*. 44(17): 4868-4877.

2. Lautenschlager, K., Hwang, C., Ling, F., Lui, W. T., Boon, N., Köster, O., Egli, T. and

Hammes, F. (2014). Abundance and composition of indigenous bacterial communities in a

multi-step biofiltration-based drinking water treatment. *Water Research.* 62: 40-52.

3. Prest, E. I., Hammes, F., van Loosdrecht, M. C. M. and Vrouwenvelder, J. S. (2016).

Biological stability of drinking water: controlling factors, methods and challenges.

Frontiers in Microbiology. 7(45): doi: 10.3389/fmicb.2016.00045.

4. Ho, L., Braun, K., Fabris, R., Hoefel, D., Morran, J., Monis, P. and Drikas, M., 2012.

Comparison of drinking water treatment process streams for optimal bacteriological water

quality. *Water research*, *46*(12), pp.3934-3942.

5. Zhang, Y., Oh, S. and Liu, W. T. (2017). Impact of drinking water treatment and

distribution on the microbiome continuum: an ecological disturbance's perspective.

Environmental Microbiology. 19(8): 3163-3174.

6. Gomez-Alvarez, V., Pfaller, S., Pressman, J.G., Wahman, D.G. and Revetta, R.P., (2016).

Resilience of microbial communities in a simulated drinking water distribution system

subjected to disturbances: role of conditionally rare taxa and potential implications for

antibiotic-resistant bacteria. *Environmental Science: Water Research and Technology*.

2(4): 645-657.

7. Pinto, A. J., Xi, C. and Raskin, L. (2012). Bacterial community structure in the drinking

water microbiome is governed by filtration processes. *Environmental Science and*

Technology. 46: 8851-8859.

8. Pinto, A., Schroeder, J., Lunn, M., Sloan, W. and Raskin, L. (2014). Spatial-temporal

survey and occupancy-abundance modelling to predict bacterial community dynamics in

the drinking water microbiome. *mBIO*. 5(3): e01135-14.

9. Bautista-de los Santos, Q. M., Schroeder, M. C., Sevillano-Rivera, M. C., Sungthong, R.,

Ijaz, U. Z., Sloan, W. T. and Pinto, A. J. (2016). Emerging investigators series: microbial

communities in full-scale drinking water distribution systems – a meta-analysis.

Environmental Science: Water Research and Technology. doi: 10.1039/c6ew00030d.

10. Potgieter, S., Pinto, A., Sigudu, M., Du Preez, H., Ncube, E. and Venter, S. (2018). Long-

term spatial and temporal microbial community dynamics in a large-scale drinking water

distribution system with multiple disinfectant regimes. *Water Research*. 139: 406-419.

11. Hammes, F., Berger, C., Koster, O. and Egli, T. (2010). Assessing biological stability of

drinking water without disinfectant residuals in a full-scale water supply system. *Journal*

of Water Supply: Research and Technology-AQUA. 59: 31-40.

12. Lautenschlager, K., Hwang, C., Ling, F., Liu, W. T., Boon, N., Köster, O., Vrouwenvelder,

H., Egli, T. and Hammes, F. (2013). A microbiology-based multi-parametric approach

towards assessing biological stability in drinking water distribution networks. *Water*

Research. 47: 3015-3025.

13. Liu, G., Verbeck, J. Q. J. C. and Van Dijk, J. C. (2013). Bacteriology of drinking water

distribution systems: an integral and multidimensional review. *Applied and Environmental*

Microbiology. 97: 9265-9276.

14. Gillespie, S., Lipphaus, P., Green, J., Parsons, S., Weir, P., Juskowiak, K., Jefferson, B.,

Jarvis, P. and Nocker, A. (2014). Assessing microbiological water quality in drinking water

distribution systems with disinfectant residual using flow cytometry. *Water Research.* 65:

224-234.

15. Nescerecka, A., Rubulis, J., Vital, M., Juhna, T and Hammes, F. (2014). Biological

instability in a chlorinated drinking water distribution network. *PloS One.* 9: e96354.

16. Gomez-Alvarez, V., Revetta, R. P. and Santo Domingo, J. W. (2012). Metagenomic

analysis of drinking water receiving different disinfection treatments. *Appied and*

Enivronmental Microbiology. 78(17): 6095-6102.

17. Hwang, C., Ling, F., Andersen, G. L., LeChevallier, M. W. and Liu, W. (2012). Microbial

community dynamics of an urban drinking water distribution system subjected to phases of

chloramination and chlorination treatments. *Applied and Environmental Microbiology.*

78(22): 7856-7865.

18. Dai, Z., Sevillano-Rivera, M.C., Calus, S.T., Bautista-de los Santos, Q.M., Eren, A.M., Van

Der Wielen, P.W., Ijaz, U.Z. and Pinto, A.J., 2020. Disinfection exhibits systematic impacts

on the drinking water microbiome. *Microbiome*, *8*(1), pp.1-19.

19. Sevillano-Rivera, M., Dai, Z., Calus, S., Bautista-de Los Santos, Q.M., Eren, A.M., van der

Wielen, P.W., Ijaz, U.Z. and Pinto, A.J., 2020. Differential prevalence and host-association

of antimicrobial resistance traits in disinfected and non-disinfected drinking water

systems. *Science of the Total Environment*, *749*, p.141451.

20. Kirisits, M.J., Emelko, M.B. and Pinto, A.J., 2019. Applying biotechnology for drinking

water biofiltration: advancing science and practice. *Current opinion in biotechnology*, *57*,

pp.197-204.

21. Ma, B., LaPara, T.M., N. Evans, A. and Hozalski, R.M., 2020. Effects of geographic

location and water quality on bacterial communities in full-scale biofilters across North

America. *FEMS microbiology ecology*, *96*(2), p. fiz210.

22. Roeselers, G., Coolen, J., van der Wielen, P. W. J. J., Jaspers, M. C., Atsma, A., deGraaf,

B. and Schuren, F. (2015). Microbial biogeography of drinking water: patterns in

phylogenetic diversity across space and time. *Environmental Microbiology*. 17(7): 2505-

2514.

23. Gülay, A., Musovic, S., Albrechtsen, H. J., Al-Soud, W. A., Sørensen, S. J. and Smets, B.

F. (2016). Ecological patterns, diversity and core taxa of microbial communities in

groundwater-fed rapid gravity filters. *The ISME journal*. 10(9): 2209.

24. Poitelon, J. B., Joyeux, M., Welté, B., Duguet, J. P., Prestel, E. and DuBow, M. S. (2010).

Variations of bacterial 16S rDNA phylotypes prior to and after chlorination for drinking

water production from two surface water treatment plants. *Journal of Industrial*

Microbiology and Biotechnology. 37(2): 117-128.

25. Xu, J., Tang, W., Ma, J. and Wang, H., 2017. Comparison of microbial community shifts

in two parallel multi-step drinking water treatment processes. *Applied Microbiology and*

Biotechnology, *101*(13), pp.5531-5541.

26. Nescerecka, A., Juhna, T. and Hammes, F. (2018). Identifying the underlying causes of

biological instability in a full-scale drinking water supply system. *Water Research*. 135:

11-21.

27. Waak, M.B., Hozalski, R.M., Hallé, C. and LaPara, T.M., 2019. Comparison of the

microbiomes of two drinking water distribution systems—with and without residual

chloramine disinfection. *Microbiome*, *7*(1), pp.1-14.

28. Camper, A. K., Lechevallier, M. W., Broadaway, S. C. and McFeters, G. A. (1985).

Evaluation of procedures to desorb bacteria from granular activated carbon. *Journal of*

Microbiological Method. 3: 187-198.

29. Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., Schloss, P. D. (2013).

Development of a dual-index strategy and curation pipeline for analyzing amplicon-

sequencing data on the MiSeq Illumina sequencing platform. *Applied and Environmental*

Microbiology. 79: 5112e5120.

30. Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A. and Holmes,

S. P. (2016). DADA2: high-resolution sample inference from Illumina amplicon data.

Nature Methods. 13(7): 581.

31. Callahan, B. J., McMurdie, P. J. and Holmes, S. P. (2017). Exact sequence variants should

replace operational taxonomic units in marker-gene data analysis. *The ISME journal*.

11(12): 2639.

32. Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B.,

Lesniewski, R. A., Oakley, B. B., Parks, D. H., Robinson, C. J., Sahl, J. W. (2009).

Introducing mothur: open-source, platform-independent, community-supported software

for describing and comparing microbial communities. *Applied and Environmental*

Microbiology. 75(23): 7537e7541.

33. Chambers, J. M., Freeny, A. and Heiberger, R. M. (1992). *Analysis of variance; designed*

experiments. Chapter 5 of *Statistical Models in S* eds J. M. Chambers and T. J. Hastie,

Wadsworth and Brooks/Cole.

34. R Core Team (2015). R: a language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

35. Evans, J., Sheneman, L. and Foster, J.A. (2006). Relaxed neighbour-joining: a fast distance-

based phylogenetic tree construction method. *Journal of Molecular Evoltion*. 62: 785e792.

36. Lozupone, C., Lladser, M. E., Knights, D., Stombaugh, J., Knight, R. (2011). UniFrac: an

effective distance metric for microbial community comparison. *The ISME Journal*.

5(2):169e172.

37. Excoffier, L., (1993). Analysis of Molecular Variance (AMOVA) Version 1.55. Genetics

and Biometry Laboratory, University of Geneva, Switzerland.

38. Anderson, M. J., (2001). A new method for non-parametric multivariate analysis of

variance. *Australian Ecology*. 26 (1): 32e46.

39. McMurdie, P.J. and Holmes, S. (2013). Phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. *PLoS One.* 8(4): e61217.

40. Wickham, H. (2009). ggplot2: Elegant graphics for data analysis. Springer-Verlag, New

York. http://ggplot2.org.

41. Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,

P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H.,

2019. Vegan: community ecology package.

42. Shade, A., Jones, S. E., Caporaso, J. G., Handelsman, J., Knight, R., Fierer, N., Gilbert, J.

A. (2014). Conditionally rare taxa disproportionately contribute to temporal changes in

microbial diversity. *mBio ASM*. 5(4): 1e9.

43. Prathumratana, L., Sthiannopkao, S. and Kim, K.W. (2008). The relationship of climatic

and hydrological parameters to surface water quality in the lower Mekong River.

Environment International. 34(6): 860-866.

44. Delpla, I., Jung, A.V., Baures, E., Clement, M. and Thomas, O. (2009). Impacts of climate

change on surface water quality in relation to drinking water production. *Environment*

International. 35(8): 1225-1233.

45. Lin, W., Yu, Z., Zhang, H. and Thompson, I. P. (2014). Diversity and dynamics of

microbial communities at each step of treatment plant for potable water generation. *Water*

Research. 52: 218-230.

46. Prest, E. I., El-Chakhtoura, J., Hammes, F., Saikaly, P.E., Van Loosdrecht, M.C.M. and

Vrouwenvelder, J. S. (2014). Combining flow cytometry and 16S rRNA gene

pyrosequencing: a promising approach for drinking water monitoring and characterization.

Water Research. 63: 179-189.

47. Wang, H., Proctor, C. R., Edwards, M. A., Pryor, M., Santo Domingo, J. W., Ryu, H., et al.

(2014). Microbial community response to chlorine conversion in a chloraminated drinking

water distribution system. *Environmental Science and Technology*. 48: 10624–10633. doi:

10.1021/es502646d.

48. Eichler, S., Christen, R., Höltje, C., Westphal, P., Bötel, J., Brettar, I., Mehling, A. and

Höfle, M. G. (2006). Composition and dynamics of bacterial communities of a drinking

water supply system as assessed by RNA-and DNA-based 16S rRNA gene fingerprinting.

Applied and Environmental Microbiology. 72(3): 1858-1872.

49. Wang, H., Masters, S., Edwards, M. A., Falkinham, J. O., and Pruden, A. (2014). Effect of

disinfectant, water age, and pipe materials on bacterial and eukaryotic community structure

in drinking water biofilm. *Environmental Science and Technology*. 48: 1426–1435. doi:

10.1021/es402636u

50. Fish, K. and Boxall, J. (2018). Biofilm Microbiome (Re) Growth Dynamics in Drinking

Water Distribution Systems Are Impacted by Chlorine Concentration. *Frontiers in*

Microbiology. 9: 2519.

51. Fish, K.E., Reeves-McLaren, N., Husband, S. and Boxall, J., 2020. Unchartered waters: the

unintended impacts of residual chlorine on water quality and biofilms. *npj Biofilms and*

Microbiomes, *6*(1), pp.1-12.

52. Atlas, R.M., Horowitz, A., Krichevsky, M. and Bej, A.K., 1991. Response of microbial

populations to environmental disturbance. *Microbial Ecology*, *22*(1), pp.249-256.

53. Shade, A., Peter, H., Allison, S.D., Baho, D., Berga, M., Bürgmann, H., Huber, D.H.,

Langenheder, S., Lennon, J.T., Martiny, J.B. and Matulich, K.L., 2012. Fundamentals of

microbial community resistance and resilience. *Frontiers in microbiology*, *3*, p.417.

54. Neden, D.G., Jones, R.J., Smith, J.R., Kirmeyer, G.J. and Foust, G.W., 1992. Comparing

chlorination and chloramination for controlling bacterial regrowth. *Journal‑American*

Water Works Association, *84*(7), pp.80-88.

55. Norton, C. D and LeChevallier, M. W. (1997). Chloramination: its effect on distribution

system water quality. *J. AWWA*. 89(7): 66-77.

56. Fish, K.E., Osborn, A.M. and Boxall, J., 2016. Characterising and understanding the impact

of microbial biofilms and the extracellular polymeric substance (EPS) matrix in drinking

water distribution systems. *Environmental science: water research & technology*, *2*(4),

pp.614-630.

57. Potgieter, S.C., Dai, Z., Venter, S.N., Sigudu, M. and Pinto, A.J., 2020. Microbial Nitrogen

Metabolism in Chloraminated Drinking Water Reservoirs. *mSphere*, *5*(2).

FIGURES

Fig. 1: (A) Site map of the location of the drinking water treatment plants (R_DWTP and S_DWTP) and their corresponding distribution systems (R_DWDS and S_DWDS). System R is indicated in red and System S in blue. The two treatment plants are represented as squares, the two-secondary disinfection boosting stations, where chloramine is added, are represented as triangles and all sample locations are represented as circles. (B) Schematic of the layout of the DWTP and DWDS showing all sample locations sampled monthly for the duration of the study. Within the two DWTPs source water (SW), filter inflow (FI), filter bed media (FB) and filter effluent (FE) samples were collected. All other sample locations are indicated on the figure and described in the text.

Fig. 2: Maximum-likelihood phylogenetic tree (left) showing the groupings of the top ASV with a relative abundance > 0.5 (33 ASVs).

Phylogenetic tree was constructed based on representative ASV DNA sequences with a bootstrap analysis of 1000 replicates (bootstrap values

indicated as percentages). The log percentage of average relative abundance of those top ASVs for each sample site in each system are shown in

heatmaps (right) followed by their taxonomic association. The average relative abundance for each sample location was averaged over duration

of the study for each system. The log percentage relative abundance of each ASV is indicated in the legends on the left of the figure. See Table

for mean relative abundances (MRA) of top dominant ASVs.

Fig. 3: (A) Average pairwise beta diversity comparisons of both structure-based metric: Bray-Curtis and membership-based metric: Jaccard.

Comparisons are between consecutive locations within each of the two systems for corresponding months. Sample abbreviations on the x-axis

refer to comparisons of samples following the flow of bulk water through both systems, i.e. source water and filter inflow (SW_FI), filter inflow

and filter bed media (FI_FB), filter bed media and filter effluent (FB_FE), filter effluent and chlorinated water leaving the DWTP (FE_CHLA),

chlorinated water leaving the DWTP and chlorinated water entering the secondary disinfection boosting station (CHLA_CHLB), chlorinated

water entering the secondary disinfection boosting station and chloraminated water (CHLB_CHM), chloraminated water and distribution system

site 1 (CHM_DS1) and finally distribution system site 1 and distribution system site 2 (DS1_DS2). Sample comparisons from System R

indicated in red and System S in blue. (B) Direct pairwise beta diversity comparisons (Bray-Curtis and Jaccard) between corresponding samples

from the two systems. Pairwise beta diversity comparisons include samples from the same month. Sample abbreviations on the x-axis refer to

source waters (SW), filter inflows (FI), filter bed medias (FB), filter effluents (FE), chlorinated waters leaving the DWTP (CHLA), chlorinated

waters entering the secondary disinfection boosting station (CHLB), chloraminated waters (CHM), distribution system sites 1 (DS1) and

distribution system sites 2 (DS2). Mean and standard deviations of each comparison is shown in Table S7. See Fig. S4 for corresponding

weighted and unweighted Unifrac plots

659   **Fig. 4**: Principal coordinate analysis plot (based on Bray-Curtis dissimilarity) showing the spatial and temporal variability of the bacterial

660   community structure among all samples from both systems (A), within the two DWTPs (B) and within the two corresponding DWDSs (C).

661   Spatial groupings are shown where data points are colored based on sample location and shaped based on the system they originate from

662   (System R samples are indicated as circles and System S samples as triangles). Color and shapes are indicated in the legends on the right of all

663   plots.

666
Fig. 5. Relationship between measured water quality parameters and changes in the microbial

community profile of locations distribution systems of the two systems where disinfectant

residuals were applied. The plots represent the same distance-based redundancy analysis (dbRDA)

based on Bray-Curtis distances of 16S rRNA amplicon sequencing data showing the clustering of

individual samples (left) and their impacting water quality parameters (right). Chloraminated

samples are were found to cluster in the purple ellipse and chlorinated samples in the green ellipse.

Significance codes based on ANOVA are indicated as follows 0 '***' 0.001 '**' 0.01 '*' 0.05 '.'

0.1 ' ' 1.