Short term supplementation of celecoxib shifted butyrate production on a simulated

model of the gut microbial ecosystem and ameliorated *in vitro* inflammation

Emma Hernandez-Sanabria[1], Evelien Heiremans[1], Marta Calatayud Arroyo[1], Ruben

Props[1], Laurent Leclercq[2], Jan Snoeys [2], and Tom Van de Wiele[1]

[1] Center for Microbial Ecology and Technology (CMET), Ghent University, Coupure

Links 653, 9000 Ghent, Belgium. [2] Janssen Research & Development, A Division of

Janssen Pharmaceutica NV, Turnhoutseweg 30, Beerse, Antwerp, 2340, Belgium.

Running title: Celecoxib impacts butyrate in an *in vitro* gut microbial ecosystem

Correspondence: Tom Van de Wiele, Ghent University, Faculty of Bioscience

Engineering, Center for Microbial Ecology and Technology (CMET), Coupure Links

653, B-9000 Gent, Belgium, phone: +32 (0)9 264 59 76, fax: +32 (0)9 264 62 48. E-mail:

Tom.VandeWiele@UGent.be

ABSTRACT

Celecoxib has been demonstrated effective in the prevention and treatment of chronic

inflammatory disorders through inhibition of altered cyclooxygenase-2 (COX-2)

pathways. Despite the benefits for preventing colorectal cancer (CRC), continuous

administration may increase risk of cardiovascular events. Understanding microbiome-

drug-host interactions is fundamental for improving drug disposition and safety responses

of colon-targeted formulations, but little information is available on the bidirectional

interaction between individual microbiomes and celecoxib. Here we conducted *in vitro*

batch incubations of faecal microbiota to evaluate the short-term impact of celecoxib on

activity and composition of colon bacterial communities. Celecoxib-exposed microbiota

shifted metabolic activity and community composition, whereas total transcriptionally

active bacterial population was not significantly changed. Butyrate production decreased

by 50% in a donor-dependent manner, suggesting that celecoxib impacts *in vitro*

fermentation. Microbiota-derived acetate has been associated with inhibition of cancer

markers and our results suggest uptake of acetate for bacterial functions when celecoxib

was supplied, which potentially favoured bacterial competition for acetyl-CoA. We

further assessed whether colon microbiota modulates anti-inflammatory efficacy of

celecoxib using both a simplified inflammation model, and a novel *in vitro* simulation of

the enterohepatic metabolism. Celecoxib was responsible for only 5% of the variance in

bacterial community composition but celecoxib-exposed microbiota preserved barrier

function and decreased concentrations of IL-8 and CXCL16 in a donor-dependent manner

in our two cell models simulating inflammatory milieu in the gut. Our results suggest that

celecoxib-microbiome-host interactions may not only elicit adaptations in community

composition but also in microbiota functionality and may need to be considered for

guaranteeing efficient COX-2 inhibition.

IMPORTANCE: As inter-individual changes in the microbiome composition and

functionality may be a confounder on pharmacotherapy, we obtained mechanistic

understanding on how short-term celecoxib exposure impacts the functional activities of

colon communities. Celecoxib-exposed microbiota shifted metabolic activity without

impacting numbers of total active bacteria, but only community composition. Thus,

increased relative abundance of particular genera during celecoxib supplementation may

just indicate changes in maintenance energy. Focus on the influence of acetyl-CoA on

cancer cells and verifying whether changes in acetate:propionate:butyrate ratios rather

than in taxonomic diversity can be used as markers of decreased inflammation may be

the next frontiers for predicting successful NSAID therapy, and ultimately for developing

microbiome-based therapies.

Keywords: celecoxib, gut microbiome, *in vitro* models, butyrate, colorectal cancer

INTRODUCTION

Colorectal cancer (CRC) is the second most common cancer in the developed world (1), and it is expected to increase by 60% in the next 20 years (2). Increased synthesis of prostaglandins is a feature of chronic inflammation and has been proposed to promote tumorigenesis in the colon (3). Prostaglandin E_2 (PGE_2) is a key mucosal inflammatory mediator produced when membrane phospholipids are metabolized by cyclooxygenase (COX) enzymes. Previous research showed that COX-2 expression was increased in CRC (4), causing proliferation, enhanced angiogenesis and apoptosis suppression (5). In this way, COX-2 inhibitors such as celecoxib are among the most promising chemopreventive agents for CRC (6). Despite these benefits, their long-term use may shift cardiovascular homeostasis, inhibiting biosynthesis of vascular COX-2 dependent prostacyclin (PGI_2) (7) and increasing cardiovascular events risk (8).

Advances in omics techniques have provided insights on the role of the gut microbiome in cancer development (9). Currently, the driver-passenger model suggests that intestinal "driver" bacteria contribute to initiate CRC by damaging epithelial DNA (10). This microenvironment alteration impairs barrier function and favours opportunistic "passenger" bacteria, placing selective pressure on the microbiome (10).

When colon-targeted drugs are orally administered, they encounter the gut microbiota before reaching host tissues. Microbial drug metabolism may impact drug stability and activity, influencing toxicity and inflammation on the host (11). Conversely, drugs may also alter the gut microbiota, modulating community structure, and potentially resulting in dysbiosis (12). Understanding microbiome-drug-host interactions will be crucial to assess the potential impact on ADME and safety responses of intestinal targets. Although the gut microbiome plays an enormous role in patient health, and in pharmacokinetic and pharmacodynamic response (13), pharmaceutical companies have only recently started to

explore the microbiome as potential contributor to drug outcomes. Similarly, regulatory agencies are not currently considering gut bacteria within the official evaluations before approval of colon-targeted drugs. Besides the microbial modulation of drug bioavailability and pharmacokinetics (11), low solubility and the small aqueous volume in the colon may additionally hinder drug efficacy (14). As interest in developing novel colon-targeted drug delivery systems for CRC has increased, insight into the presystemic biotransformation by the gut microbiota is fundamental. *In vitro* studies have shown that bacteria can metabolize celecoxib (15), potentially jeopardizing its anti-inflammatory effects. Additionally, whether celecoxib presence can modulate the host-microbiome crosstalk remains unresolved. Prior reports correlating gut microbiome changes with celecoxib treatment relied on DNA amplicon sequencing (16). As the presence of dead bacterial cells may obscure the interpretation from microbiome data, we analysed the bacterial community composition based on cDNA, as a proxy for the metabolically active community.

Colonic transit times (17-24 hrs) (17, 18) are longer compared to those of the small intestine (3-4 hrs), and overall systemic absorption of poorly soluble drugs may be improved with colon-delivery systems (17). Although celecoxib can be absorbed throughout the GI tract, we simulated delivery in a proximal colon-targeted formulation because of the environment vicinity between proximal colon and small intestine. Thus, the aim of this research was to screen the short-term bidirectional interactions between celecoxib and the microbiome of the *in vitro* proximal colon, and to elucidate the ultimate impact of the microbiota on the anti-inflammatory efficacy of celecoxib.

RESULTS

Celecoxib impacts bacterial fermentative metabolism in a donor-dependent manner.

Short-chain Fatty Acids (SCFAs) were used to monitor functional activities of the colon

communities following exposure to celecoxib (CX) or its vehicle (PEG). Total SCFA

production from separate faecal microbiota incubations of 8 different individuals was

significantly lower for CX incubations compared to those with PEG ($P < 0.05$, Table 1),

but pH remained stable throughout the experiment (Table 1). While the 60:20:20 ratio of

the three major SCFA (acetate, propionate, and butyrate) was consistent across donors at

the initial time point (Supplementary Table 1A), a shift was evident at the end of the

incubation. Both relative and absolute concentrations of butyrate and propionate were

higher in samples treated with only PEG compared to those with CX. In fact, celecoxib

significantly lowered butyrate after 16 hours ($P < 0.05$, Table 1), except in donors 1, 4

and 8 (Supplementary Table 1B). In contrast, total active bacterial load remained

unchanged when celecoxib was supplemented (Table 1, Supplementary Table 2). These

findings might indicate that bacteria either alter their metabolic activity when exposed to

CX or that changes in bacterial composition lead to shifts in fermentation profiles.

119   Bacterial abundance remains stable, but microbiome diversity and composition

120   shifted following celecoxib supplementation

Total DNA may also include that of dead bacterial cells, so we analysed community

composition based on cDNA, as a proxy for the metabolically active bacterial community

Quantitative RT-PCR revealed that total 16S rRNA gene copy numbers from

metabolically active bacteria were not significantly different between treatments, across

donors (Supplementary Table 2). In contrast, community composition and structure were

influenced, as celecoxib was associated with reduced diversity at the end of the incubation

(Figure 1, Table 2A and 2B), with significant donor effect (PERMANOVA $P < 0.001$,

Figure 2). CX shifted alpha diversity and community composition metrics across donors

(Supplementary Table 3, Figure 3). Although interindividual variability was significant,

metabolically active taxa with the highest relative abundances in control incubations

were: *Bacteroides* sp., *Faecalibacterium* sp., Ruminococcaceae UGC002, and

Megasphaera sp. In contrast, *Alistipes* sp., *Dialister* sp., *Escherichia-Shigella* sp.,

Lachnospira sp., *Megamonas* sp., *Parabacteroides* sp., *Roseburia* sp., and *Sutterella* sp.

were highly abundant when CX was supplied (Figure 3).

Short-term exposure to celecoxib increases number of community metabolic

networks

Association networks were constructed to highlight potential shifts among bacterial

relative abundances after short-term exposure to CX or its vehicle. The number of links

in the network was lower when the vehicle PEG alone was provided (Figure 4A), but the

central position of *Phascolarctobacterium*, *Parabacteroides*, *Coprococcus* 2 and

Unclassified Victivallales and the linkages among these genera suggest concomitant

changes in their relative abundances in presence of PEG. The central position of the

acetate-producing *Erysipelatoclostridium* in the celecoxib network (Figure 4B) indicated

that its relative abundance was associated with *Subdoligranulum*, a butyrate and lactate-

producing genus, in a donor-dependent manner ($P < 0.05$, Supplementary Table 4).

Networks connecting bacterial load, metabolites and relative abundances were assembled

to monitor functional alterations associated with changes in genera abundances. Activity-

based relative abundance of *Bacteroides* was positively associated with butyrate,

propionate and valerate levels and total 16S rRNA gene copy number in presence of PEG

(Figure 5A, Supplementary Table 4). These results confirmed that *Bacteroides* was one

of the most abundant active genera, as first suggested in Figure 3. Higher number of links

in comparison with PEG were detected after 16 h of exposure to celecoxib (Figure 5).

Indeed, one network linking acetate with the relative abundance of diverse taxa was only

present when celecoxib was supplied and contained the largest number of active genera

participating. Relative abundance of *Faecalibacterium* decreased over time with

celecoxib supplementation, corresponding with the decline in butyrate levels (Figure 3).

However, this genus tended to be positively associated with acetate production (Figure

5B). Relative abundance of active *Lachnoclostridium* was significantly associated with

butyrate in the celecoxib treatment (Figure 5B). Here, *Oscillospira*, *Allisonella*,

Hungatella and *Streptococcus* were linked because they possibly participated in butyrate

production as well. *Megamonas* and *Megasphaera* were other genera with high relative

abundances positively associated with branched-chain SCFA and isovalerate. The

associations between the latter metabolites and *Colinsella* remained unchanged upon

celecoxib exposure, suggesting that this community function is either independent from

drug supplementation or directly associated with PEG addition (Figure 5A and 5B).

Similarly, associations between *Allisonella* (butyrate consumer), *Streptococcus* (butyrate

producer) and butyrate were common for both treatments. Results from the network

analysis indicate that shifts in the community metabolic activity upon short-term

supplementation of celecoxib, potentially impacted butyrate production.

Inter-individual differences play a role in reduction of *in vitro* inflammation when

celecoxib is presented to the microbiome.

Understanding the microbial interactions with the host epithelium is essential to assess

whether the anti-inflammatory efficacy of celecoxib remains unaltered following

exposure to the simulated microbiome. We incorporated the immune component in a

simplified inflammation model using THP-1 (macrophage-like) cells. We further verified

the microbiome impact using a multicellular model simulating the enterohepatic

ecosystem, including enterocyte-like and goblet-like cell lines, plus THP-1 cells. This set-

up enabled closer communication between cell lines, as would occur when macrophages

infiltrate the underlying intestinal mucosa.

Direct contact of macrophage-like cells with filter-sterilized supernatants from the CX-

exposed microbiota resulted in a variable outcome in IL-8 and CXCL16 concentrations.

Compared with the PEG controls, a decrease in IL-8 and CXCL16 was only observed in

4 out of 8 individuals (Figure 6A and 6B).

High IL-8 concentrations were found when non-bacterially exposed celecoxib was

applied in the enterohepatic model (Figure 6C). In contrast, the lowest concentration of

IL-8 and thus the most significant decrease in inflammatory response was found with the

celecoxib-exposed supernatant of donor 6 ($P < 0.05$, Table 3). Celecoxib that was not in

contact with bacteria promoted the highest decrease in transepithelial electrical resistance

(TEER) ($P < 0.05$, Table 3), in comparison with supernatants from celecoxib-exposed

bacteria. This outcome indicates less damage to epithelial barrier function and decreased

inflammatory response when gut microbiota is exposed to celecoxib.

DISCUSSION

As the lipophilic characteristics of celecoxib favour metabolic elimination (19), exploring

only the pathways for celecoxib distribution in the host is insufficient for assessing drug

bioavailability. *In vitro* models have been applied to simulate drug-bug associations in

the human gut (20, 21), but the impact of NSAIDs on the metabolic activities of the

microbiome has been overlooked. Previous studies reported the influence of different

NSAIDs on the gut microbiome (22), yet no changes in bacterial richness, microbiome

composition or beta diversity were found when celecoxib was supplied (16). These results

focused on screening community structure using DNA-based amplicon sequencing,

which may also include DNA of dead bacterial cells. As such approach may deliver a

partial overview, we used cDNA as an indicator of the metabolically active bacterial

population and we performed community composition analyses based on information

from transcribed RNA (23). In this way, we intended to highlight the main bacterial

genera whose metabolic activities were impacted by celecoxib in the RNA pool. Activity-

based cell load was not significantly different, and the only alterations were on the

bacterial metabolic activities. Although composition, diversity and richness were not

significantly different as result of the treatment, differences in the most abundant genera

between PEG and celecoxib were observed. PERMANOVA analysis suggested that

donor was the main driver for dissimilarities in community structure.

The shifts in microbial metabolism observed in our study may indicate that celecoxib can

inhibit functional activities of bacterial groups, rather than of single genera, ultimately

impacting microbial fermentation (24). This was confirmed by the total active bacterial

load, which remained unchanged between treatments. Celecoxib users have been reported

to have enrichment of Enterobacteriaceae and Acidaminococcaceae (22). Here, we

validated that the relative abundance of transcriptionally active *Escherichia-Shigella*

(Enterobacteriaceae) was increased with celecoxib supplementation, whereas the relative

abundance of active *Acidaminococcus* (Acidaminococcaceae) was positively associated

with acetate in the same treatment. These results highlight the applicability of our *in vitro*

model used for this study. Coriobacteriaceae (*Colinsella*) has been proposed to participate

in xenobiotic metabolism by decreasing levels of serine and glycine (24), which can be

both fermented to acetate in the gut (25). We observed that the relative abundance of this

genus was positively associated with isovalerate levels when celecoxib was added. As

indicated in the corresponding KEGG module, *Colinsella aerofaciens* has the capacity of

using isovalerate for synthesising leucin and serine (26), confirming its potential

xenobiotic role. On the other hand, the capacity of *Alistipes* for producing succinate (27),

may explain its association with propionate in the PEG treatment. *Dialister* is another

succinate producer (28) present at high relative abundance, but it does not seem to

participate in the propionate network when PEG was supplied. This may suggest that,

even based on cDNA, high relative abundance may not be an indicator of metabolic

activity but just of maintenance metabolism.

The significantly lower butyrate concentration post-celecoxib delivery may have several

implications. Butyrate has tumour inhibitory properties (29) but its paradoxical

correlation with polyp formation (30, 31) suggests that the CRC-butyrate interplay is not

just a direct effect of altered SCFAs concentrations, and may also be associated with

changes in microbiome functionality. Indeed, the protecting properties of celecoxib

towards CRC (32, 33) may be partially derived from lowering bacteria-derived butyrate.

Nutrients can be utilized in their final form by both host and gut microbes, leading to

competition for resources (34). In fat, blocking fatty acid supply and host recapture has

been proposed as strategy for influencing cancer cell bioenergetics (35, 36), but few

studies have addressed the potential role of acetate in cancer (37). Acetyl-CoA is the main

carbon source in mammalian cells, and the ligation of acetate and CoA by acetyl-CoA

synthetase (ACSS) contributes to the supply of this metabolite (37). Cancer cells use

acetyl Co-A during aerobic metabolism regardless of stress conditions (37). Butyrate is

also metabolised to acetyl-CoA (29), inhibiting tumour growth via silencing acetyl-CoA

synthetases (35) or potentially through acetate uptake. As acetyl-CoA is a bacterial

substrate for fatty acid production (38), our network analysis suggests that acetate was

used for bacterial metabolic functions when celecoxib was supplied. Thus, celecoxib may

favour bacterial competition for acetyl Co-A, generating potential host protective effects.

Indeed, microbiota-derived acetate has been associated with inhibition of cancer-

associated deacetylation (39). Increasing acetate concentration shifts SCFA ratios (40),

decreasing butyrate and explaining why the acetate:propionate:butyrate ratios tended to

change when celecoxib was supplied. *Faecalibacterium* sp. is a well-known butyrate

producer and acetate consumer (40) and its presence in the acetate network may indicate

that the relative abundance of this genus increased when celecoxib was provided and

acetate levels increased.

Synthesis of reduced products including H_2, succinate, and butyrate creates redox balance

during colonic fermentation, whereas formation of oxidised products, such as acetate, is

associated with bacterial ATP production (38). Fermentative activities of

Lachnoclostridium yield ethanol, acetate, CO_2 and H_2 (41, 42), explaining the numerous

links with other genera in the celecoxib network. These potential cross-feeding

interactions (43, 44), may support its positive association with the total 16s rRNA gene

copy number. Acetate is a co-substrate for butyrate production (45), but butyrate

production may be reduced when NADH used for H_2 formation during glycolysis is not

available to convert acetoacetyl-CoA to butyryl-CoA (38). Acetate-producing

Lachnoclostridium and *Oscillospira* may have been involved in the shift from butyrate to

acetate production when celecoxib was supplied. This is because these genera possess

transporters involved in uptake of oligosaccharides and glycerol-P (43). Additionally,

Peptococcaceae uses the acetyl-CoA pathway in a reverse direction to oxidize butyrate

(46). Acetyl Co-A is consequently routed into acetate formation (45), which is

energetically advantageous for bacterial use (38). In fact, butyrate producers may shift

metabolic pathways towards acetate under stress conditions (47). As in our study, other

acetate producers such as Bacteroidetes and Prevotellaceae positively correlated to

butyrate-producing communities using the acetyl CoA pathway for butyrate production

(46), supporting the observed shift in butyrate towards acetate (48).

Although the role of SCFA on inflammation and CRC progression has been documented,

most of the studies lack natural commensal microbiota cross-talking with host cells (29,

31). As interindividual differences in microbiome play a crucial role on drug responses

(12), elucidating whether microbiome fermentation products synergistically act with

NSAIDs for impacting inflammation is paramount. Celecoxib was responsible for only

5% of the variance in bacterial community composition among the 8 donors, but the effect

of celecoxib-exposed bacteria on the proinflammatory IL-8 was sufficient to be detected

in a donor-dependent manner. Moreover, samples derived from 50% of the donors

showed increased CXCL-16, which is a marker of favourable prognosis in CRC (49).

This suggests that the interaction between celecoxib and microbiome to resolve

inflammation may be not solely related to community composition, and functionality

must be accounted for. While butyrate seemed to be reduced with short-term treatment,

Acetate:Propionate:Butyrate ratio and relative levels of acetate (of the total SCFA) may

be other functional bacterial metrics to be considered for assessing successful reduction

on inflammation when NSAIDs are administered.

As indicated in our study, gut microbiome activities and its interplay with the host can be

modulated through celecoxib. Our *in vitro* model retained interindividual variability,

allowing for mechanistic research on drug-microbiota-host interplay, fundamental for

developing individualised therapeutic strategies. Divergence in the molecular approaches

(i.e. next generation sequencing, qPCR) used for investigating CRC-associated

microbiome signatures contributes to the lack of consensus on the composition of

microbiota in CRC patients. Information on the role of SCFA in cancer signalling remains

inconclusive, and thus, future studies assessing the crosstalk of the microbiota and colon

cells in CRC are imperative. Certainly, individual microbiome may need to be considered

before supplementing celecoxib as a colon targeted drug for CRC. However, focusing on

functionality rather than taxonomic diversity may be the next frontier for developing

predictive markers of CRC-induced dysbiosis and NSAID efficacy.

MATERIALS AND METHODS

Faecal incubations simulating the proximal colon environment

Batch incubations of faecal samples were conducted using phosphate-buffered nutritional medium (pH 5.9, 0.1 M) containing (per litre): 12.25 g of KH_2PO_4 (Carl Roth GmbH, Karlsruhe, Germany), 1.78 g of Na_2HPO_4 (Carl Roth GmbH, Karlsruhe, Germany), 3 g of yeast extract (Oxoid Ltd, Basingstoke, Hampshire, England), 1 g of proteose peptone (Oxoid Ltd, Basingstoke, Hampshire, England), 0.25 g of gum Arabic from acacia tree (Sigma Aldrich Co., St-Louis, USA), 0.5 g of apple pectin (Sigma Aldrich Co., St-Louis, USA), 0.25 g of xylan (Carl Roth GmbH, Karlsruhe, Germany), 1 g of potato starch (Prodigest, Zwijnaarde, Belgium), and 4 g of mucin from porcine stomach Type II (Sigma Aldrich Co., St-Louis, USA).

Faecal material was obtained from eight healthy, non-obese, non-smoking volunteers (4M and 4F, 30 ± 2.8 yo), following an omnivorous diet and without antibiotic exposure for the prior 6 months. Stool samples were collected and prepared as previously described (50). Celecoxib (Janssen Pharmaceutica, Beerse, Belgium) was dissolved in 70:30 (v/v) PEG_{400}:H_2O as vehicle, at a stock concentration of 100 mg/mL. For each donor, triplicate incubations of 10% faecal slurry, 100 mg/mL of celecoxib or 70:30 (v/v) PEG_{400} (Acros, Geel, Belgium):H_2O, and nutritional medium (final volume = 175 mL) were performed for 16 hours at 37°C and 120 rpm (New Brunswick Scientific InnOva 4080 Incubator Shaker, Wezembeek-Oppem, Belgium), under anaerobic conditions (90%-10% N_2/CO_2). Once the incubation was completed, samples were collected, flash-frozen using liquid nitrogen and preserved at −80 °C for further analysis.

Surveying community functionality and composition

Short-chain Fatty Acids (SCFAs) production. SCFA were used as benchmarks of

community activity. Extraction and analyses have been described by Truchado et al. (51).

Differences in SCFA concentrations among treatments were compared using a repeated

measures mixed model (52), with the lsmeans adjustment and Bonferroni correction for

multiple comparisons (SAS, 2012). Statistical significance was assumed at $P < 0.05$.

Quantification of total metabolically active bacterial community. Samples collected

from the faecal incubations at 0h (start) and after 16h (end of experiment) were thawed

on ice, centrifuged for 5 minutes at 4°C and $5000 \times g$, and disrupted two times for 25

seconds at 1400 rpm (Power Lyser 24, MO BIO Laboratories, Carlsbad, USA), with a 5-

second interval, where the samples were placed on ice. RNA was further extracted using

the Nucleospin RNA plus kit (Macherey Nagel, Düren, Germany), following the

manufacturer's instructions. Reverse transcription was completed using the Reverse

Transcriptase Core kit (Eurogentec, Seraing, Belgium).

For the quantification of total bacterial 16S rRNA gene copy numbers, a standard curve

was constructed using serial dilutions of plasmid DNA from a clone identified

as *Bifidobacterium breve* LMG 33264. Briefly, universal bacterial primers 27F and

1492R were used to amplify the full-length 16S rRNA gene from the plasmid DNA of

a *Bifidobacterium breve* clone. The resultant PCR product was purified using the using

the InnuPREP PCRpure Kit (AJ innuscreen, Berlin, Germany), cloned in *E. coli* with the

TA Cloning Kit (Life Technologies, Carlsbad, VS) and plasmid extracted with the

PureYield Plasmid Miniprep System (Promega, Leiden, The Netherlands). The mass

concentration of the plasmid was measured using an ND-1000 spectrophotometer

(NanoDrop Technologies, Wilmington, DE), converted to the molecule concentration

(53) and copy numbers of total bacteria in 50 ng of cDNA and per ml of faecal incubation

were determined by relating the threshold cycle (C_T) values to the standard curves based

on the following equation: $Y = -3.193 \times \log X + 35.003$ (Y, C_T value; X, copy number of

16S rRNA gene) ($r^2 = 0.996$) (53).

Quantitative reverse transcription-PCR (qRT-PCR) analysis of the metabolically active

community was achieved with the StepOnePlus Real-Time PCR system (Applied

Biosystems, Foster City, CA USA). Amplification reactions were in: 10× Taq buffer with

KCl, 25 mM of $MgCl_2$, 10 mM of dNTP mix, 10 µM of each forward and reverse primers,

20 mg/mL of BSA (Thermo Scientific, Waltham, USA), 1.25 U of Taq Polymerase, 50

ng of cDNA and 0.1× SYBR Green I ([Invitrogen, Carlsbad, USA] provided at 10 000×,

stock solutions of 20× were prepared in DMSO). qRT-PCR conditions included 10

minutes at 95°C, followed by 40 cycles of 15 s at 95°C and 30 s at 60°C. The efficiencies

(E) of RT-PCR were calculated from the given slopes in StepOnePlus software using the

following equation: $E = [10^{(-1/\text{slope})} - 1] \times 100\%$. Data generated from reactions with

efficiencies between 90 and 110% were used for further analysis (53).

Community composition and dynamics.

The V5-V6 hypervariable region of the 16S rRNA gene from the cDNA was amplified

using primers 341F and 785R. High-throughput amplicon sequencing was performed

with the Illumina MiSeq platform according to the manufacturer's guidelines at LGC

Genomics GmbH (Berlin, Germany). Library preparation and purification as well as

bioinformatic processing are described in the supplemental material.

Data was imported into R using *phyloseq* (54) and taxon abundances were rescaled by

calculating the taxon proportions and multiplying them by the minimum sample size (n =

2201) present in the data set (55). Inverse Simpson was the metric used for assessing

alpha diversity and Pielou's index was used as indicator of community evenness (56).

Differences in alpha diversity and evenness measures were compared among treatments

using a repeated measures mixed model in SAS (version 9.4, SAS Institute, Cary, USA).

Beta diversity based on Chao and Bray-Curtis indices was used to examine dissimilarity

and determine the impact of treatment and time on microbial community structure (52).

PCoA was employed to visualize the differences among samples, using the vegan

package in R (57). Stratified permutational multivariate analysis of variance

(PERMANOVA) with 999 permutations was conducted to indicate the significance of

each covariate (time or treatment) on the microbial community.

ANOVA was applied to reveal whether the distribution of the genera was different

between treatments over time (57). Because of the over-dispersion in the OTU data, a

zero-inflated negative binomial distribution count model was used to assess the effect of

donor, time and treatment and the interactions between donor*time*treatment on each

individual genus. The model was selected based on the Akaike Information Criterion

(AIC). Differences among library size sample were accounted for with the offset option

in proc GLIMMIX in SAS (52).

Sparse inverse covariance selection was employed to infer an association network within

treatments, using the package SpiecEasi (v0.1.2) (58) in R (v3.5.1). Neighbourhood

selection method (S-E(MB)) was used to construct a network of bacterial genera

displaying core networks with either functional or phylogenetic similarities. The

calculated adjacency matrix was examined with the software Gephi 0.9.2 (59), using the

Fruchterman-Reingold algorithm for visualizing significant correlations ($R \geq 0.7$). This

method computes a layout in which the length of the connections indicates the absolute

value of the correlation. The nodes representing genera whose relative bacterial

abundances have several significant correlations among them were placed closer to each

other in the network (50).

Bipartite networks highlighted functional associations among bacterial genera and

metabolites (52), using a pair-wise similarity matrix obtained from a Regularised

Canonical Correlation Analysis (60). Values of the similarity matrix were computed as

the correlation between relative abundances of bacterial genera and metabolic variables,

projected onto the space spanned by the first 3 components retained in the analysis

($r \geq 0.75$). Genera in the plot were close to correlated variables in the treatment where

they were more abundant (52).

Assessing the anti-inflammatory potential of bacteria-exposed celecoxib

Simplified inflammation model. The human monocytic leukaemia cell line THP-1

(THP-1, ECACC 88081201, Public Health England, UK) was grown in Roswell Park

Memorial Institute (RPMI) 1640 culture medium (Thermo Fisher Scientific, Waltham,

USA) supplemented with 10% heat-inactivated foetal bovine serum (iFBS) (Greiner Bio-

One, Wemmel, Belgium) and 1% antibiotic-antimycotic solution (10,000 units/mL of

penicillin, 10,000 µg/mL of streptomycin, and 25 µg/mL of Amphotericin B) (Thermo

Fisher Scientific, Waltham, USA), at 37°C in 10 % CO_2 in a humidified incubator. THP-

1 cells were sub-cultured twice a week and used between passage 20 and 30.

Macrophage-like phenotype was obtained by treating cells with 25 ng/mL phorbol 12-

myristate 13-acetate (PMA, Sigma Aldrich, St-Louis, USA) for 48 h in 24-wells culture

plates (Corning, Sigma Aldrich, St-Louis, USA) at a density of 1 x 10^5 cells/cm^2.

Differentiated, plastic-adherent cells were washed twice with culture medium

(supplemented RPMI 1640 without PMA) and incubated for 24 h. Medium was removed

and replaced by supernatant from the eight incubations, passed through a sterile 0.2 µm

filter and diluted five times with RPMI 1640 without supplementation. Celecoxib solution

at 100 mg/mL dissolved in 70:30 (v/v) PEG:H_2O or 70:30 (v/v) PEG:H_2O alone were

diluted 1:5 with medium and added as controls of non-bacteria exposed celecoxib and

vehicle. This dilution step was performed to simulate low solubility and the small aqueous

volume in the colon (14). LPS (100 ng/mL) dissolved in RPMI 1640 was added to the

cells to simulate systemic inflammation, as in CRC. The control used in all measurements

was PMA-differentiated THP-1 macrophages exposed to the culture medium.

Development of an *in vitro* model of the enterohepatic system. A novel model

including intestinal epithelial (Caco-2, ECACC 86010202), goblet (HT29-MTX-E12

ECACC 12040401, Public Health England, UK), hepatic (HepG2 ATCC HB-8065,

Molsheim, France), and immune-like cells (THP-1, ECACC 88081201, Public Health

England, UK) was developed to study host-drug-microbiome interactions (61, 62). Cell

layers were grown in a double-chamber insert (Corning, Sigma Aldrich, St-Louis, USA),

and maintained for 23 days. Briefly, 100 µl of a THP-1 cell suspension (10^6 cells/mL) in

20 mg/mL collagen from rat tail type I (Sigma Aldrich, St. Louis, USA) were spread in

the basolateral surface of inverted inserts. After 45 min of incubation at 37ºC and 10%

CO_2, a 9:1 cell suspension of Caco-2/HT29-MTX cells (density of 7.5×10^4 cells/cm^2) was

added to the apical side of the insert in 1.5 ml of Dulbecco's Modified Eagle's medium

(DMEM) supplemented with 10% (v/v) of iFBS and 1% of antibiotic-antimycotic

solution (v/v). The basolateral compartment of the double-chamber system was filled with

2 ml of supplemented RPMI 1640, and the medium was refreshed every 48 h. Inserts were

transferred to 6-well plates with pre-seeded HepG2 cells at day 19 post-seeding, when the

cell culture media in the basal compartment was replaced by RPMI 1640, and the co-

culture was maintained 24 h prior to the assay. HepG2 cells were seeded at a density of

10^5 cells/cm^2 and maintained 5 days in supplemented minimal essential medium (MEM)

supplemented with 10% (v/v) of iFBS and 1% (v/v) of antibiotic-antimycotic solution.

Five days prior the assay all the cell culture media was depleted of antibiotic/antimycotic

solution. Supernatants from volunteers (2F, 29 yo) showing extreme microbiota

metabolism responses in the short-term incubations were used for validating whether

celecoxib effectively decreases inflammation even after exposure to the gut microbiota.

Volunteer 6 displayed the most significant decrease in butyrate after 16 h of incubation,

while volunteer 8 did not present any changes. Triplicate supernatants from each

volunteer were passed through a sterile 0.2 µm filter, diluted five times with DMEM

without supplementation, pH was adjusted to pH 7-7.2 with filter-sterilized NaOH 0.5M

(Carl Roth GmbH & Co), and placed in the apical compartment of the model. Results

from the simplified inflammation model indicated that there were no significant

differences between control and treatment among donors. Thus, we only included

celecoxib solution dissolved in 70:30 (v/v) PEG:H_2O or 70:30 (v/v) PEG:H_2O alone in

the enterohepatic model. Celecoxib solution at 100 mg/mL and a 70:30 (v/v) PEG:H_2O

were diluted 1:5 with DMEM without supplementation, and added to the apical

compartment, as controls of non-bacterially modified celecoxib and vehicle. This dilution

step was performed to simulate low solubility and the small aqueous volume in the colon

(14). LPS (100 ng/mL) was added to the basal compartment to simulate systemic

inflammation. Plates were incubated for 16 h at 37°C, 10% CO_2 and 95% humidity. TEER

was evaluated with the Millicell ERS-2 Voltohmmeter at the beginning and at the end of

the assay (Merck-Millipore, Overijse, Belgium).

Evaluating inflammatory response. Concentration of IL-8 and CXCL16 were measured

in the simplified inflammation model with the Human IL-8 (CXCL-8) and Human

CXCL16 Mini ABTS ELISA Development Tests (PeproTech, London, UK) and the

ABTS ELISA Buffer Kit (PeproTech, London, UK), following the manufacturer's

instructions. Samples were not diluted, and determination was performed at 405 nm, with

correction for 650 nm (Infinite 200 PRO, Tecan, Männedorf, Switzerland). IL-8 was

additionally determined in the presystemic metabolism model, using samples from the

basal compartment diluted ten times with PBS (ABTS ELISA Buffer Kit, PeproTech,

London, UK), and measured as indicated above.

Data Availability

Raw 16S rRNA reads have been made available on the SRA under accession number ID

PRJNA540406.   Original   R   scripts   are   available   in   GitHub

(https://github.ugent.be/ehernand/CX_sequencing).

ACKNOWLEDGEMENTS

E.H.-S. is a postdoctoral fellow supported by Flanders Innovation and Entrepreneurship

(Agentschap Innoveren & Ondernemen). M.C. was supported by an FWO postdoctoral

fellowship (FWO/12R2717N). E.H.-S. and T.Vd.W. conceived the simulations of the

colon environment. E.H.-S. and M.C. devised cell models. J.S. and L.L. supplied material

and advised on the use of celecoxib. E.H.-S., E.H., and M.C. conducted all laboratory

work. R.P. completed the amplicon sequencing library pre-processing and guided

sequencing data mining. E.H.-S. performed the statistical analyses and interpretation of

all data generated, produced figures and tables, and wrote the manuscript. All authors

read and approved the manuscript. The authors gratefully acknowledge the volunteers

participating in this study. We thank Dr. Davide Gottardi for his scientific advice and

Racha El Hage, Fabian Mermans, Olivier Dezutter, and Dr. Chiara Ilgrande for their

technical assistance. The authors declare that they have no competing interests.

REFERENCES

1.   Kumar M, Cash BD. 2017. Screening and surveillance of colorectal cancer using
CT colonography. Current treatment options in gastroenterology 15:168-183.

2.   Arnold M, Sierra MS, Laversanne M, Soerjomataram I, Jemal A, Bray F. 2017.
Global patterns and trends in colorectal cancer incidence and mortality. Gut
66:683-691.

3.   Wang J, Cho NL, Zauber AG, Hsu M, Dawson D, Srivastava A, Mitchell-
Richards KA, Markowitz SD, Bertagnolli MM. 2018. Chemopreventive efficacy
of the cyclooxygenase-2 (Cox-2) inhibitor, celecoxib, is predicted by adenoma
expression of Cox-2 and 15-PGDH. Cancer Epidemiology and Prevention
Biomarkers 27:728-736.

4.   Bhala N, Emberson J, Merhi A, Abramson S, Arber N, Baron J, Bombardier C,
Cannon C, Farkouh M, FitzGerald G. 2013. Vascular and upper gastrointestinal
effects of non-steroidal anti-inflammatory drugs: meta-analyses of individual
participant data from randomised trials. Elsevier.

5.   Kawanishi S, Ohnishi S, Ma N, Hiraku Y, Murata M. 2017. Crosstalk between
DNA damage and inflammation in the multiple steps of carcinogenesis.
International journal of molecular sciences 18:1808.

6.   Wang D, DuBois RN. 2010. The role of COX-2 in intestinal inflammation and
colorectal cancer. Oncogene 29:781.

7.   Funk CD, FitzGerald GA. 2007. COX-2 inhibitors and cardiovascular risk.
Journal of cardiovascular pharmacology 50:470-479.

8.   Cuzick J, Otto F, Baron JA, Brown PH, Burn J, Greenwald P, Jankowski J, La
Vecchia C, Meyskens F, Senn HJ. 2009. Aspirin and non-steroidal anti-
inflammatory drugs for cancer prevention: an international consensus statement.
The lancet oncology 10:501-507.

9.   Johnson CH, Spilker ME, Goetz L, Peterson SN, Siuzdak G. 2016. Metabolite and
microbiome interplay in cancer immunotherapy. Cancer research 76:6146-6152.

10.   Tjalsma H, Boleij A, Marchesi JR, Dutilh BE. 2012. A bacterial driver–passenger
model for colorectal cancer: beyond the usual suspects. Nature Reviews
Microbiology 10:575.

11.   Spanogiannopoulos P, Bess EN, Carmody RN, Turnbaugh PJ. 2016. The
microbial pharmacists within us: a metagenomic view of xenobiotic metabolism.
Nature Reviews Microbiology 14:273.

12.   Wilson ID, Nicholson JK. 2017. Gut microbiome interactions with drug
metabolism, efficacy, and toxicity. Translational Research 179:204-222.

13.   Sousa T, Paterson R, Moore V, Carlsson A, Abrahamsson B, Basit AW. 2008.
The gastrointestinal microbiota as a site for the biotransformation of drugs.
International journal of pharmaceutics 363:1-25.

14.   Philip AK, Philip B. 2010. Colon targeted drug delivery systems: a review on
primary and novel approaches. Oman medical journal 25:79.

15.   Srisailam K, Veeresham C. 2010. Biotransformation of celecoxib using microbial
cultures. Applied biochemistry and biotechnology 160:2075-2089.

16. Bokulich NA, Battaglia T, Aleman JO, Walker J, Blaser MJ, Holt PR. 2016.
Celecoxib does not alter intestinal microbiome in a longitudinal diet-controlled
study. Clinical Microbiology and Infection 22:464-465.

17. HARDY JG, WILSON CG, WOOD E. 1985. Drug delivery to the proximal colon.
Journal of Pharmacy and Pharmacology 37:874-877.

18. Macfarlane G, Macfarlane S, Gibson G. 1998. Validation of a three-stage
compound continuous culture system for investigating the effect of retention time
on the ecology and metabolism of bacteria in the human colon. Microbial ecology
35:180-187.

19. Paulson SK, Vaughn MB, Jessen SM, Lawal Y, Gresk CJ, Yan B, Maziasz TJ,
Cook CS, Karim A. 2001. Pharmacokinetics of celecoxib after oral administration
in dogs and humans: effect of food and site of absorption. Journal of
Pharmacology and Experimental Therapeutics 297:638-645.

20. McConnell EL, Fadda HM, Basit AW. 2008. Gut instincts: explorations in
intestinal physiology and drug delivery. International journal of pharmaceutics
364:213-226.

21. Esch EW, Bahinski A, Huh D. 2015. Organs-on-chips at the frontiers of drug
discovery. Nature reviews Drug discovery 14:248.

22. Rogers MA, Aronoff DM. 2016. The influence of non-steroidal anti-inflammatory
drugs on the gut microbiome. Clinical Microbiology and Infection 22:178. e1-
178. e9.

23. Blazewicz SJ, Barnard RL, Daly RA, Firestone MK. 2013. Evaluating rRNA as
an indicator of microbial activity in environmental communities: limitations and
uses. The ISME journal 7:2061.

24. Montrose DC, Zhou XK, McNally EM, Sue E, Yantiss RK, Gross SS, Leve ND,
Karoly ED, Suen C, Ling L. 2016. Celecoxib alters the intestinal microbiota and
metabolome in association with reducing polyp burden. Cancer Prevention
Research:canprevres. 0095.2016.

25. Smith EA, Macfarlane G. 1997. Dissimilatory amino acid metabolism in human
colonic bacteria. Anaerobe 3:327-337.

26. Okuda S, Yamada T, Hamajima M, Itoh M, Katayama T, Bork P, Goto S,
Kanehisa M. 2008. KEGG Atlas mapping for global analysis of metabolic
pathways. Nucleic acids research 36:W423-W426.

27. Rautio M, Eerola E, Vaisanen-Tunkelrott M-L, Molitoris D. 2003.
Reclassification of bacteroides putredinis (Weinberg et al., 1937) in a new genus
alistipes gen. nov., as alistipes purtredinis comb. nov., and description of alistipes
finegoldii sp. nov., from human sources. Systematic and applied microbiology
26:182.

28. Morotomi M, Nagai F, Sakon H, Tanaka R. 2008. Dialister succinatiphilus sp.
nov. and Barnesiella intestinihominis sp. nov., isolated from human faeces.
International journal of systematic and evolutionary microbiology 58:2716-2720.

29. Donohoe DR, Collins LB, Wali A, Bigler R, Sun W, Bultman SJ. 2012. The
Warburg effect dictates the mechanism of butyrate-mediated histone acetylation
and cell proliferation. Molecular cell 48:612-626.

30. Belcheva A, Irrazabal T, Robertson SJ, Streutker C, Maughan H, Rubino S,
Moriyama EH, Copeland JK, Surendra A, Kumar S. 2014. Gut microbial
metabolism drives transformation of MSH2-deficient colon epithelial cells. Cell
158:288-299.

31. Kang M, Martin A. Microbiome and colorectal cancer: Unraveling host-
microbiota interactions in colitis-associated colorectal cancer development, p. *In*
(ed), Elsevier,

32. Arber N, Eagle CJ, Spicak J, Rácz I, Dite P, Hajer J, Zavoral M, Lechuga MJ,
Gerletti P, Tang J. 2006. Celecoxib for the prevention of colorectal adenomatous
polyps. New England Journal of Medicine 355:885-895.

33. Umar A, Steele VE, Menter DG, Hawk ET. Mechanisms of nonsteroidal anti-
inflammatory drugs in cancer prevention, p 65-77. *In* (ed), Elsevier,

34. Wasielewski H, Alcock J, Aktipis A. 2016. Resource conflict and cooperation
between human host and gut microbiota: implications for nutrition and health.
Annals of the New York Academy of Sciences 1372:20-28.

35. Röhrig F, Schulze A. 2016. The multifaceted roles of fatty acid synthesis in
cancer. Nature Reviews Cancer 16:732.

36. Schug ZT, Voorde JV, Gottlieb E. 2016. The metabolic fate of acetate in cancer.
Nature Reviews Cancer 16:708.

37. Schug ZT, Peck B, Jones DT, Zhang Q, Grosskurth S, Alam IS, Goodwin LM,
Smethurst E, Mason S, Blyth K. 2015. Acetyl-CoA synthetase 2 promotes acetate
utilization and maintains cancer cell growth under metabolic stress. Cancer cell
27:57-71.

38. Macfarlane S, Macfarlane GT. 2003. Regulation of short-chain fatty acid
production. Proceedings of the Nutrition Society 62:67-72.

39. Wu W, Sun M, Chen F, Cao AT, Liu H, Zhao Y, Huang X, Xiao Y, Yao S, Zhao
Q. 2017. Microbiota metabolite short-chain fatty acid acetate promotes intestinal
IgA response to microbiota which is mediated by GPR43. Mucosal immunology
10:946.

40. Duncan SH, Barcenilla A, Stewart CS, Pryde SE, Flint HJ. 2002. Acetate
utilization and butyryl coenzyme A (CoA): acetate-CoA transferase in butyrate-
producing bacteria from the human large intestine. Applied and environmental
microbiology 68:5186-5190.

41. Yutin N, Galperin MY. 2013. A genomic update on clostridial phylogeny: G ram-
negative spore formers and other misplaced clostridia. Environmental
microbiology 15:2631-2641.

42. Warnick TA, Methe BA, Leschine SB. 2002. Clostridium phytofermentans sp.
nov., a cellulolytic mesophile from forest soil. International journal of systematic
and evolutionary microbiology 52:1155-1160.

43. Gophna U, Konikoff T, Nielsen HB. 2017. Oscillospira and related bacteria–From
metagenomic species to metabolic features. Environmental microbiology 19:835-
841.

44. Mills S, Stanton C, Lane JA, Smith GJ, Ross RP. 2019. Precision nutrition and
the microbiome, Part I: Current state of the science. Nutrients 11:923.

45. Louis P, Flint HJ. 2017. Formation of propionate and butyrate by the human
colonic microbiota. Environmental microbiology 19:29-41.

46. Vital M, Howe AC, Tiedje JM. 2014. Revealing the bacterial butyrate synthesis
pathways by analyzing (meta) genomic data. MBio 5:e00889-14.

47. Geirnaert A, Steyaert A, Eeckhaut V, Debruyne B, Arends JB, Van Immerseel F,
Boon N, Van de Wiele T. 2014. Butyricicoccus pullicaecorum, a butyrate
producer with probiotic potential, is intrinsically tolerant to stomach and small
intestine conditions. Anaerobe 30:70-74.

48. Vital M, Gao J, Rizzo M, Harrison T, Tiedje JM. 2015. Diet is a major factor
governing the fecal butyrate-producing community structure across Mammalia,
Aves and Reptilia. The ISME journal 9:832.

49. Kee J-Y, Ito A, Hojo S, Hashimoto I, Igarashi Y, Tsuneyama K, Tsukada K,
Irimura T, Shibahara N, Takasaki I. 2014. CXCL16 suppresses liver metastasis of
colorectal cancer by promoting TNF-α-induced apoptosis by tumor-associated
macrophages. BMC cancer 14:949.

50. De Weirdt R, Hernandez-Sanabria E, Fievez V, Mees E, Geirnaert A, Van
Herreweghen F, Vilchez-Vargas R, Van den Abbeele P, Jauregui R, Pieper DH.
2017. Mucosa-associated biohydrogenating microbes protect the simulated colon
microbiome from stress associated with high concentrations of poly-unsaturated
fat. Environmental microbiology 19:722-739.

51. Truchado P, Hernandez-Sanabria E, Salden BN, Van den Abbeele P, Vilchez-
Vargas R, Jauregui R, Pieper DH, Possemiers S, Van de Wiele T. 2017. Long
chain arabinoxylans shift the mucosa-associated microbiota in the proximal colon
of the simulator of the human intestinal microbial ecosystem (M-SHIME). Journal
of Functional Foods 32:226-237.

52. El Hage R, Hernandez-Sanabria E, Calatayud Arroyo M, Props R, Van de Wiele
T. 2019. Propionate-producing consortium restores antibiotic-induced dysbiosis
in a dynamic in vitro model of the human intestinal microbial ecosystem. Frontiers
in Microbiology 10:1206.

53. Hernandez-Sanabria E, Goonewardene LA, Wang Z, Zhou M, Moore SS. 2013.
Influence of sire breed on the interplay among rumen microbial populations
inhabiting the rumen liquid of the progeny in beef cattle. PloS one 8:e58461.

54. McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible
interactive analysis and graphics of microbiome census data. PloS one 8:e61217.

55. McMurdie PJ, Holmes S. 2014. Waste not, want not: why rarefying microbiome
data is inadmissible. PLoS computational biology 10:e1003531.

56. Grunert O, Hernandez-Sanabria E, Vilchez-Vargas R, Jauregui R, Pieper DH,
Perneel M, Van Labeke M-C, Reheul D, Boon N. 2016. Mineral and organic

growing media have distinct community structure, stability and functionality in
soilless culture systems. Scientific reports 6:18837.

57.   Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara R, Simpson
666   GL, Solymos P, Stevens MHH, Wagner H. 2013. Package 'vegan'. Community
667   ecology package, version 2.

668   58.   Kurtz ZD, Müller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA. 2015.
669   Sparse and compositionally robust inference of microbial ecological networks.
670   PLoS computational biology 11:e1004226.

671   59.   Bastian M, Heymann S, Jacomy M. 2009. Gephi: an open source software for
672   exploring and manipulating networks. Icwsm 8:361-362.

673   60.   Lê Cao K-A, Boitard S, Besse P. 2011. Sparse PLS discriminant analysis:
674   biologically relevant feature selection and graphical displays for multiclass
675   problems. BMC bioinformatics 12:253.

676   61.   Paul W, Marta C. 2018. Resolving host–microbe interactions in the gut: the
677   promise of in vitro models to complement in vivo research. Current opinion in
678   microbiology 44:28-33.

679   62.   Calatayud MA, de Wiele Van T, Hernandez-Sanabria E. 2018. Assessing the
680   Viability of a Synthetic Bacterial Consortium on the In Vitro Gut Host-microbe
681   Interface. Journal of visualized experiments: JoVE.

685 Figure 1. Inverse Simpson index (D_2) was used to assess alpha diversity dynamics at the beginning and at the end of the incubation (0 and 16 h

686 post treatment) on each volunteer. Horizontal bars indicate means and distribution of each data point is indicated. Alpha diversity for celecoxib

687 treatment (CX) is indicated in green boxes, while purple is for this metric observed when the control treatment with the vehicle (PEG) was supplied.

688

Figure 2. Interindividual differences drive bacterial community composition. Principal

coordinate analysis (PCoA) revealed differences in community structure between

volunteers. The variance explained by the experimental factors ($P < 0.01$) was calculated

using PERMANOVA analysis and it is indicated on the top right. Donor was the factor

significantly contributing to the variations among individual microbiomes

Figure 3. Relative abundances of the bacterial genera occurring when celecoxib (CX) or a control treatment (PEG) were supplied. The upper panel

describes the community composition at the beginning of the treatment (0 h), while the lower panel presents the 12 taxa with the highest activity-

based (cDNA-based) relative abundances after 16 h of treatment.

Figure 4. Bacterial network of communities exposed to short-term administration of

celecoxib (CX) or its vehicle (control treatment, PEG). (A) The faecal microbiota exposed

only to PEG displayed a lower number of associations, while the network architecture

was enriched when CX was provided (B). This representation uncovers relationships

between clusters of genera across donors. Nodes representing bacterial genera with

similar abundances tended to cluster closely. Node diameter indicates the relative

abundance of each genus. Neighbourhood selection method (MB) was the graphical

model inference procedure selected. The associations described may be considered 'core'

functional relationships preserved across donors.

Figure 5: Community metabolic activity indicates that networks of bacterial interactions

were few when only the vehicle (PEG) was supplied (A) and increased following short-

term supplementation of celecoxib (B). These bipartite networks are based on the

regularised canonical correlations between relative bacterial abundances and relative

concentrations of the main short-chain fatty acids. Interactions have been filtered for an

absolute correlation above 0.5 and are coloured following the key shown. Significant

interactions are indicated by shorter lines, and genera with similar abundances within

treatment tended to cluster closely.

Figure 6. Concentration of pro-inflammatory cytokines was decreased in a donor-
dependent manner when celecoxib was supplied. Volunteers are numbered consecutively
as D1-D8. (A) IL-8 was significantly lower in the simplified inflammation model
subjected to the bacteria-exposed celecoxib from 3 of the volunteers screened (D1, D2
and D6). Two additional volunteers (D3 and D7) showed a trend for decreased IL-8, but
it was not significantly lower than the control treatment. (B) CXCL16 was significantly
lower in the simplified inflammation model subjected to the bacteria-exposed celecoxib
from only two volunteers (D2 and D6) and tended to be decreased in other two (D3 and
D4). (C) The two volunteers with the most extreme metabolic responses in butyrate
production were selected to be screened in a multicompartment cell model resembling the
enterohepatic system. Volunteer 6 (D6) had the most significant decrease in butyrate
levels and was considered "responder" to the CX treatment. Volunteer 8 (D8) did not
show any significant difference in butyrate between the control and the treatment after 16
h of incubation and it was considered as "non-responder". IL-8 concentration was
significantly lower in the model subjected to the supernatant of D6 containing bacteria-
exposed celecoxib ("responder volunteer"). The opposite effect was observed in D8
("non-responder volunteer"), suggesting that interindividual differences in the
functionality of the microbiome may play a role in the response to anti-inflammatory
drugs.

Table 1. Bacterial functionality metrics following short-term incubation with either

celecoxib or carrier (n = 8). CX, celecoxib; PEG, carrier. SEM, standard error of the mean.

Variable	Treatment (Mean ± SEM)		*P* value
	CX	PEG	
Acetate (mM)	11.51 ± 1.44	13.31 ± 1.50	0.39
Propionate (mM)	3.29 ± 0.41	3.86 ± 0.42	0.33
Butyrate (mM)	1.57 ± 0.33	3.20 ± 0.33	0.0007
Valerate (mM)	0.61 ± 0.15	0.78 ± 0.15	0.45
Isobutyrate (mM)	0.30 ± 0.07	0.34 ± 0.07	0.70
Isovalerate (mM)	2.70 ± 0.52	3.30 ± 0.54	0.42
Caproate (mM)	0.15 ± 0.04	0.12 ± 0.04	0.61
Isocaproate (mM)	0.001 ± 0.002	0.01 ± 0.002	0.009
Total SCFA (mM)	20.14 ± 2.37	24.91 ± 2.45	0.16
Total Branched SCFA (mM)	2.56 ± 0.49	3.25 ± 0.52	0.34
St:Br ratio	17.76 ± 4.48	12.71 ± 4.31	0.42
% of Acetate	60.87 ± 2.60	53.58 ± 2.77	0.06
% of Propionate	16.17 ± 0.65	15.75 ± 0.70	0.67
% of Butyrate	6.57 ± 0.64	12.99 ± 0.68	<0.0001
Acetate:Propionate:Butyrate ratio	61:16:7	54:16:13	-
pH	5.76 ± 0.04	5.75 ± 0.04	0.86
Total copy number of 16S rRNA gene	3.37E09 ± 3.33E08	4.35E09 ± 3.33E08	0.04

743 Table 2A. Impact of donor and supplementation of celecoxib (CX) or vehicle (PEG) on

744 initial bacterial community metrics (time 0 h), n = 8.

| Metric | Treatment | | SEM | *P* value | | Treatment*Donor interaction |
	CX	PEG		Treatment	Donor	
Evenness	0.61	0.60	0.0009	0.83	0.0001	0.23
Shannon (*H'*)	2.71	2.72	0.03	0.81	<0.0001	0.13
Simpson	0.85	0.85	0.007	0.43	<0.0001	0.04
Inverse Simpson (D_2)	8.00	8.48	0.53	0.54	<0.0001	0.76
Fisher's diversity	13.41	13.16	0.25	0.49	<0.0001	0.34
Total Richness (S)	92.50	95.83	3.83	0.55	0.002	0.77
Chao	144.12	128.55	9.54	0.27	0.01	0.59

747 Table 2B. Impact of donor and supplementation on bacterial community metrics,

748 following a short-term incubation (16 h) with either celecoxib (CX) or carrier (PEG), n =

749 8.

Metric	Treatment		SEM	P value		
	CX	PEG		Treatment	Donor	Treatment*Donor interaction
Evenness	0.55	0.59	0.01	0.06	0.003	0.38
Shannon (*H'*)	2.54	2.70	0.06	0.08	0.001	0.33
Simpson	0.84	0.86	0.009	0.14	0.009	0.10
Inverse Simpson (D_2)	7.59	8.13	0.63	0.55	0.02	0.23
Fisher's diversity	13.28	13.17	0.25	0.49	<0.0001	0.34
Total Richness (S)	92.56	97.12	2.78	0.88	0.005	0.07
Chao	94.01	114.65	5.34	0.01	0.008	0.89

752　Table 3. Epithelial integrity metrics of a multicompartment cell model simulating the

753　enterohepatic metabolism, after 16 h of incubation with CX-exposed bacteria. 0 h = prior

754　to LPS exposure, 16 h = after 16 h of incubation with LPS. Different superscripts indicate

755　significantly different means across treatments, within a particular time point. N = 3.

Treatment	Time point	TEER	SEM	*P* value		
				Treatment	Time	Treatment*Time interaction
CX alone (100 mg/mL)	0 h 16 h	385.00 271.67 [a]				
PEG	0 h 16 h	330.67 338.67 [a]				
Donor 6 + CX	0 h 16 h	470.00 425.33 [b]	34.61	0.007	0.01	0.40
Donor 8 + CX	0 h 16 h	398.33 311.33 [a]				
Control medium without LPS	0 h 16 h	421.33 337.33 [a]				
Control with LPS	0 h 16 h	428.67 433.00 [a]				