Short-term celecoxib supplementation impacts compositional and functional

features of *in vitro* gut microbial ecosystem

Emma Hernandez-Sanabria[1], Evelien Heiremans[1], Marta Calatayud Arroyo[1], Ruben

Props[1], Laurent Leclercq[2], Jan Snoeys [2], and Tom Van de Wiele[1]

[1] Center for Microbial Ecology and Technology (CMET), Ghent University, Coupure

Links 653, 9000 Ghent, Belgium. [2] Janssen Research & Development, A Division of

Janssen Pharmaceutica NV, Turnhoutseweg 30, Beerse, Antwerp, 2340, Belgium.

Running title: Celecoxib impacts *in vitro* gut microbial ecosystem

Correspondence: Tom Van de Wiele, Ghent University, Faculty of Bioscience

Engineering, Center for Microbial Ecology and Technology (CMET), Coupure Links

653, B-9000 Gent, Belgium, phone: +32 (0)9 264 59 76, fax: +32 (0)9 264 62 48. E-mail:

Tom.VandeWiele@UGent.be

ABSTRACT

Celecoxib has been demonstrated effective in preventing colorectal cancer (CRC), through inhibition of altered cyclooxygenase-2 (COX-2) inflammatory pathways. Alas, continuous administration may increase cardiovascular events risk. Understanding microbiome-drug-host interactions is fundamental for improving ADME and safety responses of colon-targeted formulations, but little information is available on the two-way interaction between individual microbiome and celecoxib. Here we evaluated short-term impact of celecoxib on *in vitro* activity and composition of colon bacterial communities using faecal incubations. Celecoxib-exposed microbiota shifted metabolic activity and community composition whereas total transcriptionally active bacterial population was not significantly changed. Butyrate production decreased by 50% in a donor-dependent manner, suggesting that celecoxib impacts *in vitro* fermentation. Microbiota-derived acetate has been associated with inhibition of cancer markers and our results suggested uptake of acetate for bacterial functions when celecoxib was supplied, which potentially favoured bacterial competition for acetyl-CoA. We further assessed whether colon microbiota modulates anti-inflammatory efficacy of celecoxib in an *in vitro* enterohepatic cell model. Celecoxib-exposed microbiota preserved barrier function and decreased inflammatory response in a donor-dependent manner.

IMPORTANCE: As inter-individual changes in the microbiome composition and functionality may be a confounder on pharmacotherapy, we obtained mechanistic understanding on how short-term celecoxib exposure impacts the functional activities of colon communities. Celecoxib-exposed microbiota shifted metabolic activity without impacting numbers of total active bacteria, but only community composition. Thus, increased relative abundance of particular genera during celecoxib supplementation may just indicate changes in maintenance activity. Focus on the influence of acetyl-CoA on

42 cancer cells and verifying whether changes in acetate:propionate:butyrate ratios rather

43 than in taxonomic diversity can be used as markers of decreased inflammation may be

44 the next frontiers for predicting successful NSAID therapy, and ultimately for developing

45 microbiome-based therapies.

46 Keywords: celecoxib, gut microbiome, *in vitro* models, butyrate, colorectal cancer

INTRODUCTION

Colorectal cancer (CRC) is the second most common cancer in the developed world (1), and it is expected to increase by 60% in the next 20 years (2). Advances in omics techniques have provided insights on the role of the gut microbiome in cancer development (3). Currently, the driver-passenger model suggests that intestinal "driver" bacteria contribute to initiate CRC by damaging epithelial DNA (4). This microenvironment alteration impairs barrier function and favours opportunistic "passenger" bacteria, placing selective pressure on the microbiome (4).

Previous research showed that COX-2 expression was increased in CRC (5), causing proliferation, enhanced angiogenesis and apoptosis suppression (6). In this way, COX-2 inhibitors such as celecoxib are among the most promising chemopreventive agents for CRC (7). Despite these benefits, their long-term use may shift cardiovascular homeostasis, inhibiting biosynthesis of vascular COX-2 dependent prostacyclin (PGI_2) (8) and increasing cardiovascular events risk (9).

When colon-targeted drugs are orally administered, they encounter the gut microbiota before reaching host tissues. Microbial drug metabolism may impact drug stability and activity, influencing toxicity and inflammation on the host (10). Conversely, drugs may also alter the gut microbiota, modulating community structure, and potentially resulting in dysbiosis (11). Understanding microbiome-drug-host interactions will be crucial to assess the potential impact on ADME and safety responses of intestinal targets. Although the gut microbiome plays an enormous role in patient health, and in pharmacokinetic and pharmacodynamic response (12), pharmaceutical companies have only recently started to explore the microbiome as potential contributor to drug outcomes. Similarly, regulatory agencies are not currently considering gut bacteria within the official evaluations before approval of colon-targeted drugs. Besides the microbial modulation of drug

bioavailability and pharmacokinetics (10), low solubility and the small aqueous volume

in the colon may additionally hinder drug efficacy (13). As interest in developing novel

colon-targeted drug delivery systems for CRC has increased, insight into the presystemic

biotransformation by the gut microbiota is fundamental. *In vitro* studies have shown that

bacteria can metabolize celecoxib (14), potentially jeopardizing its anti-inflammatory

effects. Additionally, whether celecoxib presence can modulate the host-microbiome

crosstalk remains unresolved. Prior reports correlating gut microbiome changes with

celecoxib treatment relied on DNA amplicon sequencing (15). As the presence of dead

bacterial cells may obscure the interpretation from microbiome data, we analysed the

bacterial community composition based on cDNA, as a proxy for the metabolically active

community.

Colonic transit times (17-24 hrs) (16, 17) are longer compared to those of the small

intestine (3-4 hrs), and overall systemic absorption of poorly soluble drugs may be

improved with colon-delivery systems (16). Although celecoxib can be absorbed

throughout the GI tract, we simulated delivery in a proximal colon-targeted formulation

because of the environment vicinity between proximal colon and small intestine. Thus,

the aim of this research was to screen the short-term two-way interactions between

celecoxib and microbiome of the *in vitro* proximal colon, and to elucidate the ultimate

impact of the microbiota on the anti-inflammatory efficacy of celecoxib.

RESULTS

Celecoxib impacts bacterial fermentative metabolism in a donor-dependent manner.

Short-chain Fatty Acids (SCFAs) were used to monitor functional activities of the colon

communities following exposure to celecoxib (CX) or its vehicle (PEG). Total SCFA

production from separate faecal microbiota incubations of 8 different individuals was

significantly lower for CX incubations compared to those with PEG ($P < 0.05$, Table 1),

but pH remained stable throughout the experiment (Table 1). While the 60:20:20 ratio of

the three major SCFA (acetate, propionate, and butyrate) was consistent across donors at

the initial time point (Supplementary Table 1A), a shift was evident at the end of the

incubation. Both relative and absolute concentrations of butyrate and propionate were

higher in samples treated with only PEG compared to those with CX. In fact, celecoxib

significantly lowered butyrate after 16 hours ($P < 0.05$, Table 1), except in donors 1, 4

and 8 (Supplementary Table 1B). In contrast, total active bacterial load remained

unchanged when celecoxib was supplemented (Table 1, Supplementary Table 2). These

findings might indicate that bacteria either alter their metabolic activity when exposed to

CX or that changes in bacterial composition lead to shifts in fermentation profiles.

Bacterial abundance remains stable, but microbiome diversity and composition

shifted following celecoxib supplementation

Quantitative RT-PCR revealed that total 16S rRNA gene copy numbers from

metabolically active bacteria were not significantly different between treatments, across

donors (Supplementary Table 2). In contrast, community composition and structure were

influenced, as celecoxib was associated with reduced diversity at the end of the incubation

(Figure 1, Table 2A and 2B), with significant donor effect (PERMANOVA $P < 0.001$,

Figure 2). CX shifted alpha diversity and community composition metrics across donors

(Supplementary Table 3, Figure 3). Although interindividual variability was significant,

metabolically active taxa with the highest relative abundances in control incubations

were: *Bacteroides* sp., *Faecalibacterium* sp., Ruminococcaceae UGC002, and

Megasphaera sp. In contrast, *Alistipes* sp., *Dialister* sp., *Escherichia-Shigella* sp.,

Lachnospira sp., *Megamonas* sp., *Parabacteroides* sp., *Roseburia* sp., and *Sutterella* sp.

were highly abundant when CX was supplied (Figure 3).

Short-term exposure to celecoxib increases number of community metabolic networks

Association networks were constructed to highlight potential shifts among bacterial relative abundances after short-term exposure to CX or its vehicle. The number of links in the network was lower when the vehicle PEG alone was provided (Figure 4A), but the central position of *Phascolarctobacterium*, *Parabacteroides*, *Coprococcus* 2 and Unclassified Victivallales and the linkages among these genera suggest concomitant changes in their relative abundances in presence of PEG. The central position of the acetate-producing *Erysipelatoclostridium* in the celecoxib network (Figure 4B) indicated that its relative abundance was associated with *Subdoligranulum*, a butyrate and lactate-producing genus, in a donor-dependent manner ($P < 0.05$, Supplementary Table 4).

Networks connecting bacterial load, metabolites and relative abundances were assembled to monitor functional alterations associated with changes in genera abundances. Activity-based relative abundance of *Bacteroides* was positively associated with butyrate, propionate and valerate levels and total 16S rRNA gene copy number in presence of PEG (Figure 5A, Supplementary Table 4). These results confirmed that *Bacteroides* was one of the most abundant active genera, as first suggested in Figure 3. Higher number of links in comparison with PEG were detected after 16 h of exposure to celecoxib (Figure 5). Indeed, one network linking acetate with the relative abundance of diverse taxa was only present when celecoxib was supplied and contained the largest number of active genera participating. Relative abundance of *Faecalibacterium* decreased over time with celecoxib supplementation, corresponding with the decline in butyrate levels (Figure 3). However, this genus tended to be positively associated with acetate production (Figure 5B). Relative abundance of active *Lachnoclostridium* was significantly associated with butyrate in the celecoxib treatment (Figure 5B). Here, *Oscillospira*, *Allisonella*,

Hungatella and *Streptococcus* were linked because they possibly participated in butyrate

production as well. *Megamonas* and *Megasphaera* were other genera with high relative

abundances positively associated with branched-chain SCFA and isovalerate. The

associations between the latter metabolites and *Colinsella* remained unchanged upon

celecoxib exposure, suggesting that this community function is either independent from

drug supplementation or directly associated with PEG addition (Figure 5A and 5B).

Similarly, associations between *Allisonella*, *Streptococcus* and butyrate were common for

both treatments. Results from the network analysis indicate that community metabolic

activity shifts upon short-term supplementation of celecoxib, potentially impacting

butyrate production.

Inter-individual differences play a role in reduction of *in vitro* inflammation when

celecoxib is presented to the microbiome.

Direct contact of macrophage-like cells with filter-sterilized supernatants from the CX-

exposed microbiota resulted in a variable outcome in IL-8 and CXCL16 concentrations.

Compared with the PEG controls, a decrease in IL-8 and CXCL16 was only observed in

4 out of 8 individuals (Figure 6A and 6B).

High IL-8 concentrations were found when non-bacterially exposed celecoxib was

applied in the enterohepatic model (Figure 6C). In contrast, the lowest concentration of

IL-8 and thus the most significant decrease in inflammatory response was found with the

celecoxib-exposed supernatant of donor 6 ($P < 0.05$, Table 3). Celecoxib that was not in

contact with bacteria promoted the highest decrease in transepithelial electrical resistance

(TEER) ($P < 0.05$, Table 3), in comparison with supernatants from celecoxib-exposed

bacteria. This outcome indicates less damage to epithelial barrier function and decreased

inflammatory response when gut microbiota is exposed to celecoxib.

DISCUSSION

As the lipophilic characteristics of celecoxib favour metabolic elimination (18), exploring only the pathways for celecoxib distribution in the host is insufficient for assessing drug bioavailability. *In vitro* models have been applied to simulate drug-bug associations in the human gut (19, 20), but the impact of NSAIDs on the metabolic activities of the microbiome has been overlooked. Previous studies reported the influence of different NSAIDs on the gut microbiome (21), yet no changes in bacterial richness, microbiome composition or beta diversity were found when celecoxib was supplied (15). These results focused on screening community structure using DNA-based amplicon sequencing, which may also include DNA of dead bacterial cells. As such approach may deliver a partial overview, we used cDNA as an indicator of the metabolically active bacterial population and we performed community composition analyses based on information from transcribed RNA (22). In this way, we intended to highlight the main bacterial genera whose metabolic activities were impacted by celecoxib in the RNA pool. Activity-based cell load was not significantly different, and the only alterations were on the bacterial metabolic activities. Although composition, diversity and richness were not significantly different as result of the treatment, differences in the most abundant genera between PEG and celecoxib were observed. PERMANOVA analysis suggested that donor was the main driver for dissimilarities in community structure.

The shifts in microbial metabolism observed in our study may indicate that celecoxib can inhibit functional activities of bacterial groups, rather than of single genera, ultimately impacting microbial fermentation (23). This was confirmed by the total active bacterial load, which remained unchanged between treatments. Celecoxib users have been reported to have enrichment of Enterobacteriaceae and Acidaminococcaceae (21). Here, we

validated that the relative abundance of transcriptionally active *Escherichia-Shigella*

(Enterobacteriaceae) was increased with celecoxib supplementation, whereas the relative

abundance of active *Acidaminococcus* (Acidaminococcaceae) was positively associated

with acetate in the same treatment. These results highlight the applicability of our *in vitro*

model used for this study. Coriobacteriaceae (*Colinsella*) has been proposed to participate

in xenobiotic metabolism by decreasing levels of serine and glycine (23), which can be

both fermented to acetate in the gut (24). We observed that the relative abundance of this

genus was positively associated with isovalerate levels when celecoxib was added. As

indicated in the corresponding KEGG module, *Colinsella aerofaciens* has the capacity of

using isovalerate for synthesising leucin and serine (25), confirming its potential

xenobiotic role. On the other hand, the capacity of *Alistipes* for producing succinate (26),

may explain its association with propionate in the PEG treatment. *Dialister* is another

succinate producer (27) present at high relative abundance, but it does not seem to

participate in the propionate network when PEG was supplied. This may suggest that,

even based on cDNA, high relative abundance may not be an indicator of metabolic

activity but just of maintenance metabolism.

The significantly lower butyrate concentration post-celecoxib delivery may have several

implications. Butyrate has tumour inhibitory properties (28) but its paradoxical

correlation with polyp formation (29, 30) suggests that the CRC-butyrate interplay is not

just a direct effect of altered SCFAs concentrations, and may also be associated with

changes in microbiome functionality. Indeed, the protecting properties of celecoxib

towards CRC (31, 32) may be partially derived from lowering bacteria-derived butyrate.

Nutrients can be utilized in their final form by both host and gut microbes, leading to

competition for resources (33). In fat, blocking fatty acid supply and host recapture has

been proposed as strategy for influencing cancer cell bioenergetics (34, 35), but few

studies have addressed the potential role of acetate in cancer (36). Acetyl-CoA is the main

carbon source in mammalian cells, and the ligation of acetate and CoA by acetyl-CoA

synthetase (ACSS) contributes to the supply of this metabolite (36). Cancer cells use

acetyl Co-A during aerobic metabolism regardless of stress conditions (36). Butyrate is

also metabolised to acetyl-CoA (28), inhibiting tumour growth via silencing acetyl-CoA

synthetases (34) or potentially through acetate uptake. As acetyl-CoA is a bacterial

substrate for fatty acid production (37), our network analysis suggests the use of acetate

for bacterial metabolic functions when celecoxib was supplied. Thus, celecoxib may

favour bacterial competition for acetyl Co-A, generating potential host protective effects.

Indeed, microbiota-derived acetate has been associated with inhibition of cancer-

associated deacetylation (38). Increasing acetate concentration shifts SCFA ratios (39),

decreasing butyrate and explaining why the acetate:propionate:butyrate ratios tended to

change when celecoxib was supplied. *Faecalibacterium* sp. is a well-known butyrate

producer and acetate consumer (39) and its presence in the acetate network may indicate

that the relative abundance of this genus increased when celecoxib was provided and

acetate levels increased.

Synthesis of reduced products including H_2, succinate, and butyrate is used to create redox

balance during colonic fermentation, whereas formation of oxidised products, such as

acetate, is associated with bacterial ATP production (37). As ethanol, acetate, CO_2 and

H_2 are the major products of *Lachnoclostridium* (40, 41), its central position in the

celecoxib network suggests that this genus was highly interactive with other genera,

despite not being highly relatively abundant. Because of its acetate-producing capacity,

it may cross-feed other bacteria, explaining its positive association with the total 16s

rRNA gene copy number. Acetate is a co-substrate for butyrate production (42),

contributing to maintain gut redox balance and explaining the significant association

between acetate-producing *Lachnoclostridum* and butyrate. Indeed, *Oscillospira* and

Lachnoclostridium may have been involved in butyrate production in the celecoxib

treatment, as they both possess transporters involved in uptake of oligosaccharides and

glycerol-P (43). When NADH is consumed during H_2 formation in glycolysis, it is not

available to convert acetoacetyl-CoA to butyryl-CoA and butyrate production may reduce

(37). Acetyl Co-A is consequently routed into acetate formation (42), which is

energetically advantageous for bacterial use (37). In fact, butyrate producers may shift

metabolic pathways towards acetate under stress conditions (44). As in our study, other

acetate producers such as *Bacteroides* sp. and *Prevotella* sp. positively correlated to

butyrate-producing   communities   expressing   the   butyryl-CoA:acetateCoA-

transferase (*but*) gene, supporting the observed *but*-acetate link (45).

Although the role of SCFA on inflammation and CRC progression has been documented,

most of the studies lack natural commensal microbiota cross-talking with host cells (28,

30). As interindividual differences in microbiome play a crucial role on drug responses

(11), elucidating whether microbiome fermentation products synergistically act with

NSAIDs for impacting inflammation is paramount. Celecoxib was responsible for only

5% of the variance in bacterial community composition among the 8 donors, but the effect

of celecoxib-exposed bacteria on the proinflammatory IL-8 was sufficient to be detected

in a donor-dependent manner. Moreover, samples derived from 50% of the donors

showed increased CXCL-16, which is a marker of favourable prognosis in CRC (46).

This suggests that the interaction between celecoxib and microbiome to resolve

inflammation may be not solely related to community composition, and functionality

must be accounted for. While butyrate seemed to be reduced with short-term treatment,

Acetate:Propionate:Butyrate ratio and relative levels of acetate (of the total SCFA) may

be other functional bacterial metrics to be considered for assessing successful reduction

on inflammation when NSAIDs are administered.

As indicated in our study, gut microbiome activities and its interplay with the host can be

modulated through celecoxib. Our *in vitro* model retained interindividual variability,

allowing for mechanistic research on drug-microbiota-host interplay, fundamental for

developing individualised therapeutic strategies. Divergence in the molecular approaches

(i.e. next generation sequencing, qPCR) used for investigating CRC-associated

microbiome signatures contributes to the lack of consensus on the composition of

microbiota in CRC patients. Information on the role of SCFA in cancer signalling remains

inconclusive, and thus, future studies assessing the crosstalk of the microbiota and colon

cells in CRC are imperative. Certainly, individual microbiome may need to be considered

before supplementing celecoxib as a colon targeted drug for CRC. However, focusing on

functionality rather than taxonomic diversity may be the next frontier for developing

predictive markers of CRC-induced dysbiosis and NSAID efficacy.

MATERIALS AND METHODS

Faecal incubations simulating the proximal colon environment

Batch incubations of faecal samples were conducted using phosphate-buffered nutritional

medium (pH 5.9, 0.1 M) containing (per litre): 12.25 g of KH_2PO_4 (Carl Roth GmbH,

Karlsruhe, Germany), 1.78 g of Na_2HPO_4 (Carl Roth GmbH, Karlsruhe, Germany), 3 g

of yeast extract (Oxoid Ltd, Basingstoke, Hampshire, England), 1 g of proteose peptone

(Oxoid Ltd, Basingstoke, Hampshire, England), 0.25 g of gum Arabic from acacia tree

(Sigma Aldrich Co., St-Louis, USA), 0.5 g of apple pectin (Sigma Aldrich Co., St-Louis,

USA), 0.25 g of xylan (Carl Roth GmbH, Karlsruhe, Germany), 1 g of potato starch

(Prodigest, Zwijnaarde, Belgium), and 4 g of mucin from porcine stomach Type II (Sigma

Aldrich Co., St-Louis, USA).

Faecal material was obtained from eight healthy, non-obese, non-smoking volunteers (4M

and 4F, 30 ± 2.8 yo), following an omnivorous diet and without antibiotic exposure for

the prior 6 months. Stool samples were collected and prepared as previously described

(47). Celecoxib (Janssen Pharmaceutica, Beerse, Belgium) was dissolved in 70:30 (v/v)

PEG_{400}:H_2O as vehicle, at a stock concentration of 100 mg/mL. For each donor, triplicate

incubations of 10% faecal slurry, 100 mg/mL of celecoxib or 70:30 (v/v) PEG_{400} (Acros,

Geel, Belgium):H_2O, and nutritional medium (final volume = 175 mL) were performed

for 16 hours at 37°C and 120 rpm (New Brunswick Scientific InnOva 4080 Incubator

Shaker, Wezembeek-Oppem, Belgium), under anaerobic conditions (90%-10% N_2/CO_2).

Once the incubation was completed, samples were collected, flash-frozen using liquid

nitrogen and preserved at -80 °C for further analysis.

Surveying community functionality and composition

Volatile Fatty Acids (VFAs) production. VFA were used as benchmarks of community

activity and extracted as per Truchado et al. (48). Differences in VFA concentrations

among treatments were compared using a repeated measures mixed model (49), with the

lsmeans adjustment and Bonferroni correction for multiple comparisons (SAS, 2012).

Statistical significance was assumed at $P < 0.05$.

Quantification of total metabolically active bacterial community. Samples were

thawed on ice, centrifuged for 5 minutes at 4°C and $5000 \times g$, and disrupted two times

for 25 seconds at 1400 rpm (Power Lyser 24, MO BIO Laboratories, Carlsbad, USA),

with a 5-second interval, where the samples were placed on ice. RNA was further

extracted using the Nucleospin RNA plus kit (Macherey Nagel, Düren, Germany),

following the manufacturer's instructions. Reverse transcription was completed using the

Reverse Transcriptase Core kit (Eurogentec, Seraing, Belgium).

For the quantification of total bacterial 16S rRNA gene copy numbers, a standard curve

was constructed using serial dilutions of plasmid DNA from a clone identified

as *Bifidobacterium breve* LMG 33264. Briefly, universal bacterial primers 27F and

1492R were used to amplify the full-length 16S rRNA gene from the plasmid DNA of

a *Bifidobacterium breve* clone. The resultant PCR product was purified using the using

the InnuPREP PCRpure Kit (AJ innuscreen, Berlin, Germany), cloned in *E. coli* with the

TA Cloning Kit (Life Technologies, Carlsbad, VS) and plasmid extracted with the

PureYield Plasmid Miniprep System (Promega, Leiden, The Netherlands). The mass

concentration of the plasmid was measured using an ND-1000 spectrophotometer

(NanoDrop Technologies, Wilmington, DE), converted to the molecule concentration

(50) and copy numbers of total bacteria in 50 ng of cDNA and per ml of faecal incubation

were determined by relating the threshold cycle (C_T) values to the standard curves based

on the following equation: $Y = -3.193 \times \log X + 35.003$ (Y, C_T value; *X*, copy number of

16S rRNA gene) ($r^2 = 0.996$) (50).

Quantitative reverse transcription-PCR (qRT-PCR) analysis of the metabolically active

community was achieved with the StepOnePlus Real-Time PCR system (Applied

Biosystems, Foster City, CA USA). Amplification reactions were in: 10× Taq buffer with

KCl, 25 mM of $MgCl_2$, 10 mM of dNTP mix, 10 µM of each forward and reverse primers,

20 mg/mL of BSA (Thermo Scientific, Waltham, USA), 1.25 U of Taq Polymerase, 50

ng of cDNA and 0.1× SYBR Green I ([Invitrogen, Carlsbad, USA] provided at 10 000×,

stock solutions of 20× were prepared in DMSO). qRT-PCR conditions included 10

minutes at 95°C, followed by 40 cycles of 15 s at 95°C and 30 s at 60°C. The efficiencies

(*E*) of RT-PCR were calculated from the given slopes in StepOnePlus software using the

following equation: $E = [10^{(-1/\text{slope})} - 1] \times 100\%$. Data generated from reactions with

efficiencies between 90 and 110% were used for further analysis (50).

Community composition and dynamics.

The V5-V6 hypervariable region of the 16S rRNA gene from the cDNA was amplified

using primers 341F and 785R. High-throughput amplicon sequencing was performed

with the Illumina MiSeq platform according to the manufacturer's guidelines at LGC

Genomics GmbH (Berlin, Germany). Library preparation and purification as well as

bioinformatic processing are described in the supplemental material.

Data was imported into R using *phyloseq* (51) and taxon abundances were rescaled by

calculating the taxon proportions and multiplying them by the minimum sample size (n =

2201) present in the data set (52). Inverse Simpson was the metric used for assessing

alpha diversity and Pielou's index was used as indicator of community evenness (53).

Differences in alpha diversity and evenness measures were compared among treatments

using a repeated measures mixed model in SAS (version 9.4, SAS Institute, Cary, USA).

Beta diversity based on Chao and Bray-Curtis indices was used to examine dissimilarity

and determine the impact of treatment and time on microbial community structure (49).

PCoA was employed to visualize the differences among samples, using the vegan

package in R (54). Stratified permutational multivariate analysis of variance

(PERMANOVA) with 999 permutations was conducted to indicate the significance of

each covariate (time or treatment) on the microbial community.

ANOVA was applied to reveal whether the distribution of the genera was different

between treatments over time (54). Because of the over-dispersion in the OTU data, a

zero-inflated negative binomial distribution count model was used to assess the effect of

donor, time and treatment and the interactions between donor*time*treatment on each

individual genus. The model was selected based on the Akaike Information Criterion

(AIC). Differences among library size sample were accounted for with the offset option

in proc GLIMMIX in SAS (49).

Sparse inverse covariance selection was employed to infer an association network within

treatments, using the package SpiecEasi (v0.1.2) (55) in R (v3.5.1). Neighbourhood

selection method (S-E(MB)) was used to construct a network of bacterial genera

displaying core networks with either functional or phylogenetic similarities. The

calculated adjacency matrix was examined with the software Gephi 0.9.2 (56), using the

Fruchterman-Reingold algorithm for visualizing significant correlations (R ≥ 0.7). This

method computes a layout in which the length of the connections indicates the absolute

value of the correlation. The nodes representing genera whose relative bacterial

abundances have several significant correlations among them were placed closer to each

other in the network (47).

Bipartite networks highlighted functional associations among bacterial genera and

metabolites (49), using a pair-wise similarity matrix obtained from a Regularised

Canonical Correlation Analysis (57). Values of the similarity matrix were computed as

the correlation between relative abundances of bacterial genera and metabolic variables,

projected onto the space spanned by the first 3 components retained in the analysis

(r ≥ 0.75). Genera in the plot were close to correlated variables in the treatment where

they were more abundant (49).

Assessing the anti-inflammatory potential of bacteria-exposed celecoxib

Simplified inflammation model. The human monocytic leukaemia cell line THP-1

(THP-1, ECACC 88081201, Public Health England, UK) was grown in Roswell Park

Memorial Institute (RPMI) 1640 culture medium (Thermo Fisher Scientific, Waltham,

USA) supplemented with 10% heat-inactivated foetal bovine serum (iFBS) (Greiner Bio-One, Wemmel, Belgium) and 1% antibiotic-antimycotic solution (10,000 units/mL of penicillin, 10,000 µg/mL of streptomycin, and 25 µg/mL of Amphotericin B) (Thermo Fisher Scientific, Waltham, USA), at 37°C in 10 % CO_2 in a humidified incubator. THP-1 cells were sub-cultured twice a week and used between passage 20 and 30.

Macrophage-like phenotype was obtained by treating cells with 25 ng/mL phorbol 12-myristate 13-acetate (PMA, Sigma Aldrich, St-Louis, USA) for 48 h in 24-wells culture plates (Corning, Sigma Aldrich, St-Louis, USA) at a density of 1 x 10^5 cells/cm^2. Differentiated, plastic-adherent cells were washed twice with culture medium (supplemented RPMI 1640 without PMA) and incubated for 24 h. Medium was removed and replaced by supernatant from the eight incubations, passed through a sterile 0.2 µm filter and diluted five times with RPMI 1640 without supplementation. Celecoxib solution at 100 mg/mL dissolved in 70:30 (v/v) PEG:H_2O or 70:30 (v/v) PEG:H_2O alone were diluted 1:5 with medium and added as controls of non-bacteria exposed celecoxib and vehicle. This dilution step was performed to simulate low solubility and the small aqueous volume in the colon (13). LPS (100 ng/mL) dissolved in RPMI 1640 was added to the cells to simulate systemic inflammation, as in CRC. The control used in all measurements was PMA-differentiated THP-1 macrophages exposed to the culture medium.

Development of an *in vitro* model of the enterohepatic system. A novel model including intestinal epithelial (Caco-2, ECACC 86010202), goblet (HT29-MTX-E12 ECACC 12040401, Public Health England, UK), hepatic (HepG2 ATCC HB-8065, Molsheim, France), and immune-like cells (THP-1, ECACC 88081201, Public Health England, UK) was developed to study host-drug-microbiome interactions (58, 59). Cell layers were grown in a double-chamber insert (Corning, Sigma Aldrich, St-Louis, USA), and maintained for 23 days. Briefly, 100 µl of a THP-1 cell suspension (10^6 cells/mL) in

20 mg/mL collagen from rat tail type I (Sigma Aldrich, St. Louis, USA) were spread in the basolateral surface of inverted inserts. After 45 min of incubation at 37ºC and 10% CO_2, a 9:1 cell suspension of Caco-2/HT29-MTX cells (density of 7.5×10^4 cells/cm^2) was added to the apical side of the insert in 1.5 ml of Dulbecco's Modified Eagle's medium (DMEM) supplemented with 10% (v/v) of iFBS and 1% of antibiotic-antimycotic solution (v/v). The basolateral compartment of the double-chamber system was filled with 2 ml of supplemented RPMI 1640, and the medium was refreshed every 48 h. Inserts were transferred to 6-well plates with pre-seeded HepG2 cells at day 19 post-seeding, when the cell culture media in the basal compartment was replaced by RPMI 1640, and the co-culture was maintained 24 h prior to the assay. HepG2 cells were seeded at a density of 10^5 cells/cm^2 and maintained 5 days in supplemented minimal essential medium (MEM) supplemented with 10% (v/v) of iFBS and 1% (v/v) of antibiotic-antimycotic solution. Five days prior the assay all the cell culture media was depleted of antibiotic/antimycotic solution. Supernatants from volunteers (2F, 29 yo) showing extreme microbiota metabolism responses in the short-term incubations were used for validating whether celecoxib effectively decreases inflammation even after exposure to the gut microbiota. Volunteer 6 displayed the most significant decrease in butyrate after 16 h of incubation, while volunteer 8 did not present any changes. Triplicate supernatants from each volunteer were passed through a sterile 0.2 µm filter, diluted five times with DMEM without supplementation, pH was adjusted to pH 7-7.2 with filter-sterilized NaOH 0.5M (Carl Roth GmbH & Co), and placed in the apical compartment of the model. Results from the simplified inflammation model indicated that there were no significant differences between control and treatment among donors. Thus, we only included celecoxib solution dissolved in 70:30 (v/v) PEG:H_2O or 70:30 (v/v) PEG:H_2O alone in the enterohepatic model. Celecoxib solution at 100 mg/mL and a 70:30 (v/v) PEG:H_2O

were diluted 1:5 with DMEM without supplementation, and added to the apical

compartment, as controls of non-bacterially modified celecoxib and vehicle. This dilution

step was performed to simulate low solubility and the small aqueous volume in the colon

(13). LPS (100 ng/mL) was added to the basal compartment to simulate systemic

inflammation. Plates were incubated for 16 h at 37°C, 10% CO_2 and 95% humidity. TEER

was evaluated with the Millicell ERS-2 Voltohmmeter at the beginning and at the end of

the assay (Merck-Millipore, Overijse, Belgium).

Evaluating inflammatory response. Concentration of IL-8 and CXCL16 were measured

in the simplified inflammation model with the Human IL-8 (CXCL-8) and Human

CXCL16 Mini ABTS ELISA Development Tests (PeproTech, London, UK) and the

ABTS ELISA Buffer Kit (PeproTech, London, UK), following the manufacturer's

instructions. Samples were not diluted, and determination was performed at 405 nm, with

correction for 650 nm (Infinite 200 PRO, Tecan, Männedorf, Switzerland). IL-8 was

additionally determined in the presystemic metabolism model, using samples from the

basal compartment diluted ten times with PBS (ABTS ELISA Buffer Kit, PeproTech,

London, UK), and measured as indicated above.

Data Availability

Raw 16S rRNA reads have been made available on the SRA under accession number ID

PRJNA540406. Original R scripts are available in GitHub

(https://github.ugent.be/ehernand/CX_sequencing).

ACKNOWLEDGEMENTS

E.H.-S. is a postdoctoral fellow supported by Flanders Innovation and Entrepreneurship

(Agentschap Innoveren & Ondernemen). M.C. was supported by an FWO postdoctoral

fellowship (FWO/12R2717N). E.H.-S. and T.Vd.W. conceived the simulations of the

colon environment. E.H.-S. and M.C. devised cell models. J.S. and L.L. supplied material

and advised on the use of celecoxib. E.H.-S., E.H., and M.C. conducted all laboratory

work. R.P. completed the amplicon sequencing library pre-processing and guided

sequencing data mining. E.H.-S. performed the statistical analyses and interpretation of

all data generated, produced figures and tables, and wrote the manuscript. All authors

read and approved the manuscript. The authors gratefully acknowledge the volunteers

participating in this study. We thank Dr. Davide Gottardi for his scientific advice and Dr.

Racha El Hage, Fabian Mermans, Olivier Dezutter, and Dr. Chiara Ilgrande for their

technical assistance. The authors declare that they have no competing interests.

REFERENCES

1. Kumar M, Cash BD. 2017. Screening and surveillance of colorectal cancer using CT colonography. Current treatment options in gastroenterology 15:168-183.
2. Arnold M, Sierra MS, Laversanne M, Soerjomataram I, Jemal A, Bray F. 2017. Global patterns and trends in colorectal cancer incidence and mortality. Gut 66:683-691.
3. Johnson CH, Spilker ME, Goetz L, Peterson SN, Siuzdak G. 2016. Metabolite and microbiome interplay in cancer immunotherapy. Cancer research 76:6146-6152.
4. Tjalsma H, Boleij A, Marchesi JR, Dutilh BE. 2012. A bacterial driver–passenger model for colorectal cancer: beyond the usual suspects. Nature Reviews Microbiology 10:575.
5. Bhala N, Emberson J, Merhi A, Abramson S, Arber N, Baron J, Bombardier C, Cannon C, Farkouh M, FitzGerald G. 2013. Vascular and upper gastrointestinal effects of non-steroidal anti-inflammatory drugs: meta-analyses of individual participant data from randomised trials. Elsevier.
6. Kawanishi S, Ohnishi S, Ma N, Hiraku Y, Murata M. 2017. Crosstalk between DNA damage and inflammation in the multiple steps of carcinogenesis. International journal of molecular sciences 18:1808.
7. Wang D, DuBois RN. 2010. The role of COX-2 in intestinal inflammation and colorectal cancer. Oncogene 29:781.
8. Funk CD, FitzGerald GA. 2007. COX-2 inhibitors and cardiovascular risk. Journal of cardiovascular pharmacology 50:470-479.
9. Cuzick J, Otto F, Baron JA, Brown PH, Burn J, Greenwald P, Jankowski J, La Vecchia C, Meyskens F, Senn HJ. 2009. Aspirin and non-steroidal anti-inflammatory drugs for cancer prevention: an international consensus statement. The lancet oncology 10:501-507.
10. Spanogiannopoulos P, Bess EN, Carmody RN, Turnbaugh PJ. 2016. The microbial pharmacists within us: a metagenomic view of xenobiotic metabolism. Nature Reviews Microbiology 14:273.

11.   Wilson ID, Nicholson JK. 2017. Gut microbiome interactions with drug
metabolism, efficacy, and toxicity. Translational Research 179:204-222.
12.   Sousa T, Paterson R, Moore V, Carlsson A, Abrahamsson B, Basit AW. 2008.
The gastrointestinal microbiota as a site for the biotransformation of drugs.
International journal of pharmaceutics 363:1-25.
13.   Philip AK, Philip B. 2010. Colon targeted drug delivery systems: a review on
primary and novel approaches. Oman medical journal 25:79.
14.   Srisailam K, Veeresham C. 2010. Biotransformation of celecoxib using microbial
cultures. Applied biochemistry and biotechnology 160:2075-2089.
15.   Bokulich NA, Battaglia T, Aleman JO, Walker J, Blaser MJ, Holt PR. 2016.
Celecoxib does not alter intestinal microbiome in a longitudinal diet-controlled
study. Clinical Microbiology and Infection 22:464-465.
16.   HARDY JG, WILSON CG, WOOD E. 1985. Drug delivery to the proximal colon.
Journal of Pharmacy and Pharmacology 37:874-877.
17.   Macfarlane G, Macfarlane S, Gibson G. 1998. Validation of a three-stage
compound continuous culture system for investigating the effect of retention time
on the ecology and metabolism of bacteria in the human colon. Microbial ecology
35:180-187.
18.   Paulson SK, Vaughn MB, Jessen SM, Lawal Y, Gresk CJ, Yan B, Maziasz TJ,
Cook CS, Karim A. 2001. Pharmacokinetics of celecoxib after oral administration
in dogs and humans: effect of food and site of absorption. Journal of
Pharmacology and Experimental Therapeutics 297:638-645.
19.   McConnell EL, Fadda HM, Basit AW. 2008. Gut instincts: explorations in
intestinal physiology and drug delivery. International journal of pharmaceutics
364:213-226.
20.   Esch EW, Bahinski A, Huh D. 2015. Organs-on-chips at the frontiers of drug
discovery. Nature reviews Drug discovery 14:248.
21.   Rogers MA, Aronoff DM. 2016. The influence of non-steroidal anti-inflammatory
drugs on the gut microbiome. Clinical Microbiology and Infection 22:178. e1-
178. e9.
22.   Blazewicz SJ, Barnard RL, Daly RA, Firestone MK. 2013. Evaluating rRNA as
an indicator of microbial activity in environmental communities: limitations and
uses. The ISME journal 7:2061.
23.   Montrose DC, Zhou XK, McNally EM, Sue E, Yantiss RK, Gross SS, Leve ND,
Karoly ED, Suen C, Ling L. 2016. Celecoxib alters the intestinal microbiota and
metabolome in association with reducing polyp burden. Cancer Prevention
Research:canprevres. 0095.2016.
24.   Smith EA, Macfarlane G. 1997. Dissimilatory amino acid metabolism in human
colonic bacteria. Anaerobe 3:327-337.
25.   Okuda S, Yamada T, Hamajima M, Itoh M, Katayama T, Bork P, Goto S,
Kanehisa M. 2008. KEGG Atlas mapping for global analysis of metabolic
pathways. Nucleic acids research 36:W423-W426.
26.   Rautio M, Eerola E, Vaisanen-Tunkelrott M-L, Molitoris D. 2003.
Reclassification of bacteroides putredinis (Weinberg et al., 1937) in a new genus
alistipes gen. nov., as alistipes purtredinis comb. nov., and description of alistipes
finegoldii sp. nov., from human sources. Systematic and applied microbiology
26:182.
27.   Morotomi M, Nagai F, Sakon H, Tanaka R. 2008. Dialister succinatiphilus sp.
nov. and Barnesiella intestinihominis sp. nov., isolated from human faeces.
International journal of systematic and evolutionary microbiology 58:2716-2720.

28.   Donohoe DR, Collins LB, Wali A, Bigler R, Sun W, Bultman SJ. 2012. The
554   Warburg effect dictates the mechanism of butyrate-mediated histone acetylation
555   and cell proliferation. Molecular cell 48:612-626.
556   29.   Belcheva A, Irrazabal T, Robertson SJ, Streutker C, Maughan H, Rubino S,
557   Moriyama EH, Copeland JK, Surendra A, Kumar S. 2014. Gut microbial
558   metabolism drives transformation of MSH2-deficient colon epithelial cells. Cell
559   158:288-299.
560   30.   Kang M, Martin A. Microbiome and colorectal cancer: Unraveling host-
561   microbiota interactions in colitis-associated colorectal cancer development, p. *In*
562   (ed), Elsevier,
563   31.   Arber N, Eagle CJ, Spicak J, Rácz I, Dite P, Hajer J, Zavoral M, Lechuga MJ,
564   Gerletti P, Tang J. 2006. Celecoxib for the prevention of colorectal adenomatous
565   polyps. New England Journal of Medicine 355:885-895.
566   32.   Umar A, Steele VE, Menter DG, Hawk ET. Mechanisms of nonsteroidal anti-
567   inflammatory drugs in cancer prevention, p 65-77. *In* (ed), Elsevier,
568   33.   Wasielewski H, Alcock J, Aktipis A. 2016. Resource conflict and cooperation
569   between human host and gut microbiota: implications for nutrition and health.
570   Annals of the New York Academy of Sciences 1372:20-28.
571   34.   Röhrig F, Schulze A. 2016. The multifaceted roles of fatty acid synthesis in
572   cancer. Nature Reviews Cancer 16:732.
573   35.   Schug ZT, Voorde JV, Gottlieb E. 2016. The metabolic fate of acetate in cancer.
574   Nature Reviews Cancer 16:708.
575   36.   Schug ZT, Peck B, Jones DT, Zhang Q, Grosskurth S, Alam IS, Goodwin LM,
576   Smethurst E, Mason S, Blyth K. 2015. Acetyl-CoA synthetase 2 promotes acetate
577   utilization and maintains cancer cell growth under metabolic stress. Cancer cell
578   27:57-71.
579   37.   Macfarlane S, Macfarlane GT. 2003. Regulation of short-chain fatty acid
580   production. Proceedings of the Nutrition Society 62:67-72.
581   38.   Wu W, Sun M, Chen F, Cao AT, Liu H, Zhao Y, Huang X, Xiao Y, Yao S, Zhao
582   Q. 2017. Microbiota metabolite short-chain fatty acid acetate promotes intestinal
583   IgA response to microbiota which is mediated by GPR43. Mucosal immunology
584   10:946.
585   39.   Duncan SH, Barcenilla A, Stewart CS, Pryde SE, Flint HJ. 2002. Acetate
586   utilization and butyryl coenzyme A (CoA): acetate-CoA transferase in butyrate-
587   producing bacteria from the human large intestine. Applied and environmental
588   microbiology 68:5186-5190.
589   40.   Yutin N, Galperin MY. 2013. A genomic update on clostridial phylogeny: Gram-
590   negative spore formers and other misplaced clostridia. Environmental
591   microbiology 15:2631-2641.
592   41.   Warnick TA, Methe BA, Leschine SB. 2002. Clostridium phytofermentans sp.
593   nov., a cellulolytic mesophile from forest soil. International journal of systematic
594   and evolutionary microbiology 52:1155-1160.
595   42.   Louis P, Flint HJ. 2017. Formation of propionate and butyrate by the human
596   colonic microbiota. Environmental microbiology 19:29-41.
597   43.   Gophna U, Konikoff T, Nielsen HB. 2017. Oscillospira and related bacteria–From
598   metagenomic species to metabolic features. Environmental microbiology 19:835-
599   841.
600   44.   Geirnaert A, Steyaert A, Eeckhaut V, Debruyne B, Arends JB, Van Immerseel F,
601   Boon N, Van de Wiele T. 2014. Butyricicoccus pullicaecorum, a butyrate

602		producer with probiotic potential, is intrinsically tolerant to stomach and small
603		intestine conditions. Anaerobe 30:70-74.
604	45.	Vital M, Gao J, Rizzo M, Harrison T, Tiedje JM. 2015. Diet is a major factor
605		governing the fecal butyrate-producing community structure across Mammalia,
606		Aves and Reptilia. The ISME journal 9:832.
607	46.	Kee J-Y, Ito A, Hojo S, Hashimoto I, Igarashi Y, Tsuneyama K, Tsukada K,
608		Irimura T, Shibahara N, Takasaki I. 2014. CXCL16 suppresses liver metastasis of
609		colorectal cancer by promoting TNF-α-induced apoptosis by tumor-associated
610		macrophages. BMC cancer 14:949.
611	47.	De Weirdt R, Hernandez-Sanabria E, Fievez V, Mees E, Geirnaert A, Van
612		Herreweghen F, Vilchez-Vargas R, Van den Abbeele P, Jauregui R, Pieper DH.
613		2017. Mucosa-associated biohydrogenating microbes protect the simulated colon
614		microbiome from stress associated with high concentrations of poly-unsaturated
615		fat. Environmental microbiology 19:722-739.
616	48.	Truchado P, Hernandez-Sanabria E, Salden BN, Van den Abbeele P, Vilchez-
617		Vargas R, Jauregui R, Pieper DH, Possemiers S, Van de Wiele T. 2017. Long
618		chain arabinoxylans shift the mucosa-associated microbiota in the proximal colon
619		of the simulator of the human intestinal microbial ecosystem (M-SHIME). Journal
620		of Functional Foods 32:226-237.
621	49.	El Hage R, Hernandez-Sanabria E, Calatayud Arroyo M, Props R, Van de Wiele
622		T. 2019. Propionate-producing consortium restores antibiotic-induced dysbiosis
623		in a dynamic in vitro model of the human intestinal microbial ecosystem. Frontiers
624		in Microbiology 10:1206.
625	50.	Hernandez-Sanabria E, Goonewardene LA, Wang Z, Zhou M, Moore SS. 2013.
626		Influence of sire breed on the interplay among rumen microbial populations
627		inhabiting the rumen liquid of the progeny in beef cattle. PloS one 8:e58461.
628	51.	McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible
629		interactive analysis and graphics of microbiome census data. PloS one 8:e61217.
630	52.	McMurdie PJ, Holmes S. 2014. Waste not, want not: why rarefying microbiome
631		data is inadmissible. PLoS computational biology 10:e1003531.
632	53.	Grunert O, Hernandez-Sanabria E, Vilchez-Vargas R, Jauregui R, Pieper DH,
633		Perneel M, Van Labeke M-C, Reheul D, Boon N. 2016. Mineral and organic
634		growing media have distinct community structure, stability and functionality in
635		soilless culture systems. Scientific reports 6:18837.
636	54.	Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara R, Simpson
637		GL, Solymos P, Stevens MHH, Wagner H. 2013. Package 'vegan'. Community
638		ecology package, version 2.
639	55.	Kurtz ZD, Müller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA. 2015.
640		Sparse and compositionally robust inference of microbial ecological networks.
641		PLoS computational biology 11:e1004226.
642	56.	Bastian M, Heymann S, Jacomy M. 2009. Gephi: an open source software for
643		exploring and manipulating networks. Icwsm 8:361-362.
644	57.	Lê Cao K-A, Boitard S, Besse P. 2011. Sparse PLS discriminant analysis:
645		biologically relevant feature selection and graphical displays for multiclass
646		problems. BMC bioinformatics 12:253.
647	58.	Paul W, Marta C. 2018. Resolving host–microbe interactions in the gut: the
648		promise of in vitro models to complement in vivo research. Current opinion in
649		microbiology 44:28-33.

650   59.   Calatayud MA, de Wiele Van T, Hernandez-Sanabria E. 2018. Assessing the
651   Viability of a Synthetic Bacterial Consortium on the In Vitro Gut Host-microbe
652   Interface. Journal of visualized experiments: JoVE.

 FIGURES

Figure 1. Inverse Simpson index (D_2) was used to assess alpha diversity dynamics at 0

and 16 h post treatment on each donor. Horizontal bars indicate means and distribution

of each data point is indicated.

Figure 2. Interindividual differences drive community composition. Principal coordinate

analysis (PCoA) indicated differences in community structure between donors. The

variance explained by the experimental factors ($P < 0.01$) was calculated using

PERMANOVA analysis and it is indicated on the top right.

Figure 3. Relative abundances of the bacterial genera occurring when celecoxib (CX) or

a control treatment (PEG) were supplied. The upper panel describes the community

composition at the beginning of the treatment (0 h), while the lower panel presents the 12

taxa with the highest activity-based (cDNA-based) relative abundances after 16 h of

treatment.

Figure 4. Bacterial network of communities exposed to short-term administration of

celecoxib (CX) or its vehicle (control treatment, PEG). (A) The faecal microbiota exposed

only to PEG displayed a lower number of edges or associations, while the network

architecture was enriched when CX was provided (B). This representation uncovers

relationships between clusters of genera across donors. Nodes representing bacterial

genera with similar abundances tended to cluster closely. Node diameter indicates the

relative abundance of each genus. Neighbourhood selection method (MB) was the

graphical model inference procedure selected. The associations described may be

considered 'core' functional relationships preserved across donors.

Figure 5: Community metabolic activity indicates that bacterial interactions networks

were less when only the vehicle (PEG) was supplied (A) and increased following short-

term supplementation of celecoxib (B) (B). These bipartite networks are based on the

regularised canonical correlations between relative bacterial abundances and relative

concentrations of the main short-chain fatty acids. Interactions have been filtered for an

absolute correlation above 0.5 and are coloured following the key shown. Significant

interactions are shorter lines, and genera with similar abundances within treatment tended

to cluster closely.

Figure 6. Concentration of pro-inflammatory cytokines was decreased in a donor-

dependent manner when celecoxib was supplied. Donors are numbered consecutively as

D1-D8. (A) IL-8 was significantly lower in the simplified inflammation model subjected

to the bacteria-exposed celecoxib from 3 of the donors screened (D1, D2 and D6). Two

additional donors (D3 and D7) showed a trend for decreased IL-8, but it was not

significantly lower than the control treatment. (B) CXCL16 was significantly lower in the

simplified inflammation model subjected to the bacteria-exposed celecoxib from only two

donors (D2 and D6) and tended to be decreased in other two (D3 and D4). (C) The two

volunteers with the most extreme metabolic responses in butyrate production were

selected to be screened in a multicompartment cell model resembling the enterohepatic

system. Donor 6 (D6) had the most significant decrease in butyrate levels, while Donor 8

(D8) did not show any significant difference in butyrate between the control and the

treatment, after 16 h of exposure. IL-8 concentration was significantly lower in the model

exposed to the supernatant of D6 containing bacteria-exposed celecoxib. The opposite

effect was observed in D8, suggesting that functional interindividual differences in the

microbiome play a role in the response to anti-inflammatory drugs.

Table 1. Bacterial functionality metrics following short-term incubation with either

celecoxib or carrier (n = 8). CX, celecoxib; PEG, carrier. SEM, standard error of the

mean.

Variable	Treatment (Mean ± SEM)		*P* value
	CX	PEG	
Acetate (mM)	11.51 ± 1.44	13.31 ± 1.50	0.39
Propionate (mM)	3.29 ± 0.41	3.86 ± 0.42	0.33
Butyrate (mM)	1.57 ± 0.33	3.20 ± 0.33	0.0007
Valerate (mM)	0.61 ± 0.15	0.78 ± 0.15	0.45
Isobutyrate (mM)	0.30 ± 0.07	0.34 ± 0.07	0.70
Isovalerate (mM)	2.70 ± 0.52	3.30 ± 0.54	0.42
Caproate (mM)	0.15 ± 0.04	0.12 ± 0.04	0.61
Isocaproate (mM)	0.001 ± 0.002	0.01 ± 0.002	0.009
Total SCFA (mM)	20.14 ± 2.37	24.91 ± 2.45	0.16
Total Branched SCFA (mM)	2.56 ± 0.49	3.25 ± 0.52	0.34
St:Br ratio	17.76 ± 4.48	12.71 ± 4.31	0.42
% of Acetate	60.87 ± 2.60	53.58 ± 2.77	0.06
% of Propionate	16.17 ± 0.65	15.75 ± 0.70	0.67
% of Butyrate	6.57 ± 0.64	12.99 ± 0.68	<0.0001
Acetate:Propionate:Butyrate ratio	61:16:7	54:16:13	-
pH	5.76 ± 0.04	5.75 ± 0.04	0.86
Total copy number of 16S rRNA gene	3.37E09 ± 3.33E08	4.35E09 ± 3.33E08	0.04

708   Table 2A. Impact of donor and supplementation of celecoxib (CX) or carrier (PEG) on

709   initial bacterial community metrics (time 0 h), n = 8.

Metric	Treatment		SEM	P value		Treatment*Donor interaction
	CX	PEG		Treatment	Donor	
Evenness	0.61	0.60	0.0009	0.83	0.0001	0.23
Shannon (*H'*)	2.71	2.72	0.03	0.81	<0.0001	0.13
Simpson	0.85	0.85	0.007	0.43	<0.0001	0.04
Inverse Simpson (D_2)	8.00	8.48	0.53	0.54	<0.0001	0.76
Fisher's diversity	13.41	13.16	0.25	0.49	<0.0001	0.34
Total Richness (S)	92.50	95.83	3.83	0.55	0.002	0.77
Chao	144.12	128.55	9.54	0.27	0.01	0.59

712   Table 2B. Impact of donor and supplementation on bacterial community metrics,

713   following a short-term incubation (16 h) with either celecoxib (CX) or carrier (PEG), n

714   = 8.

Metric	Treatment		SEM	P value		
	CX	PEG		Treatment	Donor	Treatment*Donor interaction
Evenness	0.55	0.59	0.01	0.06	0.003	0.38
Shannon (*H'*)	2.54	2.70	0.06	0.08	0.001	0.33
Simpson	0.84	0.86	0.009	0.14	0.009	0.10
Inverse Simpson (D_2)	7.59	8.13	0.63	0.55	0.02	0.23
Fisher's diversity	13.28	13.17	0.25	0.49	<0.0001	0.34
Total Richness (S)	92.56	97.12	2.78	0.88	0.005	0.07
Chao	94.01	114.65	5.34	0.01	0.008	0.89

717 Table 3. Epithelial integrity metrics of a multicompartment cell model simulating the

718 enterohepatic metabolism, after 16 h of incubation with CX-exposed bacteria. 0 h =

719 prior to LPS exposure, 16 h = after 16 h of incubation with LPS. Different superscripts

720 indicate significantly different means across treatments, within a particular time point.

721 N = 3.

Treatment	Time point	TEER	SEM	*P* value		
				Treatment	Time	Treatment*Time interaction
CX alone (100 mg/mL)	0 h	385.00				
	16 h	271.67 [a]				
PEG	0 h	330.67				
	16 h	338.67 [a]				
Donor 6 + CX	0 h	470.00				
	16 h	425.33 [b]	34.61	0.007	0.01	0.40
Donor 8 + CX	0 h	398.33				
	16 h	311.33 [a]				
Control medium without LPS	0 h	421.33				
	16 h	337.33 [a]				
Control with LPS	0 h	428.67				
	16 h	433.00 [a]				

(A)

(B)

(A)

Ruminococcaceae UCG-010

Haemophilus

Lactonifactor

Lachnospiraceae UCG-001

Unclassified Family XIII

Lachnoclostridium

Paraprevotella

Gastranaerophilales ge

Ruminococcaceae UCG-009

Lachnospiraceae FCS020 group

Blautia

Phascolarctobacterium

Unclassified Victivallales

Unclassified Veillonellaceae

Parabacteroides

Phascolarctobacterium

Coprococcus 2

Barnesiella

Uncultured Ruminococcaceae

Propionibacterium

Morganella

Prevotellaceae UCG-001

Proteus

(B)

Lachnospiraceae NK4A136 group

Gastranaerophilales ge

Eisenbergiella

Unclassified Christensenellaceae

Family XIII ge

Burkholderia-Paraburkholderia

Erysipelatoclostridium

Uncultured Ruminococcaceae

Ruminiclostridium 9

Unclassified Veillonellaceae

Hungatella

Fusobacterium

Ruminiclostridium 5

Subdoligranulum

Enterococcus

Ruminococcaceae UCG-003

Anaerofilum *Mitsuokella*

Faecalicoccus

Unclassified Family XIII

Defluviitaleaceae UCG-011

Candidatus *Soleaferrea*

Ruminococcaceae UCG-005

SUPPLEMENTARY MATERIAL

Illumina library preparation and purification

The V5-V6 hypervariable region of the 16S rRNA gene from the cDNA was amplified

using primers 341F and 785R. Illumina sequencing adapters and dual-index barcodes

were added to the amplicon, using a limited-cycle PCR that included an initial

denaturation step at 95 ºC for 3 min, 15 cycles of a denaturation step at 95ºC for 30 s, an

annealing step at 55 ºC for 10s, an extension step at 72 ºC for 45 s, and a final extension

at 72 ºC for 5 min. Following, a clean-up step was performed using the AMPure XP beads

(Beckman-Coulter, Krefeld, Germany) to remove free primers and primer-dimer species

from amplicons. A second PCR to attach the specific Illumina multiplexing sequencing

primers and index primers, was performed. Thermal cycling included an initial

denaturation step at 95 ºC for 3 min, 8 cycles of a denaturation step at 95ºC for 30 s, an

annealing step at 55 ºC for 30 s, an extension step at 72 ºC for 30 s, and a final extension

at 72 ºC for 5 min.

PCR products were verified by gel electrophoresis, purified using the Promega Wizard

PCR clean-up kit (Promega, Madison, WI, USA) following the manufacturer's

instructions and quantified with the QuantiFluor dsDNA System kit (Promega, Leiden,

The Netherlands). High-throughput amplicon sequencing was performed with the

Illumina MiSeq platform according to the manufacturer's guidelines at LGC Genomics

GmbH (Berlin, Germany).

Contigs were created by merging paired-end reads based on the Phred quality score (of

both reads) heuristic as described by (17) in Mothur v.1.33.3 (18). Contigs were aligned

to the SILVA database and filtered from those with (i) ambiguous bases, (ii) more than

10 homopolymers, and (iii) those not corresponding to the targeted region. Chimera

removal and operational taxonomic unit (OTU) clustering of the sequencing reads was

performed using UCHIME, with the nearest neighbour clustering algorithm implemented

in mothur, at 0.03 distance (19). Phylotype representatives were then generated by

clustering at 97% similarity (1 mismatch), with a confidence level of at least 80 with

Cyanobacteria, Eukaryota, and Archaea lineages removed. For taxonomic classification,

sequence composition of the dataset was compared using the RDP Classifier tool (20),

and the RDP trainset version 9. Quality of the sequencing and post-processing pipeline

was verified by incorporating mock samples (n = 12 species) in triplicate into the same

sequencing run. After examining read counts, if any OTU was not classified up to genus

level, the consensus sequence was blasted using the SILVA database (21) to obtain an

approximate taxonomic classification. Singletons that remained unclassified were culled.

Supplementary Table 1A. Mean starting relative percentages of the three major SCFA derived from microbial fermentation, following

supplementation with either celecoxib (CX) or the carrier (PEG). N = 6, means between treatments were not significantly different.

SCFA (%)	Ratio found in colon and stool Cummings J.H et al. (1987)	Donor 3		Donor 4		Donor 5		Donor 6		Donor 7		Donor 8	
		CX	PEG	CX	PEG	CX	PEG	CX	PEG	CX	PEG	CX	PEG
Acetate	60	60.02	65.71	35.45	39.30	63.90	65.32	59.25	63.42	40.88	41.84	49.98	53.97
Propionate	20	23.20	21.64	14.84	14.46	12.63	12.48	12.83	11.70	14.20	13.97	10.61	10.81
Butyrate	20	15.49	12.65	10.17	9.48	15.85	15.52	8.20	5.84	6.92	7.20	6.23	6.15

41 Supplementary Table 1B. Relative percentages of the three major SCFA derived from microbial fermentation, following supplementation with

42 either celecoxib (CX) or the carrier (PEG) and incubated for 16 h. N = 8; **, $P < 0.05$. ***, $P < 0.0001$

SCFA (%)	Ratio found in colon and stool Cummings J.H et al. (1987)	Donor 1		Donor 2		Donor 3		Donor 4		Donor 5		Donor 6		Donor 7		Donor 8	
		CX	PEG	CX	PEG	CX	PEG	CX	PEG	CX	PEG	CX	PEG	CX	PEG	CX	PEG
Acetate	60	49.41	47.15	74.64	62.51**	22.46	28.97	33.32	34.76	73.82	57.87***	84.89	63.64***	68.12	67.87	73.99	69.94
Propionate	20	17.30	18.05	22.40	18.27	21.01	18.89	6.90	9.06	14.96	16.13	12.58	18.93**	16.14	13.88	14.18	13.48
Butyrate	20	15.13	17.15	2.27	17.89***	4.83	11.76***	7.51	7.89	6.04	19.48***	2.46	13.01***	8.15	10.29**	7.66	8.07

 Supplementary Table 2. Total metabolically active bacterial population at the beginning (A) and at after 16h (B) of incubation with either

 celecoxib (CX) or carrier (PEG). Means presented were calculated from 3 replicate incubations run in triplicate per donor. N = 8, NS = not

 significantly different.

Donor	0 h				16 h			
	Treatment		SEM	*P* value	Treatment		SEM	*P* value
	CX	PEG			CX	PEG		
D1	-	-	-	-	9.78E+09	2.11E+10		<0.001
D2	-	-	-	-	4.90E+09	4.72E+09		NS
D3	1.54E+08	4.34E+07	4.11E+07	NS	8.59E+08	2.86E+08		NS
D4	1.49E+09	1.20E+09	3.88E+08	NS	1.79E+09	1.71E+09	9.41E+08	NS
D5	3.11E+09	9.08E+09	2.54E+09	NS	3.00E+09	4.73E+09		NS
D6	5.70E+08	3.47E+09	4.25E+08	0.0007	2.49E+08	2.78E+08		NS
D7	2.32E+08	3.43E+08	7.87E+07	NS	1.08E+09	1.64E+09		NS
D8	6.53E+08	1.12E+09	1.36E+08	0.04	5.30E+09	3.21E+08		0.04

59 Supplementary Table 3. Community diversity metrics using the Hill diversity of order

60 2, or inverse Simpson index (D_2) after exposure to celecoxib (CX) or the carrier (PEG),

61 n = 3

Donor	Time	Inverse Simpson		SEM	*P* value		
		CX	PEG		Time	Treatment	Time*Treatment
D1	0h	-	-	4.41	-	0.59	-
	16h	7.20	11.16				
D2	0h	-	-	1.85	-	0.56	-
	16h	6.08	7.89				
D3	0h	3.32	4.33	0.42	0.006	0.01	0.04
	16h	3.90	7.43				
D4	0h	6.93	9.13	0.60	0.04	0.03	0.003
	16h	12.80	7.14				
D5	0h	10.42	9.51	1.16	0.70	0.04	0.73
	16h	6.54	6.48				
D6	0h	5.41	4.16	0.58	0.66	0.01	0.17
	16h	6.81	7.50				
D7	0h	8.86	9.99	0.99	0.23	0.01	0.79
	16h	4.57	6.26				
D8	0h	13.11	13.76	1.79	0.80	0.47	0.55
	16h	12.85	11.20				

65 Supplementary Table 4. Impact of donor, treatment and time on bacterial relative abundances following short-term supplementation of either

66 celecoxib or the carrier. CX, celecoxib. PEG, carrier. Tpt, time point effect; Trt, treatment effect. N = 8.

Taxa	Time point	CX		PEG		*P* value				
		Mean	SEM	Mean	SEM	Donor	Tpt	Trt	Trt*Tpt interaction	Donor*Trt*Tpt interaction
Allisonella	0h	0.0001	0.00003	0.0002	0.0001	0.02	0.07	0.40	0.52	0.52
	16h	0.0002	0.0001	0.0003	0.0001					
Anaerostipes	0h	0.0015	0.0004	0.0015	0.0003	0.002	0.11	0.31	0.40	0.001
	16h	0.0008	0.0002	0.0013	0.0002					
Coprococcus 2	0h	0.0016	0.0005	0.0010	0.0003	<0.0001	0.20	0.63	0.30	0.02
	16h	0.0008	0.0003	0.0010	0.0002					
Erysipelotrichaceae UCG 003	0h	0.0017	0.0003	0.0014	0.0002	0.003	0.16	0.11	0.46	<0.0001
	16h	0.0015	0.0003	0.001	0.0002					
Flavonifractor	0h	0.0003	0.0001	0.0002	0.0001	0.001	0.15	0.43	0.08	0.0004
	16h	0.0001	0.00003	0.0002	0.0001					
Fusobacterium	0h	0.0016	0.0005	0.0012	0.0004	<0.0001	0.85	0.65	0.63	0.66
	16h	0.0015	0.0005	0.0015	0.0004					
Lachnospiraceae FCS020 group	0h	0.0001	0.00004	0.0002	0.0001	0.01	0.16	0.13	0.79	0.70
	16h	0.0001	0.00002	0.0001	0.00003					
Lachnospiraceae UCG 001	0h	0.0012	0.0003	0.0011	0.0002	0.11	0.004	0.99	0.63	0.24
	16h	0.0004	0.0002	0.0005	0.0001					
Marvinbryantia	0h	0.0001	0.00003	0.00004	0.00001	0.17	<0.0001	0.87	0.11	0.43
	16h	0.000004	0.000001	0.00001	0.000002					

Taxa	Time point	CX		PEG		P value				
		Mean	SEM	Mean	SEM	Donor	Tpt	Trt	Trt*Tpt interaction	Donor*Trt*Tpt interaction
Parabacteroides	0h	0.0095	0.0017	0.0075	0.0014	<0.0001	0.54	0.11	0.82	<0.0001
	16h	0.0109	0.0014	0.0080	0.0009					
Paraprevotella	0h	0.0005	0.0002	0.0005	0.0002	0.0001	0.13	0.65	0.77	0.02
	16h	0.0003	0.0001	0.0004	0.0001					
Parasutterella	0h	0.0013	0.0003	0.0013	0.0003	<0.0001	0.33	0.46	0.36	0.02
	16h	0.0020	0.0004	0.0013	0.0003					
Peptococcus	0h	0.0003	0.0001	0.0001	0.00004	0.01	0.06	0.25	0.41	0.68
	16h	0.0001	0.00004	0.0001	0.00003					
Phascolarctobacterium	0h	0.0035	0.0008	0.0028	0.0006	<0.0001	0.56	0.77	0.53	<0.0001
	16h	0.0027	0.0006	0.0029	0.0005					
Ruminiclostridium 5	0h	0.0006	0.0001	0.0005	0.0001	0.09	0.02	0.49	0.83	0.03
	16h	0.0003	0.0001	0.0003	0.0001					
Ruminiclostridium 6	0h	0.0013	0.0003	0.0012	0.0003	<0.0001	0.09	0.80	0.54	0.02
	16h	0.0007	0.0002	0.0009	0.0002					
Ruminococcaceae NK4A214 group	0h	0.0004	0.0001	0.0003	0.0001	<0.0001	0.10	0.57	0.49	0.28
	16h	0.0002	0.0001	0.0002	0.0001					
Ruminococcaceae UCG 004	0h	0.0002	0.0001	0.0002	0.0001	0.001	0.06	0.73	0.60	0.75
	16h	0.0001	0.00003	0.0001	0.00003					
Ruminococcaceae UCG 010	0h	0.0004	0.0001	0.0003	0.0001	0.0004	0.07	0.95	0.46	0.03
	16h	0.0002	0.0001	0.0002	0.0001					
Ruminococcus 2	0h	0.0017	0.0005	0.0015	0.0003	<0.0001	0.33	0.49	0.83	0.01
	16h	0.0014	0.0004	0.0011	0.0002					

Taxa	Time point	CX		PEG		_P_ value				
		Mean	SEM	Mean	SEM	Donor	Tpt	Trt	Trt*Tpt interaction	Donor*Trt*Tpt interaction
Streptococcus	0h	0.00002	0.00001	0.00002	0.00001	0.01	0.06	0.79	0.23	0.17
	16h	0.00005	0.00002	0.00003	0.00001					
Unclassified Christensenellaceae	0h	0.0001	0.00002	0.00003	0.00001	0.002	0.79	0.32	0.21	0.79
	16h	0.00004	0.00001	0.00004	0.00001					
Veillonella	0h	0.0009	0.0003	0.0007	0.0002	0.03	0.42	0.34	0.80	0.01
	16h	0.0013	0.0003	0.0009	0.0003					
Victivallis	0h	0.0008	0.0002	0.0004	0.0001	<0.0001	0.46	0.09	0.30	0.01
	16h	0.0005	0.0001	0.0004	0.0001					

68