Stress-responsive *Entamoeba* topoisomerase II: a potential anti-amoebic target

Sneha Susan Varghese, Sudip K. Ghosh[*]

Department of Biotechnology, Indian Institute of Technology Kharagpur, Kharagpur, West Bengal, India 721 302.

[*] Corresponding author

Email: sudip@bt.iitkgp.ac.in (SKG)

Short Title: *Entamoeba* topoisomerase II: a potential anti-amoebic target

Author contributions

SSV and SKG: designed research, SSV: performed experiments, SSV and SKG: wrote the paper.

Abstract

Topoisomerases are a group of ubiquitous enzymes, involved in all key DNA processes and are often stress responsive across the biological world. As a result, these enzymes are targets for various anticancer and antibacterial agents. The causative organism of amoebiasis, *E. histolytica* (*Eh*), has seven unexplored genes annotated as putative topoisomerases. One of the seven topoisomerases in this parasite was found to be highly up-regulated during heat shock and oxidative stress. The bioinformatic analysis shows that it is a eukaryotic type IIA topoisomerase. Its ortholog in *E. invaden*s (*Ei*), the encystation model of *E. histolytica* was also highly up-regulated during the late hours of encystation. Immunoprecipitated endogenous EhTopo II showed topoisomerase II activity *in vitro*. Immunolocalization studies show that this enzyme colocalized with newly forming nuclei during encystation, which is one of the significant events of mature cysts. Double-stranded RNA mediated downregulation of the Topo II both in *Eh* and *Ei* reduced the viability of actively growing trophozoites and also reduced the encystation efficiency in *Ei*. Drugs, targeting eukaryotic topoisomerase II e.g., etoposide, ICRF193, and amsacrine, show 3-5 times higher EC_{50} in *Eh* than that of mammalian cells, although EhTopo II is eukaryotic. Sequence comparison with human TopoIIα showed that key amino acid residues involved in the interactions with ICRF193 and etoposide are naturally substituted in EhTopo II. Interestingly, ciprofloxacin an inhibitor of prokaryotic DNA gyrase showed about six times less EC_{50} value in case of *Eh* in comparison to human cells. The parasite's notable susceptibility to prokaryotic topoisomerase drugs in comparison to human cells opens up future scope to study this invaluable enzyme in the light of an antiamoebic target.

Author Summary

Amoebiasis, a globally prevalent parasitic infection, claims about 100,000 lives annually. The disease is caused by the protozoan parasite *E. histolytica,* which can alternate between a pathogenic trophozoite and a dormant cyst stage, within and outside the human host respectively. Stage conversion to the resistant cyst i.e., encystation is the primary reason for the disease transmission. There are no vaccines available to date, and broad spectrum nitroimidazole derivatives are still the primary mode of treatment for this epidemic. However, the toxicity and side effects of these drugs as well as increasing drug resistance in the parasites emphasize the urgency for better treatment methods. Topoisomerases are vital enzymes required for all DNA process and are extensively studied as targets for antibacterial and anticancer drugs. *E. histolytica* has seven putative topoisomerases. Here, we identified that topoisomerase II of *Entamoeba* is crucial not only for the growth of the trophozoite form but also for the mature cyst formation. Additionally, we found that this enzyme is highly upregulated during other stress like oxidative stress and heat shock. In comparison to mammalian cells, the EC_{50} of eukaryotic topoisomerase II drugs like etoposide, ICRF 193 and amsacrine was about five times higher. Contrarily, EC_{50} of gyrase inhibitors like fluoroquinolones (e.g., ciprofloxacin) was approximately six times lower in *E. histolytica* than human cells. In short, our study provides insights into using *Entamoeba* topoisomerase II as an anti-amoebic target, as it is vital for growth and stage conversion. Further, we address the scope of designing fluoroquinolone derivatives that can specifically target the parasite with low toxicity to the human cells.

Introduction

Amoebiasis is a leading cause of death due to a parasitic protozoan infection worldwide, highly prevalent in regions with poor sanitation and hygiene. The causative organism, *Entamoeba histolytica*, alternates between two life stages- pathogenic trophozoite and resistant cyst stage [1]. Stage conversion to the dormant form is called encystation and is the critical process for host-to-host transmission of the disease. Much of the information about encystation comes from studying the closely related reptilian parasite *E. invadens*, as it can encyst *in vitro* under reduced osmotic pressure and nutrient depletion [2, 3]. Hence, stress is a precursor for encystation, and this process witnesses the activation of several specific stress-responsive proteins and signaling pathways [4, 5]. One of the key features during this stage conversion is the appearance of four distinct nuclei during the later hours of encystation [6, 7]. Association of the sexual process, like meiosis, with encystation in *Entamoeba,* has been a point of debate for long. Low levels of allelic heterozygosities [8], evidence of homologous recombination [9], the presence of meiotic genes and their upregulation during encystation [10], and the formation and aggregation of haploid nuclei in multinucleated giant cells during encystation [11] indicate the occurrence of meiosis-like processes and homologous recombination during glucose deprived encystation.

Across the biological world, all key DNA processes are associated with a group of ubiquitous enzymes called topoisomerases. These enzymes maintain DNA topology by creating breaks in the DNA and allowing strand passage. Thus, these are involved in removing DNA supercoils, chromosomal condensation, strand-breakage during recombination, and disentangling intertwined DNA, in naming a few [12, 13]. Based on their mode of action, there are two broad types of topoisomerase, viz Type I and Type II. The former function as monomers, creating single strand breaks while the latter forms multimers and introduces double-strand breaks in the

DNA. Owing to the vital functions of these enzymes, they have been extensively evaluated as

targets for many antitumors, antibacterial, and antifungal drugs [14-17]. Also, topoisomerases in

several pathogenic protozoans like *Plasmodium,* and *Leishmania* are being evaluated as targets

for antiparasitic agents [18-21].

Treatment for amoebiasis relies primarily on nitroimidazole based drugs, especially

metronidazole [22]. Although effective both in bowel lumen and tissue, this broad spectrum drug

is reported to eradicate only 50% of luminal infection [23]. Successful induction of

metronidazole-resistant parasitic strain at the laboratory level points to the probability of drug

resistance in the near future [24, 25]. Moreover, the side effects of metronidazole include nausea,

abdominal pain, diarrhea, and in severe cases, results in neurotoxicity, optic neuropathy,

peripheral neuropathy, and encephalopathy [26, 27]. Hence, there is a need for newer and

effective alternative solutions.

In this study, stress-responsive, eukaryotic Type IIA topoisomerase was identified in *Eh* and *Ei*,

upregulated both at RNA as well as protein level during later stages of encystation, and various

other stresses, like heat shock and oxidative stress. RNAi mediated silencing of that gene has

shown a significant impact on the overall viability and encystation efficiency of *Entamoeba*.

Further, a higher potency of prokaryotic topoisomerase II drugs on the parasite when compared

to human cells makes it a significant drug target worth further exploration.

2. Results

2.1. Phylogenetic classification of *Entamoeba* topoisomerase genes

Topoisomerases are broadly categorized as type I and II, based on their structure and function. Type I functions as a monomer, creating single strand breaks and based on differences in the mechanism of action, they are subcategorized as IA and IB. Prokaryotic topo I, all topo III and reverse gyrase belong to the former while latter comprises of eukaryotic topo I and archeal topoV. Although all type II topoisomerases are multimers and create double strand breaks, based on their structural differences, these are sub-classified as type IIA (eukaryotic topo II, topo IV and DNA gyrase) and type IIB (topo VI).

Phylogenetic analysis shows that all genes annotated as topoisomerases in *Entamoeba* are eukaryotic (Fig 1). Interestingly, orthologs of *Eh* and *Ei* were very closely related, and hence observations for one may be extrapolated to its ortholog. EHI_073170/EIN_344850 and EHI_087330/EIN_173080 showed a very distant relation with Topo VI, which is reported only in plants and archaeabacteria. From the remaining pool of five genes, EHI_038920/EIN_052260 and EHI_042880/EIN_174490 shared maximum similarity with topoisomerase IIIα and IIIβ (Type IA), respectively. Further, EHI_125320/EIN_371990 and EHI_194510/EIN_229710 aligned closely with SPO11 family, which function similar to Topoisomerase II and creates double-strand breaks during meiosis. EHI_120640/EIN_145900 resembled eukaryotic Topoisomerase IIα (Type IIA). Multiple sequence alignment and NCBI-CDD shows that the putative topoisomerase II has three core domains: N-terminal ATPase domain, a central domain carrying the active site and a C-terminal variable domain carrying the NLS sequence (Fig 2 and S7 Fig). Homology modeling predicts that the protein is dimeric (S5 Fig). Interestingly, neither of the *Entamoeba* species had a recognizable eukaryotic topo I (Type IB).

Fig 1. The phylogenetic classification of *Entamoeba* topoisomerase genes. Phylogenetic tree generated based on maximum likelihood method using MEGA 7.0 software. Different topoisomerases from various prokaryotic and eukaryotic organisms were used to construct the tree. The seven putative topoisomerases of *E. histolytica* and *E. invadens* is highlighted with **blue** and **red** arrows, respectively. **At:** *Arabidopsis thaliana*; **Bs**: *Bacillus subtilis*; **Ce**: *Caenorhabditis elegans*; **Cr**: *Chlamydomonas reinhardtii*; **Ds**: *Dictyostelium discoideum*; **Dm**: *Drosophila melanogaster*; **Ec**: *Escherichia coli*; **Hs**: *Homo sapien* **Mm**: *Mus musculus*; **Os**: *Oryza sativa*; **Sc**: *Saccharomyces cerevisiae*.

Fig 2. Domain organization on putative Topo II of *Entamoeba*. Comparison of predicted domains on putative topoisomerase II of *Eh* and *Ei* with the domains of human topoisomerase IIα. The putative TopoII of *Entamoeba* has all the three crucial domains: N-terminal domain with ATPase activity, Central catalytic domain housing the active Tyr residue, and highly variable C-terminal domain that nestles the Nuclear Localization Sequence.

2.2. Expression profile of putative *Entamoeba* Topoisomerase genes during different stresses and encystation

Real-time RT-PCR analysis of these five genes shows that only EHI_120640 and its ortholog EIN_145900 were highly upregulated during heat shock and oxidative stress, although the gene expression did not significantly alter during 16h of glucose starvation. As a similar upregulation of EIN_145900 gene expression was also observed during later periods of encystation, this could be important for stress response in *Entamoeba* (Fig 3). Much like during 16h of glucose starvation, early periods of encystation did not show upregulation of EIN_145900. Similar observations for all these genes during encystation were noted from the microarray data [28] (S1 Fig). As a result, further study focused on this stress responsive putative topoisomerase II. EIN_229710, predicted as SPO11 has already been reported to be a meiosis-specific gene and upregulated during encystation of *E. invadens* [10].

Fig 3. **Expression profile of putative *Entamoeba* Topoisomerase genes during different stresses and encystation**. sq-RT PCR and corresponding Real-time PCR analysis showing relative fold change in transcription of different putative topoisomerases under different stress conditions of **a**) *E. histolytica* and **b**) *E. invadens* and, **c**) *E. invadens* during different hours on encystation. EHI_120640 and its ortholog, EIN_145900 are markedly upregulated during various stress and later hours of encystation. **C-**Control; **GS-**Glucose starvation; **OS-** Oxidative stress; **HS-** Heat shock

2.3. Cloning, expression, and purification of recombinant putative EhTopo II fragment

Although different combinations of bacterial expression vector and strains were tried, full-length expression of putative EhTopo II (4050bp) was not successful. Hence a 1127bp fragment (from 760bp-1887bp) was successfully cloned into pET21a expression vector, and recombinant EhTopo II fragment was successfully expressed with a 6x Histidine tag at the C-terminal in *E. coli* Bl21(DE3) strain using 1mM IPTG. The insoluble, recombinant protein so obtained was solubilized using 0.25% S-lauryl sarcosine, purified by Ni-NTA affinity chromatography and resolved by SDS-PAGE (S2 Fig). Purified recombinant EhTopo II fragment was used to raise antibody in rabbits. Anti-EhTopoII antibody was purified from crude sera and confirmed by Western blot analysis. As the antigenic fragment of EhTopo II shared 70% sequence similarity with its ortholog in *Ei,* the anti-EhTopo II antibody could successfully bind to EiTopo II as well (S3 Fig).

2.4. Topoisomerase II protein expression is upregulated during different stresses and encystation in *Entamoeba*

Topoisomerase II protein was expressed under all the studied stress conditions in both *E. invadens* and *E. histolytica* with a marked upregulation during heat shock and oxidative stress (Fig 4a and 4b). The temporal expression of the protein during encystation followed a pattern similar to the transcript profile wherein the protein expression gradually declined in the early hours and showed a drastic upregulation during the later stages, which is also the period of tetranuclei formation (Fig 4c). Hence this confirms *Entamoeba* Topo II is upregulated at both transcription and protein synthesis.

Fig 4. Upregulation of Topoisomerase II protein expression during different stress and encystation in *Entamoeba.* Western blot analysis for the expression profile of Topo II protein during different stresses in **a**) *E. histolytica,* **b**) *E. invadens* and **c**) during different hours of encystation in *E. invadens,* respectively. Similar findings as that of Real-time RT-PCR were observed. ARF was used internal control for normalisation. **C-** Untreated control; **GS-** Glucose starvation; **OS-**Oxidative stress; **HS-** Heat shock.

2.5. *Entamoeba* trophozoites express functional topoisomerase II

The function of topoisomerases is to relieve the topological strain that occurs in the DNA during

various processes like replication, transcription, chromosomal segregation, etc. This is achieved

by relaxing positive or negative supercoils in the DNA and depending on the type of enzyme,

various cofactors are required for this process. We have predicted the presence of eukaryotic

Topo III and Topo II in *Entamoeba*. TopoIII requires only $MgCl_2$ while Topo II requires both

ATP and $MgCl_2$ as cofactors for its enzymatic activity. To determine whether *Entamoeba*

trophozoites express functional topoisomerase II, we studied its ability to relax negative

supercoils. Varying concentration of nuclear extracts, prepared from actively proliferating

cultures, could successfully relax negatively supercoiled plasmids in the presence of co-factors

ATP and $MgCl_2$. At 40μg of crude nuclear extract, *Eh* was able to relax 100% of the supercoiled

plasmid in comparison to 70-80% relaxation by *Ei* at the same concentration (Fig 5a and 5b).

This study shows the presence of active Topo II in *Entamoeba* trophozoites

Endogenous EhTopo II was immunoprecipitated from the crude nuclear extract using anti-

EhTopo II antibody. The native EhTopo II showed relaxation activity only in the presence of

both ATP and $MgCl_2$. As no relaxation was observed in the absence of either ATP or $MgCl_2$ or

both, it can be concluded that the pulled down Topo II was not contaminated with any type I

topoisomerases (Fig 5c).

195
196
197

Fig 5. *Entamoeba* trophozoites express functional topoisomerase II. Agarose gel electrophoresis (1%) depicting the relaxation of negatively supercoiled plasmids by varying amounts of crude nuclear (Lane 1-4 contains 0, 20µg, 40µg and 60µg of crude nuclear extract from the respective source) of **a)** *Eh* and **b)** *Ei*. **c)** endogenous EhTopoII (immunoprecipitated from nuclear extracts of *Eh,* using anti-TopoII antibody). Relaxation activity occurs only in the presence of both ATP and $MgCl_2$. **R** and **S** represent the negatively supercoiled plasmid and the relaxed form, respectively.

2.6. Localization of Topoisomerase II on newly forming tetranuclei during encystation in

Entamoeba

Confocal micrograph analysis showed that Topo II localizes in the nucleus of *Eh* trophozoites

during normal growth as well as during oxidative stress and heat shock. Contrarily, during

glucose starvation, the protein was observed to move out into the cytoplasm (Fig 6a), and this

observation during glucose starvation was consistent in *Ei* as well (Fig 6b). During the early

hours of encystation (12 hours) EiTopo II expression reduced in comparison to trophozoite. This

observation is in consensus with real-time RT-PCR and western blot analysis. As topoisomerase

II is a key player in cell growth and proliferation, degradation of the enzyme during glucose

starvation and early encystation could be a part of the initial survival response of the organism to

energy deficiency.   However, with the progression of encystation into later hours,

Topoisomerase II co-localized with the newly forming tetranuclei suggesting that this enzyme

may be responsible for relieving the topological strains that occur in the DNA during this nuclear

event (Fig 6c).

Fig 6. Topo II localization during stress and encystation in *Entamoeba*. The localization pattern of Topo II during **a)** different stress conditions in *Eh* and **b)** *Ei*. The enzyme primarily localized in the nucleus, except during glucose starvation**. c)** Localization of Topo II during different hours of encystation in *E. invadens*. The early hours of encystation show the decline of topoisomerase II from the cell. However, during the later stages of encystation, the protein reappears and localizes on the newly forming nuclei. *Entamoeba* TopoII is stained red with TRITC, and the nucleus is stained blue with DAPI.

2.5. TopoII silencing in *Entamoeba* significantly affects the viability of trophozoite and cyst

To understand the *in vivo* role of *Entamoeba* TopoII during normal growth as well as encystation, we employed the dsRNA mediated silencing strategy reported by Samanta and Ghosh, 2012 [29]. Gene-specific dsRNA was cloned into the pL4440 vector which is flanked by T7 promoter on either side, expressed and purified from RNase III-deficient *E. coli* HT115 cells. Approximately, 60-70% reduction in the transcription of Entamoeba *Topo II* was observed in samples soaked with 200µg/mL gene-specific dsRNA in comparison to control (untreated) (S4 Fig).

Topoisomerase is considered a proliferation marker and is an absolute requirement for rapidly proliferating eukaryotic cells. Interestingly, silencing of topoisomerase II in actively growing, log-phase trophozoites significantly (P<0.001) reduces the viability by about 25% and 30% in *Eh* and *Ei*, respectively (Fig 7a). A similar reduction in viability as well as encystation efficiency was observed upon silencing the gene in samples under encystation condition. The encystation efficiency and viability of *EiTopoII* silenced cells reduced by approximately 42% and 28% respectively in comparison to the two control conditions (Fig 7b and 7c).

Fig 7. TopoII silencing in *Entamoeba* significantly affects the viability of trophozoite and cyst. a) The % viability of *Eh* and *Ei* trophozoites, on silencing the respective Topoisomerase II for 24h. A significant reduction of 30% and 25% (p<0.001) respectively, in the viability of actively growing trophozoites, were observed. **b)** Cell viability (%) and **c)** encystation efficiency of *Ei* after 48h of encystation during dsRNA mediated Topo II downregulation (200µg/ml). A significant reduction (p<0.0001) in the viability and encystation efficiency of about 30% and 42% respectively observed upon silencing *TopoII*

2.6. Less toxicity of eukaryotic topoisomerase II inhibitors in *E. histolytica*

As topoisomerase II is an extensively studied class of enzyme and is essential for cell growth and proliferation, many drugs designed against the eukaryotic topoisomerase II are available as anticancer and antifungal agents. We tested the potency of three different kinds of eukaryotic Topoisomerase II inhibitors viz. etoposide, amsacrine, and ICRF-193 on *E. histolytica*. The former two stabilize the cleavable complex, prevent the religation of double-strand breaks and hence stimulate enzyme-mediated DNA breakage. ICRF 193, a bizdioxopiperazine, locks the enzyme in a closed-clamp and prevents ATP hydrolysis necessary to regenerate the active form of the enzyme. Etoposide, ICRF-193, and amsacrine were lethal against *E. histolytica* with EC_{50} of 200µM, 15µM and, 80µM, respectively (Fig 8). However, the susceptibility of *Eh* to eukaryotic drugs is approximately 3-5 times lower than that of the human hosts, especially towards etoposide (Table 1).

Two crucial regions in hTopo II viz, PLRGKXLNVR (motif I) and Q/MXLMM (motif II) are reported to interact with etoposide [30]. Topo II aminoacid sequence alignment of different *Entamoeba* species with that of hTopo IIα shows that the motif I is mostly conserved. The Q/MXLMM motif on hTopoIIα interacts with etoposide primarily via the M_{762} and M_{766} residues. Mutation of these residues has shown to interfere with the etoposide binding property and increase resistance to this drug [31, 32]. The motif II is mostly conserved among different species of *Entamoeba;* however, it is different from the motif II in hTopoIIα. The key drug interacting methionine residues are absent in all *Entamoeba* species and are naturally substituted with V_{723} and A_{727} in *E. histolytica* (Fig 9)*.

The drug binding pocket for bisdioxopiperazines like ICRF 193 comprises of 14 residues (7 from each monomer) in eukaryotic Topo II [33, 34]. The histidine residue in this pocket is replaced

263   with glutamine not only in the human parasite but in all other species of *Entamoeba* as well (Fig

264   10). Alterations in these residues could be the reason for low toxicity of eukaryotic Topo II drugs

265   on *E. histolytica.*

267   **Fig 8. Less toxicity of eukaryotic topoisomerase II inhibitors in *E. histolytica*.** Dose-dependent effect of eukaryotic Topoisomerase II drugs, namely **a)** etoposide **b)** ICRF 193 and **c)** amsacrine (AMSA) on *E. histolytica* viability following 24h treatment.

Einvadens	DSAKTLAVSGLSVVGRDYYGVF**PLRGKPLNAR**EISSAKVKNNQEFENIAKIMGLRYGKKY	490
Emoshkovskii	DSAKTLAVSGLAVVGRDYYGVF**PLRGKPLNAR**ELATTKVKENQEFENIAKIMGLKYGKVY	497
Edispar	DSAKTLAVSGLSVVGRDYYGVF**PLRGKPLNAR**EIAPSKVKENQEFENIAKIMGLRYGKVY	414
Ehistolytica	DSAKTLAVSGLSVVGRDYYGVF**PLRGKPLNAR**EIAPSKVKENHEFENIAKIMGLRYGKVY	494
Enuttalli	DSAKTLAVSGLSVVGRDYYGVF**PLRGKPLNAR**EIAPSKVKENHEFENIAKIMGLRYGKVY	494
Scerevisiae	DSALSLAVAGLAVVGRDYYGCY**PLRGKMLNVR**EASADQILKNAEIQAIKKIMGLQHRKKY	510
Hsapiens	DSAKTLAVSGLGVVGRDKYGVF**PLRGKILNVR**EASHKQIMENAEINNIIKIVGLQYKKNY	522

```
              *** :***:**.***** ** :***** **.** :    :: :* *:: * **:**:: * *
```

Einvadens	SYHHGE**VSLQS**TIVNMAQNFCGANNINLLLPSGQFGSRLQGGKDQAAARYIYTRLSTITR	772
Emoshkovskii	SYHHGE**QSLQA**TIVNMAQNFCGSNNINWLLPSGQFGSRLGGGKDQAAARYIFTRLSSISR	779
Edispar	SYHHGE**VSLQA**TIVNMAQNFCGSNNVNWLLPSGQFGSRLGGGKDQAAARYIYTRLSSISR	696
Ehistolytica	SYHHGE**VSLQA**TIVNMAQNFCGSNNVNWLLPSGQFGSRLGGGKDQAAARYIYTRLSSISR	776
Enuttalli	SYHHGE**VSLQA**TIVNMAQNFCGSNNVNWLLPSGQFGSRLGGGKDQAAARYIYTRLSSISR	776
Scerevisiae	AYHHGE**QSLAQ**TIIGLAQNFVGSNNIYLLLPNGAFGTRATGGKDAAAARYIYTELNKLTR	792
Hsapiens	SYHHGE**MSLMM**TIINLAQNFVGSNNLNLLQPIGQFGTRLHGGKDSASPRYIFTMLSSLAR	815

```
              :***** ** **:..**** *:**:   * * * **:*   **** *: ***:* *.:.:*
```

269   **Fig 9. Prediction of etoposide binding motif on *Entamoeba* Topo II.** Multiple Sequence Alignment of

270   hTopoIIα and ScTopoII with the ortholog in different *Entamoeba* species showing the PLRGKXLNVR (Motif I) and Q/MXLMM (Motif II) motifs involved in etoposide interaction (highlighted yellow). Most of the interacting residues were conserved between the host and parasite proteins. However, in the motif II,

271   Met$_{762}$ and Met$_{766}$ of hTopoIIα were substituted with Val$_{723}$ and A$_{727}$ in EhTopo II, and these aminoacid replacements were present in all other *Entamoeba* species as well (marked with red arrow). [30]

Fig 10. Prediction of ICRF-193 binding motif on *Entamoeba* Topo II. Multiple Sequence Alignment of hTopoIIα and ScTopoII with the ortholog in different *Entamoeba* species showing the residues involved in the interaction with bizdioxopiperazine derivatives like ICRF-193, in eukaryotic Topo II (highlighted blue). The His_{41} involved in the drug binding pocket of hTopoIIα is substituted with Gln_{20} in the EhTopo II. Substitution of His with Gln in the drug binding pocket was conserved in different species of *Entamoeba* (marked with red arrow) [33,34]

2.7. Susceptibility of *E. histolytica* to fluoroquinolones targeting prokaryotic Topo IIA

Drugs that target prokaryotic topoisomerase II like DNA gyrases and Topoisomerase IV are primarily fluoroquinolone derivatives. Ciprofloxacin and clinafloxacin are new generation fluoroquinolones with a broad spectrum of antibacterial activity. However, these drugs have a much lower potency against eukaryotic topoisomerase II. Interestingly, both of these drugs were toxic to the survival of *Eh* with an effect comparable to that of etoposide (Fig 11). Further, ciprofloxacin has a profound, direct inhibitory effect on the enzymatic activity of native EhTopo II in comparison to etoposide, affecting the enzyme at a much lower concentration than that of the latter (Fig 12). It is also interesting to note that ciprofloxacin has much lower EC_{50} against *Eh* in comparison to that reported in mammalian cell line [35, 36]. Like in case of *Eh*, ciprofloxacin

289 can reduce the viability of *Ei* and also decrease the encystation efficiency (Fig 13). This opens up

290 scope to further explore newer and better fluoroquinolones in the light of an anti-amoebic drug

291 against this parasite (Table 1).

Fig 11. Susceptibility of *E. histolytica* to gyrase inhibitors. Effect of prokaryotic TopoII inhibitors (fluoroquinolones) **a)** ciprofloxacin and **b)** clinafloxacin on the viability of *Eh* post 24h drug treatment.

Fig 12. Ciprofloxacin inhibits EhTopo II enzymatic activity. Agarose gel (1%) depicting the inhibitory activity of etoposide and ciprofloxacin on EhTopoII ability to relax negative supercoils. (Lane 1-7: Etoposide treatments; Lane 8-12: different controls; Lane 13-17: Ciprofloxacin treatments.) Ciprofloxacin inhibits the enzymatic reaction at much lower concentrations (160 µM) than the eukaryotic inhibitor, etoposide.

Fig 13. Effect of etoposide and ciprofloxacin on encystation. a) The % cell viability and **b)** encystation efficiency of *Ei* after 48h of drug treatment (160µM etoposide and 200µM ciprofloxacin) during encystation. Both treatments reduced the viability as well as the efficiency of encystation.

Table 1. Comparison of EC_{50} of various TopoII inhibitors against *Eh* to that of mammalian cell lines. (N/D: Not determined)

Inhibitor	EC_{50} *E. histolytica*	EC_{50} Mammalian cell line
Etoposide	200µM	40-80µM[37,38]
ICRF 193	15µM	2-5 µM[39]
Amsacrine	80µM	20-30µM[40]
Ciprofloxacin	250µM	1740µM[35,36]
Clinafloxacin	270µM	N/D

3. Discussion

Topoisomerases, being a crucial player in all DNA processes, is often involved in the stress response and survival of all organisms across the biological world [41, 42]. Like all other organisms, *E. histolytica* also has a pool of genes coding for various types of topoisomerases. The bioinformatic analysis shows that the putative topoisomerases of *E. histolytica* resembled Topo IIIα, Topo IIIβ, Topo II, and SPO11 and are very closely related to their orthologs in *E. invadens*. Although all topoisomerases genes were eukaryotic, (Phylogenetic analysis) recognizable eukaryotic Topo I could not be identified in *Entamoeba*. Among these genes, we observe that topoisomerase II is crucial not only for normal growth and proliferation of the trophozoites but appears to be the most important for the parasites' response to various stress conditions like heat shock as well as oxidative stress. Similar levels of upregulation of topoisomerase II under these stress conditions are a reported phenomenon in human cancer cells and mediate the excision of chromosomal DNA loops into High Molecular weight (HMW) fragments as a stress response signal under oxidative stress [43, 44].

Stress is one of the most important environmental signals for encystation in *Entamoeba*, leading to the formation of environmentally resistant cysts. As a response to energy deficient environment during glucose starvation and early encystation, trophozoites switch from an actively proliferating stage to a dormant form during which many genes involved in the metabolism and proliferation are downregulated. Likewise, we show that Topoisomerase II, a proliferation marker, also faces a steady decline in the early periods of encystation in *Entamoeba*. Similar stage-specific expression of topoisomerase II has been observed in other parasites as well. For example, in *P. falciparum* the ring stage has a relatively lower level of the protein expression in comparison to trophozoite and schizont stage [18] while in *Leishmania*

infantum expression levels varied considerably between infective intracellular amastigotes,

proliferative and non-proliferative stages of promastigotes [45]. However, we report that, with

the progression of encystation, transcription and translation of topoisomerase II is upregulated in

Entamoeba around the period of tetranuclear formation, which is one of the key phenomena in a

maturing cyst. Further, during this stage of encystation, the enzyme colocalizes on the newly

forming nuclei in *E. invadens* suggesting that it may be necessary to remove the topological

strains that occur in the DNA during this nuclear event. Also, the significant drop in viability and

encystation efficiency upon its downregulation suggests that *Entamoeba* topoisomerase II is

indeed important for the proper stage conversion of this parasite.

Over the years, topoisomerase II has emerged as a good drug target and consequently led to the

development of an extensive array of medicines as anti-cancer and anti-bacterial agents. As

already mentioned, it is extensively explored as a potential drug target in many other parasites as

well. We showed that *E. histolytica* has a lower susceptibility to eukaryotic topoisomerase

targeting drugs, especially etoposide while fluoroquinolones that target prokaryotic gyrases and

topoisomerase IV were significantly toxic to the enzyme activity as well as parasite viability.

Sequence analysis of EhTopo II with human TopoIIα show that specific key residues involved in

the drug interaction with etoposide and ICRF 193 are naturally substituted in *Eh* and this may be

a contributing factor to the poor performance of these drugs towards the parasite in comparison

to mammalian cells. On the other hand, gyrase inhibitors like fluoroquinolones have reported

lower potency against higher eukaryotes, including humans [46]. Chemical modifications to the

side chains of fluoroquinolones have shown to enhance or decrease its potency towards different

organisms [47, 48]. Similar susceptibility to fluoroquinolones, in comparison to mammalian

cells, is also reported in case of other parasites like *P. falciparum* and *L. donovanii* and is being

explored as a drug target as well [18-21]. Another bacterial gyrase inhibitor GSK299423 (a

piperidinylalkylquinoline) has been reported to be extremely effective against *P. falciparum* with

100 times higher potency than against mammalian cell lines [49]. In addition, computational

design of newer fluoroquinolones with enhanced potency against parasitic topoisomerase is

already gaining traction and thus opens up possibilities for better anti-parasitic drugs [50, 51].

Hence, high potency of fluoroquinolones to *E. histolytica*, low toxicity towards human cells and

ease of chemical modification in its side chains are advantages for designing these drugs to

specifically target EhTopo II.

Limited treatment methods and potential drug resistance have accelerated the need for newer

therapies to tackle amoebiasis. The topoisomerase II of *Eh* is crucial for the stress response and

formation of mature cysts and shares only 45% identity with the human counterpart. As we

already show that ciprofloxacin is toxic not only to the proliferating trophozoite stage but also

severely reduces encystation, designing newer variations of fluoroquinolones that can

specifically target *E. histolytica* with minimum side effects to the host could be the answer to not

only treat the disease but also prevent the host-to-host transmission of amoebiasis.

4. Materials and Methods:

4.1. In silico analysis:

4.1.1. Multiple sequence analysis:

Genes annotated as putative topoisomerases from *Eh* and *Ei* were identified from AmoebaDB. Protein sequences of topoisomerases from different organisms were retrieved from the UniProt database (http://www.uniprot.org/) and were aligned with putative topoisomerases of *Entamoeba* using ClustalOmega (http://www.ebi.ac.uk/).

4.1.2. Phylogenetic tree construction:

Phylogenetic analysis was performed to understand the categories of topoisomerases the seven in *Entamoeba* fall into. Protein sequence from 50 different topoisomerases covering all known categories, belonging to a range of prokaryotic and eukaryotic organisms like bacteria, fungi, protozoa, plant, and animals were retrieved using UniProt database and aligned with the putative topoisomerases of *Eh* and *Ei* using Clustal W. Using this alignment, phylogenetic tree was constructed with MEGA 7 [52] software by the Maximum Likelihood method and evolutionary distance between sequence pairs was analyzed by the WAG model. Reliability of the model was assessed by bootstrapping with 1000 iterations.

4.1.3. Conserved domain identification and analysis:

Conserved motifs on putative *Entamoeba* topoisomerases were identified using MEME software and further analyzed using NCBI-CDS (http://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi) [53], and nuclear localization sequences (NLS) was predicted using cNLS mapper (http://nls-mapper.iab.keio.ac.jp/cgi-bin/NLS_Mapper_form.cgi) [54].

4.2. Growth, stress induction, and encystation of *Entamoeba:*

Trophozoites of *E. histolytica* strain HM-1:1MSS and *E. invadens* IP-1 were grown at 37 °C and 25 °C respectively in Trypticase-Yeast Extract-Iron-Serum (TYI-S-33) medium containing 10% heat-inactivated adult bovine serum, 125μl/100ml streptomycin-penicillin G and 3% vitamin mix [55,56]. As in vitro encystation of *Eh* is not feasible yet, *Ei* is used as the model organism for the same. Late log phase cells were chilled to detach and harvested by centrifugation at 1500rpm for 5min at 4°C. $5x10^5$ cells/ml were transferred to 47% Low Glucose (LG) medium (TYI without glucose diluted 2.12 times with water, 2% heat inactivated adult bovine serum, 2.5% vitamin mix, 125μL/100ml antibiotic). Cysts cultured in LG medium were collected after 12, 24, 36, and 48h.

Cells were subjected to glucose starvation by overnight incubation in TYI-S-33 media devoid of glucose. Trophozoites of *Eh* and *Ei* were incubated at 42°C and 37ºC respectively for 1hr for heat shock. Oxidative stress was induced by incubating the cells in media containing 1.0mM of H_2O_2 for one hour [57, 58].

4.3. RNA isolation, cDNA synthesis, and real-time RT-PCR:

The total RNA was isolated using Trizol reagent (Ambion, USA) following standard protocol. The RNA quantity was measured using a NanoDrop spectrophotometer. The isolated RNA were treated with DNase I (Fermentas, USA) at 37°C for 1hr followed by enzyme inactivation at 60°C for 5 min. Genomic DNA contamination was checked by PCR amplification without RT, of actin or ARF gene. Total RNA and genomic DNA were used as template for negative and positive control, respectively.

cDNA was synthesized from 2μg of purified RNA template in 20 μl reaction volume using OligodT by First Strand cDNA synthesis kit (BioBharati, India) at 42°C for 50 min. The enzyme

was deactivated by incubating at 70°C for 15 min. Real-time RT-PCR was carried out following

our previously reported standard protocol in Eppendorf Master Cycler RealPlex, using 200ng

cDNA, 0.3µM gene specific primers (S1 Table) and PowerUpTM SYBR Green Master Mix

(Thermo Fischer Scientific, USA).The fold change of transcript expression was calculated using

the $\Delta\Delta C_T$method with respect to housekeeping genes like ADP-ribosylation factor (ARF) or

actin. The specificity of the amplicon was validated through melting curve analysis.

4.4. Cloning, expression, and purification of truncated EhTopo II

4.4.1. Cloning of *EhTopoII* fragment into a bacterial expression system

Despite using several combinations of vector-strain bacterial expression systems, the full length

(4050bp) expression of putative EhTopoII (EHI_120640) was not successful. Hence, a 1127bp

long fragment (from 760bp-1887bp) containing no tandem repeats of rare codons was PCR

amplified from *Eh* genomic DNA with the specific primers EhTopoIIfgS*Bam*HI and

EhTopoIIfgAS*Xho*I (S1 Table) using High Fidelity Taq polymerase (Thermo Fischer Scientific,

USA) and subcloned into pGEMT-Easy (Promega, USA) via TA ligation. This 1127bp fragment

shared 77% DNA sequence identity with the ortholog in *Ei* (EIN_145900) with no gaps and

carried regions of the putative catalytic site. Positive clones identified by blue-white screening

and confirmed by *Bam*HI-*Xho*I double digestion were sequenced and further cloned into the

pET21a expression vector. Restriction digestion was done for clone confirmation, and

recombinant vector was transformed into *E. coli* BL21 (DE3) expression strain.

4.4.2. Expression and purification of recombinant truncated EhTopo II protein

Overnight cultures of *E. coli* BL21(DE3) strain carrying recombinant pET21a vectors were

diluted to OD_{600} 0.05 and grown at 37°C till the OD_{600} was 0.6-0.8 at which recombinant protein

expression was induced using 1mM IPTG for 4 hours at 37°C. The recombinant protein was

solubilized using 0.25% S-lauryl sarcosine and further purified using Ni-NTA affinity

chromatography. The purified recombinant protein fragment was resolved on 12% SDS-PAGE.

4.5. Generation and purification of Anti-EhTopo II polyclonal antibody

Polyclonal antibody against recombinant EhTopo II fragment was commercially raised in rabbits

(Abgenex, India) and purified from the crude sera using Protein-A sepharose (Invitrogen, USA)

affinity chromatography. The sera were loaded over a pre-equilibrated Protein-A sepharose

column following 1:1 v/v dilution in PBS (pH 8). After sufficient washing to remove all unbound

proteins, the antibody was eluted using 100mM glycine (pH 2.8), and at least 20 fractions of

1mL each were collected into tubes containing 100µL of 1M Tris (pH 9). The absorbance of the

fractions at 280nm was measured, and the antibody-containing samples were pooled, dialyzed

over several changes of 1x PBS and concentrated.

4.6. Total protein extraction from *Entamoeba*

The *Entamoeba* cells grown at different conditions were harvested, washed with 1x PBS and

resuspended in lysis buffer containing 20mM Tris (pH 7.5), 1mM EDTA, 200mM NaCl, 1mM

PMSF, 1µg/mL of leupeptin and pepstatin,0.3µM of aprotinin, 15µM E-64, 1% Triton-X and

0.1% SDS and incubated for 30mins. The samples were then sonicated, centrifuged for 10mins at

10,000 rpm at 4ºC, and the supernatant was collected for further experiments. This method was

followed for samples from all conditions, including different kinds of stress and different hours

of encystation.

4.7. Western blot analysis

Recombinant as well as native protein samples were run on 12% SDS-PAGE and blotted onto

PVDF membrane at 70V for 2h. The membrane was blocked for 2h using 3% BSA in PBST

(0.1% Tween 20 in 1xPBS) and probed with anti-EhTopo II antibody (1:2000) for 2h at 4ºC.

Further, it was washed extensively with PBST and incubated with HRP conjugated goat anti-

rabbit IgG antibody (1:5000) for 90min at room temperature. After washing thrice with PBST,

the membrane was developed using Luminata Classico Western HRP substrate (Merck

Millipore, USA) and detected in ImageQuant LAS500 imager (GE Life Science, USA).

2.8. Nuclear extract preparation from *Entamoeba*

The harvested log phase trophozoites were washed thrice with 1x PBS and resuspended in 5x

volume ice-cold TEMP buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA, 4 mM MgCl2, 0.5 mM

PMSF), containing 2 μg/ml leupeptin and aprotenin, 1μg/ml pepstatin and μM E-64. After 30

min incubation on ice, the cells are lyzed using 0.5% Triton-X treatment for 10 mins. The lysate

is overlaid on a 1.5M sucrose cushion and spun at 12,000xg for 10mins. The pure nuclei pellet so

obtained is resuspended in TEP buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA, 0.5 mM PMSF)

containing 0.35M NaCl and all protease inhibitors and incubated on ice for 45 mins followed by

centrifugation at 12,000xg for 15 min. The supernatant is the nuclear extract which was assayed

for topoisomerase II activity.

4.9. Immunoprecipitation of native Topoisomerase II

Functional characterization of *Entamoeba* Topoisomerase II was carried out by

immunoprecipitation of the protein from crude nuclear extracts of log phase trophozoites using

anti-EhTopo II antibody. The nuclear extract was pre-cleared with pre-immune rabbit IgG (10

μg/ml) and 20 μl of protein-agarose beads (BioBharati, India) 1h at 4ºC. The pre-cleared

supernatant was collected by centrifugation and incubated with purified anti-EhTopo II IgG

(10μg/ml) at4 °C following which, 20 μl of protein-A agarose beads were added to precipitate

the antigen-antibody complex and incubated at 4°C for 2h. Agarose beads with bound immune

complexes were then harvested by centrifugation at 4°C and washed with nuclear extract buffer.

The immunoprecipitated topoisomerase II was confirmed by Western blotting and used for

further assays.

4.10. Topoisomerase II assay

The relaxation assay was carried out according to the protocol reported by Chakraborty and

Majumder, 1987 [59]. The standard relaxation assay mixture contained: 25mM Tris-HCl, pH

7.5, 10mM MgCl$_2$, 0.1mM EDTA, 1mM DTT, 2mM ATP, 50mM NaCl, 10% glycerol, 500ng of

supercoiled pure pL4440 plasmid substrate and different concentrations of nuclear extract or

immunoprecipitated protein solution as enzyme source. The reactions were carried out for 30min

at 25°C and 37°C for extracts from *Ei* and *Eh* respectively and arrested by adding dye containing

1% SDS and 10mM EDTA. The samples were electrophorized on 1% agarose gel at 1V/cm

overnight at room temperature and stained with ethidium bromide.

4.11. Staining and confocal microscopy

Entamoeba cells were harvested, washed with PBS (pH 7.6) and fixed with 2% p-formaldehyde

for 30mins at room temperature. The fixed cells were washed thoroughly and permeabilized with

0.1% Triton-X for 5mins blocked for 1h with 3% BSA and incubated overnight at 4ºC with anti-

EhTopo II antibody (1:100) in 1xPBS containing 1% BSA. It was followed with further washes

and incubation at room temperature with TRITC conjugated anti-rabbit IgG (1:400) (Sigma,

USA) for 1h. The nucleus was stained using 10µg/mL DAPI. The stained cells were visualized

and analyzed using Olympus FluoView FV1000 confocal microscope and software.

4.12. dsRNA mediated RNA interference studies

4.12.1. Cloning and expression of *EhTopoII* and *EiTopoII* specific dsRNA

In order to achieve downregulation at the RNA level, a 300bp and 250bp fragment within

EhTopoII and *EiTopoII,* respectively were selected as targets for RNA interference. Specificity

of the dsRNA was ensured by the selected regions within *Entamoeba TopoII* that did not contain

19mer homology to any other genes in *Eh* and *Ei*. These regions were amplified using specific

primers EiTopoIIdsF*Hin*dIII-EiTopoIIdsR*Xho*I and EhTopoIIdsF*Xba*I-EhTopoIIdsR*Xho*I (S1

Table), subcloned into pGEMT-Easy TA vector and then into expression vector pL4440, a gift

from Andrew Fire (Addgene) (http://n2t.net/addgene:1654) which is flanked by T7 promoter on

either side of its MCS. Recombinant plasmids carrying the inserts were transformed into the

RNase III-deficient, *E. coli* HT115 cells. The dsRNA expression was induced by 1mM IPTG

when OD_{600} reached 0.6, followed by incubation at 37ºC for 4h.

4.12.2. Extraction and purification of dsRNA

The gene specific dsRNA was extracted using water saturated phenol: chloroform: isoamyl

alcohol (25:24:1) followed by phase separation at high speed centrifugation. The RNA from the

aqueous layer was precipitated with equal volumes of isopropanol, further washed with 70%

ethanol, air dried and resuspended in nuclease-free water. Single-stranded RNA and DNA

contaminants were eliminated by treating the sample with 0.2µg/µL RNaseA (Sigma, USA) and

0.1U/µL DNase (Thermo Fischer Scientific, USA) for 1h at 37ºC. The samples were further

purified using Trizol, following standard protocol. The dsRNA from a 150 bp region in the

flavin mononucleotide based fluorescence protein of *B.subtilis* (BS1), with no sequence

similarity to *Entamoeba* genome, was used as negative control dsRNA.

4.12.3. dsRNA mediated downregulation of *Entamoeba* by soaking

Eh and *Ei* trophozoites were incubated in TYI or LG media containing 200 μg/ml of the

respective, purified TopoII-specific dsRNA for various time points, based on the experiment, and

the silencing efficiency was calculated using real-time RT-PCR [29]. All RNAi studies were

carried out with two controls: one without dsRNA and one with non-specific dsRNA (from *B.*

subtilis, as mentioned above).

4.13. Determination of cell viability and encystation efficiency

Effect of *Entamoeba Topo II* silencing and potency of various topoisomerase II drugs was

studied by assessing their effect on viability. Cell viability was determined by trypan blue dye

exclusion assay. Harvested cells, following various treatments, are resuspended in 1x PBS and

mixed with equal volumes of 0.4% trypan blue (Himedia, India). Dead cells take up the dye and

appear blue, while live ones remain white. The cell count is calculated at the start of the

experiment as well as following dye treatment, using a Neubauer hemocytometer.

After 48h of encystation, the total cell number (cyst+trophozoite) was calculated, and the

harvested cells were treated with 0.5% sarcosine for 10mins to lyze the trophozoites. The number

of cysts was then calculated and the encystation efficiency is represented as (cyst x

100)/(cyst+trophozoite).

4.14. Structure modeling and analysis

The three-dimensional protein structure of EhTopo II was predicted by homology modeling

using SWISS-Model server (https://swissmodel.expasy.org/). The template structure was

obtained from PDB based on sequence similarity and, refined using ModRefiner server

(http://zhanglab.ccmb.med.umich.edu/ModRefiner/). The generated structure was validated by

the Ramachandran plot using PROCHECK. PyMol software was used for the visualization of the

protein structure.

Acknowledgement

The authors thank Indian Institute of Technology Kharagpur for the research facilities and

Fellowship to SSV. The authors like to acknowledge FIST, DST, Govt. of India for the confocal

facility. The *E. coli* HT115 (DE3) strain was a kind gift from the Caenorhabditis Genetics

Centre, funded by the NIH National Centre for Research Resources, USA.

Reference:

1. McConnachie EW. The morphology, formation and development of cysts of *Entamoeba. Parasitol.*. 1969; 59: 41-53.

2. Avron B, Stolarsky T, Chayen A, Mirelman D. Encystation of *Entamoeba invadens* IP_1 is induced by lowering the osmotic pressure and depletion of nutrients from the medium. J Parasitol. 1986; 33: 522–525.

3. Sanchez LB, Enea V, Eichinger D. Identification of a developmentally regulated transcript expressed during encystation of *Entamoeba invadens*. Mol Biochem Parasitol. 1994; 67: 125-135.

4. Samanta SK, Varghese SS, Krishnan D, Baidya M, Nayak D, Mukherjee S, Ghosh SK. A novel encystation specific protein kinase regulates chitin synthesis in *Entamoeba invadens*. Mol Biochem Parasitol. 2018; 220: 19-27.

5. Singh T, Agarwal T, Ghosh SK. Identification and functional analysis of a stress-responsive MAPK15 in *Entamoeba invadens*. Mol Biochem Parasitol. 2018; 222: 34-44.

6. Sirijintakam P and Bailey GB. The relationship of DNA synthesis and cell cycle events to encystation by *Entamoeba invadens*. Arch Invest Med. 1980; 11(1): 3-10.

7. Silberman JD, Clark CG, Diamond LS, Sogin ML. Phylogeny of the genera *Entamoeba* and *Endolimax* as deduced from small-subunit ribosomal RNA sequences. Mol Biol Evol. 1999; 16: 1740-1751.

8. Sehgal D, Mittal V, Ramachandran S, Dhar SK, Bhattacharya A, Bhattacharya S. Nucleotide sequence organization and analysis of the nuclear ribosomal DNA circle of the protozoan parasite *Entamoeba histolytica*. Mol Biochem Parasitol. 1994; 67(2), 205-214.

9.   López-Casamichana M, Orozco E, Marchat LA and López-Camarillo C. Transcriptional

profile of the homologous recombination machinery and characterization of the EhRAD51

recombinase in response to DNA damage in *Entamoeba histolytica*. BMC Mol Biol. 2008; 9:

35.

10. Singh N, Bhattacharya S, Paul J. *Entamoeba invadens*: dynamics of DNA synthesis during

differentiation from trophozoite to cyst. Exp Parasitol. 2011; 127(2): 329-333.

11. Krishnan D and Ghosh SK. Cellular events of multinucleated giant cells formation during the

encystation of *Entamoeba invadens*. Front Cell Infect Microbiol. 2018; 8: 262.

12. Champoux JJ. Domains of Topoisomerase I and associated functions. Prog Nucleic Acid Res

1998; 60:111-132.

13. Nitiss J.L. DNA topoisomerase II and its growing repertoire of biological functions. Nat Rev

Cancer. 2009; 9: 327-337.

14. Topcu Z. DNA topoisomerases as targets for anticancer drugs. J Clin Pharm Ther. 2001; 26:

405-411.

15. Pourquier P. and Pommier Y. Topoisomerase I mediated DNA damages. Adv Cancer Res.

2010; 80: 189-216.

16. Hasinoff BB, Wu X, Patel D, Kanagasabai R, Karmahapatra S, Yalowich JC. Mechanisms of

the reduced cardiotoxicity of Pixantrone. J Pharmacol Exp Ther. 2016; 356(2): 397-409.

17. Janockova J, Korabecny J, Plsikova J, Babkova K, Konkolova E, Kucerova D et al. *In*

vitro investigating of anticancer activity of new 7-MEOTA-tacrine heterodimers.

J Enzyme Inhib Med Chem 2019; 34(1): 877-897.

18. Cheesman S, Horrocks P, Tosh K, Kilbey B. Intraerythrocytic expression of topoisomerase II

from *Plasmodium falciparum* is developmentally regulated. Mol Biochem Parasitol. 1998;

92: 39-46.

19. Majumder HK, Sengupta T, Dasgupta A, Das A. Topoisomerases of kinetoplastid parasites

as potential chemotherapeutic targets. Trends Parasitol 2004; 20(8): 381-387.

20. Chowdhury SR, Kumar A, Godinho JP, Silva ST, Zuma AA, Majumder HK.et al. Voacamine

alters *Leishmania* ultrastructure and kills parasite by poisoning unusual bi-subunit

topoisomerase IB. Biochem Pharmacol. 2017; 138: 19-30.

21. Chowdhury SR and Majumder HK. DNA topoisomerases in unicellular pathogens: structure,

function and druggability. Trends Biochem Sci. 2019; 44(5): 415-432.

22. Freeman CD, Klutman NE, Lamp KC. Metronidazole A therapeutic review and update.

Drugs. 1997; 54: 679–708.

23. Teimey LM, McPhee SJ, Papadakis MA. Current medical diagnosis and treatment 1999

(Appleton & Lange, Stamford, Conn).

24. Samarawickrema NA, Brown DM, Upcroft JA, Thammapalerd N, Upcroft P. Involvement

of superoxide dismutase and pyruvate: ferredoxin oxidoreductase in mechanisms of

metronidazole resistance in *Entamoeba histolytica.* J Antimicrob Chemother. 1997; 40:

833–840.

25. Wassmann C, Hellberg A, Tannich E, Bruchhaus I. Metronidazole resistance in the

protozoan parasite *Entamoeba histolytica* is associated with increased expression of iron-

containing superoxide dismutase and peroxiredoxin and decreased expression of

ferredoxin 1 and flavin reductase. J Biol Chem 1999; 274, 26051–26056.

26. Dingsdag SA and Hunter N. Metronidazole: an update on metabolism, structure-cytotoxicity

and resistance mechanisms. J Antimicrob Chemother 2017; 73: 265-279.

27. Goolsby TA, Jakeman B, Gaynes RP. Clinical relevance of metronidazole and peripheral

neuropathy: a systematic review of the literature. Int J Antimicrob Agents. 2017; 51: 319-

325.

28. Jeelani G, Sato D, Husain A, Cadiz AE, Sugimoto M. Metabolic profiling of the protozoan

parasite *Entamoeba invadens* revealed activation of unpredicted pathway during encystation.

PLoS ONE 2012; 7(5): e37740.

29. Samanta SK and Ghosh SK. The chitin biosynthesis pathway in Entamoeba and the role of

glucosamine-6-P isomerase by RNA interference. Mol Biochem Parasitol 2012; 186(1): 60-

68.

30. Wu CC, Li TK, Farh L, Lin LY, Lin TS, Yu YJ. Structural Basis of Type II Topoisomerase

Inhibition by the Anticancer Drug Etoposide. Science. 2011; 333 (6041), 459-462.

31. Bax BD, Chan PF, Eggleston DS, Fosberry A, Gentry DR. Type IIA topoisomerase

inhibition by a new class of antibacterial agents. Nature. 2010; 466: 935.

32. Meresse P, Dechaux E, Monneret C, Bertounesque E. Etoposide: Discovery and medicinal

chemistry. Curr Med Chem. 2004; 11: 2443.

33. Patel S, Jazrawi E, Creighton AM, Austin CA, Fisher LM. Probing the interaction of the

cytotoxic bisdioxopiperazine ICRF-193 with the closed enzyme clamp of human

topoisomerase IIα. Mol Pharmacol. 2000; 58: 560-8.

34. Classen S, Olland S, Berger JM. Structure of the topoisomerase II ATPase region and its

mechanism of inhibition by the chemotherapeutic agent ICRF-187. Proc Natl Acad Sci

U.S.A. 2003; 100(19): 10629-10634.

35. Mahmoudi N, Ciceron L, Franetich JF, Farhati K, Silvie O, Eling W, et al. *In vitro* activities

of 25 quinolones and fluoroquinolones against liver and blood stage *Plasmodium* spp.

Antimicrob Agents Chemother. 2003; 47: 2636–2639.

36. Nenortas E, Kulikowicz T, Burri C, Shapiro TA. Antitrypanosomal activities of

fluoroquinolones with pyrrolidinyl substitutions. Antimicrob Agents Chemother. 2003; 47;

3015–3017.

37. Yang X, Sladek TL, Liu X, Butler BR, Froelich CJ, Thor AD. Reconstitution of Caspase 3

Sensitizes MCF-7 Breast Cancer Cells to Doxorubicin- and Etoposide-induced Apoptosis.

Cancer Res. 2001; 61(1): 348-354.

38. Jiang H, Geng D, Liu H, Li Z, Cao J. Co-delivery of etoposide and curcumin by lipid

nanoparticulate drug delivery system for the treatment of gastric tumors. Drug Delivery

2016; 23(9): 3665-3673.

39. Ishida R, Sato M, Narita T, Utsumi KR, Nishimoto T, Morita T et al. Inhibition of DNA

topoisomerase II by ICRF-193 induces polyploidization by uncoupling chromosome

dynamics from other cell cycle events. J Cell Biol. 1994; 126 (6): 1341-1351.

40. Arlin Z, Mehta R, Feldman E. et al. Amsacrine treatment of patients with supraventricular

arrhythmias and acute leukemia. Cancer Chemother Pharmacol. 1987; 19: 163.

41. Ciavarra RP, Goldman C, Wen KK, Tedeschi B, Castora FJ. Heat stress induces

hsc70/nuclear topoisomerase I complex formation in vivo: evidence for hsc70-mediated,

ATP-independent reactivation in vitro. Proc Natl Acad Sci USA. 1994; 91: 1751-1755.

42. Simkova K, Moreau F, Pawlak P, Vriet C Baruah A, Alexander C et al. Integration of stress-

related and reactive oxygen species-mediated signals by Topoisomerase VI in *Arabidopsis*

thaliana. Proc Natl Acad Sci. USA. 2012; 109(40): 16360-16365.

43. Matsuo K, Kohno K, Sato S, Uchiumi T, Tanimura H, Yamada Y et al. Enhanced expression

of the DNA topoisomerase II gene in response to heat shock stress in human epidermoid

cancer KB Cells. Cancer Res. 1993; 53(5): 1085-1090.

44. Li TK, Chen AY, Yu C, Mao Y, Wang H, Liu LF. Activation of topoisomerase II-mediated

excision of chromosomal DNA loops during oxidative stress. Genes Dev. 1999; 13: 1553-

1560.

45. Hanke T, Ramiro MJ, Trigueros S, Roca J, Larraga V. Cloning, functional analysis and post-

transcriptional regulation of a type II DNA topoisomerase from *Leishmania infantum*. A new

potential target for anti-parasite drugs. Nucleic Acids Res. 2003; 31: 4917-4928.

46. Fief CA, Hoang KG, Phipps SD, Wallace JL, Deweese JE. Examining the impact of

antimicrobial fluoroquinolones on human DNA topoisomerase IIα and IIβ. ACS Omega.

2019; 4(2): 4049-4055.

47. Dang Z, Yang Y, Ji R, Zhang, S. Synthesis and antibacterial activity of novel

fluoroquinolones containing substituted piperidines. Bioorg Med Chem Lett 2007; 17(16):

4523-4526.

48. Pokrovskaya V, Belakhov V, Hainrichson M, Yaron S, Baasoy T. Design, synthesis and

evaluation of novel fluoroquinolone-aminoglycoside hybrid antibiotics. J Med Chem. 2009;

52(8): 2243-2254.

49. Mudeppa DG, Kumar S, Kokkonda S, White J, Rathod PK. Topoisomerase II from human

malaria parasites: expression, purification, and selective inhibition. J Biol Chem 2015;

290(33): 20313-20324.

50. Anquetin G, Greiner J, Mahmoudi N, Santillana-Hayat M, Gozalbes R, Farhati K et al.

Design, synthesis and activity against *Toxoplasma gondii*, *Plasmodium* spp., and

Mycobacterium tuberculosis of new 6-fluoroquinolones. Eur J Med Chem. 2006; 41(12):

1478-1493.

51. Zhao X, Zhao Y, Ren Z, Li Y. Combined QSAR/QSPR and molecular docking study on

fluoroquinolones to reduce biological enrichment. Comput Biol Chem. 2019; 79: 177-184.

52. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: molecular evolutionary

genetics analysis version 6.0. Mol Biol Evol. 2013; 30(12): 2725–2729.

53. Marchler-Bauer A and Bryant SH. CD-Search: protein domain annotations on the

fly. Nucleic Acids Res. 2004; 32: W327–W331.

54. Kosugi S, Hasbe M, Tomita M, Yanagawa H. Systematic identification of yeast cell cycle-

dependent nucleocytoplasmic shuttling proteins by prediction of composite motifs. Proc Natl

Acad Sci USA. 2009: 106: 10171-10176.

55. Diamond LS. Axenic cultivation of *Entamoeba histolytica*. Science. 1961; 134: 336–337.

56. Diamond LS, Harlow DR, Cunnick CC. A new medium for the axenic cultivation of

Entamoeba histolytica and other *Entamoebas*. Trans R Soc Trop Med Hyg. 1978; 72: 431-

432

57. Field J, Dellen K, Ghosh SK and Samuelson J. Responses of *Entamoeba invadens* to heat

shock and encystation are related. J Eukaryot Microbiol. 2000; 47: 511-514.

58. Tovy A, Hertz R, Siman-Tov R, Syan S, Faust D, Guillen N et al. Glucose starvation

boosts *Entamoeba histolytica* virulence. PLoS Negl Trop Dis. 2011; 5(8): e1247.

59. Chakraborty AK and Majumder HK. Decatenation of kinetoplast DNA by an ATP-dependent

DNA topoisomerase from the kinetoplast hemoflagellate *Leishmania donovani* Mol Biochem

Parasitol. 1987; 26: 215-224.

Supporting information caption

S1 Table: List of primers

S2 Table: List of putative topoisomerase in *Entamoeba*

S1 Fig: Microarray data for putative topoisomerase of *E. invadens* during encystation

S2 Fig: Expression and purification of recombinant truncated EhTopo II.

S3 Fig: Western blot analysis assaying anti-Topo II antibody

S4 Fig: Expression of *Entamoeb*a TopoII specific dsRNA and dsRNA mediated silencing of

TopoII by soaking

S5 Fig: Homology modeling of EhTopoII structure and its validation

722
723