Title: Artisanal and farmer bread making practices differently shape fungal species community

composition in French sourdoughs

Running title: Bread making practices as a driver of yeast species community composition

Authors

Elisa Michel[†,°], Estelle Masson[‡], Sandrine Bubbendorf[‡], Léocadie Lapicque[‡], Thibault Nidelet[†], Diego

Segond[†], Stéphane Guézenec[†], Thérèse Marlin[†], Hugo Devillers[†], Olivier Rué[¶], Bernard Onno[°], Judith

Legrand[§*], Delphine Sicard[†] *and the participating bakers[#]

Authors affiliation

[†] SPO, Univ Montpellier, INRAE, Institut Agro, Montpellier, France

[°] Oniris, Laboratoire MicrobioTech, UMR GEPEA 6144, Rue de la Géraudière CS 82225, 44322 Nantes

Cedex 3, France

[‡]Laboratoire de Psychologie : Cognition, Comportement, Communication – EA 1285, Université de

Bretagne Occidentale, 20 rue Duquesne, CS 93837, F-29238 Brest 03, France

[§]Génétique Quantitative et Evolution-Le Moulon, Université Paris-Sud, INRAE, CNRS,

AgroParisTech, Université Paris-Saclay, 91190 Gif-sur-Yvette, France

[¶]Université Paris-Saclay, INRAE, MaIAGE, 78350, Jouy-en-Josas, France

[¶]Université Paris-Saclay, INRAE, BioinfOmics, MIGALE bioinformatics facility, 78350, Jouy-en-

Josas, France

22
23

*Judith Legrand and Delphine Sicard should be considered both last authors.

Corresponding author

Delphine Sicard, SPO, Univ Montpellier, INRAE, Institut Agro, Montpellier, France +33 624720501

delphine.sicard@inrae.fr

Key words: domestication, yeast, *Kazachstania*, *Saccharomyces cerevisiae*, fermentation

No competing interests

Financial supports: French National Research Agency (ANR-13-ALID-0005) and INRAE

Abstract

Preserving microbial diversity in food systems is one of the many challenges to be met to

achieve food security and quality. Although industrialization led to the selection and spread of specific

fermenting microbial strains, there are still ongoing artisanal processes that may allow the conservation

of a wider species and genetic diversity. We examined whether the diversity of artisanal practices could

lead to an increased level in fungal species diversity for bread making. We used an interdisciplinary

participatory research approach including bakers, psycho-sociologists and microbiologists to analyze

French bread making practices and describe fungal communities in naturally fermented sourdough of

27 bakers and 12 farmer bakers. Bread making practices were classified in two groups: the farmer-like

practices group and the artisanal-like practices group. The well-known bakery yeast, *Saccharomyces*

cerevisiae, was dominant (i.e. with a relative abundance over 50%) in only 24% of sourdoughs while

other yeast species belonging to the *Kazachstania* genus were dominant in 54% of sourdoughs. Bread

making practices were found to drive the distribution of fungal species across sourdoughs. The most

striking bread making practice effect was the occurrence of *Kazachstania humilis* in sourdoughs made

with artisanal-like practices and the occurrence of *Kazachstania bulderi* in sourdoughs made with

farmer-like practices. Phenotypic divergences between sourdough and non-sourdough strains were

found for *K. humilis* but not for *K. bulderi*. Overall, our results showed that preserving bread making

practice diversity allows the preservation of a higher species and phenotypic diversity in microbial

communities.

Introduction

Humans started to ferment food before the Neolithic using naturally fermenting microbial communities. In the 19th century, the industrialization and the increase of knowledge in microbiology resulted in changes in fermented food practices with the use of starters. This selection led to a reduction in species diversity and genetic diversity for fermented food processing and limited *in situ* conservation of microbial communities in industrialized systems [1–3]. Domestication of the yeast *Saccharomyces cerevisiae* for the production of beer, wine, cheese, leavened bread, that of the fungi *Penicillium roqueforti* or *Penicillium camemberti* for cheese production or that of the fungus *Aspergillus oryzae* for rice or soybean fermented products are well studied cases [1–10]. The recent renewed interest in artisanal practices that make use of naturally fermenting microbial communities could promote the conservation of microbial diversity. However, the effect of artisanal practices on the distribution of microbial species across sourdoughs remains poorly documented.

Among fermented foods, bread is still a symbol deeply engrained in the history, religious rites and medicine of several cultures. Bread likely originated 14 000 years ago, suggesting that bread was made long before plant domestication [11]. Since the Neolithic, bread history is intimately associated with the domestication of cereals, bread making associated tools and the advent of Mediterranean civilizations [12]. Investigation of the morphology of plant remains which were incorporated in Neolithic bread identified wheat, barley, millet, linseed [12]. Leavened bread was traditionally made with flour, water and a fermenting agent, which was either a fermenting beverage or a fermenting dough, called sourdough. This sourdough was generally initiated from a mixture of flour and water, naturally colonized by lactic acid bacteria (LAB) and yeasts. Sourdough was then either maintained over time or initiated again and again, depending on the craftsman [13, 14]. In the 19th century, the use of yeast starters made of *S. cerevisiae,* often called « baker's yeast », spread as an alternative to sourdough. Nowadays, *S. cerevisiae* industrial starters are more frequently used than sourdoughs, although the latter are gaining interest. A recent study showed that industrial populations of *S. cerevisiae* have followed a different evolutionary path than sourdough populations of *S. cerevisiae* [10]. Both have been

domesticated by humans which have improved their fermentation performance in a sourdough-

mimicking medium. Industrial and sourdough strains of *S. cerevisiae* differ genetically and

phenotypically, indicating that sourdough use contributes to the conservation of bread related *S.*

cerevisiae lineages [10].

Beyond *S. cerevisiae*, sourdough can also host other yeast species. Yeasts, whether ascomycetes

or basidiomycetes, are generally characterized as fungi, that asexually reproduce by budding or fission,

which results in growth that is comprised mainly of single cells. Their sexual states are not enclosed in

fruiting body. To date, more than 40 yeast species have been detected in sourdough [1, 15, 16]. The

most frequently encountered species are *Wickerhamomyces anomalus* and *Kazachstania humilis*.

Several other species in the genus *Kazachstania* (*Kazachstania barnettii*, *Kazachatania exigua*,

Kazachstania bulderi, *Kazachstania unispora*) as well as several species in the polyphyletic genus

Pichia have also been recurrently detected. The factors that determine the presence in sourdough of

these species are still unknown. A recent large-scale study of 500 sourdoughs from four continents found

no effect of geography or factors related to bread making practices such as age of sourdough, storage

location, feeding frequency, or grain intake [17]. However, most of the sourdoughs in this study were

made by private citizens who probably did not maintain the sourdough microbial community in the same

way as professional bakers. To our knowledge, no studies have been conducted to date to investigate

the effect of bakers' bread making practices on sourdough yeast community composition.

In France, sourdough breads are made both by bakers and farmers who also grow and mill their

own wheat. The number of farmer-bakers has increased in the 2000s with two motives: to grow wheat

varieties meeting their needs and to assert their independence from industry [18]. Although farmer-

bakers are less numerous than bakers, they participate in the renewed interest in local wheat varieties

and artisanal know-how, which may contribute to the conservation of both socio-cultural diversity and

microbial diversity.

Here, we used a participatory research approach involving psycho-sociologists, biologists, bio-

statisticians, bakers and farmer-bakers to study whether and how bakers and farmer-bakers contribute

to the preservation of socio-cultural and fungal species diversity in sourdough microbial community.

Materials and methods

111   A total of 27 bakers and 12 farmer-bakers participated to the study. They were all making bread with

112   organic flour except five. All of them sent sourdough to the lab for microbiological and metabolic

113   analysis. Among them, 36 described their bread making practices as well.

A questionnaire survey, face-to-face interviews and focus-groups to collect bread making practices

Data on bread making practices were collected through a questionnaire survey, interviews and focus

groups. The collected variables were related to *i)* the ingredients origin : wheat varieties types (ancient

populations also called landraces / modern varieties), whether they produced flour from their own wheat,

whether they had their own mill or use an external mill, water origin, *ii)* the sourdough recipe: its age,

its hydration state, the origin of the chief sourdough (sample of dough or sourdough), the number of

back-sloppings before bread making and per week, the temperature of water used for back-sloppings,

iii) their bread making practices: the number of bread makings per week, the percentage of sourdough,

flour and salt in bread dough, the kneading methods, the total duration of fermentation and the addition

of baker's yeast in dough.

Sourdoughs samples, enumeration and strain isolation

126 Sourdoughs were collected before kneading and referred to as final sourdoughs (Table S1). On the day

127 of collection, they were sent to the lab where yeast and bacteria were enumerated and isolated as in [19,

128 20], and sourdoughs stored at -20°C in sterile vials for non- culture based analysis. Ethics and rights

129 associated with sourdough collection and strains isolation have been respected.

Sourdough acidity and metabolic analyses

For each sourdough, three independent 1-g replicates were analyzed. pH and Total Titrable Acidity were

measured as described in [20]. Organic acids, alcohol and sugars concentrations (expressed as g/kg of

sourdough) were analyzed by liquid chromatography using an HPLC HP 1100 LC system (Agilent

technologies, Santa clara, CA, USA) equipped with a refractive index detector (RID Agilent G1382A)

and a UV detector (Agilent G1314A). Two different columns were used, a Rezex ROA-organic acids

column and a Rezex RPM-monosaccharide column (SDVB – Pb+2 8%, 300x7.8mm, Phenomenex,

Torrance, CA, USA). The details of the experiments are described in supplementary information

(Method S1).

Yeast species identification

The Internal transcribed spacer 1 (ITS1) ribosomal DNA of each 1216 yeast isolates was amplified by

PCR from chromosomal DNA, either by using primers ITS1F and ITS2 [21, 22], or primers NSA3 and

58A2R [19, 22]. For isolates unidentified with the ITS1 region alone, DNA was extracted with the

MasterPure yeast DNA purification kit (Epicentre, Epibio). PCR reactions targeting partial genes, the

D1D2 region of the large subunit of rRNA (LSU), a part of the RNA polymerase II large subunit

encoding gene (*RPB1)*, a part of the RNA polymerase II encoding gene (*RPB2)*, a part of the actin

encoding gene (*ACT1)* and transcriptional elongation factor (TEF) were performed. To discriminate

three specific isolates, PCR on genes *GHD1*, *FSY1*, *URA3*, *DRC1*, *MET2* were performed [23–26]

(Table S2). All PCR products were sent to be sequenced with Sanger sequencing (Eurofins, Germany).

Species were identified using NCBI [27], YeastIP [28] and a personal database, which was constructed

after *ITS1*, *RPB2*, LSU sequencing of all 33 yeast species reportedly found in sourdoughs in the literature

[19].

Sourdough DNA extraction, MiSeq sequencing, bioinformatics

The ITS1 region was targeted with the PCR primers ITS1-F (5′- CTTGGTCATTTAGAGGAAGTAA - 3′) and ITS2 (5'-GCTGCGTTCTTCATCGATGC-3').

The sequencing run was performed with MiSeq Reagent Kit v3. 2015 [20]. Sequences were analyzed through FROGS "Find Rapidly OTU with Galaxy Solution" [29] and home-made pipelines. Overlapped reads were merged with Flash [30] with a minimum overlap of 10 nucleotides, a maximum overlap of 300 nucleotides and a maximum mismatch density of 0.1. Primers were removed with Cutadapt [31] and data were cleaned with Sickle with quality-threshold and length-threshold equal to 20 [32]. Reads were clustered with Swarm (d=3) [33] and chimeras deleted with VSEARCH [34]. Sequences were then filtered on minimum abundance of 0.005% of all sequences. From the OTU abundance table and for each OTU, the taxonomic affiliation using UNITE Version 7.1, Release 2016-11-20 [35], YeastIP [28] and our own databases [19] was obtained by blasting OTUs representative sequences against each database.

Phenotypic analysis of yeast strains

Fermentation performance of the two most frequently encountered *Kazachstania* species was assessed as described in [10] for *S. cerevisiae*. Fifteen sourdough strains of *K. bulderi* and 16 sourdough strains of *K. humilis* were included in the analysis. *Kazachstania bulderi* strains were coming from sourdoughs B4, B12, B15, B17 B20, B21, while *K. humilis* strains were coming from sourdoughs B2, B5, B6, B7, B10, B17. From one to three strains per sourdough were analyzed in the experiment. In addition, four strains of *K. bulderi* (strain MUCL 38021 isolated from silage in Namur, Belgium, strain MUCL 54694 isolated from silage in Erezée, Belgium, strain NRRL Y-27205 and strain CLIB 604 isolated from maize silage in the Netherland) and three strains of *K. humilis* (strain CBS 7754 isolated from food dressing, strain CLIB 1323 isolated from bantu beer, and strain CBS 2664 isolated from alpechin) which were coming from non-sourdough habitats, were added as control to test the effect of habitat of origin. Each strain was phenotyped at least in triplicate leading to a total of 145 fermentations distributed over two

177   blocks. Briefly, fermentations were carried out at 24 °C with constant magnetic stirring (300 rpm) during

178   24 h. CO_2 release was measured by weight loss every 40 min using an automated robotic system [36].

179   At the end of fermentation, population size and cell viability were determined by flow cytometer (C6

180   cytometer, Accuri, BD Biosciences) as described in [37].

Data analyses

A multiple correspondence analysis (MCA) and hierarchical clustering (complete linkage clustering

method) on principal components based on the first two axes of the MCA were performed using the

FactoMineR R package [38].

To analyse fungal community, weighted Unifrac distances between sourdough communities were

computed from a rooted phylogenetic tree based on the OTU sequences using the R-packages Phyloseq

and GUniFrac [39, 40]. Phylogenetic sequences were aligned with Clustalo and phylogenetic trees were

built with the parsimony algorithm, with 100 replicates bootstraps, pairwise ktuple-distances with

Seaview [41]. The results presented in the main text were obtained using the phylogenetic tree rooted

on the OTU identified as *Sporidiobolales species*. Different roots were tested. The roots were chosen

among the OTUs that were affiliated to the most distant taxa (*Sporidiobolales species*, *Bullera*

globospora, *Trichosporon asahii*, *Udeniomyces pyricola*). Tree architecture did not change with the

root. The tree did not fit the expected phylogeny and, notably, some *Ascomycota* were located among

the *Basidiomycota*. However, the dominant sourdough species belonging to the *Saccharomycetaceae*

family were clustered in the expected clades or subclades, except that *Kazachstania servazzi* and

Kazachstania unispora were grouped in a clade closer to *Saccharomyces* species than to other

Kazachstania species. Using the Unifrac distances matrix, we performed a principal coordinate analysis

(PCoA) and clustered sourdough communities using the first two axes of the PCoA, and the complete

linkage clustering method (hclust R function). To check the sensitivity of our analysis to this

misclassification, we performed the same analyses without the sourdoughs that had one misclassified

species representing more than 10% of their reads, *i.e.* sourdoughs B20, B41, B42, and B44 and found

the same clustering [40].

For each sourdough, the species richness, Chao1, Shannon and Simpson indexes were computed. Chao1

was used as an indicator of species richness corrected by the number of OTUs present in the community

but not observed. Shannon and Simpson index values were converted to the effective number of species

per sourdough. This number was estimated from the Shannon diversity index as $exp^{Shannon index}$ and

from the Simpson diversity index as $\frac{1}{1-Simpson index}$ [42, 43]. For probability estimates, the exact 95%

confidence intervals were computed using a binomial distribution.

To study the links between β-diversity and differences in bakery practices, we performed a

permutational multivariate analysis of variance (PERMANOVA) on the Unifrac distance matrix for

each bakery practice variable. We included in the analysis sourdough fungal communities of the 30

bakers who had less than 8 missing values among the 29 bread making practices variables and adjusted

the p-value using false discovery rate method correction to account for multiple testing [44]. In addition,

we performed independence exact Fisher tests between fungal community PCoA groups and each of the

bread making practices variables. Multiple testing was accounted for using the false discovery rate

method [44].

In addition, we tested the link between the baker practices group, the fungal community group or the

yeast dominant species and the variation of each quantitative variable (microbial density, pH, TTA,

metabolite concentration) with the following mixed effect model: $Y_{ijk} = \mu + \alpha_i + B_j + \varepsilon_{ijk}$ with ε_{ijk}

$\sim N(0,\sigma^2)$, where α_i is the effect of the fungal community group i modelled as a fixed effect and B_j is

the effect of sourdough j modelled as a random effect and k represents the measurement replicates. For

sourdough hydration rate, the variable was arcsin transformed but sourdough effect was not included in

the model because no repetition was obtained from any sourdough. The model parameters were

estimated using the lmertest R package [45]. To test the fixed effects, we used likelihood ratio tests.

Multiple comparisons of means were performed using Tukey tests with the multcomp package. p-values

were all adjusted for multiple testing with the FDR method. The geographical structuration was tested

with a Mantel test on the Unifrac distances matrix and the geographical distances matrix computed with

the package geosphere [46] and ade4 [47].

The phenotypic diversity of *K. bulderi* and *K. humilis* strains coming from sourdough and non-

sourdough habitats were analyzed. Population size and mortality rate after 27h of fermentation measured

by flow cytometer were used as proxies for absolute fitness. The cumulative CO_2 production curve was

calculated and the kinetics of CO_2 production rate over time was estimated by successive linear

smoothing over five points. Four fermentation parameters were then estimated. The maximum CO_2

release (CO_2max, in g/L) was estimated by the maximum of the cumulative CO_2 production curve. The

fermentation latency phase time was estimated by the time between inoculation and the beginning of the

fermentation calculated as 1g/L of CO_2 release (t1g, in h). The maximum CO_2 production rate (Vmax

in g/L/h) was estimated by the maximum of the CO_2 production rate kinetic. The time of the Vmax

(tVmax in h) was calculated as the time between inoculation and the Vmax. Hence, the phenotype of

each strain was characterized by six quantitative variables called "phenotype variables" below: its

population size, its mortality rate and the four fermentation parameters. To determine whether the origin

of the strain (sourdough or non-sourdough) had an impact on strain phenotype, each log-transformed

quantitative variable was analyzed separately using a mixed linear model as described below. The

experimental design was unbalanced between the two blocks with very few non-sourdough strains in

one of the two blocks. Therefore, for each phenotype variable, we first estimated the block effect with

a subset of 8 strains cultivated in both blocks using a linear model with two fixed effects: the strain and

the block. Additive models were used as the interaction terms were not significantly different from zero

after adjusting p-values with the Benjamini-Hochberg method. Second, each phenotype variable was

corrected for the block effect and analyzed with the mixed effect model: $Y_{ijkl} = \mu + \alpha_i + \beta_j + \gamma_{ij} +$

$Z_k + \epsilon_{ijkl}$ where Y_{ijkl} represents the log-transformed phenotype variable corrected for the block effect

for the strain *k,* from species *i (i=1,2),* sampled in environment *j (j=1,2),* observed for replicate *l.* μ

represents the mean of the phenotype variable, α_i the additive effect of species *i,* β_j the additive effect

of environment j, and γ_{ij} their interaction. Z_k represents the gaussian random effect of strain k with $Z_k \sim$

$N(0, \sigma_s^2)$ and ϵ_{ijkl} the gaussian residuals with $\epsilon_{ijkl} \sim N(0, \sigma^2)$. For each species i, the impact of the

environment was quantified using the contrast $\Delta_i = \beta_S + \gamma_{i,S} - \beta_{NS} - \gamma_{i,NS}$ with NS standing for

"non-sourdough" and "S" for "'sourdough" and tests ($H_0 = \Delta_i = 0, H_1: \Delta_i \neq 0$) were performed and

p-values were adjusted using the Benjamini-Hochberg method. As log-transformed data were analyzed,

the exponential of this contrast can be interpreted as the ratio between the sourdough mean and the non-

sourdough mean. Confidence intervals and tests were performed using the doBy R package.

All statistics and plots have been done with R (ggplot2 [48], leaflet package [49], with minor esthetical

adjustment with Inkscape). Data and scripts are shared on Zenodo.

http://doi.org/10.5281/zenodo.2600170

264 Results

265 Two groups of bread making practices

A total of 39 French bakers producing natural sourdough bread and distributed all over France

participated to the study (Table S1). The bread making practices of 35 of them were collected through

one or several methods: personal interviews (with 12 bakers), focus groups (three groups), observation

during break-making workshops (two workshops), and an online/phone survey (36 bakers). The general

process of sourdough bread making was presented in Figure 1. We analyzed 28 variables, describing

variations of the practices at all steps of the bread making process, from wheat grains to baked bread

(Figure S1). Four bakers (B6, B10, B19, B20) who did not provide enough information about their

practices were excluded from the multivariate analysis. According to a hierarchical clustering on

principal components (HCPC), the 32 other bakers clustered into two groups corresponding to two main

types of bread making practices (Figure 2). The first group, hereafter termed "farmer-like" practices

group, included six bakers and 11 farmer-bakers using the following practices: low bread production

(<500 kg per week, 81% of the bakers of the "farmer-like" group), use of wheat landraces (56%), manual

kneading (63%), working at ambient temperature (88%), long fermentation periods (more than 4 hours

for 88%), and no use of commercial baker's yeast (88%). In addition, they tend to make their chief

sourdough from dough after kneading (75%). The second group, hereafter called "artisanal-like"

practices group consisted of 12 bakers and four farmer-bakers having more intensive practices,

characterized by a large bread production (>500 kg per week, 81%), mechanical kneading (100%), use

of modern wheat varieties (63%), working at ambient temperature (56%), using *S. cerevisiae* starters in

addition to sourdough for bread making or for pastries and buns making (81%). In this second group,

bakers tended to make their chief sourdough from a final sourdough.

Composition of sourdough fungal communities

Sourdough is a mix of flour and water naturally fermented by bacteria and yeasts. Sourdough yeast

density ranged from $8.1\ 10^4$ to $5.8\ 10^8$ CFU per gram of sourdough, with a mean value of $2.9\ 10^7$ CFU

per gram, as commonly found in sourdoughs from all over the world [1, 16, 50, 51]. We isolated 20 to

40 yeast strains from each sourdough by picking colonies randomly and identified species using ITS

sequence as well as other barcodes when the ITS alone was not able to discriminate between closely

related species. Among the 39 collected sourdoughs, one (sourdough B14) did not give any colony in

the laboratory, suggesting that his sourdough microbiota was no longer alive. A total of 1216 strains

were characterized from the other 38 sourdoughs. In addition, we developed an ITS1 meta-barcoding

MiSeq sequencing method on sourdough (see sup M&M). After filtering 5,360,620 raw ITS1 sequences

for quality, abundance (0.005%) and chimera, 3,542,801 sequences were further analyzed. Overall, the

sequences clustered in 113 OTUs. The number of reads per sourdough ranged from 8421 to 194,557.

Therefore, we carried out our analysis on the rarefied matrix. Among all OTUs, 10 were assigned to the

order *Triticodae* (especially to the species *Triticum aestivum*), 50 were assigned to a filamentous fungi

genus including plant pathogen species such as *Alternaria*, *Aspergillus*, *Fusarium*, or *Gibberella*, while

4 OTUs remained unidentified. Among the 40 yeast OTUs, 96% of total reads were assigned to the

phylum *Ascomycota*, 87.5% to the order *Saccharomycetales* and 85.7% to the family

Saccharomycetaceae. Only 4% of the total reads were assigned to the phylum *Basidiomycota*. Overall,

three OTUs assigned to the species *Kazachstania humilis*, *Kazachstania bulderi* and *Saccharomyces*

cerevisiae represented 20.3%, 15.5% and 24.1% respectively of the total number of reads and 28.1%,

23.7% and 18.2% respectively of the number of reads identified as yeast species (Figure 3).

Both non-culture-based and culture-based methods allowed the identification of the same dominant

species (defined as a species with an over 50% frequency) for all sourdoughs but five (B09, B20, B22,

B25, B41) (Figure 3). In two cases, the discrepancy was explained by the detection of *Cladosporium*

genus at high frequency with metabarcoding while this species could not be isolated in the laboratory

(Figure 3). In two other cases, it was explained by a high number of *S. cerevisiae* isolated in the

laboratory compared to what was observed using metabarcoding sequencing. In the last case, the

identification of *Pichia kudriavzevii* required additional sequencing as it shares an identical ITS with

Candida xylopsoci. Because metabarcoding allows a deeper characterization of the fungal species

diversity with few discrepancy cases, the distribution of fungal species across sourdoughs will be further

described using metabarcoding data only. Previous analysis of the same sourdoughs revealed that

Fructilactobacillus sanfranciscensis was the dominant bacterial species in all analyzed sourdoughs but

two, where the dominant species was either *Latilactobacillus curvatus* or *Companilactobacillus*

heilongjiangensis [20, 52, 53]. Therefore, we decided to study the species composition of fungal

community only.

Fungal species diversity within and between sourdoughs

All sourdoughs but two had a dominant yeast species with a relative abundance over 50% and

many species with a lower relative abundances (Figure 3). Within sourdoughs, fungal species richness

ranged from 10 to 33, with a 23 median (Table S3). The effective number of species per sourdough

calculated from the Shannon diversity index ranged from 1 to 7 (Table S3), with 70% of sourdoughs

having an index below two (Table S3). The bread making practice group (artisanal-like/farmer-like) did

not influence significantly the level of fungal α-diversity in sourdough (Wilcoxon rank exact test,

$W_{shannon}$ = 156, p-value = 0.16, $W_{simpson}$ = 165, p-value = 0.08). Between-sourdough ß-diversity

were analyzed using weighted Unifrac distances, computed from a phylogenetic tree built from the

distances between OTUs using *Sporidiobolales* species as root (Figure S2). Unifrac distances computed

with four differently rooted trees were highly positively correlated (Figure S3). Unifrac distances

between sourdoughs ranged from 0.0005 and 0.71, with a median of 0.49 and a mean of 0.52. The

clustering of sourdoughs according to their Unifrac distances is shown Figure 5. There was no significant

correlation between the Unifrac distances and geographical distances between sourdoughs (Mantel test,

P=0.35).

We then analyzed specifically the occurrence of yeast species in sourdoughs as yeasts, together

with lactic acid bacteria, are the main functional players in a sourdough ecosystem and for bread quality.

Over the 40 yeast species detected in the 38 sourdoughs, 12 had a relative abundance over 50% in at

least one sourdough, four had a relative abundance between 20% and 50% and 24 had a relative

abundance below 10%. All dominant species (relative abundance over 50%) were fermentative yeast

species, except in one sourdough that had a C*ladosporium* species. We found all the sourdough yeast

genera (*Saccharomyces*, *Kazachstania*, *Pichia*, *Torulaspora* and *Hyphopichia)* commonly reported in

the literature except the *Wickerhamomyces* genus that we did not detect in our samples [1, 15, 16].

The baker's yeast species, *Saccharomyces cerevisiae* is not the most widespread yeast

species in French organic sourdoughs

Saccharomyces cerevisiae was found in 53% of all sourdoughs (95% confidence intervals=36%

- 69%) but was dominant (relative abundance over 50%) in only 24% (95% confidence intervals=11%

- 40%) (Figure 3). In two cases, *S. cerevisiae* co-occurred with another yeast species at similar relative

abundance. In the first case, *S. cerevisiae* was present at a relative abundance of 40% with *Candida sake*

at a 41% relative abundance. In the second case, it was found at a relative abundance of 47% with *Pichia*

kudriavzevii at a relative abundance of 52%. In all the other cases, *S. cerevisiae* had a relative abundance

below 21% and was found with other dominant yeast species, such as *Kazachstania australis*,

Kazachstania humilis, *Saccharomyces uvarum* or *Torulaspora delbrueckii*. This suggests that *S.*

cerevisiae did not displace other species and can indeed be out-competed by other species in sourdoughs.

Sourdough yeast species mostly belong to the *Kazachstania* genus

Kazachstania was the most represented yeast genus over all sourdoughs, when considering both the number of reads over all sourdoughs and the number of detected species. This genus represented 57% of the total number of reads while *Saccharomyces* represented 26% of the total number of reads. In addition, eight species of the *Kazachstania* genus were found in sourdough, while the *Saccharomyces* genus was represented by two species (*S. uvarum* and *S. cerevisiae*) (Figure 3). The *Kazachstania* genus is one of the closest genetically related genus to *Saccharomyces* and contained Crabtree positive yeasts, able to ferment glucose even when oxygen is present if the amount of sugar is sufficient (Hagman & Piskur 2015). *Kazachstania* species dominated in 54% (95% confidence intervals=36%-69%) of sourdoughs while *Saccharomyces* species dominated in 27% only (95% confidence intervals=13%-43%). *Kazachstania humilis*, followed by *K. bulderi* were the most commonly dominant *Kazachstania* species, and found in respectively 21% (95% confidence intervals=10%-37%) and 15% of sourdoughs (95% confidence intervals=6%-31%) (Figure 3). A recently described *Kazachstania* species, *Kazachstania bozae*, was also identified in five sourdoughs (4.5%-29%) and found dominant in three (1.7%-22%) [64]. Strains of this species were closely related to a strain previously isolated from boza, a Bulgarian fermented drink, as estimated with ITS and LSU (D1D2) barcodes (Source: NCBI, GenBank: KC118125.1 and KX369579.1). In addition, *Kazachstania saulgeensis,* a recently described species [65, 66], was dominant in one sourdough (0.07%-14%). *Kazachstania unispora* and *Kazachstania servazzi* which had previously been detected in sourdough were also found [17, 17, 53, 57, 58, 63, 67–71]. Some *Kazachstania* species were detected for the first time as dominant in sourdoughs, whereas they had been previously found in other environments, like soil (*K. australis*), sauerkraut (*K. barnettii*) [72–74]. None of the previous studies on sourdough have observed as many *Kazachstania* species in sourdough.

The composition of sourdough fungal communities was associated with differences in bread making practices

We tested whether sourdough fungal community beta diversity could be explained by bread making practices. To do so, we performed univariate PERMANOVA analysis on the 30 bakers with fewer than 8 missing values for the 29 bread making practices variables (Table S4). The univariate analysis revealed that the weighted Unifrac distance was structured according to the use of commercial yeast in bakery ($P<0.05$). It also varied significantly with sourdough age, chief sourdough origin (dough, sourdough or both), the quantity of bread produced per week, the milling method (cylinder, millstone, Astrie, Tyrol), the type of wheat variety (ancient, modern or a mix thereof) and the fermentation duration. However, after FDR correction for taking into account multiple testing, none of these variables significantly explained Unifrac distances.

In order to understand further the relationship between sourdough fungal community composition and bread making practices, we clustered sourdoughs according to their fungal community composition, on the basis of the PCoA of their weighted Unifrac distances. Then, we tested the link between the fungal community group and the bread making practice group (farmer/artisanal practices group) as well as the link between the fungal community group and each of the different bread making practices (Figure 5). Sourdoughs were clustered into three fungal community groups. Group 1 encompassed all sourdoughs (but two) having *Kazachstania* species as dominant species (*K. humilis, K. barnettii, K. bulderi, K. saulgeensis, K. bozae*). Group 2 contained sourdoughs with *Saccharomyces sp.*, *K. servazzi* or *K. unispora* as dominant species. Group 3 harbored sourdoughs with *S. cerevisiae* together with other species such as *Pichia kudriavzevii*, *Candida sake*, or a Dipodascaceae sp. Group 1 sourdoughs were mostly made by bakers having farmer's bread making practices while group 2 and group 3 sourdoughs were mostly made by bakers using artisanal practices (exact Fisher test, $P=0.035$). The fungal community groups were significantly associated with two specific bread making practice variables: the quantity (in kg) of bread made per week (Exact Fisher test, $P=0.001$) and the use of commercial yeast (Exact Fisher test, $P=0.05$). All sourdoughs in group 2 but one were found in bakeries making between

500 kg and 1000 kg of bread per week, while groups 1 and 3 sourdoughs originated from bakeries

producing very different amounts of bread (ranging from amounts below 250 kg to over 1000 kg). In

addition, group 1 sourdoughs were more frequently found in bakeries that do not use commercial yeast

while groups 2 and 3 were more frequently found in bakeries using the commercial yeast *S. cerevisiae*

(Exact Fisher test, P=0.01). Interestingly, group 1 sourdoughs harbored *S. cerevisiae* either at a relative

abundancy below 1% or not at all, while all groups 2 and 3 sourdoughs had *S. cerevisiae* at a relative

abundancy over 20%, except in three cases where it was either absent or at a relative abundancy below

6%.

To test more specifically the link between bread making practices and the distribution of

Kazachstania species, we analyzed more in-depth group 1 sourdoughs. Within this group, eight

sourdoughs had *K. humilis* as dominant species, six had *K. bulderi*, three had *K. bozae* and the remainder

had still other *Kazachstania* species. All sourdoughs made with artisanal practices carried *K. humilis* as

dominant species or, in one case, the *K. bozae*. By contrast, sourdoughs made with farmer practices had

as dominant species *K. bulderi*, *K. australis*, *K. barnettii*, *K. saulgeensis* or *K. bozae* (exact Fisher test,

P=0.004).

Fungal community composition was partly related to sourdough acidity, maltose

concentration and hydration

The composition of fungal community may affect sourdough metabolic content (sugars, acids, alcohols)

via fungal strains metabolite consumption and production. Inversely, the presence and concentration of

different compounds (sugars, acids, alcohols) may affect differently the fitness depending on the strains

and consequently be one of the drivers of fungal community composition. For example, lactic acid

bacteria (LAB) are the main producers of acidity in sourdough, but yeasts also produce acetic acid and

also indirectly affect acidity through positive or negative interaction with bacteria.

To investigate the relation between sourdough fungal communities and metabolic compounds, we

quantified sourdough hydration, yeast density, bacteria density, sourdough pH, total titrable acidity

(TTA), sourdough concentration in seven sugars (maltose, glucose, fructose, raffinose, arabinose,

mannose, xylose), four alcohols (glycerol, ethanol, mannitol, meso-erythtritol), six acids (lactate,

acetate, glutarate, pyruvate, malate, succinate) and calculated the fermentative quotient (lactate over

acetate ratio). For each variable, there was a wide range of variation (Table S5). The principal component

analysis based on all variables showed no evidence of sourdough grouping (Figure S4). As expected in

fermentation, yeast density was positively correlated to ethanol (r=0.74, P<0.001), glycerol (r=0.67,

P<0.001), and acetate (r=0.6, P<0.001) concentration. However, it was not significantly correlated to

sugar concentrations. This might be explained by the co-occurrence of bacteria which have their own

metabolism and interact by competition and/or cross feeding with sourdough yeasts.

We then tested whether the variation of each quantitative variable was associated with the bread

making practices groups (farmer-like practices and artisanal practices). There was no significant effect

of the bread making practice group except for sourdough hydration that was significantly higher in

sourdoughs made using farmer-like practices ($F_{1,94}$=11,69, P<0.001). On average, sourdoughs made with

farmer-like practices had 55% water while sourdoughs made with artisanal-like practices had in average

49% of water.

In addition, we tested whether variations in quantitative variables were associated with the fungal

community groups (Table S5). Group 3 microbial community sourdoughs (defined by PCoA clustering

on Unifrac distance, see below), which contains *S. cerevisiae* in co-dominance with a second yeast

species (*Candida sake*, *Pichia kudriavzevii* or a *Dipodascus* species), had a significantly higher mean

pH (mean pH$_{group3}$=4.2 against pH $_{group1}$=3.8, Tukey Contrasts, P<0.001), lower TTA (mean TTA

$_{group3}$=7.7 against TTA $_{group1}$=17.1, Tukey Contrasts, P=0.002), and a higher maltose concentration (mean

Maltose $_{group3}$=52.8 mg/gr of sourdough against Maltose $_{group1}$=24.1 mg/gr of sourdough, Tukey

Contrasts P=0.002) than group1, having a *Kazachstania* dominant species. Compared to group 2 having

in most cases *S. cerevisiae* as dominant species, it also had higher pH (pH $_{group2}$=3.9, Tukey Contrasts,

P=0.003), and higher maltose concentration (Maltose $_{group2}$=23.7, Tukey contrast, P=0.003). These data

may reflect a lower fermentative activity for group 3 fungal community having two co-dominant species,

and/or a negative interaction effect of group 3 fungal community on the activity of lactic acid bacteria

(LAB), which are the main producers of sourdough acids. Previous studies on the bacteria content of

the same sourdoughs showed that *F. sanfranciscensis* was most generally the dominant species, although

C. heilongjiangensis, L. curvatus or *Levilactobacillus brevis* were also found as dominant species [20,

52, 53]. We found no significant correlation between LAB and yeast densities (r= -0.15, p= 0.45, Figure

S4) but the link between fungal and bacterial community might be species and strains dependent.

Additional studies on the interactions between fungal and bacterial communities need to be performed

to better understand how they may drive sourdough acidity and sugar content.

We also analyzed whether the variations of each quantitative variable was associated with the

dominant yeast species. We only considered the 26 sourdoughs having either *S. cerevisiae* (9

sourdoughs), *K. humilis* (8 sourdoughs), *K. bulderi* (6 sourdoughs) or *K. bozae* (3 sourdoughs) as

dominant species, since the other yeast species were found dominant only once. The differences in

dominant species was not significantly associated to variation in sourdough sugar, acids or alcohol

concentration. However, on average, sourdoughs dominated by *K. bulderi* were more hydrated (63%

water content in average) than sourdoughs dominated by *K. humilis*, *K. bozae*, and *S. cerevisiae*, having

respectively 49%, 47%, 53 % water content in average (P<0.001 for the 3 Tukey Contrasts).

Kazachstania bulderi was found to be dominant only in sourdoughs made using farmer-like practices, a

bread making practice group that was also found to be associated with more hydrated sourdoughs.

Additional experiments should be carried out to test whether this species has indeed a better fitness in

more hydrated sourdoughs or whether its presence in more hydrated sourdoughs is related to covariation

with other farmer practices.

In conclusion, no clear evidence was found of the impact of bread making practices or of the

dominant yeast species on the metabolic composition of sourdough. On the other hand, our results

showed metabolic differences between sourdoughs having one or two co-dominant yeast species.

Phenotypic signatures of domestication

A previous analysis on *S. cerevisiae* revealed that sourdough strains had higher average fitness and fermentation performance than strains from other environments in a sourdough-mimicking medium [10]. Here, we investigated whether evidence of a domestication syndrome could also be found in *K. humilis* and *K. bulderi*, the two *Kazachstania* species most commonly found in French sourdoughs. We tested whether fitness (log of population size and mortality at the end of fermentation) and fermentation performance (CO_2max, Vmax, t1g, tVmax) differed between sourdough strains and strains from elsewhere.

A principal component analysis of 38 strains of *K. bulderi* and *K. humilis*, based on quantitative variation in the six phenotypic variables described below was carried out. The first two axis explained 80.5% of the variation and clearly separated strains by species (Figure 6). The *K. bulderi* strains were located at the right of the PCA and were characterized by high population size and low mortality at the end of fermentation, while the *K. humilis* strains were located at the left and were characterized by a rapid onset of fermentation (t1g), high maximum fermentation rate (Vmax), and a short time to reach Vmax (tVmax). Non-sourdough strains of *K. humilis* were located outside the cloud of sourdough strains while non-sourdough strains of *K. bulderi* were distributed within and outside the cloud of sourdough strains.

Statistical comparisons of sourdough and non-sourdough strains of *K. bulderi* and *K. humilis* for each phenotypic variable revealed phenotypic divergence for *K. humilis* but not for *K. bulderi*. While the *K. bulderi* sourdough strains did not ferment significantly faster than the non-sourdough strains, the *K. humilis* sourdough strains showed significantly higher Vmax, lower t1g, and lower tVmax than the non-sourdough *K. humilis* strains (Figure 7, Table S6). On average, they started fermentation two hours before the others and reached Vmax three hours before the others. In addition, their Vmax were on average 34% higher than the others.

Discussion

Sourdough microbial diversity has been intensively studied worldwide. Despite a cultural and historical

interest on bread in France, French sourdough fungal diversity was only partly characterized before this

study [19, 53, 71]. A recent large-scale (> 500 starters) study of sourdough microbial diversity revealed

the fungal diversity that can be detected over the globe across home-made sourdoughs [17]. All the

yeast species detected at a relative abundance over 1% in this international collection of sourdoughs

were detected in French sourdoughs except the species *Wickerhamomyces anomalus*, *Pichia*

membranifaciens, *Naumovozyma castellii* and *Saccharomyces bayanus*. Inversely, French baker's

sourdough harbored some yeast species that were never found elsewhere, such as *K. bozae*, *K. australis*,

K. saulgeensis [65].

Beyond the genus of the baker yeast species, *S. cerevisiae*, the most represented yeast genus in French

sourdough was *Kazachstania*. Eight *Kazachstania* species were found as dominant yeast species in at

least one French sourdough: *K. humilis*, *K. bulderi*, *K. barnettii*, *K. unispora*, *K. servazzii*, *K. bozae*, *K.*

australis and *K. saulgeensis*. Three *Kazachstania* species (*K. exigua, K. lodderae, K. naganishi*) already

reported in sourdough were not found in our collection of French sourdough. *Kazachstania lodderae*

and *K. naganishi* are rarely found in sourdough. By contrast, *K. exigua* is a frequently cited sourdough

species in the literature. This species has been previously found in France, Finland [77], Italy [78, 79],

Denmark [80], Ethiopia [81], USA [17] and is the first species to have been isolated from a sourdough

(in San Francisco, [29]). However, its taxonomic characterization may have been hampered by the fact

that it probably originated from hybridization between unknown yeast species [82]. To date, the genus

Kazachstania is composed of more than 40 species, of which 11 are present in sourdough. It is possible

that an adaptive radiation linked to the adaptation to different sourdoughs or to different anthropized

niches has taken place, as it has been observed for example in cichlids during their adaptation to different

lakes or in Penicilium domesticated fungi. Indeed, five of the *Kazachstania* species present in the

sourdough have so far only been detected in human-related niches. These are *K. saulgeensis*, *K.*

barnettii, *K. bozae*, *K. bulderi* and *K. humilis*. These species are genetically closer to each other than

they are to *K. australis*, *K. servazzii* and *K. unispora* which have also been found in nature and are

grouped in another part of the *Kazachstania* pylogenetic tree. Genomic analysis of these species would

shed light on their evolution and the genetic changes that would have been selected during potential

domestication. So far, eight of the 11 *Kazachstania* species found in the sourdough have at least one

genome assembly available in public databases, including *K. saulgeensis* and *K. barnettii* [44, 45], and

the assemblies of *K. bulderi* and *K. humilis*, which were recently published in public databases (Bio

project: PRJEB44438). The genomic and phenomic analysis of the large collection of *Kazachstania*

strains obtained by our study, together with the world collection of *Kazachstania* strains, may shed light

on the radiation and domestication processes of these species.

We found that yeast community composition was partly related with bread making practices. Bread

making practice divergence also led to different phenotypic signatures. Strains of the *K. humilis* species,

which was typical of sourdough made by artisanal-like practices, had higher fermentation rate while

strains of *K. bulderi* which was typical of sourdough made by farmer-like bread making practices had

not. The species *K. humilis* has been found in many countries, viz. Austria, Canada, China, Denmark,

Ethiopia, Finland, Germany, Greece, Italy, Morocco, the Netherland, Spain, UK, USA, and France [1,

15–17, 19, 53, 55–60]. It is also the most frequently encountered *Kazachstania* species in sourdoughs

around the world. This species is therefore frequently found in bakeries, where short fermentations are

often favored. This may explain why sourdough strains of *K. humilis* seem to have been selected for

increased fermentation rate. Increased fermentation rate was also found in bakery strains of *S. cerevisiae*

when compared to non-bakery strains.

In contrast, we did not find evidence for improved fermentation performance in *K. bulderi*, which

was the third most represented species in French sourdough. *Kazachstania bulderi* was found in bakeries

with farmer-like practices. These bakeries often bake bread once or twice a week and store their

sourdough for several days. They also often use long fermentation and thus may not have selected an

increased fermentation rate. Farmer-bakers typically store their sourdough for several days and therefore

make a lower number of backslopping. In addition, they make bread with longer fermentation times than

artisanal bakers. It is therefore possible that they did not select to accelerate the speed of fermentation

and instead let natural selection in the sourdough environment act alone. Alternatively, the lack of

phenotypic divergence between sourdough and non-sourdough strains may reflect the limitation of our

sampling. *Kazachstania bulderi* has been reported for the first time in anaerobic maize silage in the

Netherlands and in fermented liquid feed for piglets [61, 62], more recently in French, Belgium and

Spain sourdoughs [19, 53, 57, 63] but to our knowledge was never found in wild environment. Here, we

compare sourdough strains with strains coming from silage and animal feed. It is unknown whether

silage and animal feed strains are wild strains or feral strains that have escaped from other domesticated

environments. This may explain why we did not detect any phenotypic divergence between sourdough

and non-sourdough strains of *K. bulderi*. Other than fermentation phenotypes, there was no evidence of

fitness differences between sourdough and non-sourdough strains of *K. humilis* and *K. bulderi* in

sourdough mimicking media. Additional experiments in real dough should be performed to further test

the effect of natural selection in this environment.

Other evolutionary process than selection could also explain the distribution of yeast species across

sourdoughs. Interviews with the bakers working with sourdough hosting *K. bulderi* and *K. bozae*

suggested the role of dispersion of these species in French sourdoughs. Indeed, these bakers have been

connected over the years either through seed exchanges, sourdough mixing or gifts, bread making

training in common or working in one another's bakery. Some yeast species have been found in the

bakery house environment and baker's hands and may thus be dispersed through baker's tools or baker's

travels [58, 67, 75, 76]. However, it is still unclear whether wheat seeds and flour are a source of

sourdough yeasts.

To our knowledge, this is the first evidence of the influence of artisanal practices on taxonomic

diversity in microbial communities. On the other hand, several studies have shown that making

fermented products could lead to the selection of divergent phenotypes and genotypes. This is the case

of sourdough and industrial populations of the baker's yeast *S. cerevisiae* that diverge from each other

and have a better fermentation performance than non-baker's strains. As for beer populations, sourdough

populations have acquired a better capacity to assimilate maltose, linked at least in part to an increase

in the number of copies of the genes involved in the assimilation of maltose. Several studies on wine

populations of *S. cerevisiae*, *Torulaspora delbrueckii* and *Lachancea thermotolerans* also showed that

wine populations are genetically differentiated from strains from other environments and present

beneficial phenotypes in grape must and for wine quality. The analysis of filamentous cheese fungi *P.*

roqueforti and *P. camemberti* populations also revealed cheese making genetically differentiated

populations. Interestingly, different genetic groups associated with different cheese making practices

were found. Strains of the blue cheese fungus, *P. roqueforti*, isolated from Roquefort cheese were more

diverse and were genetically and phenotypically different than strains used to make other blue cheeses

[83, 84, 85]. Two varieties of the white cheese making fungus, *P. camemberti*, with different phenotypic

features, were associated with different kind of cheese (camembert and brie). All together these studies

show that the diversity of practices used to make fermented products allows to maintain genetic,

phenotypic and taxonomic diversity.

However, fungal domestication also involved strong bottlenecks. For example, the low level of genetic

diversity found in blue-cheese *P. roqueforti* strains and soft-cheese *P. camemberti* strains revealed the

risk of diversity erosion in fermented product making [6, 7, 84]. This risk is accentuated by fertility

depression among fungal domesticated strains [86]. This risk is also associated with the massive use of

few industrial strains or the need to standardize products to meet the specifications of industrial

production or protected designation of origin (PDO) [6]. Here, we show that despite the recurrent use of

S. cerevisiae as industrial starter species in bakeries and homes, and the occurrence of this species in a

wide range of habitats such as soil, trees, and humans [9, 54], this species does not appear to have

overwhelmingly colonized French traditional sourdoughs (Figure 3, Figure 4). This result confirmed a

recent analysis which revealed that *S. cerevisiae* sourdough strains had a different evolutionary history

from industrial strains [10]. The dynamic of microbial species colonization and invasion in food

environment remain largely unknown. A recent study suggested that domesticated *Penicillium* strains

can evolved phenotypically in a few weeks [87]. Additional experiments at the level of microbial

community will shed light on the dynamic of microbial community establishment in food production

and on the ability of industrial strains to invade food microbial community.

Conclusion and perspectives

In conclusion, a great diversity of bread making practices and fungal community composition was found in our sample of French sourdoughs. Surprisingly, the well-known baker's yeast *Saccharomyces cerevisiae* was found dominant only in one fourth of the sampled sourdoughs. By contrast, several species of the neighboring genus *Kazachstania* were detected at high frequency, revealing a major role for this mostly unknown genus in the study of fungal domestication and in bread making. Therefore, our results highlight the necessity of maintaining socio-cultural diversity to maintain microbial diversity in food systems. These findings could not have been evidenced without the collaboration of bakers and scientists, showing the importance of participatory research projects to gain new insight into biodiversity preservation.

Acknowledgements

This work has emanated from research conducted with the financial support of the French National Research Agency, ANR-13-ALID-0005 and INRAE. We thank all the bakers and farmer-bakers for constructing the project with us and for sharing their sourdoughs and knowledge. We thank Yoann Robert, Candice Aulard, Josette Bessière for lab assistance and Matthieu Barret for MiSeq sequencing. We also would like to thank Philippe Roussel for sharing his knowledge about French bread.

References

1.   Carbonetto B, Ramsayer J, Nidelet T, Legrand J, Sicard D. Bakery yeasts, a new model for studies in ecology and evolution. *Yeast* ; **0**.

2.   Gibbons JG, Rinker DC. The genomics of microbial domestication in the fermented food environment. *Current Opinion in Genetics & Development* 2015; **35**: 1–8.

3.   Steensels J, Gallone B, Voordeckers K, Verstrepen KJ. Domestication of Industrial Microbes.

Current Biology 2019; **29**: R381–R393.

4.   Gibbons JG, Salichos L, Slot JC, Rinker DC, McGary KL, King JG, et al. The evolutionary

imprint of domestication on genome variation and function of the filamentous fungus Aspergillus

oryzae. *Curr Biol* 2012; **22**: 1403–1409.

5.   Watarai N, Yamamoto N, Sawada K, Yamada T. Evolution of Aspergillus oryzae before and

after domestication inferred by large-scale comparative genomic analysis. *DNA Research* 2019; **26**:

465–472.

6.   Ropars J, Didiot E, Rodríguez de la Vega RC, Bennetot B, Coton M, Poirier E, et al.

Domestication of the Emblematic White Cheese-Making Fungus Penicillium camemberti and Its

Diversification into Two Varieties. *Curr Biol* 2020; **30**: 4441-4453.e4.

7.   Ropars J, López-Villavicencio M, Snirc A, Lacoste S, Giraud T. Blue cheese-making has

shaped the population genetic structure of the mould Penicillium roqueforti. *PLOS ONE* 2017; **12**:

e0171387.

8.   Legras J-L, Galeote V, Bigey F, Camarasa C, Marsit S, Nidelet T, et al. Adaptation of *S.*

cerevisiae to Fermented Food Environments Reveals Remarkable Genome Plasticity and the

Footprints of Domestication. *Mol Biol Evol* 2018; **35**: 1712–1727.

9.   Peter J, De Chiara M, Friedrich A, Yue J-X, Pflieger D, Bergström A, et al. Genome evolution

across 1,011 *Saccharomyces* cerevisiae isolates. *Nature* 2018; **556**: 339–344.

10.   Bigey F, Segond D, Friedrich A, Guezenec S, Bourgais A, Huyghe L, et al. Evidence for Two

Main Domestication Trajectories in *Saccharomyces* cerevisiae Linked to Distinct Bread making

Processes. *Current Biology* 2020; S0960982220316912.

11.   Arranz-Otaegui A, Carretero LG, Ramsey MN, Fuller DQ, Richter T. Archaeobotanical

evidence reveals the origins of bread 14,400 years ago in northeastern Jordan. *PNAS* 2018; **115**: 7925–

7930.

12.   Samuel D. Bread in archaeology. *Civilisations Revue internationale d'anthropologie et de*

sciences humaines 2002; 27–36.

13.   Roussel P, Chiron H. Les pains français: évolution, qualité, production. 2005. Maé-Erti.

14.   Decock P, Cappelle S. Bread technology and sourdough technology. *Trends in Food Science*

& Technology 2005; **16**: 113–120.

15.   Van Kerrebroeck S, Maes D, De Vuyst L. Sourdoughs as a function of their species diversity

and process conditions, a meta-analysis. *Trends in Food Science & Technology* 2017; **68**: 152–159.

16.   Arora K, Ameur H, Polo A, Di Cagno R, Rizzello CG, Gobbetti M. Thirty years of knowledge

on sourdough fermentation: A systematic review. *Trends in Food Science & Technology* 2021; **108**:

71–83.

17.   Landis EA, Oliverio AM, McKenney EA, Nichols LM, Kfoury N, Biango-Daniels M, et al.

The diversity and function of sourdough starter microbiomes. *eLife* 2021; **10**: e61644.

18.   Demeulenaere É, Rivière P, Hyacinthe A, Baltassat R, Baltazar S, Gascuel J-S, et al. La

sélection participative à l'épreuve du changement d'échelle. À propos d'une collaboration entre

paysans sélectionneurs et généticiens de terrain. *Natures Sciences Societes* 2017; **Vol. 25**: 336–346.

19.   Urien C, Legrand J, Montalent P, Casaregola S, Sicard D. Fungal Species Diversity in French

Bread Sourdoughs Made of Organic Wheat Flour. *Front Microbiol* 2019; **10**.

20.   Michel E, Monfort C, Deffrasnes M, Guezenec S, Lhomme E, Barret M, et al.

Characterization of relative abundance of lactic acid bacteria species in French organic sourdough by

cultural, qPCR and MiSeq high-throughput sequencing methods. *International Journal of Food*

Microbiology 2016; **239**: 35–43.

21.   White TJ, Bruns T, Lee S, Taylor J. 38 - AMPLIFICATION AND DIRECT SEQUENCING

OF FUNGAL RIBOSOMAL RNA GENES FOR PHYLOGENETICS. In: Innis MA, Gelfand DH,

Sninsky JJ, White TJ (eds). *PCR Protocols*. 1990. Academic Press, San Diego, pp 315–322.

22.   Martin KJ, Rygiewicz PT. Fungal-specific PCR primers developed for analysis of the ITS

region of environmental DNA extracts. *BMC Microbiology* 2005; **5**: 28.

23.   Peris D, Sylvester K, Libkind D, Gonçalves P, Sampaio J, Alexander WG, et al. Population

structure and reticulate evolution of *Saccharomyces* eubayanus and its lager-brewing hybrids. *Mol*

Ecol 2014; **23**: 2031–2045.

24.   Nguyen H-V, Gaillardin C. Evolutionary relationships between the former species

Saccharomyces uvarum and the hybrids *Saccharomyces* bayanus and *Saccharomyces* pastorianus;

reinstatement of *Saccharomyces* uvarum (Beijerinck) as a distinct species. *FEMS Yeast Res* 2005; **5**:

471–483.

25.   Libkind D, Hittinger C, of the … V-E. Microbe domestication and the identification of the

wild genetic stock of lager-brewing yeast. *Proceedings of the …* 2011.

26.   Sampaio JP, Gonçalves P. Natural populations of *Saccharomyces* kudriavzevii in Portugal are

associated with oak bark and are sympatric with *S. cerevisiae* and S. paradoxus. *Appl Environ*

Microbiol 2008; **74**: 2144–2152.

27.   Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res*

2016; **44**: D7–D19.

28.   Weiss S, Samson F, Navarro D, Casaregola S. YeastIP: a database for identification and

phylogeny of Saccharomycotina yeasts. *FEMS Yeast Research* 2013; **13**: 117–125.

29.   Escudié F, Auer L, Bernard M, Mariadassou M, Cauquil L, Vidal K, et al. FROGS: Find,

Rapidly, OTUs with Galaxy Solution. *Bioinformatics* 2018; **34**: 1287–1294.

30.   Magoč T, Salzberg SL. FLASH: fast length adjustment of short reads to improve genome

assemblies. *Bioinformatics* 2011; **27**: 2957–2963.

31.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 2011; **17**: 10–12.

32.   Sickle 1.33 – Windowed Adaptive Trimming for Fastq files using Quality – My Biosoftware –

Bioinformatics Softwares Blog.

33.   Mahé F, Rognes T, Quince C, Vargas C de, Dunthorn M. Swarm: robust and fast clustering

method for amplicon-based studies. *PeerJ* 2014; **2**: e593.

34.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source tool

for metagenomics. *PeerJ* 2016; **4**: e2584.

35.   Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Bahram M, et al. Towards a

unified paradigm for sequence-based identification of fungi. *Molecular Ecology* 2013; **22**: 5271–5277.

36.   Bloem A, Rollero S, Seguinot P, Crépin L, Perez M, Picou C, et al. Workflow Based on the

Combination of Isotopic Tracer Experiments to Investigate Microbial Metabolism of Multiple Nutrient

Sources. *J Vis Exp* 2018.

37.   Delobel P, Pradal M, Blondin B, Tesniere C. A 'fragile cell' sub-population revealed during

cytometric assessment of *Saccharomyces* cerevisiae viability in lipid-limited alcoholic fermentation.

Lett Appl Microbiol 2012; **55**: 338–344.

38.   Lê S, Josse J, Husson F. FactoMineR: An R Package for Multivariate Analysis. *Journal of*

Statistical Software 2008; **25**: 1–18.

39.   McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and

Graphics of Microbiome Census Data. *PLOS ONE* 2013; **8**: e61217.

40.   Lozupone C, Knight R. UniFrac: a new phylogenetic method for comparing microbial

communities. *Appl Environ Microbiol* 2005; **71**: 8228–8235.

41.   Gouy M, Guindon S, Gascuel O. SeaView version 4: A multiplatform graphical user interface

for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 2010; **27**: 221–224.

42.   Chao A, Chiu C-H, Jost L. Phylogenetic diversity measures based on Hill numbers. *Philos*

Trans R Soc Lond B Biol Sci 2010; **365**: 3599–3609.

43.   Jost L. Partitioning diversity into independent alpha and beta components. *Ecology* 2007; **88**:

2427–2439.

44.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*

1995; **57**: 289–300.

45.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed

Effects Models. *Journal of Statistical Software* 2017; **82**: 1–26.

46.   Hijmans RJ, Karney (GeographicLib) C, Williams E, Vennes C. geosphere: Spherical

Trigonometry. 2021.

47.   Dray S, Dufour A-B. The ade4 Package: Implementing the Duality Diagram for Ecologists.

Journal of Statistical Software 2007; **22**: 1–20.

48.   Pedersen HW Danielle Navarro, and Thomas Lin. Welcome | ggplot2.

49.   Cheng J, Karambelkar B, Xie Y. Leaflet: Create interactive web maps with the

javascript'leaflet'library. *R package version* 2018; **2**.

50.   Clément H, Prost C, Chiron H, Ducasse MB, Della Valle G, Courcoux P, et al. The effect of

organic wheat flour by-products on sourdough performances assessed by a multi-criteria approach.

Food Res Int 2018; **106**: 974–981.

51.   De Vuyst L, Van Kerrebroeck S, Leroy F. Microbial Ecology and Process Technology of

Sourdough Fermentation. *Adv Appl Microbiol* 2017; **100**: 49–160.

52.   Lhomme E, Orain S, Courcoux P, Onno B, Dousset X. The predominance of Lactobacillus

sanfranciscensis in French organic sourdoughs and its impact on related bread characteristics.

International Journal of Food Microbiology 2015; **213**: 40–48.

53.   Lhomme E, Urien C, Legrand J, Dousset X, Onno B, Sicard D. Sourdough microbial

community dynamics: An analysis during French organic bread making processes. *Food Microbiol*

2016; **53**: 41–50.

54.   Goddard MR, Greig D. *Saccharomyces* cerevisiae: a nomadic yeast with no niche? *FEMS*

Yeast Res 2015; **15**: fov009.

55.   Valmorri S, Tofalo R, Settanni L, Corsetti A, Suzzi G. Yeast microbiota associated with

spontaneous sourdough fermentations in the production of traditional wheat sourdough breads of the

Abruzzo region (Italy). *Antonie Van Leeuwenhoek* 2010; **97**: 119–129.

56.   Wang X, Du H, Zhang Y, Xu Y. Environmental Microbiota Drives Microbial Succession and

Metabolic Profiles during Chinese Liquor Fermentation. *Appl Environ Microbiol* 2018; **84**: e02369-

17.

57.   Chiva R, Celador-Lera L, Uña JA, Jiménez-López A, Espinosa-Alcantud M, Mateos-

Horganero E, et al. Yeast Biodiversity in Fermented Doughs and Raw Cereal Matrices and the Study

of Technological Traits of Selected Strains Isolated in Spain. *Microorganisms* 2020; **9**: 47.

58.   Reese AT, Madden AA, Joossens M, Lacaze G, Dunn RR. Influences of Ingredients and

Bakers on the Bacteria and Fungi in Sourdough Starters and Bread. *mSphere* 2020; **5**.

59.   Syrokou MK, Themeli C, Paramithiotis S, Mataragas M, Bosnea L, Argyri AA, et al.

Microbial Ecology of Greek Wheat Sourdoughs, Identified by a Culture-Dependent and a Culture-Independent Approach. *Foods* 2020; **9**: 1603.

60.   Sugihara TF, Kline L, Miller MW. Microorganisms of the San Francisco sour dough bread process. I. Yeasts responsible for the leavening action. *Appl Microbiol* 1971; **21**: 456–458.

61.   Middelhoven WJ, Kurtzman CP, Vaughan-Martini A. *Saccharomyces* bulderi sp. nov., a yeast that ferments gluconolactone. *Antonie Van Leeuwenhoek* 2000; **77**: 223–228.

62.   Gori K, Bjørklund MK, Canibe N, Pedersen AØ, Jespersen L. Occurrence and identification of yeast species in fermented liquid feed for piglets. *Microb Ecol* 2011; **61**: 146–153.

63.   Comasio A, Verce M, Van Kerrebroeck S, De Vuyst L. Diverse Microbial Composition of Sourdoughs From Different Origins. *Front Microbiol* 2020; **11**: 1212.

64.   Gouliamova D, Dimitrov R. *KAZACHSTANIA* CHRYSOLINAE AND *KAZACHSTANIA* BOZAE TWO NEW YEAST SPECIES OF THE GENUS *KAZACHSTANIA*. TRANSFER OF FOUR *KAZACHSTANIA* SPECIES TO GRIGOROVIA GEN. NOV. AS NEW COMBINATIONS. 2020.

65.   Jacques N, Sarilar V, Urien C, Lopes MR, Morais CG, Uetanabaro APT, et al. Three novel ascomycetous yeast species of the *Kazachstania* clade, *Kazachstania* saulgeensis sp nov., *Kazachstania* serrabonitensis sp nov and *Kazachstania* australis sp nov Reassignment of Candida humilis to *Kazachstania* humilis f.a. comb. nov and Candida pseudohumilis to *Kazachstania* pseudohumilis f.a. comb. nov. *International Journal of Systematic and Evolutionary Microbiology* 2016; **66**: 5192--5200.

66.   Sarilar V, Sterck L, Matsumoto S, Jacques N, Neuvéglise C, Tinsley CR, et al. Genome sequence of the type strain CLIB 1764T (=CBS 14374T) of the yeast species *Kazachstania* saulgeensis isolated from French organic sourdough. *Genomics Data* 2017; **13**: 41–43.

67.   Minervini F, Lattanzi A, De Angelis M, Di Cagno R, Gobbetti M. Influence of Artisan Bakery- or Laboratory-Propagated Sourdoughs on the Diversity of Lactic Acid Bacterium and Yeast Microbiotas. *Appl Environ Microbiol* 2012; **78**: 5328–5340.

68.   Salovaara H, Savolainen J. Yeast type isolated from Finnish sour rye dough starters. *Acta Alimentaria Polonica* 1984; **10**.

69.   Decimo M, Quattrini M, Ricci G, Fortina MG, Brasca M, Silvetti T, et al. Evaluation of microbial consortia and chemical changes in spontaneous maize bran fermentation. *AMB Express* 2017; **7**: 205.

70.   Fraberger V, Unger C, Kummer C, Domig KJ. Insights into microbial diversity of traditional Austrian sourdough. *LWT* 2020; **127**: 109358.

71.   Lhomme E, Lattanzi A, Dousset X, Minervini F, De Angelis M, Lacaze G, et al. Lactic acid bacterium and yeast microbiotas of sixteen French traditional sourdoughs. *Int J Food Microbiol* 2015; **215**: 161–170.

72.   Jacques N, Sarilar V, Urien C, Lopes MR, Morais CG, Uetanabaro APT, et al. Three novel ascomycetous yeast species of the *Kazachstania* clade, *Kazachstania* saulgeensis sp. nov., *Kazachstania*serrabonitensis sp. nov. and *Kazachstania* australis sp. nov. Reassignment of Candida humilis to *Kazachstania* humilis fa comb. nov. and Candida pseudohumilis to *Kazachstania* pseudohumilis fa comb. nov. *International journal of systematic and evolutionary microbiology* 2016; **66**: 5192–5200.

73.   Kurtzman CP. Phylogenetic circumscription of *Saccharomyces*, Kluyveromyces and other members of the Saccharomycetaceae, and the proposal of the new genera Lachancea, Nakaseomyces, Naumovia, Vanderwaltozyma and Zygotorulaspora. *FEMS yeast research* 2003; **4**: 233–245.

74.   Boynton PJ, Greig D. The ecology and evolution of non-domesticated *Saccharomyces* species. *Yeast* 2014; **31**: 449–462.

75.   Minervini F, Lattanzi A, De Angelis M, Celano G, Gobbetti M. House microbiotas as sources of lactic acid bacteria and yeasts in traditional Italian sourdoughs. *Food Microbiology* 2015; **52**: 66–76

76.   Minervini F, Lattanzi A, Dinardo FR, De Angelis M, Gobbetti M. Wheat endophytic lactobacilli drive the microbial and biochemical features of sourdoughs. *Food Microbiol* 2018; **70**: 162–171.

77.   Infantes M, Schmidt JL. Characterisation of the yeast flora of natural sourdoughs located in various French areas. Sciences des Aliments (France) [Internet]. 1992, available on: https://agris.fao.org/agris-

search/search.do?recordID=FR9204966search/search.do?recordID=FR9204966

78. Pulvirenti A, Solieri L, Gullo M, Vero LD, Giudici P. Occurrence and dominance of yeast

species in sourdough. *Letters in Applied Microbiology* 2004; **38**: 113-7.

79. Minervini F, Lattanzi A, De Angelis M, Di Cagno R, Gobbetti M. Influence of Artisan Bakery-

or Laboratory-Propagated Sourdoughs on the Diversity of Lactic Acid Bacterium and Yeast

Microbiotas. *Appl Environ Microbiol* 2012; **78**: 5328-40.

80. Nuobariene L, Hansen ÅS, Arneborg N. Isolation and identification of phytase-active yeasts

from sourdoughs. LWT - *Food Science and Technology* 2012; **48**:190-6.

81. Desiye A, Abegaz K. Isolation, characterization and identification of lactic acid bacteria and

yeast involved in fermentation of Teff (EragrostisTef) Batter. 2013; /paper/Isolation%2C-

characterization-and-identification-of-Desiye Abegaz/3373ba786cbed5f07d2c6330a5362ea56198adc2

82. Bon E, Neuvéglise C, Lépingle A, Wincker P, Artiguenave F, Gaillardin C, et al. Genomic

Exploration of the Hemiascomycetous Yeasts: 6. *Saccharomyces* exiguus. *FEBS Letters* 2000; **487**:

42-6.

83. Cheeseman K, Ropars J, Renault P, Dupont J, Gouzy J, Branca A, et al. Multiple recent

horizontal transfers of a large genomic region in cheese making fungi. *Nat Commun* 2014; **5**: 2876.

84. Dumas E, Feurtey A, Rodríguez de la Vega RC, Le Prieur S, Snirc A, Coton M, et al.

Independent domestication events in the blue-cheese fungus Penicillium roqueforti. *Molecular Ecology*

2020; **29**: 2639-60.

85. Ropars J, Rodríguez de la Vega RC, López-Villavicencio M, Gouzy J, Sallet E, Dumas É, et al.

Adaptive Horizontal Gene Transfers between Multiple Cheese-Associated Fungi. *Current Biology* 2015;

25: 2562-9.

86. Ropars J, Lo Y, Dumas E, Snirc A, Begerow D, Rollnik T, et al. Fertility depression among

cheese-making Penicillium roqueforti strains suggests degeneration during domestication. *Evolution*

2016; **70**: 2099–2109.

87.   Bodinaku I, Shaffer J, Connors AB, Steenwyk JL, Biango-Daniels MN, Kastman EK, et al.

Rapid Phenotypic and Metabolomic Domestication of Wild Penicillium Molds on Cheese. *mBio* 2019;

10: e02445-19.

 Figure Legends

Figure 1. The sourdough bread making process. Sourdough is a mix of flour and water naturally

fermented by bacteria and yeasts. It is initiated by mixing flour, water and occasionally other ingredients.

It is then "fed" by regularly adding flour and water, a process termed back-slopping. Once considered

mature by bakers based on their acidity, flavour and bubbling activity, the sourdough is called "chief",

or "mother" sourdough, and can then be used for bread making. The bread making process starts from

this "chief sourdough", or from a piece of dough or sourdough sampled from the preceding bread making

process, or initiated from a mix of flour and water naturally colonized by yeasts and LAB following

several back-sloppings. Once or several times, the chief sourdough is refreshed by adding flour and

water to constitute the final sourdough, which is used for bread making. This final sourdough is mixed

with flour, water, and other ingredients (salt, seeds, yeasts starters, etc.) during kneading to constitute

the dough. After kneading, primary fermentation occurs during the first rising. The dough is then divided

and shaped. The pieces of dough are then left to rise during a second fermentation and finally oven-

baked.

Figure 2. Multiple Correspondence Analysis (MCA) based on 28 categorical variables describing

bread making practices.

A) Representation of bakers. Each point represents a bakery. The purple area on the left brings together

baker with "artisanal" practices and the light blue area on the right the bakers with "farmer" practices.

The dot's colors indicate the PCoA cluster of the sourdough fungal community (see Figure 5). Black

dots for group 1, empty dot for group 2, grey for group 3. The fungal community of the sourdough of

baker 14 was not studied. B) Representation of the 20 first categories that contributed the most to the

MCA axes. The category, which corresponds to a class of a variable, is written next to the triangle. C)

Distribution of each variable for each bread making practices group. Only variables that mostly

explained differences between bread making practice groups are shown: use of commercial yeast,

kneading method, chief origin, kg of bread production per week, number of bread making per week,

percent of water in dough, number of back-sloppings before making bread, water origin, sourdough age

and flour percentage in dough. The categories of each of these variables are indicated on the right.

Figure 3. Yeast species diversity was analyzed for 38 out of the 39 sourdoughs with both cultural and

metabarcoding methods. Left: species were identified by traditional microbial isolation and

identification using ITS sequencing. Right: species were identified using ITS1 metabarcoding. The

three most frequently-encountered species are shown in contrasting colors surrounded by black (blue:

Saccharomyces cerevisiae, red: *Kazachstania bulderi*, yellow: *Kazachstania humilis*). The bread

making practice of the baker who supplied the sourdough is indicated on the left ("artisanal" in purple,

"farmer" in light blue).

Figure 4. Distribution of yeast species diversity across French sourdoughs. Each bar represents the yeast

species diversity of one sourdough and is placed on the map where the baker is located. Sourdoughs

from "farmer" practice are surrounded in light blue and sourdoughs from "artisanal" practice are

surrounded by purple.

Figure 5. Representation of sourdough depending on the weighted Unifrac distances between

microbial communities. A) shows the clustering of sourdoughs according to their Unifrac distances on

a tree (left) or on a PCoA (right). Sourdough fungal community can be clustered in three groups

according to their weighted Unifrac distances. B to E) shows on the left: a tree constructed from the

Unifrac distance matrix, in the center: sourdoughs represented on the first 2 axes of the PCoA and on

the right: the distribution of modalities of a variable for each group of sourdoughs (number of each

group on the x-axis). B) shows the distribution of bread making practices among fungal community

groups, C) shows the distribution of the dominant or most frequent species among fungal community

(K: *Kazachstania*, C: *Candida*, T: *Torulaspora*, P: *Pichia*), D) shows the use of commercial

Saccharomyces cerevisiae starter and E) shows the quantity of bread per week.

Figure 6. Principal component analysis of 37 *K. humilis* and *K. bulderi* strains based on the

quantitative variation of maximum CO2 production (CO2max), fermentation latency phase (t1g),

maximum CO2 production rate (Vmax), time to reach the maximum production rate (tVmax), log of

population size and mortality at the end of the fermentation. The correlations between variables are

presented on the left while the figure on the right shows the projection of strains on the first two axes

representing 70.64% of the variation. The strains are colored according to their habitat of origin

(sourdough/non-sourdough). Their species is indicated by symbol.

Figure 7. Ratio between the sourdough strains mean and the non-sourdough strains mean values of

each quantitative variable measuring fermentation performance: maximum CO2 production

(CO2max), fermentation latency phase (t1g), maximum CO2 production rate (Vmax), time to reach

the maximum production rate (tVmax), log of population size. Confidence intervals are indicated by

bars.

Figure 1

Figure 2

924 Figure 4
925

926
927

Figure 5

932 Figure 6
933

935   Figure 7