Maltotriose consumption by hybrid *Saccharomyces pastorianus* is heterotic and results from regulatory cross-talk between parental sub-genomes

Heterotic origin of maltotriose consumption in Saccharomyces pastorianus

Nick Brouwers[a] , Anja Brickwedde[a], Arthur R. Gorter de Vries[a], Marcel van den Broek[a], Susan M. Weening[a], Lieke van den Eijnden[a], Jasper A. Diderich[a], Feng-Yan Bai[b], Jack T. Pronk[a] and Jean-Marc G. Daran[a]*

[a]Department of Biotechnology, Delft University of Technology, Van der Maasweg 9, 2629 HZ Delft, The Netherlands.
[b]State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing, 100101, China.
* Corresponding author: Jean-Marc G. Daran, j.g.daran@tudelft.nl, Department of Biotechnology, Delft University of Technology, van der Maasweg 9, 2629 HZ Delft, The Netherlands

Nick Brouwers:	N.Brouwers-1@tudelft.nl	
Anja Brickwedde:	a.brickwedde@web.de	
Arthur R. Gorter de Vries:	A.R.GorterdeVries@tudelft.nl	0000-0002-0841-6583
Marcel van den Broek:	Marcel.vandenBroek@tudelft.nl	
Susan M. Weening:	S.M.Weening@tudelft.nl	
Lieke van den Eijnden:	A.F.vandenEijnden@student.tudelft.nl	
Jasper A. Diderich:	J.A.Diderich@tudelft.nl	
Feng-Yan Bai:	baify@im.ac.cn	
Jack T. Pronk:	J.T.Pronk@tudelft.nl	0000-0002-5617-4611
Jean-Marc G. Daran:	J.G.Daran@tudelft.nl	0000-0003-3136-8193

Keywords: *S. eubayanus*, α-oligo-glucoside metabolism , heterosis, domestication, Nanopore sequencing, brewing, hybridization.

Abstract

S. pastorianus strains are hybrids of *S. cerevisiae* and *S. eubayanus* that have been domesticated for several centuries in lager-beer brewing environments. As sequences and structures of *S. pastorianus* genomes are being resolved, molecular mechanisms and evolutionary origin of several industrially relevant phenotypes remain unknown. This study investigates how maltotriose metabolism, a key feature in brewing, may have arisen in early *S. eubayanus* x *S. cerevisiae* hybrids. To address this question, we generated a near-complete genome assembly of Himalayan *S. eubayanus* strains of the Holarctic subclade. This group of strains have been proposed to be the origin of the *S. eubayanus* subgenome of current *S. pastorianus* strains. The Himalayan *S. eubayanus* genomes harbored several copies of a *SeAGT1* α-oligoglucoside transporter gene with high sequence identity to genes encountered in *S. pastorianus*. Although Himalayan *S. eubayanus* strains are unable to grown on maltose and maltotriose, their maltose-hydrolase and *SeMALT1* and *SeAGT1* maltose-transporter genes complemented the corresponding null mutants of *S. cerevisiae*. Expression, in a Himalayan *S. eubayanus* strain, of a functional *S. cerevisiae* maltose-metabolism regulator gene (*MALx3*) enabled growth on oligoglucosides. The hypothesis that the maltotriose-positive phenotype in *S. pastorianus* is a result of heterosis was experimentally tested by constructing a *S. cerevisiae* x *S. eubayanus* laboratory hybrid with a complement of maltose-metabolism genes that resembles that of current *S. pastorianus* strains. The ability of this hybrid to consume maltotriose in brewer's wort demonstrated regulatory cross talk between sub-genomes and thereby validated this hypothesis. These results provide experimental evidence of the evolutionary origin of an essential phenotype of lager-brewing strains and valuable knowledge for industrial exploitation of laboratory-made *S. pastorianus*-like hybrids.

Introduction

Saccharomyces pastorianus is an interspecific hybrid of S. cerevisiae and S. eubayanus (Libkind et al. 2011; Nakao et al. 2009; Walther et al. 2014; Van den Broek et al. 2015). S. pastorianus strains are widely used for production of lager beer, which is currently the most produced alcoholic beverage worldwide. Lager brewing requires alcoholic fermentation at relatively low temperatures. S. pastorianus was hypothesized to have emerged by spontaneous hybridization and to have colonized early lager-brewing processes due to a combination of cold-tolerance inherited from S. eubayanus and superior fermentation kinetics inherited from S. cerevisiae (Krogerus et al. 2015a; Hebly et al. 2015; Krogerus et al. 2017a). Lager beer is brewed from barley wort, whose sugar composition consists, by weight, of approximately 15% glucose, 60% maltose and 25% maltotriose (Zastrow et al. 2001). During wort fermentation, maltotriose is generally only utilized after glucose and maltose are depleted, while its consumption is also relatively slow and often incomplete (Alves et al. 2008; Vidgren et al. 2009; Brickwedde et al. 2017).

Complete sugar utilization is desirable for lager beer fermentation to optimize concentrations of ethanol and flavor compounds and to avoid residual sweetness (Zheng et al. 1994). While S. pastorianus and S. cerevisiae strains are capable of consuming maltotriose, none of the wild isolates of S. eubayanus characterized thus far have been shown to possess this trait (Hebly et al. 2015; Brickwedde et al. 2018; Baker and Hittinger 2018). These observations led to the hypothesis that the ability of S. pastorianus to ferment maltotriose was inherited from S. cerevisiae (Brickwedde et al. 2018; Hebly et al. 2015; Krogerus et al. 2017b, 2015b; Nikulin et al. 2018).

The genetic information for maltose utilization is well conserved in Saccharomyces species and depends on three gene families. MALT genes encode plasma-membrane proton symporters with varying substrate specificities and affinities (Cheng and Michels 1991; Vidgren et al. 2010), MALS genes encode α-glucosidases that hydrolyze α-oligoglucosides into glucose, while MALR genes encode a regulator required for transcriptional induction of MALT and MALS genes by maltose

(Naumov et al. 1994; Charron et al. 1989). In *Saccharomyces* species, maltose-utilization genes are

generally organized in *MAL* loci. These loci contain a *MALT* gene (called *ScMALx1* and *SeMALTx* in *S.*

cerevisiae and *S. eubayanus*, respectively), a *MALS* gene referred to as *ScMALx2* or *SeMALSx* and a

MALR gene referred to as *ScMALx3* or *SeMALRx* (Hayford and Jespersen 1999; Brickwedde et al.

2018). In the absence of glucose and presence of maltose, the MalR regulator binds a bi-directional

promoter, thereby simultaneously activating expression of *MALT* and *MALS* genes (Bell et al. 1997).

The Sc*MAL1*-Sc*MAL4* and Sc*MAL6* loci of *S. cerevisiae* as well as the Se*MAL1*-Se*MAL4* loci of *S.*

eubayanus are located in subtelomeric regions (Barnett 1976; Salazar et al. 2017; Brickwedde et al.

2018; Baker et al. 2015). While all *S. cerevisiae* *Sc*Malx1 transporters transport maltose, only *Sc*Mal11

is able to also transport maltotriose (Alves et al. 2008). *ScMAL11* (also known as *ScAGT1*) shares only

57% nucleotide identity with other *ScMALx1* genes (Han et al. 1995). The four *SeMALT* (*SeMALT1-4*)

genes identified in the genome of the Patagonian type strain FM1318/CBS 12357 of *S. eubayanus*

were shown to encode functional maltose transporters, but none of these genes enabled maltotriose

transport (Brickwedde et al. 2018). While no clear *ScAGT1* ortholog was found in *S. eubayanus* CBS

12357^{T}, such an ortholog was recently found in the genomes of two North American isolates assigned

to the Holarctic subclade of *S. eubayanus* (Baker and Hittinger 2018).

S. pastorianus inherited *MAL* genes from both *S. cerevisiae* and *S. eubayanus* (Van den Broek et al.

2015; Nakao et al. 2009; Vidgren and Gibson 2018). However, the *S. cerevisiae*-derived maltotriose-

transporter gene *ScAGT1* is truncated and, therefore, non-functional in *S. pastorianus* (Vidgren et al.

2009). Instead, maltotriose consumption by *S. pastorianus* strains was attributed to *SeAGT1* and

SpMTY1/SpMTT1 genes (Salema-Oom et al. 2005; Dietvorst et al. 2005; Cousseau et al. 2013; Nguyen

et al. 2011). In *S. pastorianus*, *SeAGT1* is located on *S. eubayanus* CHRXV and was therefore, already

before the identification of an *AGT1* ortholog in Holarctic *S. eubayanus* strains (Baker and Hittinger

2018), assumed to originate from *S. eubayanus* (Nakao et al. 2009). *SpMTY1,* also referred to as

SpMTT1, is located on *S. cerevisiae* CHRVII and has less than 92% sequence identity with other

Saccharomyces maltose transporters (Dietvorst et al. 2005). However, *SpMTY1* contains sequence

patches with high similarity to maltose transporters from *S. eubayanus* and *S. paradoxus* (Brouwers

et al. 2018). Recently, two independent laboratory evolution studies with *S. eubayanus*

demonstrated that recombination of different *SeMALT* genes yielded chimeric, neo-functionalized

genes that encoded maltotriose transporters (Brouwers et al. 2019; Baker and Hittinger 2019).

SpMTY1 may have resulted from successive introgressions of maltose-transporter genes from *S.*

cerevisiae, *S. eubayanus* and *S. paradoxus*.

Recently made *S. cerevisiae* x *S. eubayanus* laboratory hybrids showed similar lager-brewing

performance as *S. pastorianus* strains, also with respect to maltotriose utilization (Hebly et al. 2015;

Mertens et al. 2015; Krogerus et al. 2015b; Nikulin et al. 2018). In these hybrids, maltotriose

consumption depended on the presence of a functional *Sc*Agt1 transporter encoded by the *S.*

cerevisiae subgenome (Gorter de Vries et al. 2018). However, in view of the non-functionality of

ScAGT1 in current *S. pastorianus* strains, these laboratory hybrids did not fully recapitulate the

genetic landscape of *S. pastorianus* with respect to maltotriose fermentation (Hebly et al. 2015;

Mertens et al. 2015; Nakao et al. 2009).

Studies on laboratory hybrids based on *S. eubayanus* strains whose genomes are more closely related

to the *S. eubayanus* subgenome of *S. pastorianus* strains than that of the Patagonian type strain

CBS12357 might generate new insights into the evolution of maltotriose utilization in *S. pastorianus*.

To date, Himalayan *S. eubayanus* isolates show the highest sequence identity with the *S. eubayanus*

sub-genome of *S. pastorianus*, with up to 99.82% identity, as opposed to 99.56% for *S. eubayanus*

CBS12357[T] (Bing et al. 2014).

Here, we investigated if and how the genomes of Himalayan *S. eubayanus* strains could have

contributed to maltotriose utilization in the earliest hybrid ancestors of current *S. pastorianus*

strains. To this end, we generated chromosome-level genome assemblies of these strains by long-

read DNA sequencing. Since the Himalayan strains were unable to utilize maltotriose, we functionally

characterized the assembled *MAL* genes and identified genetic determinants that prevented

maltotriose utilization. Subsequently, a laboratory hybrid of a representative Himalayan *S. eubayanus*

strain and a maltotriose-deficient ale strain of *S. cerevisiae* was generated to investigate the genetics

of maltotriose utilization in a hybrid context. We discuss the implications of the experimental results

for the proposed role and origin of *SeAGT1* in *S. pastorianus* and for the potential of hybridization to

enable maltotriose consumption in novel *Saccharomyces* hybrids.

Materials and methods

Strains and maintenance

All strains used in this study are listed in Table 1. Stock cultures of *S. eubayanus* and *S. cerevisiae* strains were grown in YPD (10 g L^{-1} yeast extract, 20 g L^{-1} peptone and 20 g L^{-1} glucose) until late exponential phase, complemented with sterile glycerol to a final concentration of 30 % (v/v) and stored at -80 °C as 1 mL aliquots until further use.

Media and cultivation

S. eubayanus batch cultures were grown on synthetic medium (SM) containing 3.0 g L^{-1} KH_2PO_4, 5.0 g L^{-1} $(NH_4)_2SO_4$, 0.5 g L^{-1} $MgSO_4.7H_2O$, 1 mL L^{-1} trace element solution, and 1 mL L^{-1} vitamin solution (Verduyn et al. 1992). The pH was set to 6.0 with 2 M KOH prior to autoclaving at 120 °C for 20 min. Vitamin solutions were sterilized by filtration and added to the sterile medium. Concentrated sugar solutions were autoclaved at 110 °C for 20 min or filter sterilized and added to the sterile flasks to give a final concentration of 20 g L^{-1} glucose (SMG), maltose (SMM) or maltotriose (SMMt). With the exception of IMZ752 and IMZ753, *S. cerevisiae* batch cultures were grown on SM supplemented with 150 mg L^{-1} uracil (Pronk 2002) to compensate for loss of plasmid pUDC156 that carried the *Spcas9* endonuclease gene, and supplemented with 20 g L^{-1} glucose (SM_uG), maltose (SM_uM) or maltotriose (SM_uMt). All batch cultures were grown in 250 mL shake flasks with a working volume of 50 mL. The cultures were inoculated at an initial OD_{660nm} of 0.1 and incubated under an air atmosphere and shaken at 200 rpm and at 20 °C in a New Brunswick™ Innova44 incubator (Eppendorf Nederland B.V, Nijmegen, The Netherlands).

S. eubayanus strains transformed with plasmids pUDP052 ($gRNA_{SeSGA1}$), pUDP091 ($gRNA_{SeMALT1}$) and pUDP090 ($gRNA_{SeAGT1}$) were selected on modified SMG medium ($SM_{Ace}G$) in which $(NH_4)_2SO_4$ was replaced by 6.6 g L^{-1} K_2SO_4 and 10 mM acetamide (Solis-Escalante et al. 2013). SM-based solid media contained 2 % Bacto Agar (BD Biosciences, Franklin Lakes, NJ). *S. cerevisiae* strains expressing either *SeMALT, SeMALS* or *ScMALR* were selected on $SM_{Ace}G$. For plasmid propagation, *E. coli* XL1-Blue-derived strains (Agilent Technologies, Santa Clara, CA) were grown in Lysogeny Broth medium (LB, 10

g L^{-1} tryptone, 5 g L^{-1} yeast extract, 5 g L^{-1} NaCl) supplied with 100 mg L^{-1} ampicillin. Synthetic wort

medium (SWM) for growth studies contained 14.4 g·L^{-1} glucose, 2.3 g·L^{-1} fructose, 85.9 g·L^{-1} maltose,

26.8 g·L^{-1} maltotriose, 5 g·L^{-1} $(NH_4)_2SO_4$, 3 g·L^{-1} KH_2PO_4, 0.5 g·L^{-1} $MgSO_4.7H_2O$, 1 mL·L^{-1} trace element

solution, 1 mL·L^{-1} vitamin solution, supplemented with the anaerobic growth factors ergosterol and

Tween 80 (0.01 g·L^{-1} and 0.42 g·L^{-1} respectively), as previously described (Verduyn et al. 1992).

Industrial wort (containing 14.4 g L^{-1} glucose, 85.9 g L^{-1} maltose, 26.8 g L^{-1} maltotriose, 2.3 g L^{-1}

fructose and 269 mg L^{-1} FAN) was provided by Heineken Supply Chain B.V. (Zoeterwoude, the

Netherlands). The wort was supplemented with 1.5 mg L^{-1} of Zn^{2+} by addition of $ZnSO_4.7H_2O$,

autoclaved for 30 min at 121 ⬚C, filtered using Nalgene 0.2 μm SFCA bottle top filters (Thermo

Scientific) and diluted with sterile demineralized water. Sporulation medium consisted of 2 % (w/v)

KAc in MilliQ water set to pH 7.0 with KOH, autoclaved at 121 °C for 20 min.

Microaerobic growth experiments

Microaerobic cultures were grown in 250-mL airlock-capped Neubor infusion bottles (38 mm neck,

Dijkstra, Lelystad, Netherlands) containing 200 mL three-fold diluted industrial wort supplemented

with 0.4 mL L^{-1} Pluronic antifoam (Sigma-Aldrich, St. Louis, MO). Bottle caps were equipped with a 0.5

mm x 16 mm Microlance needle (BD Biosciences) sealed with cotton to prevent pressure build-up.

Sampling was performed aseptically with 3.5 mL syringes using a 0.8 mm x 50 mm Microlance needle

(BD Biosciences). Microaerobic cultures were inoculated at an OD_{660nm} of 0.1 from stationary-phase

precultures in 50 mL Bio-One Cellstar Cellreactor tubes (Sigma-Aldrich) containing 30 mL of the same

medium, grown for 4 days at 12 °C. Bottles were incubated at 12 °C and shaken at 200 rpm in a New

Brunswick Innova43/43R shaker (Eppendorf Nederland B.V.). At regular intervals, 3.5 mL samples

were collected in 24 deep-well plates (EnzyScreen BV, Heemstede, Netherlands) using a LiHa liquid

handler (Tecan, Männedorf, Switzerland) to measure OD_{660nm} and external metabolites. 30 μL of each

sample was diluted 5-fold in demineralized water in a 96 well plate and OD_{660nm} was measured with a

Magellan Infinite 200 PRO spectrophotometer (Tecan). From the remaining sample, 150 μL was

vacuum filter sterilized using 0.2 μm Multiscreen filter plates (Merck, Darmstadt, Germany) for HPLC

measurements.

Analytical methods

Optical densities of yeast cultures were measured with a Libra S11 spectrophotometer (Biochrom,

Cambridge, United Kingdom) at a wavelength of 660 nm. Biomass dry weight was measured by

filtering 10-mL culture samples over pre-weighed nitrocellulose filters with a pore size of 0.45 μm.

Filters were washed with 10 mL water, dried in a microwave oven (20 min at 350 W) and reweighed.

Sugars were measured using a high pressure liquid chromatography Agilent Infinity 1260 series

(Agilent Technologies) using a Bio-Rad Aminex HPX-87H column at 65 °C with 5 mM sulfuric acid at a

flow rate of 0.8 mL min^{-1}. Compounds were measured using a RID at 35 °C. Samples were centrifuged

at 13,000 g for 5 min to collect supernatant or 0.2 μm filter-sterilized before analysis.

Plasmid construction

Plasmids used and constructed in this study are listed in Table 2, oligonucleotide primers used in this

study are listed in Supplementary Table 1. Coding regions of *SeMALT1*, *SeMALT2*, *SeMALT3* and

SeAGT1 were amplified from CDFM21L.1 genomic DNA with Phusion High-Fidelity DNA polymerase

(Thermo Scientific), according to the supplier's instructions with primers pairs 12355/12356,

12357/12358, 12359/12360 and 12361/12362, respectively. The coding sequence of *ScMAL31* was

amplified from CEN.PK113-7D genomic DNA with Phusion High-Fidelity DNA polymerase (Thermo

Scientific), according to the supplier's instructions with primer pairs 9942/9943. Each primer carried a

40 bp extension complementary to the plasmid backbone of p426-TEF-amdS (Marques et al. 2017),

which was PCR amplified using Phusion High-Fidelity DNA polymerase (Thermo Scientific) and

primers 7812/5921. Each transporter fragment was assembled with the p426-TEF-amdS backbone

fragment using NEBuilder HiFi DNA Assembly (New England Biolabs, Ipswich, MA), resulting in

plasmids pUD444 (*ScMAL31*), pUD794 (*SeMALT1*), pUD795 (*SeMALT2*), pUD796 (*SeMALT3*) and

pUD797 (*SeAGT1*). All plasmids were verified for correct assembly by Sanger sequencing (Baseclear,

Leiden, The Netherlands).

SeMALS1 and *SeMALS2* were amplified from CDFM21L.1 genomic DNA with Phusion High-Fidelity

DNA polymerase (Thermo Scientific), with primers pairs 14451/14453 and 14452/14453,

respectively. Each primer pair carried a 30 bp extension complimentary to the plasmid backbone of

pUDE044 (de Kok et al. 2011) which was PCR amplified using Phusion High-Fidelity DNA polymerase

(Thermo Scientific) and primers 14449/14450. Resulting amplicons were assembled using NEBuilder

HiFi DNA Assembly (New England Biolabs), resulting in plasmids pUDE843 (*SeMALS1*) and pUDE844

(*SeMALS2*) that were verified by Sanger sequencing (Baseclear).

S. cerevisiae ScMAL13, the *ScPGK1* promoter and the *ScTEF2* terminator were amplified from

CEN.PK113-7D genomic DNA with Phusion High-Fidelity DNA polymerase (Thermo Scientific), with

primer pairs 12915/12916, 9421/9422 and 10884/10885, respectively. Fragments were gel purified

and used with pYTK074, pYTK082 and pYTK083 in Golden Gate assembly according to the yeast

toolkit protocol (Lee et al. 2015) resulting in pUDE780, which was verified by Sanger sequencing

(Baseclear).

Guide-RNA (gRNA) sequences for deletion of *SeMALT1* and *SeAGT1* in CDFM21L.1 were designed as

described previously (Gorter de Vries et al. 2017). The DNA sequences encoding these gRNAs were

synthesized at GeneArt (Thermo Scientific) and were delivered in pUD631 and pUD634, respectively.

The gRNA spacer sequences (Se*MALT1* 5' CCCCGATATTCTTTACACTA 3', Se*AGT1* 5'-

AGCTTTGCGAAAATATCCAA -3') and the structural gRNA sequence were flanked at their 5' ends by

the Hammerhead ribozyme (HH) and at their 3' ends by the Hepatitis Delta Virus ribozyme (HDV)

(Gao and Zhao 2014). The HH-gRNA-HDV fragment was flanked on both ends with a BsaI site for

further cloning (Juergens et al. 2018; Gorter de Vries et al. 2017). Plasmids pUDP091 (gRNA$_{SeMALT1}$)

and pUDP090 (gRNA$_{SeAGT1}$) were constructed by Golden Gate cloning by digesting pUDP004 and the

gRNA-carrying plasmid (pUD631 and pUD634, respectively) using BsaI and ligating with T4 ligase

(Engler et al. 2008). Correct assembly was verified by restriction analysis with PdmI (Thermo

Scientific) and Sanger sequencing (Baseclear).

Strain construction

S. cerevisiae IMZ616, which cannot grow on α-glucosides (Marques et al. 2018), was used as a host to

test functionality of individual *S. eubayanus* (putative) maltose transporter genes (Brickwedde et al.

2018). *S. cerevisiae* IMX1702 was constructed by integrating *ScTDH3$_p$-ScMAL12-ScADH1$_t$* and *ScTEF1$_p$-*

SeMALT1-ScCYC1$_t$ at the *ScSGA1* locus of strain IMZ616. A fragment containing the *ScTDH3$_p$-*

ScMAL12-ScADH1$_t$ transcriptional unit was PCR amplified using Phusion High-Fidelity DNA

polymerase (Thermo Scientific) from pUDE044 with primers 9596/9355, which included a 5'

extension homologous to the upstream region of the *ScSGA1* locus and an extension homologous to

the co-transformed transporter fragment, respectively. The DNA fragment carrying the *S. eubayanus*

SeMALT1 maltose symporter (*ScTEF1$_p$-SeMALT1-ScCYC1$_t$*) was PCR amplified from pUD794 using

primers 9036/9039, which included a 5' extension homologous to the co-transformed transporter

fragment and an extension homologous to the downstream region of the *ScSGA1* locus, respectively.

To facilitate integration in strain IMZ616, the two PCR fragments were co-transformed with plasmid

pUDR119 (*amdS*), which expressed a gRNA targeting *ScSGA1* (spacer sequence: 5'-

ATTGACCACTGGAATTCTTC-3') (van Rossum et al. 2016). The plasmid and repair fragments were

transformed using the LiAc yeast transformation protocol (Gietz and Schiestl 2007) and transformed

cells were plated on SM$_{Ace}$G. Correct integration was verified by diagnostic PCR with primers pairs

4226/4224. Strains *S. cerevisiae* IMX1704, IMX1706 and IMX1708 were constructed following the

same principle, but instead of using pUD794 to generate the transporter fragment, pUD795, pUD796

and pUD797 were used to PCR amplify *ScTEF1$_p$-SeMALT2-ScCYC1$_t$*, *ScTEF1$_p$-SeMALT3-ScCYC1$_t$* and

ScTEF1$_p$-SeAGT1-ScCYC1$_t$ respectively. IMX1313 was constructed in a similar way using only *ScTEF1$_p$-*

ScMAL31-ScCYC1$_t$ amplified with primer pair 9036/11018 which contain 5'and 3' extensions

homologous to the upstream and downstream region of the *ScSGA1* locus. Correct integration was

verified by diagnostic PCR with primer pair 4226/4224 (Supplementary Figure 1). All PCR-amplified

gene sequences were Sanger sequenced (BaseClear). IMX1313 was grown on YPD to loose pUDR119

(*URA3*) and pUDC156 (*amdS*). An isolate unable to grown on SMG without uracil and with acetamide

was selected and named IMX1313Δ. This strain was able to grow on SMG supplemented with 150 mg

L^{-1} uracil.

To assess functionality of CDFM21L.1 *SeMALS1*, IMX1313Δ was transformed with 100 ng pUDE843

(*ScTDH3$_p$-SeMALS1-ScADH1$_t$*) by electroporation (Gorter de Vries et al. 2017), resulting in strain

IMZ752. Transformants were selected on SMG plates after 5 days of incubation at 20 °C and

validated by PCR (DreamTaq polymerase, Thermo Scientific) using primer pair 14454/14455

(Supplementary Figure 1). Similarly, functionality of the *SeMALS2* maltase gene of CDFM21L.1 was

assessed by transforming IMX1313Δ with pUDE844 (*ScTDH3$_{pr}$-SeMALS2-ScADH1$_{ter}$*), resulting in strain

IMZ753.

S. eubayanus IMK820 (*SemalT1Δ*) was constructed by transforming CDFM21L.1 with 200 ng of

pUDP091 and 1 µg of 120 bp repair fragment obtained by mixing an equimolar amount of primers

12442/12443, as previously described (Gorter de Vries et al. 2017). As a control, the same

transformation was performed without including the repair DNA fragment. Transformants were

selected on SM$_{Ace}$G plates. *S. eubayanus* IMK823 (*Seagt1Δ*) was constructed similarly, using pUDP090

and primer pair 11320/11321. Deletion of *SemalT1* was verified by PCR with primer pair

11671/11672 and Sanger sequencing. The *Seagt1* deletion was verified by PCR using primer pair

12273/12274, and by Illumina whole-genome sequencing and read alignment to the reference

genome of CDFM21L.1 (Bioproject accession number PRJNA528469).

Strains IMX1765, IMX1939 and IMX1940 were constructed by inserting *ScPGK1$_p$-ScMAL13-ScTEF2$_t$* at

the *SeSGA1* locus of CDFM21L.1, IMK820 and IMK823, respectively. A repair fragment containing

ScPGK1$_p$-ScMAL13-ScTEF2$_t$ was amplified from pUDE780 with primer pair 12917/12918. Strains

CDFML21L.1, IMK820 and IMK823 were transformed by electroporation by addition of 350 ng of

repair fragment and 560 ng pUDP052 (*amdS*) into the cells as previously described (Gorter de Vries et

al. 2017). Transformants were plated on SM$_{Ace}$G and incubated at 20 °C. IMX1762 was constructed

similarly using a repair fragment with *ScTDH3$_p$-ScMAL12-ScADH1$_t$* amplified from pUDE044 with

primer pair 12319/12320. Strains were verified by PCR using primer pair 12635/12636 and Sanger

sequencing.

Hybrid Construction

The *S. cerevisiae* X *S. eubayanus* hybrid HTSH020 was constructed by spore-to-spore mating . The *S.*

eubayanus strain CDFM21L.1 and the *S. cerevisiae* strain CBC-1 were grown in 20 mL YPD at 20 °C

until late exponential phase. Cells were centrifuged for 5 min at 1000 g and washed twice in

demineralized water. Cells were re-suspended in 20 mL sporulation medium and incubated for 64 h

at 20 °C. Presence of spores was verified by microscopy. Asci were harvested by centrifugation for 5

min at 1000 g, and washed with demineralized water, resuspended in 100 μL demineralized water

containing 100 U/mL of Zymolyase (MP Bio, Santa Ana, CA) and incubated for 10 min at 30 °C. Spores

were washed and plated on the edge of a YPD agar plate. Spores from the two strains were brought

in contact with each other with an MSM System 400 micromanipulator (Singer Instruments, Watchet,

United Kingdom). Zygote formation was observed after 6-8 h. Emerging colonies were re-streaked

twice on SM 2% maltose at 12 °C. Successful hybridization was verified by multiplex PCR using

DreamTaq DNA polymerase (Thermo Scientific), by amplifying the *S. cerevisiae* specific *MEX67* gene

with primer pairs 8570/8571 and by amplifying the *S. eubayanus* specific gene *SeFSY1* with primers

8572/8573 (Supplementary Figure 2), as previously described (Pengelly and Wheals 2013).

Illumina sequencing

Genomic DNA of *S. eubayanus* strains CDFM21L.1 and ABFM5L.1, *S. cerevisiae* strain CBC-1 and *S.*

cerevisiae x *S. eubayanus* strain HTSH020 was isolated as previously described (Van den Broek et al.

2015). Paired-end sequencing (2 X 150 bp) was performed on a 350 bp PCR-free insert library using

Illumina HiSeq2500 (San Diego, CA) by Novogene(HK) Company Ltd (Hong Kong, China). Genomic

DNA of the strains CBC-1 and HTSH020 was sequenced in house on a MiSeq sequencer (Illumina)

with 300 bp paired-end reads using PCR-free library preparation. Sequence data are available at NCBI

under Bioproject accession number PRJNA528469.

MinION long read sequencing

For long-read sequencing, a 1D sequencing library (SQK-LSK108) was prepared for CDFM21L.1 and

CBC-1 and loaded onto an FLO-MIN106 (R9.4) flow cell, connected to a MinION Mk1B unit (Oxford

Nanopore Technology, Oxford, United Kingdom), according to the manufacturer's instructions.

MinKNOW software (version 1.5.12; Oxford Nanopore Technology) was used for quality control of

active pores and for sequencing. Raw files generated by MinKNOW were base-called using Albacore

(version 1.1.0; Oxford Nanopore Technology). Reads with a minimum length of 1000 bp were

extracted in fastq format. For CDFM21L.1, 3.26 Gb of sequence with an average read length of 8.07

kb was obtained and for CBC-1 3.04 Gb sequence with an average read length of 7.27 kb. Sequencing

data are available at NCBI under Bioproject accession number PRJNA528469.

De novo *assembly*

De novo assembly of the Oxford Nanopore MinION dataset was performed using Canu (v1.4, setting:

genomesize=12m) (Koren et al. 2017). Assembly correctness was assessed using Pilon (Walker et al.

2014) and further corrected by "polishing" of sequencing/assembly errors by aligning Illumina reads

with BWA (Li and Durbin 2010) using correction of only SNPs and short indels (−fix bases parameter).

For HTSH020, an artificial reference genome was made by combining the assembly of CBC-1 and

CDFM21L.1. The genome assemblies were annotated using the MAKER2 annotation pipeline (version

2.31.9) (Holt and Yandell 2011), using SNAP (version 2013−11-29) (Korf 2004) and Augustus (version

3.2.3) (Stanke et al. 2006) as *ab initio* gene predictors. *S. cerevisiae* S288C EST and protein sequences

were obtained from SGD (*Saccharomyces* Genome Database, http://www.yeastgenome.org/) and

were aligned using BLASTX on the obtained polished sequence assembly (BLAST version 2.2.28+)

(Camacho et al. 2009). Predicted translated protein sequences of the final gene model were aligned

to the *S. cerevisiae* S288C protein Swiss-Prot database using BLASTP (http://www.uniprot.org/).

Custom-made Perl scripts were used to map systematic names to the annotated gene names. Error

rates in nanopore-sequencing data were estimated from the q score (Phred scaled) per read, as

calculated by the base caller Albacore (version 1.1.0) (Oxford Nanopore Technology). Average q score

was used to calculate the error $P = 10^{q/10}$.

RNA isolation

CDFM21L.1, IMX1765, IMX1939 and HTS020 were grown in SMG, SMM and SMMt until mid-

exponential phase (OD_{660nm} of 12 for SMG/SMM and of OD_{660nm} 15 for SMMt). Culture samples

corresponding to ca. 200 mg of biomass dry weight were directly quenched in liquid nitrogen. The

samples were processed and total RNA extracted as previously described (Tai et al. 2005). Prior to

cDNA synthesis, purity, concentration and integrity of the RNA in the samples was assessed with

Nanodrop (Thermo Scientific), Qubit (Thermo Scientific) and Tapestation 220 with RNA Screen Tape

(Agilent Technologies), respectively, according the manufacturers' recommendations. cDNA libraries

were prepared using the TruSeq RNA V2 kit (Illumina). Paired-end sequencing (2 X 150 bp) was

performed on a 300 bp PCR-free insert library on a HISeq 2500 (Illumina) at Novogene (HK) Company

Ltd (Hong Kong, China). Duplicate biological samples were processed, generating an average

sequence quantity of 23.7M reads per sample. Reads were aligned to the CDFM21L.1 reference

assembly (GEO (https://www.ncbi.nlm.nih.gov/geo/) accession number GSE133146) using a two-pass

STAR (Dobin and Gingeras 2016) procedure. In the first pass, splice junctions were assembled and

used to inform the second round of alignments. Introns between 15 and 4000 bp were allowed, and

soft clipping was disabled to prevent low-quality reads from being spuriously aligned. Ambiguously-

mapped reads were removed from the dataset. Expression level for each transcript were quantified

using htseq-count (Anders et al. 2015) in union mode. Fragments per kilo-base of feature (gene) per

million reads mapped (FPKM) values were calculated by "Applying the fpkm method" from the edgeR

package (Robinson et al. 2010; McCarthy et al. 2012). Differential expression analysis was performed

using DESeq (Anders et al. 2013).

Results

Sequencing of Himalayan *S. eubayanus* strains revealed variations of sub-telomeric regions and presence of novel putative maltose transporter genes.

It has been proposed that the *S. eubayanus* genetic pool of *S. pastorianus* was inherited from an ancestor of the Asian *S. eubayanus* lineage (Bing et al. 2014). With 99.82% identity, the Himalayan *S. eubayanus* strains CDFM21L.1 and ABFM5L.1 that belong to the Holarctic lineage (Peris et al. 2016), present the closest characterized relatives of the *S. eubayanus* ancestor of lager brewing yeasts. However, this distance was based on a limited sequencing space (Bing et al. 2014) and the analysis did not investigate the presence of specific *S. eubayanus* genetic markers found in *S. pastorianus* hybrids. Therefore, the genome of the Himalayan *S. eubayanus* strain CDFM21L.1 was sequenced with a combination of long-read and short-read techniques (Oxford Nanopore MinION and Illumina technologies, respectively) to generate a near-complete draft reference genome sequence. The resulting CDFM21L.1 genome assembly comprised 19 contigs, including its mitochondrial genome. All chromosomes were completely assembled from telomere to telomere, except for chromosome XII, which was fragmented into 3 contigs due to the repetitive rDNA region and manually assembled into a single scaffold. With a total size of 12,034,875 bp, this assembly represents the first near-complete draft genome of a *S. eubayanus* strain of the Holarctic clade (Peris et al. 2016).

Chromosome-level assemblies were hitherto only available for the Patagonia B-clade strain CBS 12357[T] (Libkind et al. 2011; Brickwedde et al. 2018). We identified three major structural differences in CDFM21L.1 relative to CBS12357[T] using Mauve (Darling et al. 2004): (i) a paracentric inversion in the sub-telomeric region of chromosome VII, involving approximately 8 kbp, (ii) a translocation of approximately 12 kbp from the left sub-telomeric region of chromosome VIII to the right sub-telomeric region of chromosome VI, and (iii) a reciprocal translocation between approximately 20 kbp from the right sub-telomeric region of chromosome V and approximately 60 kbp from the center

of chromosome XII (Figure 1A). All structural variation involved sub-telomeric regions, in accordance

with their known relative instability (Brown et al. 2010; Moller et al. 2018; Gordon et al. 2009).

An alignment comparison of the CDFM21L.1 and CBS 12357^T genomes with MUMmer revealed that

557 kb were unique to CDFM21L.1 and reciprocally 428 kb were unique to CBS 12357^T. Sequences

unique to CBS 12357^T (3.6% of its genome) and to CDFM21L.1 (4.6% of its genome) were located

primarily in sub-telomeric regions and in repetitive regions, such as rDNA on chromosome XII (Figure

1B). Out of the 32 sub-telomeric regions, 23 exhibited absence of synteny. Conserved synteny was

observed for sub-telomeric regions on CHRIII (left), CHRIV (left and right), CHRVI (left), CHRIX (right),

CHRXI (right), CHRXII (right), CHRXIV (left) and CHRXV (right) (Supplementary File 1).

The 428 kb of sequence that were absent in the Himalyan *S. eubayanus* strain included 99 annotated

ORFs (Supplementary File 1). Of the 99 ORFs that were (partly) affected, 11 were completely absent

in CDFM21L.1, involving mostly genes implicated in iron transport facilitation (Supplementary File 1).

The 557 kb of sequence that was not present in CBS 12357^T included 113 annotated ORFs

(Supplementary File 1). Of these 113 ORFs, 15 were completely absent in CBS 12357^T. These 15 ORFs

showed an overrepresentation of genes involved in transmembrane transport (Fishers exact test, P-

value 4.8E^{-5}) (Figure 1C).

Of the 15 ORFs unique to CDFM21L.1, three were identical orthologs of *S. cerevisiae MAL11/AGT1*.

These three ORFs were found in the sub-telomeric regions of chromosomes VII, XIV and XV. Their

sequence similarity with the *S. cerevisiae* CEN.PK113.7D and *S. pastorianus* CBS 1483 *MAL11/AGT1*

genes was 82.7% and 99.89%, respectively. In addition to these *SeAGT1* genes, CDFM21L.1 genome

sequence harbored three maltose transporters (*SeMALTx*), two maltases (*SeMALSx*) and three

regulators (*SeMALRx*) encoding genes. In contrast to the situation in *S. eubayanus* CBS 12357^T, none

of the *SeMAL* genes formed a canonical *MAL* locus in CDFM21L.1 (Figure 2). A systematic sequence

inspection of these CDFM21L.1 *SeMAL* genes revealed mutations that prematurely interrupted the

reading frames of *SeMALR1* (CHRV), *SeMALT2* (CHRXII) and *SeMALT3* (CHRXII).

In addition to *S. eubayanus* CDFM21L.1 strain, a second Himalayan *S. eubayanus* isolate (ABFM5L.1)

was sequenced. These two strains were 99.97% genetically identical at the nucleotide level, their

MAL genes were syntenic and the premature stop codons in *SeMALR1* (CHRV), *SeMALT2* (CHRXII) and

*SeMALT*3 (CHRXII) were conserved. Two additional mutations were identified in one of the three

SeAGT1 genes. A nucleotide variation on position 53 and 939 (T instead of an A and A instead of a G))

resulted in a glycine to valine and arginine to lysine change, respectively.

Paradoxically, Himalayan *S. eubayanus* strains do not utilize maltose and maltotriose.

Identification of *SeAGT1* in the two Himalayan *S. eubayanus* strains suggests an ability to not only

grow on maltose but also on maltotriose. Strains from the Holarctic clade have previously been

hypothesized to be the donor of the *S. eubayanus* sub-genome in *S. pastorianus* hybrids (Bing et al.

2014; Peris et al. 2016). However, no physiological data regarding their ability to grow on the sugars

present in wort are available. To assess their growth characteristics, the *Asian S. eubayanus* strains

CDFM21L.1 and ABFM5L.1, the Patagonian *S. eubayanus* type strain CBS 12357^T and the *S.*

pastorianus strain CBS 1483 were grown on diluted industrial brewer's wort at 12 °C. As reported

previously, *S. pastorianus* strain CBS 1483 could utilize all three sugars but did not fully consume

maltotriose (Figure 3) (Brickwedde et al. 2017). Also in accordance with previous observations (Hebly

et al. 2015), CBS 12357^T consumed glucose and maltose completely, but left maltotriose untouched.

However, in marked contrast to *S. eubayanus* CBS 12357^T, neither CDFM21L.1 nor ABFM5L.1

consume maltose after growth on glucose. Moreover, like CBS 12357^T, maltotriose was not

metabolized by these two *S. eubayanus* strains. While in CBS 12357^T an ability to grow on maltose

and an inability to grow on maltotriose could be readily attributed to its *MAL* genes complement,

CDFM21L.1 and ABFM5L.1 failed to grow on maltose even though they appeared to contain

complete genes encoding maltose (*SeMALT1* and *SeAGT1*) and maltotriose (*SeAGT1*) transporters.

Growth defects on maltose and maltotriose are caused by deficiency of the regulatory *Se*MalR

proteins in *S. eubayanus* CDFM21L.1.

The recent characterization of maltose metabolism in CBS 12357^T showed that the coding regions of

transcriptionally silent maltose-transporter genes in *S. eubayanus* can potentially encode functional

proteins (Brickwedde et al. 2018). The inability of the *S. eubayanus* Himalayan isolates to grow on α-

oligosaccharides precluded direct testing of transporter-gene functionality by deletion studies.

Instead, these genes were expressed in *S. cerevisiae* IMZ616, which is devoid of all native maltose

metabolism genes (Marques et al. 2018). The CDFM21L.1 transporter genes *SeMALT1, SeMALT2,*

SeMALT3 or *SeAGT1* were integrated at the *ScSGA1* locus in IMZ616 along with the *S. cerevisiae*

maltase gene *ScMAL12* (Brickwedde et al. 2018), yielding a series of strains overexpressing a single

transporter (IMX1702 (*SeMALT1*), IMX1704 (*SeMALT2*), IMX1706 (*SeMALT3*) and IMX1708 (*SeAGT1*)).

These strains, as well as the negative and positive control strains IMZ616 and IMX1365 (IMZ616

expressing *ScAGT1* and *ScMAL12*), were grown on SM media supplemented with either maltose or

maltotriose. On maltose, not only the positive control strain IMX1365, but also IMX1702 (*SeMALT1*)

and IMX1708 (*SeAGT1*) were able to grow on maltose, consuming 30 and 60%, respectively, of the

initially present maltose after 100 h (Figure 4A). As anticipated, the *SeMALT2* and *SeMALT3* alleles

with premature stop codons did not support growth on maltose. Of the two strains that grew on

maltose, only IMX1708 (*SeAGT1*) also grew on maltotriose. These results demonstrate that *SeAGT1*

from a Holarctic *S. eubayanus* encoded a functional maltotriose transporter and, consequently, that

the inability of Holarctic strains to grow on maltose and maltotriose was not caused by transporter

dysfunctionality.

In addition to transport, metabolism of α-oligoglucosides requires maltase activity. Functionality of

the putative *SeMALS1* and *SeMALS2* maltase genes was tested by constitutive expression in strain

IMZ616, together with a functional *Sc*MAL31 transporter genes, yielding strains IMZ752 and IMZ753,

respectively. The maltase-negative strain IMX1313 was used as negative control. In SM medium with

maltose, both IMZ752 (*SeMALS1*) and IMZ753 (*SeMALS2*) grew and completely consumed maltose

within 65 h, demonstrating functionality of both hydrolase genes (Figure 4C).

In *S. cerevisiae* transcriptional regulation of *MALx2* and *MALx1* genes is tightly controlled by a

transcription factor encoded by *MALx3* genes. Malx3 binds an activating site located in the

bidirectional promoters that control expression of *MALx2* and *MALx1* genes (Meurer et al. 2017;

Levine et al. 1992). To test whether absence of maltose consumption in Himalayan *S. eubayanus*

strains was caused by a lack transcriptional upregulation of *SeMALT* and *SeMALS*, the *S. cerevisiae*

ScMAL13 gene was integrated at the *SeSGA1* locus in *S. eubayanus* CDFM21L.1, under the control of

a constitutive *ScPGK1* promoter and *ScTEF2* terminator. *ScMAL13* expression in CDFM21L.1 enabled

growth on maltose and maltotriose (Figure 5A), indicating that a lack of transcriptional upregulation

was indeed the cause of the parental strain's inability to grow on these oligoglucosides. However,

consumption of maltose and maltotriose was incomplete and consumed sugars were almost

exclusively respired, as no ethanol was measured after 60 h of cultivation.

The possibility to grow an engineered variant of *S. eubayanus* CDFM21L.1 on α-oligoglucosides

offered an opportunity to study transporter function in its native context. Complementary functional

characterization by gene deletion of *SeMALT1* and *SeAGT1* was performed using CRISPR-Cas9

genome editing method (Brickwedde et al. 2018; Gorter de Vries et al. 2017). Deletion of *SeMALT1*

and *SeAGT1* in CDFM21L.1 resulted in strains IMK820 and IMK823, respectively. Complete deletion of

SeAGT1 required disruption of six alleles. To confirm the complete removal of all copies, the genome

of IMK823 was sequenced. Mapping reads onto the reference *S. eubayanus* CDFM21L.1 genome

assembly confirmed that all six alleles were removed simultaneously. Subsequently, the regulator

expression cassette (*ScPGK1$_p$-ScMAL13-ScTEF2$_t$*) was integrated in IMK820 and IMK823 at the *SeSGA1*

locus yielding strains IMX1939 and IMX1940, respectively (Figure 5B). The four deletion strains

IMK820 (*SemalT1Δ*), IMK823 (*Seagt1Δ*), IMX1939 (*SemalT1Δ Sesga1Δ::ScMAL13*) and IMX1940

(*Seagt1Δ Sesga1Δ::ScMAL13*) were characterized on SMG, SMM or SMMt. All four strains were able

to grow on glucose (Supplementary Figure 3). While strains IMK820, IMK823 and IMX1940 were

unable to grow on maltose or maltotriose (Figure 5C), strain IMX1939 (*SemalT1Δ*

Sesga1Δ::SeMAL13), which harbored functional *SeAGT1* copies, grew on maltose as well as on

maltotriose. However, after 64 h of growth, these sugars were only partially consumed. Only 1.2 g L^{-1}

ethanol was produced from maltose and no ethanol formation was observed during growth on

maltotriose. The low ethanol concentration and the relatively high OD_{660nm} suggest that, under the

experimental conditions strain IMX1939 exhibited a Crabtree negative phenotype and exclusively

respired maltotriose. *S. eubayanus* IMX1940 (*Seagt1Δ Sesga1Δ::SeMAL13*) did not consume

maltotriose after 84 h of incubation. Moreover, despite the presence of *SeMALT1*, which encoded a

functional maltose transporter upon expression in *S. cerevisiae* IMZ616, strain IMX1940 was also

unable to consume maltose.

In addition to a functional Malx3 transcription factor, transcriptional activation of *MAL* genes also

requires presence of a cis-regulatory motif in the promoter of regulated genes. Transcriptome

analysis of *S. eubayanus* CBS 12357[T] recently showed that absence of a canonical cis-regulatory motif

in *SeMALT1* and *SeMALT3* of *S. eubayanus* CBS 12357[T] caused a deficiency in their expression

(Brickwedde et al. 2018). To further explore regulation of *SeMAL* and *SeAGT1* genes, we investigated

the impact of carbon sources on genome-wide transcriptome and, specifically, on transcriptional

activation of genes involved in maltose metabolism. Duplicate cultures of *S. eubayanus* strain

IMX1765 (*ScPGK1_p-ScMAL13-ScTEF2_t*) were grown on SMG, SMM and SMMt at 20 °C and sampled in

mid-exponential phase. After mRNA isolation and processing, cDNA libraries and reads were

assembled onto the newly annotated *S. eubayanus* CDFM21L.1 genome to calculate FPKM

(fragments per kilobase of feature (gene) per million reads mapped) expression values. The

heterologous regulator *ScMAL13*, expressed from the constitutive *ScPGK1* promoter, displayed the

same expression in glucose- and maltose-grown cultures. Although *ScMAL13* was efficiently

expressed on glucose, none of the nine *S. eubayanus* maltose genes (the three identical *SeAGT1*

copies being undistinguishable) were transcriptionally induced under these conditions (Figure 6),

confirming that the hierarchical regulatory role of glucose catabolite repression (Vidgren et al. 2011;

Meurer et al. 2017) also takes place in *S. eubayanus*. During growth on maltose, all nine genes were

significantly upregulated relative to glucose-grown cultures but large variations in expression level

were observed. The maltase genes *SeMALS1* and *SeMALS2* and the transporter gene *SeAGT1* showed

the highest upregulation, with fold changes of 148, 161 and 2355 respectively. Although upregulated

SeMALT1 displayed a fold change of 13, its normalized expression in maltose-grown cultures was

886-fold lower than that of *SeAGT1*. This weaker upregulation might explain why, despite the ability

of its coding region to support synthesis of a functional maltose transporter, *SeMALT1* alone could

not restore growth on maltose. The transcriptome data also revealed that the absence of maltose

induction in CDFM21L.1 was not associated with defective cis-regulatory elements in *SeMALR*

promoter sequences since the regulator genes were properly activated, instead these results would

suggest that the *Se*MalR regulator are not functional.

Hybridization of two maltotriose-deficient *S. eubayanus* and *S. cerevisiae* lineages results in

heterosis through regulatory crosstalk.

The genetic make-up of *S. pastorianus* lager-brewing yeasts strongly advocates that they originate

from hybridization of *S. cerevisiae* and *S. eubayanus* parental lineages that were both unable to

metabolize maltotriose (Nakao et al. 2009). This hypothesis is consistent with the recurrent mutation

in the *S. cerevisiae AGT1* allele of *S. pastorianus* strains as well as with the inability of Himalayan

strains of *S. eubayanus* to grow on these oligoglucosides.

Spores of the Himalayan *S. eubayanus* CDFM21L.1 were hybridized with *S. cerevisiae* CBC-1. This top-

fermenting *S. cerevisiae* is recommended for cask and bottle conditioning and unable to consume

maltotriose (Lallemand, Montreal, Canada). Analysis of the CBC-1 assembly, obtained by a

combination of long and short read sequencing, linked its maltotriose-negative phenotype to a total

absence of the *MAL11/AGT1* gene. The resulting laboratory interspecific hybrid HTSH020 was

characterized at 12 °C on synthetic wort, a defined medium whose composition resembles that of

brewer's wort. While *S. eubayanus* CDFM21L.1 only consumed glucose and *S. cerevisiae* CBC-1

consumed glucose and maltose after 103 h (Figure 7A), the interspecific hybrid HTSH020 completely

consumed glucose, maltose and partially consumed maltotriose after 105 h, thus resembling

characteristics of *S. pastorianus* strains (e.g. CBS1483 (Brickwedde et al. 2017)). In addition to this

gain of function, the hybrid HTSH020 outperformed both its parents on maltose consumption, since

it depleted this sugar in 70 h instead of 95 h for strain CBC-1. Since *S. cerevisiae* grows generally

slower at 12 °C, the experiments were also performed at 20 °C where HTSH020 consumed all maltose

16 h earlier than CBC-1 (Supplementary Figure 4).

Transcriptome analysis of the hybrid strain HTSH020 grown on SM with different carbon sources

showed that *SeAGT1* expression was repressed during growth on glucose, with a normalized

expression level of 7 FPKM (Figure 7B). When grown on SM maltose, *SeAGT1, SeMALS1* and *SeMALS2*

were significantly induced, with fold increases of 816, 109 and 116, respectively (Figure 7B). Although

SeMALT1 and *SeMALT2* were induced, these transporters do not contribute to maltose metabolism

due to truncation of their ORFs. These transcriptome data demonstrated that *SeAGT1* and *SeMALS*

genes are induced by regulatory crosstalk between regulators encoded from the CBC-1 *S. cerevisiae*

sub-genome and maltotriose transporter genes harbored by the *S. eubayanus* genome. This

laboratory hybridization experiment may be the closest reproduction yet of how, centuries ago,

maltotriose-fermentation capacity arose in the first hybrid ancestor of *S. pastorianus*.

Discussion

The ability to consume maltose and maltotriose represents a key performance indicator of *S.*

pastorianus lager-brewing strains (Vidgren et al. 2009). This study demonstrates how mating of *S.*

cerevisiae and *S. eubayanus* strains that cannot themselves ferment maltotriose, can yield

maltotriose-fermenting hybrids. This laboratory study illustrates how, centuries ago, maltotriose-

fermentation capacity may have arisen in the first hybrid ancestor of *S. pastorianus*.

While the origin of the *S. eubayanus* parent of *S. pastorianus* strains is still under debate

(Dunn and Sherlock 2008; Okuno et al. 2016; Monerawela and Bond 2017), phylogenetic analysis

suggested a Far East Asian origin (Bing et al. 2014). However, this interpretation was based on a

limited sequencing space and was constrained by the quality of available sequence assemblies. Since

an ortholog of *SeAGT1* had previously only been found in the *S. eubayanus* sub-genome of *S.*

pastorianus strains, this finding revived the discussion on the geographical origin of the ancestral *S.*

eubayanus parent (Baker and Hittinger 2018). The high-quality, annotated genome assemblies of the

Himalayan *S. eubayanus* strains CDFM21L.1 and ABFM5L.1 presented in the present study revealed

several copies of *SeAGT1*, whose very high sequence identity with *S. pastorianus SeAGT1* are

consistent with the previously proposed Asian origin of the *S. eubayanus* sub-genome of *S.*

pastorianus (Bing et al. 2014; Baker and Hittinger 2018; Peris et al. 2016). Next, genome-sequence

comparison of the Patagonian B sub-clade *S. eubayanus* strain CBS12357[T] and the Holarctic sub-clade

strain CDFM21L.1 revealed homoplasy of *SeAGT1*, probably reflecting that these sub-clades evolved

in different ecological niches.

The *S. eubayanus* wild stock whose genome sequence most closely corresponds to the *S.*

eubayanus sub-genome of *S. pastorianus* originates from the Tibetan plateau of the Himalaya (Bing

et al. 2014). However, the first *S. cerevisiae x S. eubayanus* hybrid, from which current lager yeasts

evolved by centuries of domestication, likely originates from a region between Bavaria and Bohemia

in Central Europe. So far, European *S. eubayanus* isolates have not been reported. This may indicate

that the original hybridization event occurred elsewhere or that the ancestral European lineage went

extinct. The recent detection, in a metagenomics analysis of samples from the Italian Alps, of ITS1

sequences corresponding to *S. eubayanus* could indicate that a wild European lineage exists after all

(Alsammar et al. 2019).

Functional characterization by heterologous complementation of an *S. cerevisiae* mutant

strain established that the *Se*Agt1 transporters from the Himalayan *S. eubayanus* strains CDFM21L.1

and ABFM5L supported uptake of maltose and maltotriose. After showing that these strains also

encoded a functional maltase gene, their inability to grow on maltose and maltotriose was attributed

to an inability to transcriptionally upregulate maltose metabolism genes likely caused by loss of

function mutations the regulator. In *S. cerevisiae* and, to some extent, ,in *S. eubayanus* strains of the

Patagonian B sub-clade such as CBS12357[T] (Baker et al. 2015; Brickwedde et al. 2018; Vidgren et al.

2005), MAL loci exhibit a specific organization in which a transporter (*MALT*) and a hydrolase (*MALS*)

gene are expressed from the same bidirectional promoter and are located adjacent to a regulator

gene (*MALR*) (Charron et al. 1989). In contrast, of the seven genomic regions harboring *MAL* genes in

the two Asian *S. eubayanus* strains, none showed this canonical organization (Figure 2) and the

subtelomeric regions carrying *SeAGT1* did not harbour sequences similar to hydrolase or regulator

genes. Subtelomeric regions harboring the other *MAL* genes indicated intensive reorganization as a

result of recombination. In particular, subtelomeric regions on CHRII, CHRV and CHRXII provide clear

indications for recombination events that scattered genes from ancestral MAL1 and MAL2 loci over

several chromosomes. A similar interpretation could explain the reorganization MAL3 on CHRXIII

(Figure 2). Similar events may have contributed to loss of function of the MAL regulators (MalR), as

exemplified by the occurrence of a non-synonymous mutation in *SeMALR1* resulting in loss of

function. These rearrangements did, however, not inactivate the *cis*-regulatory sequences of the

MAL genes, since complementation with a functional *ScMAL13* allele caused induction of most

SeMAL genes (Figure 6, Figure 7B) and, thereby, the heterotic maltotriose-positive phenotype of the

hybrid strain HTSH020. Together with the high copy number of *SeAGT1*, this heterotic

complementation may have been the main driver for colonization of low-temperature brewing

processes by the early hybrid ancestors of current *S. pastorianus* strains.

Recent work on adaptation to brewing environments of laboratory *S. cerevisiae* × *S. eubayanus*

hybrids showed loss maltotriose utilization during serial transfer in wort (Gorter de Vries et al. 2018).

A similar loss of maltotriose utilization is frequently encountered in *S. cerevisiae* ale strains (Gallone

et al. 2016), as well as in some Saaz-type *S. pastorianus* strains (Gibson et al. 2013). This is thus in

contrast with retention of a maltotriose assimilation phenotype by Frohberg-type *S. pastorianus*

strains. This may have been facilitated by the occurrence of multiple copies of the *SeAGT1* gene in

the *S. eubayanus* ancestor, which could act as a sequence buffer to counteracting adverse effects of

gene copy loss. The recent release of the first long-read sequencing assembly of *S. pastorianus*

enabled a precise chromosomal mapping of the maltose-metabolism genes (Salazar et al. 2019) and

showed that the Frohberg type *S. pastorianus* strains CBS 1483 harbored one copy of *SeAGT1* on the

S. eubayanus CHRXV section (as in CDFM21L.1) of the chimeric chromosome formed from *Se*CHRXV

and *Se*CHRVIII (Salazar et al. 2019).

Differential retention and loss of maltotriose consumption in *S. pastorianus* lineages may reflect

different brewing process conditions during domestication. In modern brewing processes based on

high-gravity wort, cell division is largely constrained to the glucose and maltose phases, which occur

before depletion of nitrogen sources (Lei et al. 2012). It may be envisaged that, in early lager-brewing

processes, unstandardized mashing processes generated wort with a higher maltotriose content,

which would have allowed for continued yeast growth during the maltotriose consumption phase.

During serial transfer on sugar mixtures, the selective advantage of consuming a specific sugar from a

mixture correlates with the number of generations on that sugar during each cycle (Papapetridis et

al. 2018; Wisselink et al. 2009). Such conditions would therefore have conferred a significant

selective advantage to a maltotriose-assimilating *S. cerevisiae* x *S. eubayanus* hybrid, especially if,

similar to current ale yeasts, the *S. cerevisiae* parent was unable to ferment maltotriose.

The heterotic phenotype that was reconstructed in the interspecies *S. cerevisiae* x *S.*

eubayanus hybrid HTS020 resulted from combination of dominant and recessive genetic variations

from both parental genomes. *S. eubayanus* contributed the *SeAGT1* gene and its functional *cis-*

regulatory sequences, but also harbored recessive mutations in *MALR* genes that allowed full

expression of the heterotic phenotype. These mutations were complemented with a set of *S.*

cerevisiae genes including a functional *MALR* and a non-functional *ScAGT1* gene to match the

mutations found in *S. pastorianus* (Nakao et al. 2009). Although some *S. pastorianus* strains harbor

an additional maltotriose transporter encoded by *SpMTT1 (*Dietvorst *et al. 2005)*, this gene was

recently proposed to have emerged after the original hybridization event as a result of repeated

recombination between *MALT* genes from both sub-genomes (Brouwers et al. 2018).

Maltotriose fermentation is likely not the only heterotic phenotype of S. *pastorianus* strains.

Flocculation or formation of complex aroma profiles (Bolat et al. 2013; Vidgren and Gibson 2018) are

phenotypes that are not fully understood and difficult to reproduce, that also might result from

heterosis (Gorter de Vries et al. 2018).

Laboratory-made *S. cerevisiae* x *S. eubayanus* hybrids hold great potential for brewing process

intensification and for increasing product diversity. In addition to increasing our understanding of the

evolutionary history of lager yeast genomes evolutionary, this study has implications for the design

of new hybrids. Hitherto, laboratory crosses of *S. cerevisiae* x *S. eubayanus* strains were designed

based on combination of dominant traits of the parental strains. Our results show that recessive

traits can be just as important as contributors to the genetic diversity of such hybrids.

Accession Numbers

653 The sequencing data were deposited at NCBI (https://www.ncbi.nlm.nih.gov/) under the Bioproject

654 PRJNA528469 and the transcriptomics data were deposited at GEO (Genome Expression Omnibus

655 https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE133146.

Supplementary Data
658 Supplementary Data are available

Authors contributions
661 JMD conceived the study and designed the experiments. NB, AB, LvbE, ARGdV, SMW and JD

662 performed the experimental work. NB and MvdB performed bioinformatics analysis. NB, AB, FYB, JTP

663 and JMGD supervised the study and wrote the manuscript. All authors read and approved the final

664 manuscript.

Acknowledgements

We thank Flip de Groot for his technical assistance in gene editing in *S. eubayanus*, Niels G.A.

Kuijpers, Viktor M. Boer and Jan-Maarten Geertman for their critical input.

Funding

This work was supported by the BE-Basic R&D Program (http://www.be-basic.org/), which was

granted an FES subsidy from the Dutch Ministry of Economic Affairs, Agriculture and Innovation

(EL&I).

Conflict of Interest

The authors declare no conflict of interest.

References

Alsammar HF, Naseeb S, Brancia LB, Gilman RT, Wang P, Delneri D (2019) Targeted metagenomics approach to capture the biodiversity of Saccharomyces genus in wild environments. Environmental Microbiology Reports 11 (2):206-214. doi:10.1111/1758-2229.12724

Alves SL, Herberts RA, Hollatz C, Trichez D, Miletti LC, De Araujo PS, Stambuk BU (2008) Molecular analysis of maltotriose active transport and fermentation by *Saccharomyces cerevisiae* reveals a determinant role for the AGT1 permease. Appl Environ Microbiol 74 (5):1494-1501

Anders S, McCarthy DJ, Chen Y, Okoniewski M, Smyth GK, Huber W, Robinson MD (2013) Count-based differential expression analysis of RNA sequencing data using R and Bioconductor. Nature protocols 8 (9):1765

Anders S, Pyl PT, Huber W (2015) HTSeq—a Python framework to work with high-throughput sequencing data. Bioinformatics 31 (2):166-169

Baker E, Wang B, Bellora N, Peris D, Hulfachor AB, Koshalek JA, Adams M, Libkind D, Hittinger CT (2015) The genome sequence of *Saccharomyces eubayanus* and the domestication of lager-brewing yeasts. Mol Biol Evol 32 (11):2818-2831

Baker EP, Hittinger CT (2018) Evolution of a novel chimeric maltotriose transporter in Saccharomyces eubayanus from parent proteins unable to perform this function. bioRxiv. doi:10.1101/431171

Baker EP, Hittinger CT (2019) Evolution of a novel chimeric maltotriose transporter in Saccharomyces eubayanus from parent proteins unable to perform this function. PLoS Genet 15 (4):e1007786. doi:10.1371/journal.pgen.1007786

Barnett JA (1976) The Utilization of Sugars by Yeasts1. In: Advances in carbohydrate chemistry and biochemistry, vol 32. Elsevier, pp 125-234

Bell PJ, Higgins VJ, Dawes IW, Bissinger PH (1997) Tandemly repeated 147 bp elements cause
structural and functional variation in divergent MAL promoters of Saccharomyces cerevisiae.
Yeast 13 (12):1135-1144
Bing J, Han P-J, Liu W-Q, Wang Q-M, Bai F-Y (2014) Evidence for a Far East Asian origin of lager beer
yeast. Curr Biol 24 (10):R380-R381
Bolat I, Romagnoli G, Zhu F, Pronk JT, Daran JM (2013) Functional analysis and transcriptional
regulation of two orthologs of ARO10, encoding broad-substrate-specificity 2-oxo-acid
decarboxylases, in the brewing yeast Saccharomyces pastorianus CBS1483. FEMS Yeast Res
13 (6):505-517. doi:10.1111/1567-1364.12051
Brickwedde A, Brouwers N, van den Broek M, Gallego Murillo JS, Fraiture JL, Pronk JT, Daran J-MG
(2018) Structural, physiological and regulatory analysis of maltose transporter genes in
Saccharomyces eubayanus CBS 12357T. Front Microbiol 9:1786
Brickwedde A, van den Broek M, Geertman J-MA, Magalhães F, Kuijpers NG, Gibson B, Pronk JT,
Daran J-MG (2017) Evolutionary engineering in chemostat cultures for improved maltotriose
fermentation kinetics in Saccharomyces pastorianus lager brewing yeast. Front Microbiol
8:1690
Brouwers N, Gorter de Vries AR, van den Broek M, Weening SM, Elink Schuurman TD, Kuijpers NGA,
Pronk JT, Daran J-MG (2018) In vivo recombination of Saccharomyces eubayanus maltose-
transporter genes yields a chimeric transporter that enables maltotriose fermentation.
bioRxiv. doi:10.1101/428839
Brouwers N, Gorter de Vries AR, van den Broek M, Weening SM, Elink Schuurman TD, Kuijpers NGA,
Pronk JT, Daran JG (2019) In vivo recombination of Saccharomyces eubayanus maltose-
transporter genes yields a chimeric transporter that enables maltotriose fermentation. PLoS
Genet 15 (4):e1007853. doi:10.1371/journal.pgen.1007853
Brown CA, Murray AW, Verstrepen KJ (2010) Rapid expansion and functional divergence of
subtelomeric gene families in yeasts. Curr Biol 20 (10):895-903
Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL (2009) BLAST+:
architecture and applications. BMC bioinformatics 10 (1):421
Charron MJ, Read E, Haut SR, Michels CA (1989) Molecular evolution of the telomere-associated MAL
loci of Saccharomyces. Genetics 122 (2):307-316
Cheng Q, Michels CA (1991) MAL11 and MAL61 encode the inducible high-affinity maltose
transporter of Saccharomyces cerevisiae. J Bacteriol 173 (5):1817-1820
Cousseau F, Alves Jr S, Trichez D, Stambuk B (2013) Characterization of maltotriose transporters from
the Saccharomyces eubayanus subgenome of the hybrid Saccharomyces pastorianus lager
brewing yeast strain Weihenstephan 34/70. Lett Appl Microbiol 56 (1):21-29
Darling ACE, Mau B, Blattner FR, Perna NT (2004) Mauve: multiple alignment of conserved genomic
sequence with rearrangements. Genome research 14 (7):1394-1403. doi:10.1101/gr.2289704
de Kok S, Yilmaz D, Suir E, Pronk JT, Daran J-MG, van Maris AJA (2011) Increasing free-energy (ATP)
conservation in maltose-grown Saccharomyces cerevisiae by expression of a heterologous
maltose phosphorylase. Metab Eng 13 (5):518-526
Dietvorst J, Londesborough J, Steensma H (2005) Maltotriose utilization in lager yeast strains: MTT1
encodes a maltotriose transporter. Yeast 22 (10):775-788
Dobin A, Gingeras TR (2016) Optimizing RNA-Seq mapping with STAR. In: Data Mining Techniques for
the Life Sciences. Springer, pp 245-262
Dunn B, Sherlock G (2008) Reconstruction of the genome origins and evolution of the hybrid lager
yeast Saccharomyces pastorianus. Genome Res 18 (10):1610-1623.
doi:10.1101/gr.076075.108
Engler C, Kandzia R, Marillonnet S (2008) A One Pot, One Step, Precision Cloning Method with High
Throughput Capability. PLOS ONE 3 (11):e3647. doi:10.1371/journal.pone.0003647
Entian K-D, Kötter P (2007) 25 yeast genetic strain and plasmid collections. Methods in Microbiology
36:629-666

Gallone B, Steensels J, Prahl T, Soriaga L, Saels V, Herrera-Malaver B, Merlevede A, Roncoroni M,
Voordeckers K, Miraglia L, Teiling C, Steffy B, Taylor M, Schwartz A, Richardson T, White C,
Baele G, Maere S, Verstrepen KJ (2016) Domestication and Divergence of *Saccharomyces*
cerevisiae Beer Yeasts. Cell 166 (6):1397-1410 e1316. doi:10.1016/j.cell.2016.08.020

Gao Y, Zhao Y (2014) Self-processing of ribozyme-flanked RNAs into guide RNAs in vitro and in vivo
for CRISPR-mediated genome editing. Journal of integrative plant biology 56 (4):343-349

Gibson BR, Storgårds E, Krogerus K, Vidgren V (2013) Comparative physiology and fermentation
performance of Saaz and Frohberg lager yeast strains and the parental species
Saccharomyces eubayanus. Yeast 30 (7):255-266. doi:10.1002/yea.2960

Gietz RD, Schiestl RH (2007) High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG
method. Nature protocols 2 (1):31

Gordon JL, Byrne KP, Wolfe KH (2009) Additions, losses, and rearrangements on the evolutionary
route from a reconstructed ancestor to the modern *Saccharomyces cerevisiae* genome. PLoS
Genet 5 (5):e1000485. doi:10.1371/journal.pgen.1000485

Gorter de Vries A, Voskamp MA, van Aalst ACA, Kristensen LH, Jansen L, van den Broek M, Salazar
AN, Brouwers N, Abeel T, Pronk JT, Daran J-MG (2018) Laboratory evolution of a
Saccharomyces cerevisiae x S. eubayanus hybrid under simulated lager-brewing conditions:
genetic diversity and phenotypic convergence. bioRxiv

Gorter de Vries AR, Groot PA, Broek M, Daran J-MG (2017) CRISPR-Cas9 mediated gene deletions in
lager yeast *Saccharomyces pastorianus*. Microb Cell Fact 16 (1):222

Han EK, Cotty F, Sottas C, Jiang H, Michels CA (1995) Characterization of *AGT1* encoding a general α-
glucoside transporter from *Saccharomyces*. Mol Microbiol 17 (6):1093-1107

Hayford A, Jespersen L (1999) Characterization of Saccharomyces cerevisiae strains from
spontaneously fermented maize dough by profiles of assimilation, chromosome
polymorphism, PCR and MAL genotyping. Journal of applied microbiology 86 (2):284-294

Hebly M, Brickwedde A, Bolat I, Driessen MR, de Hulster EA, van den Broek M, Pronk JT, Geertman J-
M, Daran J-MG, Daran-Lapujade P (2015) *S. cerevisiae × S. eubayanus* interspecific hybrid, the
best of both worlds and beyond. FEMS Yeast Res 15 (3)

Holt C, Yandell M (2011) MAKER2: an annotation pipeline and genome-database management tool
for second-generation genome projects. BMC bioinformatics 12 (1):491

Juergens H, Varela JA, Gorter de Vries AR, Perli T, Gast VJ, Gyurchev NY, Rajkumar AS, Mans R, Pronk
JT, Morrissey JP (2018) Genome editing in Kluyveromyces and Ogataea yeasts using a broad-
host-range Cas9/gRNA co-expression plasmid. FEMS yeast research 18 (3):foy012

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM (2017) Canu: scalable and accurate
long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res:gr.
215087.215116

Korf I (2004) Gene finding in novel genomes. BMC bioinformatics 5 (1):59

Krogerus K, Magalhães F, Vidgren V, Gibson B (2015a) New lager yeast strains generated by
interspecific hybridization. J Ind Microbiol Biotechnol 42 (5):769-778

Krogerus K, Magalhães F, Vidgren V, Gibson B (2015b) New lager yeast strains generated by
interspecific hybridization. Journal of Industrial Microbiology & Biotechnology 42 (5):769-
778. doi:10.1007/s10295-015-1597-6

Krogerus K, Magalhães F, Vidgren V, Gibson B (2017a) Novel brewing yeast hybrids: creation and
application. Appl Environ Microbiol 101 (1):65-78

Krogerus K, Magalhães F, Vidgren V, Gibson B (2017b) Novel brewing yeast hybrids: creation and
application. Appl Microbiol Biotechnol 101 (1):65-78

Lee ME, DeLoache WC, Cervantes B, Dueber JE (2015) A Highly Characterized Yeast Toolkit for
Modular, Multipart Assembly. ACS Synth Biol 4 (9):975-986. doi:10.1021/sb500366v

Lei H, Zhao H, Yu Z, Zhao M (2012) Effects of wort gravity and nitrogen level on fermentation
performance of brewer's yeast and the formation of flavor volatiles. Appl Biochem
Biotechnol 166 (6):1562-1574. doi:10.1007/s12010-012-9560-8
Levine J, Tanouye L, Michels CA (1992) The UAS(MAL) is a bidirectional promotor element required
for the expression of both the MAL61 and MAL62 genes of the Saccharomyces MAL6 locus.
Current genetics 22 (3):181-189
Li H, Durbin R (2010) Fast and accurate long-read alignment with Burrows–Wheeler transform.
Bioinformatics 26 (5):589-595
Libkind D, Hittinger CT, Valério E, Gonçalves C, Dover J, Johnston M, Gonçalves P, Sampaio JP (2011)
Microbe domestication and the identification of the wild genetic stock of lager-brewing
yeast. Proc Natl Acad Sci U S A:201105430
Marques WL, Mans R, Henderson RK, Marella ER, ter Horst J, de Hulster E, Poolman B, Daran J-M,
Pronk JT, Gombert AK (2018) Combined engineering of disaccharide transport and
phosphorolysis for enhanced ATP yield from sucrose fermentation in Saccharomyces
cerevisiae. Metabolic engineering 45:121-133
Marques WL, Mans R, Marella ER, Cordeiro RL, van den Broek M, Daran J-MG, Pronk JT, Gombert AK,
van Maris AJA (2017) Elimination of sucrose transport and hydrolysis in *Saccharomyces*
cerevisiae: a platform strain for engineering sucrose metabolism. FEMS Yeast Res 17
(1):fox006
McCarthy DJ, Chen Y, Smyth GK (2012) Differential expression analysis of multifactor RNA-Seq
experiments with respect to biological variation. Nucleic acids research 40 (10):4288-4297
Mertens S, Steensels J, Saels V, De Rouck G, Aerts G, Verstrepen KJ (2015) A large set of newly
created interspecific yeast hybrids increases aromatic diversity in lager beers. Appl Environ
Microbiol:AEM. 02464-02415
Meurer M, Chevyreva V, Cerulus B, Knop M (2017) The regulatable MAL32 promoter in
Saccharomyces cerevisiae: characteristics and tools to facilitate its use. Yeast 34 (1):39-49.
doi:doi:10.1002/yea.3214
Moller M, Habig M, Freitag M, Stukenbrock EH (2018) Extraordinary Genome Instability and
Widespread Chromosome Rearrangements During Vegetative Growth. Genetics 210 (2):517-
529. doi:10.1534/genetics.118.301050
Monerawela C, Bond U (2017) Brewing up a storm: The genomes of lager yeasts and how they
evolved. Biotechnology Advances 35 (4):512-519. doi:10.1016/j.biotechadv.2017.03.003
Nakao Y, Kanamori T, Itoh T, Kodama Y, Rainieri S, Nakamura N, Shimonaga T, Hattori M, Ashikari T
(2009) Genome sequence of the lager brewing yeast, an interspecies hybrid. DNA Res 16
(2):115-129
Naumov GI, Naumova ES, Michels C (1994) Genetic variation of the repeated *MAL* loci in natural
populations of *Saccharomyces cerevisiae* and *Saccharomyces paradoxus*. Genetics 136
(3):803-812
Nguyen H-V, Legras J-L, Neuvéglise C, Gaillardin C (2011) Deciphering the hybridisation history
leading to the lager lineage based on the mosaic genomes of *Saccharomyces bayanus* strains
NBRC1948 and CBS380T. PLoS One 6 (10):e25821
Nikulin J, Krogerus K, Gibson B (2018) Alternative Saccharomyces interspecies hybrid combinations
and their potential for low-temperature wort fermentation. Yeast (Chichester, England) 35
(1):113-127. doi:10.1002/yea.3246
Okuno M, Kajitani R, Ryusui R, Morimoto H, Kodama Y, Itoh T (2016) Next-generation sequencing
analysis of lager brewing yeast strains reveals the evolutionary history of interspecies
hybridization. DNA Res 23 (1):67-80. doi:10.1093/dnares/dsv037
Papapetridis I, Verhoeven MD, Wiersma SJ, Goudriaan M, van Maris AJA, Pronk JT (2018) Laboratory
evolution for forced glucose-xylose co-consumption enables identification of mutations that
improve mixed-sugar fermentation by xylose-fermenting Saccharomyces cerevisiae. FEMS
Yeast Research 18 (6). doi:10.1093/femsyr/foy056

Pengelly RJ, Wheals AE (2013) Rapid identification of *Saccharomyces eubayanus* and its hybrids.
FEMS Yeast Res 13 (2):156-161
Peris D, Langdon QK, Moriarty RV, Sylvester K, Bontrager M, Charron G, Leducq J-B, Landry C, Libkind
D, Hittinger CT (2016) Complex ancestries of lager-brewing hybrids were shaped by standing
variation in the wild yeast Saccharomyces eubayanus. PLoS Genet 12 (7):20.
doi:10.1371/journal.pgen.1006155
Pronk JT (2002) Auxotrophic Yeast Strains in Fundamental and Applied Research. Applied and
Environmental Microbiology 68 (5):2095. doi:10.1128/AEM.68.5.2095-2100.2002
Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics 26 (1):139-140
Salazar AN, Gorter de Vries AR, van den Broek M, Brouwers N, de la Torre Cortès P, Kuijpers NGA,
Daran J-MG, Abeel T (2019) Nanopore sequencing and comparative genome analysis confirm
lager-brewing yeasts originated from a single hybridization. bioRxiv:603480.
doi:10.1101/603480
Salazar AN, Gorter de Vries AR, van den Broek M, Wijsman M, de la Torre Cortés P, Brickwedde A,
Brouwers N, Daran J-MG, Abeel T (2017) Nanopore sequencing enables near-complete de
novo assembly of *Saccharomyces cerevisiae* reference strain CEN. PK113-7D. FEMS Yeast Res
17 (7)
Salema-Oom M, Pinto VV, Gonçalves P, Spencer-Martins I (2005) Maltotriose utilization by industrial
Saccharomyces strains: characterization of a new member of the α-glucoside transporter
family. Appl Environ Microbiol 71 (9):5044-5049
Solis-Escalante D, Kuijpers NG, Nadine B, Bolat I, Bosman L, Pronk JT, Daran J-MG, Pascale D-L (2013)
amdSYM, a new dominant recyclable marker cassette for *Saccharomyces cerevisiae*. FEMS
Yeast Res 13 (1):126-139
Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B (2006) AUGUSTUS: ab initio
prediction of alternative transcripts. Nucleic acids research 34 (suppl_2):W435-W439
Tai SL, Boer VM, Daran-Lapujade P, Walsh MC, de Winde JH, Daran J-M, Pronk JT (2005) Two-
dimensional transcriptome analysis in chemostat cultures combinatorial effects of oxygen
availability and macronutrient limitation in Saccharomyces cerevisiae. Journal of Biological
Chemistry 280 (1):437-447
Van den Broek M, Bolat I, Nijkamp JF, Ramos E, Luttik MAH, Koopman F, Geertman JM, De Ridder D,
Pronk JT, Daran J-MG (2015) Chromosomal copy number variation in *Saccharomyces*
pastorianus evidence for extensive genome dynamics in industrial lager brewing strains. Appl
Environ Microbiol:AEM. 01263-01215
van Rossum HM, Kozak BU, Niemeijer MS, Duine HJ, Luttik MA, Boer VM, Kötter P, Daran J-MG, van
Maris AJA, Pronk JT (2016) Alternative reactions at the interface of glycolysis and citric acid
cycle in *Saccharomyces cerevisiae*. FEMS Yeast Res 16 (3)
Verduyn C, Postma E, Scheffers WA, Van Dijken JP (1992) Effect of benzoic acid on metabolic fluxes in
yeasts: a continuous-culture study on the regulation of respiration and alcoholic
fermentation. Yeast 8 (7):501-517
Vidgren V, Gibson B (2018) Trans-regulation and localization of orthologous maltose transporters in
the interspecies lager yeast hybrid. FEMS yeast research 18 (6):foy065.
doi:10.1093/femsyr/foy065
Vidgren V, Huuskonen A, Virtanen H, Ruohonen L, Londesborough J (2009) Improved fermentation
performance of a lager yeast after repair of its *AGT1* maltose and maltotriose transporter
genes. Appl Environ Microbiol 75 (8):2333-2345
Vidgren V, Kankainen M, Londesborough J, Ruohonen L (2011) Identification of regulatory elements
in the AGT1 promoter of ale and lager strains of brewer's yeast. Yeast 28 (8):579-594.
doi:doi:10.1002/yea.1888

Vidgren V, Multanen JP, Ruohonen L, Londesborough J (2010) The temperature dependence of
maltose transport in ale and lager strains of brewer's yeast. FEMS Yeast Res 10 (4):402-411.
doi:10.1111/j.1567-1364.2010.00627.x
Vidgren V, Ruohonen L, Londesborough J (2005) Characterization and functional analysis of the MAL
and MPH Loci for maltose utilization in some ale and lager yeast strains. Appl Environ
Microbiol 71 (12):7846-7857. doi:10.1128/AEM.71.12.7846-7857.2005
Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J,
Young SK (2014) Pilon: an integrated tool for comprehensive microbial variant detection and
genome assembly improvement. PloS one 9 (11):e112963
Walther A, Hesselbart A, Wendland J (2014) Genome sequence of Saccharomyces carlsbergensis, the
world's first pure culture lager yeast. G3 (Bethesda) 4 (5):783-793.
doi:10.1534/g3.113.010090
Wisselink HW, Toirkens MJ, Wu Q, Pronk JT, van Maris AJA (2009) Novel evolutionary engineering
approach for accelerated utilization of glucose, xylose, and arabinose mixtures by engineered
Saccharomyces cerevisiae strains. Appl Environ Microbiol 75 (4):907-914
Zastrow C, Hollatz C, De Araujo P, Stambuk B (2001) Maltotriose fermentation by *Saccharomyces*
cerevisiae. J Ind Microbiol Biotechnol 27 (1):34-38
Zheng X, D'Amore T, Russell I, Stewart G (1994) Factors influencing maltotriose utilization during
brewery wort fermentations. J Am Soc Brew Chem 52 (2):41-47

Name	Species	Relevant genotype	Origin
CDFM21L.1	*S. eubayanus*	Wildtype Mal[-] Mtt[-]	(Bing et al. 2014)
ABFM5L.1	*S. eubayanus*	Wildtype Mal[-] Mtt[-]	(Bing et al. 2014)
CBS 12357	*S. eubayanus*	Wildtype Mal[+] Mtt[-]	(Libkind et al. 2011) Westerdijk institute[a]
CBS 1483	*S. pastorianus*	Wildtype Mal[+] Mtt[+]	Westerdijk institute[a]
CEN.PK113-7D	*S. cerevisiae*	*MATa MAL1x MAL2x MAL3x MAL4x MAL2-8C SUC2 LEU2 URA3*	(Entian and Kötter 2007)
IMZ616	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mp2/3hΔ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ pUDC156 (Spcas9 URA3 ARS4 CEN6)*	(Marques et al. 2018)
IMX1365	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ pUDC156 (URA3 cas9) sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-ScAGT1-ScCYC1$_t$*	(Brickwedde et al. 2018)
IMX1702	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ pUDC156 (URA3 cas9) sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-SeMALT1-ScCYC1$_t$*	This study
IMX1704	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ pUDC156 (URA3 cas9) sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-SeMALT2-ScCYC1$_t$*	This study
IMX1706	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ pUDC156 (URA3 cas9) sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-SeMALT3-ScCYC1$_t$*	This study
IMX 1708	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ pUDC156 (URA3 cas9) sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-SeAGT1-ScCYC1$_t$*	This study
IMX1313	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ sga1Δ::ScTEF1$_p$-ScMAL31-ScCYC1$_t$ pUDC156 (URA3 cas9)*	This study
IMX1313Δ	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ sga1Δ::ScTEF1$_p$-ScMAL31-ScCYC1$_t$*	This study
IMZ752	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ sga1Δ::ScTEF1$_p$-ScMAL31-ScCYC1$_t$ pUDE843 (ori (ColE1) bla 2μ ScTDH3$_p$-SeMALS1-ScADH1$_t$ URA3)*	This study
IMZ753	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ sga1Δ::ScTEF1$_p$-ScMAL31-ScCYC1$_t$ pUDE844 (ori (ColE1) bla 2μ ScTDH3$_p$-SeMALS2-ScADH1$_t$ URA3)*	This study
IMK820	*S. eubayanus*	*MATa/MATα Semalt1Δ/Semalt1Δ*	This study
IMK823	*S. eubayanus*	*MATa/MATα Seagt1Δ/Seagt1Δ (X3)*	This study
IMX1939	*S. eubayanus*	*MATa/MATα Semalt1Δ/Semalt1Δ Sesga1Δ::ScMAL13/Sesga1Δ::ScMAL13*	This study
IMX1940	*S. eubayanus*	*MATa/MATα Seagt1Δ/Seagt1Δ Sesga1Δ::ScMAL13/Sesga1Δ::ScMAL13*	This study
IMX1762	*S. eubayanus*	*MATa/MATα Sesga1Δ::ScMAL12/Sesga1Δ::ScMAL12*	This study
IMX1765	*S. eubayanus*	*MATa/MATα Sesga1Δ::ScMAL13/Sesga1Δ::ScMAL13*	This study
CBC-1	*S. cerevisiae*	*MATa/MATα Mal[+] Mtt[-]*	Lallemand
HTSH020	*S. cerevisiae X S. eubayanus*	*MATa/MATα Mal[+] Mtt[+]*	This study

Table 2 Plasmids used in this study

Plasmid	Relevant genotype	Origin
p426-TEF-amdS	*ori* (*ColE1*) *bla* 2μ amdSYM *TEF1_{pr}-CYC1_{ter}*	(Marques et al. 2017)
pUD444	*ori* (*ColE1*) *bla* 2μ amdSYM *ScTEF1_{pr}-ScMAL31-ScCYC1_{ter}*	This study
pUD794	*ori* (*ColE1*) *bla* 2μ amdSYM *ScTEF1_{pr}-SeMALT1-ScCYC1_{ter}*	This study
pUD795	*ori* (*ColE1*) *bla* 2μ amdSYM *ScTEF1_{pr}-SeMALT2-ScCYC1_{ter}*	This study
pUD796	*ori* (*ColE1*) *bla* 2μ amdSYM *ScTEF1_{pr}-SeMALT3-ScCYC1_{ter}*	This study
pUD797	*ori* (*ColE1*) *bla* 2μ amdSYM *ScTEF1_{pr}-SeAGT1-ScCYC1_{ter}*	This study
pUDE044	*ori* (*ColE1*) *bla* 2μ *ScTDH3_{pr}-ScMAL12-ScADH1_{ter} URA3*	(de Kok et al. 2011)
pUDE843	*ori* (*ColE1*) *bla* 2μ *ScTDH3_{pr}-SeMALS1-ScADH1_{ter} URA3*	This study
pUDE844	*ori* (*ColE1*) *bla* 2μ *ScTDH3_{pr}-SeMALS2-ScADH1_{ter} URA3*	This study
pUDE780	*ori* (*ColE1*) *bla* 2μ *ScPGK1_{pr}-ScMAL13-ScTEF2_{ter} URA3*	This study
pUDP002	*ori* (*ColE1*) *bla* panARSopt *Hyg ScTDH3_{pr}*-BsaI-BsaI-*ScCYC1_{ter} AaTEF1_{pr}-Spcas9^{D147Y P411T}-ScPHO5_{ter}*	(Juergens et al. 2018)
pUDP004	*ori* (*ColE1*) *bla* panARSopt amdSYM *ScTDH3_{pr}*-BsaI-BsaI-*ScCYC1_{ter} AaTEF1_{pr}-Spcas9^{D147Y P411T}-ScPHO5_{ter}*	(Gorter de Vries et al. 2017)
pUDP052	ori (ColE1) *bla* panARSopt amdSYM *ScTDH3_{pr}*- gRNA_{SGA1}-*ScCYC1_{ter} AaTEF1_{pr}-Spcas9^{D147Y P411T}-ScPHO5_{ter}*	(Brickwedde et al. 2018)
pUDP091	ori (ColE1) *bla* panARSopt amdSYM *ScTDH3_{pr}*- gRNA_{SeMALT1}-*ScCYC1_{ter} AaTEF1_{pr}-Spcas9^{D147Y P411T}-ScPHO5_{ter}*	This study
pUDP090	ori (ColE1) *bla* panARSopt amdSYM *ScTDH3_{pr}*- gRNA_{SeAGT1}-*ScCYC1_{ter} AaTEF1_{pr}-Spcas9^{D147Y P411T}-ScPHO5_{ter}*	This study
pUDR119	ori (ColE1) *bla* 2μ amdSYM *SNR52_{pr}*-gRNA_{ScSGA1}-*SUP4_{ter}*	(van Rossum et al. 2016)
pYTK074	ori (ColE1) *cat URA3*	(Lee et al. 2015)
pYTK082	*cat* 2μ	(Lee et al. 2015)
pYTK083	ori (ColE1) *bla*	(Lee et al. 2015)
pUD631	ori (ColE1) *bla* gRNA_{SeMALT1}	(Brickwedde et al. 2018)
pUD634	ori (ColE1) *bla* gRNA_{SeAGT1}	This study
pUDC156	ori (ColE1) *bla ARS4 CEN6 URA3 Spcas9*	(Marques et al. 2018)

Legends to Figures

Figure 1. Genome comparison between CDFM21L.1 and CBS 12357[T]. (A) Translocations in

CDFM21L.1 relative to CBS12357[T], (i) a paracentric inversion in the sub-telomeric region of

chromosome VII, involving approximately 8 kbp, (ii) a translocation of approximately 12 kbp from the

left sub-telomeric region of chromosome VIII to the right sub-telomeric region of chromosome VI,

and (iii) a reciprocal translocation between approximately 20 kbp from the right sub-telomeric region

of chromosome V and approximately 60 kbp from the center of chromosome XII.Red lines depicted

chromosomes separation. Genome syntheny is indicated with colored blocks. **(B)** Relative

chromosome position of gene presence differences between CDFM21L.1 (blue) and CBS 12357[T] (red).

(C) Representation of the assembled CDFM21L.1 *S. eubayanus* chromosomes, the black boxes denote

newly added sequences. New annotated open reading frames and gene entries modified relative to

the CBS 12357T draft genome (Brickwedde et al. 2018).

Figure 2. Organization of sub-telomeric regions involving *MAL* genes and *SeAGT1* in CDFM21L.1. (A)

Chromosome sections are represented as lines and red boxes denote telomeres. CDFM21L.1 genome

harbors three *SeMALT* genes of which *SeMALT2* and *SeMALT3* have a mutation resulting in an early

stop codon and truncated protein (denoted with *). Three copies of *SeAGT1* were found close to the

telomeres on CHRVII, XIV and XV. Furthermore there are two intact *SeMALS* genes on CHRII (1) and

XII (2) and three *SeMALR* genes on CHRV (1) and XIII (2) whose copy on CHRV is also mutated

(*SeMALR1**). The gene and interval sizes are approximately to scale. **(B)** Phylogeny of *Saccharomyces*

SeAGT1 genes described in *S. cerevisiae*, *S. eubayanus*, and lager-brewing hybrid *S. pastorianus*. (C)

Nucleotide percentage identities between *AGT1* orthologs from *S. cerevisiae*, *S. eubayanus*, and

lager-brewing hybrid *S. pastorianus*. Green color indicates highest similarity between SeAGT1 and

SeAGT1 genes from *S. pastorianus* strains CBS 1483 and WS3470. Red color indicates similarity

between SeAGT1 from North American strains with SeAGT1 genes from Asian *S. eubayanus* and *S.*

pastorianus strains CBS 1483 and WS3470.

Figure 3. Characterization of sugar consumption of *S. pastorianus* CBS1483 (black) and *S. eubayanus* CBS 12357^T (blue), CDFM21L.1 (red), and ABFM5L.1 (green) on wort. For every sample, glucose (■), maltose (▲), maltotriose (▼) and ethanol (◆) were measured from the supernatant. Strains were grown at 12 °C for 110 hours in infusion Neubor flasks. Samples were filtered through a 0.22 µm filter and analyzed on HPLC. Data represents average and standard deviation of three biological replicates.

Figure 4. Overexpression of *SeMALT*, *SeAGT1* and *SeMALS* genes in a maltose negative background *S. cerevisiae* strain. Maltose negative background strain IMZ616 (■), IMX1365 overexpressing *ScMAL11* (▲), IMX1702 overexpressing *SeMALT1* (▼), IMX1704 overexpressing *SeMALT2* (◆) IMX1706 overexpressing *SeMALT3* (●) and IMX1708 overexpressing SeAGT1 (□) were grown on SM 2% maltose or maltotriose at 20 °C. Growth on maltose **(A)** and on maltotriose **(B)** was monitored based on optical density (OD$_{660nm}$) and concentrations of maltose and maltotriose in culture supernatants were measured by HPLC. Data are presented as average and standard deviation of two biological replicates. **(C)** IMX1313 overexpressing only *ScMAL31* (○), IMZ752 overexpressing *ScMAL31* and *SeMALS1* (△) and IMZ753 overexpressing *ScMAL31* and *SeMALS2* (▽) grown on SM maltose 2%. Growth was monitored based on optical density measurement at 660 nm (OD$_{660nm}$) and maltose in culture supernatants was measured by HPLC. Data represents average and standard deviation of two biological replicates.

Figure 5. Integration of *ScMAL13* in CDFM21L.1 restores growth on maltose and maltotriose and enables native *SeMALT1* and *SeAGT1* characterization in knockout strains IMK820 and IMK823. (A) Characterization of *S. eubayanus* IMX1765 overexpressing *ScMAL13* (▼) and CDFM21L.1 (■) on SM with glucose, maltose or maltotriose at 20 °C. OD$_{660nm}$ was measured (black) and sugar (black) and ethanol (red) concentrations were determined from the supernatant by HPLC. **(B)** Overview of

constructed knockout strains. Knockouts of *SeMALT1* (IMK820) and *SeAGT1* (IMK823) were made

with CRISPR-Cas9. Subsequently the *SeSGA1* locus was replaced by *ScPGK1$_p$- ScMAL13-ScTEF2$_t$* using

CRISPR-Cas9 in both strains resulting in IMX1939 and IMX1940, respectively. **(C)** *S. eubayanus* strains

IMK820 (■), IMK823 (▲), IMX1939 (▼) and IMX1940 (◆) were characterized on SM with maltose or

maltotriose at 20 °C. OD_{660nm} was measured (black) and sugar (black) and ethanol (red)

concentrations were determined from the supernatant by HPLC. All data represents average and

standard deviation of biological duplicates.

Figure 6. Expression levels of maltose metabolism genes in IMX1765. Normalized transcript levels of

maltose metabolism genes from IMX1765mid-exponential phase grown on glucose (red), maltose

(blue)and maltotriose (green) at 20 °C were calculated from duplicate RNA sequencing experiments

(2x 150 bp) using the FPKM method . All data represents average and standard deviation of two

biological duplicates.

Figure 7. Hybridization of maltotriose deficient *S. cerevisiae* and *S. eubayanus* leading to crosstalk

restoring maltotriose utilization, explains *S. pastorianus* phenotype. (A) Characterization of *S.*

cerevisiae CBC-1 (▼), *S.* eubayanus CDFM21L.1 (▲) and hybrid HTSO20 (■) on mock wort at 12 °C.

Consumption of maltose (red), maltotriose (blue) and production of ethanol (green) was measured

from supernatant by HPLC. Data represents average and standard deviation from biological

triplicates. **(B)** Normalized transcript levels of maltose metabolism genes from HTSO20 mid-

exponential phase grown on glucose (red), maltose (blue)and maltotriose (green) at 20 °C were

calculated from duplicate RNA sequencing experiments (2x 150 bp) using the FPKM method . All

data represents average and standard deviation of two biological duplicates.

1000

1001

FPKM