Maltotriose consumption by hybrid *Saccharomyces pastorianus* is heterotic and results from regulatory cross-talk between parental sub-genomes

Heterotic origin of maltotriose consumption in Saccharomyces pastorianus

Nick Brouwers[a] , Anja Brickwedde[a], Arthur R. Gorter de Vries[a], Marcel van den Broek[a], Susan M. Weening[a], Lieke van den Eijnden[a], Jasper A. Diderich[a], Feng-Yan Bai[b], Jack T. Pronk[a] and Jean-Marc G. Daran[a]*

[a]Department of Biotechnology, Delft University of Technology, Van der Maasweg 9, 2629 HZ Delft, The Netherlands.
[b]State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing, 100101, China.
* Corresponding author: Jean-Marc G. Daran, j.g.daran@tudelft.nl, Department of Biotechnology, Delft University of Technology, van der Maasweg 9, 2629 HZ Delft, The Netherlands

Nick Brouwers:	N.Brouwers-1@tudelft.nl	
Anja Brickwedde:	a.brickwedde@web.de	
Arthur R. Gorter de Vries:	A.R.GorterdeVries@tudelft.nl	0000-0002-0841-6583
Marcel van den Broek:	Marcel.vandenBroek@tudelft.nl	
Susan M. Weening:	S.M.Weening@tudelft.nl	
Lieke van den Eijnden:	A.F.vandenEijnden@student.tudelft.nl	
Jasper A. Diderich:	J.A.Diderich@tudelft.nl	
Feng-Yan Bai:	baify@im.ac.cn	
Jack T. Pronk:	J.T.Pronk@tudelft.nl	0000-0002-5617-4611
Jean-Marc G. Daran:	J.G.Daran@tudelft.nl	0000-0003-3136-8193

Keywords: *S. eubayanus*, α-oligo-glucoside metabolism, heterosis, domestication, experimental evolution sequencing, brewing, hybridization.

Abstract

S. pastorianus strains are hybrids of S. cerevisiae and S. eubayanus that have been domesticated for several centuries in lager-beer brewing environments. As sequences and structures of S. pastorianus genomes are being resolved, molecular mechanisms and evolutionary origin of several industrially relevant phenotypes remain unknown. This study investigates how maltotriose metabolism, a key feature in brewing, may have arisen in early S. eubayanus x S. cerevisiae hybrids. To address this question, we generated a near-complete genome assembly of Himalayan S. eubayanus strains of the Holarctic subclade. This group of strains have been proposed to be the origin of the S. eubayanus subgenome of current S. pastorianus strains. The Himalayan S. eubayanus genomes harbored several copies of a SeAGT1 α-oligoglucoside transporter gene with high sequence identity to genes encountered in S. pastorianus. Although Himalayan S. eubayanus strains are unable to grown on maltose and maltotriose, their maltose-hydrolase and SeMALT1 and SeAGT1 maltose-transporter genes complemented the corresponding null mutants of S. cerevisiae. Expression, in a Himalayan S. eubayanus strain, of a functional S. cerevisiae maltose-metabolism regulator gene (MALx3) enabled growth on oligoglucosides. The hypothesis that the maltotriose-positive phenotype in S. pastorianus is a result of heterosis was experimentally tested by constructing a S. cerevisiae x S. eubayanus laboratory hybrid with a complement of maltose-metabolism genes that resembles that of current S. pastorianus strains. The ability of this hybrid to consume maltotriose in brewer's wort demonstrated regulatory cross talk between sub-genomes and thereby validated this hypothesis. These results provide experimental evidence of the evolutionary origin of an essential phenotype of lager-brewing strains and valuable knowledge for industrial exploitation of laboratory-made S. pastorianus-like hybrids.

Importance

S.pastorianus, a S.cerevisiae X S.eubayanus hybrid, is used for production of lager beer, the most produced alcoholic beverage worldwide It emerged by spontaneous hybridization and have colonized

early lager-brewing processes. Despite accumulation and analysis of genome sequencing data of

S.pastorianus parental genomes, the genetic blueprint of industrially relevant phenotypes remains

unknown. Assimilation of wort abundant sugar maltotriose has been postulated to be inherited from

S.cerevisiae parent. Here, we demonstrate that although Asian *S.eubayanus* isolates harbor a

functional maltotriose transporter *SeAGT1* gene, they are unable to grow on α-oligoglucosides, but

expression of *S. cerevisae* regulator *ScMAL13* was sufficient to restore growth on trisaccharides. We

hypothesized that *S. pastorianus* maltotriose phenotype results from regulatory interaction between

S.cerevisae maltose transcription activator and the promoter of *SeAGT1*. We experimentally

confirmed the heterotic nature of the phenotype and thus this results provide experimental evidence

of the evolutionary origin of an essential phenotype of lager-brewing strains.

Introduction

Saccharomyces pastorianus is an interspecific hybrid of S. cerevisiae and S. eubayanus (1-4). S. pastorianus strains are widely used for production of lager beer, which is currently the most produced alcoholic beverage worldwide. Lager brewing requires alcoholic fermentation at relatively low temperatures. S. pastorianus was hypothesized to have emerged by spontaneous hybridization and to have colonized early lager-brewing processes due to a combination of cold-tolerance inherited from S. eubayanus and superior fermentation kinetics inherited from S. cerevisiae (5-7). Lager beer is brewed from barley wort, whose sugar composition consists, by weight, of approximately 15% glucose, 60% maltose and 25% maltotriose (8). During wort fermentation, maltotriose is generally only utilized after glucose and maltose are depleted, while its consumption is also relatively slow and often incomplete (9-11).

Complete sugar utilization is desirable for lager beer fermentation to optimize concentrations of ethanol and flavor compounds and to avoid residual sweetness (12). While S. pastorianus and S. cerevisiae strains are capable of consuming maltotriose, none of the wild isolates of S. eubayanus characterized thus far have been shown to possess this trait (6, 13, 14). These observations led to the hypothesis that the ability of S. pastorianus to ferment maltotriose was inherited from S. cerevisiae (6, 13, 15-17).

The genetic information for maltose utilization is well conserved in Saccharomyces species and depends on three gene families. MALT genes encode plasma-membrane proton symporters with varying substrate specificities and affinities (18, 19), MALS genes encode α-glucosidases that hydrolyze α-oligoglucosides into glucose, while MALR genes encode a regulator required for transcriptional induction of MALT and MALS genes by maltose (20, 21). In Saccharomyces species, maltose-utilization genes are generally organized in MAL loci. These loci contain a MALT gene (called ScMALx1 and SeMALTx in S. cerevisiae and S. eubayanus, respectively), a MALS gene referred to as ScMALx2 or SeMALSx and a MALR gene referred to as ScMALx3 or SeMALRx (13, 22). In the absence

of glucose and presence of maltose, the MalR regulator binds a bi-directional promoter, thereby

simultaneously activating expression of *MALT* and *MALS* genes (23).

The Sc*MAL1*-Sc*MAL4* and Sc*MAL6* loci of *S. cerevisiae* as well as the *SeMAL1-SeMAL4* loci of *S.*

eubayanus are located in subtelomeric regions (13, 24-26). While all *S. cerevisiae* Sc*Malx1*

transporters transport maltose, only *Sc*Mal11 is able to also transport maltotriose (9). *ScMAL11* (also

known as *ScAGT1*) shares only 57% nucleotide identity with other *ScMALx1* genes (27). The four

SeMALT (*SeMALT1-4*) genes identified in the genome of the Patagonian type strain FM1318/CBS

12357 of *S. eubayanus* were shown to encode functional maltose transporters, but none of these

genes enabled maltotriose transport (13). While no clear *ScAGT1* ortholog was found in *S. eubayanus*

CBS 12357^{T}, such an ortholog was recently found in the genomes of two North American isolates

assigned to the Holarctic subclade of *S. eubayanus* (14).

S. pastorianus inherited *MAL* genes from both *S. cerevisiae* and *S. eubayanus* (2, 4, 28). However, the

S. cerevisiae-derived maltotriose- transporter gene *ScAGT1* is truncated and, therefore, non-

functional in *S. pastorianus* (10). Instead, maltotriose consumption by *S. pastorianus* strains was

attributed to *SeAGT1* and *SpMTY1/SpMTT1* genes (29-32). In *S. pastorianus*, *SeAGT1* is located on *S.*

eubayanus CHRXV and was therefore, already before the identification of an *AGT1* ortholog in

Holarctic *S. eubayanus* strains (14), assumed to originate from *S. eubayanus* (2). *SpMTY1,* also

referred to as *SpMTT1,* is located on *S. cerevisiae* CHRVII and has less than 92% sequence identity

with other *Saccharomyces* maltose transporters (30). However, *SpMTY1* contains sequence patches

with high similarity to maltose transporters from *S. eubayanus* and *S. paradoxus* (33). Recently, two

independent laboratory evolution studies with *S. eubayanus* demonstrated that recombination of

different *SeMALT* genes yielded chimeric, neo-functionalized genes that encoded maltotriose

transporters (34, 35). *SpMTY1* may have resulted from successive introgressions of maltose-

transporter genes from *S. cerevisiae*, *S. eubayanus* and *S. paradoxus*.

Recently made *S. cerevisiae* x *S. eubayanus* laboratory hybrids showed similar lager-brewing

performance as *S. pastorianus* strains, also with respect to maltotriose utilization (6, 16, 17, 36). In

these hybrids, maltotriose consumption depended on the presence of a functional *Sc*Agt1

transporter encoded by the *S. cerevisiae* subgenome (37). However, in view of the non-functionality

of *ScAGT1* in current *S. pastorianus* strains, these laboratory hybrids did not fully recapitulate the

genetic landscape of *S. pastorianus* with respect to maltotriose fermentation (2, 6, 36).

Studies on laboratory hybrids based on *S. eubayanus* strains whose genomes are more closely related

to the *S. eubayanus* subgenome of *S. pastorianus* strains than that of the Patagonian type strain

CBS12357 might generate new insights into the evolution of maltotriose utilization in *S. pastorianus*.

To date, Himalayan *S. eubayanus* isolates show the highest sequence identity with the *S. eubayanus*

sub-genome of *S. pastorianus*, with up to 99.82% identity, as opposed to 99.56% for *S. eubayanus*

CBS12357^{T} (38).

Here, we investigated if and how the genomes of Himalayan *S. eubayanus* strains could have

contributed to maltotriose utilization in the earliest hybrid ancestors of current *S. pastorianus*

strains. To this end, we generated chromosome-level genome assemblies of these strains by long-

read DNA sequencing. Since the Himalayan strains were unable to utilize maltotriose, we functionally

characterized the assembled *MAL* genes and identified genetic determinants that prevented

maltotriose utilization. Subsequently, a laboratory hybrid of a representative Himalayan *S. eubayanus*

strain and a maltotriose-deficient ale strain of *S. cerevisiae* was generated to investigate the genetics

of maltotriose utilization in a hybrid context. We discuss the implications of the experimental results

for the proposed role and origin of *SeAGT1* in *S. pastorianus* and for the potential of hybridization to

enable maltotriose consumption in novel *Saccharomyces* hybrids.

Results

Sequencing of Himalayan *S. eubayanus* strains revealed variations of sub-telomeric regions and presence of novel putative maltose transporter genes.

It has been proposed that the *S. eubayanus* genetic pool of *S. pastorianus* was inherited from an ancestor of the Asian *S. eubayanus* lineage (38). With 99.82% identity, the Himalayan *S. eubayanus* strains CDFM21L.1 and ABFM5L.1 that belong to the Holarctic lineage (39), present the closest characterized relatives of the *S. eubayanus* ancestor of lager brewing yeasts. However, this distance was based on a limited sequencing space (38) and the analysis did not investigate the presence of specific *S. eubayanus* genetic markers found in *S. pastorianus* hybrids. Therefore, the genome of the Himalayan *S. eubayanus* strain CDFM21L.1 was sequenced with a combination of long-read and short-read techniques (Oxford Nanopore MinION and Illumina technologies, respectively) to generate a near-complete draft reference genome sequence. The resulting CDFM21L.1 genome assembly comprised 19 contigs, including its mitochondrial genome. All chromosomes were completely assembled from telomere to telomere, except for chromosome XII, which was fragmented into 3 contigs due to the repetitive rDNA region and manually assembled into a single scaffold. With a total size of 12,034,875 bp, this assembly represents the first near-complete draft genome of a *S. eubayanus* strain of the Holarctic clade (39).

Chromosome-level assemblies were hitherto only available for the Patagonia B-clade strain CBS 12357^T (1, 13). We identified three major structural differences in CDFM21L.1 relative to CBS12357^T using Mauve (40): (i) a paracentric inversion in the sub-telomeric region of chromosome VII, involving approximately 8 kbp, (ii) a translocation of approximately 12 kbp from the left sub-telomeric region of chromosome VIII to the right sub-telomeric region of chromosome VI, and (iii) a reciprocal translocation between approximately 20 kbp from the right sub-telomeric region of chromosome V and approximately 60 kbp from the center of chromosome XII (Figure 1A). All structural variation involved sub-telomeric regions, in accordance with their known relative instability (41-43).

An alignment comparison of the CDFM21L.1 and CBS 12357^T genomes with MUMmer revealed that

557 kb were unique to CDFM21L.1 and reciprocally 428 kb were unique to CBS 12357^T. Sequences

unique to CBS 12357^T (3.6% of its genome) and to CDFM21L.1 (4.6% of its genome) were located

primarily in sub-telomeric regions and in repetitive regions, such as rDNA on chromosome XII (Figure

1B). Out of the 32 sub-telomeric regions, 23 exhibited absence of synteny. Conserved synteny was

observed for sub-telomeric regions on CHRIII (left), CHRIV (left and right), CHRVI (left), CHRIX (right),

CHRXI (right), CHRXII (right), CHRXIV (left) and CHRXV (right) (Supplementary File 1).

The 428 kb of sequence that were absent in the Himalyan *S. eubayanus* strain included 99 annotated

ORFs (Supplementary File 1). Of the 99 ORFs that were (partly) affected, 11 were completely absent

in CDFM21L.1, involving mostly genes implicated in iron transport facilitation (Supplementary File 1).

The 557 kb of sequence that was not present in CBS 12357^T included 113 annotated ORFs

(Supplementary File 1). Of these 113 ORFs, 15 were completely absent in CBS 12357^T. These 15 ORFs

showed an overrepresentation of genes involved in transmembrane transport (Fishers exact test, P-

value 4.8E^{-5}) (Figure 1C).

Of the 15 ORFs unique to CDFM21L.1, three were identical orthologs of *S. cerevisiae MAL11/AGT1*.

These three ORFs were found in the sub-telomeric regions of chromosomes VII, XIV and XV. Their

sequence similarity with the *S. cerevisiae* CEN.PK113.7D and *S. pastorianus* CBS 1483 *MAL11/AGT1*

genes was 82.7% and 99.89%, respectively. In addition to these *SeAGT1* genes, CDFM21L.1 genome

sequence harbored three maltose transporters (*SeMALTx*), two maltases (*SeMALSx*) and three

regulators (*SeMALRx*) encoding genes. In contrast to the situation in *S. eubayanus* CBS 12357^T, none

of the *SeMAL* genes formed a canonical *MAL* locus in CDFM21L.1 (Figure 2). A systematic sequence

inspection of these CDFM21L.1 *SeMAL* genes revealed mutations that prematurely interrupted the

reading frames of *SeMALR1* (CHRV), *SeMALT2* (CHRXII) and *SeMALT*3 (CHRXII).

In addition to *S. eubayanus* CDFM21L.1 strain, a second Himalayan *S. eubayanus* isolate (ABFM5L.1)

was sequenced. These two strains were 99.97% genetically identical at the nucleotide level, their

MAL genes were syntenic and the premature stop codons in *SeMALR1* (CHRV), *SeMALT2* (CHRXII) and

*SeMALT*3 (CHRXII) were conserved. Two additional mutations were identified in one of the three

SeAGT1 genes. A nucleotide variation on position 53 and 939 (T instead of an A and A instead of a G))

resulted in a glycine to valine and arginine to lysine change, respectively.

Paradoxically, Himalayan *S. eubayanus* strains do not utilize maltose and maltotriose.

Identification of *SeAGT1* in the two Himalayan *S. eubayanus* strains suggests an ability to not only

grow on maltose but also on maltotriose. Strains from the Holarctic clade have previously been

hypothesized to be the donor of the *S. eubayanus* sub-genome in *S. pastorianus* hybrids (38, 39).

However, no physiological data regarding their ability to grow on the sugars present in wort are

available. To assess their growth characteristics, the *Asian S. eubayanus* strains CDFM21L.1 and

ABFM5L.1, the Patagonian *S. eubayanus* type strain CBS 12357^T and the *S. pastorianus* strain CBS

1483 were grown on diluted industrial brewer's wort at 12 ˚C. As reported previously, *S. pastorianus*

strain CBS 1483 could utilize all three sugars but did not fully consume maltotriose (Figure 3) (11).

Also in accordance with previous observations (6), CBS 12357^T consumed glucose and maltose

completely, but left maltotriose untouched. However, in marked contrast to *S. eubayanus* CBS

12357^T, neither CDFM21L.1 nor ABFM5L.1 consume maltose after growth on glucose. Moreover, like

CBS 12357^T, maltotriose was not metabolized by these two *S. eubayanus* strains. While in CBS 12357^T

an ability to grow on maltose and an inability to grow on maltotriose could be readily attributed to its

MAL genes complement, CDFM21L.1 and ABFM5L.1 failed to grow on maltose even though they

appeared to contain complete genes encoding maltose (*SeMALT1* and *SeAGT1*) and maltotriose

(*SeAGT1*) transporters.

Growth defects on maltose and maltotriose are caused by deficiency of the regulatory *Se*MalR

proteins in *S. eubayanus* CDFM21L.1.

The recent characterization of maltose metabolism in CBS 12357^T showed that the coding regions of

transcriptionally silent maltose-transporter genes in *S. eubayanus* can potentially encode functional

proteins (13). The inability of the *S. eubayanus* Himalayan isolates to grow on α-oligosaccharides

precluded direct testing of transporter-gene functionality by deletion studies. Instead, these genes

were expressed in *S. cerevisiae* IMZ616, which is devoid of all native maltose metabolism genes (44).

The CDFM21L.1 transporter genes *SeMALT1*, *SeMALT2*, *SeMALT3* or *SeAGT1* were integrated at the

ScSGA1 locus in IMZ616 along with the *S. cerevisiae* maltase gene *ScMAL12* (13), yielding a series of

strains overexpressing a single transporter (IMX1702 (*SeMALT1*), IMX1704 (*SeMALT2*), IMX1706

(*SeMALT3*) and IMX1708 (*SeAGT1*)). These strains, as well as the negative and positive control strains

IMZ616 and IMX1365 (IMZ616 expressing *ScAGT1* and *ScMAL12*), were grown on SM media

supplemented with either maltose or maltotriose. On maltose, not only the positive control strain

IMX1365, but also IMX1702 (*SeMALT1*) and IMX1708 (*SeAGT1*) were able to grow on maltose,

consuming 30 and 60%, respectively, of the initially present maltose after 100 h (Figure 4A). As

anticipated, the *SeMALT2* and *SeMALT3* alleles with premature stop codons did not support growth

on maltose. Of the two strains that grew on maltose, only IMX1708 (*SeAGT1*) also grew on

maltotriose. These results demonstrate that *SeAGT1* from a Holarctic *S. eubayanus* encoded a

functional maltotriose transporter and, consequently, that the inability of Holarctic strains to grow

on maltose and maltotriose was not caused by transporter dysfunctionality.

In addition to transport, metabolism of α-oligoglucosides requires maltase activity. Functionality of

the putative *SeMALS1* and *SeMALS2* maltase genes was tested by constitutive expression in strain

IMZ616, together with a functional *Sc*MAL31 transporter genes, yielding strains IMZ752 and IMZ753,

respectively. The maltase-negative strain IMX1313 was used as negative control. In SM medium with

maltose, both IMZ752 (*SeMALS1*) and IMZ753 (*SeMALS2*) grew and completely consumed maltose

within 65 h, demonstrating functionality of both hydrolase genes (Figure 4C).

In *S. cerevisiae* transcriptional regulation of *MALx2* and *MALx1* genes is tightly controlled by a

transcription factor encoded by *MALx3* genes. Malx3 binds an activating site located in the

bidirectional promoters that control expression of *MALx2* and *MALx1* genes (45, 46). To test whether

absence of maltose consumption in Himalayan *S. eubayanus* strains was caused by a lack

transcriptional upregulation of *SeMALT* and *SeMALS*, the *S. cerevisiae ScMAL13* gene was integrated

at the *SeSGA1* locus in *S. eubayanus* CDFM21L.1, under the control of a constitutive *ScPGK1*

promoter and *ScTEF2* terminator. *ScMAL13* expression in CDFM21L.1 enabled growth on maltose and

maltotriose (Figure 5A), indicating that a lack of transcriptional upregulation was indeed the cause of

the parental strain's inability to grow on these oligoglucosides. However, consumption of maltose

and maltotriose was incomplete and consumed sugars were almost exclusively respired, as no

ethanol was measured after 60 h of cultivation.

The possibility to grow an engineered variant of *S. eubayanus* CDFM21L.1 on α-oligoglucosides

offered an opportunity to study transporter function in its native context. Complementary functional

characterization by gene deletion of *SeMALT1* and *SeAGT1* was performed using CRISPR-Cas9

genome editing method (13, 47). Deletion of *SeMALT1* and *SeAGT1* in CDFM21L.1 resulted in strains

IMK820 and IMK823, respectively. Complete deletion of *SeAGT1* required disruption of six alleles. To

confirm the complete removal of all copies, the genome of IMK823 was sequenced. Mapping reads

onto the reference *S. eubayanus* CDFM21L.1 genome assembly confirmed that all six alleles were

removed simultaneously. Subsequently, the regulator expression cassette ($ScPGK1_p$-$ScMAL13$-

$ScTEF2_t$) was integrated in IMK820 and IMK823 at the *SeSGA1* locus yielding strains IMX1939 and

IMX1940, respectively (Figure 5B). The four deletion strains IMK820 (*SemalT1Δ*), IMK823 (*Seagt1Δ*),

IMX1939 (*SemalT1Δ Sesga1Δ::ScMAL13*) and IMX1940 (*Seagt1Δ Sesga1Δ::ScMAL13*) were

characterized on SMG, SMM or SMMt. All four strains were able to grow on glucose (Supplementary

Figure 3). While strains IMK820, IMK823 and IMX1940 were unable to grow on maltose or

maltotriose (Figure 5C), strain IMX1939 (*SemalT1Δ Sesga1Δ::SeMAL13*), which harbored functional

SeAGT1 copies, grew on maltose as well as on maltotriose. However, after 64 h of growth, these

sugars were only partially consumed. Only 1.2 g L^{-1} ethanol was produced from maltose and no

ethanol formation was observed during growth on maltotriose. The low ethanol concentration and

the relatively high OD$_{660nm}$ suggest that, under the experimental conditions strain IMX1939 exhibited

a Crabtree negative phenotype and exclusively respired maltotriose. *S. eubayanus* IMX1940 (*Seagt1Δ*

Sesga1Δ::SeMAL13) did not consume maltotriose after 84 h of incubation. Moreover, despite the

presence of *SeMALT1*, which encoded a functional maltose transporter upon expression in *S.*

cerevisiae IMZ616, strain IMX1940 was also unable to consume maltose.

In addition to a functional Malx3 transcription factor, transcriptional activation of *MAL* genes also

requires presence of a cis-regulatory motif in the promoter of regulated genes. Transcriptome

analysis of *S. eubayanus* CBS 12357^T recently showed that absence of a canonical cis-regulatory motif

in *SeMALT1* and *SeMALT3* of *S. eubayanus* CBS 12357^T caused a deficiency in their expression (13). To

further explore regulation of *SeMAL* and *SeAGT1* genes, we investigated the impact of carbon

sources on genome-wide transcriptome and, specifically, on transcriptional activation of genes

involved in maltose metabolism. Duplicate cultures of *S. eubayanus* strain IMX1765 (*ScPGK1*$_P$-

ScMAL13-ScTEF2$_t$) were grown on SMG, SMM and SMMt at 20 °C and sampled in mid-exponential

phase. After mRNA isolation and processing, cDNA libraries and reads were assembled onto the

newly annotated *S. eubayanus* CDFM21L.1 genome to calculate FPKM (fragments per kilobase of

feature (gene) per million reads mapped) expression values. The heterologous regulator *ScMAL13*,

expressed from the constitutive *ScPGK1* promoter, displayed the same expression in glucose- and

maltose-grown cultures. Although *ScMAL13* was efficiently expressed on glucose, none of the nine *S.*

eubayanus maltose genes (the three identical *SeAGT1* copies being undistinguishable) were

transcriptionally induced under these conditions (Figure 6), confirming that the hierarchical

regulatory role of glucose catabolite repression (45, 48) also takes place in *S. eubayanus*. During

growth on maltose, all nine genes were significantly upregulated relative to glucose-grown cultures

but large variations in expression level were observed. The maltase genes *SeMALS1* and *SeMALS2*

and the transporter gene *SeAGT1* showed the highest upregulation, with fold changes of 148, 161

and 2355 respectively. Although upregulated *SeMALT1* displayed a fold change of 13, its normalized

expression in maltose-grown cultures was 886-fold lower than that of *SeAGT1*. This weaker

upregulation might explain why, despite the ability of its coding region to support synthesis of a

functional maltose transporter, *SeMALT1* alone could not restore growth on maltose. The

transcriptome data also revealed that the absence of maltose induction in CDFM21L.1 was not

associated with defective cis-regulatory elements in *SeMALR* promoter sequences since the regulator

genes were properly activated, instead these results would suggest that the *Se*MalR regulator are not

functional.

294 Hybridization of two maltotriose-deficient *S. eubayanus* and *S. cerevisiae* lineages results in

295 heterosis through regulatory crosstalk.

The genetic make-up of *S. pastorianus* lager-brewing yeasts strongly advocates that they originate

from hybridization of *S. cerevisiae* and *S. eubayanus* parental lineages that were both unable to

metabolize maltotriose (2). This hypothesis is consistent with the recurrent mutation in the *S.*

cerevisiae AGT1 allele of *S. pastorianus* strains as well as with the inability of Himalayan strains of *S.*

eubayanus to grow on these oligoglucosides.

Spores of the Himalayan *S. eubayanus* CDFM21L.1 were hybridized with *S. cerevisiae* CBC-1. This top-

fermenting *S. cerevisiae* is recommended for cask and bottle conditioning and unable to consume

maltotriose (Lallemand, Montreal, Canada). Analysis of the CBC-1 assembly, obtained by a

combination of long and short read sequencing, linked its maltotriose-negative phenotype to a total

absence of the *MAL11/AGT1* gene. The resulting laboratory interspecific hybrid HTSH020 was

characterized at 12 °C on synthetic wort, a defined medium whose composition resembles that of

brewer's wort. While *S. eubayanus* CDFM21L.1 only consumed glucose and *S. cerevisiae* CBC-1

consumed glucose and maltose after 103 h (Figure 7A), the interspecific hybrid HTSH020 completely

consumed glucose, maltose and partially consumed maltotriose after 105 h, thus resembling

characteristics of *S. pastorianus* strains (e.g. CBS1483 (11)). In addition to this gain of function, the

hybrid HTSH020 outperformed both its parents on maltose consumption, since it depleted this sugar

in 70 h instead of 95 h for strain CBC-1. Since *S. cerevisiae* grows generally slower at 12 °C, the

experiments were also performed at 20 °C where HTSH020 consumed all maltose 16 h earlier than

CBC-1 (Supplementary Figure 4).

Transcriptome analysis of the hybrid strain HTSH020 grown on SM with different carbon sources

showed that *SeAGT1* expression was repressed during growth on glucose, with a normalized

expression level of 7 FPKM (Figure 7B). When grown on SM maltose, *SeAGT1, SeMALS1* and *SeMALS2*

were significantly induced, with fold increases of 816, 109 and 116, respectively (Figure 7B). Although

SeMALT1 and *SeMALT2* were induced, these transporters do not contribute to maltose metabolism

due to truncation of their ORFs. These transcriptome data demonstrated that *SeAGT1* and *SeMALS*

genes are induced by regulatory crosstalk between regulators encoded from the CBC-1 *S. cerevisiae*

sub-genome and maltotriose transporter genes harbored by the *S. eubayanus* genome. This

laboratory hybridization experiment may be the closest reproduction yet of how, centuries ago,

maltotriose-fermentation capacity arose in the first hybrid ancestor of *S. pastorianus*.

Discussion

The ability to consume maltose and maltotriose represents a key performance indicator of *S.*

pastorianus lager-brewing strains (10). This study demonstrates how mating of *S. cerevisiae* and *S.*

eubayanus strains that cannot themselves ferment maltotriose, can yield maltotriose-fermenting

hybrids. This laboratory study illustrates how, centuries ago, maltotriose-fermentation capacity may

have arisen in the first hybrid ancestor of *S. pastorianus*.

While the origin of the *S. eubayanus* parent of *S. pastorianus* strains is still under debate (49-

51), phylogenetic analysis suggested a Far East Asian origin (38). However, this interpretation was

based on a limited sequencing space and was constrained by the quality of available sequence

assemblies. Since an ortholog of *SeAGT1* had previously only been found in the *S. eubayanus* sub-

genome of *S. pastorianus* strains, this finding revived the discussion on the geographical origin of the

ancestral *S. eubayanus* parent (14). The high-quality, annotated genome assemblies of the Himalayan

S. eubayanus strains CDFM21L.1 and ABFM5L.1 presented in the present study revealed several

copies of *SeAGT1*, whose very high sequence identity with *S. pastorianus SeAGT1* are consistent with

the previously proposed Asian origin of the S. *eubayanus* sub-genome of *S. pastorianus* (14, 38, 39).

Next, genome-sequence comparison of the Patagonian B sub-clade *S. eubayanus* strain CBS12357[T]

and the Holarctic sub-clade strain CDFM21L.1 revealed homoplasy of *SeAGT1*, probably reflecting

that these sub-clades evolved in different ecological niches.

The *S. eubayanus* wild stock whose genome sequence most closely corresponds to the *S.*

eubayanus sub-genome of *S. pastorianus* originates from the Tibetan plateau of the Himalaya (38).

However, the first *S. cerevisiae x S. eubayanus* hybrid, from which current lager yeasts evolved by

centuries of domestication, likely originates from a region between Bavaria and Bohemia in Central

Europe. So far, European *S. eubayanus* isolates have not been reported. This may indicate that the

original hybridization event occurred elsewhere or that the ancestral European lineage went extinct.

The recent detection, in a metagenomics analysis of samples from the Italian Alps, of ITS1 sequences

corresponding to *S. eubayanus* could indicate that a wild European lineage exists after all (52).

Functional characterization by heterologous complementation of an *S. cerevisiae* mutant

strain established that the *Se*Agt1 transporters from the Himalayan *S. eubayanus* strains CDFM21L.1

and ABFM5L supported uptake of maltose and maltotriose. After showing that these strains also

encoded a functional maltase gene, their inability to grow on maltose and maltotriose was attributed

to an inability to transcriptionally upregulate maltose metabolism genes likely caused by loss of

function mutations the regulator. In *S. cerevisiae* and, to some extent, ,in *S. eubayanus* strains of the

Patagonian B sub-clade such as CBS12357[T] (13, 26, 53), MAL loci exhibit a specific organization in

which a transporter (*MALT*) and a hydrolase (*MALS*) gene are expressed from the same bidirectional

promoter and are located adjacent to a regulator gene (*MALR*) (21). In contrast, of the seven

genomic regions harboring *MAL* genes in the two Asian *S. eubayanus* strains, none showed this

canonical organization (Figure 2) and the subtelomeric regions carrying *SeAGT1* did not harbour

sequences similar to hydrolase or regulator genes. Subtelomeric regions harboring the other *MAL*

genes indicated intensive reorganization as a result of recombination. In particular, subtelomeric

regions on CHRII, CHRV and CHRXII provide clear indications for recombination events that scattered

genes from ancestral MAL1 and MAL2 loci over several chromosomes. A similar interpretation could

explain the reorganization MAL3 on CHRXIII (Figure 2). Similar events may have contributed to loss of

function of the MAL regulators (MalR), as exemplified by the occurrence of a non-synonymous

mutation in *SeMALR1* resulting in loss of function. These rearrangements did, however, not

inactivate the *cis*-regulatory sequences of the *MAL* genes, since complementation with a functional

ScMAL13 allele caused induction of most *SeMAL* genes (Figure 6, Figure 7B) and, thereby, the

heterotic maltotriose-positive phenotype of the hybrid strain HTSH020. Together with the high copy

number of *SeAGT1*, this heterotic complementation may have been the main driver for colonization

of low-temperature brewing processes by the early hybrid ancestors of current *S. pastorianus* strains.

Recent work on adaptation to brewing environments of laboratory *S. cerevisiae* × *S. eubayanus*

hybrids showed loss maltotriose utilization during serial transfer in wort (37). A similar loss of

maltotriose utilization is frequently encountered in *S. cerevisiae* ale strains (54), as well as in some

Saaz-type *S. pastorianus* strains (55). This is thus in contrast with retention of a maltotriose

assimilation phenotype by Frohberg-type *S. pastorianus* strains. This may have been facilitated by the

occurrence of multiple copies of the *SeAGT1* gene in the *S. eubayanus* ancestor, which could act as a

sequence buffer to counteracting adverse effects of gene copy loss. The recent release of the first

long-read sequencing assembly of *S. pastorianus* enabled a precise chromosomal mapping of the

maltose-metabolism genes (56) and showed that the Frohberg type *S. pastorianus* strains CBS 1483

harbored one copy of *SeAGT1* on the *S. eubayanus* CHRXV section (as in CDFM21L.1) of the chimeric

chromosome formed from *Se*CHRXV and *Se*CHRVIII (56).

Differential retention and loss of maltotriose consumption in *S. pastorianus* lineages may reflect

different brewing process conditions during domestication. In modern brewing processes based on

high-gravity wort, cell division is largely constrained to the glucose and maltose phases, which occur

before depletion of nitrogen sources (57). It may be envisaged that, in early lager-brewing processes,

unstandardized mashing processes generated wort with a higher maltotriose content, which would

have allowed for continued yeast growth during the maltotriose consumption phase. During serial

transfer on sugar mixtures, the selective advantage of consuming a specific sugar from a mixture

correlates with the number of generations on that sugar during each cycle (58, 59). Such conditions

would therefore have conferred a significant selective advantage to a maltotriose-assimilating *S.*

cerevisiae x *S. eubayanus* hybrid, especially if, similar to current ale yeasts, the *S. cerevisiae* parent

was unable to ferment maltotriose.

The heterotic phenotype that was reconstructed in the interspecies *S. cerevisiae* x *S.*

eubayanus hybrid HTS020 resulted from combination of dominant and recessive genetic variations

from both parental genomes. *S. eubayanus* contributed the *SeAGT1* gene and its functional *cis*-

regulatory sequences, but also harbored recessive mutations in *MALR* genes that allowed full

expression of the heterotic phenotype. These mutations were complemented with a set of *S.*

cerevisiae genes including a functional *MALR* and a non-functional *ScAGT1* gene to match the

mutations found in *S. pastorianus* (2). Although some *S. pastorianus* strains harbor an additional

maltotriose transporter encoded by *SpMTT1 (30)*, this gene was recently proposed to have emerged

after the original hybridization event as a result of repeated recombination between *MALT* genes

from both sub-genomes (33).

Maltotriose fermentation is likely not the only heterotic phenotype of S. *pastorianus* strains.

Flocculation or formation of complex aroma profiles (28, 60) are phenotypes that are not fully

understood and difficult to reproduce, that also might result from heterosis (37).

Laboratory-made *S. cerevisiae* x *S. eubayanus* hybrids hold great potential for brewing process

intensification and for increasing product diversity. In addition to increasing our understanding of the

evolutionary history of lager yeast genomes evolutionary, this study has implications for the design

of new hybrids. Hitherto, laboratory crosses of *S. cerevisiae* x *S. eubayanus* strains were designed

based on combination of dominant traits of the parental strains. Our results show that recessive

traits can be just as important as contributors to the genetic diversity of such hybrids.

Materials and methods

Strains and maintenance

All strains used in this study are listed in Table 1. Stock cultures of *S. eubayanus* and *S. cerevisiae* strains were grown in YPD (10 g L^{-1} yeast extract, 20 g L^{-1} peptone and 20 g L^{-1} glucose) until late exponential phase, complemented with sterile glycerol to a final concentration of 30 % (v/v) and stored at -80 °C as 1 mL aliquots until further use.

Media and cultivation

S. eubayanus batch cultures were grown on synthetic medium (SM) containing 3.0 g L^{-1} KH_2PO_4, 5.0 g L^{-1} $(NH_4)_2SO_4$, 0.5 g L^{-1} $MgSO_4.7H_2O$, 1 mL L^{-1} trace element solution, and 1 mL L^{-1} vitamin solution (61). The pH was set to 6.0 with 2 M KOH prior to autoclaving at 120 °C for 20 min. Vitamin solutions were sterilized by filtration and added to the sterile medium. Concentrated sugar solutions were autoclaved at 110 °C for 20 min or filter sterilized and added to the sterile flasks to give a final concentration of 20 g L^{-1} glucose (SMG), maltose (SMM) or maltotriose (SMMt). With the exception of IMZ752 and IMZ753, *S. cerevisiae* batch cultures were grown on SM supplemented with 150 mg L^{-1} uracil (62) to compensate for loss of plasmid pUDC156 that carried the *Spcas9* endonuclease gene, and supplemented with 20 g L^{-1} glucose (SM$_u$G), maltose (SM$_u$M) or maltotriose (SM$_u$Mt). All batch cultures were grown in 250 mL shake flasks with a working volume of 50 mL. The cultures were inoculated at an initial OD_{660nm} of 0.1 and incubated under an air atmosphere and shaken at 200 rpm and at 20 °C in a New Brunswick™ Innova44 incubator (Eppendorf Nederland B.V, Nijmegen, The Netherlands).

S. eubayanus strains transformed with plasmids pUDP052 (gRNA$_{SeSGA1}$), pUDP091 (gRNA$_{SeMALT1}$) and pUDP090 (gRNA$_{SeAGT1}$) were selected on modified SMG medium (SM$_{Ace}$G) in which $(NH_4)_2SO_4$ was replaced by 6.6 g L^{-1} K_2SO_4 and 10 mM acetamide (63). SM-based solid media contained 2 % Bacto Agar (BD Biosciences, Franklin Lakes, NJ). *S. cerevisiae* strains expressing either *SeMALT, SeMALS* or *ScMALR* were selected on SM$_{Ace}$G. For plasmid propagation, *E. coli* XL1-Blue-derived strains (Agilent Technologies, Santa Clara, CA) were grown in Lysogeny Broth medium (LB, 10 g L^{-1} tryptone, 5 g L^{-1}

yeast extract, 5 g L^{-1} NaCl) supplied with 100 mg L^{-1} ampicillin. Synthetic wort medium (SWM) for

growth studies contained 14.4 g·L^{-1} glucose, 2.3 g·L^{-1} fructose, 85.9 g·L^{-1} maltose, 26.8 g·L^{-1}

maltotriose, 5 g·L^{-1} (NH$_4$)$_2$SO$_4$, 3 g·L^{-1} KH$_2$PO$_4$, 0.5 g·L^{-1} MgSO$_4$.7H$_2$O, 1 mL·L^{-1} trace element solution, 1

mL·L^{-1} vitamin solution, supplemented with the anaerobic growth factors ergosterol and Tween 80

(0.01 g·L^{-1} and 0.42 g·L^{-1} respectively), as previously described (61).

Industrial wort (containing 14.4 g L^{-1} glucose, 85.9 g L^{-1} maltose, 26.8 g L^{-1} maltotriose, 2.3 g L^{-1}

fructose and 269 mg L^{-1} FAN) was provided by Heineken Supply Chain B.V. (Zoeterwoude, the

Netherlands). The wort was supplemented with 1.5 mg L^{-1} of Zn^{2+} by addition of ZnSO$_4$.7H$_2$O,

autoclaved for 30 min at 121 ℃, filtered using Nalgene 0.2 µm SFCA bottle top filters (Thermo

Scientific) and diluted with sterile demineralized water. Sporulation medium consisted of 2 % (w/v)

KAc in MilliQ water set to pH 7.0 with KOH, autoclaved at 121 °C for 20 min.

Microaerobic growth experiments

Microaerobic cultures were grown in 250-mL airlock-capped Neubor infusion bottles (38 mm neck,

Dijkstra, Lelystad, Netherlands) containing 200 mL three-fold diluted industrial wort supplemented

with 0.4 mL L^{-1} Pluronic antifoam (Sigma-Aldrich, St. Louis, MO). Bottle caps were equipped with a 0.5

mm x 16 mm Microlance needle (BD Biosciences) sealed with cotton to prevent pressure build-up.

Sampling was performed aseptically with 3.5 mL syringes using a 0.8 mm x 50 mm Microlance needle

(BD Biosciences). Microaerobic cultures were inoculated at an OD$_{660nm}$ of 0.1 from stationary-phase

precultures in 50 mL Bio-One Cellstar Cellreactor tubes (Sigma-Aldrich) containing 30 mL of the same

medium, grown for 4 days at 12 °C. Bottles were incubated at 12 °C and shaken at 200 rpm in a New

Brunswick Innova43/43R shaker (Eppendorf Nederland B.V.). At regular intervals, 3.5 mL samples

were collected in 24 deep-well plates (EnzyScreen BV, Heemstede, Netherlands) using a LiHa liquid

handler (Tecan, Männedorf, Switzerland) to measure OD$_{660nm}$ and external metabolites. 30 µL of each

sample was diluted 5-fold in demineralized water in a 96 well plate and OD$_{660nm}$ was measured with a

Magellan Infinite 200 PRO spectrophotometer (Tecan). From the remaining sample, 150 µL was

vacuum filter sterilized using 0.2 μm Multiscreen filter plates (Merck, Darmstadt, Germany) for HPLC

measurements.

Analytical methods

Optical densities of yeast cultures were measured with a Libra S11 spectrophotometer (Biochrom,

Cambridge, United Kingdom) at a wavelength of 660 nm. Biomass dry weight was measured by

filtering 10-mL culture samples over pre-weighed nitrocellulose filters with a pore size of 0.45 μm.

Filters were washed with 10 mL water, dried in a microwave oven (20 min at 350 W) and reweighed.

Sugars were measured using a high pressure liquid chromatography Agilent Infinity 1260 series

(Agilent Technologies) using a Bio-Rad Aminex HPX-87H column at 65 °C with 5 mM sulfuric acid at a

flow rate of 0.8 mL min^{-1}. Compounds were measured using a RID at 35 °C. Samples were centrifuged

at 13,000 g for 5 min to collect supernatant or 0.2 μm filter-sterilized before analysis.

Plasmid construction

Plasmids used and constructed in this study are listed in Table 2, oligonucleotide primers used in this

study are listed in Supplementary Table 1. Coding regions of *SeMALT1*, *SeMALT2*, *SeMALT3* and

SeAGT1 were amplified from CDFM21L.1 genomic DNA with Phusion High-Fidelity DNA polymerase

(Thermo Scientific), according to the supplier's instructions with primers pairs 12355/12356,

12357/12358, 12359/12360 and 12361/12362, respectively. The coding sequence of *ScMAL31* was

amplified from CEN.PK113-7D genomic DNA with Phusion High-Fidelity DNA polymerase (Thermo

Scientific), according to the supplier's instructions with primer pairs 9942/9943. Each primer carried a

40 bp extension complementary to the plasmid backbone of p426-TEF-amdS (64), which was PCR

amplified using Phusion High-Fidelity DNA polymerase (Thermo Scientific) and primers 7812/5921.

Each transporter fragment was assembled with the p426-TEF-amdS backbone fragment using

NEBuilder HiFi DNA Assembly (New England Biolabs, Ipswich, MA), resulting in plasmids pUD444

(*ScMAL31*), pUD794 (*SeMALT1*), pUD795 (*SeMALT2*), pUD796 (*SeMALT3*) and pUD797 (*SeAGT1*). All

plasmids were verified for correct assembly by Sanger sequencing (Baseclear, Leiden, The

Netherlands).

SeMALS1 and *SeMALS2* were amplified from CDFM21L.1 genomic DNA with Phusion High-Fidelity

DNA polymerase (Thermo Scientific), with primers pairs 14451/14453 and 14452/14453,

respectively. Each primer pair carried a 30 bp extension complimentary to the plasmid backbone of

pUDE044 (65) which was PCR amplified using Phusion High-Fidelity DNA polymerase (Thermo

Scientific) and primers 14449/14450. Resulting amplicons were assembled using NEBuilder HiFi DNA

Assembly (New England Biolabs), resulting in plasmids pUDE843 (*SeMALS1*) and pUDE844 (*SeMALS2*)

that were verified by Sanger sequencing (Baseclear).

S. cerevisiae ScMAL13, the *ScPGK1* promoter and the *ScTEF2* terminator were amplified from

CEN.PK113-7D genomic DNA with Phusion High-Fidelity DNA polymerase (Thermo Scientific), with

primer pairs 12915/12916, 9421/9422 and 10884/10885, respectively. Fragments were gel purified

and used with pYTK074, pYTK082 and pYTK083 in Golden Gate assembly according to the yeast

toolkit protocol (66) resulting in pUDE780, which was verified by Sanger sequencing (Baseclear).

Guide-RNA (gRNA) sequences for deletion of *SeMALT1* and *SeAGT1* in CDFM21L.1 were designed as

described previously (47). The DNA sequences encoding these gRNAs were synthesized at GeneArt

(Thermo Scientific) and were delivered in pUD631 and pUD634, respectively. The gRNA spacer

sequences (Se*MALT1* 5' CCCCGATATTCTTTACACTA 3', Se*AGT1* 5'- AGCTTTGCGAAAATATCCAA -3') and

the structural gRNA sequence were flanked at their 5' ends by the Hammerhead ribozyme (HH) and

at their 3' ends by the Hepatitis Delta Virus ribozyme (HDV) (67). The HH-gRNA-HDV fragment was

flanked on both ends with a BsaI site for further cloning (47, 68). Plasmids pUDP091 (gRNA$_{SeMALT1}$) and

pUDP090 (gRNA$_{SeAGT1}$) were constructed by Golden Gate cloning by digesting pUDP004 and the gRNA-

carrying plasmid (pUD631 and pUD634, respectively) using BsaI and ligating with T4 ligase (69).

Correct assembly was verified by restriction analysis with PdmI (Thermo Scientific) and Sanger

sequencing (Baseclear).

 Strain construction

S. cerevisiae IMZ616, which cannot grow on α-glucosides (44), was used as a host to test functionality

of individual *S. eubayanus* (putative) maltose transporter genes (13). *S. cerevisiae* IMX1702 was

constructed by integrating *ScTDH3$_p$-ScMAL12-ScADH1$_t$* and *ScTEF1$_p$-SeMALT1-ScCYC1$_t$* at the *ScSGA1*

locus of strain IMZ616. A fragment containing the *ScTDH3$_p$-ScMAL12-ScADH1$_t$* transcriptional unit

was PCR amplified using Phusion High-Fidelity DNA polymerase (Thermo Scientific) from pUDE044

with primers 9596/9355, which included a 5′ extension homologous to the upstream region of the

ScSGA1 locus and an extension homologous to the co-transformed transporter fragment,

respectively. The DNA fragment carrying the *S. eubayanus SeMALT1* maltose symporter (*ScTEF1$_p$-*

SeMALT1-ScCYC1$_t$) was PCR amplified from pUD794 using primers 9036/9039, which included a 5′

extension homologous to the co-transformed transporter fragment and an extension homologous to

the downstream region of the *ScSGA1* locus, respectively. To facilitate integration in strain IMZ616,

the two PCR fragments were co-transformed with plasmid pUDR119 (*amdS*), which expressed a gRNA

targeting *ScSGA1* (spacer sequence: 5′-ATTGACCACTGGAATTCTTC-3′) (70). The plasmid and repair

fragments were transformed using the LiAc yeast transformation protocol (71) and transformed cells

were plated on SM$_{Ace}$G. Correct integration was verified by diagnostic PCR with primers pairs

4226/4224. Strains *S. cerevisiae* IMX1704, IMX1706 and IMX1708 were constructed following the

same principle, but instead of using pUD794 to generate the transporter fragment, pUD795, pUD796

and pUD797 were used to PCR amplify *ScTEF1$_p$-SeMALT2-ScCYC1$_t$*, *ScTEF1$_p$-SeMALT3-ScCYC1$_t$* and

ScTEF1$_p$-SeAGT1-ScCYC1$_t$ respectively. IMX1313 was constructed in a similar way using only *ScTEF1$_p$-*

ScMAL31-ScCYC1$_t$ amplified with primer pair 9036/11018 which contain 5′and 3′ extensions

homologous to the upstream and downstream region of the *ScSGA1* locus. Correct integration was

verified by diagnostic PCR with primer pair 4226/4224 (Supplementary Figure 1). All PCR-amplified

gene sequences were Sanger sequenced (BaseClear). IMX1313 was grown on YPD to loose pUDR119

(*URA3*) and pUDC156 (*amdS*). An isolate unable to grown on SMG without uracil and with acetamide

was selected and named IMX1313Δ. This strain was able to grow on SMG supplemented with 150 mg

L^{-1} uracil.

To assess functionality of CDFM21L.1 *SeMALS1*, IMX1313Δ was transformed with 100 ng pUDE843

(*ScTDH3$_p$-SeMALS1-ScADH1$_t$*) by electroporation (47), resulting in strain IMZ752. Transformants were

selected on SMG plates after 5 days of incubation at 20 °C and validated by PCR (DreamTaq

polymerase, Thermo Scientific) using primer pair 14454/14455 (Supplementary Figure 1). Similarly,

functionality of the *SeMALS2* maltase gene of CDFM21L.1 was assessed by transforming IMX1313Δ

with pUDE844 (*ScTDH3$_{pr}$-SeMALS2-ScADH1$_{ter}$*), resulting in strain IMZ753.

S. eubayanus IMK820 (*SemalT1Δ*) was constructed by transforming CDFM21L.1 with 200 ng of

pUDP091 and 1 μg of 120 bp repair fragment obtained by mixing an equimolar amount of primers

12442/12443, as previously described (47). As a control, the same transformation was performed

without including the repair DNA fragment. Transformants were selected on SM$_{Ace}$G plates. *S.*

eubayanus IMK823 (*Seagt1Δ*) was constructed similarly, using pUDP090 and primer pair

11320/11321. Deletion of *SemalT1* was verified by PCR with primer pair 11671/11672 and Sanger

sequencing. The *Seagt1* deletion was verified by PCR using primer pair 12273/12274, and by Illumina

whole-genome sequencing and read alignment to the reference genome of CDFM21L.1 (Bioproject

accession number PRJNA528469).

Strains IMX1765, IMX1939 and IMX1940 were constructed by inserting *ScPGK1$_p$-ScMAL13-ScTEF2$_t$* at

the *SeSGA1* locus of CDFM21L.1, IMK820 and IMK823, respectively. A repair fragment containing

ScPGK1$_p$-ScMAL13-ScTEF2$_t$ was amplified from pUDE780 with primer pair 12917/12918. Strains

CDFML21L.1, IMK820 and IMK823 were transformed by electroporation by addition of 350 ng of

repair fragment and 560 ng pUDP052 (*amdS*) into the cells as previously described (47).

Transformants were plated on SM$_{Ace}$G and incubated at 20 °C. IMX1762 was constructed similarly

using a repair fragment with *ScTDH3$_p$-ScMAL12-ScADH1$_t$* amplified from pUDE044 with primer pair

12319/12320. Strains were verified by PCR using primer pair 12635/12636 and Sanger sequencing.

Hybrid Construction

The *S. cerevisiae* X *S. eubayanus* hybrid HTSH020 was constructed by spore-to-spore mating . The *S.*

eubayanus strain CDFM21L.1 and the *S. cerevisiae* strain CBC-1 were grown in 20 mL YPD at 20 °C

until late exponential phase. Cells were centrifuged for 5 min at 1000 g and washed twice in

demineralized water. Cells were re-suspended in 20 mL sporulation medium and incubated for 64 h

at 20 °C. Presence of spores was verified by microscopy. Asci were harvested by centrifugation for 5

min at 1000 g, and washed with demineralized water, resuspended in 100 μL demineralized water

containing 100 U/mL of Zymolyase (MP Bio, Santa Ana, CA) and incubated for 10 min at 30 °C. Spores

were washed and plated on the edge of a YPD agar plate. Spores from the two strains were brought

in contact with each other with an MSM System 400 micromanipulator (Singer Instruments, Watchet,

United Kingdom). Zygote formation was observed after 6-8 h. Emerging colonies were re-streaked

twice on SM 2% maltose at 12 °C. Successful hybridization was verified by multiplex PCR using

DreamTaq DNA polymerase (Thermo Scientific), by amplifying the *S. cerevisiae* specific *MEX67* gene

with primer pairs 8570/8571 and by amplifying the *S. eubayanus* specific gene *SeFSY1* with primers

8572/8573 (Supplementary Figure 2), as previously described (72).

Illumina sequencing

Genomic DNA of *S. eubayanus* strains CDFM21L.1 and ABFM5L.1, *S. cerevisiae* strain CBC-1 and *S.*

cerevisiae x *S. eubayanus* strain HTSH020 was isolated as previously described (4). Paired-end

sequencing (2 X 150 bp) was performed on a 350 bp PCR-free insert library using Illumina HiSeq2500

(San Diego, CA) by Novogene(HK) Company Ltd (Hong Kong, China). Genomic DNA of the strains CBC-

1 and HTSH020 was sequenced in house on a MiSeq sequencer (Illumina) with 300 bp paired-end

reads using PCR-free library preparation. Sequence data are available at NCBI under Bioproject

accession number PRJNA528469.

MinION long read sequencing

For long-read sequencing, a 1D sequencing library (SQK-LSK108) was prepared for CDFM21L.1 and

CBC-1 and loaded onto an FLO-MIN106 (R9.4) flow cell, connected to a MinION Mk1B unit (Oxford

Nanopore Technology, Oxford, United Kingdom), according to the manufacturer's instructions.

MinKNOW software (version 1.5.12; Oxford Nanopore Technology) was used for quality control of

active pores and for sequencing. Raw files generated by MinKNOW were base-called using Albacore

(version 1.1.0; Oxford Nanopore Technology). Reads with a minimum length of 1000 bp were

extracted in fastq format. For CDFM21L.1, 3.26 Gb of sequence with an average read length of 8.07

kb was obtained and for CBC-1 3.04 Gb sequence with an average read length of 7.27 kb. Sequencing

data are available at NCBI under Bioproject accession number PRJNA528469.

De novo *assembly*

De novo assembly of the Oxford Nanopore MinION dataset was performed using Canu (v1.4, setting:

genomesize=12m) (73). Assembly correctness was assessed using Pilon (74) and further corrected by

"polishing" of sequencing/assembly errors by aligning Illumina reads with BWA (75) using correction

of only SNPs and short indels (–fix bases parameter). For HTSH020, an artificial reference genome

was made by combining the assembly of CBC-1 and CDFM21L.1. The genome assemblies were

annotated using the MAKER2 annotation pipeline (version 2.31.9) (76), using SNAP (version 2013–11-

29) (77) and Augustus (version 3.2.3) (78) as *ab initio* gene predictors. *S. cerevisiae* S288C EST and

protein sequences were obtained from SGD (*Saccharomyces* Genome Database,

http://www.yeastgenome.org/) and were aligned using BLASTX on the obtained polished sequence

assembly (BLAST version 2.2.28+) (79). Predicted translated protein sequences of the final gene

model were aligned to the *S. cerevisiae* S288C protein Swiss-Prot database using BLASTP

(http://www.uniprot.org/). Custom-made Perl scripts were used to map systematic names to the

annotated gene names (Supplementary Table 2). Error rates in nanopore-sequencing data were

estimated from the q score (Phred scaled) per read, as calculated by the base caller Albacore (version

1.1.0) (Oxford Nanopore Technology). Average q score was used to calculate the error $P = 10^{q/10}$.

RNA isolation

CDFM21L.1, IMX1765, IMX1939 and HTS020 were grown in SMG, SMM and SMMt until mid-

exponential phase (OD_{660nm} of 12 for SMG/SMM and of OD_{660nm} 15 for SMMt). Culture samples

corresponding to ca. 200 mg of biomass dry weight were directly quenched in liquid nitrogen. The

samples were processed and total RNA extracted as previously described (80). Prior to cDNA

synthesis, purity, concentration and integrity of the RNA in the samples was assessed with Nanodrop

(Thermo Scientific), Qubit (Thermo Scientific) and Tapestation 220 with RNA Screen Tape (Agilent

Technologies), respectively, according the manufacturers' recommendations. cDNA libraries were

prepared using the TruSeq RNA V2 kit (Illumina). Paired-end sequencing (2 X 150 bp) was performed

on a 300 bp PCR-free insert library on a HISeq 2500 (Illumina) at Novogene (HK) Company Ltd (Hong

Kong, China). Duplicate biological samples were processed, generating an average sequence quantity

of 23.7M reads per sample. Reads were aligned to the CDFM21L.1 reference assembly (GEO

(https://www.ncbi.nlm.nih.gov/geo/) accession number GSE133146) using a two-pass STAR (81)

procedure. In the first pass, splice junctions were assembled and used to inform the second round of

alignments. Introns between 15 and 4000 bp were allowed, and soft clipping was disabled to prevent

low-quality reads from being spuriously aligned. Ambiguously-mapped reads were removed from the

dataset. Expression level for each transcript were quantified using htseq-count (82) in union mode.

Fragments per kilo-base of feature (gene) per million reads mapped (FPKM) values were calculated

by "Applying the fpkm method" from the edgeR package (83, 84). Differential expression analysis was

performed using DESeq (85).

Accession Numbers

The sequencing data were deposited at NCBI (https://www.ncbi.nlm.nih.gov/) under the Bioproject PRJNA528469 and the transcriptomics data were deposited at GEO (Genome Expression Omnibus https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE133146.

Supplementary Data

Supplementary Data are available

Authors contributions

JMD conceived the study and designed the experiments. NB, AB, LvbE, ARGdV, SMW and JD performed the experimental work. NB and MvdB performed bioinformatics analysis. NB, AB, FYB, JTP and JMGD supervised the study and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Flip de Groot for his technical assistance in gene editing in *S. eubayanus*, Niels G.A. Kuijpers, Viktor M. Boer and Jan-Maarten Geertman for their critical input.

Funding

This work was supported by the BE-Basic R&D Program (http://www.be-basic.org/), which was granted an FES subsidy from the Dutch Ministry of Economic Affairs, Agriculture and Innovation (EL&I).

Conflict of Interest

The authors declare no conflict of interest.

References

1. **Libkind D, Hittinger CT, Valério E, Gonçalves C, Dover J, Johnston M, Gonçalves P, Sampaio JP.** 2011. Microbe domestication and the identification of the wild genetic stock of lager-brewing yeast. Proc Natl Acad Sci U S A:201105430.

2. **Nakao Y, Kanamori T, Itoh T, Kodama Y, Rainieri S, Nakamura N, Shimonaga T, Hattori M, Ashikari T.** 2009. Genome sequence of the lager brewing yeast, an interspecies hybrid. DNA Res **16**:115-129.

3. **Walther A, Hesselbart A, Wendland J.** 2014. Genome sequence of *Saccharomyces carlsbergensis*, the world's first pure culture lager yeast. G3 (Bethesda) **4**:783-793.

4. **Van den Broek M, Bolat I, Nijkamp JF, Ramos E, Luttik MAH, Koopman F, Geertman JM, De Ridder D, Pronk JT, Daran J-MG.** 2015. Chromosomal copy number variation in *Saccharomyces pastorianus* evidence for extensive genome dynamics in industrial lager brewing strains. Appl Environ Microbiol:AEM. 01263-01215.

5. **Krogerus K, Magalhães F, Vidgren V, Gibson B.** 2015. New lager yeast strains generated by interspecific hybridization. J Ind Microbiol Biotechnol **42**:769-778.

6. **Hebly M, Brickwedde A, Bolat I, Driessen MR, de Hulster EA, van den Broek M, Pronk JT, Geertman J-M, Daran J-MG, Daran-Lapujade P.** 2015. *S. cerevisiae × S. eubayanus* interspecific hybrid, the best of both worlds and beyond. FEMS Yeast Res **15**.

7. **Krogerus K, Magalhães F, Vidgren V, Gibson B.** 2017. Novel brewing yeast hybrids: creation and application. Appl Environ Microbiol **101**:65-78.

8. **Zastrow C, Hollatz C, De Araujo P, Stambuk B.** 2001. Maltotriose fermentation by *Saccharomyces cerevisiae*. J Ind Microbiol Biotechnol **27**:34-38.

9. **Alves SL, Herberts RA, Hollatz C, Trichez D, Miletti LC, De Araujo PS, Stambuk BU.** 2008. Molecular analysis of maltotriose active transport and fermentation by *Saccharomyces cerevisiae* reveals a determinant role for the AGT1 permease. Appl Environ Microbiol **74**:1494-1501.

10. **Vidgren V, Huuskonen A, Virtanen H, Ruohonen L, Londesborough J.** 2009. Improved fermentation performance of a lager yeast after repair of its *AGT1* maltose and maltotriose transporter genes. Appl Environ Microbiol **75**:2333-2345.

11. **Brickwedde A, van den Broek M, Geertman J-MA, Magalhães F, Kuijpers NG, Gibson B, Pronk JT, Daran J-MG.** 2017. Evolutionary engineering in chemostat cultures for improved maltotriose fermentation kinetics in *Saccharomyces pastorianus* lager brewing yeast. Front Microbiol **8**:1690.

12. **Zheng X, D'Amore T, Russell I, Stewart G.** 1994. Factors influencing maltotriose utilization during brewery wort fermentations. J Am Soc Brew Chem **52**:41-47.

13. **Brickwedde A, Brouwers N, van den Broek M, Gallego Murillo JS, Fraiture JL, Pronk JT, Daran J-MG.** 2018. Structural, physiological and regulatory analysis of maltose transporter genes in *Saccharomyces eubayanus* CBS 12357T. Front Microbiol **9**:1786.

14. **Baker EP, Hittinger CT.** 2018. Evolution of a novel chimeric maltotriose transporter in *Saccharomyces eubayanus* from parent proteins unable to perform this function. bioRxiv doi:10.1101/431171.

15. **Krogerus K, Magalhães F, Vidgren V, Gibson B.** 2017. Novel brewing yeast hybrids: creation and application. Appl Microbiol Biotechnol **101**:65-78.

16. **Krogerus K, Magalhães F, Vidgren V, Gibson B.** 2015. New lager yeast strains generated by interspecific hybridization. Journal of industrial microbiology & biotechnology **42**:769-778.

17.   **Nikulin J, Krogerus K, Gibson B.** 2018. Alternative Saccharomyces interspecies hybrid
combinations and their potential for low-temperature wort fermentation. Yeast (Chichester,
England) **35**:113-127.
18.   **Cheng Q, Michels CA.** 1991. *MAL11* and *MAL61* encode the inducible high-affinity maltose
transporter of *Saccharomyces cerevisiae*. J Bacteriol **173**:1817-1820.
19.   **Vidgren V, Multanen JP, Ruohonen L, Londesborough J.** 2010. The temperature dependence
of maltose transport in ale and lager strains of brewer's yeast. FEMS Yeast Res **10**:402-411.
20.   **Naumov GI, Naumova ES, Michels C.** 1994. Genetic variation of the repeated *MAL* loci in
natural populations of *Saccharomyces cerevisiae* and *Saccharomyces paradoxus*. Genetics
136:803-812.
21.   **Charron MJ, Read E, Haut SR, Michels CA.** 1989. Molecular evolution of the telomere-
associated *MAL* loci of *Saccharomyces*. Genetics **122**:307-316.
22.   **Hayford A, Jespersen L.** 1999. Characterization of Saccharomyces cerevisiae strains from
spontaneously fermented maize dough by profiles of assimilation, chromosome
polymorphism, PCR and MAL genotyping. Journal of applied microbiology **86**:284-294.
23.   **Bell PJ, Higgins VJ, Dawes IW, Bissinger PH.** 1997. Tandemly repeated 147 bp elements
cause structural and functional variation in divergent MAL promoters of Saccharomyces
cerevisiae. Yeast **13**:1135-1144.
24.   **Barnett JA.** 1976. The Utilization of Sugars by Yeasts1, p 125-234, Advances in carbohydrate
chemistry and biochemistry, vol 32. Elsevier.
25.   **Salazar AN, Gorter de Vries AR, van den Broek M, Wijsman M, de la Torre Cortés P,**
Brickwedde A, Brouwers N, Daran J-MG, Abeel T. 2017. Nanopore sequencing enables near-
complete de novo assembly of *Saccharomyces cerevisiae* reference strain CEN. PK113-7D.
FEMS Yeast Res **17**.
26.   **Baker E, Wang B, Bellora N, Peris D, Hulfachor AB, Koshalek JA, Adams M, Libkind D,**
Hittinger CT. 2015. The genome sequence of *Saccharomyces eubayanus* and the
domestication of lager-brewing yeasts. Mol Biol Evol **32**:2818-2831.
27.   **Han EK, Cotty F, Sottas C, Jiang H, Michels CA.** 1995. Characterization of *AGT1* encoding a
general α-glucoside transporter from *Saccharomyces*. Mol Microbiol **17**:1093-1107.
28.   **Vidgren V, Gibson B.** 2018. Trans-regulation and localization of orthologous maltose
transporters in the interspecies lager yeast hybrid. FEMS yeast research **18**:foy065.
29.   **Salema-Oom M, Pinto VV, Gonçalves P, Spencer-Martins I.** 2005. Maltotriose utilization by
industrial *Saccharomyces strains*: characterization of a new member of the α-glucoside
transporter family. Appl Environ Microbiol **71**:5044-5049.
30.   **Dietvorst J, Londesborough J, Steensma H.** 2005. Maltotriose utilization in lager yeast
strains: *MTT1* encodes a maltotriose transporter. Yeast **22**:775-788.
31.   **Cousseau F, Alves Jr S, Trichez D, Stambuk B.** 2013. Characterization of maltotriose
transporters from the *Saccharomyces eubayanus* subgenome of the hybrid *Saccharomyces*
pastorianus lager brewing yeast strain Weihenstephan 34/70. Lett Appl Microbiol **56**:21-29.
32.   **Nguyen H-V, Legras J-L, Neuvéglise C, Gaillardin C.** 2011. Deciphering the hybridisation
history leading to the lager lineage based on the mosaic genomes of *Saccharomyces bayanus*
strains NBRC1948 and CBS380T. PLoS One **6**:e25821.
33.   **Brouwers N, Gorter de Vries AR, van den Broek M, Weening SM, Elink Schuurman TD,**
Kuijpers NGA, Pronk JT, Daran J-MG. 2018. In vivo recombination of *Saccharomyces*
eubayanus maltose-transporter genes yields a chimeric transporter that enables maltotriose
fermentation. bioRxiv doi:10.1101/428839.
34.   **Brouwers N, Gorter de Vries AR, van den Broek M, Weening SM, Elink Schuurman TD,**
Kuijpers NGA, Pronk JT, Daran JG. 2019. In vivo recombination of Saccharomyces eubayanus
maltose-transporter genes yields a chimeric transporter that enables maltotriose
fermentation. PLoS Genet **15**:e1007853.

35. **Baker EP, Hittinger CT.** 2019. Evolution of a novel chimeric maltotriose transporter in
Saccharomyces eubayanus from parent proteins unable to perform this function. PLoS Genet
15:e1007786.
36. **Mertens S, Steensels J, Saels V, De Rouck G, Aerts G, Verstrepen KJ.** 2015. A large set of
newly created interspecific yeast hybrids increases aromatic diversity in lager beers. Appl
Environ Microbiol:AEM. 02464-02415.
37. **Gorter de Vries A, Voskamp MA, van Aalst ACA, Kristensen LH, Jansen L, van den Broek M,**
Salazar AN, Brouwers N, Abeel T, Pronk JT, Daran J-MG. 2018. Laboratory evolution of a
Saccharomyces cerevisiae x S. eubayanus hybrid under simulated lager-brewing conditions:
genetic diversity and phenotypic convergence. bioRxiv.
38. **Bing J, Han P-J, Liu W-Q, Wang Q-M, Bai F-Y.** 2014. Evidence for a Far East Asian origin of
lager beer yeast. Curr Biol **24**:R380-R381.
39. **Peris D, Langdon QK, Moriarty RV, Sylvester K, Bontrager M, Charron G, Leducq J-B, Landry**
C, Libkind D, Hittinger CT. 2016. Complex ancestries of lager-brewing hybrids were shaped
by standing variation in the wild yeast Saccharomyces eubayanus. PLoS Genet **12**:20.
40. **Darling ACE, Mau B, Blattner FR, Perna NT.** 2004. Mauve: multiple alignment of conserved
genomic sequence with rearrangements. Genome research **14**:1394-1403.
41. **Brown CA, Murray AW, Verstrepen KJ.** 2010. Rapid expansion and functional divergence of
subtelomeric gene families in yeasts. Curr Biol **20**:895-903.
42. **Moller M, Habig M, Freitag M, Stukenbrock EH.** 2018. Extraordinary Genome Instability and
Widespread Chromosome Rearrangements During Vegetative Growth. Genetics **210**:517-
529.
43. **Gordon JL, Byrne KP, Wolfe KH.** 2009. Additions, losses, and rearrangements on the
evolutionary route from a reconstructed ancestor to the modern *Saccharomyces cerevisiae*
genome. PLoS Genet **5**:e1000485.
44. **Marques WL, Mans R, Henderson RK, Marella ER, ter Horst J, de Hulster E, Poolman B,**
Daran J-M, Pronk JT, Gombert AK. 2018. Combined engineering of disaccharide transport
and phosphorolysis for enhanced ATP yield from sucrose fermentation in Saccharomyces
cerevisiae. Metabolic engineering **45**:121-133.
45. **Meurer M, Chevyreva V, Cerulus B, Knop M.** 2017. The regulatable MAL32 promoter in
Saccharomyces cerevisiae: characteristics and tools to facilitate its use. Yeast **34**:39-49.
46. **Levine J, Tanouye L, Michels CA.** 1992. The UAS(MAL) is a bidirectional promotor element
required for the expression of both the MAL61 and MAL62 genes of the Saccharomyces
MAL6 locus. Curr Genet **22**:181-189.
47. **Gorter de Vries AR, Groot PA, Broek M, Daran J-MG.** 2017. CRISPR-Cas9 mediated gene
deletions in lager yeast *Saccharomyces pastorianus*. Microb Cell Fact **16**:222.
48. **Vidgren V, Kankainen M, Londesborough J, Ruohonen L.** 2011. Identification of regulatory
elements in the AGT1 promoter of ale and lager strains of brewer's yeast. Yeast **28**:579-594.
49. **Dunn B, Sherlock G.** 2008. Reconstruction of the genome origins and evolution of the hybrid
lager yeast Saccharomyces pastorianus. Genome Res **18**:1610-1623.
50. **Okuno M, Kajitani R, Ryusui R, Morimoto H, Kodama Y, Itoh T.** 2016. Next-generation
sequencing analysis of lager brewing yeast strains reveals the evolutionary history of
interspecies hybridization. DNA Res **23**:67-80.
51. **Monerawela C, Bond U.** 2017. Brewing up a storm: The genomes of lager yeasts and how
they evolved. Biotechnology Advances **35**:512-519.
52. **Alsammar HF, Naseeb S, Brancia LB, Gilman RT, Wang P, Delneri D.** 2019. Targeted
metagenomics approach to capture the biodiversity of Saccharomyces genus in wild
environments. Environmental Microbiology Reports **11**:206-214.
53. **Vidgren V, Ruohonen L, Londesborough J.** 2005. Characterization and functional analysis of
the MAL and MPH Loci for maltose utilization in some ale and lager yeast strains. Appl
Environ Microbiol **71**:7846-7857.

54.   **Gallone B, Steensels J, Prahl T, Soriaga L, Saels V, Herrera-Malaver B, Merlevede A,**
Roncoroni M, Voordeckers K, Miraglia L, Teiling C, Steffy B, Taylor M, Schwartz A,
Richardson T, White C, Baele G, Maere S, Verstrepen KJ. 2016. Domestication and
Divergence of *Saccharomyces cerevisiae* Beer Yeasts. Cell **166**:1397-1410 e1316.
55.   **Gibson BR, Storgårds E, Krogerus K, Vidgren V.** 2013. Comparative physiology and
fermentation performance of Saaz and Frohberg lager yeast strains and the parental species
Saccharomyces eubayanus. Yeast **30**:255-266.
56.   **Salazar AN, Gorter de Vries AR, van den Broek M, Brouwers N, de la Torre Cortès P,**
Kuijpers NGA, Daran J-MG, Abeel T. 2019. Nanopore sequencing and comparative genome
analysis confirm lager-brewing yeasts originated from a single hybridization. bioRxiv
doi:10.1101/603480:603480.
57.   **Lei H, Zhao H, Yu Z, Zhao M.** 2012. Effects of wort gravity and nitrogen level on fermentation
performance of brewer's yeast and the formation of flavor volatiles. Appl Biochem
Biotechnol **166**:1562-1574.
58.   **Papapetridis I, Verhoeven MD, Wiersma SJ, Goudriaan M, van Maris AJA, Pronk JT.** 2018.
Laboratory evolution for forced glucose-xylose co-consumption enables identification of
mutations that improve mixed-sugar fermentation by xylose-fermenting Saccharomyces
cerevisiae. FEMS Yeast Research **18**.
59.   **Wisselink HW, Toirkens MJ, Wu Q, Pronk JT, van Maris AJA.** 2009. Novel evolutionary
engineering approach for accelerated utilization of glucose, xylose, and arabinose mixtures
by engineered *Saccharomyces cerevisiae* strains. Appl Environ Microbiol **75**:907-914.
60.   **Bolat I, Romagnoli G, Zhu F, Pronk JT, Daran JM.** 2013. Functional analysis and
transcriptional regulation of two orthologs of ARO10, encoding broad-substrate-specificity 2-
oxo-acid decarboxylases, in the brewing yeast Saccharomyces pastorianus CBS1483. FEMS
Yeast Res **13**:505-517.
61.   **Verduyn C, Postma E, Scheffers WA, Van Dijken JP.** 1992. Effect of benzoic acid on metabolic
fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic
fermentation. Yeast **8**:501-517.
62.   **Pronk JT.** 2002. Auxotrophic Yeast Strains in Fundamental and Applied Research. Applied and
Environmental Microbiology **68**:2095.
63.   **Solis-Escalante D, Kuijpers NG, Nadine B, Bolat I, Bosman L, Pronk JT, Daran J-MG, Pascale**
D-L. 2013. amdSYM, a new dominant recyclable marker cassette for *Saccharomyces*
cerevisiae. FEMS Yeast Res **13**:126-139.
64.   **Marques WL, Mans R, Marella ER, Cordeiro RL, van den Broek M, Daran J-MG, Pronk JT,**
Gombert AK, van Maris AJA. 2017. Elimination of sucrose transport and hydrolysis in
Saccharomyces cerevisiae: a platform strain for engineering sucrose metabolism. FEMS Yeast
Res **17**:fox006.
65.   **de Kok S, Yilmaz D, Suir E, Pronk JT, Daran J-MG, van Maris AJA.** 2011. Increasing free-
energy (ATP) conservation in maltose-grown *Saccharomyces cerevisiae* by expression of a
heterologous maltose phosphorylase. Metab Eng **13**:518-526.
66.   **Lee ME, DeLoache WC, Cervantes B, Dueber JE.** 2015. A Highly Characterized Yeast Toolkit
for Modular, Multipart Assembly. ACS Synth Biol **4**:975-986.
67.   **Gao Y, Zhao Y.** 2014. Self-processing of ribozyme-flanked RNAs into guide RNAs in vitro and
in vivo for CRISPR-mediated genome editing. Journal of integrative plant biology **56**:343-349.
68.   **Juergens H, Varela JA, Gorter de Vries AR, Perli T, Gast VJ, Gyurchev NY, Rajkumar AS,**
Mans R, Pronk JT, Morrissey JP. 2018. Genome editing in Kluyveromyces and Ogataea yeasts
using a broad-host-range Cas9/gRNA co-expression plasmid. FEMS yeast research **18**:foy012.
69.   **Engler C, Kandzia R, Marillonnet S.** 2008. A One Pot, One Step, Precision Cloning Method
with High Throughput Capability. PLOS ONE **3**:e3647.

70. **van Rossum HM, Kozak BU, Niemeijer MS, Duine HJ, Luttik MA, Boer VM, Kötter P, Daran J-**
MG, van Maris AJA, Pronk JT. 2016. Alternative reactions at the interface of glycolysis and
citric acid cycle in *Saccharomyces cerevisiae*. FEMS Yeast Res **16**.

71. **Gietz RD, Schiestl RH.** 2007. High-efficiency yeast transformation using the LiAc/SS carrier
DNA/PEG method. Nature protocols **2**:31.

72. **Pengelly RJ, Wheals AE.** 2013. Rapid identification of *Saccharomyces eubayanus* and its
hybrids. FEMS Yeast Res **13**:156-161.

73. **Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM.** 2017. Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome
Res:gr. 215087.215116.

74. **Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,**
Wortman J, Young SK. 2014. Pilon: an integrated tool for comprehensive microbial variant
detection and genome assembly improvement. PloS one **9**:e112963.

75. **Li H, Durbin R.** 2010. Fast and accurate long-read alignment with Burrows–Wheeler
transform. Bioinformatics **26**:589-595.

76. **Holt C, Yandell M.** 2011. MAKER2: an annotation pipeline and genome-database
management tool for second-generation genome projects. BMC bioinformatics **12**:491.

77. **Korf I.** 2004. Gene finding in novel genomes. BMC bioinformatics **5**:59.

78. **Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B.** 2006. AUGUSTUS: ab initio
prediction of alternative transcripts. Nucleic acids research **34**:W435-W439.

79. **Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.** 2009.
BLAST+: architecture and applications. BMC bioinformatics **10**:421.

80. **Tai SL, Boer VM, Daran-Lapujade P, Walsh MC, de Winde JH, Daran J-M, Pronk JT.** 2005.
Two-dimensional transcriptome analysis in chemostat cultures combinatorial effects of
oxygen availability and macronutrient limitation in Saccharomyces cerevisiae. Journal of
Biological Chemistry **280**:437-447.

81. **Dobin A, Gingeras TR.** 2016. Optimizing RNA-Seq mapping with STAR, p 245-262, Data
Mining Techniques for the Life Sciences. Springer.

82. **Anders S, Pyl PT, Huber W.** 2015. HTSeq—a Python framework to work with high-throughput
sequencing data. Bioinformatics **31**:166-169.

83. **Robinson MD, McCarthy DJ, Smyth GK.** 2010. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics **26**:139-140.

84. **McCarthy DJ, Chen Y, Smyth GK.** 2012. Differential expression analysis of multifactor RNA-
Seq experiments with respect to biological variation. Nucleic acids research **40**:4288-4297.

85. **Anders S, McCarthy DJ, Chen Y, Okoniewski M, Smyth GK, Huber W, Robinson MD.** 2013.
Count-based differential expression analysis of RNA sequencing data using R and
Bioconductor. Nature protocols **8**:1765.

86. **Entian K-D, Kötter P.** 2007. 25 Yeast genetic strain and plasmid collections. Methods in
Microbiology **36**:629-666.

Table 1: *Saccharomyces* strains used in this study. The abbreviation *malΔ* indicates *mal11-mal12*::loxP *mal21-mal22*::loxP
mal31-32::loxP. Mal and Mtt denote the maltose and maltotriose phenotype respectively.

Name	Species	Relevant genotype	Origin
CDFM21L.1	*S. eubayanus*	Wildtype Mal[-] Mtt[-]	(38)
ABFM5L.1	*S. eubayanus*	Wildtype Mal[-] Mtt[-]	(38)
CBS 12357	*S. eubayanus*	Wildtype Mal[+] Mtt[-]	(1) Westerdijk institute[a]
CBS 1483	*S. pastorianus*	Wildtype Mal[+] Mtt[+]	Westerdijk institute[a]
CEN.PK113-7D	*S. cerevisiae*	*MATα MAL1x MAL2x MAL3x MAL4x MAL2-8C SUC2 LEU2 URA3*	(86)
IMZ616	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mp2/3hΔ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ* pUDC156 (*Spcas9 URA3 ARS4 CEN6*)	(44)
IMX1365	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ* pUDC156 (*URA3 cas9*) *sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-ScAGT1-ScCYC1$_t$*	(13)
IMX1702	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ* pUDC156 (*URA3 cas9*) *sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-SeMALT1-ScCYC1$_t$*	This study
IMX1704	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ* pUDC156 (*URA3 cas9*) *sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-SeMALT2-ScCYC1$_t$*	This study
IMX1706	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ* pUDC156 (*URA3 cas9*) *sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-SeMALT3-ScCYC1$_t$*	This study
IMX 1708	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ* pUDC156 (*URA3 cas9*) *sga1Δ:: ScTDH3$_p$-ScMAL12-ScADH1$_t$ ScTEF1$_p$-SeAGT1-ScCYC1$_t$*	This study
IMX1313	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ sga1Δ::ScTEF1$_p$-ScMAL31-ScCYC1$_t$* pUDC156 (*URA3 cas9*)	This study
IMX1313Δ	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ sga1Δ::ScTEF1$_p$-ScMAL31-ScCYC1$_t$*	This study
IMZ752	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ sga1Δ::ScTEF1$_p$-ScMAL31-ScCYC1$_t$* pUDE843 (*ori* (*ColE1*) *bla* 2μ *ScTDH3$_p$-SeMALS1-ScADH1$_t$ URA3*)	This study
IMZ753	*S. cerevisiae*	*MATa ura3-52 LEU2 MAL2-8C malΔ mph2/3Δ suc2Δ ima1Δ ima2Δ ima3Δ ima4Δ ima5Δ sga1Δ::ScTEF1$_p$-ScMAL31-ScCYC1$_t$* pUDE844 (*ori* (*ColE1*) *bla* 2μ *ScTDH3$_p$-SeMALS2-ScADH1$_t$ URA3*)	This study
IMK820	*S. eubayanus*	*MATα/MATα Semalt1Δ/Semalt1Δ*	This study
IMK823	*S. eubayanus*	*MATα/MATα Seagt1Δ/Seagt1Δ* (X3)	This study
IMX1939	*S. eubayanus*	*MATα/MATα Semalt1Δ/Semalt1Δ Sesga1Δ::ScMAL13/Sesga1Δ::ScMAL13*	This study
IMX1940	*S. eubayanus*	*MATα/MATα Seagt1Δ/Seagt1Δ Sesga1Δ::ScMAL13/Sesga1Δ::ScMAL13*	This study
IMX1762	*S. eubayanus*	*MATα/MATα Sesga1Δ::ScMAL12/Sesga1Δ::ScMAL12*	This study
IMX1765	*S. eubayanus*	*MATα/MATα Sesga1Δ::ScMAL13/Sesga1Δ::ScMAL13*	This study
CBC-1	*S. cerevisiae*	*MATα/MATα* Mal[+] Mtt[-]	Lallemand
HTSH020	*S. cerevisiae X S. eubayanus*	*MATα/MATα* Mal[+] Mtt[+]	This study

899 [a] Westerdijk Fungal Biodiversity Institute www.westerdijkinstitute.nl/

Table 2 Plasmids used in this study

Plasmid	Relevant genotype	Origin
p426-TEF-amdS	ori ($ColE1$) bla 2μ amdSYM $TEF1_{pr}$-$CYC1_{ter}$	(64)
pUD444	ori ($ColE1$) bla 2μ amdSYM $ScTEF1_{pr}$-$ScMAL31$-$ScCYC1_{ter}$	This study
pUD794	ori ($ColE1$) bla 2μ amdSYM $ScTEF1_{pr}$-$SeMALT1$-$ScCYC1_{ter}$	This study
pUD795	ori ($ColE1$) bla 2μ amdSYM $ScTEF1_{pr}$-$SeMALT2$-$ScCYC1_{ter}$	This study
pUD796	ori ($ColE1$) bla 2μ amdSYM $ScTEF1_{pr}$-$SeMALT3$-$ScCYC1_{ter}$	This study
pUD797	ori ($ColE1$) bla 2μ amdSYM $ScTEF1_{pr}$-$SeAGT1$-$ScCYC1_{ter}$	This study
pUDE044	ori ($ColE1$) bla 2μ $ScTDH3_{pr}$-$ScMAL12$-$ScADH1_{ter}$ URA3	(65)
pUDE843	ori ($ColE1$) bla 2μ $ScTDH3_{pr}$-$SeMALS1$-$ScADH1_{ter}$ URA3	This study
pUDE844	ori ($ColE1$) bla 2μ $ScTDH3_{pr}$-$SeMALS2$-$ScADH1_{ter}$ URA3	This study
pUDE780	ori ($ColE1$) bla 2μ $ScPGK1_{pr}$-$ScMAL13$-$ScTEF2_{ter}$ URA3	This study
pUDP002	ori ($ColE1$) bla panARSopt Hyg $ScTDH3_{pr}$-BsaI-BsaI-$ScCYC1_{ter}$ $AaTEF1_{pr}$-$Spcas9^{D147Y\ P411T}$-$ScPHO5_{ter}$	(68)
pUDP004	ori ($ColE1$) bla panARSopt amdSYM $ScTDH3_{pr}$-BsaI-BsaI-$ScCYC1_{ter}$ $AaTEF1_{pr}$-$Spcas9^{D147Y\ P411T}$-$ScPHO5_{ter}$	(47)
pUDP052	ori ($ColE1$) bla panARSopt amdSYM $ScTDH3_{pr}$- gRNA$_{SGA1}$-$ScCYC1_{ter}$ $AaTEF1_{pr}$-$Spcas9^{D147Y\ P411T}$-$ScPHO5_{ter}$	(13)
pUDP091	ori ($ColE1$) bla panARSopt amdSYM $ScTDH3_{pr}$- gRNA$_{SeMALT1}$-$ScCYC1_{ter}$ $AaTEF1_{pr}$-$Spcas9^{D147Y\ P411T}$-$ScPHO5_{ter}$	This study
pUDP090	ori ($ColE1$) bla panARSopt amdSYM $ScTDH3_{pr}$- gRNA$_{SeAGT1}$-$ScCYC1_{ter}$ $AaTEF1_{pr}$-$Spcas9^{D147Y\ P411T}$-$ScPHO5_{ter}$	This study
pUDR119	ori ($ColE1$) bla 2μ amdSYM $SNR52_{pr}$-gRNA$_{ScSGA1}$-$SUP4_{ter}$	(70)
pYTK074	ori ($ColE1$) cat URA3	(66)
pYTK082	cat 2μ	(66)
pYTK083	ori ($ColE1$) bla	(66)
pUD631	ori ($ColE1$) bla gRNA$_{SeMALT1}$	(13)
pUD634	ori ($ColE1$) bla gRNA$_{SeAGT1}$	This study
pUDC156	ori ($ColE1$) bla ARS4 CEN6 URA3 Spcas9	(44)

Legends to Figures

Figure 1. Genome comparison between CDFM21L.1 and CBS 12357[T]. (A) Translocations in

CDFM21L.1 relative to CBS12357[T], (i) a paracentric inversion in the sub-telomeric region of

chromosome VII, involving approximately 8 kbp, (ii) a translocation of approximately 12 kbp from the

left sub-telomeric region of chromosome VIII to the right sub-telomeric region of chromosome VI,

and (iii) a reciprocal translocation between approximately 20 kbp from the right sub-telomeric region

of chromosome V and approximately 60 kbp from the center of chromosome XII.Red lines depicted

chromosomes separation. Genome syntheny is indicated with colored blocks. **(B)** Relative

chromosome position of gene presence differences between CDFM21L.1 (blue) and CBS 12357[T] (red).

(C) Representation of the assembled CDFM21L.1 *S. eubayanus* chromosomes, the black boxes denote

newly added sequences. New annotated open reading frames and gene entries modified relative to

the CBS 12357T draft genome (13).

Figure 2. Organization of sub-telomeric regions involving *MAL* genes and *SeAGT1* in CDFM21L.1. (A)

Chromosome sections are represented as lines and red boxes denote telomeres. CDFM21L.1 genome

harbors three *SeMALT* genes of which *SeMALT2* and *SeMALT3* have a mutation resulting in an early

stop codon and truncated protein (denoted with *). Three copies of *SeAGT1* were found close to the

telomeres on CHRVII, XIV and XV. Furthermore there are two intact *SeMALS* genes on CHRII (1) and

XII (2) and three *SeMALR* genes on CHRV (1) and XIII (2) whose copy on CHRV is also mutated

(*SeMALR1**). The gene and interval sizes are approximately to scale. (**B**) Phylogeny of *Saccharomyces*

SeAGT1 genes described in *S. cerevisiae*, *S. eubayanus*, and lager-brewing hybrid *S. pastorianus*. (C)

Nucleotide percentage identities between *AGT1* orthologs from *S. cerevisiae*, *S. eubayanus*, and

lager-brewing hybrid *S. pastorianus*. Green color indicates highest similarity between SeAGT1 and

SeAGT1 genes from *S. pastorianus* strains CBS 1483 and WS3470. Red color indicates similarity

between SeAGT1 from North American strains with SeAGT1 genes from Asian *S. eubayanus* and *S.*

pastorianus strains CBS 1483 and WS3470.

Figure 3. Characterization of sugar consumption of _S. pastorianus_ CBS1483 (black) and _S._
eubayanus CBS 12357^T (blue), CDFM21L.1 (red), and ABFM5L.1 (green) on wort. For every sample,
glucose (■), maltose (▲), maltotriose (▼) and ethanol (◆) were measured from the supernatant.
Strains were grown at 12 °C for 110 hours in infusion Neubor flasks. Samples were filtered through a
0.22 μm filter and analyzed on HPLC. Data represents average and standard deviation of three
biological replicates.

Figure 4. Overexpression of _SeMALT_, _SeAGT1_ and _SeMALS_ genes in a maltose negative background
S. cerevisiae strain. Maltose negative background strain IMZ616 (■), IMX1365 overexpressing
ScMAL11 (▲), IMX1702 overexpressing _SeMALT1_ (▼), IMX1704 overexpressing _SeMALT2_ (◆)
IMX1706 overexpressing _SeMALT3_ (●) and IMX1708 overexpressing SeAGT1 (□) were grown on SM
2% maltose or maltotriose at 20 °C. Growth on maltose **(A)** and on maltotriose **(B)** was monitored
based on optical density (OD$_{660nm}$) and concentrations of maltose and maltotriose in culture
supernatants were measured by HPLC. Data are presented as average and standard deviation of two
biological replicates. **(C)** IMX1313 overexpressing only _ScMAL31_ (O), IMZ752 overexpressing
ScMAL31 and _SeMALS1_ (△) and IMZ753 overexpressing _ScMAL31_ and _SeMALS2_ (▽) grown on SM
maltose 2%. Growth was monitored based on optical density measurement at 660 nm (OD$_{660nm}$) and
maltose in culture supernatants was measured by HPLC. Data represents average and standard
deviation of two biological replicates.

Figure 5. Integration of _ScMAL13_ in CDFM21L.1 restores growth on maltose and maltotriose and
enables native _SeMALT1_ and _SeAGT1_ characterization in knockout strains IMK820 and IMK823. (A)
Characterization of _S. eubayanus_ IMX1765 overexpressing _ScMAL13_ (▼) and CDFM21L.1 (■) on SM
with glucose, maltose or maltotriose at 20 °C. OD$_{660nm}$ was measured (black) and sugar (black) and
ethanol (red) concentrations were determined from the supernatant by HPLC. **(B)** Overview of

constructed knockout strains. Knockouts of *SeMALT1* (IMK820) and *SeAGT1* (IMK823) were made

with CRISPR-Cas9. Subsequently the *SeSGA1* locus was replaced by *ScPGK1$_p$- ScMAL13-ScTEF2$_t$* using

CRISPR-Cas9 in both strains resulting in IMX1939 and IMX1940, respectively. **(C)** *S. eubayanus* strains

IMK820 (■), IMK823 (▲), IMX1939 (▼) and IMX1940 (◆) were characterized on SM with maltose or

maltotriose at 20 °C. OD$_{660nm}$ was measured (black) and sugar (black) and ethanol (red)

concentrations were determined from the supernatant by HPLC. All data represents average and

standard deviation of biological duplicates.

Figure 6. Expression levels of maltose metabolism genes in IMX1765. Normalized transcript levels of

maltose metabolism genes from IMX1765 mid-exponential phase grown on glucose (red), maltose

(blue)and maltotriose (green) at 20 °C were calculated from duplicate RNA sequencing experiments

(2x 150 bp) using the FPKM method . All data represents average and standard deviation of two

biological duplicates.

Figure 7. Hybridization of maltotriose deficient *S. cerevisiae* and *S. eubayanus* leading to crosstalk

restoring maltotriose utilization, explains *S. pastorianus* phenotype. (A) Characterization of *S.*

cerevisiae CBC-1 (▼), *S.* eubayanus CDFM21L.1 (▲) and hybrid HTS020 (■) on mock wort at 12 °C.

Consumption of maltose (red), maltotriose (blue) and production of ethanol (green) was measured

from supernatant by HPLC. Data represents average and standard deviation from biological

triplicates. **(B)** Normalized transcript levels of maltose metabolism genes from HTS020 mid-

exponential phase grown on glucose (red), maltose (blue)and maltotriose (green) at 20 °C were

calculated from duplicate RNA sequencing experiments (2x 150 bp) using the FPKM method . All

data represents average and standard deviation of two biological duplicates.

FPKM