1   **Title**
2
3   **Deep data mining reveals variable abundance and distribution of microbial reproductive**
4   **manipulators within and among diverse host species**
5
6   **Authors**
7

8   Paloma Medina[1]*, Shelbi L Russell[2], Kavya Aswadhati[1], and Russell Corbett-Detig[1]*
9
Affiliations
[1] Genomics Institute and Department of Biomolecular Engineering UC Santa Cruz, Santa
Cruz, CA 95064
[2] Department of Molecular, Cellular and Developmental Biology, UC Santa Cruz, Santa Cruz,
CA 95064
* Correspondence To: pamedina@ucsc.edu, rucorbet@ucsc.edu

Abstract

Bacterial symbionts that manipulate the reproduction of their hosts are important factors in
invertebrate ecology and evolution. Studying the genomic and phenotypic diversity of
reproductive manipulators can improve efforts to control infectious diseases and contribute to
our understanding of host-symbiont evolution. Despite the vast genomic and phenotypic
diversity of reproductive manipulators, only a handful of strains are used as biological control
agents because little is known about the broad scale infection frequencies and densities of
these bacteria in nature. Here we develop a data mining approach to quantify the number of
arthropod and nematode host species infected with *Wolbachia* and other reproductive
manipulators such as *Rickettsia* and *Spiroplasma.* Across the entire Sequence Read Archive
(SRA) database, we found reproductive manipulators infected 2,083 arthropod and 119
nematode samples, representing 240 and 8 species, respectively. After accounting for sampling
and infection frequency differences among species, we estimated that *Wolbachia* infects
approximately 44% of all arthropod species and 34% of all nematode species. In contrast, we
estimated other reproductive manipulators infect 1-8% of arthropod and nematode species.
Next, we explored another important biological parameter: the relative bacterial density, or titer,
within hosts. We found variation in titer within and between arthropod species to be large, and
that host species explains approximately 36% of variation in titer across our dataset. This
suggests bacterial strain and/or host species plays a role in shaping bacterial densities within
and between host species. By leveraging the model system *Drosophila melanogaster,* we also
found a number of host SNPs associated with titer in genes potentially relevant to host
interactions with *Wolbachia*, suggesting bacterial induced host genome evolution. Our study
demonstrates that data mining is a powerful tool to understand host-symbiont co-evolution and
opens an array of previously inaccessible questions for further analysis.

Introduction
Bacterial symbionts of eukaryotic hosts exhibit an impressive array of phenotypes that interact
with host biology. Included among these symbionts are bacteria that alter host reproduction in

order to increase their likelihood of transmission to the next host generation [1–4], a strategy
termed reproductive manipulation. Depending on the nature of host reproduction, reproductive
manipulators are generally transmitted vertically by associating with either the oocyte or
developing embryos [5]. Manipulative phenotypes range from strategies that prevent the
survival of uninfected offspring, such as cytoplasmic incompatibility, to strategies such as
feminization, male killing, and parthenogenesis, that actually change the sex ratio to favor
females for overall increased infection rates [1–4]. These drastic manipulations of normal host
biology mean that reproductive manipulators can induce reproductive isolation, drive changes in
sexuality, and alter the reproductive ecology of their hosts [6–8].
54
55   While reproductive manipulators undergo vertical transmission to reach the next host
56   generation, many species and strains also exhibit low rates of horizontal transmission between
57   contemporary host species [5]. This combination of transmission strategies coupled with
58   reproductive manipulation serves to quickly increase the prevalence of reproductive
59   manipulators in a host population and to spread them rapidly among host species.
60
Given the impressive abilities of these bacteria to colonize new hosts and impact host fertility
and development for their own reproduction, much effort has been spent attempting to estimate
the infection frequencies of reproductive manipulators. Due to its wide distribution and presence
in model organisms such as *Drosophila, Wolbachia* is one of the most well studied symbionts in
general, and is the best studied reproductive manipulator in particular. Estimates of the
frequency of *Wolbachia* infections amongst arthropods range from 11% [9] to 76% [10]. The
large variation in estimates is likely an effect of methodological differences including sampling
bias and variation in PCR assay sensitivity. Other reproductive manipulators in the *Cardinium,*
Arsenophonus, Rickettsia, and *Spiroplasma* clades are reported to occur in 4% to 7% of all
species [11]. However, these estimates are even less certain than those for *Wolbachia* because
less infection frequency data is available for these clades, and similar sampling and assay
biases might impact frequency estimates. Given that the probability of sampling an infected
individual is positively correlated to the prevalence of the symbiont and the number of
individuals sampled from the host population, undersampling also imposes a barrier to confident
detection of low frequency infections.
76
Reproductive manipulators also experience a range of fitness tradeoffs during host growth and
development. A symbiont must be present at sufficiently high frequencies within a host to
promote successful transmission to subsequent host generations. However, exceedingly high
abundance of symbiont cells relative to host cells, termed titer, may impose a significant fitness
cost for the host and their symbionts [12, 13]. Reproductive manipulators may be more virulent
to their hosts when they achieve high titers, and natural selection on symbiont proliferation and
host regulation should favor the evolution of intermediate symbiont frequencies [14, 15]. This
tradeoff between reliable transmission and fitness costs to hosts is therefore an essential
component of understanding reproductive manipulator co-evolution. Nonetheless, little is known
about the relative abundances of reproductive manipulators within host individuals in large part
due to the challenges of collecting these data in high throughput ways.
88

Whole genome sequencing and bioinformatic approaches offer appealing alternatives to
conventional PCR-based survey methods to estimate reproductive manipulator infection rates.
These methods are not biased by primer selection, are less sensitive to false positives due to
contamination, and enable testing of large numbers of samples. Similarly, genome-sequencing
is a potentially powerful tool for interrogating symbiont titer within host individuals. With Illumina
shotgun sequencing, when the genome of a potential host individual is sequenced, the host's
symbionts are also sequenced. This makes the publicly-available databases a treasure trove for
sampling reproductive manipulators with bioinformatic approaches. In fact, the NCBI Sequence
Read Archive (SRA) [16] contains ~60,000 sequencing runs for samples tagged as Arthropoda
alone. Searching, or mining, genomic sequencing data has been shown to be a cost effective
and powerful strategy to detect *Wolbachia* infections [9, 17]. However, prior studies did not
include other reproductive manipulators [9, 17] or were focused on a single host species [17].
Indeed, one of the most compelling arguments for using non-targeted, publicly available data is
to mitigate ascertainment biases towards selecting species already known to harbor
reproductive manipulator infections.
104
Characterizing the prevalence and distribution of reproductive manipulators could be especially
valuable to biomedical researchers using reproductive manipulators to control the spread of
human pathogenic viruses such as Zika, Dengue and Chikungunya [18–20]. Currently, only the
wMel strain of *Wolbachia* is being used as a biological control agent [20, 21]. However, the large
genetic and phenotypic diversity of reproductive manipulators could suggest different species or
strains of bacteria to more effectively combat the spread of arboviruses. Additionally, a strategy
that opportunizes on cytoplasmic incompatibility, termed the "sterile male technique", can be
used to control mosquito population sizes [22, 23]. As this strategy requires low infection
frequencies in females to work [24], and different strains may have compatible rescue abilities
[25], it is necessary to understand the diversity and distribution of these bacteria in nature.
Cataloguing the distribution and titer of reproductive manipulators builds a foundation to explore
reproductive manipulators as powerful and versatile biological control agents.
Here, we establish and validate a computational pipeline to determine the infection status of
host samples downloaded from the SRA database and estimate the titer of symbionts within
positively infected host individuals. We use this approach to (i) quantify *Wolbachia* infection
frequency and infection titer of hosts, (ii) quantify the infection frequency and titer of other
genera including a range of reproductive manipulator symbiont clades, (iii) compare infection
frequencies and titers (from i and ii) among host groups, and (iv) identify *Drosophila*
melanogaster SNPs associated with *Wolbachia* titer. From this work, we classified 2,083
arthropod and 119 nematode samples as infected with a reproductive manipulator, including 95
species with previously unknown infections. Additionally, we show substantial variation in
symbiont titer within and between arthropod host species. Moreover, we show that symbiont
titer does not vary systematically across bacterial clades which indicates that other evolutionary,
ecological, or physiological processes may explain the disparate global distributions of
reproductive manipulators. We present extensive validation of our methods, including
orthogonal *in vivo* approaches, to help validate our large-scale data mining approach for
addressing fundamental questions in symbiont genomics and evolutionary biology.

Methods
Reproductive manipulator reference genome panel
We built our BLAST database using RefSeq genome assemblies for *Arsenophonus,*
Spiroplasma, Rickettsia, and *Wolbachia.* We included three *Arsenophonus*, six *Wolbachia,* 27
Spiroplasma, five *Cardinium,* and 61 *Rickettsia* genome assemblies (Supplementary Table S1).
These genomes were selected to span the known diversity of these bacterial groups. On
average, each genome assembly was about 1.3 Mb. We used these 102 genome assemblies to
build a BLAST database using the blastdb command from the NCBI blast package (version
2.7.1). We also included additional *Wolbachia* reference genomes to estimate symbiont titer,
which are listed in Supplementary Table S1, bringing the total number of references to 141.

Arthropod and nematode SRA dataset
We downloaded all Arthropoda sequencing read data from the NCBI Sequence Read Archive
(SRA) database [16]. We filtered samples under the Arthropoda and Nematoda taxonomy for
those sequenced on Illumina sequencing platforms. We filtered nominally for whole genome
shotgun libraries, but for completeness we further removed samples that were marked as
"reduced representation", "chipseq", and other terms that preclude a fully random, shotgun
library approach. In total, we tested 27,256 arthropod and 5,229 nematode samples for
reproductive manipulator infections. We consolidated all subspecies into a single species which
resulted in a total of 1,299 arthropod and 128 nematode species. SRA metadata for samples we
classified can be found in Supplementary Tables S2 and S3.

Determining Infection Status using a BLAST-based approach.
In order to classify a host sample as positively or negatively infected with a reproductive
manipulator, we analyzed local alignments between host DNA sequence data and reproductive
manipulator reference genomes. We binned each reference genome into 5kb segments and
computed the proportion of bins with a significant hit (breadth of coverage). We also computed
the variance of BLAST hits across each bin and estimated the coverage of the reproductive
manipulator genome (Supplementary Method S1). We determined a sample to be a candidate
for a positive infection if it had a 90% breadth of coverage and >1x estimated coverage on a
reproductive manipulator reference genome.

Validation of bacterial detection pipeline
To estimate the sensitivity and the specificity of our reproductive manipulator annotation
method, we compared our results to a previous extensive survey conducted by [17], which was
PCR validated. We determined the reproductive manipulator infection status of 158 individuals
from the *Drosophila Genetics Research Panel* (DGRP) [26] using our computational pipeline.
These 158 *D. melanogaster* samples had matching PCR and WGS determined infection
statuses generated by [17] and had their sequence data available on the NCBI SRA database.
In addition, there were 16 samples we excluded from our analysis because four of them were
not available on the SRA, eight did not have corresponding PCR and WGS *Wolbachia* infection
statuses, and four had discrepancies in infection status between sequence iterations of DGRP
lines (Supplementary Method S2). See Supplementary Method S2, Table S4, and Table S5 for

subsampling validation methods, pipeline accuracy to divergent reference strains, and a
comparison of our approach to other methods [9].

Beta-binomial estimation of reproductive manipulator prevalence.
Beta-binomial distributions have been used to fit *Wolbachia* infectious status across species
previously [27]. The beta-binomial model considers N random variables, X_j, which are all
binomially distributed (*i.e.*, infected vs. not-infected), but each with different parameters q_j and
n_j, so that $X_j \sim Bin(q_j, n_j)$ (Figure S1). Using the approach developed in [27], we determined (1)
moment estimators *u* and *s*, (2) beta distribution parameters α and $\square$, and (3) the global
infection rate *x.* After we fit a beta distribution to the data, we took the integral from *c* to 1, where
c is the minimum infection rate of a species to be considered positively infected. For example, a
value of 0.001 means if one individual in 1000 is classified as positive, the species would be
classified as being positively infected. For consistency with previous work, we set *c* to equal
0.001 (as in [27]). See Supplementary Method S3, Figure S2, and Table S6 for beta-binomial
rationale and downsampling results, which aimed to reduce sampling intensity bias.

Estimating Symbiont Titer
We estimated the ratio of symbiont genome compliments to host genome compliments,
hereafter referred to as titer (Supplementary Method S4). First, we computed the number of
symbiont genome compliments through our BLAST-based method with all available *Wolbachia,*
Arsenophonus, Spiroplasma, and *Rickettsia* genomes available on NCBI (Supplementary
Method S1, Supplementary Table S1). Next, we computed the number of host genome
compliments by aligning DNA sequencing reads to single copy orthologous arthropod proteins
from OrthoDBv9 [28] and taking the average of maximum depth across proteins with hits. We
report the symbiont haploid : host pseudo haploid titer computation throughout this manuscript.
See Supplementary Methods S4 and Figures S3 and S4 for validation of our approach to
estimate host genome compliments.

Drosophila oocyte sampling, imaging, and analysis
We obtained *Drosophila melanogaster* and *Drosophila simulans* fly stocks infected with the
wMel and wRi strains of *Wolbachia* from the Sullivan Lab. Stage 9/10a oocytes were dissected
from these flies, stained and mounted on glass slides, and imaged with a SP5 Leica confocal
microscope. We analyzed the fluorescence due to *Wolbachia* as described in [29] (See
Supplementary Method S5).

Estimating the Biological and Methodological Contributions to Titer Variation
We fit several models to determine how much host/symbiont genetics contributes to variation in
titer. We fit Generalized Linear Models models to log transformed arthropod titer data using the
glm() functions in R version 3.5.0. Specifically, we fit a GLM of the form: glm(log(titer) ~
arthropod_species). We also compared that model to one including reproductive manipulator
clade (e.g. "*Wolbachia*") using a GLM of the form: glm(log(titer) ~ arthropod_species +
reproductive_manipulator_clade. We used a likelihood ratio test to compare these two model fits
using anova(model1, model2, test="Chisq"). In addition, to test for an effect of sequencing

strategy (see also Supplementary Method S8), we fit a GLM of the form: glm(log(titer) ~
arthropod_species + pool_status).
220
221   **Software Availability**
222   The pipeline and individual scripts used to classify microbial reproductive manipulator infections
223   and estimate titer can be accessed through GitHub (www.github.com/pamedina/prevalence).
224
225   **Results and Discussion**
226
Reproductive Manipulators Infect Arthropod and Nematode Hosts in the SRA
We developed a powerful bioinformatic pipeline to identify reproductive manipulator infections
within sequencing datasets. Briefly, our approach compares sequencing reads from a given
sample to a set of reference genomes from reproductive manipulator species to determine if a
given host is positively infected (Figure 1, see Methods). We extensively characterized the
sensitivity and accuracy of our bioinformatic pipeline using previously known *Wolbachia*
infection statuses of the Drosophila Genetic Reference Panel (DGRP [26, 30]) and other
samples known to harbor genetically divergent *Wolbachia* [31] (see Supplementary Method S2).
Using the confident infection status lines in the DGRP as our validation set, our method is
completely concordant with both PCR and previous bioinformatic methods. Additionally, our
approach remains accurate even for references that exhibit 5-15% pairwise sequence
divergence (Supplementary Tables S7 and S8, Figures S5 and S6), indicating that our pipeline
is an accurate and robust method to determine reproductive manipulator infections in the
majority of host samples.
241
Using our bioinformatic classification pipeline (see Methods, Figure 1), we tested nearly all
arthropod and nematode samples on the SRA database that had been whole genome shotgun
sequenced (as of January 22, 2020) and found 240 arthropod species and 8 nematode species
had samples infected with a reproductive manipulator out of 1,299 and 128 species tested,
respectively (Table 1, Supplementary Figure S7 and S8, Table S9 and S10, File S1 and S2).
We identified 95 arthropod species with previously unreported reproductive manipulator
infections (Supplementary Table S11 and Method S6).
249
Wolbachia was the most frequent reproductive manipulator in arthropod samples and species
(1,978 samples, 179 species, Table 1)**.** We also found numerous arthropod species infected
with other clades of reproductive manipulators (Supplementary Table S9). Specifically,
Cardinium, Rickettsia, *Spiroplasma*, and *Arsenophonus* infect two, 54, 69, and nine arthropod
species in our dataset, respectively. The substantially lower infection rates for other
reproductive manipulators than for *Wolbachia* are consistent with prior work [32–34]
(Supplementary Table S10, Figure S8).
257
Wolbachia was the sole reproductive manipulator found in nematodes besides an instance of
Cardinium, as expected based on previous work [11, 35]. Almost all nematode species infected
with *Wolbachia* are filarial worms (Supplementary Table S10), a result supported by previous
studies showing that filarial nematodes and *Wolbachia* are in an obligatory, mutualistic

relationship [36–38]. There was one non-filarial nematode species infected with *Wolbachia*:
Pratylenchus penetrans, a plant-parasitic nematode (order Tylenchida). This species has been
shown previously to be infected with *Wolbachia* [35, 39]. These results, in addition to our
validation experiments, indicate that our approach can accurately detect symbiont infections.
266
We analyzed co-infections at two different levels: species harboring samples infected with more
than one reproductive manipulator clade and single samples infected by more than one
bacterial clade. We found high rates of co-infections of different reproductive manipulators
among arthropod hosts species. Species co-infections occurred in 29 of the 240 arthropod
species that harbored any infection ($p < 0.001$ permutation test, Supplementary Method S7,
Figure S9, and Table S12). Our observed rate of co-infection is consistent with previous work
that identified eight out of 44 species were co-infected ([11], $p = 0.34$, Fisher's Exact Test).
However, we found no evidence for an excess of individual samples infected by more than one
reproductive manipulator relative to expectations obtained by permutation ($p = 0.155$, 27
individuals, Supplementary Figure S10). These results may indicate that a subset of hosts, at
the species level, are more likely to acquire symbionts than others or that they are more
permissive to prolonged infections. This is supported by the observation that some host species
harbor genetic variation that influences the abundance of symbionts within individuals [14, 40].
Alternatively, any direct or indirect mutualism between symbionts might facilitate the build-up of
reproductive manipulators within a single species or host maternal lineage.
282
Global Rates of Infection Range from 1-44% Across Symbiont Taxa
Our study aims to estimate the infection rate of reproductive manipulators in arthropod and
nematode species overall, however, different taxa have been sampled to varying degrees and
infection frequencies vary (Supplementary Table S9) so the number cannot be directly
calculated from infection counts of individuals. To evaluate and correct for this ascertainment
bias, we use an approach based on modelling observed infections using a beta-binomial
distribution to estimate the total proportion of species infected with a reproductive manipulator
bacterial species as has been done previously [27] (Supplementary Method S3, Figure S1).
Using this approach, we estimated that 44% (95% CI 29-61%), and 34% (95% CI 5-69%), of
arthropod and nematode species, respectively, are infected with *Wolbachia* (Table 2). We note
that these values are consistent with expectations from previous work [10, 41–47]. Moreover,
because the SRA has been populated with samples mostly without considering *Wolbachia*
infection status, our approach should provide a relatively unbiased estimate of global infection
frequency. Nonetheless, it is possible that other sampling biases, *e.g.,* medical relevance of
focal species, might impact our estimates if medically relevant groups have unusual infection
frequencies relative to an "average" arthropod or nematode.
299
Among the other reproductive manipulators, we estimated *Arsenophonus*, *Rickettsia*, and
Spiroplasma infected 1% (95% CI 0-5%), 3% (95% CI 1-8%), and 8% (95% CI 3-15%) of
arthropod species, respectively (Table 2). We were not able to estimate global infection rates of
Cardinium because of the extremely low rate of positive infection in our dataset. Owing to the
fact that we do not have positive and negative controls readily available for each of these other
reproductive manipulator clades, it is difficult to completely rule out infections that failed to map

to the known reference(s) for each group and therefore induce a higher rate of false negatives.
However, our results from mapping *Wolbachia* reads to extremely diverse reference genomes
(e.g., 5-15% divergence) suggests that the rate of false negatives is low, provided the
divergence within these other bacterial groups does not exceed our tested values and, as we
note above, our raw frequency estimates are in line with previous work based on other methods
[27, 48, 49].
312
Taxonomic Distribution of Reproductive Manipulator Infections
The taxonomic distribution of *Wolbachia* spans 11 arthropod orders, out of the 47 tested (Figure
2). Across all of the arthropod species that we studied here, the orders with the greatest number
of species sampled are Hymenoptera, Diptera, Lepidoptera, Coleoptera, and Hemiptera. These
orders had 152, 195, 400, 139, and 91 species sampled, respectively. Positively infected
samples are dispersed widely across the phylogeny, including hosts as distant as Coleoptera
and Araneae. Horizontal transmission among species may contribute towards explaining
Wolbachia's large taxonomic distribution despite only modest sequence level divergence among
Wolbachia strains.
322
Wolbachia infection frequencies vary substantially across insect orders. For example,
Hemiptera, Hymenoptera, Diptera, and Coleoptera vary by approximately 30%, with 54%, 42%,
63%, 78% of species estimated to be infected, respectively (Table 3). Conversely, we
estimated the lowest *Wolbachia* infection frequency for Lepidoptera (36%, 95% CI 9.28-70.4%,
Table 3). Lepidoptera's infection frequency, in particular, is slightly lower than the frequencies
found in other orders of insects ($p < 0.08$, permutation test). Low frequencies might be the result
of high fitness cost to infected hosts or from *Wolbachia's* possible low transmission fidelity in
these groups. Notably, *Wolbachia* infections feminize or kill males in Lepidopteran hosts [50].
Both of these phenotypes can impose fitness costs on populations harboring these symbionts
because fewer males are available for mating and, in the case of feminization, males tend to
prefer genetic females over feminized males, which have lower mating rates and receive less
sperm [51]. Lower infection frequencies of bacteria with sex-ratio-distorting phenotypes might
suggest feminization and male killing have a higher fitness cost, or that host populations can
more easily subvert sex-ratio-distorting phenotypes compared to other reproductive phenotypes
like cytoplasmic incompatibility and parthenogenesis. Sex-ratio-distorting phenotypes might
therefore play a role in limiting infection polymorphism in host populations.
339
Symbiont Titer
Replication control is important for vertically transmitted symbionts because the fitness of the
symbiont is dependent on the fitness of the host. Reproductive manipulator infections must
remain sufficiently high to ensure vertical transovarial transmission while being low enough to
minimize pathogenic cost to a host [52]. Thus, replication control is important for reproductive
manipulators to be maintained in host populations. However, the extent to which reproductive
manipulator titers vary between orders of arthropods and among reproductive manipulator
clades is largely unknown and might be an important component of the fitness costs
reproductive manipulator species impose on their arthropod hosts.
349

In Vivo and *In Silico* Validation to Estimate Symbiont Titer

To evaluate the accuracy of our titer estimation method (Figure 1F, 1G, 1E and Supplementary Method S4, Supplementary Table S13), we compared the relative density of *Wolbachia* within *D. melanogaster* and *D. simulans*, which are two of the most commonly studied species known to host *Wolbachia*. The titer of *Wolbachia* has been demonstrated to influence the degree of *Wolbachia's* virulence on its host [53]. We hypothesized that titers of *Wolbachia* infecting *D. melanogaster* would be significantly lower than the titer of strains infecting *D. simulans* for three reasons. First, the strength of reproductive manipulation, which may be influenced by titer, is stronger in *D. simulans* than *D. melanogaster* [54, 55]. Second, the wRi strain commonly found in *D. simulans* has been reported to exhibit higher titers during embryogenesis than wMel does in *D. melanogaster* [56]. Third, wMel transinfected into *D. simulans* exhibited significantly higher titers than in its native host, *D. melanogaster [12]*.

Consistent with our expectations, we found that the strains inhabiting *D. simulans* exhibit significantly higher titer relative to those in *D. melanogaster* (p < 1e-05, one-tailed Mann Whitney U Test, Figure 3). In fact, for some extreme samples, titer in *D. simulans* shows an astounding >30:1 bacterial to host haploid genome complement ratio (Figure 3). Accounting for relative genome sizes, this indicates that *Wolbachia* can contribute approximately 30% as much DNA as the host does to the sequence data for that sample. To confirm these findings, we compared the amount of fluorescence due to *Wolbachia* DNA staining between stage 9/10a developing oocyte cysts from *D. melanogaster* flies infected with wMel and *D. simulans* flies infected with wRi, and found an average 2.1x more signal in *D. simulans* than *D. melanogaster* (n=15 and 19, respectively, Wilcoxon Rank-Sum p < 5.12e-05; Figure 3, Supplementary Method S5). Thus, the bioinformatically predicted titer differences between *Wolbachia* strains in *D. melanogaster* and *D. simulans* are also reflected using a completely orthogonal method to assay titer in the female germline. Importantly, this suggests that our *in silico* approach can yield information about *Wolbachia* abundance in tissues that are most relevant to understanding *Wolbachia* transmission (i.e., the female germline).

Comparative Study of Titer Across Symbiont Taxa

While symbiont titer does not necessarily predict the type or intensity of reproductive manipulation phenotypes in insect hosts [53], it does generally predict the virulence, or cost of infection, to a host [13, 14]. High titers are necessary to ensure an adequate number of bacteria make it to the germline for reliable vertical transmission. However, too high of titer may negatively impact host fitness. Since endosymbiotic bacteria like *Wolbachia* are maternally inherited, strong fitness costs to the host would also impact the fitness of *Wolbachia*. Therefore, evolutionary theory predicts symbiont titer will evolve towards a "goldilocks" zone, minimizing fitness costs to host and maximizing vertical transmission of symbiont bacteria [57]. The mechanism for this could either be via selection on symbiont growth or physiological activity [58, 59] or selection on host sanctions and regulatory mechanisms employed on the symbionts [14, 15, 59, 60].

We sought to identify the largest contributing factors to symbiont titer variation. Variation in titer may be a consequence of genetics (including host, symbiont, and combined host-symbiont

genotype interactions) or a result of other biological variation among genetically similar
individuals (*e.g.*, due to ecology). We found significant inter-sample variability within species
(Figure 4A, Supplementary Table S14). For example, there is a more than 200-fold range in
Wolbachia titer across *Drosophila melanogaster* samples, consistent with previous estimates of
titer in another *Drosophila* species *[61]*. Similarly, our results are in the range of previous
estimates of *Wolbachia* titer in *Aedes Aegypti [62]*. To determine the degree to which host
species influences variation in titer, we fit a generalized linear model to the data (Supplementary
Table S14). We found that *Wolbachia* titer in arthropods varied significantly between arthropod
species ($p < 2e-16$, GLM, Figure 4A), suggesting a strong genetic component shaping symbiont
titer among arthropod species. Indeed, we found that this model can explain 36% of variation in
titer. The remaining 64% of titer variation may be explained by within species differences, such
as inter-individual variability and uncontrolled extrinsic factors, such as local diet or habitat. Our
results therefore support the idea that symbiont titer varies considerably both within and
between species.
408
We hypothesize that *Wolbachia* titer in arthropod hosts could be significantly different compared
to other reproductive manipulator bacterial titers because *Wolbachia* is found at substantially
higher frequencies across arthropod species and many studies have reported that *Wolbachia*
generally elicits smaller fitness effects on its hosts than do *Rickettsia, Arsenophonus,* and
Spiroplasma [63–68]. Although an expanded model including reproductive manipulator as an
explanatory variable provides a better fit to the data ($p = 0.004$, likelihood ratio test), this model
explains only an additional 0.5% of the variance in symbiont titer relative to a model where only
host species is included as an explanatory variable (see Methods). No reproductive manipulator
clade significantly contributed to titer variation ($p > 0.05$, likelihood ratio test). Our results
therefore indicate that titer does not systematically differ between clades of reproductive
manipulators, contrary to our expectations based on their infection frequencies and pronounced
differences in their phenotypic effects. Finding that *Wolbachia's* titer is relatively consistent with
that of other reproductive manipulators suggests that co-evolved symbiont titer may not
contribute to its widespread host range. These results suggest that other co-evolutionary
mechanisms, perhaps encoded by the host genome or regulating bacterial physiology,
contribute to *Wolbachia*'s widespread taxonomic host distribution and generally decreased
virulence relative to other widespread reproductive manipulator taxa.
426
If several individuals, some infected and some not, were pooled to produce a single library, this
could affect titer estimates. Most concerningly, the SRA metadata is not always complete. To
confirm that pooling does not strongly affect our estimated titer and therefore bias our results,
we performed a literature search and annotated each sample as pooled or not pooled when this
information can be confidently determined from the source publication (Supplementary Table
S15). Then, we discluded samples that we could not discernably categorize. We fit a GLM
model of the same form as above but with pooled sequencing status as an additional factor, we
found that this model does not fit significantly better than a GLM fitting titer and arthropod host
species ($p = 0.4$, likelihood ratio test). This suggests our estimates from the whole SRA are
robust, and that pooling samples does not significantly impact the variation we see in titer

across arthropod hosts in our dataset. Our approach therefore appears robust to vagaries
associated with aggregating diverse and distinct public datasets for computational analysis.
439
Host Genotypes Associated with *Wolbachia* Titer
Considering the strong influence *Wolbachia* strains can have on their hosts' reproductive
outcomes, it is likely that the host genome experiences strong selective pressures to respond to
or control the bacteria. Indeed, previous work [14] identified a gene in the wasp host, *Nasonia*
vitripennis, that effectively controls *Wolbachia* titer and has been under recent positive selection,
likely since the wVitA strain horizontally transferred into the species. To address this question
using our public data-sourced method, we used the DGRP to perform a genome-wide
association study (GWAS) of titer in *D. melanogaster* using the DGRP GWAS webtool [26].
448
Although the present analysis is exploratory, there are several noteworthy results that may
suggest mechanisms of host control over *Wolbachia* titer. We identified 16 candidate single
nucleotide polymorphisms (SNPs) in the *D. melanogaster* genome associated with *Wolbachia*
titer (Supplementary Table S16). One of the most strongly associated SNPs is found in a gene
associated with the endoplasmic reticulum (ER) membrane (*XBP1*) [69]. Consistent with this
functional prediction, a recent study on a wMel-infected *D. melanogaster* cell line found that
Wolbachia resides within ER-derived membrane near and within the ER itself [70]. Modifying
this membrane might therefore enable the host to impact *Wolbachia* titer on the cellular level.
Additionally, another strongly associated SNP is found in a gene associated with actin binding
and microtubule transport (CG43901). Recent work has shown that *Wolbachia* uses host actin
for localization in host tissues [71, 72] and modifications to actin-binding proteins can clear
Wolbachia infections in host individuals [73]. Finally, a SNP in CG17048 is also strongly
associated with *Wolbachia* titer in the DGRP. This gene's role in protein ubiquitination is
consistent with previous results from a genome-wide RNAi screen that found this process
contributes disproportionately to modifications of *Wolbachia* titer in cell culture [74]. These are
therefore appealing candidate genes for evaluating the potential of natural host variation to
control symbiont infections.
466
Functional work will be necessary to validate our specific predictions, nonetheless these titer-
associated genetic polymorphisms suggest that the host genome is capable of evolving to
control *Wolbachia* infections. Although the present association work is focused on a well-
characterized genetic mapping panel in *D. melanogaster*, our results illustrate more generally
the potential impacts of high throughput data mining for identifying both reproductive
manipulator infections and candidate host genetic factors that may be involved in controlling the
infections within host tissues.
474
Conclusion
Here, we presented a reference-based approach to detect bacterial infections in short read data
and highlighted the ways it can be used to generate insights into host-symbiont interactions.
Indeed, our work is the first to estimate the global distribution of multiple reproductive
manipulators across all sequenced arthropod and nematode hosts using full-genome high
throughput methods. Moreover, we show that publically-available short read data can be used

to interrogate other biological attributes of host-symbiont associations, such as titer. We found that symbiont titer, measured as relative genome copy number, is highly variable among arthropod host species, and that titer can vary 200 fold between and within host species. Furthermore, our database catalogs some of the vast phenotypic and genetic diversity of reproductive manipulators, which could be used directly as biological control agents to control the spread of infectious disease or indirectly to inform on the natural infection state of host targets.

While our approach relies on datasets gathered for a wide array of purposes and therefore requires a level of approximation, we have shown that accurate and robust predictions can still be obtained using this method. Moreover, as publicly available sequence data continues to accumulate at exceptional rates, this framework will become increasingly powerful relative to gathering purpose-built datasets to assay symbiont infection statuses and frequencies. More generally, our method and related approaches could be used to detect other microbial symbionts, such as medically relevant pathogens, or even viruses, for which a reference genome sequence is available. Hence, future work will build on this framework of leveraging increasingly vast datasets to conduct direct and precise hypothesis testing of fundamental questions in host and symbiont ecology and evolution.

Acknowledgements
We thank the Corbett-Detig Lab and Sullivan Lab members for helpful feedback on this work. During this research, PM was supported by NIH training grant (T32 HG008345). This work was supported in part by an Alfred P. Sloan Fellowship and by NIH/NIGMS R35GM128932 to RC-D.

Contributions
Conceived and designed research: PM, RC-D, SLR. Performed the computational screen: PM. Analyzed sequence data and developed software: PM. Oocyte imaging and analysis: SLR. Novel infection literature analysis: KA. Wrote the manuscript: PM, SLR, RC-D. Edited the manuscript: RC-D, SLR, PM.

References

1. Dedeine F, Vavre F, Fleury F, Loppin B, Hochberg ME, Bouletreau M. Removing symbiotic Wolbachia bacteria specifically inhibits oogenesis in a parasitic wasp. *Proc Natl Acad Sci U S A* 2001; **98**: 6247–6252.
2. Stevens L, Giordano R, Fialho RF. Male-Killing, Nematode Infections, Bacteriophage Infection, and Virulence of Cytoplasmic Bacteria in the GenusWolbachia. *Annu Rev Ecol Syst* 2001; **32**: 519–545.
3. Stouthamer R, Breeuwer JA, Hurst GD. Wolbachia pipientis: microbial manipulator of arthropod reproduction. *Annu Rev Microbiol* 1999; **53**: 71–102.
4. Werren JH. Biology of Wolbachia. *Annu Rev Entomol* 1997; **42**: 587–609.
5. Russell SL. Transmission mode is associated with environment type and taxa across bacteria-eukaryote symbioses: a systematic review and meta-analysis. *FEMS Microbiol Lett* 2019; **366**.
6. Bandi C, Dunn AM, Hurst GD, Rigaud T. Inherited microorganisms, sex-specific virulence and reproductive parasitism. *Trends Parasitol* 2001; **17**: 88–94.
7. Charlat S, Hurst GDD, Merçot H. Evolutionary consequences of Wolbachia infections. *Trends Genet* 2003; **19**: 217–223.
8. Hurst GD, Werren JH. The role of selfish genetic elements in eukaryotic evolution. *Nat Rev Genet* 2001; **2**: 597–606.
9. Pascar J, Chandler CH. A bioinformatics approach to identifying Wolbachia infections in arthropods. *PeerJ* 2018; **6**: e5486.
10. Jeyaprakash A, Hoy MA. Long PCR improves Wolbachia DNA amplification: wsp sequences found in 76% of sixty-three arthropod species. *Insect Mol Biol* 2000; **9**: 393–405.
11. Duron O, Bouchon D, Boutin S, Bellamy L, Zhou L, Engelstädter J, et al. The diversity of reproductive parasites among arthropods: Wolbachia do not walk alone. *BMC Biol* 2008; **6**: 27.
12. Serbus LR, Ferreccio A, Zhukova M, McMorris CL, Kiseleva E, Sullivan W. A feedback loop between Wolbachia and the Drosophila gurken mRNP complex influences Wolbachia titer. *J Cell Sci* 2011; **124**: 4299–4308.

13. Chong RA, Moran NA. Intraspecific genetic variation in hosts affects regulation of obligate heritable symbionts. *Proc Natl Acad*
Sci U S A 2016; **113**: 13114–13119.
14. Funkhouser-Jones LJ, van Opstal EJ, Sharma A, Bordenstein SR. The Maternal Effect Gene Wds Controls Wolbachia Titer in
Nasonia. *Curr Biol* 2018; **28**: 1692–1702.e6.
15. Mergaert P. Role of antimicrobial peptides in controlling symbiotic bacterial populations. *Nat Prod Rep* 2018; **35**: 336–356.
16. Leinonen R, Sugawara H, Shumway M, Collaboration INSD. The sequence read archive. *Nucleic Acids Res* 2010; **39**: D19–
D21.
17. Richardson MF, Weinert LA, Welch JJ, Linheiro RS, Magwire MM, Jiggins FM, et al. Population genomics of the Wolbachia
endosymbiont in Drosophila melanogaster. *PLoS Genet* 2012; **8**: e1003129.
18. Moreira LA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AT, Hedges LM, et al. A Wolbachia symbiont in Aedes aegypti limits
infection with dengue, Chikungunya, and Plasmodium. *Cell* 2009; **139**: 1268–1278.
19. Hoffmann AA, Montgomery BL, Popovici J, Iturbe-Ormaetxe I, Johnson PH, Muzzi F, et al. Successful establishment of
Wolbachia in Aedes populations to suppress dengue transmission. *Nature* 2011; **476**: 454–457.
20. Walker T, Johnson PH, Moreira LA, Iturbe-Ormaetxe I, Frentiu FD, McMeniman CJ, et al. The wMel Wolbachia strain blocks
dengue and invades caged Aedes aegypti populations. *Nature* 2011; **476**: 450–453.
21. Frentiu FD, Zakir T, Walker T, Popovici J, Pyke AT, van den Hurk A, et al. Limited dengue virus replication in field-collected
Aedes aegypti mosquitoes infected with Wolbachia. *PLoS Negl Trop Dis* 2014; **8**: e2688.
22. Zabalou S, Riegler M, Theodorakopoulou M, Stauffer C, Savakis C, Bourtzis K. Wolbachia-induced cytoplasmic incompatibility
as a means for insect pest population control. *Proc Natl Acad Sci U S A* 2004; **101**: 15042–15045.
23. Nikolouli K, Colinet H, Renault D, Enriquez T, Mouton L, Gibert P, et al. Sterile insect technique and Wolbachia symbiosis as
potential tools for the control of the invasive species Drosophila suzukii. *J Pest Sci* 2018; **91**: 489–503.
24. Landmann F, Orsi GA, Loppin B, Sullivan W. Wolbachia-mediated cytoplasmic incompatibility is associated with impaired
histone deposition in the male pronucleus. *PLoS Pathog* 2009; **5**: e1000343.
25. Bourtzis K, Nirgianaki A, Markakis G, Savakis C. Wolbachia infection and cytoplasmic incompatibility in Drosophila species.
Genetics 1996; **144**: 1063–1073.
26. Huang W, Massouras A, Inoue Y, Peiffer J, Ràmia M, Tarone AM, et al. Natural variation in genome architecture among 205
Drosophila melanogaster Genetic Reference Panel lines. *Genome Res* 2014; **24**: 1193–1208.
27. Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH. How many species are infected with Wolbachia?--A
statistical analysis of current data. *FEMS Microbiol Lett* 2008; **281**: 215–220.
28. Zdobnov EM, Tegenfeldt F, Kuznetsov D, Waterhouse RM, Simão FA, Ioannidis P, et al. OrthoDB v9.1: cataloging evolutionary
and functional annotations for animal, fungal, plant, archaeal, bacterial and viral orthologs. *Nucleic Acids Res* 2017; **45**: D744–
D749.
29. Russell SL, Lemseffer N, White PM, Sullivan WT. Wolbachia and host germline components compete for kinesin-mediated
transport to the posterior pole of the Drosophila oocyte. *PLoS Pathog* 2018; **14**: e1007216.
30. Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. The Drosophila melanogaster Genetic Reference
Panel. *Nature* 2012; **482**: 173–178.
31. Gerth M, Gansauge M-T, Weigert A, Bleidorn C. Phylogenomic analyses uncover origin and spread of the Wolbachia
pandemic. *Nat Commun* 2014; **5**: 5117.
32. Zchori-Fein E, Perlman SJ. Distribution of the bacterial symbiont Cardinium in arthropods. *Mol Ecol* 2004; **13**: 2009–2016.
33. Russell JA, Funaro CF, Giraldo YM, Goldman-Huertas B, Suh D, Kronauer DJC, et al. A veritable menagerie of heritable
bacteria from ants, butterflies, and beyond: broad molecular surveys and a systematic review. *PLoS One* 2012; **7**: e51027.
34. Sazama EJ, Ouellette SP, Wesner JS. Bacterial Endosymbionts Are Common Among, but not Necessarily Within, Insect
Species. *Environ Entomol* 2019.
35. Wasala SK, Brown AMV, Kang J, Howe DK, Peetz AB, Zasada IA, et al. Variable Abundance and Distribution of Wolbachia and
Cardinium Endosymbionts in Plant-Parasitic Nematode Field Populations. *Front Microbiol* 2019; **10**: 964.
36. Casiraghi M, Bain O, Guerrero R, Martin C, Pocacqua V, Gardner SL, et al. Mapping the presence of Wolbachia pipientis on the
phylogeny of filarial nematodes: evidence for symbiont loss during evolution. *Int J Parasitol* 2004; **34**: 191–203.
37. Bandi C, Anderson TJ, Genchi C, Blaxter ML. Phylogeny of Wolbachia in filarial nematodes. *Proc Biol Sci* 1998; **265**: 2407–
2413.
38. Casiraghi M, Anderson TJ, Bandi C, Bazzocchi C, Genchi C. A phylogenetic analysis of filarial nematodes: comparison with the
phylogeny of Wolbachia endosymbionts. *Parasitology* 2001; **122 Pt 1**: 93–103.
39. Brown AMV, Wasala SK, Howe DK, Peetz AB, Zasada IA, Denver DR. Genomic evidence for plant-parasitic nematodes as the
earliest Wolbachia hosts. *Sci Rep* 2016; **6**: 34955.
40. Login FH, Balmand S, Vallier A, Vincent-Monégat C, Vigneron A, Weiss-Gayet M, et al. Antimicrobial peptides keep insect
endosymbionts under control. *Science* 2011; **334**: 362–365.
41. Werren JH, Windsor DM. Wolbachia infection frequencies in insects: evidence of a global equilibrium? *Proc Biol Sci* 2000; **267**:
1277–1285.
42. Werren JH, Windsor D, Guo LR. Distribution of Wolbachia among neotropical arthropods. *Proc R Soc Lond B Biol Sci* 1995;
262: 197–204.
43. West SA, Cook JM, Werren JH, Godfray HCJ. Wolbachia in two insect host--parasitoid communities. *Mol Ecol* 1998; **7**: 1457–
1465.
44. Kikuchi Y, Fukatsu T. Diversity of Wolbachia endosymbionts in heteropteran bugs. *Appl Environ Microbiol* 2003; **69**: 6082–
6090.
45. Nirgianaki A, Banks GK, Frohlich DR, Veneti Z, Braig HR, Miller TA, et al. Wolbachia infections of the whitefly Bemisia tabaci.
Curr Microbiol 2003; **47**: 93–101.
46. Tagami Y, Miura K. Distribution and prevalence of Wolbachia in Japanese populations of Lepidoptera. *Insect Mol Biol* 2004; **13**:
359–364.
47. Gotoh T, Noda H, Hong X-Y. Wolbachia distribution and cytoplasmic incompatibility based on a survey of 42 spider mite
species (Acari: Tetranychidae) in Japan. *Heredity* 2003; **91**: 208–216.
48. Zug R, Hammerstein P. Still a host of hosts for Wolbachia: analysis of recent data suggests that 40% of terrestrial arthropod

species are infected. *PLoS One* 2012; **7**: e38544.

49. Weinert LA, Araujo-Jnr EV, Ahmed MZ, Welch JJ. The incidence of bacterial endosymbionts in terrestrial arthropods. *Proc Biol Sci* 2015; **282**: 20150249.

50. Werren JH, Baldo L, Clark ME. Wolbachia: master manipulators of invertebrate biology. *Nat Rev Microbiol* 2008; **6**: 741–751.

51. Moreau J, Bertin A, Caubet Y, Rigaud T. Sexual selection in an isopod with Wolbachia-induced sex reversal: males prefer real females. *J Evol Biol* 2001; **14**: 388–394.

52. McGraw EA, Merritt DJ, Droller JN, O'Neill SL. Wolbachia density and virulence attenuation after transfer into a novel host. *Proc Natl Acad Sci U S A* 2002; **99**: 2918–2923.

53. Duron O, Labbé P, Berticat C, Rousset F, Guillot S, Raymond M, et al. High Wolbachia density correlates with cost of infection for insecticide resistant Culex pipiens mosquitoes. *Evolution* 2006; **60**: 303–314.

54. Boyle L, O'Neill SL, Robertson HM, Karr TL. Interspecific and intraspecific horizontal transfer of Wolbachia in Drosophila. *Science* 1993; **260**: 1796–1799.

55. Reynolds KT, Hoffmann AA. Male age, host effects and the weak expression or non-expression of cytoplasmic incompatibility in Drosophila strains infected by maternally transmitted Wolbachia. *Genet Res* 2002; **80**: 79–87.

56. Veneti Z, Clark ME, Karr TL, Savakis C, Bourtzis K. Heads or tails: host-parasite interactions in the Drosophila-Wolbachia system. *Appl Environ Microbiol* 2004; **70**: 5366–5372.

57. Perrot-Minnot MJ, Werren, J. H. (Department of Biology, University of Rochester, Rochester, NY 14620 (USA)). Wolbachia infection and incompatibility dynamics in experimental selection lines. *J Evol Biol* 1999; **12**.

58. Enomoto S, Chari A, Clayton AL, Dale C. Quorum Sensing Attenuates Virulence in Sodalis praecaptivus. *Cell Host Microbe* 2017; **21**: 629–636.e5.

59. Yoder JB, Tiffin P. Sanctions, Partner Recognition, and Variation in Mutualism. *The American Naturalist* . 2017. , **190**: 491–505

60. Voronin D, Cook DAN, Steven A, Taylor MJ. Autophagy regulates Wolbachia populations across diverse symbiotic associations. *Proc Natl Acad Sci U S A* 2012; **109**: E1638–46.

61. Unckless RL, Boelio LM, Herren JK, Jaenike J. Wolbachia as populations within individual insects: causes and consequences of density variation in natural populations. *Proc Biol Sci* 2009; **276**: 2805–2811.

62. Crawford JE, Clarke DW, Criswell V, Desnoyer M, Cornel D, Deegan B, et al. Efficient production of male Wolbachia-infected Aedes aegypti mosquitoes enables large-scale suppression of wild populations. *Nat Biotechnol* 2020; **38**: 482–492.

63. Hurst GDD, Frost CL. Reproductive parasitism: maternally inherited symbionts in a biparental world. *Cold Spring Harb Perspect Biol* 2015; **7**.

64. Correa CC, Ballard JWO. Wolbachia Associations with Insects: Winning or Losing Against a Master Manipulator. *Front Ecol Evol* 2016; **3**: 506.

65. Harumoto T, Anbutsu H, Lemaitre B, Fukatsu T. Male-killing symbiont damages host's dosage-compensated sex chromosome to induce embryonic apoptosis. *Nat Commun* 2016; **7**: 12781.

66. Hayashi M, Watanabe M, Yukuhiro F, Nomura M, Kageyama D. A Nightmare for Males? A Maternally Transmitted Male-Killing Bacterium and Strong Female Bias in a Green Lacewing Population. *PLoS One* 2016; **11**: e0155794.

67. Parratt SR, Frost CL, Schenkel MA, Rice A, Hurst GDD, King KC. Superparasitism Drives Heritable Symbiont Epidemiology and Host Sex Ratio in a Wasp. *PLoS Pathog* 2016; **12**: e1005629.

68. Pang R, Chen M, Yue L, Xing K, Li T, Kang K, et al. A distinct strain of Arsenophonus symbiont decreases insecticide resistance in its insect host. *PLoS Genet* 2018; **14**: e1007725.

69. Ryoo HD, Li J, Kang M-J. Drosophila XBP1 expression reporter marks cells under endoplasmic reticulum stress and with high protein secretory load. *PLoS One* 2013; **8**: e75774.

70. Fattouh N, Cazevieille C, Landmann F. Wolbachia endosymbionts subvert the endoplasmic reticulum to acquire host membranes without triggering ER stress. *PLoS Negl Trop Dis* 2019; **13**: e0007218.

71. Landmann F, Bain O, Martin C, Uni S, Taylor MJ, Sullivan W. Both asymmetric mitotic segregation and cell-to-cell invasion are required for stable germline transmission of Wolbachia in filarial nematodes. *Biology Open* . 2012. , **1**: 536–547

72. Sheehan KB, Martin M, Lesser CF, Isberg RR, Newton ILG. Identification and Characterization of a Candidate Wolbachia pipientis Type IV Effector That Interacts with the Actin Cytoskeleton. *MBio* 2016; **7**.

73. Newton ILG, Savytskyy O, Sheehan KB. Wolbachia utilize host actin for efficient maternal transmission in Drosophila melanogaster. *PLoS Pathog* 2015; **11**: e1004798.

74. White PM, Serbus LR, Debec A, Codina A, Bray W, Guichet A, et al. Reliance of Wolbachia on High Rates of Host Proteolysis Revealed by a Genome-Wide RNAi Screen of Drosophila Cells. *Genetics* 2017; **205**: 1473–1488.

653 **Figure Descriptions**
654

Name	Description
Figure 1	A schematic of the computational pipeline used to determine the reproductive manipulator infection status and symbiont titer of a sequencing run.
Figure 2	Taxonomy of Arthropoda orders tested and number of reproductive manipulator-positive species within each order.
Figure 3	*Wolbachia* titer among *D. melanogaster* and *D. simulans* computed from our *in silico* titer estimate approach and *in vivo* assay.
Figure 4	Titer of positively infected Arthropoda host species for *Wolbachia*, *Rickettsia*, *Arsenophonus, Spiroplasma*.

Figure 1. A schematic of the computational pipeline used to determine the reproductive manipulator infection status and endosymbiont titer of a sequencing run (see also Supplementary Methods S1 and S7). The pipeline **(A)** takes in a sample's unique identification number, then **(B)** downloads two million endogenous and exogenous reads. Then, **(C)** reads are aligned to *Wolbachia, Arsenophonus, Spiroplasma, Cardinium,* and *Rickettsia* reference genomes, and **(D)** summary statistics for the sample aligned to each reference are computed. If a sample had between 0.1 and 0.9 breadth of coverage, the full dataset was downloaded and the workflow repeated to prevent false negative calls. **(F and G)** To estimate host coverage without requiring a reference genome, reads are also aligned to a set of 1066 single copy ancestral orthologs obtained from ORTHODB v9. Then, **(E)** we apply coverage breadth and depth cutoffs to classify infection status as positive, or negative. We compare the coverage of

host reads to single copy orthologs to the coverage of endosymbiont reads to reproductive manipulator reference genome to estimate endosymbiont load.

Order	Number of Species Samples	Species *Wolbachia* Positive	Species *Arsenophonus* Positive	Species *Rickettsia* Positive	Species *Spiroplasma* Positive
Diplura	2	0	0	0	0
Xiphosura	2	0	0	0	0
Amblypygi	1	0	0	0	0
Schizomida	1	0	0	0	0
Ricinulei	1	0	0	0	0
Araneae	10	2 (0.2)	0	0	0
Pseudoscorpiones	1	0	0	0	0
Opiliones	1	0	0	0	0
Scorpiones	7	0	0	1 (0.14)	0
Ixodida	3	0	0	0	0
Mesostigmata	2	0	0	0	0
Sarcoptiformes	7	0	0	0	0
Trombidiformes	4	1 (0.25)	0	1 (0.25)	0
Scutigeromorpha	1	0	0	0	0
Geophilomorpha	1	0	0	0	0
Podocopida	3	0	0	0	0
Myodocopida	1	0	0	0	0
Anostraca	1	0	0	0	0
Notostraca	3	0	0	0	0
Diplostraca	5	0	0	0	0
Decapoda	139	0	0	0	0
Isopoda	5	0	0	0	0
Amphipoda	3	0	0	0	0
Sessilia	1	0	0	0	0
Calanoida	4	0	0	0	0
Siphonostomatoida	3	0	0	0	0
Cyclopoida	2	0	0	0	0
Harpacticoida	3	0	0	0	0
Entomobryomorpha	3	0	0	0	0
Archaeognatha	1	1 (1.0)	0	0	0
Zygentoma	1	0	0	0	0
Odonata	2	1 (0.5)	0	0	0
Ephemeroptera	4	0	0	0	0
Coleoptera	139	29 (0.21)	0	18 (0.13)	2 (0.01)
Hymenoptera	152	27 (0.18)	7 (0.05)	3 (0.02)	5 (0.03)
Diptera	195	39 (0.20)	0	2 (0.01)	2 (0.01)
Plecoptera	5	0	0	0	0
Lepidoptera	400	46 (0.12)	0	5 (0.01)	33 (0.08)
Trichoptera	3	0	0	0	0
Blattodea	3	0	0	0	0
Orthoptera	7	0	0	0	1 (0.14)
Phasmatodea	2	1 (0.5)	0	0	0
Dermaptera	3	0	0	0	0
Hemiptera	91	31 (0.34)	1 (0.01)	5 (0.05)	0
Thysanoptera	2	1 (0.5)	0	0	0
Psocoptera	3	0	0	0	0
Phthiraptera	68	0	0	3 (0.04)	0

Figure 2. Phylogeny of Arthropoda orders tested and number of reproductive manipulator positive species within each order. The frequency of reproductive manipulator-positive species listed in parentheses. No frequency is listed if there was no infection within an arthropod order. We used the Tree of Life taxonomic and phylogenetic package and rotl (Michonneau et al. 2016), to group host species by their orders.

Figure 3. Wolbachia titer among *D. melanogaster* and *D. simulans* computed from our **(A)** in silico titer estimate approach (y-axis log10 scaled) and **(B-E)** in vivo fluorescence assay in stage 10a oocyte cysts. **(B)** Quantification of *Wolbachia* fluorescence, showing a significant difference in titer between host species (Wilcoxon rank sum test). **(C)** *D. melanogaster* and **(D)** *D. simulans* imaged propidium iodide DNA staining and confocal microscopy. **(E)** *Uninfected D. melanogaster* oocyte shows no evidence of *Wolbachia* staining. Scale bars = 50um.

Figure 4. Titer variation across species infected with *Wolbachia* and between diverse reproductive manipulator clades. To increase readability of both plots, categories were randomly downsampled to show 100 samples. The y-axis is log10 scaled. **(A)** Titer for *Wolbachia* positive arthropod samples were grouped by host species and plotted. Host species with at least three samples were plotted. **(B)** Titer for *Wolbachia*, *Rickettsia*, *Arsenophonus*, *Spiroplasma* infected samples. We plotted up to three samples for every species infected with *Wolbachia* to show the range of *Wolbachia* across tested arthropod species. These plots show us that titer variation within host species is significant, and this variation is not due to pooled sequencing samples. Our results suggest a symbiont and host genetic contribution to shaping within-host infection densities.

Table 1. Raw infection counts of *Wolbachia, Spiroplasma, Rickettsia* and *Arsenophonus* infecting arthropods and nematodes. The frequency of positive samples or species is listed in parentheses.

	Samples		Species	
	Arthropoda	**Nematoda**	**Arthropoda**	**Nematoda**
Total	27256	5229	1299	128
Infected With Any RM	2083 (0.08)	119 (0.02)	240 (0.18)	8 (0.06)
Infected With *Wolbachia*	1978 (0.07)	119 (0.02)	179 (0.14)	8 (0.06)

Table 2. Estimated infection frequencies and confidence intervals from our data for *Wolbachia, Spiroplasma, Rickettsia* and *Arsenophonus* infecting arthropods and nematodes. All species in dataset were downsampled to maximum 100 individuals. We used a minimum threshold of 1 in 1000 of infected individuals within a species to classify a species as positively infected (Supplementary Method S3).

Host Phylum	Reproductive Manipulator	2.5% CI	97.5% CI	Median	Mean
Arthropods	*Arsenophonus*	0.0005	0.05186	0.0081	0.0124
	Rickettsia	0.0137	0.0822	0.0360	0.0389
	Spiroplasma	0.0329	0.1571	0.0778	0.0826
	Wolbachia	0.2902	0.6170	0.4411	0.4414
Nematodes	*Wolbachia*	0.0499	0.6917	0.3291	0.3429

Table 3. *Wolbachia* global infection frequencies and confidence intervals generated for arthropod orders. All species in dataset were downsampled to maximum 100 individuals. Confidence intervals were generated using 1000 bootstrap replicates fitting a beta-binomial model to species infection frequency data among orders. A minimum infection frequency of 0.001 was used to classify a species as positively infected (Supplementary Method S3).

Order	2.50%	97.50%	Median	Mean
Coleoptera	0.5272	0.9593	0.7940	0.7787
Diptera	0.3002	0.8983	0.6410	0.6333
Hymenoptera	0.1174	0.7727	0.4084	0.4236
Hemiptera	0.1719	0.9950	0.5118	0.5475
Lepidoptera	0.0928	0.7040	0.3503	0.3640