Tumor fitness, immune exhaustion and clinical outcomes: impact of immune checkpoint inhibitors

Adrian Bubie(1), Edgar Gonzalez-Kozlova(1), Nicholas Akers(2), Augusto Villanueva (3), Bojan Losic* (4)

1: Department of Genetics and Genomic Sciences ,Icahn Institute for Data Science and Genomic Technology, Icahn School of Medicine at Mount Sinai, New York, NY 10029

2: Adaptive biotechnologies, 1551 Eastlake Avenue E, Suite 200, Seattle WA 98102

3: Division of Liver Diseases, Department of Medicine, Liver Cancer Program, Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, USA.

4: Department of Genetics and Genomic Sciences, Tisch Cancer Institute, Cancer Immunology, Diabetes, Obesity and Metabolism Institute, Icahn Institute for Data Science and Genomic Technology, Icahn School of Medicine at Mount Sinai, New York, NY 10029

*Correspondence to: bojan.losic@mssm.edu

Abstract

Recently proposed tumor fitness measures, based on profiling neoepitopes for reactive viral epitope similarity, have led to improved prediction models of response to immune checkpoint inhibitors in melanoma and small-cell lung cancer. Here we apply these checkpoint based fitness measures in the matched checkpoint-treatment naive TCGA samples where cytolytic activity imparts a known survival benefit. No significant survival predictive power beyond that of overall patient tumor mutational burden is observed in either cohort, and furthermore, there is no observed association between checkpoint based fitness and tumor T-cell infiltration, cytolytic activity (CYT), or abundance (TIL burden). We investigate the key assumption of viral epitope similarity driving immune response in the hepatitis B virally infected liver cancer TCGA cohort, and uncover suggestive evidence that tumor neoepitopes dominate viral epitopes in putative immunogenicity and plausibly drive immune response.

Significance: Tumor fitness does not predict patient survival better than tumor mutation burden, unlike a measure of CYT. In checkpoint treatment naive patients, our results suggest TIL recruitment and cytolytic activity are essentially driven by tumor neoepitopes, regardless of the presence of viral epitopes.

Introduction

Quantifying tumor fitness modulation by immune selection pressure has assumed a new urgency given the growing need to predict immunotherapy efficacy (1,2). Immune-based anti-tumoral response is driven in part by T-cell receptor (TCR) recognition of tumor neoantigens, which mainly represent mutated protein epitopes (i.e., neoepitopes) derived from somatic mutations. These neoantigens are presented by host-cell major histocompatibility complex (MHC) molecules to facilitate 'self' recognition by TCRs (**Fig. 1b**). Cancer clones that escape immune recognition can hijack CTLA-4 and PD-1 checkpoint blockade, resulting in heterogeneous tumors with a complex immunogenomic

profile which may include immune exhaustion, giving rise to mixed or unfavorable response to immune checkpoint inhibition (1).

Recently, a tumor fitness model has been proposed that incorporates two key components defining the TCR-neoantigen interaction: putative neoantigen immunogenicity and TCR recognition (3). Putative immunogenicity is modeled via a nonlinear function of *in silico* derived MHC-I binding affinities arising from the clonal somatic mutation spectrum, while the TCR recognition likelihood is set to scale with sequence similarity to known, infectious, virally derived epitopes from the Immune Epitope Database (IEDB) (19). Given these assumptions, a combined measure of tumor fitness is then defined as the inverse of the maximum clonal immunogenicity potential Ï weighted across subclones in the tumor (see Methods, **Fig. 1a**).

Although this model successfully predicts response to checkpoint inhibition in melanoma and small cell lung cancer, it is of interest to investigate its applicability, positive predictive power, and key assumptions in the arena of the analogous immunotherapy naive (endogenous) TCGA SKCM and NSCLC cohorts. Crucially, even though checkpoint predictive biomarkers are not necessarily prognostic, substantial evidence has already accumulated about the positive survival effects of cytolytic immune activity in melanoma and lung cancer (11,20), strongly suggesting immune selection pressure significantly modulates treatment naive tumor fitness. Our central hypothesis is that the predictive tumor fitness model of checkpoint response will predict survival in cohorts where cytolytic activity imparts a significant survival benefit. Indeed, the tumor fitness model (3) makes no mathematical distinction between endogenous or checkpoint-induced immune response.

An important corollary to our central hypothesis is to directly test a key assumption of the Luksza model (3), namely that of viral similarity driving differential immune recognition of neoepitopes. We hypothesized that in the setting of the simultaneous presence of viral epitopes and neoepitopes, i.e. in a virally infected tumor, we could directly determine if the predicted MHC-I binding affinity (putative immunogenicity) of viral peptides was higher, lower, or indistinguishable from that of tumor neopeptides from the same patients. Developing and deploying new in-silico methods, we carried this test out using the hepatitis B virally positive portion (HBV+) of the LIHC TCGA liver cancer cohort.

In this brief report we show that there is no significant correlation between tumor fitness and immune response signatures, nor is patient survival predicted using tumor fitness better than with ordinary tumor mutation burden (TMB), in either SKCM or NSCLC cohorts. Indeed we show that, as demonstrated in the checkpoint treatment setting (23), an expression signature capturing tumor-infiltrating T-cell activity (CYT) significantly outperforms TMB in predicting survival in SKCM, and nearly so in NSCLC. We also show that tumor neoantigens actually *dominate* viral

antigens in average putative immunogenicity within the same patients, and also better correlate with immune response, in the HBV+ HCC LIHC cohort.

Results

CYT, not Tumor fitness, correlates with immune measures in NSCLC and SKCM

We selected 305 patients with non-small cell lung cancer (NSCLC) and 337 with skin cutaneous and melanoma (SKCM) in the TCGA with matched whole exome sequencing (WES) and RNA-seq data (**Fig. 1a**). Putative tumor neoepitopes (ic50 < 500nM) with significant homology with viral sequences (e-value < 10) from the IEDB (see Methods) were found in 204/337 (60%) of SKCM and 221/306 (72%) of NSCLC patients. We found that SKCM patients with viral-like epitopes had a significantly worse survival, and correlated with lower TMB (but not CYT), while no survival associations were observed for NSCLC (**Supp. Fig. 3**). We observed significant associations between CYT and TIL burden, for both the NSCLC and SKCM cohorts (**Fig. 2a,b**), as previously reported in patients with these tumor types treated with checkpoint inhibitors (23). Significantly, however, we found that maximal clonal immunogenicity Ï (see Methods) does not correlate with TIL burden or CYT in either cohort (**Fig 2a,b**).

Tumor immunogenicity, cytolytic immune activity, and patient survival

Consistent with previous reports of strong predictive power of CYT in checkpoint treated patients (23), we found that the gene signature of immune cytolytic activity (CYT) was significantly associated with survival in the non-checkpoint treated patients in SKCM and NSCLC (**Fig. 3b,d**). In order to assess if tumor fitness predicted survival in these cohorts, we computed maximal clonal tumor fitness scores in each patient (using the same method, software versions, and parameter specifications as Luksza et al., see Methods) and found that they did not separate overall survival in either SKCM or NSCLC cohorts (**Fig. 3a,c**). As observed in other studies (4), it is challenging for neoepitope based techniques to predict patient survival better than TMB alone, particularly when regressing out powerful covariates such as clinical tumor stage. To establish the relative predictive power of tumor fitness, we constructed a series of multivariate Cox proportional hazard models for each cohort which included TMB, tumor stage, tumor fitness (Ï), and CYT. For each model, we computed the time dependent Brier score (see Methods; briefly, the weighted average of the squared distances between the observed survival status and the predicted survival probability of a model) and compared the resulting prediction error curves. We compared the relative predictive power of different regression models via bootstrap resampling (see Methods). As shown in Figure 3, tumor fitness does not significantly reduce the prediction error in NSCLC nor SKCM compared to models with TMB alone, while CYT does so in SKCM

but not in NSCLC (**Fig. 3e,f**). Computing the integrated Brier scores under cross-validation confirms these results are robust (see supplementary materials).

Neoepitopes, not viral epitopes, drive immune response in HBV-related liver cancer

To examine the relationship between TCR recognition and neoepitope sequence similarity to pathogen derived epitopes in samples that coexpress both, we examined patients with hepatocellular carcinoma co-infected with HBV from the LIHC TCGA dataset. We assessed 190 patients with matched WES and RNA-Seq data, detecting HBV transcripts in 115/190 (60%) via HBV genome assembly from RNAseq data unmapped to human genes (see Methods). Of these, 88 patients produced HBV matching protein sequences through ORFfinder. Using the RNA-seq data for patient HLA typing, we estimated the distribution of binding affinities of the viral antigens and compared it with that of neoantigens produced by tumor mutations. A total of 133,834 unique tumor neoepitopes and 30,357 HBV viral epitopes were predicted across the patients. Cumulative distribution functions of the predicted MHC-I binding affinities indicate a significant MHC-I binding affinity preference for tumor derived neoantigens over that of HBV viral cofactors (**Fig. 1b, 4**). This result is robust against subsampling among neopeptides (see Methods and **Supp. Fig. 1**). In addition, we found that tumor neoepitope burden is lowly but significantly associated with patient TIL burden (Spearman's ρ=0.24, p=0.038), but HBV epitope burden was not (restricting to MHC-I strong binders [ic50 < 200 nM]; **Supp. Fig. 2**), further suggesting that TIL recruitment is preferentially driven by relative neoepitope burden compared to viral burden. Our observation forms a counterexample to the model assumption that viral sequence similarity drives preferential epitope recognition.

Discussion

Determining the degree of 'non-selfness' of mutant peptides in the face of enormous individual variability of TCR repertoires is notoriously difficult, and yet critical, to answer at scale. Point somatic mutations located outside the MHC-binding anchor positions may appear to be more 'non-self' since they may lead to significantly increased peptide-MHC binding compared with wildtype sequence. However it has traditionally not been straightforward to translate this notion into a better survival or treatment response predictor. The relative spectra of subclonal and clonal immunogenic mutations, and similarly mutant sequence homology with experimentally validated pathogen epitopes, potentially contain important tumor evolution signatures, which may indicate intriguing similarities with pathogen evolution.

To gain further intuition into a recent neoepitope based tumor fitness scoring model (3), we applied it to samples from checkpoint-treatment-naive samples of the same tumor types in the TCGA. A key aim was to test how this model predicts survival given that it has shown promise in predicting response to immune checkpoint inhibitors. Although this setting may involve substantially different (thus far uncharacterized) tumor-immune dynamics, we reasoned that a sufficiently general notion of tumor fitness should correlate with patient survival, especially in those cohorts where TIL and CYT are already known as strong survival correlates.

Our results suggest that the proposed tumor fitness measure based on pathogen epitope similarity does not correlate with CYT in either TCGA NSCLC nor SKCM, potentially indicating that it is specific to the patients treated with checkpoint inhibitors. Furthermore, in a test of a key model assumption, in the case of HBV-related HCC in the TCCA cohort, we find that even when viral epitopes are competing with neoantigens for T cell presentation there is actually a substantial average bias towards higher MHC-I binding affinity for the neoepitopes. In addition, a significant correlation between TMB and TIL burden further suggests that the tumor dominates immune recruitment over HBV viral epitopes. It is interesting to speculate on whether other tumors with known viral cofactors, such as cervical or head and neck with Human Papillomavirus, and Epstein Barr virus and Burkitt's lymphoma, show similar biases.

Furthermore, we find that tumor fitness in the treatment-naive samples does not improve prediction error of survival beyond that of TMB and clinical tumor staging alone, while a simple measure of CYT does so in SKCM. The tumor fitness model has two key degrees of freedom that could very likely be tuned to reduce this prediction error, as they were for patients treated with checkpoint inhibitors (3), but it is instructive to simply note here that these choices do not work well in untreated samples. While these data suggest that the tumor immune dynamics in treated patients are notably different than in untreated patients, it may also simply reflect a very cancer or dataset specific sensitivity of the tumor fitness score. Indeed, as noted, the CYT measure alone seems to reduce survival prediction error, as found by others (21,22) and model authors themselves as a useful component to add to the tumor fitness score (3). Though the improvement of survival prediction by the inclusion of CYT may be, essentially, expected based on its efficacy as an independent predictor of therapy responsiveness in the checkpoint-setting (23), we demonstrate precisely in our work here this common-sense expectation to be true in the non-checkpoint setting as well.

Although our study points out key limitations in extending the paradigm of tumor fitness scoring to untreated patients, it does have the same caveats any in-silico immunogenicity estimation based analysis method has. Indeed, the accuracy of in-silico predictions of neoantigen or viral binding affinity in estimating immune reactivity are at best suboptimal compared to using mass spectroscopy. Also, in comparing viral and neo-epitopes it's important to note that several lines of evidence suggest that the replication status of HBV is dependent on the differentiation status of the cells (15,16). Thus, the virus may hardly replicate in HCC cells and further reduce the viral epitope burden within the tumor.

Overall our study indicates that while tumor fitness scores can be used to predict response to checkpoint inhibitors, these measures do not seem to predict survival in patients not treated with checkpoint inhibitors where endogenous immune activation is a significant survival factor. It is important to note that while this tumor fitness measure and its underlying assumptions have been adapted with positive predictive results in other contexts (24,25), we find that the exact form of the model is not predictive in these endogenous cases. Nevertheless, it is encouraging to observe that CYT measures are powerful in both treated and untreated patients, which may point to the existence of universally predictive and prognostic immune activation signatures.

Methods

TCGA analysis

RNA-Seq data for the TCGA patient cohorts was aligned to Hg38 with STAR (v2.5.1b) in two-pass mode (6). Gene counts for Gencode v23 (www.gencodegenes.org) gene annotations were generated using featureCounts. Read counts underwent TMM normalization and logCPM transformation using voom (9). Mapped reads were used to allelotype (MHC class-I loci) each patient (7), estimate the putative TIL burden per patient by profiling TCR and BCR sequences with MiXCR (8), and normalizing by patient library size. To predict neoantigen and associated epitope burden, we used Topiary (10) to call mutation-derived cancer T-cell epitopes from somatic variants, tumor RNA expression data, and patient class I HLA type. This tool matches mutations with gene annotations, filters out non-protein coding changes, and finally creates a window around amino acid changes, which is then fed into NetMHCCons for each patient HLA allele across tiles of 9 - 12 aa in length. Given that HLA-I processes neoantigens by degradation to non-conformational 8-11 amino-acid residues, we excluded neoepitopes with mutations obscured to T-cells within HLA-I binding pockets. In the case of frameshift mutations, in principle this window starts from the mutation minus the length of the peptide up to the first stop codon. Predictions were filtered by a binding affinity threshold of ic50 < 10000nM. A measure of tumor mutation burden (TMB) for each patient was calculated by enumerating the number of called DNA-mutations in patient associated MAF files.

Tumor fitness score

We assigned tumor immunogenicity scores following the method outlined by the model described in (3). Briefly, a protein BLAST of the neoepitopes from the SKCM and NSCLC cohorts was performed per patient against a database of viral epitopes sourced from the International Epitopes Database (IEDB)(19); the best hits with higher sequence similarity normalized by sequence database size (bitscore) for each neoepitope were retained. A second, randomized set of neoepitopes was also blasted, as a negative control group. Smith-Waterman alignments scores ($|s,e|$)

were calculated for each match, following a strict 500nM binding affinity threshold filtering for all predicted neoepitopes. For each patient, a TCR recognition potential (R) was calculated by modeling the probability for a given neoepitope to be bound to a viral match from the IEDB using:

$$R = \frac{\sum\limits_{e \in IEDB} exp(-k(\alpha - |s,e|))}{1 + \sum\limits_{e \in IEDB} exp(-k(\alpha - |s,e|))} \tag{1}$$

with constants α (binding curve displacement) and k (curve steepness) retained from the original model in matching to each TCGA patient cohort. For each neoantigen, we also calculated an MHC-I binding affinity amplitude score (A), as a ratio of the binding affinities of the mutant neoepitope and the wild type epitope, which were corrected for dissociation constant bias by multiplying by the reciprocal max among all wild type epitopes (3). Tumor immunogenicity was assigned by selecting the maximum product of these scores from among all neoantigens across patient clones, j:

$$Ï = max\left(R_j A_j\right) \tag{2}$$

This term represents the effective expected size of immune response and is used to rank the dominant immunogenic clone per patient. The inverse of this score, tumor fitness, describes the likelihood of evading immune detection.

Cytolytic T-Cell activity (CYT) scores were calculated using the geometric mean of the FPKM (fragments per kilobase mapped) expression of granzyme (*GZMA*) and perforin-1 (*PRF1*) (11). Spearman correlations between patient tumor immunogenicity scores and CYT were calculated in R.

HBV antigen calling

Raw RNA sequencing reads that did not map to the GRCh38 reference genome was assembled into contigs using Trinity (using *−no_run_chrysalis −no_run_butterfly flags*, which only invokes the Inchworm step) to perform greedy kmer-25 contig assembly (12). Resulting contigs with a sufficiently high entropy (to exclude homopolymer sequences), at least 100 bp long and supported by at least 20 reads were retained and aligned with BLAST against the HBVdb (17) reference genomes (e-value < 1e-6) for HBV (genotypes A, B, C, D, E, F, and G), effectively identifying the HBV positive patients (n=115). CD-Hit was used to consolidate highly similar contig sequences (e-values < 1e-10) (13). Finally, ORFfinder was used to predict the putative HBV protein products from each contig, which were validated and filtered (e-value < 1e-10) by a protein BLAST against an HBV protein database compiled from HBVdb for the HBV genotypes above (14). Importantly, our contig length filtering means that ORFs shorter than 300 aa are excised.

HBV epitopes were predicted using the matching protein products and patient HLA-typing calls with Topiary (see above), and peptides were filtered by binding affinity (ic50<10000nM). Comparison of viral and neo-epitope MHC-I binding affinity distributions was performed using a one-sided Kolmogorov-Smirnov test.

Additionally, to assure that the MHC-I binding affinity bias toward tumor neoepitopes was not confounded by a larger epitope set, we subsampled 10000 epitopes from both populations and compared binding affinity distributions by one-sided (neoepitope greater) Kolmogorov-Smirnov test for 1000 iterations (**Supp. Fig. 1**).

Survival Analysis

Univariate (Kaplan-Meier) curves and risk tables using patient tumor immunogenicity, Ï, and immune activity, CYT, were created with the ggsurvplot package. Patient grouping into low and high tumor immunogenicity and immune activity was done using the bottom 15% and top (85%) 15% quantiles for these measures, respectively. A log-rank test was used to evaluate significance in predicted survival outcomes between groups. The relative predictive power of various survival models was evaluated using the 'pec' R package by computing the Integrated Brier Score (IBS) across 632+ bootstrap resampling with 100 iterations (B=100). We used the Kaplan-Meier estimator for the censoring times in the bootstrap simulations. The weights in the IBS correspond to the probability of not being censored. Model error curves were compared using Wilcoxon test for IBS comprising approximately 80% of total patients, with test significance provided in graphic tables (**Figure 3e,f**). IBS significance testing was performed by sampling Brier scores with 100 iterations across random patient groups for each model and applying a one-sided (*TMB_I* less) Kolmogorov-Smirnov test.

Availability of Code and Data: All code and non-protected data is available upon reasonable request to the corresponding author.

References

1. Sharma, P., Hu-Lieskovan, S., Wargo, J. A. & Ribas, A. Primary, Adaptive, and Acquired Resistance to Cancer Immunotherapy. Cell 168, 707–723 (2017).
2. Riaz, N. et al. Tumor and Microenvironment Evolution during Immunotherapy with Nivolumab. Cell 171, 934-949.e16 (2017).
3. Łuksza, M. et al. A neoantigen fitness model predicts tumour response to checkpoint blockade immunotherapy. Nature 551, 517–520 (2017).
4. Havel, J. J., Chowell, D. & Chan, T. A. The evolving landscape of biomarkers for checkpoint inhibitor immunotherapy. Nat. Rev. Cancer 19, 133–150 (2019).

5. Rizvi, N. A. et al. Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. Science 348, 124–128 (2015).

6. Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15–21 (2013)

7. Szolek, A. HLA Typing from Short-Read Sequencing Data with OptiType. Methods Mol. Biol. 1802, 215–223 (2018)

8. Bolotin, D. A. et al. MiXCR: software for comprehensive adaptive immunity profiling. Nat. Methods 12, 380–381 (2015)

9. Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., Smyth, G. K. limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research, 7, e47 (2015)

10. Rubinsteyn, A., Nathanson, T. Topiary: predict mutated T-cell epitopes from sequencing data. Github repository, https://github.com/openvax/topiary (2018).

11. Rooney, M. S., Shukla, S. A., Wu, C. J., Getz, G. & Hacohen, N. Molecular and genetic properties of tumors associated with local immune cytolytic activity. Cell 160, 48–61 (2015).

12. Haas, B. J. et al. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. Nat. Protoc. 8, 1494–1512 (2013).

13. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-generation sequencing data. Bioinformatics 28, 3150–3152 (2012); Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformatics 22, 1658–1659 (2006).

14. Rombel, I. T., Sykes, K. F., Rayner, S. & Johnston, S. A. ORF-FINDER: a vector for high-throughput gene identification. Gene 282, 33–41 (2002).

15. Halgand, B., Desterke, C., Riviere., L., Fallot, G., Sebagh, M., Calderaro, J., et al. Hepatitis B Virus Pregenomic RNA in Hepatocellular Carcinoma: A Nosological and Prognostic Determinant. Hepatology, 1, 86-96 (2018).

16. Fu, S., Li, N., Zhou, P. C., Huang, Y., Zhou, R. R., Fan, X. G. Detection of HBV DNA and antigens in HBsAg-positive patients with primary hepatocellular carcinoma. Clin. Res. Hepatol. Gastroenterol. 4, 415-423 (2017).

17. Hayer, J., Jadeau, F., Deleage, G., Kay, A., Zoulim, F., Combet, C. HBVdb: a knowledge database for Hepatitis B Virus. Nucleic Acids Research, D1, D566-D570 (2013).

18. Mogensen, U. B., Ishwaran, H., Gerds, T. A. Evaluating Random Forests for Survival Analysis using Prediction Error Curves. J. Stat. Softw. 50,1-23 (2012) URL: http://www.jstatsoft.org/v50/i11/

19. Vita, R., Mahajan, S., Overton, J. A., Dhanda, S. K., Martini, S., et al. The Immune Epitope Database (IEDB): 2018 Update. Nucleic Acids Res. (2018).

20. Barnes, T. A., Amir, E. HYPE or HOPE: the prognostic value of infiltrating immune cells in cancer. Br J Cancer. 4,451-460 (2017)

21. Wakiyama, H., Masuda, T., Motomura, Y., Hu, Q., Tobo, T., Eguchi, H., Sakamoto, K., et al. Cytolytic Activity (CYT) Score is a Prognostic Biomarker Reflecting Host Immune Status in Hepatocellular Carcinoma (HCC). Anticancer Res. 12, 6631-6638 (2018).

22. Roufas, C., Chasiotis, D., Makris, A., Efstathiades, C., Dimopoulos, C., Zaravinos, A. The Expression and Prognostic Impact of Immune Cytolytic Activity-Related Markers in Human Malignancies: A comprehensive Meta-analysis. Front. Oncol. 8, 27 (2018).

23. Cristescu, R., Mogg, R., Ayers, M., Albright, A., Murphy, E., Yearly, J., et al. Pan-tumor genomic biomarkers for PD-1 checkpoint blockade-based immunotherapy. Science 362 (2018).

24. Zhang, J., Caruso, F. P., Sa, J. K., Justesen, S., Nam, D.-H., Sims, P., et al. The combination of neoantigen quality and T lymphocyte infiltrates identifies glioblastomas with the longest survival. Comm. Biol. 2(1), 135 (2019).

25. Wood, M. A., Paralkar, M., Paralkar, M. P., Nguyen, A., Struck, A. J., Ellrott, K., et al. Population-level distribution and putative immunogenicity of cancer neoepitopes. BMC Cancer. 18(1), 414 (2018).

Figures:

Figure 1: A) Workflow summary. Matched mutation and RNA-seq TCGA datasets underwent HLA-Typing, neoepitope calling, and VDJ-alignment for each cohort, and tumor fitness scores calculated for those in the SKCM and NSCLC (3). Unmapped reads from the LIHC cohort were mined for HBV reads, which were used to predict viral epitopes. **B)** T-cell recognition by TCR of a cancer cell by MHC-I presentation. Antigens bound to the MHC can displace other antigens due to higher binding affinity with the MHC molecule. This competitive binding affinity is represented by an ic50 score.

Figure 2: Maximal clonal immunogenicity (inverse tumor fitness), Ï, was found to be lowly and non-significantly correlated (Spearman) with all our measures of expression-based tumor immune response (TIL burden, TIL clonality and CYT) and TMB in both **A)** NSCLC, **B)** SKCM patients. Several of the immune measures were found to be highly correlated with each other in both TCGA patient sets. Values in each circle indicate the spearman correlations between the features, with color and size representative of correlation strength. Tiles with no color indicate a non-significant (p > 0.05) correlation between features.

Figure 3: Kaplan-Meier survival curves between patients grouped by highest and lowest immunogenicity (*I*) and immune activity marker expression (*CYT*) scores for the **A,B)** lung (NSCLC) and **C,D)** melanoma (SKCM) TCGA cohorts. Quantiles of *I* and *CYT* patient scores used to group cohorts with the lowest and highest groups represented by the curves; **E,F)** Time dependent Brier scores (prediction error curves) comparing multivariate cox proportional hazard survival models for the NSCLC cohort (e), and SKCM cohort (f). Reference is intercept, all other models include tumor stage covariates. TMB includes patient tumor mutation burden, TMB_I includes immunogenicity and TMB, CYT_TMB includes CYT and TMB. Vertical lines demarcate approximately 80% of the patients, for which integrated Brier scores were calculated.

Figure 4 : Distribution of MHC-I binding affinities for tumor neoantigens (blue) and HBV antigens (red). Smaller ic50 values indicate a stronger binding affinity. Tumor neoantigens are on average more likely to bind to patient MHC-I alleles than antigens from HBV proteins.

Supplemental Figures:

Supp. Figure 1: Kolmogorov-Smirnov test results (**A**) of randomly subsampled distributions of tumor neoantigen binding affinities compared to HBV antigens across 1000 iterations. **B)** Distribution of test p values strongly support the alternative hypothesis that tumor neoantigen cumulative distributions were significantly less than HBV distributions, indicating better average binding affinity scores are robust to subsampling.

Supp. Figure 2: Spearman correlations of antigen burden with mutation burden in HBV+ HCC LIHC patients. Tumor neoantigen burden is slightly but significantly positively correlated with patient TIL-Burden, while HBV antigen load is not.

Supp. Figure 3: Kaplan-Meier survival curves between patients with and without neoepitope matches against the viral IEDB database for the **A)** lung and **B)** melanoma TCGA cohorts. Tumor Mutational burdens for patients with viral neoantigen matches fall significantly below those with no matches in both the **C)** lung and, **D)** melanoma patient groups (two sided T-test). Differences in cytolytic activity (CYT) expression were found to be negligible in both the NSCLC and SCKM patients between those with and without IEDB matches (**E** and **F**, respectively)