Human Endometrial Transcriptome and Progesterone Receptor Cistrome Reveal

Important Pathways and Epithelial Regulators

Ru-pin Alicia Chi[1], Tianyuan Wang[2], Nyssa Adams[3], San-pin Wu[1], Steven L. Young[4],

Thomas E. Spencer[5,6], and Francesco DeMayo[1]

[1] Reproductive and Developmental Biology Laboratory, National Institute of

Environmental Health Sciences, Research Triangle Park, North Carolina, USA

[2] Integrative Bioinformatics Support Group, National Institute of Environmental Health

Sciences, Research Triangle Park, North Carolina, USA

[3] Interdepartmental Program in Translational Biology and Molecular Medicine, Baylor

College of Medicine, Houston, Texas, USA

[4] Department of Obstetrics and Gynecology, University of North Carolina at Chapel Hill,

Chapel Hill, North Carolina, USA

[5] Division of Animal Sciences and [6]Department of Obstetrics, Gynecology and Women's

Health, University of Missouri, Columbia, Missouri, USA

Disclosure summary: The authors have nothing to disclose.

ABSTRACT

The endometrium undergoes highly dynamic modifications in a time and compartment specific

manner during each menstrual cycle resulting in receptivity to embryo implantation during a

discrete temporal window of implantation (WOI). Attainment of receptivity is strictly governed by

the steroid hormone progesterone, which acts via the two isoforms of its nuclear Progesterone

Receptor (PGR). In order to define the molecular mechanisms regulated by PGR in uterine

receptivity, we conducted RNA-sequencing (RNA-seq) and PGR ChIP-sequencing (ChIP-seq)

in endometrial biopsies taken from fertile women during the proliferative (P) and mid-secretory

(MS) phases. Overlaying the genes with altered PGR binding and differential expression

(DEGs) during the phase transition identified 653 genes, which are involved in inflammatory

response signaling, xenobiotic metabolism, epithelial-mesenchymal transition (EMT), cell death

regulation, interleukin/STAT signaling, estrogen response, as well as MTORC1 response. RNA-

sequencing conducted using epithelium-derived RNA identified 3,052 DEGs in the epithelium, of

which 658 were uniquely regulated in the epithelium. Transcription factors IRF8 and MEF2C

were selected for validation and found to be regulated in the epithelium during the WOI at the

protein level, suggesting potentially important functions that are previously unrecognized. In

summary, we present herein data that expands our understanding of progesterone action in

human endometrium during the WOI.

Introduction

The human endometrium is a highly complex tissue. The functionalis layer consists of the stromal compartment which makes up significant portion of the endometrium; the glandular epithelium which is responsible for secreting an array of growth factors and cytokines [1]; and the luminal epithelium which lines the stromal compartment and is the first maternal interface with which the embryo interacts inside the uterus. In order to maximize the chances of a successful pregnancy, the uterus prepares for embryo implantation after each menstruation by the generation and differentiation of the endometrial functionalis, a process known as the menstrual cycle [2, 3]. This is orchestrated by the interplay of two steroid hormones, estrogen and progesterone. During the proliferative (P) phase, estrogen promotes proliferation of both the stromal and epithelial cells, steadily increasing the thickness of the functionalis [4, 5]. Upon ovulation, the ovary begins secreting progesterone, halting estrogen-induced proliferation and initiating differentiation of stromal cells (decidualization) and epithelial cells. These include depolarization, altered surface morphology, expression of specific adhesion proteins, altered steroid receptor expression, and secretion of glycogen [5, 6]. Without a successful implantation, the levels of both steroid hormones decrease during the late secretory phase, leading to endometrial involution and subsequently endometrial shedding (menstruation), initiating another cycle [7].

Abnormal embryo implantation and implantation failure are major causes of infertility and early pregnancy loss, which is linked to other pregnancy complications [8-12]. Attainment of human endometrial receptivity occurs in the mid-secretory phase (MS) after sufficient time and concentration of progesterone exposure as seen in other placental mammals [13-18]. In women

without ovaries, sequential treatment with estrogen followed by estrogen plus progesterone,

without any other ovarian hormones, is sufficient to achieve high rates of successful

implantation of embryos derived from donor oocytes [16, 19], highlighting the mportance of

hormone actions in mediating implantation.

Abnormal progesterone signaling leads not only to fertility issues but also a spectrum of

gynecological diseases [20-22], emphasizing the criticality of progesterone signaling in

maintaining normal uterine biology and initiating pregnancy. The impact of progesterone is

mediated through its nuclear receptor – Progesterone Receptor (PGR), where binding of

progesterone induces its conformational change. This leads to altered affinity for target DNA

response elements, thereby influencing the gene expression network at the transcriptional level

[23]. Although many PGR-regulated genes have been identified in both animal model systems

and human studies as important mediators of implantation, including Indian Hedgehog (*IHH*)

[24-26], Krüpple-like Factor 15 (*KLF15*) [27, 28], Heart and Neural Crest Derivatives-expressed

2 (*HAND2*) [29], Bone Morphogenesis Protein 2 (*BMP2*) [30, 31], Homeobox gene *HOXA10* [28,

32, 33], and CCAAT/Enhancer-binding Protein β (*CEBPB*) [34-36]. Yet, implantation failure

remains a great challenge in both natural pregnancies and assisted reproductive interventions.

Additionally, epithelial aspects of PGR actions are important, sometimes underappreciated

determinants of implantation and pregnancy outcome. Endometrial epithelial cells line the

uterine lumen and glands, with the latter derived from the former [37, 38]. The endometrial

epithelium undergoes dramatic cellular and molecular changes common to both mice and

humans during the WOI, including adhesion mechanisms enabling the attachment of embryo to

the luminal epithelium [39, 40], alterations in nuclear pore complex presentation [41],

downregulation of the Serum and Glucocorticoid Regulated Kinase 1 (SGK1) [42], apoptotic

cascade [43, 44], and expression of epithelial-specific receptivity markers [45]. The glandular

epithelium further facilitates implantation via the production of Leukemia Inhibitory Factor (LIF),

a critical factor in embryo-uterine communication during WOI [46-48]. Elaborate cross-talk exists

between the endometrial epithelium and stroma that is indispensable for allowing implantation,

adding further complexity to the regulatory mechanisms governing pregnancy establishment.

Although animal model systems and *in vitro* cultured cells have proven instrumental in

advancing our knowledge in reproductive functions, the high rate of implantation failure remain a

challenge [49]. The aim of this study is to use a single, comparative, human-derived, *ex vivo*

analysis to examine the dynamics of PGR action during the WOI. We employed ChIP-seq

technique to explore the modification of PGR binding landscape during the P to MS transition in

human endometrial samples. Additionally, parallel RNA-sequencing analysis enabled the

identification of differentially regulated genes, which allowed us to identify the subset of PGR-

bound genes with altered mRNA abundance and hence relevance in regulating implantation and

decidualization. Epithelial-specific RNA-sequencing allowed more precise assessment of the

endometrial epithelial transcriptomic network, providing a deeper understanding of the dynamic

transformation in the endometrium during the WOI.

Results

Characterization of PGR binding trend during the P and MS phases

To gain insights into the transcriptional regulatory function of PGR during the peri-implantation period, the physical association of PGR with DNA was assessed by PGR chromatin immunoprecipitation coupled to massively parallel DNA sequencing (ChIP-seq) using human endometrial biopsies from the P and MS phases. We identified over 10,000 genomic intervals (defined as a stretch of DNA sequence identified as exhibiting stastically significant PGR binding) as PGR bound in the endometrium. Analysis using the Peak Annotation and Visualization tool showed that majority of the PGR binding occurred within the intronic, intergenic, 5' UTR and upstream region relative to the gene body, with no significant alteration in PGR binding preference to these categories between the two phases (Fig. 1. A).

Then, we characterized the PGR binding dynamics by identifying intervals with consistent or differential PGR binding (DPRB). Collectively, we analyzed two sets of samples each containing a P and MS pair. To circumvent the batch variation observed between the two sets of samples, we defined the consistent/constitutive PGR binding sites as those with PGR binding during both P and MS, where the read counts were not significantly different between P and MS in either one or both batches. For the DPRB intervals, we first analyzed each set independently to identify differential PGR binding sites, and only those DPRB common to both datasets were considered for additional analyses. In total, we identified 12,469 genomic sites with consistent PGR binding in proximity to 11,058 genes (Supplemental Table 1); and 2,787 genomic sites with altered PGR binding in proximity to 2,249 genes (Fig. 1. B, Supplemental Table 2). There were 2,466 intervals with increased PGR binding in proximity to 1,966 genes (88%) and 321

intervals with decreased PGR binding in proximity to 307 genes (12%, Fig. 1. C), and 423 genes

were found with multiple differential PGR binding intervals in proximity.

Amongst the identified DPRB intervals, many were found in proximity to known PGR-regulated

genes previously reported in both humans and mice, including FK506 Binding Protein 5

(*FKBP5*) [28], Indian Hedgehog (*IHH*), Insulin Receptor Substrate 2 (*IRS2*) [50], *CASP8* and

FADD Like Apoptosis Regulator (*CFLAR*) [51], FOS Like 2 AP-1 Transcription Factor Subunit

(*FOSL2*) [28], Perilipin 2 (*PLIN2*), Basic Leucine Zipper ATF-Like Transcription Factor (*BATF*)

and Baculoviral IAP Repeat Containing 5 (*BIRC5*, Supplemental Table 2) [21]. In addition, many

known decidualizing and implantation mediators were found with constitutive PGR binding,

including Forkhead Box Protein O1 (*FOXO1*) [52], Homeobox A10 (*HOXA10*) [52], Heart And

Neural Crest Derivatives Expressed 2 (*HAND2*) [2], Cysteine Rich Angiogenic Inducer 61

(*CYR61*) [28] and Sex Determining Region Y-Box 17 (*SOX17*, Supplemental Table 1) [53, 54].

The biological impact of PGR transcriptional activity during the P to MS phase was determined

by examining the functional profile associated with the DPRB genes using the DAVID

Bioinformatics Database [55, 56], and selected enriched pathways are shown in Table 1.

Enrichment was observed in pathways regulating insulin resistance, focal adhesion,

complement and coagulation cascades, cytokine-cytokine receptor interactions, ECM receptor

interaction, apoptosis, as well as various signaling pathways including chemokines, Ras, FOXO,

Prolactin, AMPK and Tumor Necrosis Factor (TNF). In addition, Gene Ontology functional

annotation showed that the DPRB-associated genes are involved in the regulation of cell

migration, signal transduction, angiogenesis, vasculature development and secretion (Fig. 1. D).

Despite the decrease in PGR expression during the MS phase (Supplemental Table 2, see

below), the global PGR binding trend was elevated as 88% of the intervals differentially bound

by PGR exhibited increased binding during the MS phase (Fig. 1. C), which is likely due to the

increased serum progesterone level in this phase of the cycle. To further explore enrichment of

other transcription factor binding sites co-occupying the PGR binding intervals, the DPRB DNA

motifs were analyzed by HOMER in two parts; those that showed elevated binding during MS

(MS-gain) or reduced binding during MS (MS-loss). The MS-gain intervals, indeed, showed

significant enrichment in PGR binding motif with a *p*-value of 1.00^{-40} (Table 2). MS-gain and MS-

loss intervals exhibited distinct profiles of transcription factor binding site preferences, with

FOSL2, *FRA1*, *JUN-AP1*, *ATF3* and *BATF* binding domains as top enriched sites in MS-gain

intervals (Table 2). Nuclear Receptors *AR*, bZIP transcription factor *CHOP* and some STAT

transcription factor members *STAT1*, *STAT3* and *STAT5* binding sites were also enriched in

sites with increased PGR binding (Table 2). In contrast, enriched motifs in the MS-loss intervals

included Estrogen Response Element (ERE), and binding domains for Transcription Factor 21

(*TCF21*), Atonal BHLH Transcription Factor 1 (*ATOH1*), Zinc Finger And BTB Domain

Containing 18 (*ZBTB18*), as well as GLI Family Zinc Finger 3 (*GLI3*, Table 3). Of note, during

the P to MS transition, PGR showed an increased preference for the Basic Leucine Zipper

Domain (bZIP), as the MS-gain intervals belonged mainly to this class. On the other hand,

preference for the Basic Helix Loop Helix (bHLH) and Zinc Finger (ZF) binding domains were

lost during this phase transition, as the enriched motifs identified in the MS-loss intervals

belonged mainly to these two classes. Thus, PGR's effects on gene expression may be partially

modulated through altered affinity for the different DNA responsive elements between the

liganded and unliganded form.

Transcriptional regulatory network of the P and MS endometrium

Whilst PGR has been widely studied in both humans and rodents and many direct and indirect target genes have been identified, a comprehensive analysis revealing its global regulatory function in the cycling human endometrium is still lacking. To fully characterize the functional relevance associated with PGR binding activities during the P to MS transition, we conducted RNA-seq on whole endometrium and incorporated the global gene expression profile into the ChIP-seq analyses during these two phases.

In total, we collected six P and five MS endometrial biopsies from which whole endometrial RNA was analyzed. This revealed a total of 14,985 expressed genes within the endometrium (FPKM > 1 in at least one of the two phases), whereby 14,303 and 14,156 were expressed in each of the P and MS phase, respectively. The transcriptomic profiles were subjected to hierarchical clustering and principal component analysis (PCA) as a measure of quality control. As shown in Supplemental Figure 1. A, a distinct segregation was observed for the P- and MS-derived RNA expression profile, and this is further supported by the hierarchical clustering presented in the dendrogram shown in Supplemental Figure 1. B, where samples from the two stages clustered accordingly. This suggested that the samples were well-characterized according to stage and of appropriate quality.

Of the genes expressed in the endometrium, 4,576 were differentially expressed (DEGs, Supplemental Table 3) between the two phases (absolute fold change > 1.5; and adjusted *p* value < 0.05). In total, 2,392 genes showed increased expression while 2,184 were downregulated during MS. Several genes known to regulate uterine biology, decidualization and implantation were identified as DEGs including decidualizing markers IGF Binding Protein 1

(*IGFBP1*) and prolactin (*PRL*); hedgehog protein, Indian Hedgehog (*IHH*); transcription factors

FOXO1 and *GATA2*; Wnt signaling molecules *WNT4*, *WNT2*, *WNT5A* and their inhibitor *DKK1*;

transcriptional repressor *ZEB1*; and extracellular matrix modulator *VCAN*. To interpret the

biological impact of the DEGs during the P to MS transition, Gene Set Enrichment Analysis

(GSEA) was performed to retrieve the functional profile associated with the DEGs [57].

Consistent with current literature, elevated inflammatory response was identified as an enriched

molecular function for the DEGs associated with the P to MS transition, as indicated by the

positive enrichment in the TNFA-NFKB signaling axis, coagulation, allograft rejection, hypoxia,

the complement cascade, interferon gamma response, IL6-JAK-STAT3 signaling and apoptosis

(Table 4). On the other hand, the negatively enriched functions which represents repressed

molecular pathways during MS showed significance in cell division regulatory mechanisms –

including E2F targets, G2M checkpoint and mitotic spindle regulations (Table 4). The xenobiotic

metabolism pathway was identified as one of the most positively enriched functions in the MS

endometrium by both GSEA (Table 4, Fig. 2. A) and Ingenuity Pathway Analysis (data not

shown). To validate the RNA-seq results, we examined expression of selected xenobiotic

metabolism genes using RNA extracted from an independent set of endometrial biopsies (n = 6

for each of the P and MS phase), along with the expression of the decidualization markers *PRL*

and *IGFBP1* to confirm the sample stages (Figs. 2. B and C). In accordance with the RNA-seq

results (Fig. 2. D), the cytochrome P450 members *CYP2C18* and *CYP3A5*, solute carriers

SLC6A12 and *SLCO4A1*, and glucuronosyltransferase *UGT1A6* were all found to be

upregulated during MS (Fig. 2. E). Further, glutathione S-transferase Mu genes (*GSTM1*,

GSTM3 and *GSTM5*), sulfotransferase *SULT1C4*, and solute carrier *SLCO2A1* were found to

be repressed during the MS phase (Fig. 2. G) similarly to that observed with RNA-seq (Fig. 2.

F).

Functional profiling of DEGs with regulated PGR binding during P to MS

To search for the genes that are directly regulated by PGR and important in modulating implantation, we identified the genes that were both differentially expressed and differentially bound by PGR in the whole endometrium between the P and MS phases. Comparison of DEGs and DPRB gene lists revealed 653 genes common to both datasets (Fig. 3. A). The trend for PGR binding and altered gene expression during MS, as compared to P is summarized in Table 5 and graphically presented in Figure 3. B. This analysis found 87% of the genes showed increased PGR binding (572 out of 653), and 70% showed upregulation during the MS phase (454 out of 653). Interestingly, the majority of these genes showed a positive correlation between PGR binding change and transcriptional regulation, i.e. increased PGR binding was associated with increased gene expression and vice versa. Thus, PGR binding generally promotes rather than represses gene expression in the human endometrium (Fig. 3. C).

The physiological function of PGR in regulating endometrial biology was next examined by elucidating the enriched functions associated with the PGR-regulated DEGs during the P to MS shift. The genes, along with fold change were submitted to GSEA to examine the enrichment of biological functions (Table 6). Enrichment was observed for a wide range of biological processes including inflammatory response signaling (coagulation, TNFA signaling via NFKB, complement, hypoxia, interferon gamma response), xenobiotic metabolism, epithelial mesenchymal transition (EMT), cell death regulation (apoptosis, p53 pathway), interleukin/STAT signaling, estrogen response, and MTORC1 response. Many of these biological functions were similarly identified using the DAVID Bioinformatic Database such as the regulation of cell death, inflammatory response, cytokine production, response to hormone and response to oxygen levels (Supplemental Table 4). Additionally, "secretion by cell" was identified as a regulated pathway by DAVID (p = 6.60E-5), supporting the validity of the secretory-phase derived gene

expression profile. Other pathways identified by DAVID included cell migration, signal

transduction, angiogenesis, leucocyte migration, nitric oxide biosynthetic processes, ECM

disassembly, and various activities associated with lipid regulation and insulin response

(Supplemental Table 4).

Additionally, some genes known to regulate decidualization and implantation showed

constitutive PGR binding during both phases (*FOXO1*, *HOXA10*, *HAND2*, *SOX17* and *CYR61*),

suggesting that constitutive PGR binding may regulate endometrial functions. We thus

examined the biological significance of the DEGs with constitutive PGR binding. Overlaying the

constitutive PGR bound genes (Supplemental Table 1) and DEGs (Supplemental Table 3)

identified 2,334 common genes (Fig. 3. D). The consistent PGR binding to these genes suggest

that their altered expression is not regulated directly by altered PGR binding and may require

input from other regulatory factors. Evaluation of the biological proccecesses controlled by this

group of genes showed primarily proliferative functions (cell cycle, cell division, nuclear division,

DNA replication, Supplemental Table 5), and further analysis using GSEA confirmed that the

proliferative function is repressed (Table 7, negatively enriched pathways). Additionally, TNFA

signaling via NFKB, inflammatory response and hypoxia were identified as top positively

enriched signaling pathways associated with this group of genes. Comparison of the functional

profile defined by the DEGs that were differentially (Table 6) or constitutively (Table 7) bound by

PGR showed some common signaling pathways involving both groups of genes. However,

DEGs with DPBR appear to engage more specifically with functions including coagulation, EMT,

estrogen response and apoptosis.

To authenticate the ChIP-seq results and the regulatory role of PGR, PGR-chromatin

association was evaluated for selected genes from the apoptosis and EMT pathways, both of

which are known to regulate receptivity. In addition, we examined PGR binding near the MAF

bZIP Transcription Factor (*MAF*), a regulator of the xenobiotic metabolism pathway shown

earlier to be positively enriched during MS. Two known PGR-regulated genes in the human

endometrial cells, *IHH* and *FOSL2* were first validated and confirmed to show increased

(*FOSL2*) and decreased (*IHH*) PGR binding during the MS phase (Figs. 3. E). Apoptosis

regulating genes Epithelial Membrane Protein 1 (*EMP1*), Immediate Early Response 3 (*IER3*),

and B-Cell CLL/Lymphoma 2 Like 10 (*BCL2L10*), as well as EMT mediators GTP Binding

Protein Overexpressed In Skeletal Muscles (*GEM*) and Serpin Family E Member 1

(*SERPINE1*), all displayed elevated PGR binding during the MS phase indicated by

independent ChIP-qPCR analysis (Fig. 3. F). Additionally, independent qPCR analysis revealed

the elevated transcription of apoptotic modulators (*EMP1*, *IER3* and *BCL2L10*) and the EMT

regulator *SERPINE1*. Other genes regulating these two pathways were also found to be

transcriptionally regulated, including Glutathione Peroxidase 3 (*GPX3*), Tissue Inhibitor Of

Metalloproteinases 3 (*TIMP3*), Vanin 1 (*VNN1*), Nicotinamide N-Methyltransferase (*NNMT*) and

Transglutaminase 2 (*TGM2*, Fig. 3. G).

To identify potential regulators associated with PGR, we next used IPA to predict for activity of

upstream regulators based on the 653 common genes (DEG + DPRB), and DEGs without

differential PR binding (DEG – DPRB, 3,923 genes), and upstream regulators were compared.

This comparison showed a higher Z-score for both progesterone and FOXO1 (a known co-

factor of PGR) in the regulation of the DEG + DPRB genes compared to the DEG – DPRB

genes (Fig. 3. H), confirming that this group of genes is more closely associated with with the

progesterone-PGR signaling. Amongst the upstream regulators predicted for each gene set, the

inflammation associated transcription factor NFKB family including REL, RELB and NFKB2 all

possessed a stronger activation score in the DEGs + DPRB (Fig. 3. H), suggesting enhanced

activity based on the altered gene expression network. In addition to NFKB, the angiogenic

modulators ANGPT2 and VEGF, developmental regulators HOXD10 and SOX4, histone

modifier KAT5 and the kinase MAP2K4 were all regulators predicted to have a higher activation

score in regulating the group of genes with differential PGR binding. Interestingly, the cell cycle

regulator CCND1, transcriptional regulators FOXM1 and MITF, prostaglandin receptor PTGER2

and the kinase protein ERBB2 were all predicted to be strongly inhibited in the regulation of

DEG - DPRB, but Z-score prediction suggest that those factors were not inhibited in the

regulation of the DEGs + DPRB. This suggests that although PGR may not directly inhibit these

factors, they may engage with PGR in a co-operative manner to regulate the downstream gene

expression network. Moreover, the MET-HGF receptor ligand pair as well as fat metabolism

modulators PLIN5, LEPR and Insulin I were all found with increased activity in regulating the

DEGs + DPRB, suggesting that these signaling axes are also associated with PGR function in

the cycling human uterus. Interestingly, although Insulin (*INS*) itself was not transcriptionally

regulated during the P to MS cycle, its cognate receptor Insulin Receptor (*INSR*) showed strong

transcriptional induction (Supplemental Table 3). Additionally, many genes known to be

regulated by insulin including *TIMP3* (Fig. 3. G, Supplemental Table 3), *SOD2*, *SOCS3*, *PRLR*

and *MMP2* all showed elevated mRNA expression in the MS endometrium (Supplemental Table

3).

Epithelial transcriptome in the cycling endometrium

To further understand the complexity of the cycling uterus, we assessed transcriptional changes

in the epithelial lining of the endometrium. As the endometrium consists of a complex and

dynamically changing set of cells, gene expression profiles derived from whole endometrial

biopsies often overlook alterations of specific cell types. Four P and five MS endometrial

samples were obtained, from which the luminal and glandular epithelial RNA were extracted and

subjected to RNA-seq analysis. Principal component analysis (PCA) and hierarchical clustering

found good segregation of the gene expression profile derived from two differently staged

samples (Supplemental Fig. 2. A and B). In the epithelium, we found a comparable number of

genes expressed to that of the whole endometrium, with 14,502 genes and 13,993 genes

transcriptionally active during the P and MS phase, respectively. The same threshold for

identifying DEGs in the whole endometrium was applied to the epithelium-expressed genes,

with which 3,052 epithelial-specific DEGs were found (epi-DEGs, Supplemental Table 6). Of

those, 57% (1,764) showed elevated transcription and 43% (1,288) was transcriptionally

repressed during the MS phase. Functional enrichment analysis of the epi-DEGs using GSEA

showed a positive enrichment for the genes encoding components of the apical junction

complex (Table 8), a molecular process important in defining the polarity of the epithelium and

hence supports the authenticity of the gene expression profile obtained from an epithelial origin.

Most of the pathways identified for the epithelium, whether positively or negatively enriched,

were principally similar to that of the whole endometrium, with enrichment in pathways

regulating immune responses including coagulation, complement, TNFA signaling via NFKB,

apoptosis, as well as xenobiotic metabolism. On the other hand, cell division related processes

including E2F regulated cell cycle, G2M checkpoint, mitotic spindle and DNA repair were

negatively enriched (Table 8).

Functions specific to the endometrial epithelium during implantation

To tease out the epithelial specific molecular events, we next compared the whole endometrium DEG with the epi-DEG to identify DEGs that are unique to the epithelium. In total, 2,411 common genes were found, representing those that show differential expression in both the whole endometrium and epithelium (Fig 4. A, Supplemental Tables 3 and 6). Of those, 2,394 genes showed the same transcriptional change between the two compartments, and 17 genes, although identified as "common" DEGs, exhibited the reversed change in mRNA level between the two compartments. Altogether with the 641 genes which were exclusively regulated in the epithelium, a total of 658 genes which were "specifically" regulated in the epithelium was found. Canonical pathways regulated by this group of genes were assessed using IPA and ranked according to significance in Table 9. Synthesis of glycosaminoglycans, including dermatan sulfate and chondroitin sulfate, as well as cholesterol biosynthesis were the most significant pathways identified (-Log p value > 3). Osteoarthritis pathway, cholecystokinin/gastrin mediated signaling, IL8 signaling and TGFB signaling were all significant pathways with a positive Z-score, suggesting increased activity during MS in the endometrial epithelium. On the other hand, PTEN signaling was identified as significantly repressed in the MS epithelium (Table 9).

IPA was next used to predict for upstream regulator activities in the epithelium (Supplemental Table 7). As expected, both progesterone and PGR were identified as activated upstream regulators, with Z-score values of 2.269 and 3.812, and p values of 1.77E-46 and 1.29E-29, for progesterone and PGR, respectively. Estrogen Receptor Alpha (ESR1) was shown to be repressed while ESR2 was activated. Interestingly, RNA-seq results illustrated decreased expression of *ESR1* and upregulation of *ESR2* in the MS epithelium (Supplemental Table 6). The top activated regulators were cytokines including IL1B, TNF, IFNG, OSM and IL1A; as well as transcriptional regulators such as NUPR1, NFKB, SMARCA4 and CEBPA (Z-score > 5).

Repressed regulators included transcription factors TBX2, TAL1; small GTPase RABL6, as well

as the E1A Binding Protein P400 (EP400).

Lastly, to identify regulators with specific activities in the epithelium during the MS phase, we

cross-compared the upstream regulators identified for the whole endometrium DEGs and

epithelial DEGs. To ensure that the upstream regulators identified were meaningful and

relevant, we compared only the regulators with p-values less than 0.05, and the numerical

activation Z-score values greater than 1.5. This comparison yielded several regulator proteins

with specific actions in the epithelium, of which selected are shown in Table 10. Amongst those

were transcriptional regulators POU5F1, IRF5, IRF8 and FOXJ1; Myocyte Enhancer Factors

family MEF2C and MEF2D; transmembrane receptors TLR5, IL1R1 and FCGR2A; kinase

proteins MET and AURKB; the growth factor HBEGF; the CYP27B1 enzyme and Wnt ligand

WNT7A; as well as the Notch ligand DLL4. Interestingly, *MEF2C*, *MEF2D*, *IRF8*, *FOXJ1*,

HBEGF, *CYP27B1* and *DLL4* were found to be uniquely regulated in the epithelium, where

either transcriptional regulation was not detected in the whole endometrium or showed a

different pattern of gene expression during the P to MS transition. In addition, *HBEGF* was

detected at very low levels as indicated by an average FPKM value of 2.63 in the whole

endometrium; compared to 19.26 in the epithelium (data not shown), suggesting that the

transcription of this gene is enriched in the epithelial cells during the WOI. We examined the

protein expression of two epithelial-specific regulators IRF8 and MEF2C using formalin-fixed

and paraffin-embedded endometrial biopsies from independent patients. As shown in Figure 4.

B, IRF8 is expressed in both stromal and epithelial cells and exhibited elevated protein

expression during the MS phase. MEF2C, on the other hand, did not show substantial increase

in staining intensity, but displayed a robust cytoplasmic-to-nuclear translocation from the P to

MS stage in the glandular epithelium (Fig. 4. C). These results suggest that both IRF8 and

MEF2C, two proteins previously unreported to have a role in implantation are regulated both at

the mRNA and protein level during the peri-implantation phase of the menstrual cycle in the

epithelium, and hence may have important functions in the implantation-phase endometrium.

Discussion

Here, we investigated the cycling human endometrium at the molecular level with two major aims in mind. First, to gain better understanding of the human endometrial signaling pathways and molecular events controlled by PGR during the P to MS transition. Combining the PGR cistromic and whole endometrium transcriptomic profile allowed the identification of genes with both proximal PGR binding and transcriptional regulation during the WOI. Second, we examined the gene expression profile using RNA derived from the whole endometrium or from the epithelium, including both luminal and glandular. Comparison of the two expression profiles delineated a more sophisticated and compartment specific transcriptional network. The latter has remained a challenging task and for this reason, the endometrium has often been examined as a whole when conducting *in vivo* studies.

The biological significance of PGR transcriptional activity during the WOI

Using PGR ChIP-seq, we obtained a genome wide DNA-binding blueprint of PGR in the endometrium at the P and MS phases. Comparison of the two identified DEGs with constitutive or regulated PGR binding in proximity during this period. Using the motif finding tool HOMER, we found a distinguishing difference in PGR binding preference from P to MS. While sites with increased PGR bindings at MS were predominantly co-occupied by bZIP and STAT transcription factors, sites with reduced PGR binding during MS were shared by bHLH and ZF transcription factors. This finding may suggest a mechanism of regulation for PGR transcriptional activity whereby its preference for certain DNA motifs is gained or lost during different phases of the menstrual cycle. Alternatively, the association of PGR to these DNA motifs may not be a direct one, but rather through interaction with other transcription factors

which then associate with the promoter region. The changes in DNA motifs detected based on

altered PGR binding could in turn suggest a change in PGR preference for different transcription

factors rather than different DNA motifs. Indeed, PGR is known to control gene expression in

this way through transcription factors such as SP1 and AP1 in human endometrial cells and

mammary cells [28, 58, 59].

A more comprehensive landscape of PGR biological impact was achieved by comparing the

whole endometrium derived DEGs to genes with DPRB in proximity to identify genes whose

transcription is likely directly regulated during the menstrual cycle by PGR. We found 653 such

genes, and analysis by GSEA identified many enriched pathways one of which is the

metabolism of xenobiotics. To the best of our knowlesge, our study is the first to identify

xenobiotic metabolism as a PGR regulated patway in the cyling endometrium. Xenobiotics are

conventionally defined as entities foreign to a cell or tissue such as drugs and pollutants,

although it can also refer to entities found at levels greater than considered norm. Xenobiotic

metabolism hence refers to the modification of these entities which in turn allows their systemic

removal. Genes involved in this pathway are broadly categorized into 3 phases: phase 1 and

phase 2 enzymes increase the solubility of the xenobiotics by introducing polar moieties and

conjugating to endogenous hydrophilic molecules; and phase 3 genes encode transporters

which then traffic the xenobiotic metabolites out of the cells to be excreted [60]. Although

expression of xenobiotic metabolizing genes has been previously reported in the endometrium

[61], defined and validated endometrial expression and function are still absent. Our data

demonstrate transcriptional regulation of genes encoding phase 1 and 2 enzymes, as well as

phase 3 transporters in the endometrium. These included numerous aldehyde dehydrogenase

(ALDH) members, carboxylesterases, carbohydrate sulfotransferases, cytochrome P450

members, glutathione S-transferases, monoamine oxidases and UDP glycotransferases; and

multi-drug resistance protein member ABCC3. Independent qPCR analysis confirmed that

xenobiotic metabolism genes were transcriptionally regulated during the menstrual cycle.

Interestingly, genes encoding receptors known to mediate xenobiotic metabolism gene

expression, including *NR1I3* and *NR1I2* were virtually not expressed (FPKM < 1), while *AHR*

was lowly and non-differentially expressed in the endometrium during the phase transition (data

not shown), suggesting that transcriptional regulation of the xenobiotic metabolism network may

not occur in a classical manner, but rather through alternative regulatory mechanisms [62].

Although the impact of xenobiotic metabolism regulation during mid-secretory in the human

endometrium remains elusive, there has been evidence linking dysregulation of xenobiotic

metabolism genes to pathological conditions such as infertility and cancer [63]. Moreover, it has

been proposed that xenobiotic metabolism may act as a detoxification mechanism, providing

protection and guarding the endometrium against harmful environmental insult for appropriate

and efficient implantation, such as environmental estrogen [63].

In addition to xenobiotic metabolism, apoptosis and EMT were also pathways identified as PGR

regulated, and both have received ample attention as pathways important in endometrial

function. Apoptosis has long been known to mediate uterine homeostasis, a disruption of which

is evidently linked to implantation failure and endometriosis [64, 65]. Based on our *in silico*

analysis, PGR appeared to promote as well as suppress apoptosis in the mid-secretory

endometrium (See Supplemental Table 4). However, the onset of apoptosis in the cycling

endometrium is typically around the late-secretory to menstruation phase [66], suggesting a

possibility that during the MS phase, PGR acts to balance rather than induce cell death before

the mass apoptosis ensues during late-secretory. Indeed, we confirmed increased PGR binding

and increased transcription of both pro- and anti-apoptotic genes including *EMP1* [67], *IER3* [68]

and *BCL2L10* [69]. EMT and its reciprocal pathway, the mesenchymal-epithelial transition

(MET) are important modulators of uterine physiology. During each menstrual cycle, the

endometrium undergoes extensive remodeling which involves the building and shedding of the

functional layer. The origin of the epithelial cells has long been under debate, with some

evidence supporting MET being a major player for endometrial re-epithelialization [70, 71]. It

has been postulated that by retaining imprint of the mesenchymal origin, the endometrial

epithelial cells are prone to return to its mesenchymal state via EMT [72]. In the MS

endometrium, we found various EMT modulating genes to be transcriptionally regulated by

PGR, including *MMP2*, *SERPINE1*, *NNMT*, and *WNT5A*. Interestingly, although PGR appeared

to promote the expression of EMT genes during the WOI, a closer examination of our gene

expression data suggested that the consequences of these regulatory activities resulted in

neither decreased epithelial properties nor increased mesenchymal properties. The

mesenchymal cell marker *CDH2* was strongly repressed (seven-fold), while another marker,

VIM, although not identified as a DEG, showed a significant decrease with a fold-change that

did not qualify for differential expression in the MS endometrium (data not shown). On the other

hand, numerous epithelial cell markers including *CDH1*, *CLDN1*, *CLDN4*, *CLDN8*, *CLDN10*,

KLF4 and *KLF5* were all upregulated during MS. Additionally, *CLDN4*, *CLDN8* and *KLF4* were

also presented with increased PGR binding in proximity, suggesting that PGR may directly

promote the upregulation of these epithelial markers and maintain the epithelial-like

characteristic of these cells. It is possible that while some mesenchymal properties in the

epithelium provide for the implanting embryo (such as decreased cell to cell adhesion), but a

complete loss of the epithelial status is likely unfavorable and hence PGR acts both to increase

EMT as well as maintain the epithelial state. In support of this, EMT has been postulated as an

important modulator of noninvasive trophoblast implantation in bovines [73].

Role and function of the endometrial epithelia during the WOI

RNA-seq was conducted to evaluate the transcriptomic profile in the epithelial compartment of

the endometrium. A simple functional annotation found comparable biological functions as that

of the whole endometrium, including inflammatory responses, TNFA/NFKB signaling, xenobiotic

metabolism, apoptosis, KRAS signaling and EMT as positively enriched; and E2F signaling,

G2M checkpoint, mitotic spindle and DNA repair as repressed. IPA predicted the activity of

various upstream regulators based on the epithelial transcriptome which included cytokines and

transcriptional regulators. Some cytokines identified in our study have been known to facilitate

implantation in mammals, whilst the functions of others remain elusive. Additionally, the majority

of the epithelial transcription regulators identified in our study have yet to be studied for

functional relevance in mediating implantation in the human endometrium, including NUPR1,

TBX2, SMARCA4, CEBPA, RABL6 and EP400. Interestingly, the Estrogen Receptors ESR1

and ESR2 showed repression and activation during WOI in the epithelium, respectively.

Accordingly, RNA-seq results showed downregulation of ESR1 and upregulation of ESR2

during the phase transition in the epithelium. The repression of ESR1 activity during the window

of implantation is well documented, and a mouse model with epeithelial ESR1 deletion

illustrated a role in regulating apoptosis [74]. There is also evidence linking ESR1

overexpression and implantation failure in humans, emphasing the importance of regulated

ESR1 expression during this critical period [75]. In contrast, ESR2 has received little attention in

the endometrium based on its low expression level compared to the ovary, oviduct or mammary

gland [15]. Our results showed that there is enrichment of ESR2 expression specifically in the

epithelium, with FPKM values in the endometrium averaging 0.75 in the whole endometrium and

2.1 in the epithelium (data not shown). The increased activity of ESR2 (as predicted by IPA), the

robust upregulation of its transcript as well as enriched epithelial expression during the WOI

propose a possibility that ESR2 engage in previously unrecognized role in mediating pregnancy.

To further unravel molecular pathways with increased specificity to the epithelium, we used two additional approaches. Firstly, we compared the whole endometrial-derived DEGs to epithelial-derived DEGs and excluded the common DEGs to obtain a profile of DEGs that were only detected in the epithelium. Whilst excluding the "common" DEGs may seem counter-intuitive, since the epithelium comprises a part of the endometrium and some "epithelial" genes with substantial transcriptional changes would surface when examined in the whole endometrium, thereby excluding the common DEGs would altogether eliminate those genes. However, the purpose of the epithelial specific examination is to identify previously "missed" epithelial-specific pathways (genes) when examining the endometrium as a whole. Genes in this category may show changes that are subtle but not necessarily less important in nature, and hence our approach of excluding the "common" DEGs. The second approach was to compare the activity status of the upstream regulators calculated for each set of DEGs and identify upstream regulators with enhanced activity in the epithelium. Using the IPA software to examine the 658 epithelial-specific DEGs, the most represented canonical pathways were dermatan sulfate, chondroitin sulfate and cholesterol biosynthesis. Dermatan and chondroitin sulfate are glycosaminoglycans found mostly in the skin, blood vessels and the heart valves [76]. They are known to regulate coagulation and wound repair, as well as recruit natural killer cells into the uterus during the reproductive cycle [77]. However, the specific role of the endometrial epithelial cells in biosynthesis of these glycosaminoglycans has not yet been reported. On the other hand, progesterone has been reported to inhibit the synthesis of cholesterol in the uterine epithelium of mice, and this has been postulated as a mechanism to block epithelial cell proliferation. Our data accordingly suggest that suppression of cholesterol biosynthesis may be more specifically refined to the epithelial compartment, possibly associated with PGR-mediated inhibition of epithelial cell proliferation during the MS phase [78].

Lastly, we identified two transcription factors, IRF8 (ICSBP) and MEF2C with enhanced activity in the epithelium, whose protein levels and cellular localization were regulated in the epithelium during the WOI. MEF2C specifically showed nuclear localization during this period, and as MEF2C is a transcription factor, it's likely that the nuclear localization is associated with its increased transcriptional capacity. IRF8 is a member of the interferon (IFN) regulatory factor (IRF) family and is known to regulate gene expression in an interferon-dependent manner [79]. It is a modulator of cellular apoptosis under pathological conditions and deregulation of other family members are associated with endometrial adenocarcinoma [80-84], suggesting that IRF proteins may regulate female reproduction. Supporting this, Kashiwagi *et al.* have reported IRF8 expression in the murine endometrium in response to the implanting embryo, but not in pseudopregnancy [85], and Kusama *et al.* later reported the upregulation of IRF8 in the bovine endometrial luminal epithelium in response to the embryo derived interferon tau [86]. MEF2C belongs to the MADS box transcription enhancer 2 family, which plays a role in proliferation, invasion and differentiation in various cell types [87]. Other members of the family (MEF2A and MEF2D) are known to modulate cytotrophoblast invasion and differentiation in the human placenta [88], and MEF2C itself has been associated with endometriosis, although no apparent function has been reported in the endometrial epithelium [89]. Whilst little is known regarding the epithelial function of IRF8 and MEF2C in the endometrium during the WOI, our findings suggest that these factors could have important functions in the uterus and female reproduction.

In summary, signaling pathways controlled by progesterone and PGR are indispensable in uterine biology and homeostasis, a disruption of which manifests in a wide range of gynecological abnormalities such as endometriosis, adenomyosis, fertility defects and endometrial cancer. These pathological conditions are linked to dysregulation of many

molecular pathways amongst which are EMT, apoptosis, cell migration and inflammatory

response. In this study we provide evidence to show how some of these pathways could be

directly controlled by the progesterone signaling through the transcriptional activity of PGR. An

understanding of the precise regulatory pattern and mechanism of PGR, that is, what genes are

regulated by PGR, and how these genes are regulated by PGR provide a bridging link to explain

the molecular mechanism of disease phenotypes under aberrantly regulated PGR conditions.

One limitation of this study is that ChIP-seq does not take into consideration the control of PGR

over distal DNA response element due to the chromatin interaction in a three-dimensional

structure. It is worth noting that roughly 21-22 % of the PGR bound intervals occurred in the

"intergenic" regions of the genome (Fig. 1. A), which is defined as greater than 25 kb from the

TSS. Although it is possible that these bindings have transcriptional relevance, we cannot draw

any conclusion from this study. To address this, future studies should aim to attain a

comprehensive three-dimensional structure to elucidate the chromatin conformation in parallel

to PGR binding using techniques such as Hi-C [90, 91]. This will allow the identification of PGR

binding sites in a more global view without the limitation of chromosomal distance. Additional to

the PGR regulatory function, approaching the uterine transcriptomic analysis in a compartment

specific manner enabled the identification of numerous proteins with previously unrecognized

roles in uterine biology and pregnancy. These findings provide a direction for future studies

aimed to explore molecular factors crucial for uterine homeostasis.

Materials and Methods

Ethics Statement

This project was executed in accordance with the federal regulation governing human subject

research. All procedures were approved by the following ethics committees the University of

North Carolina at Chapel Hill IRB under file #:05-1757. Informed consent was obtained from all

patients before their participation in this study.

Human Endometrial Samples

We recruited normal volunteers with the following inclusion criteria: ages 18-37, normal

menstrual cycle characteristics, an inter-cycle interval of 25-35 days, varying no more than 2

days from cycle to cycle, a normal luteal phase length without luteal spotting, and a body mass

index (BMI) between 19 - 28. We excluded women with infertility, pelvic pain, signs and

symptoms of endometriosis, history of fibroids or history of taking medication affecting hormonal

function in the last 3 months. Endometrial samples were taken using an office biopsy instrument

(Pipelle™, Milex Products Inc., Chicago, IL) from the volunteers. Cycle day was determined

based on the last menstrual period combined with menstrual history (P samples) or date of

Luteinizing Hormone surge. Cycle phase and endometrial normality was confirmed with H&E

staining based on the Noyes criteria [92]. Details for patients with accessible data are

summarized in Supplemental Table 8.

RNA-seq and Analysis

RNA was prepared from endometrial samples using TRIzol (Thermo Fisher Scientific, Waltham, MA) under the manufacturer's suggested conditions. Absorption spectroscopy (NanoDrop 8000, Thermo Fisher Scientific, Waltham, MA) was used for quantification of RNA with a ribosomal RNA standard curve. The RNA libraries were sequenced with a HiSeq 2000 system (Illumina). The raw RNA-Seq reads (100 nt, paired-end) were initially processed by filtering with average quality scores greater than 20. Reads which passed the initial processing were aligned to the human reference genome (hg19; Genome Reference Consortium Human Build 19 from February 2009) using TopHat version 2.0.4 [93] and assembled using Cufflinks version 2.0.2 [94]. BigWig file was generated from normalized bedgraph file of each sample using bedGraphToBigWig. Scores represent normalized mapped read coverage. Expression values of RNA-Seq were expressed as FPKM (fragments per kilobase of exon per million fragments) values. Differential expression was calculated using Cuffdiff [94]. Transcripts with FPKM > 1, q-value < 0.05 and at least 1.5-fold change were defined as differentially expressed genes (DEG). The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus and are accessible through GEO Series accession number GSE132713 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE132713).

Chromatin immunoprecipitation sequencing (ChIP-seq) and qRT-PCR (ChIP-qPCR)

Two sets of biopsied tissues were derived from healthy volunteers, each set comprising of one P and one MS endometrial samples (termed P1 and MS1 for set1, and P2 and MS2 for set2). The tissues were flash frozen and sent to the Active Motif company for Factor-Path ChIP-seq analysis. The tissues were fixed, followed by sonication to shear the chromatin into smaller fragments before immunoprecipitation using the Progesterone Receptor (PGR) antibody (sc-

7208, Santa Cruz). PGR-bound DNA was subsequently purified and amplified to generate a

library for sequencing and quantitative real-time PCR (ChIP-seq and ChIP-qPCR). Sequencing

was performed using a NextSeq 500 system (Illumina). The raw ChIP-seq reads (75 nt, single-

end) were processed and aligned to the human reference genome (hg19; Genome Reference

Consortium Human Build 19 from February 2009) using Bowtie version 1.1.2 [95] with unique

mapping and up to 2 mismatches for each read (-m 1 -v 2). The duplicated reads with the same

sequence were discarded, and the bigWig files were displayed on UCSC genome browser as

custom tracks. Peak calling for each sample was performed by SICER version 1.1 with FDR of

0.001. Software MEDIP was used to identify differential peaks of PGR binding between the P

and MS samples [96]. Each region was defined as the genomic interval with at least 2-fold

difference of read count and p-value ≤ 0.01. Each differential peak was mapped to nearby gene

using software HOMER's "annotatePeaks.pl" function [97]. As we observed technical variation

between sample set1 and set2, we employed a paired-analysis strategy where differential PGR

binding intervals were independently determined for P1 VS MS1; and P2 VS MS2. Differential

PGR binding that were common to both data sets were used for downstream analysis (Fig. 1.

B). Genomic intervals with consistent (or constitutive) PGR binding were defined as motifs

bound by PGR in both P and MS phases in either set1, set2 or both; where the read count

between the two phases did not qualify for "differential" PGR binding. The motif analysis of

differential PGR binding peaks was performed using HOMER software's "findMotifsGenome.pl"

command with default setting [97]. The data discussed in this publication have been deposited

in NCBI's Gene Expression Omnibus and are accessible through GEO Series accession

number GSE132713 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE132713).

Epithelial isolation

Endometrial samples obtained from normal controls during the secretory phase of the menstrual

cycle were washed with Opti-mem media supplemented with fetal bovine serum (FBS) and

antibiotics (10 000 IU/mL penicillin, 10 000 IU/ mL streptomycin; Life Technologies, Grand

Island, New York). Tissue was recovered via centrifugation and incubated with collagenase-

containing medium (phenol red-free Dul- becco Modified Eagle Medium/F12, 0.5% collagenase

I, 0.02% DNase, and 5% FBS). Cell types were separated as described previously [98].

RNA extraction, cDNA conversion and qPCR

For validation of RNA-seq results, selected genes were examined for RNA expression using

independent patients' samples. Endometrial tissues were resected from patients and flash

frozen in liquid nitrogen (see Suppelmental Table 8 for patient details). RNA was extracted as

described above. Reverse transcription was performed to convert RNA into cDNA using the

Moloney Murine Leukemia Virus (MMLV) reverse transcriptase (Thermo Fisher) according to

the manufacturer's instructions. Quantitative real-time PCR was performed using the

SsoAdvancedTM Universal SYBR Green Supermix (1725274, Bio-Rad). Briefly, reaction

samples were prepared to a total volume of 10 µL with 250 nM of each of the forward and

reverse primers, 0.5 ng of cDNA and a final 1 X concentration of the SYBR Green Supermix.

The reaction was heated to 98 ^{O}C for 30 sec, followed by 35 cycles of denaturation at 95 ^{O}C for

5 sec and annealing and elongation for 15 sec. Temperature cycles were performed on the CFX

Connect^{TM} Real-Time PCR Detection System (Bio-Rad). The primer sequences were as follows

(from 5' to 3', F = forward and R = reverse): *CYP3A5* - GTATGAAGGTCAACTCCCTGTG (F)

and GGGCCTAAAGACCTTCGATTT (R); *FMO5* - GATTTAAGACCACTGTGTGCAG (F),

CCATGACTCCATCAAAGACATTC (R); *UGT1A6* – TGTCTCAGGAATTTGAAGCCTAC (F),

GCAATTGCCATAGCTTTCTTCTC (R); *SLCO4A1* – CCCGTCTACATTGCCATCTT (F),

GGCCCATTTCCGTGTAGATATT (R); *SLC6A12* – CTTCTACCTGTTCAGCTCCTTC (F),

CGTGCAATGCTCTGTGTTC (R); *CYP2C18* – CATTGTGGTGTTGCATGGATATG (F),

AGGATTCCAAGTCCTTTGTTAACTT (R); *SULT1C4* – TAAAGCAGGAACAACATGGACT (F),

TTCGAGGAAAGGAAATCGTTGA (R); *SLCO2A1* – CTGTACAGCGCCTACTTCAA (F),

GATGGCATTGCTGATCTCATTC (R); *GSTM1* – CAAGCACAACCTGTGTGG (F),

TTGTCCATGGTCTGGTTCTC (R); *GSTM3* – GGAGTTCACGGATACCTCTTATG (F),

GGTAGGGCAGATTAGGAAAGTC (R); *GSTM5* – CGCTTTGAGGGTTTGAAGAAG (F),

TGGGCCCTATTTGCTGTT (R); *EMP1* – GTCTTCGTGTTCCAGCTCTT (F),

AAGAATGCACAGCCAGCA (R); *IER3* – TGGAACTGCGGCAAAGTA (F),

GTAGACAGACGGAGTTGAGATG (R); *BCL2L10* – CCAAAGAACCGCAGAAGAAAC (R),

GAAGTTGTGGAGAGATGAGAGG (R); *GPX3* – TCTGGTCATTCTGGGCTTTC (F),

ACCTGGTCGGACATACTTGA (R); *TIMP3* – CCCATGTGCAGTACATCCATAC (F),

ATCATAGACGCGACCTGTCA (R); *VNN1* – CAGATCAGGGTGCGCATATT (F),

GTTTACTTCAGGGTCTGGGATG (R); *SERPINE1* – CTGAGAACTTCAGGATGCAGAT (F),

AGACCCTTCACCAAAGACAAG (R); *NNMT* – ACCTCCAAGGACACCTATCT (F),

CACACCGTCTAGGCAGAATATC (R); and *TGM2* – ACCCAGCAGGGCTTTATCTA (F),

CCCATCTTCAAACTGCCCAA (R). All primers were synthesized by Sigma-Aldrich (St Louis,

MO), and gene expression was normalized to 18s rRNA by the ΔΔCT method.

Immunohistochemistry

Sections were cut from patient's endometrial biopsies that have been formalin-fixed and paraffin

embedded at 5 µm per section. Sections were baked at 65^{O}C for roughly 5 minutes and

deparaffined using the Citrisolv clearing agent (22-143-975, Thermo Fisher, Waltham, MA,

USA) and hydrated by immersing in decreasing gradient of ethanol. Antigen retrieval was

performed using the Vector Labs Antigen Unmasking Solution as per manufacturer's protocol

(H-3300, Vector Laboratories, Burlingame, CA, USA), followed by blocking the endogenous

peroxide using 3% hydrogen peroxide diluted in distilled water. Tissues were blocked in 5%

normal donkey serum before an overnight incubation with the primary antibody at 4^{O}C (1:200 for

ICSBP antibody, sc-365042, Santa Cruz; and 1:100 for MEF2C antibody, SAB4501860, Sigma-

Aldrich). The slides were washed twice in PBS at room temperature and applied with secondary

antibody diluted 1:200 in 1% BSA prepared in PBS (biotinylated anti-mouse IgG (H+L), BA-

9200, and biotinylated anti-rabbit IgG (H+L), BA-1000, Vector Laboratories). The ABC reagent

was applied to tissue in accordance with the manufacturer's instructions (Vector Labs ABC PK-

6100, Vector Laboratories). Signal was developed using the Vector Labs DAB ImmPACT

staining kit (Vector Labs SK-4105, Vector Laboratories). Finally, the tissue sections were

counterstained with hematoxylin and dehydrated through increasing ethanol concentration

before incubation in Citrisolv and coverslipping.

Data Analysis

Various bioinformatic tools were utilized to analyze the high content data generated in this

study. Principle component analysis and hierarchical clustering were achieved using the Partek

Genomics Suites 7.0 (Partek Inc., St. Louis, MO, USA, http://www.partek.com/partek-genomics-

suite/). Functional annotation and enrichment analysis were performed using a combination of

the following three tools: Ingenuity Pathway Analysis Software (IPA, http://www.ingenuity.com/),

Gene Set Enrichment Analysis (GSEA, http://software.broadinstitute.org/gsea/index.jsp/), and

Database   for   Annotation,   Visualization   and   Integrated   Discovery   (DAVID,

http://david.ncifcrf.gov/). Distribution of PGR binding throughout the genome was conducted

using   the   Peak   Annotation   and   Visualization   tool   (PAVIS,

https://manticore.niehs.nih.gov/pavis2/) [99], and PGR-bound motif was submitted to HOMER

motif   analysis   software   to   identify   presence   of   other   DNA-response   elements

(http://homer.salk.edu/homer/). GraphPad Prism software was used to analyze single gene

expression data generated from both RNA-seq, qPCR, and PGR ChIP-qPCR. Statistical

analysis including one-way ANOVA and Student's *t* test, with a *p*-value of less than 0.05

considered as significant. For pathway analysis using IPA, a given biological category was

subjected to Fisher's exact test to measure the probability that the category was randomly

associated. The categories with *p*-values less than 0.05 were defined as significantly enriched.

References

1.   Paiva, P., et al., *Human chorionic gonadotrophin regulates FGF2 and other cytokines produced by*
human endometrial epithelial cells, providing a mechanism for enhancing endometrial
receptivity. Hum Reprod, 2011. **26**(5): p. 1153-62.

2.   Gellersen, B. and J.J. Brosens, *Cyclic decidualization of the human endometrium in reproductive*
health and failure. Endocr Rev, 2014. **35**(6): p. 851-905.

3.   Jabbour, H.N., et al., *Endocrine regulation of menstruation.* Endocr Rev, 2006. **27**(1): p. 17-46.

4.   Ramathal, C.Y., et al., *Endometrial decidualization: of mice and men.* Semin Reprod Med, 2010.
28(1): p. 17-26.

5.   Hawkins, S.M. and M.M. Matzuk, *Menstrual Cycle: Basic Biology.* Ann N Y Acad Sci, 2008. **1135**:
p. 10-8.

6.   Mesen, T.B. and S.L. Young, *Progesterone and the luteal phase: a requisite to reproduction.*
Obstet Gynecol Clin North Am, 2015. **42**(1): p. 135-51.

7.   Speroff, L. and M.A. Fritz, *Clinical Gynecologic Endocrinology and Infertility*. 2005: Lippincott
Williams & Wilkins.

8.   Bashiri, A., K.I. Halper, and R. Orvieto, *Recurrent Implantation Failure-update overview on*
etiology, diagnosis, treatment and future directions. Reprod Biol Endocrinol, 2018. **16**(1): p. 121.

9.   Simon, A. and N. Laufer, *Assessment and treatment of repeated implantation failure (RIF).* J
Assist Reprod Genet, 2012. **29**(11): p. 1227-39.

10.   Garrido-Gomez, T., et al., *Defective decidualization during and after severe preeclampsia reveals*
a possible maternal contribution to the etiology. Proc Natl Acad Sci U S A, 2017. **114**(40): p.
E8468-e8477.

11.   Conrad, K.P., M.B. Rabaglino, and E.D. Post Uiterweer, *Emerging role for dysregulated*
decidualization in the genesis of preeclampsia. Placenta, 2017. **60**: p. 119-129.

12.   Macklon, N.S., J.P. Geraedts, and B.C. Fauser, *Conception to ongoing pregnancy: the 'black box'*
of early pregnancy loss. Hum Reprod Update, 2002. **8**(4): p. 333-43.

13.   Dey, S.K., et al., *Molecular cues to implantation.* Endocr Rev, 2004. **25**(3): p. 341-73.

14.   Bhurke, A.S., I.C. Bagchi, and M.K. Bagchi, *Progesterone-Regulated Endometrial Factors*
Controlling Implantation. Am J Reprod Immunol, 2016. **75**(3): p. 237-45.

15.   Pawar, S., et al., *Minireview: Steroid-regulated paracrine mechanisms controlling implantation.*
Mol Endocrinol, 2014. **28**(9): p. 1408-22.

16.   Young, S.L., *Oestrogen and progesterone action on endometrium: a translational approach to*
understanding endometrial receptivity. Reprod Biomed Online, 2013. **27**(5): p. 497-505.

17.   Young, S.L., et al., *Effect of randomized serum progesterone concentration on secretory*
endometrial histologic development and gene expression. Hum Reprod, 2017. **32**(9): p. 1903-
1914.

18.   Wilcox, A.J., D.D. Baird, and C.R. Weinberg, *Time of implantation of the conceptus and loss of*
pregnancy. N Engl J Med, 1999. **340**(23): p. 1796-9.

19.   Sunderam, S., et al., *Assisted reproductive technology surveillance--United States, 2009.* MMWR
Surveill Summ, 2012. **61**(7): p. 1-23.

20.   Patel, B., et al., *Role of nuclear progesterone receptor isoforms in uterine pathophysiology.* Hum
Reprod Update, 2015. **21**(2): p. 155-73.

21.   Yin, P., et al., *Genome-wide progesterone receptor binding: cell type-specific and shared*
mechanisms in T47D breast cancer cells and primary leiomyoma cells. PLoS One, 2012. **7**(1): p.
e29021.

22.   Kim, J.J., T. Kurita, and S.E. Bulun, *Progesterone action in endometrial cancer, endometriosis,*
792   *uterine fibroids, and breast cancer.* Endocr Rev, 2013. **34**(1): p. 130-62.
793   23.   Tamm, K., et al., *Genes targeted by the estrogen and progesterone receptors in the human*
794   *endometrial cell lines HEC1A and RL95-2.* Reprod Biol Endocrinol, 2009. **7**: p. 150.
795   24.   Matsumoto, H., et al., *Indian hedgehog as a progesterone-responsive factor mediating*
796   *epithelial-mesenchymal interactions in the mouse uterus.* Dev Biol, 2002. **245**(2): p. 280-90.
797   25.   Takamoto, N., et al., *Identification of Indian hedgehog as a progesterone-responsive gene in the*
798   *murine uterus.* Mol Endocrinol, 2002. **16**(10): p. 2338-48.
799   26.   Franco, H.L., et al., *Epithelial progesterone receptor exhibits pleiotropic roles in uterine*
800   *development and function.* Faseb j, 2012. **26**(3): p. 1218-27.
801   27.   Pan, H., et al., *Microarray analysis of uterine epithelial gene expression during the implantation*
802   *window in the mouse.* Endocrinology, 2006. **147**(10): p. 4904-16.
803   28.   Mazur, E.C., et al., *Progesterone receptor transcriptome and cistrome in decidualized human*
804   *endometrial stromal cells.* Endocrinology, 2015. **156**(6): p. 2239-53.
805   29.   Li, Q., et al., *The antiproliferative action of progesterone in uterine epithelium is mediated by*
806   *Hand2.* Science, 2011. **331**(6019): p. 912-6.
807   30.   Lee, K.Y., et al., *Bmp2 is critical for the murine uterine decidual response.* Mol Cell Biol, 2007.
808   **27**(15): p. 5468-78.
809   31.   Li, Q., et al., *Bone morphogenetic protein 2 functions via a conserved signaling pathway*
810   *involving Wnt4 to regulate uterine decidualization in the mouse and the human.* J Biol Chem,
811   2007. **282**(43): p. 31725-32.
812   32.   Lim, H., et al., *Hoxa-10 regulates uterine stromal cell responsiveness to progesterone during*
813   *implantation and decidualization in the mouse.* Mol Endocrinol, 1999. **13**(6): p. 1005-17.
814   33.   Taylor, H.S., et al., *HOXA10 is expressed in response to sex steroids at the time of implantation in*
815   *the human endometrium.* J Clin Invest, 1998. **101**(7): p. 1379-84.
816   34.   Kannan, A., et al., *The transcription factor C/EBPbeta is a marker of uterine receptivity and*
817   *expressed at the implantation site in the primate.* Reprod Sci, 2010. **17**(5): p. 434-43.
818   35.   Mantena, S.R., et al., *C/EBPbeta is a critical mediator of steroid hormone-regulated cell*
819   *proliferation and differentiation in the uterine epithelium and stroma.* Proc Natl Acad Sci U S A,
820   2006. **103**(6): p. 1870-5.
821   36.   Plante, B.J., et al., *Cyclic regulation of transcription factor C/EBP beta in human endometrium.*
822   Reprod Biol Endocrinol, 2009. **7**: p. 15.
823   37.   Gray, C.A., et al., *Developmental biology of uterine glands.* Biol Reprod, 2001. **65**(5): p. 1311-23.
824   38.   Kelleher, A.M., F.J. DeMayo, and T.E. Spencer, *Uterine Glands: Developmental Biology and*
825   *Functional Roles in Pregnancy.* Endocr Rev, 2019.
826   39.   Singh, H. and J.D. Aplin, *Adhesion molecules in endometrial epithelium: tissue integrity and*
827   *embryo implantation.* J Anat, 2009. **215**(1): p. 3-13.
828   40.   Wang, B., et al., *Annexin A2 Acts as an Adhesion Molecule on the Endometrial Epithelium during*
829   *Implantation in Mice.* PLoS One, 2015. **10**(10): p. e0139506.
830   41.   Guffanti, E., et al., *Nuclear pore complex proteins mark the implantation window in human*
831   *endometrium.* J Cell Sci, 2008. **121**(Pt 12): p. 2037-45.
832   42.   Salker, M.S., et al., *Activation of SGK1 in Endometrial Epithelial Cells in Response to PI3K/AKT*
833   *Inhibition Impairs Embryo Implantation.* Cell Physiol Biochem, 2016. **39**(5): p. 2077-2087.
834   43.   Boeddeker, S.J. and A.P. Hess, *The role of apoptosis in human embryo implantation.* J Reprod
835   Immunol, 2015. **108**: p. 114-22.
836   44.   Joswig, A., et al., *Apoptosis in uterine epithelium and decidua in response to implantation:*
837   *evidence for two different pathways.* Reprod Biol Endocrinol, 2003. **1**: p. 44.

45.   Valdez-Morales, F.J., et al., *Changes in receptivity epithelial cell markers of endometrium after*
ovarian stimulation treatments: its role during implantation window. Reprod Health, 2015. **12**: p.
45.
46.   Kelleher, A.M., et al., *Uterine glands coordinate on-time embryo implantation and impact*
endometrial decidualization for pregnancy success. Nat Commun, 2018. **9**(1): p. 2435.
47.   Aghajanova, L., *Leukemia inhibitory factor and human embryo implantation.* Ann N Y Acad Sci,
2004. **1034**: p. 176-83.
48.   Kimber, S.J., *Leukaemia inhibitory factor in implantation and uterine biology.* Reproduction,
2005. **130**(2): p. 131-45.
49.   Norwitz, E.R., D.J. Schust, and S.J. Fisher, *Implantation and the survival of early pregnancy.* N
Engl J Med, 2001. **345**(19): p. 1400-8.
50.   Vassen, L., W. Wegrzyn, and L. Klein-Hitpass, *Human insulin receptor substrate-2 (IRS-2) is a*
primary progesterone response gene. Mol Endocrinol, 1999. **13**(3): p. 485-94.
51.   Kaya, H.S., et al., *Roles of progesterone receptor A and B isoforms during human endometrial*
decidualization. Mol Endocrinol, 2015. **29**(6): p. 882-95.
52.   Okada, H., T. Tsuzuki, and H. Murata, *Decidualization of the human endometrium.* Reprod Med
Biol, 2018. **17**(3): p. 220-227.
53.   Rubel, C.A., et al., *A Gata2-Dependent Transcription Network Regulates Uterine Progesterone*
Responsiveness and Endometrial Function. Cell Rep, 2016. **17**(5): p. 1414-1425.
54.   Wang, X., et al., *SOX17 regulates uterine epithelial-stromal cross-talk acting via a distal enhancer*
upstream of Ihh. Nat Commun, 2018. **9**(1): p. 4421.
55.   Huang da, W., B.T. Sherman, and R.A. Lempicki, *Bioinformatics enrichment tools: paths toward*
the comprehensive functional analysis of large gene lists. Nucleic Acids Res, 2009. **37**(1): p. 1-13.
56.   Huang da, W., B.T. Sherman, and R.A. Lempicki, *Systematic and integrative analysis of large*
gene lists using DAVID bioinformatics resources. Nat Protoc, 2009. **4**(1): p. 44-57.
57.   Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for*
interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A, 2005. **102**(43): p. 15545-
50.
58.   Owen, G.I., et al., *Progesterone regulates transcription of the p21(WAF1) cyclin- dependent*
kinase inhibitor gene through Sp1 and CBP/p300. J Biol Chem, 1998. **273**(17): p. 10696-701.
59.   Tseng, L., et al., *Progesterone receptor (hPR) upregulates the fibronectin promoter activity in*
human decidual fibroblasts. DNA Cell Biol, 2003. **22**(10): p. 633-40.
60.   Omiecinski, C.J., et al., *Xenobiotic metabolism, disposition, and regulation by receptors: from*
biochemical phenomenon to predictors of major toxicities. Toxicol Sci, 2011. **120 Suppl 1**: p. S49-
75.
61.   Tseng, L.H., et al., *Genome-based expression profiling as a single standardized microarray*
platform for the diagnosis of endometrial disorder: an array of 126-gene model. Fertil Steril,
2010. **94**(1): p. 114-9.
62.   Hariparsad, N., et al., *Identification of pregnane-X receptor target genes and coactivator and*
corepressor binding to promoter elements in human hepatocytes. Nucleic Acids Res, 2009. **37**(4):
p. 1160-73.
63.   Altmae, S., et al., *Endometrial gene expression analysis at the time of embryo implantation in*
women with unexplained infertility. Mol Hum Reprod, 2010. **16**(3): p. 178-87.
64.   Harada, T., et al., *Apoptosis in human endometrium and endometriosis.* Hum Reprod Update,
2004. **10**(1): p. 29-38.
65.   Antsiferova, Y. and N. Sotnikova, *Apoptosis and endometrial receptivity: Relationship with in*
vitro fertilization treatment outcome. World Journal of Obstetrics and Gynecology, 2016. **5**(1): p.
87-96.

66.   Kokawa, K., T. Shikone, and R. Nakano, *Apoptosis in the human uterine endometrium during the*
menstrual cycle. J Clin Endocrinol Metab, 1996. **81**(11): p. 4144-7.

67.   Li, H., et al., *The expression and function of epithelial membrane protein 1 in laryngeal*
carcinoma. Int J Oncol, 2017. **50**(1): p. 141-148.

68.   Akilov, O.E., et al., *Resistance of Sezary cells to TNF-alpha-induced apoptosis is mediated in part*
by a loss of TNFR1 and a high level of the IER3 expression. Exp Dermatol, 2012. **21**(4): p. 287-92.

69.   Opferman, J.T. and A. Kothari, *Anti-apoptotic BCL-2 family members in development.* Cell Death
Differ, 2018. **25**(1): p. 37-45.

70.   Maybin, J.A. and H.O. Critchley, *Menstrual physiology: implications for endometrial pathology*
and beyond. Hum Reprod Update, 2015. **21**(6): p. 748-61.

71.   Bilyk, O., et al., *Epithelial-to-Mesenchymal Transition in the Female Reproductive Tract: From*
Normal Functioning to Disease Pathology. Front Oncol, 2017. **7**: p. 145.

72.   Matsuzaki, S. and C. Darcha, *Epithelial to mesenchymal transition-like and mesenchymal to*
epithelial transition-like processes might be involved in the pathogenesis of pelvic endometriosis.
Hum Reprod, 2012. **27**(3): p. 712-21.

73.   Bai, R., et al., *Down-regulation of transcription factor OVOL2 contributes to epithelial-*
mesenchymal transition in a noninvasive type of trophoblast implantation to the maternal
endometrium. Faseb j, 2018: p. fj201701131RR.

74.   Winuthayanon, W., et al., *Uterine epithelial estrogen receptor alpha is dispensable for*
proliferation but essential for complete biological and biochemical responses. Proc Natl Acad Sci
U S A, 2010. **107**(45): p. 19272-7.

75.   Dorostghoal, M., et al., *Overexpression of Endometrial Estrogen Receptor-Alpha in The Window*
of Implantation in Women with Unexplained Infertility. Int J Fertil Steril, 2018. **12**(1): p. 37-42.

76.   Trowbridge, J.M. and R.L. Gallo, *Dermatan sulfate: new functions from an old*
glycosaminoglycan. Glycobiology, 2002. **12**(9): p. 117r-25r.

77.   Kitaya, K. and T. Yasuo, *Regulatory role of membrane-bound form interleukin-15 on human*
uterine microvascular endothelial cells in circulating CD16(-) natural killer cell extravasation into
human endometrium. Biol Reprod, 2013. **89**(3): p. 70.

78.   Stacey, K., et al., *Effects of female sex hormones on lipid metabolism in the uterine epithelium of*
the mouse. Int J Biochem, 1991. **23**(3): p. 371-6.

79.   Tailor, P., et al., *The feedback phase of type I interferon induction in dendritic cells requires*
interferon regulatory factor 8. Immunity, 2007. **27**(2): p. 228-39.

80.   Hu, X., et al., *IRF8 regulates acid ceramidase expression to mediate apoptosis and suppresses*
myelogeneous leukemia. Cancer Res, 2011. **71**(8): p. 2882-91.

81.   Liu, K. and S.I. Abrams, *Coordinate regulation of IFN consensus sequence-binding protein and*
caspase-1 in the sensitization of human colon carcinoma cells to Fas-mediated apoptosis by IFN-
gamma. J Immunol, 2003. **170**(12): p. 6329-37.

82.   Yang, D., et al., *Repression of IFN regulatory factor 8 by DNA methylation is a molecular*
determinant of apoptotic resistance and metastatic phenotype in metastatic tumor cells. Cancer
Res, 2007. **67**(7): p. 3301-9.

83.   Giatromanolaki, A., et al., *Interferon regulatory factor-1 (IRF-1) suppression and derepression*
during endometrial tumorigenesis and cancer progression. Cytokine, 2004. **26**(4): p. 164-8.

84.   Kuroboshi, H., et al., *Interferon regulatory factor-1 expression in human uterine endometrial*
carcinoma. Gynecol Oncol, 2003. **91**(2): p. 354-8.

85.   Kashiwagi, A., et al., *The Postimplantation Embryo Differentially Regulates Endometrial Gene*
Expression and Decidualization. Endocrinology, 2007. **148**(9): p. 4173-4184.

86.   Kusama, K., et al., *Endometrial factors similarly induced by IFNT2 and IFNTc1 through*
transcription factor FOXS1. PLoS One, 2017. **12**(2): p. e0171858.

87.   Pon, J.R. and M.A. Marra, *MEF2 transcription factors: developmental regulators and emerging*
935   *cancer genes.* Oncotarget, 2016. **7**(3): p. 2297-312.
936   88.   Li, L., L.P. Rubin, and X. Gong, *MEF2 transcription factors in human placenta and involvement in*
937   *cytotrophoblast invasion and differentiation.* Physiol Genomics, 2018. **50**(1): p. 10-19.
938   89.   Ohlsson Teague, E.M., et al., *MicroRNA-regulated pathways associated with endometriosis.* Mol
939   Endocrinol, 2009. **23**(2): p. 265-75.
940   90.   de Wit, E. and W. de Laat, *A decade of 3C technologies: insights into nuclear organization.* Genes
941   Dev, 2012. **26**(1): p. 11-24.
942   91.   Diaz, N., et al., *Chromatin conformation analysis of primary patient tissue using a low input Hi-C*
943   *method.* Nat Commun, 2018. **9**(1): p. 4938.
944   92.   Noyes, R.W., A.T. Hertig, and J. Rock, *Dating the endometrial biopsy.* Am J Obstet Gynecol, 1975.
945   **122**(2): p. 262-3.
946   93.   Trapnell, C., L. Pachter, and S.L. Salzberg, *TopHat: discovering splice junctions with RNA-Seq.*
947   Bioinformatics, 2009. **25**(9): p. 1105-11.
948   94.   Trapnell, C., et al., *Transcript assembly and quantification by RNA-Seq reveals unannotated*
949   *transcripts and isoform switching during cell differentiation.* Nat Biotechnol, 2010. **28**(5): p. 511-
950   5.
951   95.   Langmead, B., et al., *Ultrafast and memory-efficient alignment of short DNA sequences to the*
952   *human genome.* Genome Biol, 2009. **10**(3): p. R25.
953   96.   Zang, C., et al., *A clustering approach for identification of enriched domains from histone*
954   *modification ChIP-Seq data.* Bioinformatics, 2009. **25**(15): p. 1952-8.
955   97.   Heinz, S., et al., *Simple combinations of lineage-determining transcription factors prime cis-*
956   *regulatory elements required for macrophage and B cell identities.* Mol Cell, 2010. **38**(4): p. 576-
957   89.
958   98.   Arnold, J.T., et al., *Endometrial stromal cells regulate epithelial cell growth in vitro: a new co-*
959   *culture model.* Hum Reprod, 2001. **16**(5): p. 836-45.
960   99.   Huang, W., et al., *PAVIS: a tool for Peak Annotation and Visualization.* Bioinformatics, 2013.
961   **29**(23): p. 3097-9.

Acknowledgements

We thank Dr. Sylvia Hewitt and Dr. John Lydon for editorial assistance. This work was supported by the Intramural Research Program of the National Institute of Health: Project Z1AES103311-01 (F.J.D.), R01HD067721 (S.L.Y.) and 1R01HD096266-01 (T.E.S.).

FIGURES

971   **Figure 1.**

976   **Figure 2.**

979   **Figure 3.**

F

G

H

Figure 4.

FIGURE LEGENDS

Figure 1. Genome wide PGR binding identified by ChIP-seq in endometrial tissue of fertile women during the proliferative and mid-secretory phases.

(A). Distribution of PGR binding in the genome relative to the gene body during the P and MS phase, as analyzed by PAVIS.

(B). Paired analysis was employed to identify differential PGR bound (DPRB) genomic intervals, where differential PGR binding was calculated for each of set1 and set2 (refer to Materials and Methods: *Chromatin immunoprecipitation sequencing (ChIP-seq) and qRT-PCR (ChIP-qPCR)*). The DPRB DNA common to both batches were defined as the real differential PGR bound sites. A total of 2,787 PGR bound regions were found to be in proximity of 2,249 genes (TSS ± 25 kb).

(C). The percentage of total DPRB sites that showed increased (red) or decreased (green) PGR binding transitioning from P to MS.

(D). Gene Ontology functional annotation showing enriched biological functions associated with DPRB genes (defined as DPRB within 25 kb of transcriptional start sites), as analyzed by the online bioinformatic tool DAVID.

Figure 2. Endometrial gene expression profile during the proliferative and mid-secretory phases.

(A). Gene Set Enrichment Analysis (GSEA) identified the xenobiotic metabolism pathway as significantly and positively enriched in the differentially expressed genes (DEGs), suggesting an increased activity in this pathway during MS.

(B and C). Decidualization markers *IGFBP1* and *PRL* was examined by qRT-PCR using endometrial samples from independent patients to confirm stage of menstrual cycle.

(D - G). Selected genes from the xenobiotic metabolism pathway were validated by qRT-PCR (E and G) using independent patient RNAs and presented in parallel with results from RNA-seq (D and F), n = 6, # $p < 0.05$ and * $p < 0.01$.

Figure 3. Identification of PGR regulated genes during the menstrual cycle.

(A). Overlaying the genes with DPRB and differential expression identified 653 genes during the P to MS transition.

(B). Number of genes showing increased and decreased PGR binding and expression in the endometrium during MS.

(C). The percentage of genes showing increased or decreased expression with
increased or decreased PGR binding from P to MS.

(D). Overlaying the genes with proximal constitutive and PGR binding and differential
expression identified 2,334 such genes during the P to MS transition.

(E). PGR binding activity near two known target genes, *FOSL2* and *IHH* were examined
by PGR ChIP-qPCR to confirm the phases of endometrial sample from which chromatin
was obtained. qPCR was conducted in triplicates for each sample, and results shown
are normalized to values from the P phase, n = 2 independent patients.

(F). PGR occupancy was validated for selected genes from the xenobiotic metabolism,
apoptosis and epithelial-mesenchymal transition (EMT) pathways using ChIP-qPCR.
Experiments were performed using two sets of paired patient samples (each consisting
of one P and one MS), and a representative result is shown. * $p < 0.05$.

(G). Selected genes from the xenobiotic metabolism, apoptosis and EMT pathways
were validated using qPCR, n = 6 and * $p < 0.05$.

(H). Comparison of the upstream regulator activity (as indicated by the Z-score) for
DEGs with and without differential PGR binding. Activity status (Z-score) is plotted on
the left Y-axis (blue and purple bars, representing without DPRB and with DPRB,
respectively), and significance (*p* value) is plotted on the right Y-axis (circle and square,
representing without DPRB and with DPRB, respectively).

1042

Figure 4. Epithelial functions during implantation and protein regulation of
epithelial regulators IRF8 and MEF2C.

(A). Comparison of DEGs derived from the epithelium to DEGs derived from the whole
endometrium, with a total of 658 genes that were uniquely regulated in the epithelium.

(B – C) Immunohistochemistry staining for IRF8 (B) and MEF2C (C) in human
endometrial samples during P and MS. Results show that both proteins were expressed
in both the epithelium, with increased levels of IRF8 and increased cytoplasmic-nuclear
translocation of MEF2C during the MS phase. Experiment was conducted on three
independent patients' samples and a representative is shown, alongside the negative
control stained with secondary antibody only.

1053

1054 **Supplemental Figure 1. Whole endometrial gene expression quality assessment.**

1055 (A and B). Principal component analysis (PCA, A) and hierarchical clustering (B) using
1056 the gene expression profile for each individual sample derived from P (green) and MS
1057 (red) showed a clear segregation between the two phases.

1058

1059 **Supplemental Figure 2. Epithelial gene expression quality assessment.**

1060 **(A and B).** Principal component analysis (PCA, A) and hierarchical clustering (B) using
1061 the gene expression profile for each individual sample derived from P (orange) and MS
1062 (blue) specifically in the epithelium, showing a clear segregation between the two
1063 phases.

1064

1065

TABLES

 TABLE 1. DAVID functional analysis using KEGG pathways for genes with differential
 PGR binding as determined by PGR ChIP-seq in the P and MS endometrium.

Term	P-Value	Genes
Insulin resistance	2.30E-04	*SREBF1, PIK3CG, IL6, IRS2, PTPRF, SOCS3, PRKAG2, NFKBIA, TRIB3, MAPK10, PPARGC1B, STAT3, PTPN11, SLC2A2, GFPT2, CREB3L2, CREB3L1, MLXIP, PTPN1, SLC27A3, PIK3R3, PIK3R2, PYGB*
Glycerophospholipid metabolism	2.70E-04	*CHKA, PLD1, PLB1, PISD, GPCPD1, LPIN2, LPIN1, CHPT1, LPCAT3, CDS2, GPD1L, PNPLA7, DGKD, PLA2G2A, DGKZ, LCLAT1, PLA2G2C, PLA2G2D, AGPAT3, PLA2G5, PLA2G2F*
Focal adhesion	1.70E-03	*PGF, BCAR1, PXN, CTNNB1, MYL9, COL6A6, ITGB8, PAK3, COMP, COL27A1, RAC1, PDGFC, PIK3R3, PIK3R2, PIK3CG, COL4A3, VAV3, TNXB, ACTN4, MYLK3, HGF, MAPK10, CAPN2, FLNB, COL5A1, VEGFD, LAMA3, ITGA6, RASGRF1, FYN, COL24A1, PARVB, PARVA*
Complement and coagulation cascades	3.20E-03	*PLAT, A2M, C3, C6, F13A1, C1R, BDKRB1, SERPING1, SERPINF2, SERPINE1, TFPI, SERPINA1, SERPIND1, CFD, PROS1*
Cytokine-cytokine receptor interaction	5.30E-03	*IL1R2, IL1R1, CXCR1, KITLG, CCL8, IL13, CXCR2, CXCR3, IL10, ACVR1B, CCL20, CXCR5, CXCR4, CLCF1, IL1RAP, CSF3R, PDGFC, CD27, IFNGR1, THPO, IL6, TNFSF4, HGF, TNFSF9, TNFSF8, IFNAR1, VEGFD, CCR7, TNFSF10, TNFSF11, CXCL14, PRLR, CCR2, IL22RA2*
Chemokine signaling pathway	9.50E-03	*ADCY7, BCAR1, CXCR1, CCL8, NFKBIA, CXCR2, FOXO3, CXCR3, PXN, CCL26, DOCK2, CCL20, CXCR5, CXCR4, RAC1, PIK3R3, PIK3R2, PIK3CG, VAV3, STAT1, STAT3, CCL17, CCR7, CXCL14, CCR2, IKBKG, GRK7, GRK5*
Ras signaling pathway	1.30E-02	*FGF14, PGF, KITLG, FGF12, RASAL2, PAK3, RAC1, TEK, PDGFC, RASA3, FGF1, PIK3R3, PIK3R2, PIK3CG, PLD1, HGF, MAPK10, RALGDS, PTPN11, VEGFD, PLCE1, RASGRF1, ETS1, ETS2, IKBKG, PLA2G2A, RIN1, PLA2G2C, KSR1, PLA2G2D, PLA2G5, PLA2G2F*
Proteoglycans in cancer	1.30E-02	*WNT5A, TFAP4, TLR4, MMP2, MIR21, TIMP3, PXN, CTNNB1, ANK1, CD44, RAC1, WNT6, PIK3R3, PIK3R2, PIK3CG, HSPG2, ESR1, HGF, FLNB, FZD7, ITPR1, STAT3, PTPN11, ITPR2,*

		WNT2B, CTSL, SDC1, PLCE1, WNT9B
FoxO signaling pathway	1.80E-02	*USP7, PIK3CG, IL6, IRS2, GABARAPL1, PRKAG2, SMAD3, BNIP3, MAPK10, FOXO3, IL10, STAT3, SOD2, TNFSF10, S1PR1, SETD7, FBXO32, PIK3R3, KLF2, GADD45A, PIK3R2*
Apoptosis	2.20E-02	*PIK3CG, CFLAR, TNFSF10, TNFRSF10B, CASP8, CASP12, IKBKG, NFKBIA, CAPN2, MAP3K14, PIK3R3, PIK3R2*
Prolactin signaling pathway	2.50E-02	*PIK3CG, TNFSF11, PRLR, SOCS3, SLC2A2, SOCS1, ESR1, MAPK10, FOXO3, STAT1, PIK3R3, STAT3, PIK3R2*
AMPK signaling pathway	2.60E-02	*SREBF1, PIK3CG, IRS2, PPP2R3A, PFKFB3, PPARG, PRKAG2, PFKP, FBP1, FOXO3, PPP2CB, CREB3L2, FASN, CREB3L1, PIK3R3, TBC1D1, PPP2R2C, LIPE, PIK3R2*
TNF signaling pathway	2.90E-02	*TRAF1, PIK3CG, CFLAR, IL6, SOCS3, NFKBIA, MAPK10, CCL20, CASP8, IKBKG, MAP3K8, CREB3L2, BCL3, CREB3L1, PIK3R3, MAP3K14, PIK3R2*
ECM-receptor interaction	9.40E-02	*COL4A3, TNXB, HSPG2, COL5A1, SDC1, LAMA3, ITGA6, COL6A6, CD44, ITGB8, COMP, COL27A1, COL24A1*

1070

1071

TABLE 2. Transcription factor binding sites enrichment in MS-gain intervals.

Family	Transcription factor	p-value	Motif sequence
bZIP	FOSL2, FRA1, JUN-AP1, ATF3	1.00E-67	
bZIP	CEBP, CEBP:AP1	1.00E-64	
NR	GRE	1.00E-59	
bZIP	ATF4, CEBP:AP1	1.00E-52	
NR	AR	1.00E-41	
NR	PGR	1.00E-40	
bZIP	CHOP	1.00E-37	
STAT	STAT3	1.00E-34	
STAT	STAT1	1.00E-31	
STAT	STAT5	1.00E-30	

 TABLE 3. Transcription factor binding sites enrichment in MS-loss intervals.

1075

Family	Transcription factor	p-value	Motif sequence
NR	ER	1.00E-20	
bHLH	TCF21	1.00E-11	
bHLH	ATOH1	1.00E-09	
ZF	ZBTB18	1.00E-09	
ZF	GLI3	1.00E-08	
bHLH	AP4	1.00E+07	
bHLH	NEUROD1	1.00E-06	
NR	AR	1.00E-05	
bHLH	TCF12	1.00E-05	
bHLH	ASCL1	1.00E-05	

1076

1077

1078   **TABLE 4. Gene sets enrichment analysis of the 4,576 DEG in whole endometrium.**

1079

Enriched gene sets	Enrichment	NES*	NOM p-val	FDR q-val
TNFA SIGNALING VIA NFKB	Positive	2.37	0	0
XENOBIOTIC METABOLISM	Positive	2.24	0	6.48E-04
COAGULATION	Positive	2.19	0	4.32E-04
INFLAMMATORY RESPONSE	Positive	2.15	0	3.24E-04
COMPLEMENT	Positive	1.83	0.00159236	0.012257015
INTERFERON GAMMA RESPONSE	Positive	1.77	0.0015456	0.017625704
IL6 JAK STAT3 SIGNALING	Positive	1.70	0.00980392	0.031572554
APOPTOSIS	Positive	1.54	0.02276423	0.08806576
ANGIOGENESIS	Positive	1.46	0.0970696	0.122301854
E2F TARGETS	Negative	-3.12	0	0
G2M CHECKPOINT	Negative	-3.04	0	0
MITOTIC SPINDLE	Negative	-2.50	0	0
MYC TARGETS V1	Negative	-1.65	0.01590909	0.024345282
DNA REPAIR	Negative	-0.96	0.4974359	0.69968206

1080   * NES = normalized enrichment score

1081 **TABLE 5. Genes with altered PGR binding and expression during P to MS transition.**

1082

PR Binding	Gene expression	No. of genes (total = 653)	%
Increased	Increased	441	67.53
Increased	Decreased	131	20.06
Decreased	Increased	13	1.99
Decreased	Decreased	68	10.41

1083

1084

TABLE 6. Gene sets enrichment analysis of the 653 genes with differential expression

and differential PGR binding.

Enriched gene sets	Enrichment	NES	NOM p-val	FDR q-val
COAGULATION	Positive	1.85	0.001364257	0.0448207
INFLAMMATORY RESPONSE	Positive	1.59	0.037333332	0.2136788
TNFA SIGNALING VIA NFKB	Positive	1.59	0.026041666	0.1491825
XENOBIOTIC METABOLISM	Positive	1.57	0.03547963	0.1247853
EPITHELIAL MESENCHYMAL TRANSITION	Positive	1.55	0.038208168	0.1259948
COMPLEMENT	Positive	1.38	0.11479945	0.2892615
APOPTOSIS	Positive	1.37	0.092369474	0.2594335
HYPOXIA	Positive	1.31	0.1342711	0.3065951
INTERFERON GAMMA RESPONSE	Positive	1.29	0.17036012	0.302116
ESTROGEN RESPONSE LATE	Positive	1.15	0.31600547	0.4711447
IL2 STAT5 SIGNALING	Positive	1.13	0.31117022	0.4615895
P53 PATHWAY	Positive	1.12	0.32647464	0.4400889
MTORC1 SIGNALING	Positive	0.89	0.6025825	0.7500677
ESTROGEN RESPONSE EARLY	Positive	0.86	0.6555407	0.7511974
IL6 JAK STAT3 SIGNALING	Positive	0.76	0.7735584	0.8316847

Table 7. Gene sets enrichment analysis of the 2,334 DEGs in whole endometrium with constitutive PGR binding.

Enriched gene sets	Enrichment	NES	NOM p-val	FDR q-val
TNFA SIGNALING VIA NFKB	Positive	2.27366	0	0.002744444
INFLAMMATORY RESPONSE	Positive	2.2703116	0	0.001372222
HYPOXIA	Positive	2.1009197	0	0.004157759
INTERFERON GAMMA RESPONSE	Positive	1.9754444	0	0.01090216
IL6 JAK STAT3 SIGNALING	Positive	1.8338909	0.017621145	0.02559644
XENOBIOTIC METABOLISM	Positive	1.6081412	0.025882352	0.10718091
KRAS SIGNALING UP	Positive	1.5911685	0.03671706	0.10251326
UV RESPONSE DN	Positive	1.5904794	0.015521064	0.09035362
COMPLEMENT	Positive	1.5408636	0.04405286	0.10960495
E2F TARGETS	Negative	-3.255928	0	0
G2M CHECKPOINT	Negative	-3.1368256	0	0
MITOTIC SPINDLE	Negative	-2.8043678	0	0
HEDGEHOG SIGNALING	Negative	-1.9978325	0	0.002890576
EPITHELIAL MESENCHYMAL TRANSITION	Negative	-1.4042959	0.06571936	0.14482453

1093 **TABLE 8. Gene sets enrichment analysis of the 3,052 DEGs in the epithelium.**

1094

Enriched gene sets	Enrichment	NES	NOM p-val	FDR q-val
COAGULATION	Positive	2.4593391	0	0
COMPLEMENT	Positive	2.3820252	0	0
TNFA SIGNALING VIA NFKB	Positive	2.260513	0	0
INFLAMMATORY RESPONSE	Positive	2.2356772	0	0
XENOBIOTIC METABOLISM	Positive	2.1790328	0	0.000179012
HYPOXIA	Positive	1.9205347	0	0.008392723
APOPTOSIS	Positive	1.8877827	0.001414427	0.009475978
INTERFERON GAMMA RESPONSE	Positive	1.8587548	0	0.011746863
KRAS SIGNALING UP	Positive	1.8500544	0.004172462	0.011368177
EPITHELIAL MESENCHYMAL TRANSITION	Positive	1.6994203	0.004178273	0.03300058
ANGIOGENESIS	Positive	1.4216607	0.09548611	0.15824698
APICAL JUNCTION	Positive	1.2970207	0.15987934	0.27779698
P53 PATHWAY	Positive	1.2064549	0.21958457	0.38236877
E2F TARGETS	Negative	-3.7750194	0	0
G2M CHECKPOINT	Negative	-3.4905815	0	0
MYC TARGETS V1	Negative	-2.5305622	0	0
MITOTIC SPINDLE	Negative	-2.4694543	0	0
ESTROGEN RESPONSE LATE	Negative	-1.8183266	0	0.009814334
DNA REPAIR	Negative	-1.0963912	0.32970026	0.43156412
MTORC1 SIGNALING	Negative	-0.7685945	0.8393939	0.88358915
ESTROGEN RESPONSE EARLY	Negative	-0.7567437	0.8778878	0.8235289

1095

1096

TABLE 9. Canonical pathway analysis of the epithelium specific DEGs using IPA.

Ingenuity Canonical Pathways	-log(p-value)	z-score	Molecules in dataset
Dermatan Sulfate Biosynthesis	3.67	-	*CHST1,HS6ST1,CSGALNACT2,SULT1A1,CHST10,HS6ST3,DSEL,SULT2B1*
Cholesterol Biosynthesis I	3.46	-	*FDFT1,EBP,MSMO1,CYP51A1*
Chondroitin Sulfate Biosynthesis (Late Stages)	3.41	-	*CHST1,HS6ST1,CSGALNACT2,SULT1A1,CHST10,HS6ST3,SULT2B1*
Superpathway of Cholesterol Biosynthesis	3.03	-	*FDFT1,EBP,MSMO1,HMGCS1,CYP51A1*
Chondroitin Sulfate Biosynthesis	3.01	-	*CHST1,HS6ST1,CSGALNACT2,SULT1A1,CHST10,HS6ST3,SULT2B1*
Osteoarthritis Pathway	2.18	2.496	*CXCL8,MTOR,FRZB,SMAD3,BMP2,ITGA2,BMPR2,WNT16,SOX9,MEF2C,HES1,ACAN,MMP1*
Cholecystokinin/Gastrin-mediated Signaling	2.15	2.121	*GAST,MAPK14,RHOB,MEF2D,CREM,MEF2C,GNA13,PRKCG*
IL-8 Signaling	2.03	2.309	*CXCL8,PIK3CA,MTOR,FLT1,RHOB,HBEGF,CXCL1,GNA13,KDR,MAP4K4,PRKCG,EIF4EBP1*
TGF-β Signaling	1.97	1.342	*IRF7,MAPK14,BMP2,SMAD3,BMPR2,TGIF1,INHBA*
PTEN Signaling	1.74	-2.121	*PIK3CA,FLT1,ITGA2,PREX2,BMPR2,KDR,BCL2L11,PDGFRB*

TABLE 10. Upstream regulators with specific actions in the epithelium (identified using IPA).

1103

Upstream Regulator	Log2(FC) in epithelium	Molecule Type	Activation z-score	p-value of overlap
POU5F1	1.696	transcription regulator	2.429	3.93E-06
IRF5	1.526	transcription regulator	2.266	2.04E-04
MEF2C	1.098	transcription regulator	1.835	6.15E-03
MEF2D	1.014	transcription regulator	2.478	1.02E-03
IRF8	1.199	transcription regulator	1.608	3.35E-05
FOXJ1	-1.613	transcription regulator	-1.96	3.52E-02
TLR5	1.34	transmembrane receptor	3.062	7.08E-04
IL1R1	1.745	transmembrane receptor	2.872	2.27E-02
FCGR2A	2.025	transmembrane receptor	1.544	1.30E-04
MET	1.993	kinase	2.054	1.26E-09
AURKB	-3.136	kinase	-2.132	2.99E-03
HBEGF	1.739	growth factor	1.754	1.94E-05
WNT7A	-2.761	Wnt ligand	-1.98	2.60E-02
CYP27B1	-1.773	enzyme	-1.51	9.40E-03
DLL4	1.067	other	2.119	1.74E-05

1104

1105

1106