First detection of Infectious Spleen and kidney Necrosis Virus (ISKNV)

associated with massive mortalities in farmed tilapia in Africa

Short running tittle: First detection of ISKNV in farmed tilapia in Africa

José Gustavo Ramírez-Paredez[a]†, Richard K. Paley[bc]†, William Hunt[a], Stephen W. Feist[bc]

David M. Stone[bc], Terence R. Field[a], David J. Haydon[a], Peter A. Ziddah[d], Mary Nkansa[d],

Emanuel K. Pecku[e], Joseph A. Awuni[e], James Guilder[bc], Joshua Gray[b], Samuel Duodu[f],

Timothy S. Wallis[a], David W. Verner -Jeffreys[bc]∗

[a] Ridgeway Biologicals Limited a Ceva Santé Company, Units 1-3 Old Station Business Park,

Compton, Berkshire, England, United Kingdom

[b] Cefas Weymouth Laboratory, Weymouth, Dorset, England, United Kingdom.

[c] OIE Collaborating Centre for Emerging Aquatic Animal Diseases.

[d] Fisheries Commission, Ministry of Fisheries and Aquaculture Development, Accra, Ghana

[e] Veterinary Services Directorate, Accra, Ghana

[f] University of Ghana, Accra, Ghana

†Both authors contributed equally to this study

*Corresponding author Dr D. W. Verner-Jeffreys Cefas Weymouth Laboratory, Weymouth,

Dorset, England, United Kingdom

Summary

In late 2018, unusual patterns of very high mortality (>50% production) were reported in intensive

tilapia cage culture systems across Lake Volta in Ghana. Samples of fish and fry were collected

and analysed from two affected farms between October 2018 and February 2019. Affected fish

showed darkening, erratic swimming and abdominal distension with associated ascites.

Histopathological observations of tissues taken from moribund fish at different farms revealed

lesions indicative of viral infection. These included haematopoietic cell nuclear and cytoplasmic

pleomorphism with marginalisation of chromatin and fine granulation. Transmission electron

microscopy showed cells contained conspicuous virions with typical Iridovirus morphology i.e.

enveloped, with icosahedral and or polyhedral geometries and with a diameter c.160 nm. PCR

confirmation and DNA sequencing identified the virions as Infectious Spleen and Kidney

Necrosis Virus (ISKNV). Samples of fry and older animals were all strongly positive for the

presence of the virus by qPCR. All samples tested negative for TiLV and Nodavirus by qPCR. All

samples collected from farms prior to the mortality event were negative for ISKNV. Follow up

testing of fish and fry sampled from 5 additional sites in July 2019 showed all farms had fish that

were PCR positive for ISKNV, whether there was active disease on the farm or not, demonstrating

the disease was endemic to farms all over Lake Volta by that point. The results suggest that

ISKNV was the cause of disease on the investigated farms and likely had a primary role in the

mortality events. A common observation of coinfections with *Streptococcus agalactiae* and other

tilapia bacterial pathogens further suggests that these may interact to cause severe pathology,

particularly in larger fish. Results demonstrate that there are a range of potential threats to the

sustainability of tilapia aquaculture that need to be guarded against.

Keywords: Tilapia; Africa; Tilapia Viral Diseases; fish virus; Infectious Spleen and Kidney

Necrosis Virus; ISKNV;

Introduction

The farming of tilapia species (*Oreochromis* spp.) is one of the most important sectors in aquaculture worldwide with total global production estimated at more than 6 686 000 tonnes in 2016 (FAO, 2018). In Africa, production is still dominated by Egypt with more than 1 000 000 tonnes produced in 2017 (1). However, tilapia culture has become increasingly important in several other African countries, where it boosts the local economy and constitutes an affordable source of animal protein for human consumption.

In Ghana, Nile tilapia production in 2016 had reached more than 50 000 tonnes, from only 2 000 tonnes per year in 2006 (FAO, 2016) with more than 90% of the production derived from high stocking density floating cage systems in Lake Volta. However, as production systems have intensified in the area, the industry has been increasingly affected by a range of disease issues (Jansen, Cudjoe, & Brun, 2018; Verner-Jeffreys et al., 2018).

In 2017 (Verner-Jeffreys et al., 2018) conducted the first comprehensive disease investigation in tilapia farmed in Lake Volta Ghana. *Streptococcus agalactiae* multilocus sequence type 261 was shown to be a major cause of mortality for farmed Nile tilapia and a range of other bacterial and parasitic pathogens including *Aeromonas* spp., *Streptococcus iniae*, *Flavobacterium columnare* and a *Myxobolus* sp. were also detected.

Jansen et al. (Jansen et al., 2018) conducted follow up studies, including a broad-ranging epidemiological investigation and suggested that mortalities caused by bacteria in Lake Volta were not a major concern for the local economy as farmers had managed to sustain losses by increasing production of fingerlings, treatment with antibiotics and use of autogenous vaccines.

Historically, bacterial infections were the major threat for the health of farmed tilapia (Plumb & Hanson, 2011). However, in recent years a number of viral diseases have emerged worldwide with

devastating effects for the industry(Machimbirike et al., 2019). Tilapia Lake Virus (TiLV) is

considered the main viral challenge to tackle, as it has spread to many producing countries,

causing high mortalities in all production stages. Although TiLV has been detected elsewhere in

Africa (Hounmanou et al., 2018), the initial study by (Verner-Jeffreys et al., 2018), diagnostic

investigations undertaken by Ridgeway Biologicals Ltd. (Ramirez *et al*. unpublished data), or the

more recent survey undertaken by (Jansen et al., 2018), all failed to provide any evidence of TiLV

in diseased tilapia reared in Lake Volta. Up until September 2018, with the exception of the

detection of a nodavirus sequence by (Verner-Jeffreys et al., 2018), no significant association

between viral agents and tilapia mortality events have been demonstrated in Ghana.

From September 2018 to March 2019 outbreaks of disease with very high levels of morbidity and

mortality (60-90%), were experienced in both vaccinated and unvaccinated tilapia by farmers in

Lake Volta. In late-September 2018, a farm located below the lower dam in the region of

Asutsuare, was the first to suffer episodes of massive acute mortalities (https://goo.gl/LmqbG2

and https://bit.ly/2NwDEbD). Approximately a week after the first report, a second farm located

in the Akuse region (~5 km upstream of Asutsuare, but still below the lower dam) also

experienced acute mortalities. By mid-October, multiple floating cage-based farms in the Dodi

region (above the upper dam) reported losses of more than 10 tonnes per day

https://goo.gl/yj4oT4. In late November, farmers that had been unaffected in the Asikuma region

(downstream of Dodi but still above the upper dam) also started to suffer episodes of massive

acute mortality. (**Figure 1**).

By the end of 2018, most tilapia farmers in Lake Volta had reported mortalities that they were not

able to contain by increased production of fingerlings or treating with antibiotics (**Supplementary**

File 1, https://bit.ly/2KZFFuw and https://goo.gl/cPmpSE). Mortality events continued into and

throughout 2019. We report the results of a comprehensive disease investigation, conducted at

seven affected farms from two different regions of Lake Volta, to gain insights into the causes of

these   mortalities.

Figure 1 Map of lower region of Lake Volta in Ghana, West Africa. Red triangles indicate the regions in chronological order (A to D) where the outbreaks of mortality occurred.

2 Materials and Methods

2.1 Sampling

Farm 1, a medium size (approximately 600-800 tonnes per annum production) cage culture operation on upper Lake Volta, was visited on the 18/10/18. Samples from 10 fish (average weight 200g) were taken for bacteriology. In addition five sets of samples containing liver and brain from fish from different cages were collected for molecular diagnostics as detailed in **Table 1**.

Farm 2, a large cage culture operation (>2000 tonnes per annum), was visited for sampling on 28/11/18, 17/12/18 and 20/02/19. For the first two visits, moribund fish between 40-646g collected from cages on the main lake were sampled. In the first visit for bacteriology and virology on the second visit for virology only. For the third visit, responding to reports that they were now experiencing very heavy mortalities in their fry production units (>70%), samples of moribund fry and juveniles from both nursery cages on the main lake and from their onshore hatchery supplied with water pumped from the main lake, were analysed for virology. From this farm material was also taken for histological and molecular diagnostic investigations during the visits as detailed in **Table 1 and Supplementary File 2**. During the visits, semi-structured interviews were carried out with farm managers and/or workers. Interview questions were primarily constructed to ascertain trends in mortality levels since September 2019, any observed clinical signs in this time and whether there were any differences in impact associated with fish life stage and/or system. Additionally, relevant information on potential risk factors, mitigation measures and biosecurity practices, as well as any further farmer concerns, were discussed.

A further 5 farms in the Akosombo, Atimpoku and Dasasi regions were visited and sampled for virology, bacteriology, histology and molecular diagnostic investigation from 9-

10/07/2019. These included fry, on-growing and broodstock fish from farms of varying

capacity and with either no reported disease, fish that had survived recent mortality events on

farms or fish with ongoing clinical signs or mortalities.

All moribund fish from the visited farms were humanely euthanised with a lethal overdose of

tricaine methanesulfonate 1,000 mg/g (Pharmaq, Hampshire, UK) followed by brain

destruction prior to the necropsy.

2.2 Bacteriology

Samples for bacteriology were collected from the brain, liver, kidney and spleen with sterile

cotton swabs and inoculated onto tryptone soya agar (TSA), Columbia blood agar (CBA),

Tryptone yeast extract salts agar (TYES) (Southern Group Laboratory, Corby, UK) and

cystine heart agar with 2% bovine haemoglobin (CHAH) (Becton Dickinson, Oxford, UK).

All inoculated agar plates were incubated at 28 °C for 24-72 hours. Colonies assessed as

significant based on occurrence and dominance were subcultured to purity on similar media.

Pure relevant isolates were initially identified by morphology and Gram staining. The partial

16S rRNA genes of the Gram negative isolates identified were PCR amplified and sequenced

using the method described by (Klindworth et al., 2013). Gram positive cocci forming chains

were screened using a *Streptococcus agalactiae* specific capsular typing multiplex PCR

developed by (Shoemaker, Xu, Garcia, & Lafrentz, 2017). The Gram negative isolates strains

confirmed as *Aeromonas* spp. based on partial 16S rRNA gene sequence analysis were

further characterised based on partial *gyrB* sequencing analysis for identification at the

species level as described by (Yáñez, Catalán, Apráiz, Figueras, & Martínez-Murcia, 2013).

2.3 Histopathology and Electron Microscopy

Tissues were fixed in neutral buffered formalin (NBF) for a minimum of 24 hr before being

placed in glycerol diluted 50:50 with phosphate buffered saline (PBS) for transportation to

Cefas. On receipt, tissues were rinsed in 70% alcohol and placed again in NBF for a final

period of fixation prior to processing using standard protocols in a vacuum infiltration

processor. Tissue sections were cut at a thickness of 3-4 µm on a Finnese® microtome,

mounted on glass slides and stained with haematoxylin and eosin using an automated staining

protocol. Stained sections were examined for general histopathology by light microscopy

(Nikon Eclipse E800). Digital images and measurements were obtained using the Lucia[TM]

Screen Measurement software system (Nikon, UK).

For electron microscopy, small samples of tissues fixed in NBF as above were rinsed three

changes of 0.1 M sodium cacodylate buffer, followed by post fixation in 2.5% glutaraldehyde

in the same buffer for 1 hour prior to a second post fixation for 1 hour in 1 % osmium

tetroxide in 0.1 M sodium cacodylate buffer. Subsequently, fixed tissues were dehydrated in

an ascending acetone series acetone series and embedded in epoxy resin 812 (Agar Scientific

pre-Mix Kit 812, Agar scientific, UK) and polymerised at 60 °C overnight. Semi-thin (1 µm)

survey sections were stained with 1 % Toluidine Blue and examined by light microscope to

identify areas of interest. Ultrathin sections (70-90 nm) of the targeted areas were placed on

uncoated copper grids and stained with uranyl acetate and Reynold's lead citrate (Reynolds

1963). Grids were examined using a JEOL JEM 1400 transmission electron microscope and

digital images captured using a GATAN Erlangshen ES500W camera and Gatan Digital

Micrograph[TM] software.

2.4 Molecular diagnosis (viral and bacterial detection)

The samples collected for molecular diagnosis were washed twice in 750µl of sterile 1x PBS

to remove the RNA-*later*® and homogenised. Total nucleic acids were extracted using

nanomagnetic beads i.e. Genesig Easy DNA/RNA Extraction Kit (Primerdesign,

Southampton, UK) and stored until further use.

Multiplex PCR for detection of Streptococcus spp.

Nucleic acids extracted were used as a template on a multiplex PCR (unpublished data) to

confirm the presence of *Streptococcus* spp. that had been previously reported as fish

pathogens including *Streptococcus agalactiae* (Delannoy et al., 2013), *Streptococcus uberis*

(Luo et al., 2017) and *Streptococcus dysgalactiae* (Abdelsalam, Asheg, & Eissa, 2013).

qPCR for detection of TiLV and Nodavirus

Nucleic acids were used for the detection of tilapia lake virus and nodavirus by quantitative

PCR using the commercial kits: Path-TiLV-EASY and Path-Betanodavirus-EASY

(Primerdesign, Southampton, UK) in the platform Genesig q16® (Primerdesign,

Southampton, UK) as per the protocols suggested by the manufacturer.

Conventional PCR for detection of megalocytiviruses

The generic PCR protocol for notifiable aquatic megalocytiviruses (Red Seabream Iridoviral

disease / Infection spleen and kidney necrosis virus (J Kurita, Nakajima, Hirono, & Aoki,

1998; World Organisation for Animal Health OIE, 2018)) was initially used, to screen the

samples of fish collected from visit 2 at Farm 2. For this, genomic DNA was extracted as

follows: the RNA-*later*® was removed and the tissue samples weighed. Depending on the

weight of the tissue available the samples were diluted in RTL buffer (Qiagen) to provide

either a 1:10 w/v or a 1:5 w/v and homogenised per fish i.e. all the organs of each fish into an

individual pool using Matrix A and the FastPrep apparatus (MP Biomedicals). Following

homogenisation, the samples were diluted further with RTL buffer to give a 1:60 w/v

homogenate and total nucleic acid was extracted from 300 μl of the clarified sample using the

RNA tissue mini kit without DNase (Qiagen) and eluted in a 60 μl volume.

RT was performed at 37°C for 1 h in a 20 μl volume consisting of 1× M-MLV RT reaction

buffer (50 mM) Tris pH 8.3, 75 mM KCl, 10 mM DTT, 3 mM MgCl2) containing 1 mM

dNTP, 100 pmol random primers, 20 U M-MLV reverse transcriptase (Promega,

Southampton UK) and 4μl of the nucleic acid extracted above.

PCR was performed in duplicate in a 50 μl reaction volume with 2.5 μl of cDNA of total

nucleic acid, 25 mM dNTPs, 1 x GoTaq® buffer (2.5 mM MgCl2 solution), 5 pmol of each

primer   (C1105   5'-GGTTCATCGACATCTCCGCG-3'   and   C1106   5'-

AGGTCGCTGCGCATGCCAATC-3') and 1.25 units of GoTaq® DNA polymerase

(Promega). The cycling conditions were as follows: 40 temperature cycles (1 min at 95°C, 1

min at 55°C and 1 min at 72°C) after an initial denaturing step (5 min at 95°C) followed by a

final extension step of 10 min at 72°C.

Amplified products were electrophoresed in a 2% (w/v) agarose/TAE (40 mM Tris-acetate,

pH 7.2, 1 mM EDTA) gel containing 1.0 μg ml-1 ethidium bromide at 120 v for 30 mins and

viewed under UV light.

PCR products were excised from the gel and the DNA was extracted and purified by ethanol

precipitation. Both strands of the PCR product were sequenced using the ABI PRISM Big

Dye Terminator v3.1 cycle sequencing kit and the same primers used for the amplification.

The forward and reverse sequences were aligned and a consensus sequence generated using

the CLC software (Qiagen). Generated consensus sequences were compared with sequences

from GenBank using BLASTN (Altschul et al., 1997) and aligned using the MUSCLE

application of the MEGA software version 6 (Tamura, Stecher, Peterson, Filipski, & Kumar,

2013).

In addition, the OIE recommended PCR protocol for notifiable aquatic megalocytiviruses

(World Organisation for Animal Health OIE, 2018) developed by (J Kurita et al., 1998) was

used was used to screen total nucleic acids extracted from the rest of the samples fixed for

viral molecular analyses.

In analyses from farms where clinical disease was not observed, a second round PCR using

the nested primers C1073 5'-AATGCCGTGACCTACTTTGC-3' and C1074 5'-

GATCTTAACACGCAGCCACA -3' (15) was employed.

qPCR for detection and quantification of megalocytiviruses

The amount of virus present in the samples was also investigated by qPCR. For this the

homogenised tissues were subjected to total nucleic acids extraction (~20mg of each organ)

using the Genesig Easy DNA/RNA Extraction Kit (Primerdesign) as described earlier. The

extracted nucleic acids were tested using the commercial kit Path-ISKNV-EASY

(Primerdesign, Southampton, UK) in the platform Genesig q16® (Primerdesign) as per the

manufacturer instructions. This detects both red sea bream iridovirus and ISKNV variants. In

all cases fish were individually analysed either by pooling liver, spleen and brain or screening

individual tissues.

Retrospective analyses of archived samples by qPCR

A total of 16 samples of archived tissue homogenates from 5 different farms (that included

Farms 1 and 2) were retrospectively screened for ISKNV by qPCR with the commercial kit

Path-ISKNV-EASY as described before. From these 7 had been collected during 2017 and

the rest in March 2018 (**Supplementary File 5**). All the samples had been previously

confirmed as negative for TiLV and Nodavirus using the commercial kits Path-TiLV-EASY

and Path-Betanodavirus-EASY.

Virus isolation

Frozen spleen and kidney tissue or whole fry of fish showing clinical signs taken from 2 farm

sites on Lake Volta on 10 July 2019 were homogenised with sand and pestle and mortar in

1:10 w/v cell culture transport medium (L-15 plus 1% antibiotic antimycotic solution, Gibco).

Homogenate was clarified by centrifugation for 10 min at 3000g, inoculated at 1:100 and

1:1000 final dilutions onto GF, BF-2 and E-11 cells in 24 well cell culture plates (Gibco) and

incubated at 25°C. After 7 days cells were blind passaged and incubated for a further 7 days.

Cells were observed for cytopathic effect (CPE) by light microscopy with phase contrast

(IX83 inverted microscope, Olympus, UK).

3 Results

3.1 Farm visits

On the first visit to Farm 2 on the 27/11/18, farm staff reported very high and ongoing

mortalities (Figure 2) in fish bigger than 20g, including broodstock, but no significant losses

in fingerlings. Losses reportedly peaked at about 670 crates (equating to approximately 40

tonnes per day) shortly after this visit on the 2/12/2018 **(Figure 2)**.

Losses were so severe that accurate estimation was not possible, with more than 50 additional

labourers recruited locally just to remove dead and moribund fish during the peak period. By

the second visit to Farm 2, losses of ongrowing fish had reportedly declined back to the

background 10-20 crates per day more typically observed i.e. less than 1-2 tonnes per day

(**Figure 2**).

256
257 **Figure 2. Number of crates of dead/ moribund fish removed from Farm 2 during period**
258 **of maximum losses.** Daily mortalities were estimated based on numbers of crates of rejected
259 ongrowing tilapia collected each day by farm staff. Each crate typically contained
260 approximately 60kg tilapia collected from the cages and rejected because they were either
261 dead, moribund or displayed other adverse signs that prevented their sale.
262

During the first two visits to the farms, but particularly to Farm 2, diseased fish were

observed swimming away from the school with erratic swimming i.e. on one side, in circles,

lethargic, with no equilibrium, upside down etc. **(Supplementary File 3).**

Externally, the fish displayed a range of clinical signs, including skin nodules, frayed fins,

loss of eyes, opaque eyes, loss of scales, exophthalmia, anorexia, decolouration or darkened

skin, excess of mucous, skin haemorrhages and distended abdomen (**Figure 3**). At necropsy,

fish from both the first visits to Farm 1 and 2 presented with marked ascites, enlarged and

haemorrhagic organs including the spleen, heart, brain, gills, but most notably liver and

kidney. The gastrointestinal track was empty of solids but contained transparent fluid similar

to that also seen in the peritoneal cavity.

In contrast to the earlier visits, when Farm 2 was visited on 20/02/19, there were very high

and ongoing mortalities in the fry production systems (>70%). This was both in their onshore

hatchery (supplied with water pumped from the main lake) where eggs were hatched and held

until the fry were approximately 20g, and in the nursery cages on the main lake to which fry

had been transferred. As with the larger fish sampled, affected fry showed erratic swimming

behaviour, skin haemorrhages and severe ascites as the main clinical signs (**Figure 3**).

279
Figure 3 External lesions and clinical signs in diseased Nile tilapia in Lake Volta. A
Ongrowing fish with emaciation slight ascites (arrow), endophthalmia (left arrow head) and
skin purulent abscess (right arrow head). **B** Broodstock with microphthalmia left arrow head),
skin haemorrhages (middle arrow head) and skin ulcers (right arrow head). **C** juvenile with
exophthalmia (left arrow head), ascites (arrow) and loss of scales, excess of mucus and
haemorrhages (right arrow head). Bar = 1cm. **D** Ventral view of juvenile fish presenting
severe ascites black arrow and skin haemorrhages (arrow head). Bar= 1cm.
287
288

3.2 *Megalocytivirus* and bacterial infections in Farm 2 during period of high mortalities

Four out of the ten fish examined histologically showed mild tissue necrosis in the spleen and

renal haematopoietic tissue with the presence of large numbers of cells showing relative

eosinophilia cytoplasmic and nuclear pleomorphism with margination of chromatin in some

affected nuclei suggestive of a viral infection diffused throughout the tissue (**Figure 4**).

Figure 4 Histopathological cross sections of tissues from diseased Nile tilapia in Lake Volta. All
sections stained with H&E unless otherwise stated. **A.** Section through the intestine of an infected fry
showing the presence of large numbers of megalocytes in the lamina propria (*) but not affecting the
mucosa or underlying muscularis. Note the presence of isolated megalocytes in the pancreatic acinar
tissue (arrows). Bar = 50µm. **B.** Same fish as in A, showing infiltration and disruption of the
muscularis associated with the presence of megalocytes (arrows). Bar=50 µm.   **C.** Pseudobranch
showing extensive necrosis (*). Bar=50µm. **D.** Heart showing necrosis of the ventricular muscle (*),
particularly in the peripheral regions associated with inflammation. Bar=50µm. **E.** Spleen from a fish
from Farm 2 during a mortality episode. Affected cells show pronounced eosinophilia and are
distributed throughout the section (*). Numerous vacuolated macrophage aggregates are also present
(M). Bar = 50µm. **F.** Pronephros showing extensive distribution of megalocytes without associated
tissue necrosis (*). Bar=100µm. Inset shows pale staining of DNA in the cytoplasm of affected cells
(arrows). Feulgen stain. Bar=25µm.

The ultrastructure of affected cells revealed the presence of conspicuous viral particles. Some

cells showed the presence of numerous diffusely spread virions within the hypertrophied

nucleus of affected cells, which also showed degradation of the nuclear membrane (**Figure**

5). Virions were approximately 160 nm in diameter (**Figure 5D**). Virion morphology

showing icosahedral symmetry with an external double membrane and internal core was

consistent with that of viruses from the genus *Megalocytivirus*. In other cases, the nucleus of

affected cells appeared condensed and densely stained in histological and resin sections.

TEM showed that in these cases the nucleus was tightly packed with virions in various stages

of maturation and with some evidence of formation of 'arrays' (**Figure 5**). In two of these

cases, a concomitant Gram +ve bacterial infection was also present in the gill and liver.

Incidental findings of gill parasites, myxozoan cysts and monogeneans, both present in low

numbers as well as low grade epitheliocystis were observed. The brain of a single fish

harboured small cysts containing necrotic debris. Gram and Ziehl-Neelsen staining did not

demonstrate the presence of bacteria. Other tissues appeared normal. For the set of samples

collected at the height of the mortalities from Farm 2 on 28/11/18, *Aeromonas jandaei*,

Aeromonas veronii (from skin), *Streptococcus agalactiae* capsular type Ib biotype 2 (non

haemolytic) and *Edwardsiella tarda* (from liver and kidney) were recovered.

3.3 PCR and sequence confirmation of Megalocytivirus infection

Within the 7 individuals collected from Farm 2 on the second visit that were analysed with

the protocol proposed by (Rimmer, A. E., Becker, Tweedie, & Whittington, 2012), a single

fish (fish 1) was clearly positive by PCR for RSIV/ISKN and a second very weak product of

the correct size was also seen in tissues from Fish 6 (**Supplementary File 4**) The consensus

sequence generated from the PCR product from Fish 1 was confirmed as ISKNV sharing

100% nucleotide identity with ISKNV accession no AF371960.1. In the phylogenetic

analysis the sequence was assigned to the same lineage as the bulk of the ISKNV sequences

(**Figure 6**).

The samples collected from Farm 1 and Farm 2 at the height of the mortalities were found to

be strongly positive when they were retrospectively tested using the (J Kurita et al., 1998;

World Organisation for Animal Health OIE, 2018) current OIE recommended PCR method.

Figure 5 Micrographs of the kidney of diseased Nile tilapia in Lake Volta. A Semithin
section of affected renal tissue showing characteristic cellular hypertrophy (black arrows).
Toluidine Blue stain. Scale bar = 25μm. **B.** Electron micrograph of an individual infected cell
with numerous viral particles (*). Adjacent cells appear uninfected. Scale bar = 2μm. **C.**
Numerous lightly stained developing virions with mature virions. The outer membranes and
central electron lucent core are clearly visible. Scale bar = 500nm. **D.** Mature icosahedral
virion showing detail of the outer caspid and inner membrane with central electron lucent
core.

349
3.4 qPCR results for megalocytivirus from ongrowing tilapia samples

The qPCR results confirmed that all the fish sampled from Farm 1 in the Akuse region were

positive for megalocytivirus. Also, as expected, the samples collected from the first Farm 2

visit during the peak of mortality (28/11/18), were also positive and presented the highest

viral titres in grow out fish with some containing over 3×10^6 copies per sample reaction. In

contrast, the samples collected from grow out fish during the second visit to Farm 2 had

much lower viral copy numbers.

All the archived samples collected in 2017 and March 2018 were negative for ISKNV, when

tested by qPCR (**Supplementary File 5**).

3.5 Fry samples had characteristic megalocytivirus pathology and high copy numbers of

virus

All the fry samples collected from Farm 2 on 20/02/19, when there were reportedly very high

(>90%) losses in that part of their system, were positive for ISKNV by qPCR and these

presented the highest titres in the study with some containing up to 1.5×10^7 copies per

sample reaction (**Table 1**). All fry showed moderate to marked histological and pathological

features of infection with *megalocytivirus*. Splenic tissues were necrotic and associated with

the presence of megalocytes characterised by light sometimes granular cytoplasmic

basophilia and hypertrophied nuclei. Kidney also showed the presence of megalocytes but

usually with only mild cellular necrosis. The lamina propria in the intestine of a single fish

was packed with megalocytes (**Figure 4**), although necrosis appeared to be absent and the

epithelial layer remained intact. Gills showed only minimal focal necrosis, usually affecting

the underlying connective tissues. In some fish the choroidal rete was affected with mild

necrosis and variable numbers of megalocytes and most fish showed mild myositis with few

megalocytes in the skeletal muscle. However, a single fish showed extensive inflammation

and myofibrillar necrosis **(Figure 4)**. Connective tissues of the head and in particular around

the pharyngeal teeth were often infiltrated with megalocytes. Brain and spinal cord appeared

normal. Liver samples were not examined as they were used for virus quantification.

3.6 Other pathological observations

Megalocytivirus -like pathology was not observed in any of the samples taken from the

second visit to Farm 2, two weeks after the peak of mortalities had passed. As with the

samples taken at the first visit, there was evidence of bacterial infection in some individuals,

particularly fish 5, had marked bacterial infection of the spleen, liver and brain (meningitis).

All the samples were positive for the presence of *Streptococcus agalactiae* and negative for

Streptococcus uberis and S*treptococcus dysgalactiae* by PCR **(Supplementary File 6)**.

A range of different potential bacterial species, including *Aeromonas jandaei* and

Plesiomonas shigelloides were recovered from fish from Farm 1 **(Table 1)**, but not as pure

growths or high quantities, suggesting they had a limited role in observed disease in these

animals.

All the samples from Farm 1 and Farm 2 tested for TiLV and from Farm 2 for nodavirus by

qPCR were all negative **(Table 1 and Supplementary File 4)**.

The results for all the individual fish tested are shown in Supplementary File 2.

3.7 Follow up testing in July 2019

A further 5 farms sampled in July 2019 were tested using nested conventional PCR (15).

Fish on farms Farms 6 (fry) and 7 (lake ongrowers) were experiencing ongoing mortality and

showing typical clinical signs described above at time of sampling. All samples were

strongly positive in a single round assay and virus was isolated in cell culture from both

farms.   For   samples from farms 4 (fry) and 5 (lake ongrowers) which both were

recovering from recent mortality events, but had no remaining observable clinical disease,

between 40-100% of samples tested positive but these were in second round only of nested

assay and in some cases only in one of duplicate reactions indicating low levels of virus.

From farm 3 (all age groups) for which no mortality events had been reported between 30%

to 100% of samples were positive but all in the second round only.   On sequencing

representatives from all positive farms showed identical sequence (data not shown).

Sampling date & Farm	No. of Fish and size	Farm observations and ‡clinical signs	Bacteriology	Histopathology	Virology Results[¥] [ ] – qPCR $_{RSIV/ISKNV}$ copy number/sample reaction range
18/10/2018 †Farm 1	n=5; 100-300g	Wide scale ongoing mortalities in on growing fish in cages.	*Aeromonas veronii*; *Aeromonas jandaei*; *Plesiomonas shigelloides*; *Chryseobacterium sp*; *Acinetobacter johnsonii*;	Not done	*qPCR $_{TiLV}$ 5/5 -ve *cPCRa $_{ISKNV}$ 4/5 +ve *qPCR $_{RSIV/ISKNV}$ 5/5 +ve [$7.5 \times 10^1 – 4.60 \times 10^5$]
28/11/2018 †Farm 2	n=14; 74-401g	Wide scale ongoing mortalities in on growing fish in cages (**Figure 2**). On shore fry production unit: no unusual mortalities.	*Aeromonas veronii*; *Aeromonas jandaei*; *Streptococcus agalactiae* capsular type Ib biotype 2 (non-haemolytic); *Edwardsiella tarda*	Evidence of Gram negative and Gram positive bacterial and viral# infection (including in same fish)	qPCR $_{TiLV}$ 14/14 -ve qPCR $_{NODA}$ 14/14 -ve *cPCRa $_{ISKNV}$ 12/14 +ve *qPCR $_{RSIV/ISKNV}$ 14/14 +ve [$2.48 \times 10^1 - 3.3 \times 10^6$]
17/12/2018 †Farm 2	n=7; 40-646g	Much lower mortalities in on growing fish in cages than previous sampling visit. On shore fry production unit: no unusual mortalities.	Not done	Evidence of bacterial infection but not of viral infection	qPCR $_{TiLV}$ 7/7 -ve qPCR $_{NODA}$ 7/7 -ve **cPCRb $_{RSIV/ISKNV}$ 1/7 +ve megalocytivirus, sequence = ISKNV** *qPCR $_{RSIV/ISKNV}$ 6/7 +ve [$1 \times 10^0 - 24 \times 10^0$]
20/02/2019 †Farm 2	n=14; 6-9 cms	Limited mortalities in on growing fish in cages. Very severe mortalities in on shore fry production unit.	Not done	Very severe virus-associated pathology (in tissue samples from all fish).	qPCR $_{TiLV}$ not done qPCR $_{NODA}$ not done cPCRa $_{ISKNV}$ 7/7 +ve qPCR $_{RSIV/ISKNV}$ 13/13 +ve [$5.1 \times 10^5 – 1.51 \times 10^7$]
09/07/2019 Farm 3	Grp 1, n=3; 60-80g Grp 2, n=3; 20-40g Grp 3, n=5; 1-2g fry Grp 4, n=5 pools of ≤0.2g fry	No mortality events reported at site. Grp 1 on-growing in lake, Grps 2-4 in ponds.	Grps 1 & 2 - No significant colonies Grps 3-5 Not done	Evidence of metacercaria, bacterial gill epitheliocystis and myxospordiosis in Grps 1 & 2. - No evidence of viral infection.	cPCRb $_{RSIV/ISKNV}$ Grp 1 - 2/3 +ve (nested only) Grp 2 - 3/3 +ve (nested only) Grp 3 – 5/10 +ve (nested only) Grp 4 - 3/10 +ve (nested only)

09/07/2019 Farm 4	n=5 pools of ≤0.2g fry	Reported recent mortality event now recovering. No clinical signs	Not done	Myxosporidian cysts in cranial sub-epithelial and connective tissue	cPCRb $_{RSIV/ISKNV}$ 2/5 +ve (nested only and in one of duplicates only)
10/07/2019 Farm 5	n=5 x3 fish organ pools; ~40g	Ongrowing fish on lake, recent mortality event in adjacent cage cleared previous month, no clinical signs in sampled fish	No significant colonies	not done	cPCRb $_{RSIV/ISKNV}$ 5/5 +ve (nested only, one of which weak in 1st round)
10/07/2019 Farm 6	n=10 x3 fish pools; 0.2-0.75g	Ongoing losses of approx. 30% in fry pools. Fry observed with distended abdomen and disturbed swimming.	Not done	Evidence of viral infection, occasionally severe	cPCRb $_{RSIV/ISKNV}$ 10/10 +ve Virus isolation – BF cells
10/07/2019 Farm 7	n=5 x2 fish organ pools; 20-40g	Internal haemorrhaging, ascites, pale kidney, anaemia, reduced spleen and enlarged liver	No significant colonies	Evidence of viral infection	cPCRb $_{RSIV/ISKNV}$ 5/5 +ve Virus isolation – BF cells

Table 1 Summary of sampling and results
† See Materials and Methods for farm descriptions
‡ All moribund fish, including fry, presented similar signs: erratic swimming, lethargy, ascites, swollen, dark spleen, haemorrhagic livers and other organs.
# Haematopoietic cell nuclear and cytoplasmic pleomorphism with marginalisation of chromatin and fine granulation.
¥ qPCR $_{TiLV}$, qPCR $_{NODA}$, qPCR $_{RSIV/ISKNV}$: results of qPCR testing for tilapia lake virus, nodavirus and ISKNV respectively. cPCRa & cPCRb : conventional PCR using
OIE recommended protocols by Kurita et al., 1998 (a) (J Kurita et al., 1998) and Rimmer et al., 2012 (b) (Rimmer, A. E. et al., 2012) protocols, results in bold represent
index finding, *represents retrospective testing post index finding.

Figure 6 N-J tree showing the genetic relationship of partial MCP sequence from LV#1 to a range of RSIV, ISKNV and related Megalocytiviruses. The GeneBank accession numbers and the host species are included where available.

3.8 Virus isolation

Fish material from Farms 6 and 7 was inoculated onto BF-2, E-11 and GF cell lines.

Cytopathic effects (CPE) were observed in BF-2 and E-11 cells but not GFs on first

inoculation (Figure 7). On passage (P1) in the same cell types CPE was only observed for

BF-2 cells and intensity of CPE was diminished. Isolated virus from clarified harvested cell

culture supernatant from the P1 BF-2 cells was confirmed positive for ISKNV by PCR with

sequence identical to that obtained by PCR direct from tissue homogenate (data not shown).

472

Figure 7 Virus isolation and cytopathic effects in BF-2 cells. (A) BF-2 cells at 5 days post

inoculation (-3 dilution) showing enlarged, rounded, phase bright cells. Inset, advanced

extensive CPE from -2 dilution of BF-2 at same time point. (B) Negative control BF-2 cells

at 5 days.

4 Discussion

The results suggest ISKNV had a significant role in the high mortalities experienced by the

two farms that were investigated. Fish sampled from the second farm at the height of the

disease outbreak showed severe clinical and pathological signs typically associated with

infection by the virus (including visualisation of distinctive megalocytes with characteristic

virions). Both, these fish and those sampled earlier from Farm 2, had extremely high copy

numbers of RSIV/ISKNV *Megalocytivirus*-like virions and ISKNV was confirmed by gene

sequencing. Further investigations later in 2019 of five further farms from different areas of

lake Volta showed by then the disease was widely established (endemic) in farms across the

lake, with both symptomatic and asymptomatic fish positive by PCR for the virus. Farmers

emphasised how the disease had a devastating effect on the industry during discussions on

this latter visit. They reported how the disease continued to have an impact on the broodstock

and growout fish at a lower rate, however the mass mortalities were now predominantly

observed in juvenile fish (1-5g in weight). Survival rates to the growout stage, at that time,

were estimated to be as low as 5-20%. Juvenile mortalities were reported to be episodic in

nature, occurring a few days after the sex reversal process or translocation to lake cages, and

lasting up to 3 to 4 weeks. These stress triggered mortality outbreaks may be indicative of

either widespread persistence of virus in the environment or a latent ISKNV stage.

To improve survival rates, some farmers had trialled reductions in juvenile stocking densities.

Large reductions in stocking density were only associated with small increases in survival,

therefore this practice was not considered a viable solution. Instead, farmers resorted to a

substantial increase in the level of juvenile production and stocking rates, trading off higher

overall mortality with some guarantee of a small but not inconsequential harvest.   The

economic impact of ISKNV has been significant. The higher production costs and reduced

harvests resulted in most farms having to either temporarily or permanently halt production.

As larger farms can be the primary employers of some villages, the disease also had a direct

impact on the livelihoods of local community members. It was also reported that tilapia

market price had more than doubled due to the production shortages and that the feed sales of

Raanan Fish Feeds had reduced by 70% (https://thefishsite.com/articles/ghanaian-fish-

farmers-seek-help-from-big-business), both potential indicators of the virus having an impact

on a much larger scale.

The results would be consistent with a recent introduction of the virus onto one or more farms

prior to October 2018, that then extended upstream of the dam to other farms resulting in the

unusual, widespread and significant mortalities observed. Firstly, there were no observations

of typical ISKNV-associated pathology in any previous disease investigations, on the affected

or other farms. Secondly, the limited PCR screening in this study of archived samples of

diseased tilapia recovered from before the major mortality event including both farms, were

all negative. Moreover, (Verner-Jeffreys et al., 2018) also screened for the presence of

RSIV/ISKNV by PCR in 2016, including samples from the affected farms, without detecting

the virus, or associated pathology.

Iridioviruses are large icosahedral cytoplasmic double□stranded DNA viruses, which can

infect a wide range of hosts, including invertebrates and poikilothermic vertebrates. The

family *Iridoviridae* includes five genera: *Iridovirus, Chloriridovirus, Ranavirus,*

Lymphocystivirus and *Megalocytivirus* (Jancovich et al., 2012). Fish pathogenic iridoviruses

are representatives of *Ranavirus, Lymphocystivirus* and *Megalocytivirus* genera

(Subramaniam, Shariff, Omar, & Hair-Bejo, 2012; Whittington, Becker, & Dennis, 2010).

Infectious spleen and kidney necrosis virus (ISKNV), is a member of the genus

Megalocytivirus (Jun Kurita & Nakajima, 2012), and causes disease in a range of freshwater

and marine fish species (Subramaniam, Shariff, Omar, & Hair-Bejo, 2012; Whittington,

Becker, & Dennis, 2010). ISKNV is closely related to red sea bream iridovirus and both

viruses are listed by the OIE as notifiable pathogens (World Organisation for Animal Health

OIE, 2018).   Although tilapia is not listed as a susceptible species by the OIE at present

(World Organisation for Animal Health OIE, 2018),   recent   reports   from   the   USA

(Subramaniam et al., 2016) and Thailand (Suebsing et al., 2016) suggest that it is a

susceptible species likely to suffer significant mortality. The change in the known host range

of virus needs to be communicated to the international community to prevent future

transboundary spread through movement of infected tilapia.

It was interesting to note that many of the ISKNV positive fish were actively co-infected with

Streptoccocus agalactiae and other bacterial pathogens, presenting severe bacteraemia /

meningitis, as well as ISKNV-associated pathology. The high mortalities on the farms in the

larger on-growing fish may well have been exacerbated by these coinfections.

Although the mortalities were initially confined to on-growing fish in cage culture systems,

the later observations of very high ISKNV associated mortalities in fry, associated clinical

signs and high viral copy numbers shows the virus likely affects all life stages. As fry are

often reared in onshore facilities below the dam and then translocated to on-growing cages on

the main lake, this may have been one of the routes that disease was rapidly spread after it

first emerged. Anecdotally, at the time of writing, farmers on the sites visited report that

mortalities in on-growing facilities have declined, while fry production continues to be badly

affected. It is possible that surviving fry have been exposed to the virus and then protected

against subsequent exposure. This suggests that immunisation of fry, or use of previously

exposed individuals, could represent a practical disease management strategy. Vaccination as

a control strategy may be used to control red sea bream iridiodovirus (Nakajima et al., 1999;

Shimmoto, Kawai, Ikawa, & Oshima, 2010) and there are also encouraging reports of its

potential effectiveness for protection against ISKNV in mandarin fish (Dong, Weng, He, &

Dong, 2013). As some of these reports showed efficacy using formalin killed virus infected

cells (Dong et al., 2013), rapid development and testing of vaccines based on the direct use

of the strain(s) of ISKNV circulating in Lake Volta farms (e.g. autogenous vaccines) should

be possible.

Outbreaks of disease that cause significant morbidity and/or mortalities in an aquaculture

operation is always a major concern. This is exacerbated when this appears to represent the

incursion of a new agent into a system, or region (country or zone in a country) which has not

previously been affected. A stark example of this is the epidemic of infectious salmon

anaemia virus (ISAV) which reduced production by three quarters and resulted in severe

economic and social crisis in the developing Chilean Atlantic salmon industry between 2007

and 2010 (Godoy et al., 2008; Mardones, Martinez-Lopez, Valdes-Donoso, Carpenter, &

Perez, 2014; Vike, Stian Nylund, & Nylund, 2009).

The Ghanaian authorities have for some time been concerned that the, to date, successful,

expansion of its industry on Lake Volta may be affected by such disease incursions. Partly for

this reason, and to safeguard the genetic integrity of Lake Volta Nile tilapia strains, they have

tried to limit the culture to locally reared Nile tilapia stocks. However genetic testing by the

Ghanaian Fisheries Commission (Ziddah *et al.*, Unpublished Observations)   showed that

fish on some of the farms on Lake Volta were likely of imported GIFT strain origin (Ponzoni

et al., 2011) or hybrids of GIFT and indigenous strains.

If farmers have been illegally sourcing broodstock from Asia and other areas, that would be

an ideal method of translocating pathogens from one region to another. It should be noted

though that ISKNV has also been detected in internationally traded freshwater ornamental

species, theoretically posing another possible introduction route (Jung-Schroers et al., 2016)

It is very possible that this is not the first time disease introduction has taken place in Lake

Volta Ghana. The study by (Verner-Jeffreys et al., 2018) showed that outbreaks of *S* .

agalactiae investigated in 2016 were all caused by genetically indistinguishable isolates of

ST 261, with closest genetic identity to Asian isolates. Discussions with affected farmers at

the time suggested, that *S. agalactiae* associated mortalities were a relatively recent

phenomenon, although the disease is now clearly endemic to all the areas in the Volta area.

Other studies have shown that *S. agalactiae* ST261 has likely been translocated around the

world in association with farmed tilapia (Kawasaki et al., 2018). As a single large

epidemiological unit, it will be difficult to control transmission of virus between farms on

lake Volta. It is important to try to prevent spread from Lake Volta to surrounding

watersheds and the wider African continent by control over movement of live fish and

equipment. The development of biosecure offline hatcheries with borehole water or UV

treatment of water, to facilitate production of juveniles which survive to a size they can be

vaccinated will likely be key to vaccine control. Additionally, recent technological advances

in rapid selective breeding (Houston et al., 2020; Robledo, Palaiokostas, Bargelloni,

Martínez, & Houston, 2018) should be employed to develop ISKNV disease resistant

populations, or strains of tilapia to enable the industry to recover.

Although most attention to date has focussed on the emergence and spread of TiLV within

the tilapia industry world-wide, these results also demonstrate that there is a range of other

potential threats to the sustainability of tilapia aquaculture.

Conclusion

This is the first report of Infectious Spleen and Kidney Necrosis Virus (ISKNV) in farmed

tilapia in Africa. ISKNV was found in co-infection with *Streptococcus agalactiae* and other

bacterial pathogens in Lake Volta, Ghana. The correlations seen between the mortality

events, histopathology and viral loads in the tissues suggest that ISKNV was a major cause of

mortalities during the outbreaks. In general, the results support continued efforts to improve

the biosecurity of the industry in Ghana. There is a clear need to strengthen domestic

capability to rapidly diagnose and control emerging disease threats caused by ISKNV and

other pathogens. Further work is also needed to map the distribution of the virus and its

impact, including potential effects on wild fish species, and to implement practical control

strategies.

Acknowledgements

The farmers involved in the investigation are thanked for their support and provision of

information. Support from Defra (contracts FB002 and FX001 for the OIE Collaborating

Centre for Emerging Aquatic Animal Diseases) is gratefully acknowledged.

References

Data sharing

The data that support the findings of this study are openly available in BioRxiv doi:

https://doi.org/10.1101/680538

Abdelsalam, M., Asheg, A., & Eissa, A. E. (2013). Streptococcus dysgalactiae: An emerging

pathogen of fishes and mammals. *International Journal of Veterinary Science and*

Medicine, *1*(1), 1–6. https://doi.org/10.1016/j.ijvsm.2013.04.002

Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman,

D. J. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database

search programs. *Nucleic Acids Research*, *25*, 3389–3402.

https://doi.org/10.1093/nar/25.17.3389

Delannoy, C. M., Crumlish, M., Fontaine, M. C., Pollock, J., Foster, G., Dagleish, M. P., …

Zadoks, R. N. (2013). Human Streptococcus agalactiae strains in aquatic mammals and

fish. *BMC Microbiology*, *13*(1), 41. https://doi.org/10.1186/1471-2180-13-41

Dong, Y., Weng, S., He, J., & Dong, C. (2013). Field trial tests of FKC vaccines against

RSIV genotype Megalocytivirus in cage-cultured mandarin fish (Siniperca chuatsi) in an

inland reservoir. *Fish and Shellfish Immunology*, *35*, 1598–1603.

https://doi.org/10.1016/j.fsi.2013.09.005

FAO. (2016). *The State of World Fisheries and Aquaculture. Contributing to food security*

and nutrition for all. Rome.

FAO. (2018). *The State of World Fisheries and Aquaculture 2018 - Meeting the sustainable*

development goals. https://doi.org/978-92-5-130562-1

Godoy, M. G., Aedo, A., Kibenge, M. J. T., Groman, D. B., Yason, C. V., Grothusen, H., …

Kibenge, F. S. B. (2008). First detection, isolation and molecular characterization of

infectious salmon anaemia virus associated with clinical disease in farmed Atlantic

salmon (Salmo salar) in Chile. *BMC Veterinary Research*, *4:28*.

https://doi.org/10.1186/1746-6148-4-28

Hounmanou, Y. M. G., Mdegela, R. H., Dougnon, T. V., Achoh, M. E., Mhongole, O. J.,

Agadjihouèdé, H., … Dalsgaard, A. (2018). Tilapia lake virus threatens tilapiines

farming and food security: Socio-economic challenges and preventive measures in Sub-

Saharan Africa. *Aquaculture*, *493*, 123–129.

https://doi.org/10.1016/j.aquaculture.2018.05.001

Houston, R. D., Jin, Y. H., Jenkins, T. L., Selly, S. L. C., Martin, S. A. M., Stevens, J. R., &

Santos, E. M. (2020). Harnessing genomics to fast-track genetic improvement in

aquaculture. *Nature Reviews Genetics*. https://doi.org/10.1038/s41576-020-0227-y

Jancovich, J. K., Chinchar, V. G., Hyatt, A., Miyazaki, T., Williams, T., & Zhang, Q. Y.

(2012). Family Iridoviridae. In A. M. Q. King, M. J. Adams, E. B. Carstens, &

Lefkowitz. E.J. (Eds.), *Virus Taxonomy: Ninth Report of the International Committee on*

Taxonomy of Viruses. (pp. 193–210). San Diego, CA: Elsevier Academic Press.

Jansen, M. D., Cudjoe, K., & Brun, E. (2018). *Investigation of tilapia mortality in Ghana*.

Jung-Schroers, V., Adamek, M., Wohlsein, P., Wolter, J., Wedekind, H., & Steinhagen, D.

(2016). First outbreak of an infection with infectious spleen and kidney necrosis virus

(ISKNV) in ornamental fish in Germany. *Diseases of Aquatic Organisms*, *119*, 239–244.

https://doi.org/10.3354/dao02995

Kawasaki, M., Delamare-Deboutteville, J., Bowater, R. O., Walker, M. J., Beatson, S., Ben

Zakour, N. L., & Barnes, A. C. (2018). Microevolution of Streptococcus agalactiae ST-

261 from Australia Indicates Dissemination via Imported Tilapia and Ongoing

Adaptation to Marine Hosts or Environment. *Applied and Environmental Microbiology*,

84, e00859-1. https://doi.org/10.1128/aem.00859-18

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F. O.

(2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and

next-generation sequencing-based diversity studies. *Nucleic Acids Research*, *41*(1), e1.

https://doi.org/10.1093/nar/gks808

Kurita, J, Nakajima, K., Hirono, I., & Aoki, T. (1998). Polymerase chain reaction (PCR)

amplification of DNA of red sea bream iridovirus (RSIV). *Fish Pathology*, *33*(1), 17–23.

Retrieved from http://www.scopus.com/inward/record.url?eid=2-s2.0-

0032388179&partnerID=40&md5=3128c9b74c19de5bce885a2566e475a0%5Cnhttps://

www.jstage.jst.go.jp/article/jsfp1966/33/1/33_1_17/_pdf

Kurita, Jun, & Nakajima, K. (2012). Megalocytiviruses. *Viruses*, *4*, 521–538.

https://doi.org/10.3390/v4040521

Luo, X., Fu, X., Liao, G., Chang, O., Huang, Z., & Li, N. (2017). Isolation, pathogenicity and

characterization of a novel bacterial pathogen Streptococcus uberis from diseased

mandarin fish Siniperca chuatsi. *Microbial Pathogenesis*, *107*, 380–389.

https://doi.org/https://doi.org/10.1016/j.micpath.2017.03.049

Machimbirike, V. I., Jansen, M. D., Senapin, S., Khunrae, P., Rattanarojpong, T., & Dong, H.

T. (2019). Viral infections in tilapines: More than just tilapia lake virus. *Aquaculture*,

503, 508–518. https://doi.org/10.1016/j.aquaculture.2019.01.036

Mardones, F. O., Martinez-Lopez, B., Valdes-Donoso, P., Carpenter, T. E., & Perez, A. M.

(2014). The role of fish movements and the spread of infectious salmon anemia virus

(ISAV) in Chile, 2007-2009. *Preventive Veterinary Medicine*, *114*(1), 37–46.

https://doi.org/10.1016/j.prevetmed.2014.01.012

Nakajima, K., Maeno, Y., Honda, A., Yokoyama, K., Tooriyama, T., & Manabe, S. (1999).

Effectiveness of a vaccine against red sea bream iridoviral disease in a field trial test.

Diseases of Aquatic Organisms, *36*, 73–75. https://doi.org/10.3354/dao036073

Plumb, J. A., & Hanson, L. A. (2011). Tilpia bacterial diseases. In J. A. Plum & L. A.

Hanson (Eds.), *Health Maintenance and Principle Microbial Diseases of Cultured*

Fishes, Third edition. (pp. 445–463).

https://doi.org/https://doi.org/10.1002/9780470958353.ch16

Ponzoni, R. W., Nguyen, N. H., Khaw, H. L., Hamzah, A., Bakar, K. R. A., & Yee, H. Y.

(2011). Genetic improvement of Nile tilapia (Oreochromis niloticus) with special

reference to the work conducted by the WorldFish Center with the GIFT strain. *Reviews*

in Aquaculture, *3*(1), 27–41. https://doi.org/doi:10.1111/j.1753-5131.2010.01041.x

Rimmer, A. E., Becker, J. A., Tweedie, A., & Whittington, R. J. (2012). 2012 Development

of a quantitative polymerase chain reaction (qPCR) for the detection of dwarf gourami

iridovirus (DGIV) and other Megalocytiviruses and comparison with the Office

International des Epizooties reference PCR protocol. Aquaculture 358-359:15.

Aquaculture, *358–359*, 155–163.

Robledo, D., Palaiokostas, C., Bargelloni, L., Martínez, P., & Houston, R. (2018).

Applications of genotyping by sequencing in aquaculture breeding and genetics.

Reviews in Aquaculture, *10*(3), 670–682. https://doi.org/10.1111/raq.12193

Shimmoto, H., Kawai, K., Ikawa, T., & Oshima, S. I. (2010). Protection of red sea bream

Pagrus major against red sea bream iridovirus infection by vaccination with a

recombinant viral protein. *Microbiology and Immunology*, *54*, 135–142.

https://doi.org/10.1111/j.1348-0421.2010.00204.x

Shoemaker, C. A., Xu, D. H., Garcia, J. C., & Lafrentz, B. R. (2017). Capsular typing of

streptococcus agalactiae (Lancefield group B streptococci) from fish using multiplex

PCR and serotyping. *Bulletin of the European Association of Fish Pathologists*, *37*(5),

190–197.

Subramaniam, K., Gotesman, M., Smith, C. E., Steckler, N. K., Kelley, K. L., Groff, J. M., &

Waltzek, T. B. (2016). Megalocytivirus infection in cultured Nile tilapia Oreochromis

niloticus. *Diseases of Aquatic Organisms*, *119*(3), 253–258.

https://doi.org/10.3354/dao02985

Subramaniam, K., Shariff, M., Omar, A. R., & Hair-Bejo, M. (2012). Megalocytivirus

infection in fish. *Reviews in Aquaculture*, *4*(4), 221–233. https://doi.org/10.1111/j.1753-

5131.2012.01075.x

Suebsing, R., Pradeep, P. J., Jitrakorn, S., Sirithammajak, S., Kampeera, J., Turner, W. A., …

Kiatpathomchai, W. (2016). Detection of natural infection of infectious spleen and
kidney necrosis virus in   farmed tilapia by hydroxynapthol blue-loop-mediated
isothermal amplification assay. *Journal of Applied Microbiology*, *121*(1), 55–67.
https://doi.org/10.1111/jam.13165

Tamura, K., Stecher, G., Peterson, D., Filipski, A., & Kumar, S. (2013). MEGA6: Molecular
Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution*, *30*(12),
2725–2729. https://doi.org/10.1093/molbev/mst197

Verner-Jeffreys, D., Wallis, T., Cano, I., Ryder, D., Hayden, D., Domozoro, J., … Feis, S.
(2018). Streptococcus agalactiae Multi Locus Sequence Type 261 is associated with
mortalities in the emerging Ghanaian tilapia industry. *Journal of Fish Diseases*, *41*,
157–179. https://doi.org/10.1111/jfd.12681

Vike, S., Stian Nylund, S., & Nylund, A. (2009). ISA virus in Chile: evidence of vertical
transmission. *Archives of Virology*, *154*, 1–8.

Whittington, R. J., Becker, J. A., & Dennis, M. M. (2010). Iridovirus infections in finfish-
critical review with emphasis on ranaviruses. *Journal of Fish Diseases*, *33*(2), 95–122.
https://doi.org/10.1111/j.1365-2761.2009.01110.x

World Organisation for Animal Health OIE. (2018). Chapter 2.3.8 Red sea bream iridoviral
disease. In *Manual of Diagnostic Tests for Aquatic Animals*. Retrieved from
http://www.oie.int/en/standard-setting/aquatic-manual/access-online/

Yáñez, M. A., Catalán, V., Apráiz, D., Figueras, M. J., & Martínez-Murcia, A. J. (2013).
Phylogenetic analysis of members of the genus Aeromonas based on gyrB gene
sequences. *International Journal of Systematic and Evolutionary Microbiology*, *53*,
875–883.