First detection of Infectious Spleen and kidney Necrosis Virus (ISKNV)

associated with massive mortalities in farmed tilapia in Africa

Short running tittle: First detection of ISKNV in farmed tilapia in Africa

José Gustavo Ramírez-Paredes[a], Richard K. Paley[bc], William Hunt[a], Stephen W. Feist[bc]

David M. Stone[bc], Terry Field[a], David J. Haydon[a], Peter A. Ziddah[d] , Samuel Duodu[e],

Timothy S. Wallis[a], David W. Verner -Jeffreys[bc]*

[a] Ridgeway Biologicals Limited, Units 1-3 Old Station Business Park, Compton, Berkshire,

England, United Kingdom

[b] Cefas Weymouth Laboratory, Weymouth, Dorset, England, United Kingdom.

[c] OIE Collaborating Centre for Emerging Aquatic Animal Diseases.

[d] Fisheries Commission, Ministry of Fisheries and Aquaculture Development, Accra, Ghana

[e] University of Ghana, Accra, Ghana

*Corresponding author Dr D. W. Verner-Jeffreys Cefas Weymouth Laboratory, Weymouth,

Dorset, England, United Kingdom

Abstract

In late 2018, unusual patterns of very high mortality (>50% production) were reported in intensive tilapia cage culture systems across Lake Volta in Ghana. Affected fish showed darkening, erratic swimming and abdominal distension with associated ascites. Histopathological observations of tissues taken from moribund fish at different farms revealed the presence of lesions indicative of viral infection. These included haematopoietic cell nuclear and cytoplasmic pleomorphism with marginalisation of chromatin and fine granulation. Transmission electron microscopy showed tilapia cells contained conspicuous virions with typical Iridovirus morphology i.e. enveloped, with icosahedral and or polyhedral geometries and with a diameter c.160 nm. PCR confirmation and DNA sequencing identified the virions in the tissues of the diseased fish as Infectious Spleen and Kidney Necrosis Virus (ISKNV). Samples of fry and older animals were all strongly positive for the presence of the virus by qPCR. All samples tested negative for TiLV and Nodavirus by qPCR. All samples collected from farms in the year prior to the mortality event were negative for ISKNV. The results suggest that ISKNV was the cause of disease on the investigated farms and likely had a primary role in the mortality events. A common observation of coinfections with *Streptococcus agalactiae* and other tilapia bacterial pathogens further suggests that these may interact to cause severe pathology, particularly in larger fish. The present results demonstrate that there are a range of potential threats to the sustainability of tilapia aquaculture that need to be guarded against.

Keywords: Tilapia; Africa; Tilapia Viral Diseases; fish virus; Infectious Spleen and Kidney Necrosis Virus; ISKNV;

Introduction

The farming of tilapia species (*Oreochromis* spp.) is one of the most important sectors in aquaculture worldwide with total global production estimated at more than 6 686 000 tonnes in 2016 (1). In Africa, production is still dominated by Egypt with more than 1 000 000 tonnes produced in 2017 (1). However, tilapia culture has become increasingly important in several other African countries, where it boosts the local economy and constitutes an affordable source of animal protein for human consumption.

In Ghana, Nile tilapia production in 2016 had reached more than 50 000 tonnes, from only 2 000 tonnes per year in 2006 (2) with more than 90% of the production derived from high stocking density floating cage systems in Lake Volta. However, as production systems have intensified in the area, the industry has been increasingly affected by a range of disease issues (3, 4).

In 2017 (3) conducted the first comprehensive disease investigation in tilapia farmed in Lake Volta Ghana. *Streptococcus agalactiae* multilocus sequence type 261 was shown to be a major cause of mortality for farmed Nile tilapia and a range of other bacterial and parasitic pathogens including *Aeromonas* spp., *Streptococcus iniae*, *Flavobacterium columnare* and a *Myxobolus* sp. were also detected.

Jansen et al. (4) conducted follow up studies, including a broad-ranging epidemiological investigation and suggested that mortalities caused by bacteria in Lake Volta were not a major concern for the local economy as farmers had managed to sustain losses by increasing production of fingerlings, treatment with antibiotics and use of autogenous vaccines.

Historically, bacterial infections were the major threat for the health of farmed tilapia (5). However, in recent years a number of viral diseases have emerged worldwide with devastating effects for the industry(6). Tilapia Lake Virus (TiLV) is considered the main viral challenge to

tackle, as it has spread to many producing countries, causing high mortalities in all production

stages. Although TiLV has been detected elsewhere in Africa (7), the initial study by (3),

diagnostic investigations undertaken by Ridgeway Biologicals Ltd. (Ramirez *et al*. unpublished

data), or the more recent survey undertaken by (4), all failed to provide any evidence of TiLV in

diseased tilapia reared in Lake Volta. Up until September 2018, with the exception of the

detection of a nodavirus sequence by (3), no significant association between viral agents and

tilapia mortality events have been demonstrated in Ghana.

From September 2018 to March 2019 outbreaks of disease with very high levels of morbidity and

mortality (60-90%), were experienced in both vaccinated and unvaccinated tilapia by farmers in

Lake Volta. In late-September 2018, a farm located below the lower dam in the region of

Asutsuare, was the first to suffer episodes of massive acute mortalities (https://goo.gl/LmqbG2

and https://bit.ly/2NwDEbD). Approximately a week after the first report, a second farm located

in the Akuse region (~5 km upstream of Asutsuare, but still below the lower dam) also

experienced acute mortalities. By mid-October, multiple floating cage-based farms in the Dodi

region (above the upper dam) reported losses of more than 10 tonnes per day

https://goo.gl/yj4oT4. In late November, farmers that had been unaffected in the Asikuma region

(downstream of Dodi but still above the upper dam) also started to suffer episodes of massive

acute mortality. (**Figure 1**).

By the end of 2018, most tilapia farmers in Lake Volta had reported mortalities that they were not

able to contain by increased production of fingerlings or treating with antibiotics (**Supplementary**

File 1, https://bit.ly/2KZFFuw and https://goo.gl/cPmpSE). We report the results of a

comprehensive disease investigation, conducted at two affected farms from two different regions

of Lake Volta, to gain insights into the causes of these mortalities.

84
85 **Figure 1 Map of lower region of Lake Volta in Ghana, West Africa.** Red triangles indicate the regions in chronological order (A to D) where
86 the outbreaks of mortality occurred.

2 Materials and Methods

2.1 Sampling

Farm 1, a medium size (approximately 600-800 tonnes per annum production) cage culture operation on upper Lake Volta, was visited on the 18/10/18. Samples from 10 fish (average weight 200g) were taken for bacteriology. In addition five sets of samples containing liver and brain from fish from different cages were collected for molecular diagnostics as detailed in **Table 1**.

Farm 2, a large cage culture operation (>2000 tonnes per annum), was visited for sampling on 28/11/18, 17/12/18 and 20/02/19. For the first two visits, moribund fish between 40-646g collected from cages on the main lake were sampled. In the first visit for bacteriology and virology on the second visit for virology only. For the third visit, responding to reports that they were now experiencing very heavy mortalities in their fry production units (>70%), samples of moribund fry and juveniles from both nursery cages on the main lake and from their onshore hatchery supplied with water pumped from the main lake, were analysed for virology. From this farm material was also taken for histological and molecular diagnostic investigations during the visits as detailed in **Table 1 and Supplementary File 2**.

All moribund fish from the visited farms were humanely euthanised with a lethal overdose of tricaine methanesulfonate 1,000 mg/g (Pharmaq, Hampshire, UK) followed by brain destruction prior to the necropsy.

2.2 Bacteriology

Samples for bacteriology were collected from the brain, liver, kidney and spleen with sterile cotton swabs and inoculated onto tryptone soya agar (TSA), Columbia blood agar (CBA),

Tryptone yeast extract salts agar (TYES) (Southern Group Laboratory, Corby, UK) and

cystine heart agar with 2% bovine haemoglobin (CHAH) (Becton Dickinson, Oxford, UK).

All inoculated agar plates were incubated at 28 °C for 24-72 hours. Colonies assessed as

significant based on occurrence and dominance were subcultured to purity on similar media.

Pure relevant isolates were initially identified by morphology and Gram staining. The partial

16S rRNA genes of the Gram negative isolates identified were PCR amplified and sequenced

using the method described by (8). Gram positive cocci forming chains were screened using a

Streptococcus agalactiae specific capsular typing multiplex PCR developed by (9). The

Gram negative isolates strains confirmed as *Aeromonas* spp. based on partial 16S rRNA gene

sequence analysis were further characterised based on partial *gyrB* sequencing analysis for

identification at the species level as described by (10).

2.3 Histopathology and Electron Microscopy

Tissues were fixed in neutral buffered formalin (NBF) for a minimum of 24 hr before being

placed in glycerol diluted 50:50 with phosphate buffered saline (PBS) for transportation to

Cefas. On receipt, tissues were rinsed in 70% alcohol and placed again in NBF for a final

period of fixation prior to processing using standard protocols in a vacuum infiltration

processor. Tissue sections were cut at a thickness of 3-4 µm on a Finnese® microtome,

mounted on glass slides and stained with haematoxylin and eosin using an automated staining

protocol. Stained sections were examined for general histopathology by light microscopy

(Nikon Eclipse E800). Digital images and measurements were obtained using the Lucia™

Screen Measurement software system (Nikon, UK).

For electron microscopy, small samples of tissues fixed in NBF as above were rinsed three

changes of 0.1 M sodium cacodylate buffer, followed by post fixation in 2.5% glutaraldehyde

in the same buffer for 1 hour prior to a second post fixation for 1 hour in 1 % osmium

tetroxide in 0.1 M sodium cacodylate buffer. Subsequently, fixed tissues were dehydrated in

an ascending acetone series acetone series and embedded in epoxy resin 812 (Agar Scientific

pre-Mix Kit 812, Agar scientific, UK) and polymerised at 60 °C overnight. Semi-thin (1 µm)

survey sections were stained with 1 % Toluidine Blue and examined by light microscope to

identify areas of interest. Ultrathin sections (70-90 nm) of the targeted areas were placed on

uncoated copper grids and stained with uranyl acetate and Reynold's lead citrate (Reynolds

1963). Grids were examined using a JEOL JEM 1400 transmission electron microscope and

digital images captured using a GATAN Erlangshen ES500W camera and Gatan Digital

Micrograph[TM] software.

2.4 Molecular diagnosis (viral and bacterial detection)

The samples collected for molecular diagnosis were washed twice in 750µl of sterile 1x PBS

to remove the RNA-*later®* and homogenised. Total nucleic acids were extracted using

nanomagnetic beads i.e. Genesig Easy DNA/RNA Extraction Kit (Primerdesign,

Southampton, UK) and stored until further use.

Multiplex PCR for detection of Streptococcus spp.

Nucleic acids extracted were used as a template on a multiplex PCR (unpublished data) to

confirm the presence of *Streptococcus* spp. that had been previously reported as fish

pathogens including   *Streptococcus agalactiae* (11), *Streptococcus uberis* (12) and

Streptococcus dysgalactiae (13).

qPCR for detection of TiLV and Nodavirus

Nucleic acids were used for the detection of tilapia lake virus and nodavirus by quantitative

PCR using the commercial kits: Path-TiLV-EASY and Path-Betanodavirus-EASY

(Primerdesign, Southampton, UK) in the platform Genesig q16® (Primerdesign,

Southampton, UK) as per the protocols suggested by the manufacturer.

Conventional PCR for detection of megalocytiviruses

The Office International des Epizooties (OIE) reference PCR protocol for notifiable aquatic

megalocytiviruses (Red Seabream Iridoviral disease / Infection spleen and kidney necrosis

virus (14, 15)) with the modification proposed by (16) was initially used, to screen the

samples of fish collected from visit 2 at Farm 2. For this, genomic DNA was extracted as

follows: the RNA-*later*® was removed and the tissue samples weighed. Depending on the

weight of the tissue available the samples were diluted in RTL buffer (Qiagen) to provide

either a 1:10 w/v or a 1:5 w/v and homogenised per fish i.e. all the organs of each fish into an

individual pool using Matrix A and the FastPrep apparatus (MP Biomedicals). Following

homogenisation, the samples were diluted further with RTL buffer to give a 1:60 w/v

homogenate and total nucleic acid was extracted from 300 µl of the clarified sample using the

RNA tissue mini kit without DNase (Qiagen) and eluted in a 60 µl volume.

RT was performed at 37°C for 1 h in a 20 µl volume consisting of 1× M-MLV RT reaction

buffer (50 mM) Tris pH 8.3, 75 mM KCl, 10 mM DTT, 3 mM MgCl2) containing 1 mM

dNTP, 100 pmol random primers, 20 U M-MLV reverse transcriptase (Promega,

Southampton UK) and 4µl of the nucleic acid extracted above.

PCR was performed in duplicate in a 50 µl reaction volume with 2.5 µl of cDNA of total

nucleic acid, 25 mM dNTPs, 1 x GoTaq® buffer (2.5 mM MgCl2 solution), 5 pmol of each

primer   (C1105   5$^\prime$-GGTTCATCGACATCTCCGCG-3'   and   C1106   5$^\prime$-

AGGTCGCTGCGCATGCCAATC-3') and 1.25 units of GoTaq® DNA polymerase

(Promega). The cycling conditions were as follows: 40 temperature cycles (1 min at 95°C, 1

min at 55°C and 1 min at 72°C) after an initial denaturing step (5 min at 95°C) followed by a

final extension step of 10 min at 72°C.

Amplified products were electrophoresed in a 2% (w/v) agarose/TAE (40 mM Tris-acetate,

pH 7.2, 1 mM EDTA) gel containing 1.0 μg ml-1 ethidium bromide at 120 v for 30 mins and

viewed under UV light.

PCR products were excised from the gel and the DNA was extracted and purified by ethanol

precipitation. Both strands of the PCR product were sequenced using the ABI PRISM Big

Dye Terminator v3.1 cycle sequencing kit and the same primers used for the amplification.

The forward and reverse sequences were aligned and a consensus sequence generated using

the CLC software (Qiagen). Generated consensus sequences were compared with sequences

from GenBank using BLASTN (17) and aligned using the MUSCLE application of the

MEGA software version 6 (18).

In addition, the OIE recommended PCR protocol for notifiable aquatic megalocytiviruses

(14) developed by (15) was used was used to screen total nucleic acids extracted from the rest

of the samples fixed for viral molecular analyses.

qPCR for detection and quantification of megalocytiviruses

The amount of virus present in the samples was also investigated by qPCR. For this the

homogenised tissues were subjected to total nucleic acids extraction (~20mg of each organ)

using the Genesig Easy DNA/RNA Extraction Kit (Primerdesign) as described earlier. The

extracted nucleic acids were tested using the commercial kit Path-ISKNV-EASY

(Primerdesign, Southampton, UK) in the platform Genesig q16® (Primerdesign) as per the

manufacturer instructions. This detects both red sea bream iridovirus and ISKNV variants. In

all cases fish were individually analysed either by pooling liver, spleen and brain or screening

individual tissues.

Retrospective analyses of archived samples by qPCR

A total of 16 samples of archived tissue homogenates from 5 different farms (that included

Farms 1 and 2) were retrospectively screened for ISKNV by qPCR with the commercial kit

Path-ISKNV-EASY as described before. From these 7 had been collected during 2017 and

the rest in March 2018 (**Supplementary File 5**). All the samples had been previously

confirmed as negative for TiLV and Nodavirus using the commercial kits Path-TiLV-EASY

and Path-Betanodavirus-EASY.

3 Results

3.1 Farm visits

On the first visit to Farm 2 on the 27/11/18, farm staff reported very high and ongoing

mortalities (Figure 2) in fish bigger than 20g, including broodstock, but no significant losses

in fingerlings. Losses reportedly peaked at about 670 crates (equating to approximately 40

tonnes per day) shortly after this visit on the 2/12/2018 **(Figure 2)**.

Losses were so severe that accurate estimation was not possible, with more than 50 additional

labourers recruited locally just to remove dead and moribund fish during the peak period. By

the second visit to Farm 2, losses of ongrowing fish had reportedly declined back to the

background 10-20 crates per day more typically observed i.e. less than 1-2 tonnes per day

(**Figure 2**).

221
Figure 2. Number of crates of dead/ moribund fish removed from Farm 2 during period
of maximum losses. Daily mortalities were estimated based on numbers of crates of rejected
ongrowing tilapia collected each day by farm staff. Each crate typically contained
approximately 60kg tilapia collected from the cages and rejected because they were either
dead, moribund or displayed other adverse signs that prevented their sale.

During the first two visits to the farms, but particularly to Farm 2, diseased fish were

observed swimming away from the school with erratic swimming i.e. on one side, in circles,

lethargic, with no equilibrium, upside down etc. **(Supplementary File 3).**

Externally, the fish displayed a range of clinical signs, including skin nodules, frayed fins,

loss of eyes, opaque eyes, loss of scales, exophthalmia, anorexia, decolouration or darkened

skin, excess of mucous, skin haemorrhages and distended abdomen (**Figure 3**). At necropsy,

fish from both the first visits to Farm 1 and 2 presented with marked ascites, enlarged and

haemorrhagic organs including the spleen, heart, brain, gills, but most notably liver and

kidney. The gastrointestinal track was empty of solids but contained transparent fluid similar

to that also seen in the peritoneal cavity.

In contrast to the earlier visits, when Farm 2 was visited on 20/02/19, there were very high

and ongoing mortalities in the fry production systems (>70%). This was both in their onshore

hatchery (supplied with water pumped from the main lake) where eggs were hatched and held

until the fry were approximately 20g, and in the nursery cages on the main lake to which fry

had been transferred. As with the larger fish sampled, affected fry showed erratic swimming

behaviour, skin haemorrhages and severe ascites as the main clinical signs (**Figure 3**).

244
Figure 3 External lesions and clinical signs in diseased Nile tilapia in Lake Volta. A
Ongrowing fish with emaciation slight ascites (arrow), endophthalmia (left arrow head) and
skin purulent abseces (right arrow head). **B** Broodstock with microphthalmia left arrow head),
skin haemorrhages (middle arrow head) and skin ulcers (right arrow head). **C** juvenile with
exophthalmia (left arrow head), ascites (arrow) and loss of scales, excess of mucus and
haemorrhages (right arrow head). Bar = 1cm. **D** Ventral view of juvenile fish presenting
severe ascites black arrow and skin haemorrhages (arrow head). Bar= 1cm.

252
3.2 *Megalocytivirus* and bacterial infections in Farm 2 during period of high mortalities

Four out of the ten fish examined histologically showed mild tissue necrosis in the spleen and

renal haematopoietic tissue with the presence of large numbers of cells showing relative

eosinophilia cytoplasmic and nuclear pleomorphism with margination of chromatin in some

affected nuclei suggestive of a viral infection diffused throughout the tissue **(Figure 4)**.

Figure 4 Histopathological cross sections of tissues from diseased Nile tilapia in Lake Volta. All
sections stained with H&E unless otherwise stated. **A.** Section through the intestine of an infected fry
showing the presence of large numbers of megalocytes in the lamina propria (*) but not affecting the
mucosa or underlying muscularis. Note the presence of isolated megalocytes in the pancreatic acinar
tissue (arrows). Bar = 50µm. **B.** Same fish as in A, showing infiltration and disruption of the
muscularis associated with the presence of megalocytes (arrows). Bar=50 µm. **C.** Pseudobranch
showing extensive necrosis (*). Bar=50µm. **D.** Heart showing necrosis of the ventricular muscle (*),
particularly in the peripheral regions associated with inflammation. Bar=50µm. **E.** Spleen from a fish
from Farm 2 during a mortality episode. Affected cells show pronounced eosinophilia and are
distributed throughout the section (*). Numerous vacuolated macrophage aggregates are also present
(M). Bar = 50µm. **F.** Pronephros showing extensive distribution of megalocytes without associated
tissue necrosis (*). Bar=100µm. Inset shows pale staining of DNA in the cytoplasm of affected cells
(arrows). Feulgen stain. Bar=25µm.

The ultrastructure of affected cells revealed the presence of conspicuous viral particles. Some

cells showed the presence of numerous diffusely spread virions within the hypertrophied

nucleus of affected cells, which also showed degradation of the nuclear membrane (**Figure**

5). Virions were approximately 160 nm in diameter **(Figure 5D)**. Virion morphology

showing icosahedral symmetry with an external double membrane and internal core was

consistent with that of viruses from the genus *Megalocytivirus*. In other cases, the nucleus of

affected cells appeared condensed and densely stained in histological and resin sections.

TEM showed that in these cases the nucleus was tightly packed with virions in various stages

of maturation and with some evidence of formation of 'arrays' **(Figure 5)**. In two of these

cases, a concomitant Gram +ve bacterial infection was also present in the gill and liver.

Incidental findings of gill parasites, myxozoan cysts and monogeneans, both present in low

numbers as well as low grade epitheliocystis were observed. The brain of a single fish

harboured small cysts containing necrotic debris. Gram and Ziehl-Neelsen staining did not

demonstrate the presence of bacteria. Other tissues appeared normal. For the set of samples

collected at the height of the mortalities from Farm 2 on 28/11/18, *Aeromonas jandaei*,

Aeromonas veronii (from skin), *Streptococcus agalactiae* capsular type Ib biotype and

Edwardsiella tarda (from liver and kidney) were recovered.

3.3 PCR and sequence confirmation of Megalocytivirus infection

Within the 7 individuals collected from Farm 2 on the second visit that were analysed with

the protocol proposed by (16), a single fish (fish 1) was clearly positive by PCR for

RSIV/ISKN and a second very weak product of the correct size was also seen in tissues from

Fish 6 (**Supplementary File 4**) The consensus sequence generated from the PCR product

from Fish 1 was confirmed as ISKNV sharing 100% nucleotide identity with ISKNV

accession no <u>AF371960.1</u>. In the phylogenetic analysis the sequence was assigned to the

same lineage as the bulk of the ISKNV sequences (**Figure 6**).

The samples collected from Farm 1 and Farm 2 at the height of the mortalities were found to

be strongly positive when they were retrospectively tested using the (14, 15) recommended

PCR method.

Figure 5 Micrographs of the kidney of diseased Nile tilapia in Lake Volta. A Semithin
section of affected renal tissue showing characteristic cellular hypertrophy (black arrows).
Toluidine Blue stain. Scale bar = 25μm. **B.** Electron micrograph of an individual infected cell
with numerous viral particles (*). Adjacent cells appear uninfected. Scale bar = 2μm. **C.**
Numerous lightly stained developing virions with mature virions. The outer membranes and
central electron lucent core are clearly visible. Scale bar = 500nm. **D.** Mature icosahedral
virion showing detail of the outer caspid and inner membrane with central electron lucent
core.

312
3.4 qPCR results for megalocytivirus from ongrowing tilapia samples

The qPCR results confirmed that all the fish sampled from Farm 1 in the Akuse region were

positive for megalocytivirus. Also, as expected, the samples collected from the first Farm 2

visit during the peak of mortality (28/11/18), were also positive and presented the highest

viral titres in grow out fish with some containing over 3×10^6 copies per sample reaction. In

contrast, the samples collected from grow out fish during the second visit to Farm 2 had

much lower viral copy numbers.

All the archived samples collected in 2017 and March 2018 were negative for ISKNV, when

tested by qPCR (**Supplementary File 5**).

3.5 Fry samples had characteristic megalocytivirus pathology and high copy numbers of

virus

All the fry samples collected from Farm 2 on 20/02/19, when there were reportedly very high

(>90%) losses in that part of their system, were positive for ISKNV by qPCR and these

presented the highest titres in the study with some containing up to 1.5×10^7 copies per

sample reaction (**Table 1**). All fry showed moderate to marked histological and pathological

features of infection with *megalocytivirus*. Splenic tissues were necrotic and associated with

the presence of megalocytes characterised by light sometimes granular cytoplasmic

basophilia and hypertrophied nuclei. Kidney also showed the presence of megalocytes but

usually with only mild cellular necrosis. The lamina propria in the intestine of a single fish

was packed with megalocytes (**Figure 4**), although necrosis appeared to be absent and the

epithelial layer remained intact. Gills showed only minimal focal necrosis, usually affecting

the underlying connective tissues. In some fish the choroidal rete was affected with mild

necrosis and variable numbers of megalocytes and most fish showed mild myositis with few

megalocytes in the skeletal muscle. However, a single fish showed extensive inflammation

and myofibrillar necrosis **(Figure 4)**. Connective tissues of the head and in particular around

the pharyngeal teeth were often infiltrated with megalocytes. Brain and spinal cord appeared

normal. Liver samples were not examined as they were used for virus quantification.

3.6 Other pathological observations

Megalocytivirus -like pathology was not observed in any of the samples taken from the

second visit to Farm 2, two weeks after the peak of mortalities had passed. As with the

samples taken at the first visit, there was evidence of bacterial infection in some individuals,

particularly fish 5, had marked bacterial infection of the spleen, liver and brain (meningitis).

All the samples were positive for the presence of *Streptococcus agalactiae* and negative for

Streptococcus uberis and S*treptococcus dysgalactiae* by PCR **(Supplementary File 6)**.

A range of different potential bacterial species, including *Aeromonas jandaei* and

Plesiomonas shigelloides were recovered from fish from Farm 1 **(Table 1)**, but not as pure

growths or high quantities, suggesting they had a limited role in observed disease in these

animals.

All the samples from Farm 1 and Farm 2 tested for TiLV and from Farm 2 for nodavirus by

qPCR were all negative **(Table 1 and Supplementary File 4)**.

The results   for all the individual fish tested are   shown in Supplementary File 2.

Table 1 Summary of sampling and results

Sampling date & Farm	No. of Fish and size	Farm observations and ‡clinical signs	Bacteriology	Histopathology	Virology Results[¥] [ ] – qPCR $_{RSIV/ISKNV}$ copy number/sample reaction range
18/10/2018 †Farm 1	n=5; 100-300g	Wide scale ongoing mortalities in on growing fish in cages.	*Aeromonas veronii*; *Aeromonas jandaei*; *Plesiomonas shigelloides*; *Chryseobacterium sp*; *Acinetobacter johnsonii*;	Not done	*qPCR $_{TiLV}$ 5/5 -ve *cPCRa $_{ISKNV}$ 4/5 +ve *qPCR $_{RSIV/ISKNV}$ 5/5 +ve [$7.5 \times 10^1 – 4.60 \times 10^5$]
28/11/2018 †Farm 2	n=14; 74-401g	Wide scale ongoing mortalities in on growing fish in cages (**Figure 2**). On shore fry production unit: no unusual mortalities.	*Aeromonas veronii*; *Aeromonas jandaei*; *Streptococcus agalactiae* capsular type Ib biotype 2 (non-haemolytic); *Edwardsiella tarda*	Evidence of Gram negative and Gram positive bacterial and viral# infection (including in same fish)	qPCR $_{TiLV}$ 14/14 -ve qPCR $_{NODA}$ 14/14 -ve *cPCRa $_{ISKNV}$ 12/14 +ve *qPCR $_{RSIV/ISKNV}$ 14/14 +ve [$2.48 \times 10^1 - 3.3 \times 10^6$]
17/12/2018 †Farm 2	n=7; 40-646g	Much lower mortalities in on growing fish in cages than previous sampling visit. On shore fry production unit: no unusual mortalities.	Not done	Evidence of bacterial infection but not of viral infection	qPCR $_{TiLV}$ 7/7 -ve qPCR $_{NODA}$ 7/7 -ve **cPCRb $_{RSIV/ISKNV}$ 1/7 +ve megalocytivirus, sequence = ISKNV** *qPCR RSIV/ISKNV 6/7 +ve [$1 \times 10^0 - 24 \times 10^0$]
20/02/2019 †Farm 2	n=14; 6-9 cms	Limited mortalities in in on growing fish in cages. Very severe mortalities in on shore fry production unit.	Not done	Very severe virus-associated pathology (in tissue samples from all fish).	qPCR TiLV not done qPCR NODA not done cPCRa ISKNV 7/7 +ve qPCR RSIV/ISKNV 13/13 +ve [$5.1 \times 10^5 – 1.51 \times 10^7$]

† Farm 1: approx. 800 tonnes production per year cage unit on lake; Farm 2: approx. 2000 tonnes per year production ongrowing cage culture on lake and fry unit on land
supplied with water pumped from lake.
‡ All moribund fish, including fry, presented similar signs: erratic swimming, lethargy, ascites, swollen, dark spleen, haemorrhagic livers and other organs.
# Haematopoietic cell nuclear and cytoplasmic pleomorphism with marginalisation of chromatin and fine granulation.
¥ qPCR TiLV, qPCR NODA, qPCR RSIV/ISKNV: results of qPCR testing for tilapia lake virus, nodavirus and ISKNV respectively. cPCRa & cPCRb : conventional
PCR using OIE recommended protocols by Kurita et al., 1998 (a) (15) and Rimmer et al., 2012 (b) (16) protocols, results in bold represent index finding, *represents
retrospective   testing   post   index   finding.

364
365 **Figure 6 N-J tree showing the genetic relationship of partial MCP sequence from LV#1**
366 **to a range of RSIV, ISKNV and related Megalocytiviruses.** The GeneBank accession
367 numbers and the host species are included where available.

4 Discussion

The results suggest ISKNV had a significant role in the high mortalities experienced by the two farms that were investigated. Fish sampled from the second farm at the height of the disease outbreak showed severe clinical and pathological signs typically associated with infection by the virus (including visualisation of distinctive megalocytes with characteristic virions). Both, these fish and those sampled earlier from Farm 2, had extremely high copy numbers of RSIV/ISKNV *Megalocytivirus*-like virions and ISKNV was confirmed by gene sequencing.

Although it cannot be confirmed, without testing fish recovered from a range of the other affected farms, the results would be consistent with a recent introduction of the virus onto one or more farms prior to October 2018, that then extended upstream of the dam to other farms resulting in the unusual, widespread and significant mortalities observed. Firstly, there were no observations of typical ISKNV-associated pathology in any previous disease investigations, on the affected or other farms. Secondly, the limited PCR screening in this study of archived samples of diseased tilapia recovered from before the major mortality event including both farms, were all negative. Moreover, (3) also screened for the presence of RSIV/ISKNV by PCR in 2016, including samples from the affected farms, without detecting the virus, or associated pathology.

Iridioviruses are large icosahedral cytoplasmic double⬜stranded DNA viruses, which can infect a wide range of hosts, including invertebrates and poikilothermic vertebrates. The family *Iridoviridae* includes five genera: *Iridovirus, Chloriridovirus, Ranavirus, Lymphocystivirus* and *Megalocytivirus* (19). Fish pathogenic iridoviruses are representatives of *Ranavirus, Lymphocystivirus* and *Megalocytivirus* genera (20, 21). Infectious spleen and kidney necrosis virus (ISKNV), is a member of the genus *Megalocytivirus* (22), and causes

disease in a range of freshwater and marine fish species (Subramaniam, Shariff, Omar, &

Hair-Bejo, 2012; Whittington, Becker, & Dennis, 2010). ISKNV is closely related to red sea

bream iridovirus and both viruses are listed by the OIE as notifiable pathogens (14).

Although tilapia is not listed as a susceptible species by the OIE at present (14), recent

reports from the USA (23) and Thailand (24) suggest that it is a susceptible species likely to

suffer significant mortality. The change in the known host range of virus needs to be

communicated to the international community to prevent future transboundary spread through

movement of infected tilapia.

It was interesting to note that many of the ISKNV positive fish were actively co-infected with

Streptoccocus agalactiae and other bacterial pathogens, presenting severe bacteraemia /

meningitis, as well as ISKNV-associated pathology. The high mortalities on the farms in the

larger on-growing fish may well have been exacerbated by these coinfections.

Although the mortalities were initially confined to on-growing fish in cage culture systems,

the later observations of very high ISKNV associated mortalities in fry shows the virus likely

affects all life stages. As fry are often reared in onshore facilities below the dam and then

translocated to on-growing cages on the main lake, this may have been one of the routes that

disease was rapidly spread after it first emerged. Anecdotally, at the time of writing, farmers

on the sites visited report that mortalities in on-growing facilities have declined, while fry

production continues to be badly affected. It is possible that surviving fry have been exposed

to the virus and then protected against subsequent exposure. This suggests that immunisation

of fry, or use of previously exposed individuals, could represent a practical disease

management strategy. Vaccination as a control strategy may be used to control red sea bream

irididovirus (25, 26) and there are also encouraging reports of its potential effectiveness for

protection against ISKNV in other fish species (27).

Outbreaks of disease that cause significant morbidity and/or mortalities in an aquaculture

operation is always a major concern. This is exacerbated when this appears to represent the

incursion of a new agent into a system, or region (country or zone in a country) which has not

previously been affected. A stark example of this is the epidemic of infectious salmon

anaemia virus (ISAV) which reduced production by three quarters and resulted in severe

economic and social crisis in the developing Chilean Atlantic salmon industry between 2007

and 2010 (28–30).

The Ghanaian authorities have for some time been concerned that the, to date, successful,

expansion of its industry on Lake Volta may be affected by such disease incursions. Partly for

this reason, and also to safeguard the genetic integrity of Lake Volta Nile tilapia strains, they

have tried to limit the culture to locally reared Nile tilapia stocks. However genetic testing by

the Ghanaian Fisheries Commission (Ziddah *et al*., Unpublished Observations) showed that

fish on some of the farms on Lake Volta were likely of imported GIFT strain origin (31) or

hybrids of GIFT and indigenous strains.

If farmers have been illegally sourcing broodstock from Asia and other areas, that would be

an ideal method of translocating pathogens from one region to another. It should be noted

though that ISKNV has also been detected in internationally traded freshwater ornamental

species, theoretically posing another possible introduction route (32)

It is very possible that this is not the first time disease introduction has taken place in Lake

Volta Ghana. The study by (3) showed that outbreaks of *S . agalactiae* investigated in 2016

were all caused by genetically indistinguishable isolates of ST 261, with closest genetic

identity to Asian isolates. Discussions with affected farmers at the time suggested, that *S.*

agalactiae associated mortalities were a relatively recent phenomenon, although the disease

is now clearly endemic to all the areas in the Volta area. Other studies have shown that *S.*

agalactiae ST261 has likely been translocated around the world in association with farmed

tilapia (33).

Although most attention to date has focussed on the emergence and spread of TiLV within

the tilapia industry world-wide, these results also demonstrate that there is a range of other

potential threats to the sustainability of tilapia aquaculture.

Conclusion

This is the first report of Infectious Spleen and Kidney Necrosis Virus (ISKNV) in farmed

tilapia in Africa. ISKNV was found in co-infection with *Streptococcus agalactiae* and other

bacterial pathogens in Lake Volta, Ghana. The correlations seen between the mortality

events, histopathology and viral loads in the tissues suggest that ISKNV was a major cause of

mortalities during the outbreaks. In general, the results support continued efforts to improve

the biosecurity of the industry in Ghana. There is a clear need to strengthen domestic

capability to rapidly diagnose and control emerging disease threats caused by ISKNV and

other pathogens. Further work is also needed to map the distribution of the virus and its

impact, including potential effects on wild fish species, and to implement practical control

strategies.

Acknowledgements

The farmers involved in the investigation are thanked for their support and provision of

information. Support from Defra (contracts FB002 and FX001 for the OIE Collaborating

Centre for Emerging Aquatic Animal Diseases) is gratefully acknowledged.

References

1. FAO. 2018. The State of World Fisheries and Aquaculture 2018 - Meeting the
sustainable development goals.

2. FAO. 2016. The State of World Fisheries and Aquaculture. Contributing to food
security and nutrition for all. Rome.

3. Verner-Jeffreys D, Wallis T, Cano I, Ryder D, Hayden D, Domozoro J, Dontwi J,
Field T, Adjei-Boteng D, Wood G, Bean T, Feis S. 2018. *Streptococcus agalactiae* Multi
Locus Sequence Type 261 is associated with mortalities in the emerging Ghanaian tilapia
industry. J Fish Dis 41:157–179.

4. Jansen MD, Cudjoe K, Brun E. 2018. Investigation of tilapia mortality in Ghana.

5. Plumb JA, Hanson LA. 2011. Tilapia bacterial diseases, p. 445–463. In Plum, JA,
Hanson, LA (eds.), Health Maintenance and Principle Microbial Diseases of Cultured Fishes,
Third edition. Blackwell Publishing Ltd, Iowa.

6. Machimbirike VI, Jansen MD, Senapin S, Khunrae P, Rattanarojpong T, Dong HT.
2019. Viral infections in tilapines: More than just tilapia lake virus. Aquaculture 503:508–
518.

7. Hounmanou YMG, Mdegela RH, Dougnon T V., Achoh ME, Mhongole OJ,
Agadjihouèdé H, Gangbè L, Dalsgaard A. 2018. Tilapia lake virus threatens tilapines farming
and food security: Socio-economic challenges and preventive measures in Sub-Saharan
Africa. Aquaculture 493:123–129.

8.   Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glöckner FO.

2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-

generation sequencing-based diversity studies. Nucleic Acids Res 41:e1.

9.   Shoemaker CA, Xu DH, Garcia JC, Lafrentz BR. 2017. Capsular typing of

Streptococcus agalactiae (Lancefield group B streptococci) from fish using multiplex PCR

and serotyping. Bull Eur Assoc Fish Pathol 37:190–197.

10.   Yáñez MA, Catalán V, Apráiz D, Figueras MJ, Martínez-Murcia AJ. 2013.

Phylogenetic analysis of members of the genus *Aeromonas* based on gyrB gene sequences.

Int J Syst Evol Microbiol 53:875–883.

11.   Delannoy CM, Crumlish M, Fontaine MC, Pollock J, Foster G, Dagleish MP,

Turnbull JF, Zadoks RN. 2013. Human *Streptococcus agalactiae* strains in aquatic mammals

and fish. BMC Microbiol 13:41.

12.   Luo X, Fu X, Liao G, Chang O, Huang Z, Li N. 2017. Isolation, pathogenicity and

characterization of a novel bacterial pathogen *Streptococcus uberis* from diseased mandarin

fish Siniperca chuatsi. Microb Pathog 107:380–389.

13.   Abdelsalam M, Asheg A, Eissa AE. 2013. *Streptococcus dysgalactiae*: An emerging

pathogen of fishes and mammals. Int J Vet Sci Med 1:1–6.

14.   World Organisation for Animal Health OIE. 2018. Chapter 2.3.8 Red sea bream

iridoviral diseaseManual of Diagnostic Tests for Aquatic Animals. OIE, Paris.

15.   Kurita J, Nakajima K, Hirono I, Aoki T. 1998. Polymerase chain reaction (PCR)

amplification of DNA of red sea bream iridovirus (RSIV). Fish Pathol 33:17–23.

16.   Rimmer, A. E., Becker JA, Tweedie A, Whittington RJ. 2012. 2012 Development of a

quantitative polymerase chain reaction (qPCR) for the detection of dwarf gourami iridovirus

(DGIV) and other Megalocytiviruses and comparison with the Office International des

Epizooties reference PCR protocol. Aquaculture 358-359:15. Aquaculture 358–359:155–163.

17.   Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ.

1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search

programs. Nucleic Acids Res 25:3389–3402.

18.   Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular

Evolutionary Genetics Analysis version 6.0. Mol Biol Evol 30:2725–9.

19.   Jancovich JK, Chinchar VG, Hyatt A, Miyazaki T, Williams T, Zhang QY. 2012.

Family Iridoviridae., p. 193–210. In King, AMQ, Adams, MJ, Carstens, EB, Lefkowitz. E.J.

(eds.), Virus Taxonomy: Ninth Report of the International Committee on Taxonomy of

Viruses. Elsevier Academic Press, San Diego, CA.

20.   Subramaniam K, Shariff M, Omar AR, Hair-Bejo M. 2012. Megalocytivirus infection

in fish. Rev Aquac 4:221–233.

21.   Whittington RJ, Becker JA, Dennis MM. 2010. Iridovirus infections in finfish-critical

review with emphasis on ranaviruses. J Fish Dis 33:95–122.

22.   Kurita J, Nakajima K. 2012. Megalocytiviruses. Viruses 4:521–538.

23.   Subramaniam K, Gotesman M, Smith CE, Steckler NK, Kelley KL, Groff JM,

Waltzek TB. 2016. Megalocytivirus infection in cultured Nile tilapia *Oreochromis niloticus*.

Dis Aquat Organ 119:253–258.

24.   Suebsing R, Pradeep PJ, Jitrakorn S, Sirithammajak S, Kampeera J, Turner WA,

Saksmerprome V, Withyachumnarnkul B, Kiatpathomchai W. 2016. Detection of natural

infection of infectious spleen and kidney necrosis virus in farmed tilapia by hydroxynapthol

blue-loop-mediated isothermal amplification assay. J Appl Microbiol 121:55–67.

25.   Shimmoto H, Kawai K, Ikawa T, Oshima SI. 2010. Protection of red sea bream

Pagrus major against red sea bream iridovirus infection by vaccination with a recombinant

viral protein. Microbiol Immunol 54:135–142.

26.   Nakajima K, Maeno Y, Honda A, Yokoyama K, Tooriyama T, Manabe S. 1999.

Effectiveness of a vaccine against red sea bream iridoviral disease in a field trial test. Dis

Aquat Organ 36:73–75.

27.   Dong Y, Weng S, He J, Dong C. 2013. Field trial tests of FKC vaccines against RSIV

genotype Megalocytivirus in cage-cultured mandarin fish (*Siniperca chuatsi*) in an inland

reservoir. Fish Shellfish Immunol 35:1598–1603.

28.   Vike S, Stian Nylund S, Nylund A. 2009. ISA virus in Chile: evidence of vertical

transmission. Arch Virol 154:1–8.

29.   Mardones FO, Martinez-Lopez B, Valdes-Donoso P, Carpenter TE, Perez AM. 2014.

The role of fish movements and the spread of infectious salmon anemia virus (ISAV) in

Chile, 2007-2009. Prev Vet Med 114:37–46.

30.   Godoy MG, Aedo A, Kibenge MJT, Groman DB, Yason C V., Grothusen H,

Lisperguer A, Calbucura M, Avendaño F, Imilán M, Jarpa M, Kibenge FSB. 2008. First

detection, isolation and molecular characterization of infectious salmon anaemia virus

associated with clinical disease in farmed Atlantic salmon (Salmo salar) in Chile. BMC Vet

Res 4:28.

31.   Ponzoni RW, Nguyen NH, Khaw HL, Hamzah A, Bakar KRA, Yee HY. 2011.

Genetic improvement of Nile tilapia (*Oreochromis niloticus*) with special reference to the

work conducted by the WorldFish Center with the GIFT strain. Rev Aquac 3:27–41.

32.   Jung-Schroers V, Adamek M, Wohlsein P, Wolter J, Wedekind H, Steinhagen D.

2016. First outbreak of an infection with infectious spleen and kidney necrosis virus (ISKNV)

in ornamental fish in Germany. Dis Aquat Organ 119:239–244.

33.   Kawasaki M, Delamare-Deboutteville J, Bowater RO, Walker MJ, Beatson S, Ben

Zakour NL, Barnes AC. 2018. Microevolution of *Streptococcus agalactiae* ST-261 from

Australia indicates dissemination via imported tilapia and ongoing adaptation to marine hosts

or environment. Appl Environ Microbiol 84:e00859-1.