Title: Plant mixed lineage kinase domain-like proteins limit biotrophic

pathogen growth

One Sentence Summary: Plants have a protein family that is

structurally homologous to vertebrate mixed lineage kinase domain-

like protein, which induces necroptotic cell death, but these plant

proteins can confer immunity without host cell death.

Short title: Plant MLKLs confer disease resistance

Authors: Lisa Mahdi*[1,2], Menghang Huang*[3], Xiaoxiao Zhang*[3], Ryohei

Thomas Nakano[1,4], Leïla Brulé Kopp[1,5], Isabel M.L. Saur[1], Florence Jacob[1,6,7],

Viera Kovacova[8,9], Dmitry Lapin[1], Jane E. Parker[1,4], James M. Murphy[10,11],

Kay Hofmann[12], Paul Schulze-Lefert[1,4], Jijie Chai[1,3,4] and Takaki Maekawa[1, †]

*These authors contributed equally
[1]Max Planck Institute for Plant Breeding Research, 50829 Cologne, Germany
[2]Current address, Botanical Institute, Cologne Biocenter, University of Cologne, Cologne, 50674,
Germany.
[3]Beijing Advanced Innovation Center for Structural Biology, Tsinghua-Peking Joint Center for Life
Sciences, Center for Plant Biology, School of Life Sciences, Tsinghua University, 100084 Beijing, China.
[4]Cluster of Excellence on Plant Sciences (CEPLAS), Max Planck Institute for Plant Breeding Research,
50829 Cologne, Germany
[5]Ecole normale supérieure Paris, F-75230 Paris cedex 05
[6]Institute of Plant Sciences Paris-Saclay, Centre National de la Recherche Scientifique, Institut National
de la Recherche Agronomique, Université Paris-Sud, Université d'Evry, Université Paris-Diderot,
Sorbonne Paris-Cité, Université Paris-Saclay, 91405 Orsay, France
[7]Current address, PalmElit SAS, F-34980 Montferrier-sur-Lez, France
[8]University of Cologne, Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated
Diseases (CECAD), University of Cologne, 50931 Cologne, Germany.
[9]Current address: Institute for Biological Physics, University of Cologne, Cologne, 50937, Germany.
[10]Walter and Eliza Hall Institute of Medical Research, 1G Royal Parade, Parkville, VIC 3052, Australia;
[11]Department of Medical Biology, University of Melbourne, Parkville, VIC 3052, Australia.
[12]Institute for Genetics, University of Cologne, 50674 Cologne, Germany
[†]Corresponding author. Email: maekawa@mpipz.mpg.de (T.M.)
34

Abstract:

Mixed lineage kinase domain-like (MLKL) protein mediates necroptotic cell

death in vertebrates. We report here the discovery of a conserved protein

family across seed plants that is structurally homologous to vertebrate

MLKL. The *Arabidopsis thaliana* genome encodes three MLKLs with

overlapping functions in limiting growth of obligate biotrophic fungal and

oomycete pathogens. Although displaying a cell death activity mediated by N-

terminal helical bundles, termed HeLo domain, *At*MLKL-dependent immunity

can be separated from host cell death. Cryo-electron microscopy structures of

*At*MLKLs reveal a tetrameric configuration, in which the pseudokinase domain

and brace region bury the HeLo-domains, indicative of an auto-repressed

complex. We also show the association of two *At*MLKLs with microtubules.

These findings, coupled with resistance-enhancing activity and altered

microtubule association of a phosphomimetic mutation in the pseudokinase

domain of *At*MLKL1, point to a cell death-independent immunity mechanism.

Main text:

Regulated cell death (RCD) is intimately connected with innate immunity in plants and animals (*1-3*). A shared feature of several proteins involved in RCD in plants, animals and fungi is a four-helical bundle structure called the HeLo-domain (*4*). HeLo domain-containing MLKL (mixed lineage kinase domain-like protein) mediates necroptosis in animals (*5, 6*), a form of RCD that is proposed to combat pathogens by releasing pro-inflammatory molecules (*7, 8*). Necroptosis is initiated by the plasma membrane (PM)-resident death receptors, with downstream activation of receptor-interacting serine/threonine-protein kinases (RIPKs) leading to the phosphorylation of the pseudokinase domain of the terminal pathway effector, MLKL (*7, 8*). This process assembles monomeric MLKL into pro-necroptotic oligomers (*9*) that translocate to the PM where oligomerized HeLo-domains interfere with membrane integrity (*10-14*). The extent to which RCD in plants and animals is directly responsible for disease resistance is under debate (*3, 15-17*).

To identify novel immune regulators in plants, we searched for HeLo domain-containing proteins in plant genomes by comparing Hidden Markov Models. This analysis identified a protein family that is highly conserved across plants (Fig. 1, Table S1, Supplemental file 1), with a modular structure resembling MLKL (Fig. 1A). The kinase-like domain lacks canonical residues known to underlie phosphoryl transfer (*18*) (Fig. 1B), suggestive of a catalytically inactive pseudokinase (*8*). Hereafter, we refer to these proteins as plant MLKLs. Plant MLKLs additionally possess an extended serine-rich region of varying length after the pseudokinase domain without similarity to any known

domain (Fig. 1A, Supplemental file 1). Plant MLKLs separate into two

subgroups based on sequence similarity (Fig. 1D). *Arabidopsis thaliana*

harbours three *MLKL* genes with *AtMLKL1* and *AtMLKL2* in subfamily I and

AtMLKL3 belonging to subfamily II (Fig. 1D).

To explore similarities between plant and animal MLKLs, we expressed

*At*MLKL2 and *At*MLKL3 for structural analysis using cryo-electron microscopy

(Fig. 2, Fig. S1 and S2). In gel filtration, both proteins eluted at an estimated

molecular weight corresponding to tetramers (Fig. 2A, Fig. S2A). This

contrasts with the vertebrate MLKL protein, which displayed heterogeneity in

a similar assay (*13*). Representative 2D projection views indicate that both

*At*MLKL oligomers form a triangle-like architecture with a 2-fold symmetry (Fig.

S1AB, Fig. S2BC). Further 3D classification and refinement generated density

maps of oligomeric *At*MLKL2 and *At*MLKL3 with a global resolution of 4.1Å

(Fig. S2E) and 3.4Å (Fig. S1D) respectively, based on the gold Fourier Shell

Correlation standard (Fig. S1C, Fig. S2D). The 3D reconstructions show that

both *At*MLKL2 and *At*MLKL3 oligomers are composed of four MLKL

molecules (Fig. 2C, Fig. S3A), confirming that these form tetramers.

Tetramerization of *At*MLKL2 or *At*MLKL3 results in formation of a pyramid-like

structure. Structure alignment of the two final models indicates that tetramers

of two subfamily members are nearly identical (Fig. S3AB). As the quality of

the density of *At*MLKL3 is superior for model building, we limited our structural

analysis to *At*MLKL3.

The N-terminal HeLo domain of *At*MLKL3 forms a four-helix bundle (Fig. S3C)

which superimposed well with the HeLo domains of mouse and human MLKL

(mMLKL and hMLKL, respectively; Fig. 2D). This observation supports the

idea that *At*MLKL3 is a bona fide homolog of vertebrate MLKLs. Nevertheless,

compared to mMLKL, the HeLo domain of *At*MLKL3 packs tightly against its

pseudokinase domain (Fig. S3D). The packing is further strengthened by the

brace region of *At*MLKL3, which contains a string of five helices that

simultaneously interact with the HeLo and pseudokinase domains (Fig. S3C).

The HeLo domains and the brace regions form the core of the *At*MLKL3

tetramer, whereas the pseudokinase domains are presented at the apices of

the pyramid-like structure (Fig. 2C). Hydrophobic packing of α1 helices (Fig.

S3E) from two *At*MLKL3 molecules contributes to formation of a homodimeric

*At*MLKL3. In the tetrameric *At*MLKL3, the two α1-mediated homodimers pack

perpendicularly to each other (Fig. 2G). Four brace regions, which are

positioned nearly in the same plane, are sandwiched between the two

homodimers and exclusively mediate homodimer-homodimer interaction (Fig.

2F). The N-terminal halves of the four brace regions form two homodimer

pairs, and the C-terminal halves another two pairs (Fig. S3E). The

intermolecular interactions lead to further sequestering of the *At*MLKL3 N-

terminal HeLo domain. Taken together, our observations indicate that

*At*MLKL3 is a structural homolog of vertebrate MLKLs and its N-terminal HeLo

domain is sequestered through both intra- and intermolecular interactions.

To determine the role of plant MLKLs in immunity, we challenged

combinatorial loss-of-function mutants of *At*MLKLs (Fig. S4) with different

microbial pathogens. *Atmlkl* single mutants exhibited increased susceptibility

to the obligate biotrophic fungus *Golovinomyces orontii* compared to wild-type

plants and, strikingly, the triple mutant was as susceptible as an *eds1* mutant

(Fig. 3A), which is hyper-susceptible to a number of pathogens including the

obligate biotrophic oomycete *Hyaloperonospora arabidopsidis* (*Hpa*). The

immune response restricting *Golovinomyces* growth in Col-0 wild-type plants

is not associated with host cell death (*19*). In *A. thaliana*, immunity to *Hpa*

isolate Emwa1 is mediated by RPP4, an NLR that requires EDS1 for function

(*20*). RPP4-mediated disease resistance was partially compromised in the

Atmlkl triple mutant, as measured by increased *Hpa* sporangiophore formation

on true leaves (Fig. 3B). We did not detect marked differences between the

susceptibility of wild-type and mutant plants to hemi-biotrophic *Pseudomonas*

syringae pv. *tomato* DC3000 bacteria or the fungal necrotroph *Botrytis cinerea*

(Fig. S5AB). Furthermore, disease resistance mediated by RPS2 and cell

death mediated by RPM1 NLRs were largely retained in the *Atmlkl123* mutant

(Fig. S5CD).

RNA-seq analysis using mock- and *G. orontii*-challenged leaves of wild-type,

Atmlkl23, Atmlkl123 and *eds1* lines showed that the *AtMLKL1* transcript was

induced upon pathogen challenge (7.4 fold) and that induction was fully

dependent on *EDS1* (Fig. 3C, Fig. S6). Further analysis revealed that

*At*MLKLs regulate transcriptional reprogramming both under unchallenged

conditions and upon pathogen attack (Fig. 3C). These data imply a specific

and partly redundant role of *At*MLKLs in resistance to obligate biotrophic

pathogens.

To test whether *At*MLKLs possess cell death activity, possibly regulated by phosphorylation, we introduced single phosphomimetic substitutions at serine residues in the activation loops of *At*MLKLs (Fig. 1B) and expressed in *A. thaliana* leaf protoplasts (see method). Upon overexpression, all wild-type *At*MLKLs elicited cell death, which was as potent as an N-terminal barley NLR cell death module (*21*)(Fig. S7). We found that a phosphomimetic substitution at serine[393] but not serine[395] in *At*MLKL1 enhanced its cell-killing activity (Fig. 4A). However, an alanine substitution at serine[393] did not compromise the cell death activity, suggesting that other serine or threonine residues in the activation loop (Fig. 1B) have a compensatory function. Using transgenic *A. thaliana* lines, we observed that the enhanced susceptibility of *Atmlkl1* and *Atmlkl2* mutants to *G. orontii* was restored by stable transformations of genomic fragments encompassing wild-type *AtMLKL1* and *AtMLKL2*, respectively (Fig. 4B,C and Fig. S8-9). In line with the protoplast assay (Fig. 4A), *At*MLKL1 (S393D) transgenic plants exhibited enhanced resistance to *G. orontii* compared to Col-0 wild type, plants expressing the transgenes *At*MLKL1 or *At*MLKL1 (S395D) (Fig. 4B,C). Unexpectedly, *At*MLKL1 (S393D) transgenic plants did not exhibit cell death lesions or apparent plant growth retardation indicative of autoimmunity (Fig. 4B, Fig. S8A). Collectively our results suggest that plant MLKL function is in part regulated by activation loop conformation, similar to vertebrate MLKLs (*22*), but *At*MLKL1 activity needed to restrict *G. orontii* growth can be separated from host cell death.

We next examined whether the *At*MLKL N-terminal HeLo domain contributes

to its cell death activity, as in animals (*8, 12, 14, 23*). Taking advantage of a

chemically enforced oligomerization system (Fig. S10, see method), we found

that expression of the HeLo domains with the brace region of *At*MLKL1 and

*At*MLKL3 was sufficient for cytotoxic activity and this activity was further

enhanced upon oligomerization (Fig. S10D）. This finding mirrors the activity

of the HeLo domains with the brace region of MLKL (*14*) and is consistent

with our structure-based hypothesis that the full-length MLKL tetramer with

buried HeLo domains represents an inactive form (Fig. 2).

As animal MLKL translocates to the PM upon activation (*10-14*) we examined

whether GFP-tagged *At*MLKL variants associate with the PM using time-

resolved confocal microscopy. We detected mobile punctate signals in the

cytoplasmic space for all *At*MLKL1 variants (Fig. 4DE). Reduced mobility for a

fraction of *At*MLKL1-GFP signals when proximal to the PM (Fig. 4E) is

consistent with PM association. Intriguingly, maximum projection of the time-

lapse images revealed that *At*MLKL1 (wild type)-GFP, the S395D mutant

moved along filamentous structures (Fig. 4E, Supplemental movies 1 and 3;

Fig. S11A). These filamentous structures specifically colocalized with the

microtubule marker (*24*) mCherry-MAP4 (Fig. S11B-C), and association was

detectable upon *G. orontii* invasion (Fig. S12). Intriguingly, no filamentous

structures were detected in the S393D phosphomimetic line despite the

presence of a microtubule array (Fig. S13). These data suggest that specific

phosphorylation modulates the intracellular localization and activities of

*At*MLKL1 via cytoskeletal association.

The plant MLKL family described here is a first example of a non-receptor

immunity component consisting of multiple domains that is structurally shared

between animal and plants. MLKLs in these two kingdoms are likely products

of convergent evolution because their exon-intron structures are unrelated (~4

and 11 exons in *At*MLKLs and hMLKL, respectively). Our work serves as a

template to test structural predictions implicating the presence of a HeLo-

domain fold in a number of plant modular proteins, including the ADR1 family

of NLRs (*4, 25-27*). Despite the cell death activity of a putative HeLo-domain

of ADR1 (*28*), this family confers disease resistance without apparent host cell

death (*29*). This resembles *At*MLKL-mediated defense in wild-type plants to

G. orontii without host cell death, although *At*MLKLs have the capacity to

elicit cell death. These results raise the question of whether host defence and

cell death functions can be further disentangled in animal necroptosis

signalling.

Fig. 1: Plant genomes encode structural homologs of MLKL.

A, A diagram of plant and animal MLKLs. A variably sized serine-rich region that has no similarity to known structures is directly C-terminal to the pseudokinase domain of plant MLKLs.

B, Alignments of the four helical bundles (HeLo domain) and the pseudokinase domains of *A. thaliana* MLKLs (*At*MLKLs) and representative vertebrates. Invariant residues and conservative substitutions in > 50% of the sequences are shown in light blue and grey backgrounds, respectively. The residues responsible for the activation of human or mouse MLKL upon phosphorylation are indicated in the red box. The serine residues examined in this study are shown in red.

C, Plant MLKL-encoding genes across representative plant species. A full list of MLKL-encoding genes in the genomes of 48 seed plants is shown in Table S1.

D, Phylogenetic relationship of plant MLKLs. Neighbor-net analysis discriminates two subgroups of plant MLKLs colored in light blue and orange with bootstrap support of 0.85 (1,000 bootstrap replicates).

Fig. 2. Structure of the *At*MLKL3 tetramer.

A, Top: Gel filtration profiles of *At*MLKL3. Position of standard molecular weight is indicated by arrow. Bottom: Peak fractions on the top were verified by reducing SDS-PAGE with Coomassie Blue staining.

B, Final EM density map of the *At*MLKL3 tetramer. Local resolution of the density map is shown on the right with scale indicated by colors, in angstroms.

C, Top: Final 3D reconstruction of *At*MLKL3 tetramer shown in three orientations. Each monomer of *At*MLKL3 is shown in different colors. Bottom: Cartoon shows the overall structure of *At*MLKL3 tetramer in three orientations. Subdomains of *At*MLKL3 are shown in different colors.

D Superposition of the HeLo domains of *At*MLKL3 (yellow, PDB ID code 6KA4), human MLKL (blue, PDB ID code 2MSV) and mouse MLKL (Pink, PDB ID code 4BTF). RMSD between *At*MLKL3 and mMLKL:3.745, RMSD between *At*MLKL3 and hMLKL:4.010.

E, Electrostatic surface of HeLo domains of *At*MLKL3, human MLKL and mouse MLKL in two orientations.

F, Cartoon shows the brace region of the *At*MLKL3 tetramer. Each monomer of *At*MLKL3 is shown in different colors.

G, Cartoon shows the HeLo domains of the *At*MLKL3 tetramer in two orientations. Each monomer of *At*MLKL3 is shown in different colors.

Fig. 3: Arabidopsis MLKLs confer resistance to powdery and downy mildew pathogens.

A, *Atmlkl* mutants are susceptible to the powdery mildew *Golovinomyces orontii* (arrowheads). Quantification of *G. orontii* DNA at seven days after inoculation is relative to the corresponding Col-0 samples. (n=3, Tukey HSD *p* < 0.05).

B, The *Atmlkl* triple mutant is susceptible to the downy mildew *Hyaloperonospora arabidopsidis*. The number of *Hpa* spores per gram fresh weight of plants is presented. Data were obtained in three independent experiments, each including two biological replicates (n=6). Asterisks indicate significant differences (Tukey HSD *p* < 0.05). The 1st or 2nd true leaves were stained with trypan blue to visualise hyphae growth. Sporangiophores were formed on the true leaves of the *Atmlkl* triple mutant (black arrowheads).

C, Principal component analysis (PCA) of RNA-seq data from mock- and pathogen-challenged leaves collected at 48 hours post *G. orontii* inoculation.

*WT is a segregant line derived from the cross between *Atmlkl1* and *Atmlkl3*. The *eds1* mutant was used as a susceptible control.

Fig. 4: Cell death activity, disease resistance and subcellular localization of phosphomimetic *At*MLKL1 variants.

A, A phosphomimetic substitution in *At*MLKL1 elicited enhanced cell death in *Arabidopsis* mesophyll protoplasts. Luciferase and *At*MLKL1 expression constructs were co-transfected into protoplasts and luciferase activity was measured as a proxy of cell viability at 16 hours post transfection. The positions of serine-to-aspartate residues are indicated in Fig. 1B. The C-terminally HA-tagged variants were expressed under the control of the constitutive cauliflower mosaic virus 35S promoter. (n=3, Tukey HSD *p* < 0.01).

B, Macroscopic phenotype of transgenic *Arabidopsis* expressing phosphomimetic variants of *At*MLKL1 in response to *G. orontii* (arrowheads). Plants were photographed seven days after pathogen challenge.

C, Quantification of *G. orontii* DNA in infected leaves of transgenic plants at 14 days after pathogen challenge (n=4, Tukey HSD *p* < 0.05).

D, Schematic diagram of the confocal images shown in **B**. The white circle indicates GFP signal proximal to the PM.

E, Confocal images of the abaxial epidermis of the *Arabidopsis* transgenic lines expressing phosphomimetic and wild-type variants of *At*MLKL1. GFP signals indicated by white circles were immobile in the examined time period. The corresponding movies are available as supplemental movies 1-3. The asterisk indicates plastidial autofluorescence. Scale bars = 5 µm.

Acknowledgments: We thank the Max Planck Genome Centre Cologne for RNA-Seq and Petra Köchner, Sabine Haigis and Makoto Yoshikawa-Maekawa for technical assistance and Neysan Donnelly for editing the manuscript. We also thank Hirofumi Nakagami, Ton Timmers, Hamid Kashkar and Manolis Pasparakis for helpful suggestions. We also thank Takashi Hashimoto and Stefanie Sprunck for the mcherry-MAP4 and tagRFP-T-Lifeact vectors, respectively.

Data and materials availability: All data underlying the study are deposited at Protein Data bank (PDB): https://www.rcsb.org/) or Gene Expression Omnibus (GEO) database (https://www.ncbi.nlm.nih.gov/geo/).

Author contributions: T.M. conceptualized the project; L.M., M.H., X.Z., R.T.N., L.B.K., I.M.L.S., F.J., V.K., D.L., K.H., T.M. performed the investigations. J.P. J.M.M., P.S-L. J.C. T.M. validated the data, J.P. J.M.M., P.S-L. J.C. T.M. supervised the work, J.C. and T.M. wrote the original draft of the manuscript, J.P. P.S-L., J.M.M. reviewed and edited the manuscript.

Competing interests: The authors declare no competing interests.

Data and materials availability: All data needed to replicate the work are deposited in the Protein Data Bank (PDB) or Gene Expression Omnibus (GEO) database. Plants and plasmids described in the manuscript are available upon request.

References

1.   N. S. Coll, P. Epple, J. L. Dangl, Programmed cell death in the plant
immune system. *Cell Death Differ* **18**, 1247-1256 (2011).
2.   W. G. van Doorn, Classes of programmed cell death in plants,
compared to those in animals. *J Exp Bot* **62**, 4749-4761 (2011).
3.   F. K. Chan, N. F. Luz, K. Moriwaki, Programmed necrosis in the cross
talk of cell death and inflammation. *Annu Rev Immunol* **33**, 79-106
(2015).
4.   A. Daskalov *et al.*, Identification of a novel cell death-inducing domain
reveals that fungal amyloid-controlled programmed cell death is related
to necroptosis. *Proc Natl Acad Sci U S A* **113**, 2720-2725 (2016).
5.   L. Sun *et al.*, Mixed lineage kinase domain-like protein mediates
necrosis signaling downstream of RIP3 kinase. *Cell* **148**, 213-227
(2012).
6.   J. Zhao *et al.*, Mixed lineage kinase domain-like is a key receptor
interacting protein 3 downstream component of TNF-induced necrosis.
Proc Natl Acad Sci U S A **109**, 5322-5327 (2012).
7.   T. Vanden Berghe, A. Linkermann, S. Jouan-Lanhouet, H. Walczak, P.
Vandenabeele, Regulated necrosis: the expanding network of non-
apoptotic cell death pathways. *Nat Rev Mol Cell Biol* **15**, 135-147
(2014).
8.   E. J. Petrie, P. E. Czabotar, J. M. Murphy, The Structural Basis of
Necroptotic Cell Death Signaling. *Trends Biochem Sci* **44**, 53-63 (2019).
9.   E. J. Petrie, J. M. Hildebrand, J. M. Murphy, Insane in the membrane: a
structural perspective of MLKL function in necroptosis. *Immunol Cell*
Biol **95**, 152-159 (2017).
10. X. Chen *et al.*, Translocation of mixed lineage kinase domain-like
protein to plasma membrane leads to necrotic cell death. *Cell Res* **24**,
105-121 (2014).
11. Y. Dondelinger *et al.*, MLKL compromises plasma membrane integrity
by binding to phosphatidylinositol phosphates. *Cell Rep* **7**, 971-981
(2014).
12. J. M. Hildebrand *et al.*, Activation of the pseudokinase MLKL unleashes
the four-helix bundle domain to induce membrane localization and
necroptotic cell death. *Proc Natl Acad Sci U S A* **111**, 15072-15077
(2014).
13. H. Wang *et al.*, Mixed lineage kinase domain-like protein MLKL causes
necrotic membrane disruption upon phosphorylation by RIP3. *Mol Cell*
54, 133-146 (2014).
14. G. Quarato *et al.*, Sequential Engagement of Distinct MLKL
Phosphatidylinositol-Binding Sites Executes Necroptosis. *Mol Cell* **61**,
589-601 (2016).
15. A. Bendahmane, K. Kanyuka, D. C. Baulcombe, The Rx gene from
potato controls separate virus resistance and cell death responses.
Plant Cell **11**, 781-792 (1999).
16. N. S. Coll *et al.*, Arabidopsis type I metacaspases control cell death.
Science **330**, 1393-1397 (2010).
17. J. S. Pearson, J. M. Murphy, Down the rabbit hole: Is necroptosis truly
an innate response to infection? *Cell Microbiol* **19**, (2017).

18.   G. Manning, D. B. Whyte, R. Martinez, T. Hunter, S. Sudarsanam, The
366   protein kinase complement of the human genome. *Science* **298**, 1912-
367   1934 (2002).
368   19.   W. Wang, Y. Wen, R. Berkey, S. Xiao, Specific targeting of the
369   Arabidopsis resistance protein RPW8.2 to the interfacial membrane
370   encasing the fungal Haustorium renders broad-spectrum resistance to
371   powdery mildew. *Plant Cell* **21**, 2898-2913 (2009).
372   20.   E. A. van der Biezen, C. T. Freddie, K. Kahn, J. E. Parker, J. D. Jones,
373   Arabidopsis RPP4 is a member of the RPP5 multigene family of TIR-
374   NB-LRR genes and confers downy mildew resistance through multiple
375   signalling components. *Plant J* **29**, 439-451 (2002).
376   21.   F. Jacob *et al.*, A dominant-interfering camta3 mutation compromises
377   primary transcriptional outputs mediated by both cell surface and
378   intracellular immune receptors in Arabidopsis thaliana. *New Phytol* **217**,
379   1667-1680 (2018).
380   22.   J. M. Murphy *et al.*, The pseudokinase MLKL mediates necroptosis via
381   a molecular switch mechanism. *Immunity* **39**, 443-453 (2013).
382   23.   M. C. Tanzer *et al.*, Evolutionary divergence of the necroptosis effector
383   MLKL. *Cell Death Differ* **23**, 1185-1197 (2016).
384   24.   J. Marc *et al.*, A GFP-MAP4 reporter gene for visualizing cortical
385   microtubule rearrangements in living epidermal cells. *Plant Cell* **10**,
386   1927-1940 (1998).
387   25.   A. R. Bentham, R. Zdrzalek, J. C. De la Concepcion, M. J. Banfield,
388   Uncoiling CNLs: Structure/Function Approaches to Understanding CC
389   Domain Function in Plant NLRs. *Plant Cell Physiol* **59**, 2398-2408
390   (2018).
391   26.   C. Barragan *et al.*, RPW8/HR Repeats Predict NLR-dependent Hybrid
392   Performance. *bioRxiv*, 559864 (2019).
393   27.   L. M. Jubic, S. Saile, O. J. Furzer, F. El Kasmi, J. L. Dangl, Help
394   wanted: helper NLRs and plant immune responses. *Curr Opin Plant*
395   *Biol* **50**, 82-94 (2019).
396   28.   S. M. Collier, L. P. Hamel, P. Moffett, Cell death mediated by the N-
397   terminal domains of a unique and highly conserved class of NB-LRR
398   protein. *Mol Plant Microbe Interact* **24**, 918-931 (2011).
399   29.   V. Bonardi *et al.*, Expanded functions for a family of plant intracellular
400   immune receptors beyond specific recognition of pathogen effectors.
401   *Proc Natl Acad Sci U S A* **108**, 16463-16468 (2011).

Methods

Plant material and growth conditions

The Arabidopsis thaliana (L.) Heynh. ecotype Columbia (Col-0) was *used in*
this study. The T-DNA insertional mutants(1, 2) (SALK_041569c (AtMLKL1),
SALK_124412c (AtMLKL2) and GABI_491E02 (AtMLKL3) were obtained
from the Nottingham Arabidopsis Stock Centre (NASC). Double and triple
mutants of *Atmlkl* were generated by crossing the T-DNA insertion lines. A
segregant line derived from the cross between the *Atmlkl1* and *Atmlkl3*
mutants were used as a wild-type line in addition to Col-0 wild-type. Each
genotype was confirmed by PCR. The *eds1-2* mutant was described
previously (*3*).

The transgenic lines expressing AtMLKL1 variants or AtMLKL2 fused to a
monomeric green fluorescence protein were established in *Atmlkl1* and
Atmlkl2 mutant backgrounds, respectively. Genomic fragments including
coding region and native *cis*-regulatory sequence were amplified by PCR from
Col-0 genomic DNA and cloned into pENTR/D-TOPO (Thermo Fisher
Scientific, Waltham, MA, USA). The phosphomimetic substitutions were
introduced using the QuikChange Lightning site-directed mutagenesis kit
(Agilent Technologies, Santa Clara, CA, USA). Resulting entry vectors were
transferred into pGWB550(*4*) using LR clonase II (Thermo Fisher Scientific).
Plants were transformed by the floral dip method (*5*) with *Agrobacterium*
tumefaciens strain GV3101 harbouring pMP90RK (*6*). Plant growth conditions
were described previously (*7*). Primer sequences for genotyping and plasmid
construction are listed in Table S3.

Sequence analysis of plant and animal MLKLs

Sequence similarity within the animal and plant families was established by the generalized profile method (*8*). Sequences were aligned by the L-INS-I method of the MAFFT alignment software (*9*), followed by minor manual editing of ambiguously aligned regions. Sequence similarity between the animal and plant MLKL families was established by Hidden Markov Model (HMM)-to-HMM comparison using the HHSEARCH package (*10*). The 99,696 orthogroups (OGs) among 52 plant species have been established recently (*11*). An OG containing *At*MLKLs was used for the neighbor-net analysis of codon-aligned nucleotide sequence as descried previously (*12*). The sequence from the papaya genome was excluded in this study.

Protein expression and purification

Full length *At*MLKL3 (residues 1-701) with an engineered C-terminal 6×His tag was generated by standard PCR-based cloning strategy and its identity was confirmed by sequencing. The protein was expressed in sf21 insect cells using the vector pFastBac 1 (Invitrogen). One litre of cells (2.5×10^6 cells ml^{-1}, medium from Expression Sysems) was infected with 20 ml baculovirus at 28°C. After growth at 28°C for 48 hours, the cells were harvested, re-suspended in the buffer containing 50 mM Tris-HCl pH 8.0 and 300 mM NaCl, and lysed by sonication. The soluble fraction was purified from the cell lysate using Ni2+-nitrilotriacetate affinity resin (Ni-NTA, Qiagen). The protein was then further purified by further purified by gel filtration (Superose 6, 10/30; GE Healthcare). For cryo-EM investigation, the purified protein was concentrated to about 0.3 mg/mL in buffer containing 50 mM Tris-HCl pH 8.0, 300 mM NaCl and 3 mM DTT.

The construct of full length *At*MLKL2 (residues 1-711) with N-terminal GST

tag was cloned into the pGEX-6P-1 vector (GE Healthcare), and was

expressed in *Escherichia coli* strain BL21(DE3; Novagen) at 16 °C. After

isopropyl-β-D-thiogalactopyranoside (IPTG; Sigma) induction for 12 h, cells

were harvested and re-suspended in buffer containing 50 mM Tris-HCl pH 8.0

and 300 mM NaCl, and lysed by sonication. The soluble fraction was purified

from the cell lysate using Glutathione Sepharose 4B beads (Invitrogen). The

proteins were then digested with PreScission protease (GE Healthcare) to

remove the GST tag and further purified by gel filtration (Superose 6, 10/30;

GE Healthcare). For cryo-EM investigation, the purified protein was

concentrated to about 0.3 mg/mL in buffer containing 50 mM Tris-HCl pH 8.0,

300 mM NaCl and 3 mM DTT.

Cryo-EM sample preparation and data collection

For cryo-EM analysis, an aliquot of 3.5 µl *At*MLKL2 or *At*MLKL3 protein was

applied to a holey carbon grids (Quantifoil Cu 1.2/1.3, 200 mesh) glow-

discharged (Harrick Plasma) with a middle force for 30 s after evaluating for 2

min. The grids were blotted by a pair of 55 mm filter papers (TED PELLA,

INC.) for 3-3.5 s at 8 °C with 100% humidity and flash-frozen in liquid ethane

using a FEI Vitrobot Marked IV. Cryo-EM data were collected on Titan Krios

electron microscope operated at 300 kV and a Gatan K2 Summit direct

electron detection camera (Gatan) using eTas. Micrographs were recorded in

super-resolution mode at a nominal magnification of 22500 ×, resulting in a

physical pixel size of 1.30654 Å per pixel. Defocus values varied from -1.7 µm

to -2.3 µm for data set. The dose rate was 10.6 electron per pixel per second.

Exposures of 8.0 s were dose-fractionated into 32 sub-frames, leading to a

total accumulated dose of 50 electrons per Å2. In total, two batches of data

were collected, one for *At*MLKL3 and another for *At*MLKL2.

Image processing and 3D reconstruction

A total of 1,434 and 1,828 raw images stacks of *At*MLKL3 and *At*MLKL2

acquired under super-resolution mode, were 2x binned processed using

MotionCor2 (*13*), generating aligned, dose-weighted and summed

micrographs in a pixel size of 1.30654 Å per pixel. CTFFIND4 (*14*) was used

to estimate the contrast transfer function (CTF) parameters. After the removal

of bad micrographs via the evaluation of CTF parameters, remaining images

were processed in RELION (*15*). Approximate 2,000 particles were manually

picked and 2D-classified to generate templates for auto-picking. 983,779 and

1,135,463 autopicked particles for *At*MLKL3 and *At*MLKL2 respectively were

then used for reference-free 2D classification, to remove contaminants and

bad particles. The left good particles were subjected to 3D classification using

initial 3D reference model obtained by *ab initio* calculation from Relion3.0.

Particles from good classes that possess density map with better overall

structure features were selected for the 3D refinement. The final 3D

refinement using D2 symmetry resulted in reconstructions of *At*MLKL3 and

*At*MLKL2 tetramer at resolution of 3.4 Å and 4.1 Å, the resolutions were

determined by gold-standard Fourier shell correlation. Local resolution

distribution was evaluated using Relion.

Model building and refinement

EM density map of *At*MLKL3 was used to build the model de novo, as the

overall resolution of map density was efficient to display side chains. The

model of *At*MLKL3 was manually built into the density in COOT(*16*), and was

refined against the EM map by PHENIX(*17*) in real space with secondary

structure and geometry restraints. The refined *At*MLKL3 model was docked

into the density of *At*MLKL2. The sequence of the docked *At*MLKL3 model

was changed to that of *At*MLKL2 under COOT and the *At*MLKL2 model with

corrected sequence was subjected to refinement by PHENIX. The C-terminal

serine-rich region of *At*MLKL2 or *At*MLKL3 is much less well defined in the

density and is not included in the models. Final model of *At*MLKL3 and

*At*MLKL2 was validated using MolProbity and EMRinger in PHENIX package.

The structures of human MLKL (*18*) and mouse MLKL (*19*) were used for the

superposition of the HeLo domains as shown in Fig. 2D.Table S2 summarized

the model statistics.

RNA sequencing

Total mRNA from leaves was obtained at 48 hours after challenge with

conidia of *Golovinomyces orontii* using the RNeasy plant mini kit (Qiagen).

RNA sequencing (RNA-Seq) libraries were prepared by the Max Planck

Genome Centre Cologne (Cologne, Germany) using the Illumina TruSeq

stranded RNA sample preparation kit (Illumina). The resulting libraries were

subjected to 150-bp single-end sequencing using the Illumina HiSeq3000

(Illumina). Mapping of sequenced reads onto the *Arabidopsis thaliana* gene

model (TAIR10), principal component analysis, and differential gene

expression analysis were performed in the CLC Genomics Workbench

(Qiagen, ver. 10.1.2) using the tool 'RNA-Seq with the default parameter

setting. The data derived from Col-0 and WT were pooled as data of wild-type

lines in the analysis. The heat map of 93 genes differentially expressed

between wild type lines and *Atmlkl* mutants (|log2FC|>1 and false discovery

rate (FDR) < 0.05) were generated using the R package (ver. 1.08) with the

pheatmap function. Gene ontology enrichment analysis was performed using

the PANTHER classification system (http://pantherdb.org/) with default

settings for *Arabidopsis thaliana*. The RNA-Seq data generated in this study

have been deposited in the Gene Expression Omnibus (GEO) database

under accession number GSE129011.

Transient gene expression in *Arabidopsis* protoplasts

Isolation, transfection and luciferase activity measurement of Arabidopsis

protoplasts were performed as described previously (*20*). Protoplasts were

isolated from the leaves of two-week-old *Arabidopsis* plants grown in liquid 1 x

Murashige and Skoog medium. Coding sequences (CDS) of *AtMLKL1* and

AtMLKL2 without stop codons were initially cloned into pENTR/D-TOPO

(Thermo Fisher Scientific). The CDS of *AtMLKL3* without a stop codon was

chemically synthesized and cloned into pENTR221 (Thermo Fisher Scientific).

Two synonymous substitutions (G1371T and A1413G) were introduced into

AtMLKL3 CDS to remove restriction sites that hamper the DNA synthesis.

Entry clones were transferred into the gateway cloning-compatible pAMPAT-

GW-mYFP, pAMPAT-GW-3xHA, or pAMPAT-GW expression vectors (*21*),

which are derivatives of pAMPAT-MCS (accession number: AY436765).

Primers sequences of for plasmid construction are listed in Table S3. pENTR-

tagRFP-T-Lifeact (*22*) was transferred into pAMPAT-GW. The expression

vectors for *Hv*MLA(1-160aa), MAP4 and Tub6 were described previously (*21,*

23, 24).

Protoplast viability assay

Following protoplast transfection and regeneration, Evans blue dye dissolved

in water was added to the samples to a final concentration of 0.04% (w/v).

The stained cells were examined under a standard microscope. For

luciferase-based viability assay luciferase and *At*MLKL expression constructs

were co-transfected into protoplasts and luciferase activity was measured as

a proxy of cell viability.

Induced oligomerization

Two domains of FKBP (F36V) tagged with HA without N-myristoylation signal

were PCR amplified from pC4M-FV2E plasmid (ARIAD, Cambridge, MA,

USA). *Nco*I and *Hind*III restriction sites were added to the 5' end of forward

and reverse primers, respectively. The digested PCR fragment with *Nco*I and

*Hind*III were ligated into the same restriction sites present between the attR2

and the terminator sequences of pAMPAT-GW expression vector. The

resulting vector is named pAMPAT-GW-FV2E-HA. Coding regions

corresponding to the N- and C-terminal luciferase fragments (nLUC and

cLUC) were PCR amplified from nLUC and cLUC expression vectors (*25*) and

cloned into pENTR/D-TOPO. Respective entry clones were transferred into

pAMPAT-GW-FV2E-HA. B/B homodimerizer (also known as AP20187 ligand)

were purchased from TakaraBio, Japan. After transfection, protoplasts in

incubation buffer (i.e. WI solution (*26*)) were separated into two tubes and

added the same amount of incubation buffer supplemented with B/B

homodimerizer (250 nM at the final concentration) or ethanol as solvent

control. Primer sequences for the plasmid construction are listed in Table S2.

We were able to reconstitute luciferase activity of co-expressed N- and C-

terminal halves of luciferase fused to 2xDmrB domains in a ligand-specific

manner (Fig. S8b) and the ligand itself did not affect the luciferase reporter

assay in protoplasts (Fig. S8c).

Pathogen infection assay

The *G. orontii* infection assay was performed as described previously (*27*). *G.*

orontii DNA was quantified by qPCR at indicated time points after inoculation

of conidia and normalised using the amount of plant specific gene

(AT3G21215). The *Hpa* infection assay was performed as previously

described (*28*). Lactophenol-trypan blue staining was described previously

(*29*). The *B. cinerea* strain B05.10 was used in this study. Droplet inoculation

of six-week-old plants was performed as described previously (*30*), except

that 2 µl of the spore solution were used on each side of the leaf and two

leaves of similar age were used per plant. *B. cinerea* DNA was quantified by

qPCR as previously described (*31*). *Pseudomonas syringae* pv. *tomato* (*Pst*)

DC3000 and *Pst* DC3000 expressing *AvrRpt2* or *AvrRpm1* were used in this

study. *Pst* growth assays and ion leakage measurement following bacterial

infiltration were performed as described previously (*7*).

Immunoblot assays

Primary antibodies were monoclonal antibodies from mouse: α-GFP (JL-8, 1:5000, Takara, Shiga, Japan) or rat: α-HA (3F10, 1:1000, Sigma-Aldrich, St. Louis, MO, USA). Goat α-mouse IgG-HRP (1:10000, Santa Cruz Biotechnology, Dallas, TX, USA) or goat α-rat IgG-HRP (1:10000, Sigma-Aldrich) were used as secondary antibodies. The detailed procedure is described in (*7*).

Biolistic transient gene expression

Biolistic delivery of plasmid DNA into the abaxial epidermis of leaves was essentially performed as described previously (*32*). Leaves were detached immediately before bombardment and the bombarded leaves were transferred to 1% agar plates supplemented with 85 μM benzimidazole and incubated at 20°C for 15 h before confocal microscopy.

Confocal microscopy

Transfected protoplasts in a chamber slide (Nunc Lab-Tek, Thermo Fisher Scientific) with incubation buffer (i.e. WI solution, (*26*)) or 2-5 mm leaf discs prepared form rosette leave of 4-5-week-old plants were observed under a confocal microscope (LSM880, Carl Zeiss) equipped with a 40X water-immersion and a 63X oil-immersion objective. Lambda stack images were obtained for spectral imaging. Images were analyzed and processed with ZEN Software (Carl Zeiss) and ImageJ (NIH). In Fig. 4E, confocal images were acquired over time (for wild type, 124 seconds; for S393D, 194 seconds; for

622   S395D, 166 seconds) and used for maximum intensity projection (bottom

623   panels). Representative single frame images are shown (top panels).

Supplemental data

Supplemental Fig. 1: 3D reconstruction of *At*MLKL3 tetramer

A, Representative cryo-EM image of *At*MLKL3 tetramer. Representative particles are indicated with black circles.
B, Representative top and side views of 2D class averages of *At*MLKL3 tetramer.
C, Fourier shell correlation (FSC) curves at 0.5 and 0.143 of the 3D reconstruction of *At*MLKL3 tetramer.
D, Flowchart representing cryo-EM data processing and 3D reconstruction of *At*MLKL3 tetramer.

Supplemental Fig. 2: Purification and 3D reconstruction of *At*MLKL2 tetramer

A, Top: Gel filtration profile of *At*MLKL2 protein. Position of standard molecular weight is indicated by arrow. Bottom: Peak fractions in the top were verified by SDS-PAGE with Coomassie Blue staining.
B, Representative cryo-EM images of the *At*MLKL2 tetramer. Representative particles are indicated with black circles.
C, Representative top and side views of 2D class averages of *At*MLKL2 tetramer.
D, Fourier shell correlation (FSC) curves at 0.5 and 0.143 of the 3D reconstruction of *At*MLKL2 tetramer.
E, Flowchart representing cryo-EM data processing and 3D reconstruction of *At*MLKL2 tetramer.

Supplemental Fig. 3: Inter- and intra-domain interactions of *At*MLKL3 tetramer and structural comparison of the *At*MLKL2 and *At*MLKL3

A, Superposition of *At*MLKL2 (grey) and *At*MLKL3 (blue). RMSD between monomer of *At*MLKL2 and *At*MLKL3 :0.985.

B, The sequence of *At*MLKL3 model (blue) was docked into *At*MLKL2 map (gray).

C, Top: Cartoon representation of *At*MLKL3 monomer in two orientations. Subdomains of *At*MLKL3 are shown in different colors, the interacting regions between domains are highlighted with open frames. Bottom left: Detailed interactions of HeLo domain and pseudokinase domain for the red-framed region. Bottom right: Detailed interactions of HeLo domain and brace region for the blue-framed region.

D, Structural comparison of the monomer *At*MLKL2 (left), *At*MLKL3 (middle, PDB ID code 6KA4) and *m*MLKL (right, PDB ID code 4BTF).

E, Top left: Cartoon showing HeLo domain and brace region of *At*MLKL3 tetramer. Top right: Detailed interactions of HeLo domains for the green-

framed region. Bottom left: Detailed C-terminal interactions of brace regions for the blue-framed region. Bottom right: Detailed N-terminal interactions of brace regions for the blue-framed region.

Supplemental Fig. 4: *Atmlkl* mutants exhibit no obvious growth defect.

A, The Arabidopsis *MLKL* genes and T-DNA insertion sites of the *Atmlkl* mutants. White and black boxes indicate non-coding and coding exons, respectively. A gene model for *AtMLKL3* was deduced from other plant *MLKL* structures.

B, Representative images of four-week-old plants of Col-0, wild type*, single mutants of *Atmlkl1*, *Atmlkl2*, *Atmlkl3*, double mutants *Atmlkl1 2*, *Atmlkl1 3*, *Atmlkl2 3*, and the triple mutant *Atmlkl1 2 3*. *The wild type is a segregant line derived from the cross between *Atmlkl1* and *Atmlkl3*. The plants were initially grown on Murashige and Skoog-agar plates for two weeks and subsequently transferred to Jiffy pots rehydrated in water with a fertilizer. Plants were grown for an additional two weeks under short-day conditions.

Supplemental Fig. 5: *Arabidopsis* MLKLs do not confer resistance to *Botrytis cinerea* and *Pseudomonas syringae* DC3000.

A, *Botrytis cinerea* DNA was quantified by qPCR at three days after spore inoculation and normalized using the amount of plant specific gene (see method). Amounts are presented relative to the corresponding Col-0 samples. The wild type is a segregant line derived from the cross between *Atmlkl1* and *Atmlkl3* (see Fig. S4). Data were obtained in three independent experiments (n=3).

B, Log_{10}-transformed colony forming units of *Pseudomonas syringae* DC3000 per cm^2 of *A. thaliana* leaves at two days after pathogen infiltration. The *eds1* mutant was used as a susceptible control. Data were obtained in three independent experiments (n=3).

C, Log_{10}-transformed colony forming units of *P. syringae* DC3000 expressing AvrRpt2 per cm^2 of *A. thaliana* leaves at three days after pathogen infiltration. The *rpm1 rps2* mutant was used as a susceptible control. Data were obtained in three independent experiments (n=3).

D, Ion leakage assay in *A. thaliana* leaves upon infiltration with *P. syringae* DC3000 expressing *AvrRpm1*. Samples were collected 30 min post infiltration with the bacterial suspension (OD600=0.1). The *rpm1 rps2* mutant was used as a negative control. Data were obtained in three independent experiments (n=3).

n.s. not significant.

Supplemental Fig. 6: Heat map of 93 genes differentially expressed in *Atmlkl123* in comparison to wild type. The gene ontology (GO) term biotic or abiotic stresses (GO:0006950, asterisks) was overrepresented in the down-regulated transcripts in *Atmlkl123* compared to wild type lines. Mock- and

711 pathogen-challenged leaves were collected at 48 hours post *G. orontii*
712 inoculation. *WT is a segregant line derived from the cross between *Atmlkl1*
713 and *Atmlkl3*.

Supplemental Fig. 7: *At*MLKLs are capable of eliciting cell death in *Arabidopsis* mesophyll protoplasts.

A, C-terminally HA-tagged and **B**, C-terminally mYFP-tagged *At*MLKLs were
expressed under the constitutive cauliflower mosaic virus 35S promoter.
Expression constructs for *At*MLKLs were transfected into protoplasts and cells
were stained with Evans blue at 16 hours post transfection. Unstained cells
(Evans blue-negative cells) were counted as living cells. For **B**, The N-
terminal signalling region of barley NLR protein (HvMLA$_{1-160}$: (7) was used as
a positive control. Data were obtained in three independent transfections
(n=3) and different letters indicate statistically significant differences (Tukey
HSD, $p < 0.01$).

C. Phosphomimetic substitutions in *At*MLKL2 or *At*MLKL3 do not alter cell
death in *Arabidopsis* mesophyll protoplasts.

Expression constructs for luciferase and each of the *At*MLKL variants were
co-transfected into protoplasts and luciferase activity was measured as a
proxy for cell viability at 16 hours post transfection. The positions of serine-to-
aspartate substitutions are indicated in Fig. 1B. The C-terminally HA-tagged
variants were expressed under the 35S promoter. Data were obtained in three
independent transfections (n=3). n.s. not significant.

D, Western blot analysis of expression of C-terminally HA-tagged *At*MLKLs in
Arabidopsis mesophyll protoplasts. Total protein extracts were collected at
seven hours post transfection. Samples that were transfected with the
corresponding empty vector or the expression construct for *At*MLKL2-mYFP
were used as negative control.

Supplemental Fig. 8: Genetic complementation of the *Atmlkl1* mutant.

A, Macroscopic phenotype of transgenic *Arabidopsis* lines expressing wild-
type or phosphomimetic variants of *At*MLKL1-GFP under the native *cis*-
regulatory sequence in in *Atmlkl1* in response to *G. orontii*. Plants were
photographed seven days after pathogen challenge.

B, Western blot analysis of the C-terminally GFP-tagged *At*MLKL1 variants in
the stable transgenic lines. Total protein extracts were collected from two-
week-old plants of the transgenic lines grown on sterile Murashige and Skoog
solid media.

Supplemental Fig. 9: Genetic complementation of the *Atmlkl2* mutant.

A, Macroscopic phenotype of transgenic *Arabidopsis* lines expressing
*At*MLKL2-GFP under the native *cis*-regulatory sequence in *Atmlkl2*. Four-

week-old plants were inoculated with *G. orontii* spores and plants were
photographed at 14 days after pathogen challenge.

B, Western blot analysis of C-terminally GFP-tagged *At*MLKL2. Total protein
extracts were collected from two-week-old plants of the transgenic lines grown
on sterile Murashige and Skoog solid media.

The asterisk and arrow indicate non-specific and specific bands, respectively.

C, Quantification of *G. orontii* DNA in infected leaves. Four-week-old plants
were inoculated with spores of the powdery mildew and plants were examined
at 14 days after the pathogen challenge. Fungal DNA was quantified by qPCR
and normalised using the amount of plant specific gene (AT3G21215). The
relative amounts to corresponding Col-0 wild-type samples were presented.
Experiments were repeated three times (n=3) and the asterisk indicates a
statistically significant difference (Tukey HSD, *p* < 0.05).

Supplemental Fig. 10: N-terminal domains of *At*MLKL1 and 3 were
sufficient to elicit cell death and their enforced oligomerization
potentiated the activities.

A, Schematic of chemically induced oligomerization. AP20187 (Dimerizer), a
synthetic cell-permeable ligand, induces homodimerization of fusion proteins
containing the DmrB domain. Dimerizer–dependent oligomerization is
facilitated by a tandem fusion of DmrB domains.

B, Dimerizer–induced reconstitution of luciferase activity of the N- or C-
terminal halves of luciferase fused with 2 x DmrB domains. Data were
obtained with eight independent transfections (n=8). Luciferase activity was
measured at 16 hours post transfection. The luciferase activity was
statistically higher in the presence of dimerizer (one sample *t*-test, *p* < 0.01).

C, Dimerizer does not influence the luciferase reporter assay. Data were
obtained with seven independent transfections (n=7). Luciferase activity was
measured at 16 hours post transfection. No statistically significant differences
were detected (one sample *t*-test).

D, HeLo domain plus brace region of *At*MLKL1 and *At*MLKL3 were sufficient
to elicit cell death and their enforced oligomerization potentiated the activities.
Expression constructs for luciferase and each of the *At*MLKL fusion proteins
indicated in the figure were co-transfected into protoplasts and luciferase
activity was measured as a proxy of cell viability at 16 hours post transfection.
Relative luciferase activities compared to those of the empty vector control
with dimerizer were plotted. Data were obtained with eight independent
transfections (n=9). Asterisks indicate statistically significant differences
(Tukey HSD, *p* < 0.01)

E, Western blot analysis of the HeLo domain plus brace region constructs
expressed in *Arabidopsis* mesophyll protoplasts. The fusion proteins were
expressed as C-terminally HA-tagged proteins. Total protein extracts were
collected at seven hours post-transfection.

B-E, Mesophyll protoplasts were prepared from the triple mutant *Atmlkl123*
and AP20187 (Dimerizer) was added after transfection (see method).

Supplemental Fig. 11: Subcellular localization of *At*MLKLs in protoplasts.

A, Subcellular localization of *At*MLKLs in *Arabidopsis* mesophyll protoplasts.
The C-terminally monomeric YFP (mYFP)-tagged variants were expressed
under the 35S promoter. Representative confocal images were taken at ten
hours post transfection. Scale bars = 10 μm.

B, Co-expression of *At*MLKL1 and cytoskeleton markers in *Arabidopsis*
mesophyll protoplasts. Co-expression of *At*MLKL1-mYFP and mcherry-MAP4
(microtubule marker: (*33*) top panels), or tagRFP-T-Lifeact (Actin marker: (*34*)
middle panels). Representative confocal images were taken at 10 hours post
transfection. Scale bars = 10 μm.

C, The filamentous structures of *At*MLKL1-mYFP were undetectable at 5
minutes after application of 5 uM of the microtubule inhibitor, oryzalin.
Representative confocal image was taken at 10 hours post transfection.
Scale bar = 10 μm.

Supplemental Fig. 12: Confocal images of an epidermal cell that was
infected by *G. orontii*. The maximum intensity projection was obtained from
confocal Z-stack images and the autofluorescence of fungal structures were
discriminated from *At*MLKL1-GFP signals by spectral imaging. Arrowheads
indicate filamentous structures associated with *At*MLKL1-GFP. C; cytoplasm,
Ha: haustorium, E, haustorial encasement. Scale bar = 5 μm.

Supplemental Fig. 13: Cortical microtubule arrays in abaxial leaf
epidermal cells of *transgenic A. thaliana lines expressing*
phosphomimetic and wild type variants of *At*MLKL1. Representative
images were obtained one day after biolistic delivery of the expression
construct for the *At*ubiqutin10 promoter driven by mCherry-TUB6 (microtubule
marker (*24*)) into the indicated transgenic lines. Scale bars = 10 μm.

Supplemental movie 1

Dynamics of *At*MLKL1(wild-type)-GFP in a non-pathogen-challenged abaxial
leaf epidermal cell. The corresponding projection image is shown in Fig. 4E.
The white circle indicates a GFP signal proximal to the PM (see Fig. 4D) and
which was less mobile compared to other signals.

Supplemental movie 2

Dynamics of *At*MLKL1(S393D)-GFP in a non-pathogen-challenged abaxial
leaf epidermal cell. The corresponding projection image is shown in Fig. 4E.

The white circle indicates a GFP signal proximal to the PM (see Fig. 4D) and
which was less mobile compared to other signals.

Supplemental movie 3

838   Dynamics of *At*MLKL1(S395D)-GFP in a non-pathogen-challenged abaxial
839   leaf epidermal cell. The corresponding projection image is shown in Fig. 4E.
840   The white circle indicates a GFP signal proximal to the PM (see Fig. 4D) and
841   which was less mobile compared to other signals.

Supplemental file 1

844   108 coding sequences (CDS) of plant *MLKLs* belong to the same orthogroups
845   (*11*). A CDS derived from *Arabidopsis halleri* is incomplete due to the
846   ambiguous assembly of the corresponding genomic region.

Table S1. Number of MLKL-like genes in plant genomes

Plant species	NCBI Taxonomy ID	Family	Number of *MLKL*-like genes
Coccomyxa subellipsoidea	248742	Coccomyxaceae	0
Ostreococcus lucimarinus	242159	Bathycoccaceae	0
Volvox carteri	3067	Volvocaceae	0
Physcomitrella patens	3218	Funariaceae	0
Selaginella moellendorffii	88036	Selaginellaceae	0
Actinidia chinensis	3625	Actinidiaceae	3
Amborella trichopoda	13333	Amborellaceae	1
Spirodela polyrhiza	29656	Araceae	2
Phoenix dactylifera	42345	Arecaceae	3
Arabidopsis halleri	81970	Brassicaceae	2
Arabidopsis lyrata	59689	Brassicaceae	3
Arabidopsis thaliana	3702	Brassicaceae	2
Brassica oleracea	3712	Brassicaceae	3
Brassica rapa	3711	Brassicaceae	2
Boechera stricta	72658	Brassicaceae	3
Capsella grandiflora	264402	Brassicaceae	3
Capsella rubella	81985	Brassicaceae	3
Eutrema salsugineum	72664	Brassicaceae	4
Schrenkiella parvula	98039	Brassicaceae	3
Beta vulgaris	161934	Chenopodiaceae	2
Cucumis sativus	3659	Cucurbitaceae	2
Ricinus communis	3988	Euphorbiaceae	2
Arachis ipaensis	130454	Fabaceae	2
Medicago truncatula	3880	Fabaceae	2
Phaseolus vulgaris	3885	Fabaceae	2
Linum usitatissimum	4006	Linaceae	4
Gossypium raimondii	29730	Malvaceae	3
Theobroma cacao	3641	Malvaceae	2
Musa acuminata	4641	Musaceae	4
Erythranthe guttata	4155	Phrymaceae	2
Pinus taeda	3352	Pinaceae	1
Brachypodium distachyon	15368	Poaceae	2
Oryza sativa	4530	Poaceae	2
Sorghum bicolor	4558	Poaceae	2
Setaria italica	4555	Poaceae	2
Zea mays	4577	Poaceae	2
Aquilegia coerulea	218851	Ranunculaceae	1
Fragaria vesca	57918	Rosaceae	2

Malus domestica	3750	Rosaceae	5
Prunus persica	3760	Rosaceae	2
Coffea canephora	49390	Rubiaceae	2
Citrus clementina	85681	Rutaceae	2
Citrus sinensis	2711	Rutaceae	2
Populus trichocarpa	3694	Salicaceae	2
Capsicum annuum	4072	Solanaceae	3
Solanum melongena	4111	Solanaceae	3
Solanum lycopersicum	4081	Solanaceae	3
Solanum tuberosum	4113	Solanaceae	3
Vitis vinifera	29760	Vitaceae	2

849 **Table S2. Cryo-EM statistics and model refinement for *At*MLKL tetramer**

	*At*MLKL3 tetramer	*At*MLKL2 tetramer
Data collection and processing		
Microscope	FEI Titan Krios	FEI Titan Krios
Detector	Gatan K2 Summit	Gatan K2 Summit
Voltage(kV)	300	300
Magnification	22500x	22500x
Pixel size (Å)	1.30654	1.30654
Total electron dose (e⁻/Å²)	50	50
Defocus range (μm)	-1.5 ~ -2.3	-1.6~-2.2
Micrographs collected	1,434	1,828
Reconstruction		
Total extraced particles	983,779	1,527,009
Number of particles used for 3D reconstruction	629,287	796,901
Number of particles used for refinement	123,472	154,008
Symmetry	D2	D2
Resolution (Å) 0.143 after refinement	3.84	4.51
Resolution (Å) 0.143 after post-processing	3.39	4.08
Map sharpening B-factor (Å²)	-152	-222.79
Refinement		
Estimated Resolution (Å)	3.4	4.1
Model composition		
Number of protein atoms	16584	14238
Number of ligand atoms	0	0
MapCC (mask/box)	0.83/0.79	0.81/0.81
Rwork/Rfree (%)	34.35/34.35	39.4/39.4
R.M.S deviations		
Bonds lengths (Å)	0.011	0.010
Bonds angles (°)	0.941	0.797
Validation		
MolProbity overall score	2.22	2.56
All-atom clashscore	15.96	18.07
Rotamer outliers (%)	0.67	1.57
C-beta deviations	0.0	0.0
EMRinger score	1.56	1.42
Ramachandran plot statistics		
Preferred (%)	91.27	85.27
Allowed (%)	8.33	14.31
Outlier (%)	0.40	0.42

Table S3 Primer pairs used in this study.

Forward primer	Sequence 5′-3′	Reverse primer	Sequence 5′-3′	Use, comments
SALK_041569c_RP	GAGAGACAGCGAAAACCTGTG	SALK_041569c_LP	CTTCTCTTGCAATGCCTCAAC	Genotyping of the wild type allele of *AtMLKL1*
SALK_041569c_RP	GAGAGACAGCGAAAACCTGTG	LBb1.3	ATTTTGCCGATTTCGGAAC	Genotyping of the mutant allele of *AtMLKL1*
SALK_124412c_RP AT5G41730	TCAGCTTAGCTTTCAACACGG	SALK_124412c_LP AT5G41730	GAAGATACTCCATCCCCCTTG	Genotyping of the wild type allele of *AtMLKL2*
SALK_124412c_RP AT5G41730	TCAGCTTAGCTTTCAACACGG	LBb1.3	ATTTTGCCGATTTCGGAAC	Genotyping of the mutant allele of *AtMLKL2*
GABI_491E02_RP	TACACGAAGCCCATTCACTTC	GABI_491E02_LP	TATTACCTCTGCGGATTCACG	Genotyping of the wild type allele of *AtMLKL3*
GABI_o3144/35St	GTGGATTGATGTGATATCTCC	GABI_491E02_LP	TATTACCTCTGCGGATTCACG	Genotyping of the mutant allele of *AtMLKL3*
AT1g64300-Fw1	CACCATGGAGCAATTCAGGCAAATCGGAGAGGTT	AT1g64300-Rv1	TGTAAGCTCAGAGTCTGAAGCAGGACCAGTTA	D-topo cloning of *At*MLKL1 cDNA without STOP codon
AT5G41730-Fw1	CACCATGGAGCAATTCAGACAAATCGGAGAAGTTCTT	AT5G41730-Rv1	AGTAAGCTCAGAGTCTGAAGCATGACCAGTCTGTT	D-topo cloning of *At*MLKL2 cDNA without STOP codon
D-topo-*At*MLKL3-f	CACCATGGATCAATTTCGAGAGATAGGAGAGGTATT	D-topo-*At*MLKL3-r_without_stop	GGAAAGCTCAGAGTCAGAGGCATGT	D-topo cloning of *At*MLKL3 cDNA without STOP codon
*At*MLKL1(S393D)-f	CCTCAGTTAAGAATCAGGATTTTTCTCGGGCTTCC	*At*MLKL1(S393D)-r	GGAAGCCCGAGAAAATCCTGATTCTTAACTGAGG	Site-directed mutagenesis to obtain *At*MLKL1(S393D) variant
*At*MLKL1(S395D)-f	GTTAAGAATCAGTCTTTTGATCGGGCTTCCTCTAGAC	*At*MLKL1(S395D)-r	GTCTAGAGGAAGCCCGATCAAAAGACTGATTCTTAAC	Site-directed mutagenesis to obtain *At*MLKL1(S395D) variant
AT5G41730_(S394D)-f	GTCGTTAAAGCGCAGGATTCCTCGAAACCAG	AT5G41730_(S394D)-r	CTGGTTTCGAGGAATCCTGCGCTTTAACGAC	Site-directed mutagenesis to obtain *At*MLKL2(S394D) variant
AT5G41730_(S395D)-f	CGTTAAAGCGCAGTCTGATTCGAAACCAGGCACTCC	AT5G41730_(S395D)-r	GGAGTGCCTGGTTTCGAATCAGACTGCGCTTAACG	Site-directed mutagenesis to obtain *At*MLKL2(S395D) variant
*At*MLKL3(S392D)-f	GTTGAATTCTGTCAAGGGATTCGATAGTAAAAGTGCTTCTTTGACG	*At*MLKL3(S392D)-r	CGTCAAAGAAGCACTTTTACTATCGAATCCCTTGACAGAATTCAAC	Site-directed mutagenesis to obtain *At*MLKL3(S392D) variant
*At*MLKL3(SS393D)-f	CTGTCAAGGGATTCTCTGATAAAAGTGCTTCTTTGAC	*At*MLKL3(S393D)-r	GTCAAAGAAGCACTTTTATCAGAGAATCCCTTGACAG	Site-directed mutagenesis to obtain *At*MLKL3(S393D) variant
Genomic of	CACCCCTTG	Genomic of	TGTAAGCT	D-topo cloning of genomic fragments

*At*3g21215-f	GAATCCACC CATACCACC AG	*At*3g21215-r	GAGGAGG AGGATGGT GATGA	qPCR quantification of the Arabidopsis DNA

Supplemental References

1. N. Kleinboelting, G. Huep, A. Kloetgen, P. Viehoever, B. Weisshaar, GABI-Kat SimpleSearch: new features of the Arabidopsis thaliana T-DNA mutant database. *Nucleic Acids Res* **40**, D1211-1215 (2012).

2. J. M. Alonso *et al.*, Genome-wide insertional mutagenesis of Arabidopsis thaliana. *Science* **301**, 653-657 (2003).

3. M. Bartsch *et al.*, Salicylic acid-independent ENHANCED DISEASE SUSCEPTIBILITY1 signaling in Arabidopsis immunity and cell death is regulated by the monooxygenase FMO1 and the Nudix hydrolase NUDT7. *Plant Cell* **18**, 1038-1051 (2006).

4. T. Nakagawa *et al.*, Improved Gateway binary vectors: high-performance vectors for creation of fusion constructs in transgenic analysis of plants. *Biosci Biotechnol Biochem* **71**, 2095-2100 (2007).

5. S. J. Clough, A. F. Bent, Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J* **16**, 735-743 (1998).

6. C. Koncz, J. Schell, The promoter of TL-DNA gene 5 controls the tissue-specific expression of chimaeric genes carried by a novel type of Agrobacterium binary vector. *Molecular and General Genetics MGG* **204**, 383-396 (1986).

7. F. Jacob *et al.*, A dominant-interfering camta3 mutation compromises primary transcriptional outputs mediated by both cell surface and intracellular immune receptors in Arabidopsis thaliana. *New Phytol* **217**, 1667-1680 (2018).

8. P. Bucher, K. Karplus, N. Moeri, K. Hofmann, A flexible motif search technique based on generalized profiles. *Comput Chem* **20**, 3-23 (1996).

9. K. Katoh, K. Misawa, K. Kuma, T. Miyata, MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059-3066 (2002).

10. J. Soding, Protein homology detection by HMM-HMM comparison. *Bioinformatics* **21**, 951-960 (2005).

11. D. Lapin *et al.*, A coevolved EDS1-SAG101-NRG1 module mediates cell death signaling by TIR-domain immune receptors. *bioRxiv*, 572826 (2019).

12. T. Maekawa *et al.*, Subfamily-Specific Specialization of RGH1/MLA Immune Receptors in Wild Barley. *Mol Plant Microbe Interact* **32**, 107-119 (2019).

13. S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).

14. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *J Struct Biol* **192**, 216-221 (2015).

15. J. Zivanov *et al.*, New tools for automated high-resolution cryo-EM structure determination in RELION-3. *Elife* **7**, (2018).

16. A. Brown *et al.*, Tools for macromolecular model building and refinement into electron cryo-microscopy reconstructions. *Acta Crystallogr D Biol Crystallogr* **71**, 136-153 (2015).

17.   P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system
for macromolecular structure solution. *Acta Crystallogr D Biol*
Crystallogr **66**, 213-221 (2010).
18.   L. Su *et al.*, A plug release mechanism for membrane permeation by
MLKL. *Structure* **22**, 1489-1500 (2014).
19.   J. M. Murphy *et al.*, The pseudokinase MLKL mediates necroptosis via
a molecular switch mechanism. *Immunity* **39**, 443-453 (2013).
20.   I. M. Saur *et al.*, Multiple pairs of allelic MLA immune receptor-powdery
mildew AVRA effectors argue for a direct recognition mechanism. *Elife*
8, (2019).
21.   T. Maekawa *et al.*, Coiled-coil domain-dependent homodimerization of
intracellular barley immune receptors defines a minimal functional
module for triggering cell death. *Cell Host Microbe* **9**, 187-199 (2011).
22.   F. Vogler, S. S. Konrad, S. Sprunck, Knockin' on pollen's door: live cell
imaging of early polarization events in germinating Arabidopsis pollen.
Front Plant Sci **6**, 246 (2015).
23.   T. Hamada *et al.*, Purification and characterization of novel
microtubule-associated proteins from Arabidopsis cell suspension
cultures. *Plant Physiol* **163**, 1804-1816 (2013).
24.   S. Fujita *et al.*, An atypical tubulin kinase mediates stress-induced
microtubule depolymerization in Arabidopsis. *Curr Biol* **23**, 1969-1978
(2013).
25.   C. Gehl *et al.*, Quantitative analysis of dynamic protein-protein
interactions in planta by a floated-leaf luciferase complementation
imaging (FLuCI) assay using binary Gateway vectors. *Plant J* **67**, 542-
553 (2011).
26.   S. D. Yoo, Y. H. Cho, J. Sheen, Arabidopsis mesophyll protoplasts: a
versatile cell system for transient gene expression analysis. *Nat Protoc*
2, 1565-1572 (2007).
27.   R. Wessling, R. Panstruga, Rapid quantification of plant-powdery
mildew interactions by qPCR and conidiospore counts. *Plant Methods*
8, 35 (2012).
28.   A. Cabral *et al.*, Identification of Hyaloperonospora arabidopsidis
transcript sequences expressed during infection reveals isolate-specific
effectors. *PLoS One* **6**, e19328 (2011).
29.   S. Bai *et al.*, Structure-function analysis of barley NLR immune
receptor MLA10 reveals its cell compartment specific activity in cell
death and disease resistance. *PLoS Pathog* **8**, e1002752 (2012).
30.   R. P. Birkenbihl, C. Diezel, I. E. Somssich, Arabidopsis WRKY33 is a
key transcriptional regulator of hormonal and metabolic responses
toward Botrytis cinerea infection. *Plant Physiol* **159**, 266-285 (2012).
31.   C. Gachon, P. Saindrenan, Real-time PCR monitoring of fungal
development in Arabidopsis thaliana infected by Alternaria brassicicola
and Botrytis cinerea. *Plant Physiol Biochem* **42**, 367-371 (2004).
32.   Q. H. Shen *et al.*, Recognition specificity and RAR1/SGT1 dependence
in barley Mla disease resistance genes to the powdery mildew fungus.
Plant Cell **15**, 732-744 (2003).
33.   J. Marc *et al.*, A GFP-MAP4 reporter gene for visualizing cortical
microtubule rearrangements in living epidermal cells. *Plant Cell* **10**,
1927-1940 (1998).

34.   J. Riedl *et al.*, Lifeact: a versatile marker to visualize F-actin. *Nat*
956   *Methods* **5**, 605-607 (2008).

Figure 1

A

B

C

HeLo domain
Brace region
Pseudokinase domain

D

*At*MLKL3
hMLKL
mMLKL

α2
α1
α4
α3

α1-α2
Loop helix

α3-α4
Loop helix

E

*At*MLKL3

hMLKL

mMLKL

F

G

Figure 2

Figure 3

Figure 4

Figure S1

Figure S2

Figure S3

A

AT1G64300   *AtMLKL1*

AT5G41730   *AtMLKL2*

No gene model avaialbe in TAIR10   *AtMLKL3*

AT1G01450

B

| Col-0 | Wild type | *Atmlkl1* | *Atmlkl2* | *Atmlkl3* |

| *Atmlkl1 2* | *Atmlkl1 3* | *Atmlkl2 3* | *Atmlkl1 2 3* |

Figure S4

Figure S5

Figure S6

A

Cell viability (% Evans blue negative cells)

- Empty vector — c
- AtMLKL1-HA — b
- AtMLKL2-HA — ab
- AtMLKL3-HA — a

B

Cell viability (% Evans blue negative cells)

- Empty vector — b
- AtMLKL1-mYFP — a
- AtMLKL2-mYFP — a
- AtMLKL3-mYFP — a
- HvMLA(1-160aa)-mYFP — a

C

Cell viablity (% Relative to empty vector control)

n. s.
- AtMLKL2 (Wild type)-HA
- AtMLKL2 (S394D)-HA
- AtMLKL2 (S395D)-HA

n. s.
- AtMLKL3 (Wild type)-HA
- AtMLKL3 (S392D)-HA
- AtMLKL3 (S393D)-HA

D

Empty vector | AtMLKL1-HA | AtMLKL2-HA | AtMLKL2-mYFP | AtMLKL3-HA

Wild type | S393D | S395D | Wild type | S394D | S395D | Wild type | S392D | S393D

kDa
100
75
50

WB: anti-HA

kDa
100
75
50

CBB staining

Figure S7

A

B

G. orontii

Col-0   *Atmlkl1*

AtMLKL1 (Wild type)-GFP
in *Atmlkl1*

AtMLKL1 (S395D)-GFP
in *Atmlkl1*

AtMLKL1 (S393D)-GFP
in *Atmlkl1*

1 cm

WB: anti-GFP

CBB staining

Figure S8

Figure S9

Figure S10

A

*At*MLKL1-mYFP   *At*MLKL2-mYFP   *At*MLKL3-mYFP

B

*At*MLKL1-mYFP   mCherry-MAP4   Overlay

*At*MLKL1-mYFP   tagRFP-T-Lifeact   Overlay

C

*At*MLKL1-mYFP
+ Oryzalin (5 min)

Figure S11

AtMLKL1(Wild type)-GFP in Atmlkl1

Maximum projection (over Z)

Figure S12

AtMLKL1(Wild type)-GFP

mCherry-TUB6
(Biolistic delivery)

AtMLKL1(S393D)-GFP

mCherry-TUB6
(Biolistic delivery)

AtMLKL1(S395D)-GFP

mCherry-TUB6
(Biolistic delivery)

Figure S13