Maturation of cortical microstructure and cognitive development in

childhood and adolescence: a T1w/T2w ratio MRI study

Linn B. Norbom[a,b,*], Jaroslav Rokicki[a,b], Dag Alnæs[a], Tobias Kaufmann[a], Nhat Trung Doan[a],

Ole A. Andreassen[a], Lars T. Westlye[a,b], Christian K. Tamnes[a,c,d], for the Pediatric Imaging,

Neurocognition and Genetics Study**

[a] NORMENT, Division of Mental Health and Addiction, Oslo University Hospital & Institute

of Clinical Medicine, University of Oslo, Norway

[b] Department of Psychology, University of Oslo, Norway

[c] PROMENTA Research Center, Department of Psychology, University of Oslo, Norway

[d] Department of Psychiatry, Diakonhjemmet Hospital, Oslo, Norway

Corresponding authors P.O. box 1094 Blindern, 0317 Oslo, Norway.

*l.c.b.norbom@psykologi.uio.no

Running title: Maturation of cortical microstructure

Keywords: adolescence; brain development; children; cognition; myelination; signal intensity

Number of words abstract: 199

Number of words introduction: 603

Number of words discussion: 1223

Number of figures: 10

Number of tables: 3

Abstract

The restructuring and optimization of the cerebral cortex from early childhood and through adolescence is an essential feature of human brain development, underlying immense cognitive improvements. Beyond established morphometric cortical assessments, the T1w/T2w ratio quantifies partly separate biological processes, and might inform models of typical neurocognitive development and developmental psychopathology. In the present study, we computed vertex-wise T1w/T2w ratio across the cortical surface in 621 youths (3-21 years) sampled from the Pediatric Imaging, Neurocognition, and Genetics (PING) study and tested for associations with individual differences in age, sex, and both general and specific cognitive abilities. The results showed a near global linear age-related increase in T1w/T2w ratio across the brain surface, with a general posterior to anterior gradient in association strength. Moreover, results indicated that boys in late adolescence had regionally higher T1w/T2w ratio as compared to girls. Across individuals, T1w/T2w ratio was negatively associated with general- , and several specific cognitive abilities mainly within anterior cortical regions. Our study indicates age-related differences in T1w/T2w ratio throughout childhood, adolescence and young adulthood, in line with the known protracted myelination of the cortex. Moreover, the study supports the relevance of the T1w/T2w ratio as a neuroimaging marker for cognitive development.

Significance Statement

Knowledge about the complex development of the cerebral cortex may inform both models concerning the qualitative and quantitative evolution of human cognitive abilities, but also the search for neurodevelopmental deviations linked to maladjustment and psychopathology. Here we show, in a large sample of youths, that the less explored neuroimaging measure of T1w/T2w ratio globally increases from early childhood until young adulthood, consistent with

58 established maturational patterns of intracortical myelination. We also show that T1w/T2w

59 ratio is a relevant marker for cognitive development, as it was inversely associated with both

60 general and several specific cognitive abilities.

Introduction

Childhood and adolescence is a time of extensive brain maturation and reorganization, shaped by multiple genetically based biological processes in dynamic interplay with the environment (Blakemore, 2012; Brown & Jernigan, 2012; Kremen et al., 2013; Lebel & Deoni, 2018). Magnetic resonance imaging (MRI) indicates that structural cortical development is characterized by widespread, but regionally variable and coordinated nonlinear decreases in volume and thickness, and early increases followed by smaller decreases in surface area (Amlien et al., 2016; Khundrakpam et al., 2019; Tamnes et al., 2017). This cortical maturation is mirrored by immense cognitive improvements from early childhood to young adulthood, with more basic functions such as sensorimotor processes improving before complex functions such as top-down control of behavior (Akshoomoff et al., 2014; Casey, Tottenham, Liston, & Durston, 2005). Detailed knowledge about cortical maturational patterns in youth is critical for models of normative neurocognitive development, and for serving as a blueprint for detecting neurodevelopmental deviations linked to maladjustment and psychopathology.

In addition to cortical morphometry, less explored features such as T1-weighted (T1w) signal intensity variation, has been shown to map partly separate neurobiological properties and processes (Andrews et al., 2017; Lewis, Evans, & Tohka, 2018; Lewis, Fonov, Collins, Evans, & Tohka, 2019; Norbom et al., 2019; Salat et al., 2009). Maturational and aging related cortical thinning, for instance, have been teased apart by their inverse relationships with cortical T1w intensity, which increases during development and declines in old age (Westlye et al., 2010). More recently, T1w/T2w ratio has been suggested as a useful and possibly more specific measure of cortical microstructure (M. F. Glasser & Van Essen, 2011; Van Essen & Glasser, 2014). A handful of studies indicate the utility of T1w/T2w ratio for mapping global and regional maturational and aging related patterns (H. Grydeland et al., 2019; Håkon Grydeland, Walhovd, Tamnes, Westlye, & Fjell, 2013; Shafee, Buckner, &

Fischl, 2015), and associations with cognitive performance in youth (Håkon Grydeland et al.,

2013; Vandewouw, Young, Shroff, Taylor, & Sled, 2019) and adults (H. Grydeland, Westlye,

Walhovd, & Fjell, 2016).

Although the underlying biology of the T1w/T2w ratio is controversial (Hagiwara et

al., 2018; Ritchie, Pantazatos, & French, 2018; Uddin, Figley, Solar, Shatil, & Figley, 2019)

and likely complex, it has been proposed that it relates to differential myelination of the

cerebral cortex (Ganzetti, Wenderoth, & Mantini, 2014; M. F. Glasser & Van Essen, 2011;

Nakamura, Chen, Ontaneda, Fox, & Trapp, 2017). Intracortical myelination is a crucial

feature of postnatal brain development, allowing for efficient signal transmission and

structural support (Baumann & Pham-Dinh, 2001; Liu, Li, Zhu, Li, & Liu, 2019; Waxman &

Bennett, 1972), and generally follows a posterior to anterior maturational pattern (Yakovlev

& Lecours, 1967), with additional regional specificity (Deoni, Dean, Remer, Dirks, &

O'Muircheartaigh, 2015; Nieuwenhuys, 2013; Ziegler et al., 2019). Cortical dysmyelination

has also been linked to neurodevelopmental psychopathology (Bartzokis, 2005; Insel, 2010).

In the present study, we computed vertex-wise T1w/T2w ratio across the cortical

surface in 621 youths (3-21 years) from the Pediatric Imaging, Neurocognition, and Genetics

(PING) sample, and tested for associations between T1w/T2w ratio and individual differences

in age, sex and both general and specific cognitive abilities. We hypothesized that the

T1w/T2w ratio would generally show a positive age association, putatively reflecting

protracted intracortical myelination, but with regional variation. Next, given previous findings

of limited sex differences in morphometric cortical development (Vijayakumar et al., 2016;

Wierenga, Bos, van Rossenberg, & Crone, 2019), we hypothesized that boys and girls would

show similar age related associations with T1w/T2w ratio. Finally, we tested for associations

between T1w/T2w ratio and general and specific cognitive abilities, tentatively hypothesizing

132 that there would be a positive association, based on the direction of reported developmental

133 T1w/T2w findings.

135 **Materials and Methods**

136 *Participants*

137 The publicly available PING study database (http://ping.chd.ucsd.edu) consists of data from a

138 large sample of typically developing children and adolescents aged 3-21 years, with

139 standardized behavioral measures, whole genome genotyping, and multimodal imaging for a

140 large subgroup (Jernigan et al., 2016). Fluent English speakers within the desired age range

141 were recruited across the US through local postings and outreach facilities. Exclusion criteria

142 included somatic illness, history of head trauma, contraindications for MRI, preterm birth,

143 diagnosis of mental disorders (except ADHD), mental retardation, and severe illicit drug use

144 by mother during pregnancy. Subjects 18 years and above provided written informed consent,

145 while written parental informed consent was obtained for all other subjects, in addition to

146 child assent for subjects aged 7-17 (Jernigan et al., 2016).

147  From 998 subjects with available MRI and demographic data, T2-weighted (T2w) data

148 was missing or acquired at insufficient resolution (> 1.2mm voxel size in any direction) for

149 322 subjects. Of the remainder, we additionally excluded 55 subjects after stringent quality

150 control (see below), yielding a final sample of 621 subjects (303 girls) aged 3.2-21.0 years

151 (mean = 12.4, SD = 4.8). Ten of these subjects had missing cognitive data and were excluded

152 from relevant analyses. See Table 1 for sample demographics.

154 *MRI acquisition and processing*

155 Imaging data for our final sample was attained on 7 separate 3T scanners; 2 from GE medical

156 systems (Signa HDx and Discovery MR750), and 5 from Siemens (TrioTim). The age-ranges

were largely overlapping across scanners (Figure 1). To account for challenges associated

with multisite imaging of children, an optimized imaging protocol was implemented across

vendors and models to obtain similar contrast properties and image-derived quantitative

measures (Jernigan et al., 2016). The protocol included a 3D T1w inversion prepared RF-

spoiled gradient echo scan , and a 3D T2w variable flip angle fast spin echo scan (Jernigan et

al., 2016), both acquired with prospective motion correction (PROMO) (White et al., 2010)

and voxel sizes ranging from $\approx$ 0.9-1.2mm^3.

Imaging processing was performed using the three step Human Connectome Project

(HCP) pipeline (Matthew F. Glasser et al., 2013). In short, the "PreFreeSurfer" stage produces

an undistorted native structural volume space, aligns the T1w and T2w image, performs bias

field correction, and registers the native- to MNI space. Based on FreeSurfer version 6, the

"FreeSurfer" stage performs volume segmentation, white and pial surface reconstruction, and

standard surface registration to fsaverage. Due to some cases exhibiting large cortical

thickness and T1w/T2w ratio discrepancies between the left and right hemisphere following

the default pipeline, we removed the t2pial flag within autorecon3. The "PostFreeSurfer"

stage produces NIFTI volume- and GIFTI surface files, registers surfaces to the HCP

standard_fs_LR mesh", and creates the final brain masks and T1w/T2w ratio maps (Matthew

F. Glasser et al., 2013).

T1w/T2w ratio maps

The creation of T1w/T2w ratio maps was based on methods described by M. F. Glasser and

Van Essen (2011). A volume-to-surface mapping algorithm was applied to all voxel centers

located within the cortical ribbon to achieve image division. For each vertex within the HCP

pipeline surface "midthickness", ribbon voxels were selected within a cylinder centered on the

vertex, orthogonal to the surface, and with a height and radius equal to the local cortical

thickness. To remove voxels with significant blood vessel- and CSF partial voluming effects,

voxels were excluded if the T1w/T2w value exceeds ±1 SD of all values within the cortical

ribbon. The remaining ribbon voxels were then averaged following a Gaussian weighted

function to produce a single value for each vertex, while the medial wall was assigned values

of zero (M. F. Glasser & Van Essen, 2011). The final output includes standard and bias

corrected maps. As uncorrected maps are recommended for quantitative analyses, smoothed

non-bias corrected maps were used for all analyses.

Quality assessment of imaging data

We performed extensive quality assessment as multisite projects inevitably creates

inconsistencies and since in-scanner head motion is more frequent in younger subjects. For

subjects with both T1w and T2w data of sufficient resolution, all NIFTI images were

processed through the quality assessment pipeline MRIQC (Esteban et al., 2017). After visual

inspection, flagged subjects were either included, tagged for detailed re-assessment after

T1w/T2w ratio map creation, or excluded due to poor image quality when there was no option

to replace the image with another satisfactory run. The remaining subjects were subsequently

processed through the HCP pipeline (Matthew F. Glasser et al., 2013). Finally, we visually

inspected all T1w/T2w ratio maps and excluded subjects with poor maps.

Cognitive assessment

Subjects completed the computerized NIH Toolbox Cognition Battery (Akshoomoff et al.,

2014; Weintraub, Bauer, et al., 2013; Weintraub, Dikmen, et al., 2013) which includes seven

different tasks, designed to measure eight diverse cognitive abilities (Table 2). Test order was

standardized across participants, and eight sum scores were calculated as described elsewhere

(Akshoomoff et al., 2014). Due to missing cognitive data on specific tests, we imputed 78

sum score values in total, from 63 subjects in R, using the "mice" package (Buuren &

Groothuis-Oudshoorn, 2011), with the default of 5 multiple imputations, and choosing the

first imputation. As expected, there were high correlations between all sum scores and age

(Pearson's r ranging from 0.63 to 0.82), and we therefore residualized sum scores by linear

and quadratic age by linear regression. We then performed a principal component analysis

(PCA) in R (https://www.r-project.org/) and the first factor, explaining 33.7% of the variance,

was extracted and used as a measure of general cognitive ability. We changed the sign of the

component so that higher factor loading would represent better performance. See Figure 2 for

a correlation matrix between age, sex, general cognitive ability and individual cognitive sum

scores.

Statistical analyses

We used general linear models (GLM), as implemented in the Permutation Analysis of Linear

Models (PALM) toolbox (Winkler, Ridgway, Webster, Smith, & Nichols, 2014), to test for

linear and quadratic (defined as polynomial of degree 2) effects of age on vertex-wise

T1w/T2w ratio, covarying for sex and scanner. Second, we tested for main effects of sex on

vertex-wise T1w/T2w ratio, covarying for linear and quadratic age and scanner, and then for

interaction effects between sex and age, with main effect of sex as an additional covariate.

Third, we performed vertex-wise analyses to assess the effects of general cognitive ability, as

well as each of the eight specific sum scores, on vertex-wise T1w/T2w ratio, with sex, linear

and quadratic age, sex*age and scanner as co-variates. Finally, we examined interaction

effects between age and general cognitive ability on T1w/T2w ratio, with sex, linear and

quadratic age, sex*age, general cognitive ability and scanner as co-variates. To assess

statistical significance we used 10,000 permutations and family wise error (FWE) correction

with threshold-free cluster enhancement (TFCE) (Smith & Nichols, 2009) corrected across

each contrast (as PALM performs one tailed analyses) and a significance threshold of $p<0.05$,

after correcting for both hemispheres.

To regionally summarize relevant vertex-wise results, we averaged our t-statistic

surface maps within 34 regions of interest (ROIs) for both the left and right hemisphere. ROIs

were defined by Freesurfer's fsaverage Desikan-Killiany Atlas parcellation (Desikan et al.,

2006; Fischl et al., 2004) which was converted to GIFTI format and mapped to the HCP

"fs_LR mesh". To further visualize relevant PALM results we used individual parcellations

and locally weighted scatterplot smoothing (loess) and ggplot2 (Wickham, 2009) in R

(https://www.r-project.org/).

Results

T1w/T2w mean ratio map

Figure 3 shows the mean T1w/T2w ratio map across all subjects. Briefly, the anatomical

signal distribution corresponded well with previous T1w/T2w maps (M. F. Glasser & Van

Essen, 2011), showing a specific regional pattern in which sensorimotor regions, such as pre-

and post-central gyri and visual cortex, showed relatively higher T1w/T2w ratio, while the

frontal lobe and temporal pole showed relatively lower T1w/T2w ratio.

Associations between T1w/T2w ratio and age

Permutation testing revealed a near global positive association between linear age and

T1w/T2w ratio, indicating higher ratio in widespread regions with higher age when

controlling for sex (Figure 4). There was a general posterior to anterior gradient in association

strength, with highest mean t-statistics found in precentral, caudal middle frontal, pars

triangularis, and postcentral regions within the left hemisphere, and rostal middle frontal,

caudal middle frontal, precentral, and paracentral regions within the right hemisphere (Table

3). Sub-regions within the medial occipital lobe and minor medial parietal and frontal regions

did not show significant age associations.

There were no statistically significant associations between T1w/T2w ratio and

quadratic age, controlling for linear age and sex. Corresponding with our vertex-wise results,

ROI-based plots with non-parametric loess curves showed steady increases of T1w/T2w ratio

with higher age (Figure 5).

Associations between T1w/T2w ratio and sex

Permutation testing revealed no statistically significant main effect of sex on vertex-wise

T1w/T2w ratio. There were, however, significant positive interaction effects between sex and

age (Figure 6). Effects were more widespread in the left compared to the right hemisphere

with highest mean t-statistics found in pars orbitalis, rostral middle frontal, pars triangularis

and frontal pole in the left hemisphere, and frontal pole, superior frontal, rostral middle frontal

and rostral anterior cingulate in the right hemisphere (Table 3). Corresponding with our

vertex-wise results, frontal ROI-based plots showed a pattern where boys within the age-range

of approximately 15-21 years had relatively higher T1w/T2w ratio than girls (Figure 7).

Associations between T1w/T2w ratio and cognition

Permutation testing revealed a negative relationship between T1w/T2w ratio and general

cognitive ability in the frontal lobe extending into left temporal pole and right parietal regions,

with more widespread associations in left compared to the right hemisphere (Figure 8). This

indicates that lower T1w/T2w ratio was associated with better cognitive performance.

Strongest negative associations were found in pars orbitalis, pars triangularis, temporal pole

and caudal middle frontal regions in the left hemisphere, and pars orbitalis, inferior parietal,

precentral and supra marginal regions in the right hemisphere (Table 3).

Permutation testing revealed no significant interaction effects between general

cognitive ability and age on T1w/T2w ratio. Corresponding with our vertex-wise results, ROI-

based plots indicated that higher general cognitive ability was associated with regionally

lower T1w/T2w ratio across the age-range (Figure 9). Moreover, 5 specific cognitive sum

scores yielded regional negative associations with T1wT2w ratio, with varying cluster

extension after correction across hemispheres, largely overlapping with the regions reported

for general cognitive ability (Figure 10). Inhibition and attention showed small clusters within

the left hemisphere, with inhibition covering precentral and postcentral sub regions, and

attention covering the postcentral region only. While vocabulary only showed associations in

small frontal clusters within the left hemisphere, the vertex-wise pattern for reading was

widespread and highly similar to the pattern found for general cognitive ability, with frontal

associations stronger in the left as compared to the right hemisphere. Finally, working

memory showed regional associations within right hemisphere only, mainly in parietal

regions, covering highly similar areas as general cognitive ability. There were no significant

associations between T1w/T2w ratio and cognitive flexibility, episodic memory and

processing speed.

Discussion

The cerebral cortex is the center for most of our evolutionarily unique cognitive abilities

(Geschwind & Rakic, 2013), and its maturation is therefore an essential part of human brain

development, underpinning major environmental adaptations and vast cognitive

improvements (Casey et al., 2005). In a large sample of youths aged 3-21 years we found a

global age-related increase of T1w/T2w ratio across the brain surface, with some regional and

sex related variation. Intriguingly, across individuals, T1w/T2w ratio was inversely associated

with general and several specific cognitive abilities, mainly in anterior cortical regions.

The first goal of the present study was to investigate associations between T1w/T2w ratio and age. As hypothesized, we found a near global positive association, with a continuous age-related increase of T1w/T2w ratio from early childhood and into young adulthood. Our results converge with the few prior studies on T1w/T2w ratio in children, adolescents and young adults (Håkon Grydeland et al., 2013; Shafee et al., 2015). Grydeland et al. (2013) reported that for their youngest age group, consisting of 8 to 19 year old subjects, there was a positive linear association between T1w/T2w ratio and age across large portions of the cerebral cortex, with strongest association in posterior frontal, parietal, and temporal cortices. We too found our strongest associations in posterior frontal regions extending into parietal regions. In contrast to the present results, Grydeland et al. did not find age-associations within parts of anterior frontal and insular cortices, but this discrepancy could be due to greater sample size and the inclusion of younger children in the current study.

Our analyses revealed no significant main effects of sex on T1w/T2w ratio, but a positive age by sex interaction indicated that boys in late adolescence had relatively higher T1w/T2w ratio than girls in the frontal lobe and parts of parietal lobe. No previous studies have to our knowledge specifically investigated the relationship between T1w/T2w ratio and sex, but several studies have tested for sex differences in cortical morphometry and morphometric development. Although there are inconsistencies in this literature, recent longitudinal results indicate larger cortical volumes and areas in boys, but few differences in cortical thickness, and limited sex differences in the estimated rates of cortical morphometric development (Vijayakumar et al., 2016; Wierenga et al., 2019). A recent study investigating a longitudinal sample of 14 to 24 year old youths, using magnetization transfer (MT), an imaging measure linked to myelination, reported main effects of sex, with females having regionally higher MT than males (Ziegler et al., 2019). Less optimal cross sectional analyses of the same data, yielded a similar sex-age interactional pattern to the one reported in the

current study, within right angular gyrus. Therefore, future longitudinal studies on T1w/T2w

ratio are needed for more direct investigations.

Few previous studies have explored the behavioral and cognitive relevance of

T1w/T2w ratio during development. We found a negative association between T1w/T2w ratio

and general cognitive ability, mainly in anterior regions. We also found corresponding

associations regarding both direction and cumulative regional extension between T1w/T2w

ratio and the specific cognitive abilities: inhibition, attention, reading, vocabulary and

working memory. We did not find an interaction between cognitive ability and age on

T1w/T2w ratio. Intriguingly, this indicates that beyond developmental improvements,

children and adolescents with high general- and specific cognitive performance have

regionally lower T1w/T2w ratio especially within the frontal lobe. Considering the positive

T1w/T2w ratio - age association, this result is surprising, but indeed partly converges with the

very limited prior literature on T1w/T2w ratio and cognition in youth. Grydeland et al. (2013)

investigated the relationship between T1w/T2w ratio and intra-individual performance

variability in response time in both youths and adults. Although the general conclusion, based

on the combined lifespan sample, was that higher T1w/T2w ratio was associated with greater

performance stability, the opposite relationship was reported for the young subgroup of 8 to

19 year old subjects in clusters within occipital and parietal regions. However, a recent study

by Vandewouw et al. (2019) found lower T1w/T2w ratio within occipital, parietal and

subcortical regions in a small group of four year old children born very preterm compared to

full term children. Across the two groups, they also found positive associations between

T1w/T2w ratio in parietal lobes and intelligence, and expressive and receptive language

ability.

The biological underpinnings of the T1w/T2w ratio is debated and likely complex. On

the one hand, it has been shown that intracortical myelin content inversely co-varies with both

the T1w and T2w MRI signal intensity (M. F. Glasser & Van Essen, 2011), through

differences in lipids (Koenig, 1991), water concentration (Miot-Noirault, Barantin, Akoka, &

Le Pape, 1997), and iron which in the cortex is strongly co-located with myelin (Fukunaga et

al., 2010). The T1w- to T2w division also mathematically cancels scanner related intensity

biases, while increasing contrast (M. F. Glasser & Van Essen, 2011), and T1w/T2w ratio

concurs with post mortem data (Nakamura et al., 2017). On the other hand, recent attempts

have not revealed close links between T1w/T2w ratio and either myelin-related genes or more

established myelin measures (Hagiwara et al., 2018; Ritchie et al., 2018; Uddin et al., 2019),

demonstrating that it is not a straightforward myelin proxy. Nevertheless, our results indicate

that T1w/T2w ratio continuously increases from early childhood, through adolescence, and

into young adulthood, with a posterior to anterior gradient in association strength. This is in

line with the known protracted maturational timing and directional pattern of intracortical

myelination (Miller et al., 2012; Yakovlev & Lecours, 1967). The observed negative

associations between the T1w/T2w ratio and cognition are less intuitive if interpreted through

a cortical myelin perspective. One could speculate that having excess levels of cortical myelin

beyond a certain normative developmental range, is disadvantageous as myelin contains

factors associated with inhibition of neurite growth. Excess myelin might consequently foster

a permanent inhibitory environment for synapse formation and lower neuronal plasticity

(Snaidero & Simons, 2017).

We did not find associations between T1w/T2w ratio and age in sub regions of medial

occipital lobe and minor medial parietal and frontal regions, which was unexpected. This

could be explained by our removal of the t2pial flag during the HCP pipeline, which when

employed additionally uses the T2 image for pial surface estimation. As described, this

removal was done to correct for sizable hemispheric differences induced in both thickness and

T1w/T2w ratio values, possibly due to sequence or resolution differences as compared to the

HCP standard. After careful visual inspection for the T1w and T2w image separately, we

suspect that pial estimation within these regions was hampered by inclusion of dura in the

T1w image alone, resulting in underestimation of T1w/T2w ratio values. Other study

limitations include the cross-sectional design, which is a sub-optimal and indirect way to

investigate maturation and development. Moreover, imaging data was attained on several

different scanners, and although statistically controlled for in all analyses, this will still most

likely confound variables of interest.

To conclude, our results indicated that cortical T1w/T2w ratio almost globally

increases from early childhood until young adulthood, corresponding with established

maturational patterns of intracortical myelination. We also found that T1w/T2w ratio was

inversely associated with measures of both general and several specific cognitive abilities,

especially in anterior cortical regions. Future studies, preferably longitudinal, are needed to

validate the robustness of these developmental patterns and links with individual cognitive

differences and developmental psychopathology.

Acknowledgments and disclosures

This work was supported by the Department of Psychology, University of Oslo (LBN and

CKT), the Research Council of Norway (223273, 249795, 230345), the South-Eastern

Norway Regional Health Authority (2014097, 2016083), and the European Commission's 7th

Framework Programme (602450, IMAGEMEND). Data collection and sharing was funded by

the Pediatric Imaging, Neurocognition and Genetics Study (PING) (National Institutes of

Health Grant RC2DA029475). PING is funded by the National Institute on Drug Abuse and

the Eunice Kennedy Shriver National Institute of Child Health & Human Development. PING

data are disseminated by the PING Coordinating Center at the Center for Human

Development, University of California, San Diego. Data were also in part provided by the

407 Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van

408 Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that

409 support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for

410 Systems Neuroscience at Washington University. A preprint is published on bioRxiv.org. The

411 authors report no biomedical financial interests or potential conflicts of interest.

References

Akshoomoff, N., Newman, E., Thompson, W. K., McCabe, C., Bloss, C. S., Chang, L., . . .

Jernigan, T. L. (2014). The NIH Toolbox Cognition Battery: results from a large

normative developmental sample (PING). *Neuropsychology, 28*(1), 1-10.

doi:10.1037/neu0000001

Amlien, I. K., Fjell, A. M., Tamnes, C. K., Grydeland, H., Krogsrud, S. K., Chaplin, T. A., . .

. Walhovd, K. B. (2016). Organizing Principles of Human Cortical Development--

Thickness and Area from 4 to 30 Years: Insights from Comparative Primate

Neuroanatomy. *Cereb Cortex, 26*(1), 257-267. doi:10.1093/cercor/bhu214

Andrews, D. S., Avino, T. A., Gudbrandsen, M., Daly, E., Marquand, A., Murphy, C. M., . . .

Ecker, C. (2017). In Vivo Evidence of Reduced Integrity of the Gray-White Matter

Boundary in Autism Spectrum Disorder. *Cereb Cortex, 27*(2), 877-887.

doi:10.1093/cercor/bhw404

Bartzokis, G. (2005). Brain Myelination in Prevalent Neuropsychiatric Developmental

Disorders: Primary and Comorbid Addiction. *Adolesc Psychiatry, 29*, 55-96.

Baumann, N., & Pham-Dinh, D. (2001). Biology of oligodendrocyte and myelin in the

mammalian central nervous system. *Physiol Rev, 81*(2), 871-927.

doi:10.1152/physrev.2001.81.2.871

Blakemore, S. J. (2012). Imaging brain development: the adolescent brain. *NeuroImage,*

61(2), 397-406. doi:10.1016/j.neuroimage.2011.11.080

Brown, T. T., & Jernigan, T. L. (2012). Brain development during the preschool years.

Neuropsychol Rev, 22(4), 313-333. doi:10.1007/s11065-012-9214-1

Buuren, S. v., & Groothuis-Oudshoorn, K. (2011). mice: Multivariate imputation by chained

equations in R. *Journal of Statistical Software, 45*(3), 1-67.

doi:http://dx.doi.org/10.18637/jss.v045.i03

Casey, B. J., Tottenham, N., Liston, C., & Durston, S. (2005). Imaging the developing brain:

what have we learned about cognitive development? *Trends Cogn Sci, 9*(3), 104-110.

doi:10.1016/j.tics.2005.01.011

Deoni, S. C. L., Dean, D. C., 3rd, Remer, J., Dirks, H., & O'Muircheartaigh, J. (2015).

Cortical maturation and myelination in healthy toddlers and young children.

NeuroImage, 115, 147-161. doi:10.1016/j.neuroimage.2015.04.058

Desikan, R. S., Segonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., . . .

Killiany, R. J. (2006). An automated labeling system for subdividing the human

cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage, 31*(3),

968-980. doi:10.1016/j.neuroimage.2006.01.021

Esteban, O., Birman, D., Schaer, M., Koyejo, O. O., Poldrack, R. A., & Gorgolewski, K. J.

(2017). MRIQC: Advancing the automatic prediction of image quality in MRI from

unseen sites. *PLoS One, 12*(9), e0184661. doi:10.1371/journal.pone.0184661

Fischl, B., van der Kouwe, A., Destrieux, C., Halgren, E., Segonne, F., Salat, D. H., . . . Dale,

A. M. (2004). Automatically parcellating the human cerebral cortex. *Cereb Cortex,*

14(1), 11-22.

Fukunaga, M., Li, T. Q., van Gelderen, P., de Zwart, J. A., Shmueli, K., Yao, B., . . . Duyn, J.

H. (2010). Layer-specific variation of iron content in cerebral cortex as a source of

MRI contrast. *Proc Natl Acad Sci U S A, 107*(8), 3834-3839.

doi:10.1073/pnas.0911177107

Ganzetti, M., Wenderoth, N., & Mantini, D. (2014). Whole brain myelin mapping using T1-

and T2-weighted MR imaging data. *Front Hum Neurosci, 8*, 671.

doi:10.3389/fnhum.2014.00671

Geschwind, D. H., & Rakic, P. (2013). Cortical evolution: judge the brain by its cover.

Neuron, 80(3), 633-647. doi:10.1016/j.neuron.2013.10.045

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L.,

. . . Jenkinson, M. (2013). The minimal preprocessing pipelines for the Human

Connectome Project. *NeuroImage, 80*, 105-124.

doi:https://doi.org/10.1016/j.neuroimage.2013.04.127

Glasser, M. F., & Van Essen, D. C. (2011). Mapping human cortical areas in vivo based on

myelin content as revealed by T1- and T2-weighted MRI. *J Neurosci, 31*(32), 11597-

11616. doi:10.1523/jneurosci.2180-11.2011

Grydeland, H., Vertes, P. E., Vasa, F., Romero-Garcia, R., Whitaker, K., Alexander-Bloch, A.

F., . . . Bullmore, E. T. (2019). Waves of Maturation and Senescence in Micro-

structural MRI Markers of Human Cortical Myelination over the Lifespan. *Cereb

Cortex, 29*(3), 1369-1381. doi:10.1093/cercor/bhy330

Grydeland, H., Walhovd, K. B., Tamnes, C. K., Westlye, L. T., & Fjell, A. M. (2013).

Intracortical Myelin Links with Performance Variability across the Human Lifespan:

Results from T1- and T2-Weighted MRI Myelin Mapping and Diffusion Tensor

Imaging. *33*(47), 18618-18630. doi:10.1523/JNEUROSCI.2811-13.2013 %J The

Journal of Neuroscience

Grydeland, H., Westlye, L. T., Walhovd, K. B., & Fjell, A. M. (2016). Intracortical Posterior

Cingulate Myelin Content Relates to Error Processing: Results from T1- and T2-

Weighted MRI Myelin Mapping and Electrophysiology in Healthy Adults. *Cereb

Cortex, 26*(6), 2402-2410. doi:10.1093/cercor/bhv065

Hagiwara, A., Hori, M., Kamagata, K., Warntjes, M., Matsuyoshi, D., Nakazawa, M., . . .

Aoki, S. (2018). Myelin Measurement: Comparison Between Simultaneous Tissue

Relaxometry, Magnetization Transfer Saturation Index, and T1w/T2w Ratio Methods.

Scientific Reports, 8(1), 10554. doi:10.1038/s41598-018-28852-6

Insel, T. R. (2010). Rethinking schizophrenia. *Nature, 468*(7321), 187-193.

doi:10.1038/nature09552

Jernigan, T. L., Brown, T. T., Hagler, D. J., Akshoomoff, N., Bartsch, H., Newman, E., . . .

Dale, A. M. (2016). The Pediatric Imaging, Neurocognition, and Genetics (PING)

Data Repository. *NeuroImage, 124*, 1149-1154.

doi:https://doi.org/10.1016/j.neuroimage.2015.04.057

Khundrakpam, B. S., Lewis, J. D., Jeon, S., Kostopoulos, P., Itturia Medina, Y., Chouinard-

Decorte, F., & Evans, A. C. (2019). Exploring Individual Brain Variability during

Development based on Patterns of Maturational Coupling of Cortical Thickness: A

Longitudinal MRI Study. *Cereb Cortex, 29*(1), 178-188. doi:10.1093/cercor/bhx317

Kremen, W. S., Fennema-Notestine, C., Eyler, L. T., Panizzon, M. S., Chen, C. H., Franz, C.

E., . . . Dale, A. M. (2013). Genetics of brain structure: contributions from the

Vietnam Era Twin Study of Aging. *Am J Med Genet B Neuropsychiatr Genet,*

162b(7), 751-761. doi:10.1002/ajmg.b.32162

Lebel, C., & Deoni, S. (2018). The development of brain white matter microstructure.

NeuroImage, 182, 207-218. doi:10.1016/j.neuroimage.2017.12.097

Lewis, J. D., Evans, A. C., & Tohka, J. (2018). T1 white/gray contrast as a predictor of

chronological age, and an index of cognitive performance. *NeuroImage, 173*, 341-350.

doi:10.1016/j.neuroimage.2018.02.050

Lewis, J. D., Fonov, V. S., Collins, D. L., Evans, A. C., & Tohka, J. (2019). Cortical and

subcortical T1 white/gray contrast, chronological age, and cognitive performance.

NeuroImage, 196, 276-288. doi:10.1016/j.neuroimage.2019.04.022

Liu, S., Li, A., Zhu, M., Li, J., & Liu, B. (2019). Genetic influences on cortical myelination in

the human brain. *Genes Brain Behav, 18*(4), e12537. doi:10.1111/gbb.12537

Miller, D. J., Duka, T., Stimpson, C. D., Schapiro, S. J., Baze, W. B., McArthur, M. J., . . .

Sherwood, C. C. (2012). Prolonged myelination in human neocortical evolution. *Proc*

Natl Acad Sci U S A, 109(41), 16480-16485. doi:10.1073/pnas.1117943109

Miot-Noirault, E., Barantin, L., Akoka, S., & Le Pape, A. (1997). T2 relaxation time as a

marker of brain myelination: experimental MR study in two neonatal animal models. *J*

Neurosci Methods, 72(1), 5-14.

Nakamura, K., Chen, J. T., Ontaneda, D., Fox, R. J., & Trapp, B. D. (2017). T1-/T2-weighted

ratio differs in demyelinated cortex in multiple sclerosis. *Ann Neurol, 82*(4), 635-639.

doi:10.1002/ana.25019

Nieuwenhuys, R. (2013). The myeloarchitectonic studies on the human cerebral cortex of the

Vogt-Vogt school, and their significance for the interpretation of functional

neuroimaging data. *Brain Struct Funct, 218*(2), 303-352. doi:10.1007/s00429-012-

0460-z

Norbom, L. B., Doan, N. T., Alnaes, D., Kaufmann, T., Moberget, T., Rokicki, J., . . .

Tamnes, C. K. (2019). Probing Brain Developmental Patterns of Myelination and

Associations With Psychopathology in Youths Using Gray/White Matter Contrast.

Biol Psychiatry, 85(5), 389-398. doi:10.1016/j.biopsych.2018.09.027

Ritchie, J., Pantazatos, S. P., & French, L. (2018). Transcriptomic characterization of MRI

contrast with focus on the T1-w/T2-w ratio in the cerebral cortex. *NeuroImage, 174*,

504-517. doi:10.1016/j.neuroimage.2018.03.027

Salat, D. H., Lee, S. Y., van der Kouwe, A. J., Greve, D. N., Fischl, B., & Rosas, H. D.

(2009). Age-associated alterations in cortical gray and white matter signal intensity

and gray to white matter contrast. *NeuroImage, 48*(1), 21-28.

doi:10.1016/j.neuroimage.2009.06.074

Shafee, R., Buckner, R. L., & Fischl, B. (2015). Gray matter myelination of 1555 human

brains using partial volume corrected MRI images. *NeuroImage, 105*, 473-485.

doi:10.1016/j.neuroimage.2014.10.054

Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: addressing

problems of smoothing, threshold dependence and localisation in cluster inference.

NeuroImage, 44(1), 83-98. doi:10.1016/j.neuroimage.2008.03.061

Snaidero, N., & Simons, M. (2017). The logistics of myelin biogenesis in the central nervous

system. *Glia, 65*(7), 1021-1031. doi:10.1002/glia.23116

Tamnes, C. K., Herting, M. M., Goddings, A. L., Meuwese, R., Blakemore, S. J., Dahl, R. E.,

. . . Mills, K. L. (2017). Development of the Cerebral Cortex across Adolescence: A

Multisample Study of Inter-Related Longitudinal Changes in Cortical Volume,

Surface Area, and Thickness. *J Neurosci, 37*(12), 3402-3412.

doi:10.1523/jneurosci.3302-16.2017

Uddin, M. N., Figley, T. D., Solar, K. G., Shatil, A. S., & Figley, C. R. (2019). Comparisons

between multi-component myelin water fraction, T1w/T2w ratio, and diffusion tensor

imaging measures in healthy human brain structures. *Scientific Reports, 9*(1), 2500.

doi:10.1038/s41598-019-39199-x

Van Essen, D. C., & Glasser, M. F. (2014). In vivo architectonics: a cortico-centric

perspective. *NeuroImage, 93 Pt 2*, 157-164. doi:10.1016/j.neuroimage.2013.04.095

Vandewouw, M. M., Young, J. M., Shroff, M. M., Taylor, M. J., & Sled, J. G. (2019). Altered

myelin maturation in four year old children born very preterm. *Neuroimage Clin, 21*,

101635. doi:10.1016/j.nicl.2018.101635

Vijayakumar, N., Allen, N. B., Youssef, G., Dennison, M., Yucel, M., Simmons, J. G., &

Whittle, S. (2016). Brain development during adolescence: A mixed-longitudinal

investigation of cortical thickness, surface area, and volume. *Hum Brain Mapp, 37*(6), 2027-2038. doi:10.1002/hbm.23154

Waxman, S. G., & Bennett, M. V. (1972). Relative conduction velocities of small myelinated and non-myelinated fibres in the central nervous system. *Nat New Biol, 238*(85), 217-219.

Weintraub, S., Bauer, P. J., Zelazo, P. D., Wallner-Allen, K., Dikmen, S. S., Heaton, R. K., . . . Gershon, R. C. (2013). I. NIH Toolbox Cognition Battery (CB): introduction and pediatric data. *Monogr Soc Res Child Dev, 78*(4), 1-15. doi:10.1111/mono.12031

Weintraub, S., Dikmen, S. S., Heaton, R. K., Tulsky, D. S., Zelazo, P. D., Bauer, P. J., . . . Gershon, R. C. (2013). Cognition assessment using the NIH Toolbox. *Neurology, 80*(11 Suppl 3), S54-64. doi:10.1212/WNL.0b013e3182872ded

Westlye, L. T., Walhovd, K. B., Dale, A. M., Bjornerud, A., Due-Tonnessen, P., Engvig, A., . . . Fjell, A. M. (2010). Differentiating maturational and aging-related changes of the cerebral cortex by use of thickness and signal intensity. *NeuroImage, 52*(1), 172-185. doi:10.1016/j.neuroimage.2010.03.056

White, N., Roddey, C., Shankaranarayanan, A., Han, E., Rettmann, D., Santos, J., . . . Dale, A. (2010). PROMO: Real-time prospective motion correction in MRI using image-based tracking. *Magn Reson Med, 63*(1), 91-105. doi:10.1002/mrm.22176

Wickham, H. (2009). *ggplot2 Elegant Graphics for Data Analysis* (1 ed.): Springer-Verlag New York.

Wierenga, L. M., Bos, M. G. N., van Rossenberg, F., & Crone, E. A. (2019). Sex Effects on Development of Brain Structure and Executive Functions: Greater Variance than Mean Effects. *J Cogn Neurosci, 31*(5), 730-753. doi:10.1162/jocn_a_01375

Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M., & Nichols, T. E. (2014).

Permutation inference for the general linear model. *NeuroImage, 92*, 381-397.

doi:10.1016/j.neuroimage.2014.01.060

Yakovlev, P. A., & Lecours, I. R. (1967). The myelogenetic cycles of regional maturation of

the brain. In A. Minkowski (Ed.), *Regional Development of the brain in early life.* (pp.

3-70). Oxford: Blackwell.

Ziegler, G., Hauser, T. U., Moutoussis, M., Bullmore, E. T., Goodyer, I. M., Fonagy, P., . . .

Consortium, N. (2019). Compulsivity and impulsivity traits linked to attenuated

developmental frontostriatal myelination trajectories. *Nature Neuroscience*.

doi:10.1038/s41593-019-0394-3

	MRI sample	Cognition sample
N	621	611
Age (years)	3.2-21.0 (mean= 12.4, SD= 4.8)	3.2-21.0 (mean= 12.5, SD= 4.8)
Sex	Boys= 318, Girls = 303	Boys=310, Girls = 301
Self-reported Ethnicity	Hispanic/Latino = 140 Pacific Islander = 5 Asian = 88 African American = 102 Native American = 31 White = 426	Hispanic/Latino = 136 Pacific Islander = 4 Asian = 86 African American = 100 Native American= 31 White = 420
Handedness	Right handed= 532 Left handed= 60 Mixed =24 Not established = 3 Not available = 2	Right handed = 523 Left handed= 60 Mixed = 24 Not established = 3 Not available = 1
Ever diagnosed with ADHD	Yes = 48 No = 567 Not available = 6	Yes = 48 No = 557 Not available = 6

626   Table1. Sample demographics. The table shows sample demographics for the MRI sample and the

627   cognition sample. The total ethnicity numbers exceed total sample size as several subjects identified as

628   belonging to more than one ethnicity group.

Test	Cognitive Ability	Abbreviation
Dimensional Change Card Sort Test	Cognitive flexibility	Cognitive flexibility
Flanker Inhibitory Control and Attention Test	Inhibitory control	Inhibition
	Visual attention	Attention
Picture Sequence Memory Test	Episodic memory	Episodic memory
Pattern Comparison Processing Speed Test	Processing speed	Processing speed
Oral Reading Recognition Test	Oral reading skill	Reading
List Sorting Working Memory Test	Working memory	Working memory
Picture Vocabulary Test	Vocabulary knowledge	Vocabulary

639   Table 2. Overview of the NIH Toolbox Cognition Battery measures. The table shows the tests in order

640   as presented to subjects, the cognitive abilities assessed, and the abbreviations used in the current

641   paper.

Cortical region	Age effect (t)		Age-Sex interaction effect (t)		General cognitive ability effect (t)	
	LH	RH	LH	RH	LH	RH
Banks superior temporal sulcus	10.5	8.6	2.4	2.1	-2.0	-2.0
Caudal anterior cingulate	8.6	9.2	2.4	2.6	-2.3	-1.9
Caudal middle frontal	14.4	13.4	3.0	2.3	-3.1	-2.4
Cuneus	5.5	3.9	1.7	1.2	-1.3	-1.2
Entorhinal	7.9	5.7	1.9	1.1	-1.9	-1.1
Fusiform	8.2	7.0	1.8	1.1	-1.5	-1.1
Inferior parietal	10.3	9.4	3.1	2.9	-2.0	-2.6
Inferior temporal	8.5	6.8	2.1	1.3	-1.7	-1.6
Isthmus cingulate	8.8	8.8	1.2	1.0	-1.5	-1.4
Lateral occipital	6.3	6.2	2.0	2.1	-1.4	-1.9
Lateral orbitofrontal	11.4	11.9	3.2	2.3	-2.8	-1.8
Lingual	6.0	5.8	1.4	1.3	-1.3	-1.1
Medial orbitofrontal	10.7	10.9	2.8	2.9	-1.9	-1.7
Middle temporal	10.5	7.7	2.5	1.6	-2.3	-2.4
Parahippocampal	6.3	5.9	1.2	1.1	-1.1	-0.4
Paracentral	12.4	13.0	2.3	2.4	-2.6	-2.3
Pars opercularis	12.6	10.8	3.1	1.5	-3.1	-2.3
Pars orbitalis	13.0	12.6	3.8	2.4	-3.4	-2.7
Pars triangularis	13.0	12.3	3.4	2.3	-3.3	-2.5
Pericalcarine	3.6	2.9	2.1	2.1	-1.7	-1.3
Postcentral	13.0	10.9	2.5	2.3	-2.9	-2.4
Posterior cingulate	8.9	9.1	1.8	1.8	-2.1	-1.9
Precentral	14.6	13.3	2.7	2.2	-3.0	-2.6

Precuneus	9.7	9.7	2.4	2.2	-2.0	-2.0
Rostral anterior cingulate	9.2	9.7	2.7	3.0	-2.0	-1.9
Rostral middle frontal	12.9	13.4	3.7	3.2	-2.9	-2.5
Superior frontal	12.4	12.6	3.3	3.3	-2.5	-2.3
Superior parietal	11.0	10.6	2.8	2.6	-2.2	-2.4
Superior temporal	11.1	7.6	2.4	1.4	-2.6	-2.1
Supramarginal	12.3	10.0	2.7	2.5	-2.4	-2.6
Frontal pole	11.0	12.5	3.4	3.7	-2.0	-2.5
Temporal pole	7.8	6.7	2.4	1.1	-3.3	-2.0
Transverse temporal	12.2	10.1	1.2	0.8	-2.4	-1.7
Insula	8.5	6.0	1.8	0.9	-2.7	-1.6

Table 3. ROI based mean t-statistic of T1w/T2w ratio associations. The table shows each of the 34 ROIs, and the mean t-statistic within each region for the association between T1w/T2w ratio and age, age*sex and general cognitive ability within left and right hemisphere separately.

Figure 1. Age and sex distribution for the full MRI sample (n= 621). (A) Depicts the age and sex

distribution of the full MRI sample, while (B) depicts the age distribution within each scanner.

Figure 2. Correlation matrix for age, sex and cognition. The figure depicts Pearson's correlation

coefficients for age, sex, general cognitive ability (General Cog. Ability), and individual cognitive

sum scores. Warm colors represent positive correlations, while cold colors represent negative

correlations. Sum scores, and by extension general cognitive ability, are residualized by linear and

quadratic age.

Figure 3. Mean T1w/T2w ratio map. The figure depicts the mean of all T1w/T2w ratio maps from the

full MRI sample (n= 621), overlaid on the average midthickness surface. Warm colors represent

regions with higher T1w/T2w ratio, while cold colors represent regions with lower T1w/T2w ratio.

Figure 4. Vertex-wise associations between T1w/T2w ratio and age. The figure depicts the t statistics

map, masked by familywise error - across contrast corrected p values thresholded at a minimum $-\log_p$

of 1.6 to correct for both hemispheres. Warm colors represent a postive age association.

Figure 5. Regional loess visualizations of associations between T1w/T2w ratio and age. The plots

show age plotted against mean T1w/T2w ratio within a selection of regions from different cortical

lobes. Datapoints and smooth splines are color coded by hemisphere. T1w/T2w ratio values are

residualized by sex and scanner.

Figure 6. Vertex-wise associations between T1w/T2w ratio and the interaction between age and sex.

The figure depicts the t statistics map, masked by familywise error - across contrast corrected p values

thresholded at a minimum $-\log_p$ of 1.6 to correct for both hemispheres. Warm colors represent a

postive interaction effect.

Figure 7. Visualizations of T1w/T2w ratio versus age color coded for sex. The plots shows age plotted

against mean T1w/T2w ratio within frontal regions, chosen based on the highest regional age*sex

mean t-statistics, with datapoints and smooth splines color coded by sex. T1w/T2w ratio values are

residualized by scanner.

Figure 8. Vertex-wise associations between T1w/T2w ratio and general cogntive ability. The figure

depicts the t statistics map, masked by familywise error - across contrast corrected p values

threasholded at a minimum –log$_p$ of 1.6 to correct for both hemispheres. Cold colors represent a

negative association between T1w/T2w ratio and general cogntive ability.

Figure 9. Visualizations of T1w/T2w ratio versus age, color coded by general cognitive ability. The

plots depict age plotted against mean T1w/T2w ratio within frontal regions, chosen based on the the

highest regional mean t-statistics, with data points and smooth splines color coded by general cogntive

ability, resulting in 4 equal qunatiles with worst (yellow) to best (blue) cognitive performers.

T1w/T2w ratio is sex and scanner residualized, while general cognitive performance is residualized by

linear and quadratic age.

710 Figure 10. Vertex-wise associations between T1w/T2w ratio and individual cognitive sum scores. The

711 figure depicts t statistics maps from top to bottom for (A) inhibition, (B) attention, (C) vocabulary, (D)

712 reading, and (E) working memory. The maps are masked by familywise error - across contrast

713 corrected p values thresholded at a minimum –log$_p$ of 1.6 to correct for both hemispheres. Cold

714 colors represent a negative association between T1w/T2w ratio and each sum score.

716