1   **Title:**

3   **A tripartite flip-flop sleep circuit switches sleep states**

7   **Authors:** Elisabeth Maluck†, Inka Busack†, Judith Besseling, Florentin Masurat, Michal

8   Turek, Henrik Bringmann*

Affiliations:

Max Planck Institute for Biophysical Chemistry, Am Fassberg 11, 37077 Göttingen,

Germany

University of Marburg, Karl-von-Frisch-Str. 8, 35043 Marburg, Germany

*correspondence to: Henrik.Bringmann@mpibpc.mpg.de

†these authors contributed equally

Abstract

Inhibitory sleep-active neurons depolarize at sleep onset to shut down the activity of

wakefulness circuits. Wake-active arousal neurons in turn suppress inhibitory sleep-active

neurons, thus forming a bipartite flip-flop switch. However, how sleep states are switched

is unclear because neural circuits that directly depolarize inhibitory sleep-active neurons

are not understood. Using optogenetics, we solved the presynaptic circuit for depolarization

of the sleep-active RIS neuron in *C. elegans*. Surprisingly, we found that the PVC forward

command interneuron, which is known to control wake behavior, is a major activator of

RIS. PVC is inhibited by reverse command interneurons, which are stimulated by arousing

cues. This suggests a model for sleep switch operation in which declining arousal increases

activation of PVC, thus triggering activation of RIS. Depolarization of RIS in turn

promotes the activation of PVC, thus forming a positive feedback loop for all-or-none sleep

induction. The flip-flop sleep switch in *C. elegans* thus is tripartite and requires excitatory

sleep-promoting neurons activated by wakefulness that act as an amplifier that translates

reduced arousal into the depolarization of an inhibitory sleep-active neuron. A tripartite

flip-flop switch likely also underlies sleep state switching in other animals including in

mammals.

Introduction

Sleep is a behavior that affects many, if not all, physiological processes. Disorders and curtailment of sleep affect the lives of 10-30% of the adult population of modern societies. Sleep deprivation is detrimental to human health and is associated with an increased risk of infection, cardiovascular disease, depression, obesity, type 2 diabetes, and cancer. The high prevalence of insomnia and insufficient sleep quality thus presents a massive unmet health and economic problem[1-4]. To understand how behavior is generated, it is crucial to solve the underlying neural circuits. For example, the circuit of the crustacean stomatogastric nervous system showed that neuromodulation alters behavior without changing the connectome[5], the learning circuits in *Aplysia* revealed synaptic mechanisms of memory formation[6], and the grid cell circuit in the mammalian entorhinal cortex demonstrated how a map of the spatial environment can be encoded[7].

Sleep circuits require inhibitory sleep-active sleep-promoting neurons, which depolarize specifically at sleep onset and actively induce sleep by releasing inhibitory neurotransmitters, GABA and neuropeptides, to dampen arousal and the activity of wake circuits[8]. The depolarization of inhibitory sleep-active neurons defines the key properties of sleep behavior, which are the suppression of voluntary movements and sensory perception, reversibility, and homeostasis[9]. Inhibitory sleep-active neurons suppress wake circuits leading to a reduction of voluntary movement, sensory processing, and reduced responsiveness to stimulation. Inhibitory sleep-active neurons can be rapidly suppressed by arousing stimulation to allow for quick awakening. Forced wakefulness is followed by an increase of inhibitory sleep-active neuron depolarization, which leads to homeostatic sleep corrections. Thus, understanding sleep control requires comprehension of the circuit mechanisms that determine when and how much inhibitory sleep-active neurons depolarize[8, 10].

Circuits control the depolarization of inhibitory sleep-active neurons. For example, wake-active wake-promoting neurons promote arousal and suppress inhibitory sleep-active neurons, whereas sleep need causes tiredness and inhibitory sleep-active neuron

depolarization. Thus, inhibitory sleep-active sleep-promoting and wake-active wake-
promoting neurons form a flip-flop switch, which ensures that sleep and wake exist as
discrete states. This sleep switch is under the control of arousal that favors wake and
inhibits sleep through the suppression of sleep-active neurons by inhibitory wake-active
neurons[8, 11]. It has been proposed that sleep induction is favored by disinhibition of
inhibitory sleep-active neurons [12-14], and also excitatory sleep-active neurons exist that
might perhaps present activators of inhibitory sleep-active neurons [15]. However, the
forces and mechanisms that flip the sleep switch from wake to sleep when an organism
gets tired cannot be satisfactorily explained by the present circuit models as it is unclear
how inhibitory sleep-active neurons are turned on when the system is set to sleep.

Sleep is under circadian and homeostatic controls that determine the timing of sleep and
ensure that enough of this essential physiological state takes place[16]. Sleep homeostasis
comprises multiple mechanisms that act on different time scales. On long time scales sleep
is a function of prior wakefulness, i.e. prolonged wakefulness leads to increased sleep
propensity and sleep loss triggers compensatory increases in the intensity or duration of
sleep. This chronic sleep homeostasis likely is mediated by several parallel mechanisms.
For example, in mammals, somnogens such as adenosine accumulate during wakefulness
leading to the inhibition of wake-promoting neurons[17, 18]. In *Drosophila*, activity-
dependent plasticity of sleep-promoting neurons increases during wakefulness to increase
subsequent sleep[19, 20]. On short time scales, acute homeostasis determines whether the
system's actual state matches the system's set point and carries out corrective action if
those values do not match. For example, to homeostatically maintain sleep despite
disturbance, micro arousals need to be compensated for. In humans, homeostatic sleep
maintenance can be seen in EEG recordings in the form of k-complexes, where a
spontaneous or evoked short cortical up state is followed by a down state[21-23].
Homeostatic sleep maintenance is also found during sleep in *C. elegans*, where sleep bouts
are interrupted by short motion bouts, with the length of a motion bout correlating with the
length of the subsequent sleep bout[24, 25]. Thus, across systems, homeostatic sleep
maintenance requires constant surveillance of sleep and corrective action.

Sleep-active sleep-promoting neurons are conserved regulators of sleep and have been
found both in vertebrates as well as in invertebrates[10, 26]. Mammals possess several
populations of sleep-active neurons across the brain, most of which are inhibitory,
including neurons in the anterior hypothalamus, brain stem, and cortex [8, 14]. Excitatory
sleep-active neurons were found in the periocular midbrain that project to inhibitory sleep-
active neurons in the anterior hypothalamus, whose role could be to activate inhibitory
sleep-active neurons, but this hypothesis has yet to be tested experimentally[15]. Studying
sleep in less complex brains facilitates sleep circuit analysis. In *Drosophila*, sleep-
promoting neurons are found at several locations in the brain. A well-characterized
population of sleep-promoting neurons is formed by ExFl1 neurons residing in the dorsal
fan-shaped body. R2 neurons accumulate homeostatic sleep pressure over time to promote
activation of ExFl1 neurons, probably by an indirect mechanism [19, 20]. *C. elegans*
possesses a single inhibitory sleep-active neuron called RIS. Like its mammalian
counterparts, RIS depolarizes at sleep onset. RIS is crucial for sleep induction as its
ablation leads to a virtually complete loss of detectable sleep bouts[27-29]. The small,
invariant nervous system, its mapped connectome, and the transparency of *C. elegans*
facilitates neural circuit analysis[30]. However, the specific neural circuits that control RIS
activity are not yet understood.

C. elegans shows sleep behavior during many stages and conditions. Here we analyzed
sleep behavior during lethargus, the stage prior to each of the four molts during larval
development [10, 29, 31-33]. We used optogenetics to dissect the neural circuit that
controls the activation of the inhibitory sleep-active RIS neuron in *C. elegans*. We found
that the PVC command interneurons, which are activated by wakefulness, are sleep-
promoting by turning on the RIS sleep-active neuron. PVC activity is inhibited by reverse
arousal command interneurons that thus inhibit sleep. This tripartite flip-flop circuit can
explain how arousing stimulation inhibits RIS depolarization, how RIS depolarization is
homeostatically controlled, and how reduced arousal can induce RIS depolarization.

Results

A command interneuron circuit controls RIS activity

RIS is crucially required for sleep and activates during each sleep bout (Figure 1A) [27]. However, the presynaptic driver neurons that activate and control this neuron are not known. To identify the circuits controlling RIS activation transients, we optogenetically tested the role of neurons that are presynaptic to RIS according to the *C. elegans* connectome [30]. The neurons AVJL, CEPDL, URYVL, RIMR, PVCL, and SDQL have been shown to be presynaptic to RIS[30]. To find out how these presynaptic neurons control RIS, we activated them with ReaChR and green light, and followed RIS calcium activity during and outside of lethargus. Because there were no specific promotors available for the expression in SDQL and PVC, we expressed ReaChR using semi-specific promoters and selectively illuminated only the presynaptic neuron class. We used L1 larvae for most of the optogenetic experiments to dissect the circuit. As SDQL is born postembryonically and likely is not yet functional during the L1 stage, we used L4 larvae to assay its function[34]. We compared the effects of optogenetic stimulation outside and during lethargus, defined as the period prior to the molt during which the animals do not feed[35]. Before lethargus, we measured an activation of RIS upon depolarization of AVJ, CEP and SDQL, whereas activation of the other neurons tested caused no significant change in RIS activity. During lethargus, the activation of CEP, PVC and SDQL caused RIS activation (Figure 1B, Supplementary Figure 1A). All neurons showed consistent effects on RIS depolarization except RIM. While RIM caused only a minor and non-significant net activation of RIS in the averaged measurements, RIM activation either activated or inhibited RIS depending on the trial (Figure S1B-C).

PVC appeared to be a strong activator of RIS only during lethargus. This suggested that either the PVC to RIS connection might be specific for lethargus or that it has not yet matured during the mid L1 stage. We hence repeated the experiment and activated PVC in L2 larvae. PVC activated RIS both during and outside of lethargus, but the activation during lethargus was much stronger, suggesting that the activation of RIS by PVC is strongly enhanced during lethargus (Figure S1C). To find out which presynaptic neurons are required for inhibition or activation of RIS during lethargus, we tested the effect of

optogenetic inhibition of the presynaptic neurons on RIS activation. We used ArchT, which hyperpolarizes neurons by pumping protons out of the cell [36, 37]. As before, we specifically illuminated each presynaptic neuron class and quantified RIS activation using calcium imaging. Before lethargus, inhibition of AVJ and PVC led to an inhibition of RIS, whereas inhibition of the other neurons tested had no acute significant effect on RIS. During lethargus, only optogenetic inhibition of PVC led to significant RIS inhibition, whereas there was no effect seen for the other neurons (Figure 1C, Figure S2A). While inhibition of RIM had no significant net effect on RIS, in roughly one quarter of the measured trials, the inhibition of RIM led to RIS inhibition before and during lethargus (Figure S2B).

Our optogenetic analysis revealed a complex set of presynaptic inputs for regulation of RIS activity. The optogenetic gain-of-function experiments suggest that CEP, PVC, and SDQL present the most potent presynaptic activators of RIS. The capacity of PVC to activate RIS is strongly increased during lethargus, indicating that this neuron is involved in the lethargus-specific activation of RIS. The optogenetic loss-of-function experiments suggest that PVC is an essential presynaptic activator of RIS during lethargus, but additional neurons might contribute to RIS activation during lethargus or other conditions. RIM has the potential to both inhibit and activate RIS. While the activation could be direct, RIM is known to be an inhibitor of PVC through activation of the reverse command interneurons AVA/AVD/AVE and could thus perhaps inhibit RIS indirectly [38-40]. Consistent with the lack of net effects of RIM on RIS, RIM ablation does not change spontaneous sleep amounts (Figure S2C). Therefore, the majority of synaptic inputs into RIS is activating and inhibition appears to be predominantly indirect. The CEP, URY, and SDQL neurons present sensory receptors and might play a role in activating RIS in response to stimulation [41, 42]. Because of the strong and lethargus-specific effects of the PVC neurons on RIS activation we focused our analysis on these key neurons.

As the PVC neurons are crucial activators of RIS we tested their role in inducing sleep behavior. We used a strain that ablated PVC and other command interneurons by expressing the pro-apoptosis regulator ICE from the *nmr-1* promotor and measured sleep

and RIS activation[43]. Command interneuron ablation reduced sleep bouts during lethargus by about 76% (Figure S3A). The command interneuron-ablated worms also generally moved much slower (Figure S3B) and RIS activation was reduced by 63% (Figure S3C). Quiescence bouts did not occur at the beginning of the lethargus phase as defined by cessation of feeding, and were only observed around the middle of the lethargus phase (Figure S3D). An independently generated strain that ablates command interneurons using *egl-1* expression caused a reduction of sleep by 81% (Figure S2A). Because the command interneurons are controlled by glutamatergic signaling we also tested mutants in which this type of signaling is impaired and also observed impaired sleep consistent with ablating command interneurons (Figure S4).

To be able to more specifically manipulate PVC and to test the effects of PVC inhibition on behavior without affecting the other command interneurons, we used a more specific promoter for expression in PVC, which had been identified from single-cell RNA sequencing data [44, 45](Jonathan Packer, personal communication). There was no gene in the available datasets that was expressed only in the cluster of cells containing PVC, but the previously uncharacterized gene *zk673.11* was expressed strongly in PVC and in only a few other neurons excluding other command interneurons and therefore we used the promoter of this gene for PVC expression (Figure S5). Hyperpolarization of PVC through activation of ArchT led to an acute inhibition of RIS, an increase in locomotion, and a reduction of sleep (Figure 1D). These experiments show that RIS is controlled by a command interneuron circuit. The major command interneuron activator of RIS during L1 lethargus are the PVC forward command interneurons (Figure 1E). While PVC has previously been shown to promote forward locomotion[46], its inhibition leads to increased mobility implying that the reduction of locomotion after simultaneous ablation of most command interneurons stems from the reverse command interneurons that promote mobility, whereas PVC appears to play a predominant role in dampening motion behaviors through activation of RIS.

RIS and PVC activate each other forming a positive feedback loop

PVC presents a major activator of RIS, but how a forward command interneuron can cause

massive activation of the RIS neuron during sleep bouts is not clear. We hence tested how

optogenetic RIS activation affects PVC activity. We selectively activated RIS using

ReaChR and measured calcium activity in PVC (Figure 2A). Because the calcium

transients observable in PVC are typically small we used immobilized worms to reduce

measurement noise[47]. Upon RIS stimulation, PVC immediately displayed unexpectedly

strong calcium transients, which were slightly stronger during lethargus (Figure 2B, Figure

S6A). These results show that PVC and RIS activate each other, thus forming a positive

feedback loop. The strong calcium activation of PVC by RIS depolarization is striking

because activation transients in PVC have hitherto only been observed after mechanical

stimulation, whereas spontaneous PVC activity transients have not yet been reported[47].

The sleep-inducing RIS neuron has so far only been shown to inhibit other neurons making

PVC the first neuron which is not inhibited but activated by RIS. For example, command

interneurons such as AVE/AVA and other neurons are not activated by RIS but

inhibited[27]. While PVC is presynaptic to RIS, RIS is not presynaptic to PVC. The

activation of PVC by RIS could involve diffusional mechanisms or could be indirect

through other neurons, perhaps mediated by the inhibition of a PVC inhibitor such as

AVA/AVD/AVE. It is currently not possible to measure spontaneous calcium transients in

PVC, but our results would predict that PVC should spontaneously activate during sleep

bouts, which would characterize PVC not only as a neuron active during wakefulness as

suggested by previous studies[46, 47] but also as a sleep-active neuron. It would thus differ

from previously identified sleep-active neurons in that it does not induce sleep directly by

inhibiting wake circuits but indirectly through activation of an inhibitory sleep-active

sleep-inducing neuron. As PVC is activated by wakefulness and also promotes sleep, it

would thus be ideally suited to mediate the transition from active to quiescent states.

Together, these data suggest that PVC and RIS rely on positive feedback to achieve strong

transients during a sleep bout.

If depolarization of RIS activates PVC, what consequences does hyperpolarization of RIS

have on PVC activity? To answer this question, we measured the response of PVC to RIS

inhibition. We hyperpolarized RIS optogenetically for one minute using ArchT and

measured the activity of PVC. Interestingly, PVC showed a small but significant activity

increase during RIS inhibition, an effect which was increased during lethargus (Figure 2C).

The disinhibition of PVC by RIS inactivation is likely not direct and may reflect a general

increase in neuronal and behavioral activity that is caused by RIS inhibition and that

extends to the PVC neurons. Since PVC is a major activator of RIS, its disinhibition could

be part of a homeostatic feedback regulation.

Our results suggest that there is a positive feedback from sleep induction onto RIS

activation and that full RIS activation is only possible when sleep is successfully induced,

explaining the strong correlation of RIS depolarization and sleep bout induction[29]. This

model would predict that RIS transients are dampened if RIS is not able to induce sleep

bouts. To test this idea we analyzed RIS calcium transients in *aptf-1(-)* mutant worms in

which RIS still shows depolarization transients during lethargus, but cannot efficiently

induce quiescence behavior because its inhibitory neuropeptides are not expressed[27, 48].

In *aptf-1(-)* mutant animals, calcium transients' maxima were reduced by about 35%

(Figure 2 D-E). These results are consistent with the idea that sleep induction is a self-

enforcing process in which RIS-mediated inhibition of brain activity promotes further RIS

activation (Figure 2F).

RIS depolarization is under homeostatic control

The design of the sleep circuit suggests an intimate mutual control mechanism of RIS and

command interneurons that could allow homeostatic control of sleep. Arousing stimulation

is known to inhibit sleep-active neurons and to increase subsequent sleep[24, 25, 27, 29].

We thus hypothesized that inhibition of RIS leads to its subsequent depolarization, forming

a homeostat that allows maintaining or reinstating sleep bouts. We tested this hypothesis

by optogenetically hyperpolarizing RIS and following its activity using calcium imaging.

We inhibited RIS directly for 60 seconds by expressing the light-driven proton pump

ArchT specifically in this neuron and used green light illumination to activate ArchT. We

followed RIS calcium activity using GCaMP during the experiment and quantified

behavior. Optogenetic hyperpolarization of RIS led to a decrease in intracellular calcium

and increased behavioral activity. Approximately one minute after the end of the inhibition,

RIS showed a rebound activation transient during which calcium activity levels increased

strongly and rose well above baseline levels, concomitant with a decrease in behavioral

activity. Overall brain activity measurements showed that behavioral activity and brain

activity correlated throughout the experiment (Figure 3A). Strikingly, while the rebound

transient was also measurable outside of lethargus, the strength of the RIS rebound

depolarization was three-fold stronger during lethargus than before lethargus, indicating

that the propensity for RIS rebound activation is strongly increased during lethargus.

To test whether rebound activation of RIS mediates acute or chronic homeostasis, we tested

whether the strength of the rebound activation is a function of length of prior inhibition.

For this experiment we increased the length of the RIS inhibition and quantified the time

it took after the end of the stimulation until the rebound transient started as well as the peak

maximum of the rebound. After inhibiting RIS for five minutes, the rebound initiated

immediately after the end of the stimulation and the maximum that was reached exceeded

that observed after about one minute of RIS stimulation. Inhibiting RIS for ten minutes did

not further increase the occurrence or strength of the rebound transient. These results show

that RIS activation rebound transients rapidly saturate with increasing length of inhibition

(Figure 3B, Figure S6B-D) Thus, RIS shows a rebound activation upon inhibition that leads

to its own subsequent activation. A rebound activation was not only seen when RIS was

inhibited directly, but we also saw a reactivation of RIS after it was inhibited indirectly

using a blue light stimulus when the worms returned to sleep (Figure S7). The rebound

activation presents the translation of RIS inhibition into subsequently increased RIS

activity and thus sleep induction. Rebound activation of RIS does not seem to constitute a

chronic integrator of wake time but presents an acute homeostatic regulatory phenomenon

to induce, maintain, or reinstate sleep bouts.

If RIS is part of a homeostatic system and RIS inhibition triggers its subsequent activation,

does activation of RIS also trigger its own inhibition? To test this idea, we optogenetically

activated RIS and measured its activity before, during, and after the manipulation. We

optogenetically depolarized RIS for one minute using green light and the light-activated

cation channel ReaChR expressed specifically in RIS. Activity of RIS was followed using calcium imaging and increased strongly during optogenetic depolarization but was also reduced significantly after stimulation compared with baseline activity, indicating that RIS activity also shows a negative rebound. Again, this effect was highly increased, about five-fold, in worms that were in lethargus compared with worms outside of lethargus (Figure 3C). Thus, polarization of RIS during lethargus is under homeostatic control in both directions. Hyperpolarization leads to subsequent depolarization, perhaps through a reactivation of wake-active circuits that ultimately lead to the reactivation of PVC. Depolarization leads to subsequent hyperpolarization, perhaps through inhibition of wake-active circuits that cause PVC activation. This homeostatic regulation is highly enhanced during lethargus and can thus contribute to the repetitive pattern of RIS activation and thus the oscillating induction of consecutive sleep bouts[24].

Modest dampening of brain arousal occurs upstream of RIS

Our results demonstrate that PVC plays a major role in activating RIS involving self-enforcing positive feedback, resulting in strong RIS activation and thus sleep induction. RIS calcium transients during development outside of lethargus are small, whereas transients are high during lethargus. What determines that RIS calcium transients are limited outside of lethargus but promoted during lethargus? With other words, what tips the balance that allows the self-enforcing feedback loop of PVC and RIS to set off? As an important principle of command interneuron control, forward and reverse command interneurons inhibit each other to allow discrete forward and reverse locomotion states. The AVA/AVD/AVE/RIM interneurons initiate reverse locomotion by activating pre-motor interneurons while inhibiting the forward command circuit including AVB/PVC. By contrast, during forward movement reverse command interneurons are inhibited [46, 49].

Small changes in arousal and activity of the command interneurons can change the equilibrium of forward and reverse command interneurons [43]. Hyperactive mutants suppress sleep across species including *C. elegans* [50-52]. Many arousal cues trigger backwards escape movements and inhibit RIS [27, 29, 53]. Thus, previous studies on the

command interneuron circuit together with our results suggest that arousal inhibits RIS

through inhibiting PVC. This model of RIS activation would predict that there are changes

during lethargus that are upstream of RIS activity that change the properties of the

command circuit leading to increased PVC and thus RIS activation.

We reasoned that it should be possible to measure these changes that occur in command

interneuron activity upstream of RIS by characterizing neural activity and behavior in *aptf-*

1(-) mutant worms. We quantified behavior and command interneuron calcium levels

across lethargus in *aptf-1(-)* mutant worms. Wild type animals showed successive sleep

bouts and a 70% reduction in locomotion speed by during lethargus. By contrast, *aptf-1(-)*

mutant animals almost never showed quiescence bouts (Figure 2E), but nevertheless

locomotion speed was decreased by 25% during the lethargus phase (Figure 4). Consistent

with the behavioral activity reduction, there was a significant reduction of command

interneuron activity during lethargus also in *aptf-1(-)* mutant animals (Figure 4, Figure S8).

To further characterize the neuronal changes upstream of RIS-mediated sleep induction,

we imaged the activity of RIM as exemplary neurons during lethargus in *aptf-1(-)* mutants.

In wild type animals, RIM regularly showed activation transients before lethargus but did

not show many transients during lethargus. RIM showed not only a change in transient

frequency across the lethargus cycle but also a reduction in baseline calcium activity. In

aptf-1(-) mutant worms, RIM continued showing RIS transients during lethargus,

indicating that RIS inhibits RIM transients during sleep bouts. However, reduction of

baseline calcium activity was preserved in *aptf-1(-)*, indicating that RIM activity is

dampened during lethargus independently of RIS at the level of baseline calcium activity.

Together these experiments indicate that a dampening of behavioral and neural baseline

activity occurs during lethargus that is independent of RIS. This neuronal baseline and

behavioral dampening itself appears not to be sufficient to constitute normal sleep bouts,

but could hypothetically lead to a weaker activity of reverse arousal neurons thus allowing

stronger or longer activation of forward command interneurons including PVC and thus

RIS activation[43, 54].

An arousing stimulus inhibits RIS through RIM

Arousal plays a major role in inhibiting sleep but the circuits that mediate the effect of

arousing stimuli on RIS inhibition are not well understood. We hence studied the circuit

by which stimulation of a nociceptor, the ASH neurons, leads to a reverse escape response

and inhibition of RIS[55]. We optogenetically stimulated ASH using ReaChR and green

light and followed RIS and RIM activities. ASH activation led to the activation of the RIM

neuron and triggered a backwards response as previously described [55, 56].

Simultaneously, RIS was inhibited (Figure 5A). RIM can inhibit PVC through reverse

interneurons which it synchronizes, making it ideally suited to mediate the effects of ASH

stimulation[39, 46]. To test whether RIM is required for RIS inhibition, we ablated RIM

genetically by expression of *egl-1* from the *tdc-1* promoter, which is specific to RIM and

RIC, and repeated the optogenetic stimulation of ASH. We used L4 stage animals as the

genetic ablation was not yet effective in L1 animals. In RIM-ablated animals, activation of

ASH caused the opposite effect on RIS activity. Instead of inhibiting RIS, ASH activated

RIS, while still increasing behavioral activity (Figure 5B). Consistent with our calcium

imaging data, ASH stimulation after RIM ablation predominantly caused a forward

locomotion response (Figure 5C). Further supporting the idea that reverse interneurons

inhibit and forward PVC neurons activate RIS during stimulation, gentle tail touch

increased RIS activity more strongly when RIM was ablated (Figure S9). These results

delineate a circuit model for how sensory stimulation can control RIS activation. By

activating the reverse response that is mediated by RIM, PVC is suppressed leading to RIS

inhibition. In addition, sensory stimulation can also promote RIS activation if the forward

command circuit is activated.

Discussion

A tripartite flip-flop circuit controls sleep

Optogenetic activation and inhibition showed how the activity of presynaptic neurons

affects RIS depolarization during developmental sleep. Several presynaptic neurons can

activate RIS, but potent direct inhibitors were not identified. Loss-of-function experiments

showed that the command circuit controls activation of RIS, with PVC presenting the key

activator of RIS. PVC has long been known to mediate the forward escape response by

transmitting information from posterior sensory neurons to activate AVB pre-motor

neurons to trigger forward locomotion[46, 47, 57]. Consistent with promoting the forward

escape response, optogenetic activation of PVC leads to an, albeit modest, increase in

forward movement[57, 58]. Reverse movement, in turn, is mediated by AVA, AVE, and

AVD command premotor interneurons, which activate reverse motor neurons. Forward

PVC and reverse AVA/AVE/AVD command interneurons are presynaptic to and mutually

inhibit each other, which ensures discrete forward and reverse locomotion states analogous

to a flip-flop switch [40, 43, 46].

Our finding that PVC presents a key activator of RIS immediately suggests a model for

how RIS is controlled and provides a potential mechanism for linking sleep induction to

decreasing arousal and for homeostatically maintaining a series of sleep bouts. According

to this model, during conditions of high arousal, such as during development outside of

lethargus, larvae are constantly awake. The command interneuron circuit cycles between

forward and reverse states, leading to the activation of forward or reverse motor programs,

respectively[40, 46, 59]. Arousal promotes reverse command interneurons activity and thus

limits PVC and RIS activation. Independently of RIS, lethargus induces a modest neural

dampening that includes command interneuron activity that is evident as a decrease in

baseline calcium activity and reduction of locomotion speed. Consistent with the view that

reduction of neuronal activity can occur partially independently of RIS, the dampening of

sensory neurons during lethargus is not fully reversible upon stimulation[60, 61]. This

modest but broad activity dampening could cause a modest shift of command interneuron

activity that should lead to a modest favoring of PVC activity, perhaps through

disinhibition. Consistent with this model, sleep bouts are typically induced from long

forward movement bouts or periods of reduced locomotion activity and not from reverse

movements [54, 59, 62]. Mutations that increase arousal and suppress sleep increase the

activity of reversal neurons, whereas conditions that decrease arousal decrease the activity

of the reversal neurons and increase sleep[40, 51, 52]. Also, the ablation of reverse

command interneurons such as AVE reduces reversals and leads to ectopic quiescence[47,

63]. The increased activation of PVC in turn depolarizes RIS. Because PVC and RIS form

a self-reinforcing positive feedback loop, a modest increase of PVC activity could translate

non-linearly to a strong activation of RIS, which in turn induces the shutdown of behavior

that is required for the sleep state [24, 31, 54, 60, 61]. The activation of RIS leads to the

secretion of inhibitory neuropeptides that inhibit key arousal neurons[27, 48]. While PVC

is presynaptic to RIS and likely activates it directly, RIS is not presynaptic to PVC,

suggesting an indirect mechanism that perhaps could involve inhibition of reverse

command interneurons including AVA/AVE, which are inhibited by RIS [27] and could

thus lead to further activation of PVC through disinhibition. Thus, the process of sleep

induction, once it reaches a critical threshold, could become self-enforcing. In summary,

our model allows a circuit understanding of the mutual control of the wake-active central

pattern generator formed by the command interneurons and the sleep-active RIS neuron

and provides the mechanistic basis for understanding the correlation of sleep and

locomotion patterns observed in pan-neuronal imaging studies[54, 62]. Our model suggests

that the sleep switch is tripartite and includes self-enforcing inhibitory sleep-active sleep-

promoting neurons as well as excitatory wake-activated sleep-promoting neurons as

mediators of switch flipping. This sleep switch acts as an amplifier that can translate a

modest reduction of arousal into a massive shutdown of behavioral activity during sleep.

Dampening of neural activity independently of sleep-active neurons could be interpreted

as a neural equivalent of tiredness that leads to an increased propensity to activate sleep-

active neurons and to induce sleep bouts. Thus, the sleep switch could present an amplifier

that translates reduced brain activity into sleep bouts (Figure 6). Further experiments will

be necessary to find out how PVC activity is controlled by sleep-promoting signals such

as reduced arousal.

What causes the termination of sleep bouts? The PVC and RIS neurons might not be able

to sustain prolonged activity leading to the cessation of a sleep bout. The RIS activation

transient and thus sleep bout can be blunted by a sensory or optogenetic arousing stimulus

[27, 29, 54, 61]. Arousing stimulation, for instance by activating the nociceptive sensory

neurons, triggers a reverse escape response through backwards command and RIM

interneurons [39, 56, 59, 64]. Optogenetic RIM depolarization variably causes activation

or inhibition of RIS but the cause of this probabilistic outcome is unclear. Potentially, the

activating effect of RIM on RIS could be mediated by direct synaptic innervation and might

serve a buffering or probabilistic function [39], consistent with the almost zero net effects

of RIM depolarization on RIS activity. However, stimulation of the nociceptive ASH

neurons causes consistent inhibition of RIS that depends on RIM, suggesting that a main

physiological role of RIM is to inhibit sleep upon arousing stimulation, perhaps by

synchronizing the reverse interneurons[39]. RIM activation can inhibit sleep also in

response to acute food deprivation [65, 66]. Thus, RIM might present an arousal module

that can be activated upon sensing various external conditions that signal the need to

suppress sleep (Figure 6).

RIS inactivation leads to disinhibition of arousal and brain activity starting anew the cycle

of command interneuron activity and locomotion behavior. Depending on the arousal

levels, PVC activity should increase either immediately or after a delay until it reaches a

critical level sufficient to cause substantial RIS reactivation and thus a return to sleep. The

timing at which the rebound activation occurs can be controlled by the level of arousal,

with strong arousal leading to longer wake periods before the return to sleep, whereas

milder stimulations cause the immediate return to sleep[25]. Consistent with this circuit

model of recurrent RIS activation, RIS activity oscillates, resulting in the typical pattern of

sleep bouts that are interrupted by activity bouts[24]. This circuit design allows

homeostatic sleep maintenance of a series of consecutive sleep bouts with sensory

stimulation restarting the cycle of RIS activation thus prompting an acutely increased RIS

activation causing the return to sleep (Figure 6A)[25, 54].

How can PVC mediate seemingly contrary behaviors, the forward escape response and

sleep bout induction? Sensory mechanical stimulation causes general arousal. For example,

tail touch not only activates the forward PVC neurons to trigger a forward locomotion

response, but also activates the reverse AVA neurons, with a net dominance of the PVC

activation [46, 57]. Because the reverse command interneurons inhibit the forward

interneurons, the sensory stimulation should lead to only a limited activation of PVC that

is sufficient to trigger a forward movement, but that is insufficient to cause strong RIS

activation and thus a sleep bout. Consistent with this idea, weakening of the reverse circuit

by RIM ablation changes the behavioral response of tail touch during lethargus, and PVC

activation outside of lethargus only causes a modest increase in forward movement[57,

58]. During normal waking behavior, modest RIS activation may thus function to dampen

the behavioral response to sensory stimulation and to regulate the speed of locomotion [67,

68].

Here we have identified a circuit controlling sleep-active neuron depolarization in *C.*

elegans. This work built on the neural connectome and was facilitated by the small size

and invariance of the nervous system as well as the transparency of the organism. While

the *C. elegans* sleep circuit clearly is built from less cells than the human sleep circuit,

there are many conceptual similarities. For instance, in both *C. elegans* and humans, sleep

is controlled by inhibitory sleep-active sleep-promoting neurons that depolarize at sleep

onset to actively induce sleep by inhibiting wake circuits. A main difference is that humans

have many brain centers each consisting of thousands of sleep-active neurons[14]. The

single RIS neuron is the major inhibitory sleep-active neuron required for sleep induction

in *C. elegans*[27]. Work in mammals revealed the general principles of wake-active wake-

promoting neurons and sleep-active sleep-promoting neurons as well as their mutual

inhibition. While this information explains the flip-flop nature of sleep and wake states,

there is no satisfactory understanding of what flips the sleep switch, i.e. how wakefulness

is detected when the system is set to sleep, prompting the activation of inhibitory sleep-

active neurons [8]. Our model for the operation of the *C. elegans* sleep circuit indicates

that flipping of the sleep switch can be understood if excitatory wakefulness-activated

sleep-promoting neurons and a positive feedback of sleep-active neuron depolarization are

added to the switch model. In this tripartite flip-flop sleep switch model, the sleep-active

sleep-promoting center is activated by an excitatory sleep-promoting center that is

activated by wakefulness. Wake-activated sleep-promoting neurons and wake-active

wake-promoting neurons inhibit each other thus forming the actual flip-flop switch.

Arousing stimulation increases the activity of the wake-active wake-promoting neurons

and thus suppresses excitatory wake-activated sleep-promoting neurons and thus sleep.

According to this model, reducing arousal increases the activity of excitatory wake-

activated, sleep-promoting neurons and thus favors inhibitory sleep-active sleep-promoting neuron depolarization, causing sleep behavior (Figure 6). Therefore, reduced arousal and activity of wake-active wake-promoting neurons could present a neural correlate of tiredness. Excitatory neurons in the periocular midbrain of mammals are similar to PVC in that they may be activators of inhibitory sleep-active neurons an could perhaps be part of circuits that operate by mechanisms analogous to the tripartite sleep switch of *C. elegans* [15].

Sleep is reversible by stimulation and hyperarousal is the major cause for insomnia in humans[2, 69, 70]. Homeostatic sleep maintenance is an essential feature of sleep and is found from worms to humans[21-23, 25]. R2 ring neurons in *Drosophila* present an integrator of wake time causing subsequently increased depolarization of ExFl1 sleep-inducing neurons, thus forming a chronic sleep homeostat [71, 72]. Our model of a tripartite flip-flop circuit suggests that excitatory wake-activated sleep-promoting neurons are part of an acute sleep homeostat that translates acute brain activity into increased sleep neuron activity when the system is set to sleep. Excitatory wake-activated sleep-promoting neurons measure systemic activity, i.e. they become active as a consequence of a global brain activity increase, and directly activate the inhibitory sleep-active neuron transiently. Thus, the interplay of inhibitory sleep-active sleep-promoting and excitatory wake-active sleep-promoting neurons form an oscillator that periodically sends out sleep-inducing pulses. Macroscopically, sleep in mammals exists as cortical oscillations of global down states, known as slow waves[73]. Micro arousals trigger cortical upstates that are followed by cortical down states, known as k-complexes[21-23]. Both slow wave activity as well as k-complexes could be hypothetically explained by the concept of excitatory wake-activated sleep-promoting circuits.

Acknowledgments

We are grateful to Jonathan Packer and Robert Waterson for suggesting the use of the *zk673.11* promoter. We thank Andres Mariqc and Yun Zhang for strains. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs

(P40 OD010440). This work was funded by the Max Planck Society (Max Planck Research

Group), a European Research Council Starting Grant (ID: 637860, SLEEPCONTROL),

and the University of Marburg. Additional data can be found in the supplement.

Author Contributions

E.M, I.B, F.M., J.B., and M.T. designed, performed, and analyzed experiments. E.M and

I.B edited the manuscript. H.B. acquired funding, conceived the project, designed

experiments, supervised the work, and wrote the manuscript.

Declarations of Interest

The authors declare that they have no competing interests.

References

1.   Irwin MR. Why sleep is important for health: a psychoneuroimmunology
perspective. Annu Rev Psychol. 2015;66:143-72. Epub 2014/07/26. doi:
10.1146/annurev-psych-010213-115205. PubMed PMID: 25061767; PubMed Central
PMCID: PMCPMC4961463.
2.   Drake CL, Roehrs T, Roth T. Insomnia causes, consequences, and therapeutics:
an overview. Depression and anxiety. 2003;18(4):163-76. Epub 2003/12/09. doi:
10.1002/da.10151. PubMed PMID: 14661186.
3.   Van Cauter E, Spiegel K, Tasali E, Leproult R. Metabolic consequences of sleep
and sleep loss. Sleep Med. 2008;9 Suppl 1:S23-8. Epub 2008/12/17. doi: 10.1016/S1389-
9457(08)70013-3. PubMed PMID: 18929315; PubMed Central PMCID:
PMCPMC4444051.
4.   Knutson KL, Spiegel K, Penev P, Van Cauter E. The metabolic consequences of
sleep deprivation. Sleep medicine reviews. 2007;11(3):163-78. Epub 2007/04/20. doi:
10.1016/j.smrv.2007.01.002. PubMed PMID: 17442599; PubMed Central PMCID:
PMCPMC1991337.
5.   Marder E. Neuromodulation of neuronal circuits: back to the future. Neuron.
2012;76(1):1-11. Epub 2012/10/09. doi: 10.1016/j.neuron.2012.09.010. PubMed PMID:
23040802; PubMed Central PMCID: PMC3482119.
6.   Mayford M, Siegelbaum SA, Kandel ER. Synapses and memory storage. Cold
Spring Harbor perspectives in biology. 2012;4(6). Epub 2012/04/13. doi:
10.1101/cshperspect.a005751. PubMed PMID: 22496389; PubMed Central PMCID:
PMC3367555.

7.   Hafting T, Fyhn M, Molden S, Moser MB, Moser EI. Microstructure of a spatial
map in the entorhinal cortex. Nature. 2005;436(7052):801-6. Epub 2005/06/21. doi:
10.1038/nature03721. PubMed PMID: 15965463.
8.   Saper CB, Fuller PM, Pedersen NP, Lu J, Scammell TE. Sleep state switching.
Neuron. 2010;68(6):1023-42. Epub 2010/12/22. doi: 10.1016/j.neuron.2010.11.032.
PubMed PMID: 21172606; PubMed Central PMCID: PMC3026325.
9.   Campbell SS, Tobler I. Animal sleep: a review of sleep duration across
phylogeny. Neurosci Biobehav Rev. 1984;8(3):269-300. Epub 1984/01/01. PubMed
PMID: 6504414.
10.   Bringmann H. Sleep-Active Neurons: Conserved Motors of Sleep. Genetics.
2018;208(4):1279-89. Epub 2018/04/06. doi: 10.1534/genetics.117.300521. PubMed
PMID: 29618588; PubMed Central PMCID: PMC5887131.
11.   Lu J, Sherman D, Devor M, Saper CB. A putative flip-flop switch for control of
REM sleep. Nature. 2006;441(7093):589-94. Epub 2006/05/12. doi:
10.1038/nature04767. PubMed PMID: 16688184.
12.   Morairty S, Rainnie D, McCarley R, Greene R. Disinhibition of ventrolateral
preoptic area sleep-active neurons by adenosine: a new mechanism for sleep promotion.
Neuroscience. 2004;123(2):451-7. Epub 2003/12/31. PubMed PMID: 14698752.
13.   Saper CB, Scammell TE, Lu J. Hypothalamic regulation of sleep and circadian
rhythms. Nature. 2005;437(7063):1257-63. Epub 2005/10/28. doi: 10.1038/nature04284.
PubMed PMID: 16251950.
14.   Weber F, Dan Y. Circuit-based interrogation of sleep control. Nature.
2016;538(7623):51-9. Epub 2016/10/07. doi: 10.1038/nature19773. PubMed PMID:
27708309.
15.   Zhang Z, Zhong P, Hu F, Barger Z, Ren Y, Ding X, et al. An Excitatory Circuit in
the Perioculomotor Midbrain for Non-REM Sleep Control. Cell. 2019;177(5):1293-307
e16. Epub 2019/04/30. doi: 10.1016/j.cell.2019.03.041. PubMed PMID: 31031008.
16.   Borbely AA. A two process model of sleep regulation. Human neurobiology.
1982;1(3):195-204. Epub 1982/01/01. PubMed PMID: 7185792.
17.   Frank MG. Astroglial regulation of sleep homeostasis. Curr Opin Neurobiol.
2013;23(5):812-8. Epub 2013/03/23. doi: 10.1016/j.conb.2013.02.009. PubMed PMID:
23518138.
18.   Krueger JM, Frank MG, Wisor JP, Roy S. Sleep function: Toward elucidating an
enigma. Sleep medicine reviews. 2016;28:46-54. Epub 2015/10/09. doi:
10.1016/j.smrv.2015.08.005. PubMed PMID: 26447948; PubMed Central PMCID:
PMC4769986.
19.   Donlea JM, Pimentel D, Miesenbock G. Neuronal machinery of sleep
homeostasis in Drosophila. Neuron. 2014;81(4):860-72. Epub 2014/02/25. doi:
10.1016/j.neuron.2013.12.013. PubMed PMID: 24559676; PubMed Central PMCID:
PMC3969244.
20.   Pimentel D, Donlea JM, Talbot CB, Song SM, Thurston AJ, Miesenbock G.
Operation of a homeostatic sleep switch. Nature. 2016;536(7616):333-7. Epub
2016/08/04. doi: 10.1038/nature19055. PubMed PMID: 27487216; PubMed Central
PMCID: PMC4998959.

21. Halasz P. Hierarchy of micro-arousals and the microstructure of sleep.
Neurophysiologie clinique = Clinical neurophysiology. 1998;28(6):461-75. Epub
1999/01/23. PubMed PMID: 9894227.
22. Amzica F, Steriade M. Cellular substrates and laminar profile of sleep K-
complex. Neuroscience. 1998;82(3):671-86. Epub 1998/03/04. PubMed PMID: 9483527.
23. Cash SS, Halgren E, Dehghani N, Rossetti AO, Thesen T, Wang C, et al. The
human K-complex represents an isolated cortical down-state. Science.
2009;324(5930):1084-7. Epub 2009/05/23. doi: 10.1126/science.1169626. PubMed
PMID: 19461004; PubMed Central PMCID: PMC3715654.
24. Iwanir S, Tramm N, Nagy S, Wright C, Ish D, Biron D. The microarchitecture of
C. elegans behavior during lethargus: homeostatic bout dynamics, a typical body posture,
and regulation by a central neuron. sleep. 2013;36(3):385-95. Epub 2013/03/02. doi:
10.5665/sleep.2456. PubMed PMID: 23449971; PubMed Central PMCID: PMC3571756.
25. Nagy S, Tramm N, Sanders J, Iwanir S, Shirley IA, Levine E, et al. Homeostasis
in C. elegans sleep is characterized by two behaviorally and genetically distinct
mechanisms. eLife. 2014;3:e04380. Epub 2014/12/05. doi: 10.7554/eLife.04380.
PubMed PMID: 25474127; PubMed Central PMCID: PMC4273442.
26. Joiner WJ. Unraveling the Evolutionary Determinants of Sleep. Curr Biol.
2016;26(20):R1073-R87. Epub 2016/10/26. doi: 10.1016/j.cub.2016.08.068. PubMed
PMID: 27780049; PubMed Central PMCID: PMC5120870.
27. Turek M, Lewandrowski I, Bringmann H. An AP2 transcription factor is required
for a sleep-active neuron to induce sleep-like quiescence in C. elegans. Curr Biol.
2013;23(22):2215-23. Epub 2013/11/05. doi: 10.1016/j.cub.2013.09.028. PubMed PMID:
24184105.
28. Bennett HL, Khoruzhik Y, Hayden D, Huang H, Sanders J, Walsh MB, et al.
Normal sleep bouts are not essential for C. elegans survival and FoxO is important for
compensatory changes in sleep. BMC neuroscience. 2018;19(1):10. Epub 2018/03/11.
doi: 10.1186/s12868-018-0408-1. PubMed PMID: 29523076.
29. Wu Y, Masurat F, Preis J, Bringmann H. Sleep Counteracts Aging Phenotypes to
Survive Starvation-Induced Developmental Arrest in C. elegans. Curr Biol.
2018;28(22):3610-24 e8. Epub 2018/11/13. doi: 10.1016/j.cub.2018.10.009. PubMed
PMID: 30416057; PubMed Central PMCID: PMCPMC6264389.
30. White JG, Southgate E, Thomson JN, Brenner S. The structure of the nervous
system of the nematode Caenorhabditis elegans. Philos Trans R Soc Lond B Biol Sci.
1986;314(1165):1-340. Epub 1986/11/12. PubMed PMID: 22462104.
31. Raizen DM, Zimmerman JE, Maycock MH, Ta UD, You YJ, Sundaram MV, et
al. Lethargus is a Caenorhabditis elegans sleep-like state. Nature. 2008;451(7178):569-
72. Epub 2008/01/11. doi: nature06535 [pii]10.1038/nature06535. PubMed PMID:
18185515.
32. Trojanowski NF, Raizen DM. Call it Worm Sleep. Trends in neurosciences.
2016;39(2):54-62. Epub 2016/01/10. doi: 10.1016/j.tins.2015.12.005. PubMed PMID:
26747654; PubMed Central PMCID: PMC4738085.
33. Kayser MS, Biron D. Sleep and Development in Genetically Tractable Model
Organisms. Genetics. 2016;203(1):21-33. Epub 2016/05/18. doi:
10.1534/genetics.116.189589. PubMed PMID: 27183564; PubMed Central PMCID:
PMC4858775.

34.	Sulston JE, Horvitz HR. Post-embryonic cell lineages of the nematode,
Caenorhabditis elegans. Dev Biol. 1977;56(1):110-56. Epub 1977/03/01. doi: 0012-
1606(77)90158-0 [pii]. PubMed PMID: 838129.
35.	Cassada RC, Russell RL. The dauerlarva, a post-embryonic developmental variant
of the nematode Caenorhabditis elegans. Dev Biol. 1975;46(2):326-42. Epub 1975/10/01.
doi: 0012-1606(75)90109-8 [pii]. PubMed PMID: 1183723.
36.	Han X, Chow BY, Zhou H, Klapoetke NC, Chuong A, Rajimehr R, et al. A high-
light sensitivity optical neural silencer: development and application to optogenetic
control of non-human primate cortex. Frontiers in systems neuroscience. 2011;5:18. Epub
2011/08/04. doi: 10.3389/fnsys.2011.00018. PubMed PMID: 21811444; PubMed Central
PMCID: PMC3082132.
37.	Okazaki A, Takagi S. An optogenetic application of proton pump ArchT to C.
elegans cells. Neuroscience research. 2013;75(1):29-34. Epub 2012/10/10. doi:
10.1016/j.neures.2012.09.002. PubMed PMID: 23044183.
38.	Alkema MJ, Hunter-Ensor M, Ringstad N, Horvitz HR. Tyramine Functions
independently of octopamine in the Caenorhabditis elegans nervous system. Neuron.
2005;46(2):247-60. Epub 2005/04/26. doi: S0896-6273(05)00167-4 [pii]
10.1016/j.neuron.2005.02.024. PubMed PMID: 15848803.
39.	Gordus A, Pokala N, Levy S, Flavell SW, Bargmann CI. Feedback from network
states generates variability in a probabilistic olfactory circuit. Cell. 2015;161(2):215-27.
Epub 2015/03/17. doi: 10.1016/j.cell.2015.02.018. PubMed PMID: 25772698; PubMed
Central PMCID: PMC4821011.
40.	Kawano T, Po MD, Gao S, Leung G, Ryu WS, Zhen M. An imbalancing act: gap
junctions reduce the backward motor circuit activity to bias C. elegans for forward
locomotion. Neuron. 2011;72(4):572-86. Epub 2011/11/22. doi:
10.1016/j.neuron.2011.09.005. PubMed PMID: 22099460.
41.	Zimmer M, Gray JM, Pokala N, Chang AJ, Karow DS, Marletta MA, et al.
Neurons detect increases and decreases in oxygen levels using distinct guanylate
cyclases. Neuron. 2009;61(6):865-79. Epub 2009/03/28. doi:
10.1016/j.neuron.2009.02.013. PubMed PMID: 19323996; PubMed Central PMCID:
PMCPMC2760494.
42.	Sawin ER, Ranganathan R, Horvitz HR. C. elegans locomotory rate is modulated
by the environment through a dopaminergic pathway and by experience through a
serotonergic pathway. Neuron. 2000;26(3):619-31. Epub 2000/07/15. doi: S0896-
6273(00)81199-X [pii]. PubMed PMID: 10896158.
43.	Zheng Y, Brockie PJ, Mellem JE, Madsen DM, Maricq AV. Neuronal control of
locomotion in C. elegans is modified by a dominant mutation in the GLR-1 ionotropic
glutamate receptor. Neuron. 1999;24(2):347-61. Epub 1999/11/26. PubMed PMID:
10571229.
44.	Cao J, Packer JS, Ramani V, Cusanovich DA, Huynh C, Daza R, et al.
Comprehensive single-cell transcriptional profiling of a multicellular organism. Science.
2017;357(6352):661-7. Epub 2017/08/19. doi: 10.1126/science.aam8940. PubMed
PMID: 28818938; PubMed Central PMCID: PMC5894354.
45.	Packer JSZ, Q., Huynh CS, P., Preston E, Dueck HS, D., Tan K, Trapnell C, et al.
A lineage-resolved molecular atlas of C. elegans embryogenesis at single cell resolution.
BioArxive. 2019;(8). doi: 10.1101/565549

46.   Chalfie M, Sulston JE, White JG, Southgate E, Thomson JN, Brenner S. The
neural circuit for touch sensitivity in Caenorhabditis elegans. J Neurosci. 1985;5(4):956-
64. Epub 1985/04/01. PubMed PMID: 3981252.
47.   Li W, Kang L, Piggott BJ, Feng Z, Xu XZ. The neural circuits and sensory
channels mediating harsh touch sensation in Caenorhabditis elegans. Nat Commun.
2011;2:315. Epub 2011/05/19. doi: 10.1038/ncomms1308. PubMed PMID: 21587232;
PubMed Central PMCID: PMC3098610.
48.   Turek M, Besseling J, Spies JP, Konig S, Bringmann H. Sleep-active neuron
specification and sleep induction require FLP-11 neuropeptides to systemically induce
sleep. eLife. 2016;5. Epub 2016/03/08. doi: 10.7554/eLife.12499. PubMed PMID:
26949257; PubMed Central PMCID: PMC4805538.
49.   Brockie PJ, Mellem JE, Hills T, Madsen DM, Maricq AV. The C. elegans
glutamate receptor subunit NMR-1 is required for slow NMDA-activated currents that
regulate reversal frequency during locomotion. Neuron. 2001;31(4):617-30. Epub
2001/09/08. PubMed PMID: 11545720.
50.   Singh K, Ju JY, Walsh MB, DiIorio MA, Hart AC. Deep Conservation of Genes
Required for Both Drosphila melanogaster and Caenorhabditis elegans Sleep Includes a
Role for Dopaminergic Signaling. sleep. 2014;37(9). PubMed PMID: 25142568.
51.   Schwarz J, Bringmann H. Reduced sleep-like quiescence in both hyperactive and
hypoactive mutants of the Galphaq Gene egl-30 during lethargus in Caenorhabditis
elegans. PLoS One. 2013;8(9):e75853. Epub 2013/09/28. doi:
10.1371/journal.pone.0075853. PubMed PMID: 24073282; PubMed Central PMCID:
PMC3779211.
52.   Huang H, Hayden DJ, Zhu CT, Bennett HL, Venkatachalam V, Skuja LL, et al.
Gap Junctions and NCA Cation Channels Are Critical for Developmentally Timed Sleep
and Arousal in Caenorhabditis elegans. Genetics. 2018;210(4):1369-81. Epub
2018/10/17. doi: 10.1534/genetics.118.301551. PubMed PMID: 30323068; PubMed
Central PMCID: PMCPMC6283151.
53.   Pirri JK, Alkema MJ. The neuroethology of C. elegans escape. Curr Opin
Neurobiol. 2012;22(2):187-93. Epub 2012/01/10. doi: 10.1016/j.conb.2011.12.007.
PubMed PMID: 22226513; PubMed Central PMCID: PMCPMC3437330.
54.   Nichols ALA, Eichler T, Latham R, Zimmer M. A global brain state underlies C.
elegans sleep behavior. Science. 2017;356(6344). Epub 2017/06/24. doi:
10.1126/science.aam6851. PubMed PMID: 28642382.
55.   Kaplan JM, Horvitz HR. A dual mechanosensory and chemosensory neuron in
Caenorhabditis elegans. Proc Natl Acad Sci U S A. 1993;90(6):2227-31. Epub
1993/03/15. PubMed PMID: 8460126.
56.   Guo ZV, Hart AC, Ramanathan S. Optical interrogation of neural circuits in
Caenorhabditis elegans. Nat Methods. 2009;6(12):891-6. Epub 2009/11/10. doi:
nmeth.1397 [pii]
10.1038/nmeth.1397. PubMed PMID: 19898486.
57.   Husson SJ, Costa WS, Wabnig S, Stirman JN, Watson JD, Spencer WC, et al.
Optogenetic analysis of a nociceptor neuron and network reveals ion channels acting
downstream of primary sensors. Curr Biol. 2012;22(9):743-52. Epub 2012/04/10. doi:
10.1016/j.cub.2012.02.066. PubMed PMID: 22483941; PubMed Central PMCID:
PMC3350619.

58.   Stirman JN, Crane MM, Husson SJ, Wabnig S, Schultheis C, Gottschalk A, et al.
Real-time multimodal optical control of neurons and muscles in freely behaving
Caenorhabditis elegans. Nat Methods. 2011;8(2):153-8. Epub 2011/01/18. doi:
10.1038/nmeth.1555. PubMed PMID: 21240278; PubMed Central PMCID:
PMC3189501.
59.   Kato S, Kaplan HS, Schrodel T, Skora S, Lindsay TH, Yemini E, et al. Global
brain dynamics embed the motor command sequence of Caenorhabditis elegans. Cell.
2015;163(3):656-69. Epub 2015/10/20. doi: 10.1016/j.cell.2015.09.034. PubMed PMID:
26478179.
60.   Schwarz J, Lewandrowski I, Bringmann H. Reduced activity of a sensory neuron
during a sleep-like state in Caenorhabditis elegans. Curr Biol. 2011;21(24):R983-4. Epub
2011/12/24. doi: S0960-9822(11)01207-3[pii]10.1016/j.cub.2011.10.046. PubMed
PMID: 22192827.
61.   Cho JY, Sternberg PW. Multilevel modulation of a sensory motor circuit during
C. elegans sleep and arousal. Cell. 2014;156(1-2):249-60. Epub 2014/01/21. doi:
10.1016/j.cell.2013.11.036. PubMed PMID: 24439380; PubMed Central PMCID:
PMC3962823.
62.   Skora S, Mende F, Zimmer M. Energy Scarcity Promotes a Brain-wide Sleep
State Modulated by Insulin Signaling in C. elegans. Cell reports. 2018;22(4):953-66.
Epub 2018/02/02. doi: 10.1016/j.celrep.2017.12.091. PubMed PMID: 29386137.
63.   Katz M, Corson F, Iwanir S, Biron D, Shaham S. Glia Modulate a Neuronal
Circuit for Locomotion Suppression during Sleep in C. elegans. Cell reports.
2018;22(10):2575-83. Epub 2018/03/08. doi: 10.1016/j.celrep.2018.02.036. PubMed
PMID: 29514087; PubMed Central PMCID: PMCPMC5870883.
64.   Schrodel T, Prevedel R, Aumayr K, Zimmer M, Vaziri A. Brain-wide 3D imaging
of neuronal activity in Caenorhabditis elegans with sculpted light. Nat Methods.
2013;10(10):1013-20. Epub 2013/09/10. doi: 10.1038/nmeth.2637. PubMed PMID:
24013820.
65.   Gallagher T, Kim J, Oldenbroek M, Kerr R, You YJ. ASI regulates satiety
quiescence in C. elegans. J Neurosci. 2013;33(23):9716-24. Epub 2013/06/07. doi:
10.1523/JNEUROSCI.4493-12.2013. PubMed PMID: 23739968; PubMed Central
PMCID: PMCPMC3726252.
66.   Goetting DL, Soto R, Van Buskirk C. Food-Dependent Plasticity in
Caenorhabditis elegans Stress-Induced Sleep Is Mediated by TOR-FOXA and TGF-beta
Signaling. Genetics. 2018. Epub 2018/06/22. doi: 10.1534/genetics.118.301204. PubMed
PMID: 29925566.
67.   Spies J, Bringmann H. Automated detection and manipulation of sleep in C.
elegans reveals depolarization of a sleep-active neuron during mechanical stimulation-
induced sleep deprivation. Sci Rep. 2018;8(1):9732. Epub 2018/06/29. doi:
10.1038/s41598-018-28095-5. PubMed PMID: 29950594.
68.   Scholz M, Linder AN, Randi F, Sharma AK, Yu X, Shaevitz JW, et al. Predicting
natural behavior from whole-brain neural dynamics. Biorxive. 2018:445643v1. Epub
October 17, 2018. doi: https://doi.org/10.1101/445643
69.   Chrousos G, Vgontzas AN, Kritikou I. HPA Axis and Sleep. In: De Groot LJ,
Chrousos G, Dungan K, Feingold KR, Grossman A, Hershman JM, et al., editors.
Endotext. South Dartmouth (MA): Endotext.org; 2016.

70.   Panossian LA, Avidan AY. Review of sleep disorders. The Medical clinics of
North America. 2009;93(2):407-25, ix. Epub 2009/03/11. doi:
10.1016/j.mcna.2008.09.001. PubMed PMID: 19272516.
71.   Liu S, Liu Q, Tabuchi M, Wu MN. Sleep Drive Is Encoded by Neural Plastic
Changes in a Dedicated Circuit. Cell. 2016;165(6):1347-60. Epub 2016/05/24. doi:
10.1016/j.cell.2016.04.013. PubMed PMID: 27212237; PubMed Central PMCID:
PMC4892967.
72.   Donlea JM, Pimentel D, Talbot CB, Kempf A, Omoto JJ, Hartenstein V, et al.
Recurrent Circuitry for Balancing Sleep Need and Sleep. Neuron. 2018;97(2):378-89 e4.
Epub 2018/01/09. doi: 10.1016/j.neuron.2017.12.016. PubMed PMID: 29307711;
PubMed Central PMCID: PMCPMC5779612.
73.   Dijk DJ. Regulation and functional correlates of slow wave sleep. Journal of
clinical sleep medicine : JCSM : official publication of the American Academy of Sleep
Medicine. 2009;5(2 Suppl):S6-15. Epub 2009/12/17. PubMed PMID: 19998869; PubMed
Central PMCID: PMC2824213.
74.   Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974;77(1):71-94.
Epub 1974/05/01. PubMed PMID: 4366476.
75.   Ahringer A. Reverse genetics. WormBook, ed The C elegans Research
Community. 2006. doi: doi/10.1895/wormbook.1.47.1.
76.   Merritt C, Seydoux G. Transgenic solutions for the germline. WormBook.
2010:1-21. Epub 2010/02/20. doi: 10.1895/wormbook.1.148.1. PubMed PMID:
20169625.
77.   Redemann S, Schloissnig S, Ernst S, Pozniakowsky A, Ayloo S, Hyman AA, et
al. Codon adaptation-based control of protein expression in C. elegans. Nat Methods.
2011;8(3):250-2. Epub 2011/02/01. doi: nmeth.1565 [pii]10.1038/nmeth.1565. PubMed
PMID: 21278743.
78.   Praitis V, Casey E, Collar D, Austin J. Creation of low-copy integrated transgenic
lines in Caenorhabditis elegans. Genetics. 2001;157(3):1217-26. Epub 2001/03/10.
PubMed PMID: 11238406; PubMed Central PMCID: PMC1461581.
79.   Wilm T, Demel P, Koop HU, Schnabel H, Schnabel R. Ballistic transformation of
Caenorhabditis elegans. Gene. 1999;229(1-2):31-5. Epub 1999/03/30. PubMed PMID:
10095101.
80.   Bringmann H. Agarose hydrogel microcompartments for imaging sleep- and
wake-like behavior and nervous system development in Caenorhabditis elegans larvae. J
Neurosci Methods. 2011;201(1):78-88. Epub 2011/08/02. doi: S0165-0270(11)00418-3
[pii]10.1016/j.jneumeth.2011.07.013. PubMed PMID: 21801751.
81.   Turek M, Besseling J, Bringmann H. Agarose Microchambers for Long-term
Calcium Imaging of Caenorhabditis elegans. Journal of visualized experiments : JoVE.
2015;(100):e52742. Epub 2015/07/02. doi: 10.3791/52742. PubMed PMID: 26132740;
PubMed Central PMCID: PMC4544933.
82.   Stiernagle T. Maintenance of C. elegans. WormBook. 2006:1-11. Epub
2007/12/01. doi: 10.1895/wormbook.1.101.1. PubMed PMID: 18050451.
83.   Kim E, Sun L, Gabel CV, Fang-Yen C. Long-term imaging of Caenorhabditis
elegans using nanoparticle-mediated immobilization. PLoS One. 2013;8(1):e53419. Epub
2013/01/10. doi: 10.1371/journal.pone.0053419. PubMed PMID: 23301069; PubMed
Central PMCID: PMCPMC3536676.

84.   Nagy S, Raizen DM, Biron D. Measurements of behavioral quiescence in
Caenorhabditis elegans. Methods. 2014. Epub 2014/03/20. doi:
10.1016/j.ymeth.2014.03.009. PubMed PMID: 24642199.
879

880 **Figures**

Figure 1

Figure 1

Presynaptic neurons control the activity of the sleep-active RIS neuron

A) Sample trace of RIS activity and worm locomotion behavior outside of and in
lethargus. RIS is relatively inactive outside of lethargus and shows strong activity
transients in lethargus. Upon RIS activation, worms enter sleep bouts.

B) Presynaptic neurons activate RIS outside of and in lethargus. For statistical
calculations neural activities before the stimulation period (0 - 0.95 min) were
compared to activity levels during the stimulation period (1 - 1.95 min); *p <
0.05, **p < 0.01, ***p < 0.001, Wilcoxon Signed Rank Test.

C) PVC is essential for lethargus-specific RIS activation; Statistical calculations
were performed as described in B, but in experiments, in which SDQL was
stimulated baseline activity levels were calculated over the time interval from 0.6
- 0.95 min; *p < 0.05, Wilcoxon Signed Rank Test.

D) PVC hyperpolarization inactivates RIS and induces behavioral activity; **p <
0.01, ***p < 0.001, Wilcoxon Signed Rank Test for GCaMP and speed, Fisher's
Exact Test for sleep fraction.

E) Circuit model of the RIS presynaptic regulatory network. Presynaptic neurons
present mostly activators of RIS. PVC is essential for lethargus-specific RIS
activation. Effects of presynaptic neurons might be mostly direct, except for RIM.

Figure 2

A

1. baseline PVC GCaMP
2. RIS depolarization/hyperpolarization
   simultaneously imaging PVC GCaMP
3. recovery PVC GCaMP

B

C

D

F

E

Figure 2

RIS and PVC activate each other forming a positive feedback loop

A) The experimental set up. First, a baseline of PVC was measured through calcium
imaging. Then, RIS was optogenetically manipulated while PVC was
simultaneously calcium imaged. Finally, a recovery period was imaged of PVC.

B) RIS depolarization leads to a strong PVC depolarization outside and during
lethargus; $*p < 0.05$, $**p < 0.01$, Wilcoxon Signed Rank Test.

C) RIS hyperpolarization leads to a weak PVC depolarization outside and in
lethargus. For statistical calculations neural activities before the stimulation
period (0 - 1 min) were compared to activity levels during the stimulation period
(1 - 2 min); $*p < 0.05$, $**p < 0.01$, compared before and during stimulation,
Wilcoxon Signed Rank Test.

D) RIS does not reach the same activation levels in *aptf-1(gk794)* mutants compared
to wild-type worms. *aptf-1(gk794)* mutants do neither immobilize nor sleep
during RIS activation; $***p < 0.001$, Welch Test.

E) *aptf-1(gk794)* mutants do not show sleep during lethargus; $**p < 0.01$,
Kolmogorov-Smirnov-Test.

F) A circuit model for the positive feedback loop between RIS and PVC. During
wakefulness, reverse command interneurons inhibit PVC so that PVC does not
activate RIS. In lethargus, PVC is directly activating RIS, which then inhibits
reverse command interneurons. This may speculatively disinhibit PVC, leading to
a positive feedback.

Figure 3

Figure 3

RIS depolarization is homeostatically controlled

A)   RIS shows rebound activation following hyperpolarization. Behavioral and brain

activity measurements correlate throughout the whole experiment; $*p < 0.05$, $**p$

< 0.01, $***p < 0.001$, Wilcoxon Signed Rank Test for GCaMP and speed,

Fisher's Exact Test for sleep fraction.

B)   Dose-response curve of optogenetic RIS hyperpolarization with different stimulus

lengths. RIS activation rebound transients saturate with increasing length of

inhibition. Worms not showing a rebound activation transient after RIS

optogenetic hyperpolarization were excluded from the analysis.

C)   RIS shows a negative rebound following its optogenetic depolarization; $*p < 0.05$,

$**p < 0.01$, $***p < 0.001$, Wilcoxon Signed Rank Test for GCaMP and speed,

Fisher's Exact Test for sleep fraction.

Figure 4

Figure 4

RIM activity levels are dampened in lethargus independently of RIS

A) Sample traces of RIM transient frequencies, RIM baseline activities and worm locomotion behaviors outside of and in lethargus in wild-type worms and *aptf-1(gk794)* mutants.

B) Wild type worms, but not *aptf-1(gk974)* mutants display changes in RIM transient frequencies across lethargus. Transient frequencies were assessed manually. To be counted as a transient, RIM activity levels had to be at least twice as high as baseline activity levels. ***p < 0.001 Kolmogorov-Smirnov-Test for wild-type condition, Welch Test for mutant condition.

C) The reduction of RIM baseline activity levels in lethargus is preserved in *aptf-1(gk974)* mutants; **p < 0.01, Wilcoxon Signed Rank Test.

Figure 5

Figure 5

Arousing stimulation inhibits RIS and sleep through RIM

A) ASH depolarization in wild-type *C. elegans* leads to mobilization, awakening and
RIS inhibition; $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, Wilcoxon Signed Rank Test
for GCaMP and speed, Fisher's Exact Test for sleep fraction.

B) ASH depolarization in RIM ablated *C. elegans* leads to weaker mobilization,
awakening and RIS activation; $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, Wilcoxon
Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction.

C) The response direction following ASH activation in wild-type worms is
predominantly backward, while in RIM ablated worms it is predominantly
forward. This suggests that the forward direction is connected to RIS activation;
$***p < 0.001$, Fisher's Exact Test.

D) A speculative circuit model for RIS regulation through arousal by ASH. RIM
serves as synchronizer of AVE and AVA to regulate PVC and therefore RIS
inhibition.

Figure 6

A) the circuit for sleep control in *C. elegans*

981 **Figure 6**

982 **Model of the tripartite flip-flop sleep switch**

983 A) Model of the *C. elegans* sleep switch that is consistent with our data explaining the

984 switching of sleep states.