1      **Title:**

4      **A wake-active locomotion circuit depolarizes a sleep-active neuron to switch on**

5                                    **sleep**

10     **Authors:** Elisabeth Maluck†[1,2], Inka Busack†[1,2], Judith Besseling[1], Florentin Masurat[1],

11                 Michal Turek[1], Karl Emanuel Busch[3], Henrik Bringmann*[1,2]

**Affiliations:**

[1] Max Planck Institute for Biophysical Chemistry, Göttingen, Germany

[2] University of Marburg, Marburg, Germany

[3] University of Edinburgh, United Kingdom

* henrik.bringmann@biologie.uni-marburg.de

†these authors contributed equally

**Abstract**

Sleep-active neurons depolarize during sleep to suppress wakefulness circuits. Wake-

active wake-promoting neurons in turn shut down sleep-active neurons, thus forming a

bipartite flip-flop switch. However, how sleep is switched on is unclear because it is not

known how wakefulness is translated into sleep-active neuron depolarization when the

system is set to sleep. Using optogenetics in *C. elegans*, we solved the presynaptic circuit

for depolarization of the sleep-active RIS neuron during developmentally-regulated sleep,

also known as lethargus. Surprisingly, we found that RIS activation requires neurons that

have known roles in wakefulness and locomotion behavior. The RIM interneurons, which

are active during and can induce reverse locomotion, play a complex role and can act as

inhibitors of RIS when they are strongly depolarized and as activators of RIS when they

are modestly depolarized. The PVC command interneurons, which are known to promote

forward locomotion during wakefulness, act as major activators of RIS. The properties of

these locomotion neurons are modulated during lethargus. The RIMs become less

excitable. The PVCs become resistant to inhibition and have an increased capacity to

activate RIS. Separate activation of neither the PVCs nor the RIMs appears to be

sufficient for sleep induction; instead, our data suggests that they act in concert to

activate RIS. Forward and reverse circuit activity is normally mutually exclusive. Our

data suggest that RIS may be activated at the transition between forward and reverse

locomotion states, perhaps when both reverse (including RIM) and forward (PVC) circuit

activity overlaps. While RIS is not strongly activated outside of lethargus, altered activity

of the locomotion interneurons during lethargus favors strong RIS activation and thus

sleep. The control of sleep-active neurons by locomotion circuits suggests that sleep may

have evolved from locomotion quiescence. The flip-flop sleep switch in *C. elegans* thus

requires an additional component, wake-active sleep-promoting neurons that translate

wakefulness into the depolarization of a sleep-active neuron when the worm is sleepy.

Wake-active sleep-promoting circuits may also be required for sleep state switching in

other animals including in mammals.

## Introduction

Sleep is a behavior that affects many, if not all, physiological processes. Disorders and curtailment of sleep affect the lives of 10-30% of the adult population of modern societies. Sleep loss is associated with an increased risk of infection [1], cardiovascular disease [1], psychiatric disease including depression [2,3], obesity [4,5], type 2 diabetes [4,5], and cancer [1]. The high prevalence of insomnia and insufficient sleep quality thus presents a massive unmet health and economic problem [1,3-5]. To understand how sleep behavior is generated, it is crucial to solve the underlying neural circuits.

Sleep circuits require inhibitory sleep-active sleep-promoting neurons, which depolarize specifically at sleep onset and actively induce sleep by releasing inhibitory neurotransmitters, GABA and neuropeptides, to dampen arousal and the activity of wake circuits [6]. Sleep behavior induced by inhibitory sleep-active neurons includes the suppression of voluntary movements and sensory perception, reversibility, and homeostasis [7]. Inhibitory sleep-active neurons suppress wake circuits and can be rapidly suppressed by arousing stimulation to allow for quick awakening. Forced wakefulness is followed by an increase of sleep-active neuron depolarization, which leads to homeostatic sleep corrections. Thus, understanding sleep control requires comprehension of the circuit mechanisms that determine when and how much inhibitory sleep-active neurons depolarize [6,8].

Circuits control the depolarization of inhibitory sleep-active neurons. For example, wake-active wake-promoting neurons promote arousal and suppress inhibitory sleep-active neurons, whereas sleep need causes sleep-active neuron depolarization. Thus, sleep-active sleep-promoting and wake-active wake-promoting neurons form a flip-flop switch, which ensures that sleep and wake exist as discrete states. This sleep switch is under the control of arousal that favors wake and inhibits sleep through the suppression of sleep-active neurons by inhibitory wake-active neurons [6,9]. It has been proposed that sleep induction is favored by disinhibition of inhibitory sleep-active neurons [10-12], and also excitatory sleep-active neurons exist that might perhaps present activators of inhibitory

sleep-active neurons [13]. However, the forces and mechanisms that flip the sleep switch

from wake to sleep when an organism gets sleepy cannot be satisfactorily explained by

the present circuit models as it is unclear how sleep-active neurons are turned on when

the system is set to sleep.

Sleep is under circadian and homeostatic controls that determine the timing of sleep and

ensure that enough of this essential physiological state takes place [14]. Sleep

homeostasis comprises multiple mechanisms that act on different time scales. On long

time scales sleep is a function of prior wakefulness, i.e. prolonged wakefulness leads to

increased sleep propensity and sleep loss triggers compensatory increases in the intensity

or duration of sleep. This chronic sleep homeostasis likely is mediated by several parallel

mechanisms. For example, in mammals, somnogens such as adenosine accumulate during

wakefulness leading to the inhibition of wake-promoting neurons [15,16]. In *Drosophila*,

activity-dependent plasticity of sleep-promoting neurons increases during wakefulness to

increase subsequent sleep [17,18]. On short time scales, acute homeostasis determines

whether the system's actual state matches the system's set point and carries out corrective

action if those values do not match. For example, to homeostatically maintain sleep

despite disturbance, micro arousals need to be compensated for. In humans, homeostatic

sleep maintenance can be seen in EEG recordings in the form of k-complexes, where a

spontaneous or evoked short cortical up state is followed by a down state [19-21].

Homeostatic sleep maintenance is also found during sleep in *C. elegans,* where sleep

bouts are interrupted by short motion bouts, with the length of a motion bout correlating

with the length of the subsequent sleep bout [22,23]. Thus, across systems, homeostatic

sleep maintenance requires constant surveillance of sleep and corrective action.

Sleep-active sleep-promoting neurons are conserved regulators of sleep and have been

found both in vertebrates as well as in invertebrates [8,24]. Mammals possess several

populations of sleep-active neurons, most of which are inhibitory, across the brain. These

neurons reside in the anterior hypothalamus, brain stem, and cortex [6,12]. Excitatory

sleep-active neurons were found in the periocular midbrain that project to inhibitory

sleep-active neurons in the anterior hypothalamus, whose role could be to activate

inhibitory sleep-active neurons [13]. Studying sleep in less complex brains facilitates

sleep circuit analysis. In *Drosophila*, sleep-promoting neurons are found at several

locations in the brain. A well-characterized population of sleep-promoting neurons is

formed by ExFl1 neurons residing in the dorsal fan-shaped body. R2 neurons accumulate

homeostatic sleep pressure over time to promote activation of ExFl1 neurons, probably

by an indirect mechanism [17,18]. *C. elegans* possesses a single inhibitory sleep-active

neuron called RIS. Like its mammalian counterparts, RIS depolarizes at sleep onset. RIS

is crucial for sleep induction as its ablation leads to a virtually complete loss of detectable

sleep bouts [25-27]. The small, invariant nervous system, its mapped connectome, and

the transparency of *C. elegans* facilitates neural circuit analysis [28]. However, the

specific neural circuits that control RIS activity are not yet understood.

*C. elegans* shows sleep behavior during many stages and conditions. Here we analyzed

sleep behavior during development, also known as lethargus, the stage prior to each of

the four molts during larval development [8,27,29-31]. We used optogenetics to dissect

the neural circuits that control the activation of the sleep-active RIS neuron in *C. elegans*.

We found a third and novel important element of the flip-flop switch: Interneurons that

are active during wakefulness and that are known to control locomotion, are required for

RIS activation and sleep. These findings suggest a tripartite flip-flop circuit model that

can explain how arousing stimulation inhibits RIS depolarization, how RIS

depolarization is homeostatically controlled, and how reduced arousal can induce RIS

depolarization. Our RIS circuit model has two important implications for understanding

sleep control: 1) It suggests that sleep control has evolved from circuits controlling

locomotion quiescence. 2) Sleep induction requires an important third element, wake-

active sleep-promoting neurons, that translate wakefulness into sleep when the animal is

sleepy but awake.

**Results**

**Interneurons known to govern locomotion behavior control RIS activity**

RIS is crucially required for sleep and typically activates during sleep bouts (Fig 1A) [25]. However, the presynaptic driver neurons that activate and control this neuron are not known. To identify the circuits controlling RIS activation, we optogenetically tested the role of neurons that are presynaptic to RIS according to the *C. elegans* connectome [28]. The neurons AVJL, CEPDL, URYVL, RIMR, PVCL, and SDQL have been shown to be presynaptic to RIS [28,32]. To find out how these presynaptic neurons control RIS, we activated them with ReaChR and green light, and followed RIS calcium activity during and outside of lethargus. We confirmed the expression of ReaChR through a fused fluorescent reporter (mKate2). AVJ, CEPD, URYV, RIM, PVC, and SDQ each are a pair of two neurons, of which only one is presynaptic to RIS. As only promoters are available that express in both neurons of each pair, and as the two neurons of each pair are in close proximity, we always manipulated both neurons of the neuronal pair (except for SDQL) [28,32]. Because there were no specific promotors available for the expression in SDQL and PVC, we expressed ReaChR using semi-specific promoters and selectively illuminated only the presynaptic neuron class. We used L1 larvae for most of the optogenetic experiments to dissect the circuit. As SDQL is born postembryonically and likely is not yet functional during the L1 stage, we used L4 larvae to assay its function [33]. We compared the effects of optogenetic stimulation outside and during lethargus, defined as the period prior to the molt during which the animals do not feed [34]. Before lethargus, we measured an activation of RIS upon depolarization of AVJ, CEP and SDQL. During lethargus, the activation of CEP, PVC and SDQL caused RIS activation (Figs 1B and S1A Fig).

**Fig 1**

**Presynaptic neurons control the activity of the sleep-active RIS neuron**

A)      Sample trace of RIS activity and worm locomotion behavior outside of and during
        lethargus. RIS has no strong calcium transients outside of lethargus but shows
        strong activity transients during lethargus. Upon RIS activation, worms enter
        sleep bouts. (S1 Data, Sheet 1A)

B)      Presynaptic neurons activate or inhibit RIS outside of and during lethargus. For
        statistical calculations neural activities before the stimulation period (0 - 0.95min)

174           were compared to activity levels during the stimulation period (1 - 1.95min). *p <

175           0.05, **p < 0.01, ***p < 0.001, Wilcoxon Signed Rank Test. (S1 Data, Sheet 1B)

176   C)      RIS activity decreases upon optogenetic PVC and RIM hyperpolarization.

177           Statistical calculations were performed as described in B, but in experiments, in

178           which SDQL was stimulated baseline activity levels were calculated over the time

179           interval from 0.6 - 0.95min. Baseline activity levels were calculated starting from

180           0.6min as baseline activity levels were instable before that time point. *p < 0.05,

181           **p < 0.01, Wilcoxon Signed Rank Test. (S1 Data, Sheet 1C)

182   D)      Circuit model of the RIS presynaptic regulatory network. Activating synaptic

183           input is shown as green arrows, inhibitory synaptic input is shown as red arrows,

184           and unclear synaptic input is shown as black arrow. Gap junctions are indicated as

185           black connections. Neurons that are presynaptic to RIS present mostly activators.

186           PVC is essential for lethargus-specific RIS activation. RIM can inhibit RIS

187           through tyramine and FLP-18 and can activate RIS with glutamate.

All neurons showed consistent effects on RIS depolarization except RIM. RIM is known

to play complex roles in controlling behavior and is involved in seemingly opposing

behaviors. For example, specific RIM activation can trigger a reversal [35], whereas RIM

inhibition has been suggested to be required for reversals through an alternative circuit

[36]. We performed optogenetic depolarization experiments of RIM expressing ReaChR

using two different promoters, the *tdc-1* promoter, which is known to express strongly,

and the *gcy-13* promoter, which is known to express at a lower level [37]. Activation of

RIM with Channelrhodopsin expressed from the *tdc-1* promoter has previously been

shown to cause reversals [35], and we observed that activation of RIM using ReaChR

expressed from this promoter led to RIS inhibition (Fig 1B, RIM panel). The *tdc-1*

promoter expresses strongly in RIM, but also weakly in RIC [38]. To test whether the

inhibitory effect of *tdc-1* promoter-driven ReaChR expression on RIS was caused by

RIC, we also specifically expressed ReaChR in RIC using the *tbh-1* promoter [38].

Specific RIC activation led to RIS activation rather than inhibition (S1B Fig). Therefore,

the *tdc-1::ReaChR*-mediated RIS inhibition appears to stem from RIM activation.

Activating RIM with the weaker *gcy-13* promoter did not cause any net effects on RIS
when all trials were averaged (S1C Fig). Visual inspection of the individual trials,
however, showed that RIM activation could either inhibit or activate RIS. We hence
sorted single trials for the *gcy-13* experiment into two classes, in which RIM either
activated or inhibited RIS function (S1D Fig). The activation or inhibition of RIS by RIM
was indistinguishable during the beginning or end of lethargus (S1E Fig).

To confirm that RIM can both activate and inhibit RIS, we tested whether activation and
inhibition are mediated by different neurotransmitters. We tested the effects of RIM
activation on RIS in mutants, which lack transmitters that are known to be expressed in
RIM. The RIM neurons are well known to inhibit downstream neurons using tyramine,
which requires the *tdc-1* gene [38], and also express neuropeptides encoded by the *flp-18*
gene [39]. To test whether RIM can inhibit RIS using these known transmitters, we
analyzed mutant worms that lack functional *flp-18* and *tdc-1*. Individual inactivation of
*flp-18* and *tdc-1* reduced, and double mutation abolished inhibition of RIS by RIM (S2
Fig). Therefore, the transmitters tyramine and FLP-18 are together responsible for RIS
inhibition by RIM. We next tested activation of RIS by RIM in *eat-4(ky5)* mutant larvae,
which lack glutamatergic signaling in many neurons including RIM [40,41]. RIS
activation by RIM activation was completely gone in *eat-4(ky5)* mutant larvae (S3 Fig,
we used L4 larvae for this assay as the response was more robust). Therefore, glutamate
is required for RIS activation by RIM. Together these results suggest that RIM can act
both as an activator as well as an inhibitor of RIS by employing different
neurotransmitters, with weaker activation allowing for RIS activation and stronger
activation favoring inhibition.

The majority of synaptic inputs into RIS that we studied had activating effects, the sole
inhibitory effect was observed after strong activation of RIM, whereas weaker RIM
activation could also lead to RIS activation. The CEP, URY, and SDQL neurons present
sensory receptors and might play a role in activating RIS in response to stimulation. For
example, CEP might activate RIS as part of the basal slowing response [42,43]. The
PVCs appeared to be strong activators of RIS specifically during lethargus. This

suggested that either the PVC to RIS connection might be specific to lethargus or that it has not yet matured during the mid L1 stage. We hence repeated the experiment and activated PVC in L2 larvae. PVC activated RIS both during and outside of lethargus, but the activation during lethargus was much stronger, suggesting that the activation of RIS by PVC is strongly enhanced during lethargus (S4 Fig).

To find out which presynaptic neurons are required for inhibition or activation of RIS during lethargus, we tested the effect of optogenetic inhibition of the presynaptic neurons on RIS activation. We used ArchT, which hyperpolarizes neurons by pumping protons out of the cell [44,45]. As above, we verified the expression of ArchT in neurons of interest by using an mKate2-tagged version. As in the ReaChR experiments, we specifically illuminated each presynaptic neuron class and quantified RIS activation using calcium imaging. Before lethargus, inhibition of AVJ and PVC led to an inhibition of RIS, whereas inhibition of the other neurons tested had no acute statistically significant effect on RIS (Optogenetic RIM hyperpolarization using the stronger *tdc-1* promoter in worms outside of lethargus showed a tendency to inhibit RIS function, p-value of 0.0539 N = 11 animals, whereas the weaker *gcy-13* promoter had no detectable effect). During lethargus, optogenetic inhibition of PVC and RIM (using the stronger *tdc-1* promoter) led to significant RIS inhibition, whereas there was no effect seen for the other neurons (Fig 1C and S5A Fig, inhibition of RIM using the weaker *gcy-13* promoter only produced a tendency but no statistically significant net effect, S5B and S5C Figs).

Absence of an effect of optogenetic inhibition of presynaptic neurons could mean that either these neurons are not required for RIS activation, that the inhibition was not strong enough, or that they may act redundantly (we did not find any evidence for redundancy at least for CEP and URY, S5D Fig). Our optogenetic analysis revealed a complex set of presynaptic inputs for regulation of RIS activity (Fig 1D). The optogenetic depolarization experiments suggest that CEP, PVC, RIM and SDQL present the most potent presynaptic activators of RIS. The capacity of PVC to activate RIS is strongly increased during lethargus, indicating that this neuron is involved in the lethargus-specific activation of RIS. The optogenetic hyperpolarization experiments suggest that PVC and RIM are

essential presynaptic activators of RIS during lethargus. Therefore, we focused our
analysis on PVC and RIM neurons.

**PVC becomes resistant to inhibition during lethargus**

Neuronal activation and silencing experiments revealed PVC as a main activator of RIS.
These results predict that neuronal activity of PVC should correlate with RIS activation
and sleep bouts. To test for such correlation, we measured the activity of both neurons
simultaneously. Because the calcium transients observable in PVC are typically small
[46] and could not be detected in our assays in mobile worms (data not shown), we
immobilized the larvae and used RIS activation as a proxy for sleep bouts. We extracted
both RIS and PVC activity and aligned all data to the RIS activation maxima. This
analysis showed that PVC activated approximately one minute earlier than RIS and
reached its maximum activation approximately 1.5 minutes earlier than RIS. PVC
activity decreased slowly during the RIS transient (Fig 2A). This result is consistent with
a role for PVC in promoting RIS depolarization.

**Fig 2**
**PVC is an RIS activator that becomes resistant to inhibition during lethargus but**
**PVC activation is not sufficient for sleep induction**
A)  Simultaneous PVC and RIS GCaMP traces aligned to RIS peaks in fixed L1
lethargus worms. PVC activates before the RIS peak and stays active until the
peak. *p < 0.05, **p < 0.01, Wilcoxon Signed Rank Test. (S1 Data, Sheet 2A)
B)  PVC hyperpolarization inactivates RIS and induces behavioral activity. PVC
hyperpolarization was performed by expressing ArchT under the *zk637.11*
promoter. In contrast to the *nmr-1* promoter, the *zk637.11* promoter lacks
expression in head command interneurons. **p < 0.01, ***p < 0.001, Wilcoxon
Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction.
(S1 Data, Sheet 2B)
C)  During lethargus, PVC becomes resistant to inhibition. Outside of lethargus its
inhibition is stronger and continues beyond the end of optogenetic stimulation.

During lethargus PVC activity levels return back to baseline already during the
stimulation period. *p < 0.05, **p < 0.01, Wilcoxon Signed Rank Test. (S1 Data,
Sheet 2C)
D) PVC activation translates into mostly a forward mobilization in L1 lethargus.
*p < 0.05, ***p < 0.001, Wilcoxon Signed Rank Test for Speed. Fisher's Exact
Test for fraction of direction. (S1 Data, Sheet 2D)
304
PVC inhibition reduced RIS activity in immobilized animals, but it is unclear how PVC
inhibition affects behavior. To be able to test the effects of PVC inhibition on behavior
without affecting the other command interneurons, we chose a more specific promoter for
expression in PVC from single-cell RNA sequencing data. There was no gene in the
available datasets that was expressed only in the cluster of cells containing PVC, but the
previously uncharacterized gene *zk673.11* was expressed specifically in PVC and in only
a few other neurons excluding other command interneurons [47,48](Jonathan Packer,
personal communication, S6 Fig). Hyperpolarization of PVC using ArchT driven by the
*zk673.11* promoter led to an acute inhibition of RIS, an increase in locomotion, and a
reduction of sleep (Fig 2B). Hyperpolarization of PVC using ArchT also strongly
inhibited RIS outside of lethargus. This experiment confirmed the role of the PVCs in
activating RIS.
317
Hyperpolarization of PVC outside of lethargus appeared to have a stronger and longer
lasting effect on RIS inhibition compared with during lethargus (Figs 1C and 2B). This is
surprising as PVC is a stronger activator of RIS during lethargus in the optogenetic
activation experiments (Fig 1B). This effect could be explained if PVC responded more
severely to inhibition outside of lethargus. We tested this idea by inhibiting PVC using
ArchT and green light and simultaneously imaged PVC activity. PVC hyperpolarization
was stronger in worms outside of lethargus and PVC remained inhibited after the
optogenetic manipulation. During lethargus, PVC was only weakly inhibited at the
beginning of optogenetic stimulation and returned to baseline levels already during the
stimulation (Fig 2C). We also tested whether optogenetic excitability of PVC was
modulated during lethargus but could not find any differences in excitability of PVC

during or outside of lethargus (S7A Fig). Thus, PVC is more susceptible to inhibition outside of lethargus but becomes resistant to inhibition during lethargus. This effect can explain the stronger hyperpolarization of RIS during PVC inhibition outside of lethargus and this effect likely presents an important modulation of the circuit to favor PVC activation and thus RIS activation during lethargus.

PVC is known to promote forward movement upon posterior mechanical stimulation and optogenetic stimulation of PVC in adults has been shown to promote forward locomotion [49,50]. Our data showed that PVC also activates the RIS neuron and consistent with this observation mechanical stimulation caused RIS activation (S7B Fig). This suggested that PVC activates RIS to modulate forward locomotion speed and to promote sleep. However, it is unclear how PVC can promote forward motion and sleep, as these are two seemingly opposing behaviors. We hence tested whether optogenetic stimulation of PVC in larvae induces sleep behavior. We activated PVC using *nmr-1::ReaChR* in mobile L1 larvae during lethargus and specifically illuminated the tail of the animal, which contains the cell bodies of the PVC neurons but not the other *nmr-1*-expressing neurons. We quantified the speed as well as the direction of movement of the worm. During PVC activation during lethargus, the worms visibly accelerated movement and mostly crawled forward, but we could not see induction of sleep behavior during optogenetic stimulation (Fig 2D). Consistent with this finding, optogenetic PVC activation during and before lethargus always led to the activation of AVB interneurons, which are known to be premotor neurons required for forward locomotion [49] (S7C and S7D Figs). Together these experiments showed that PVC activates prior to RIS and is required for RIS activation. However, its activation alone does not seem to be sufficient to induce sleep behavior.

**RIS and PVC activate each other forming a positive feedback loop**

PVC presents a major activator of RIS but how a forward command interneuron can cause strong and state specific activation of the RIS neuron during sleep bouts is not clear. We hence tested how optogenetic RIS activation affects PVC activity. We

selectively activated RIS using ReaChR and measured calcium activity in PVC in
immobilized animals. Upon RIS stimulation, PVC immediately displayed unexpectedly
strong calcium transients, which were slightly stronger during lethargus (Fig 3A and S8A
Fig). These results show that PVC and RIS activate each other, thus forming a positive
feedback loop. The sleep-inducing RIS neuron has so far only been shown to inhibit other
neurons making PVC the first neuron that is not inhibited but activated by RIS. For
example, command interneurons such as AVE and AVA and other neurons are not
activated but inhibited by RIS [25].

**Fig 3**

**RIS and PVC activate each other forming a positive feedback loop**

A-E) RIS depolarization leads to a strong PVC depolarization outside of and during

lethargus. This PVC depolarization is almost abolished in *flp-11(tm2705)* and it is

significantly reduced in AVE-ablated worms. *p < 0.05, **p < 0.01, Wilcoxon

Signed Rank Test. (S1 Data, Sheets 3A, 3B, 3C-E)

F)       AVE-ablated worms show increased sleep. AVA-ablated worms do not show a

significant sleep phenotype. Shown are sleep fractions during lethargus. *p <

0.05, Kolmogorov-Smirnov-Test. (S1 Data, Sheet 3F)

G)      RIS does not reach the same activation levels in *aptf-1(gk794)* and *flp-11(tm2705)*

mutants compared to wild-type worms. *aptf-1(gk794)* and *flp-11(tm2705)* mutants

do neither immobilize nor sleep during RIS activation. ***p < 0.001, Welch Test.

(S1 Data, Sheet 3G-I)

H)      *flp-11(tm2705)* mutants have significantly fewer wide RIS peaks. *aptf-1(gk794)*

mutants display the same amount of wide RIS peaks as wild-type worms. **p <

0.01, Kolmogorov-Smirnov-Test. (S1 Data, Sheet 3G-I)

I)       *flp-11(tm2705)* and *aptf-1(gk794)* mutants do not show sleep during lethargus.

**p < 0.01, Kolmogorov-Smirnov-Test. (S1 Data, Sheet 3G-I)

J)       A circuit model for the positive feedback loop between RIS and PVC. Activating

synaptic input is shown as green arrows, inhibitory synaptic input is shown as red

arrows, and gap junctions are indicated as black connections. During wakefulness,

reverse command interneurons inhibit PVC so that PVC does not activate RIS.

During lethargus, PVC directly activates RIS, which then inhibits reverse

command interneurons through FLP-11. This may speculatively disinhibit PVC,

leading to a positive feedback.

RIS induces sleep through the release of neuropeptides with the major sleep-inducing

neuropeptides encoded by the *flp-11* gene [51]. To test whether FLP-11 neuropeptides are

required for RIS-induced PVC activation, we repeated the optogenetic RIS activation

with simultaneous PVC calcium measurement in an *flp-11* deletion mutant. RIS-induced

PVC activation was almost completely abolished in the *flp-11* deletion (reduction of

transient maximum by 79% during lethargus), indicating that FLP-11 neuropeptides are

required for RIS-induced PVC activation (Fig 3B).

While PVC is presynaptic to RIS, RIS is not presynaptic to PVC [28,32]. The activation

of PVC by RIS could involve diffusional mechanisms or could be indirect through other

neurons, perhaps mediated by the inhibition of a PVC inhibitor such as AVA/AVD/AVE.

RIS has been shown to inhibit AVA/AVE [25] and RIS is presynaptic to AVE [28,32],

suggesting that PVC activation involves inhibition of AVE. We hence repeated RIS

activation and PVC calcium imaging in a strain in which AVE was impaired through

expression of tetanus toxin [52]. The initial PVC activation maximum after AVE

impairment was reduced by 43% during lethargus, but subsequent PVC activity was

increased (Fig 3C-E). AVE is connected to other reverse command interneurons, that

collectively inhibit PVC [28,53]. This circuit design suggests that AVE might play a dual

role in controlling RIS activity. It should have a positive role in mediating activation of

PVC through RIS and thus could promote the feedback loop, but it should also have an

inhibiting role by promoting PVC inhibition. To test for a role of the arousal neurons

AVE and AVA in sleep, we inhibited AVE with tetanus toxin [52] and AVA using HisCl

[54] and quantified sleep amount. Whereas we could not find any effect of AVA

impairment on sleep amount, AVE impairment led to an average increase of sleep by

42% (Fig 3F). Together, these data suggest that PVC and RIS rely on positive feedback

for their activation that involves the release of FLP-11 neuropeptides and inhibition of

PVC by AVE.

If depolarization of RIS activates PVC, what consequences does hyperpolarization of RIS

have on PVC activity? To answer this question, we measured the response of PVC to RIS

inhibition. We hyperpolarized RIS optogenetically for one minute using ArchT and

measured the activity of PVC. Interestingly, PVC showed a small but significant activity

increase during RIS inhibition, an effect which was increased during lethargus (S8B Fig).

The disinhibition of PVC by RIS inactivation is likely not direct and may reflect a

general increase in neuronal and behavioral activity that is caused by RIS inhibition and

that extends to the PVC neurons. Since PVC is a major activator of RIS, its disinhibition

could be part of a homeostatic feedback regulation.

Our results suggest that there is a positive feedback from sleep induction onto RIS

activation and that full RIS activation is only possible when sleep is successfully induced,

explaining the strong correlation of RIS depolarization and sleep bout induction [27].

This model would predict that RIS transients are dampened if RIS is not able to induce

sleep bouts. To test this idea we analyzed RIS calcium transients in *aptf-1(-)* mutant

worms in which RIS still shows depolarization transients during lethargus, but cannot

efficiently induce quiescence [25,51]. In *aptf-1(-)* mutant animals, calcium transient

maxima were reduced by about 35% (Fig 3G-I). A major function of APTF-1 is the

expression of FLP-11 neuropeptides that are required for quiescence induction [51]. To

test whether FLP-11 neuropeptides play an essential role in shaping RIS transients, we

measured RIS calcium transients in mutant worms carrying a deletion of *flp-11*. These

mutant animals showed only a reduced number of long RIS transients that were of

reduced size (Fig 3G-I). *flp-11(-)* showed, however, many short RIS transients (S8C-F

Fig) that were not associated with sleep bouts but may reflect attempts to induce sleep

bouts. These results are consistent with the idea that sleep induction is a self-enforcing

process in which RIS-mediated inhibition of brain activity through FLP-11 neuropeptides

promotes long RIS calcium transients (Fig 3J).

We next tested what feedback interaction exists between RIM and RIS neurons. We

optogenetically depolarized or hyperpolarized RIS and measured RIM activity. RIS

activation did not significantly change RIM activity, but there was a small inhibitory

trend (S8G Fig). RIS inhibition led to an activation of RIM (S8H Fig). These results

show that while RIM can activate as well as inhibit RIS, RIS is an inhibitor of RIM.

**RIM can activate RIS but its activation is not sufficient for sleep induction**

A second important activator of RIS is RIM. We hence asked whether RIM, similar to

PVC, also is active prior to RIS depolarization and sleep bouts. We measured RIM

activity by imaging GCaMP in moving worms. All sleep bouts were extracted and RIM

activity was aligned to sleep bout onset. Averaged RIM activity peaked approximately

30s before the beginning of the sleep bout (Fig 4A). This finding is consistent with a

function for RIM in RIS activation. We then asked whether RIM is required for sleep

induction. We ablated RIM through expression of *egl-1* under the *tdc-1* promoter. We

quantified lethargus sleep in RIM-ablated worms. RIM-ablated larvae showed a normal

fraction of sleep, a slightly increased frequency of sleep bouts and a normal length of

sleep bouts (Fig 4B-D). In analogy to the PVC experiments, we analyzed the effect of

optogenetic RIM depolarization on behavior. We first tested behavior caused by

activation of RIM with ReaChR driven by the strong *tdc-1* promoter on the locomotion of

worms. Consistent with previous findings [35] and our observation that RIS is inhibited

under these conditions, RIM activation during lethargus caused mobilization and larvae

crawled mostly backwards (Fig 4E). We next tested for the effects of weaker RIM

activation using the *gcy-13* promoter. Activation of RIM caused increased mobility when

RIS was inhibited. In trials in which RIM activation led to RIS activation, there was no

significant change of speed of the worms (S1D Fig). We next wanted to test whether

excitability of RIM is altered during the lethargus state. We hence activated RIM strongly

using the *tdc-1* promoter and measured RIM activity. Outside of lethargus, RIM was

strongly excited. During lethargus, however, excitability was strongly reduced (Fig 5F

and 5G). In summary, RIM activation is not sufficient to induce sleep. RIM could,

however, contribute to strong RIS activation and sleep induction by acting in concert with

other neurons. Reduced excitability of RIM during lethargus could favor the activating

effect of RIM on RIS, while dampening the inhibiting effects of RIM on RIS.

**Fig 4**

**RIM activity peaks prior to sleep bouts, but RIM activation is not sufficient for**

**sleep induction**

A)  RIM activates prior to sleep bouts. *p < 0.05, Wilcoxon Signed Rank Test. (S1

Data, Sheet 4A)

B-D)  RIM-ablated worms have an increased sleep bout frequency while the sleep

fraction and bout duration are not significantly changed during L1 lethargus. RIM

was genetically ablated by expressing *egl-1* under the *tdc-1* promoter. *p < 0.05,

Kolmogorov-Smirnov-Test. (S1 Data, Sheet 4B-D)

E)  RIM depolarization leads to increased mobility and reverse motion. *p < 0.05,

***p < 0.001, Wilcoxon Signed Rank Test for Speed. Fisher's Exact Test for

fraction of direction. (S1 Data, Sheet 4E)

F-G)  During lethargus, RIM becomes resistant to activation. RIM was optogenetically

activated using ReaChR expressed under the *tdc-1* promoter. Outside of lethargus

its activation is stronger (F). Activity levels during the stimulation period were

quantified by subtracting baseline activity levels from levels during the

stimulation period (G). *p < 0.05, **p < 0.01, Wilcoxon Signed Rank Test for

GCaMP and Kolmogorov-Smirnov-Test for quantification of stimulation levels.

(S1 Data, Sheet 4F-G)

**Interneurons regulating locomotion act in concert to activate RIS**

Separate activation of PVC or RIM neurons caused moderate RIS activation but not the

strong activation of RIS that is typically associated with sleep bouts. Thus,

hypothetically, multiple neurons act in concert to cause strong RIS activation. Our

presynaptic neuron analysis above suggests that this hypothetical set of neurons should

include PVC and RIM interneurons, but could also include additional neurons. Our

analysis of RIM and PVC points to neurons of the command interneuron circuit for RIS

activation and hence we tested the effects of ablation of a large fraction of the

interneurons controlling locomotion. The *nmr-1* promotor expresses in AVA, AVE, AVD

and PVC command interneurons as well as in second layer RIM neurons [55]. We used a strain that ablates these locomotion-controlling interneurons by expressing the pro-apoptosis regulator ICE from the *nmr-1* promotor [55] and measured sleep and RIS activation. Command interneuron ablation reduced sleep bouts during lethargus by about 76% (Fig 5A) and RIS activation was reduced by 63% (S9A Fig). The movement of command interneuron-ablated worms also was slower (S9B Fig). Quiescence bouts did not occur at the beginning of the lethargus phase as defined by cessation of feeding, and were only observed around the middle of the lethargus phase (S9C Fig). An independently generated strain that ablates command interneurons using *egl-1* expression, also by using the *nmr-1* promoter, caused a reduction of sleep by 81% (Fig 5A).

**Fig 5**

**The locomotion interneuron circuit controls RIS activation and sleep**

A)   Command interneurons are responsible for the majority of sleep. Command interneurons were genetically ablated by expressing ICE or *egl-1* under the *nmr-1* promoter. Command interneurons-ablated worms display a massive loss-of-sleep phenotype. \*\*\*p < 0.001, Welch Test. (S1 Data, Sheet 5A)

B)   Hyperpolarization of command interneurons causes RIS inhibition and suppresses sleep. During lethargus, the hyperpolarization is followed by a strong post stimulation activation of RIS. \*\*p < 0.01, \*\*\*p < 0.001, Wilcoxon Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S1 Data, Sheet 5B)

Next, we wanted to test conditional loss of function of the command interneuron circuit on RIS activity. We expressed ArchT broadly in locomotion-controlling interneurons by using the *nmr-1* promoter. We then inhibited all command interneurons using green light and simultaneously imaged the activity of RIS. Inhibition of *nmr-1*-expressing neurons strongly inhibited RIS inhibition both outside and during lethargus. Interestingly, there was a strong post-stimulus activation of RIS, which was strongly increased only during lethargus. This activation peaked at approximately 170% of the RIS baseline. Sleep was inhibited by command interneuron inhibition and worms reached mobility speeds similar

to those outside of lethargus (Fig 5B). Mosaic analysis of an extra chromosomal array

carrying the *nmr-1::ArchT* transgene revealed that RIS was partially inhibited when

ArchT was expressed in head neurons but not in PVC, and that the effect of inhibition

was substantially stronger when ArchT was not only expressed in head neurons but also

expressed in PVC (S9D-E Fig). This experiment showed that multiple interneurons act in

concert to activate RIS and induce sleep. Among the *nmr-1*-expressing interneurons only

RIM and PVC are presynaptic to RIS [28,32]. However, additional reverse command

interneurons could also contribute to RIS regulation through indirect mechanisms.

Because the command interneuron circuit is controlled by glutamatergic signaling [55,56]

and because RIM activation of RIS requires glutamate (S3 Fig), we also analyzed the

sleep behavior of *eat-4(ky5)* mutant larvae that have impaired glutamatergic

neurotransmission. In *eat-4(ky5)* mutant larvae, sleep bout duration was significantly

reduced, whereas sleep bouts occurred with normal frequency. This indicates that

glutamate signaling might play a role in the maintenance but not in the initiation of sleep

bouts (S10A-D Fig). Consistent with these findings, glutamate signaling also plays a role

in the maintenance of NREM sleep in mice [13]. *nmr-1(ak4)* glutamate receptor mutant

larvae only displayed slightly reduced RIS activation transients, which indicates that

additional glutamate receptors are required for sleep induction (S10E-I Fig). Together,

these mutant phenotypes support the view that excitatory neurotransmitter systems that

are associated with locomotion are important for RIS activation.

**RIS inhibition causes homeostatic rebound activation**

The design of the sleep circuit suggests an intimate mutual control mechanism of RIS and

command interneurons that could allow homeostatic control of sleep. Arousing

stimulation is known to inhibit sleep-active neurons and to increase subsequent sleep

[22,23,25,27]. Consistent with these published data, we observed that the maximum RIS

GCaMP intensity increased logistically with the length of the preceding motion bout

during lethargus (S11A Fig). We thus hypothesized that stimulation inhibits RIS and

leads to its subsequent depolarization, forming a homeostat that allows maintaining or

reinstating sleep bouts. We tested this hypothesis by arousing the worms with a blue light stimulus (Fig 6A-B). During the stimulus worms mobilized and sleep was inhibited. In some of the trials, worms went back to sleep promptly after the stimulation and decreased their motion speed again within three minutes. Because worms did not remain mobile after the stimulation, we classified these trials as "non-mobilizing". In these non-mobilizing trials, RIS showed a post stimulus activation, which was 34% stronger than the baseline activity. RIS activation correlated with a significantly increased fraction of sleep. In other trials during lethargus, the worms stayed mobile for at least three minutes after stimulation and did not go back to sleep. Because worms remained mobile after the stimulation, we classified these trials as "mobilizing". In these mobilizing trials, RIS stayed inhibited and was 16% less active than the baseline before stimulation (Fig 6A). To measure global neuronal activity during the blue-light stimulation experiment, we imaged worms that expressed pan-neuronal GCaMP [57]. Trials were again divided into mobilizing and non-mobilizing trials during lethargus depending on the mobilization status after the stimulus. Non-mobilizing trials showed a global neuronal inhibition that was 93% of the baseline activity (Fig 6B). These experiments show that noxious blue-light stimulation inhibits sleep and RIS and causes a reactivation of RIS when the system returns to sleep.

**Fig 6**

**RIS inhibition causes homeostatic rebound activation**

A-B)   A blue light stimulus leads to awakening and mobilization of *C. elegans.* Worms that go back to sleep after the stimulus show an activation rebound: pan-neuronal inhibition below baseline levels and RIS activation above baseline levels; "lethargus mobilizing" refers to animals that stayed awake and active during the post-stimulus time. "lethargus non-mobilizing" refers to animals that went back to sleep after the stimulation. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, Wilcoxon Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S1 Data, Sheet 6A and 6B)

C)   RIS shows rebound activation following hyperpolarization. Behavioral and brain activity measurements correlate throughout the whole experiment. $*p < 0.05$, $**p$

$< 0.01$, ***$p < 0.001$, Wilcoxon Signed Rank Test for GCaMP and speed,

Fisher's Exact Test for sleep fraction. (S1 Data, Sheet 6C)

D-E)   Dose-response curve of optogenetic RIS hyperpolarization with different stimulus

lengths. RIS activation rebound transients saturate with increasing length of

inhibition. Worms not showing a rebound activation transient after RIS

optogenetic hyperpolarization were excluded from the analysis. Numbers of

worms not responding were as follows: 1) In experiments, in which RIS was

optogenetically inhibited for 48 seconds, all worms showed a RIS rebound

activation transient. 2) In experiments, in which RIS was optogenetically inhibited

for 5 minutes, 1 out of 7 worms did not show a RIS rebound activation transient.

3) In experiments, in which RIS was optogenetically inhibited for 10 minutes, 1

out of 13 worms did not show a RIS rebound activation transient. Curve in D was

fitted as an asymptotic function and curve in E was fitted as a BoxLucas1

function. (S1 Data, Sheet 6D, E)

In normal sleep and in the sensory stimulation experiment, periods of inactivity of RIS

were always followed by periods of RIS activation. This suggested that inhibition of RIS

causes its subsequent reactivation. We tested this hypothesis by optogenetically

hyperpolarizing RIS and following its activity using calcium imaging. We inhibited RIS

directly for 60 seconds by expressing the light-driven proton pump ArchT specifically in

this neuron and used green light illumination to activate ArchT. We followed RIS

calcium activity using GCaMP during the experiment and quantified behavior.

Optogenetic hyperpolarization of RIS led to a decrease in intracellular calcium and

increased behavioral activity. Approximately one minute after the end of the inhibition,

RIS showed a rebound activation transient during which calcium activity levels increased

strongly and rose well above baseline levels, concomitant with a decrease in behavioral

activity. Overall brain activity measurements showed that behavioral activity and brain

activity correlated throughout the experiment (Fig 6C). Rebound activation was neither

observed following PVC nor RIM inhibition (Fig 2C and S11B Fig), suggesting that

rebound activation is specific to RIS and is not a general property of all neurons [58].

Strikingly, while the rebound transient was also measurable outside of lethargus, the

strength of the RIS rebound depolarization was three-fold stronger during lethargus than
before lethargus, indicating that the propensity for RIS rebound activation is strongly
increased during lethargus.

To test whether rebound activation of RIS mediates acute or chronic homeostasis, we
tested whether the strength of the rebound activation is a function of length of prior
inhibition. For this experiment we increased the length of the RIS inhibition and
quantified the time it took after the end of the stimulation until the rebound transient
started as well as the peak maximum of the rebound. After inhibiting RIS for five
minutes, the rebound initiated immediately after the end of the stimulation and the
maximum that was reached exceeded that observed after about one minute of RIS
stimulation. Inhibiting RIS for ten minutes did not further increase the occurrence or
strength of the rebound transient. These results show that RIS activation rebound
transients rapidly saturate with increasing length of inhibition (Fig 6D-E and S11C-E
Fig). Thus, RIS shows a rebound activation following inhibition. The rebound activation
presents the translation of RIS inhibition into subsequently increased RIS activity and
thus sleep induction. Rebound activation of RIS does not seem to constitute a chronic
integrator of wake time but presents an acute homeostatic regulatory phenomenon to
induce or reinstate sleep bouts.

Rebound activation of RIS could present a cell-intrinsic property or could be generated
by a neural circuit. To discriminate between these hypotheses, we measured rebound
activation in *unc-13(s69)* mutant animals in which synaptic signaling is globally impaired
[59], or in worms that express tetanus toxin [60] specifically in RIS to abrogate synaptic
transmission specifically in this neuron. Rebound activation of RIS was abolished in
RIS::tetanus toxin (S11F-G Fig) as well as *unc-13(s69)* worms (S11F and S11H Fig).
These results indicate that rebound activation of RIS is a property of the neuronal
network.

In analogy to the activation rebound seen after optogenetic RIS inhibition, optogenetic
RIS activation might cause a negative rebound, i.e. an inhibition of RIS inhibition below

baseline levels following its optogenetic activation. Indeed, we observed such an effect.

Interestingly, the negative rebound was 3-fold stronger during lethargus compared to

outside of lethargus (S11I Fig). However, such a negative rebound was also present in

other neurons such as PVC (S7A Fig), making it difficult to judge whether this effect is

part of a specific sleep homeostatic system or rather a general response of neurons to

strong depolarization [58]. In Summary, RIS activity is homeostatically regulated with its

inhibition causing its reactivation. This rebound activation is strongly increased during

lethargus and likely is required for inducing or reinstating sleep.

**Modest dampening of brain arousal occurs upstream of RIS**

Our results demonstrate that the command interneuron circuit, including PVC, plays a

major role in activating RIS involving self-enforcing positive feedback, resulting in

strong RIS activation and thus sleep induction. RIS calcium transients during

development outside of lethargus are small, whereas transients are high during lethargus.

What determines that RIS calcium transients are limited outside of lethargus but

promoted during lethargus? As an important principle of command interneuron control,

forward and reverse command interneurons inhibit each other to allow discrete forward

and reverse locomotion states. The AVA/AVD/AVE/RIM interneurons initiate reverse

locomotion by activating pre-motor interneurons while inhibiting the forward command

circuit including AVB/PVC. By contrast, during forward movement reverse command

interneurons are inhibited [49,56].

Small changes in arousal and activity of the command interneurons can change the

equilibrium of forward and reverse command interneurons [55]. Hyperactive mutants

suppress sleep across species including *C. elegans* [61-68]. Many arousal cues trigger

backwards escape movements and inhibit RIS [25,27,69]. Thus, previous studies on the

command interneuron circuit together with our results suggest that arousal inhibits RIS

through inhibiting PVC. This model of RIS activation would predict that there are

changes during lethargus that are upstream of RIS activity that change the properties of

the command circuit leading to increased PVC and thus RIS activation.

We reasoned that it should be possible to measure these changes that occur in command

interneuron activity upstream of RIS by characterizing neural activity and behavior in

*aptf-1(-)* mutant worms. We quantified behavior and command interneuron calcium

levels across lethargus in *aptf-1(-)* mutant worms. Wild-type animals showed successive

sleep bouts and a 72% reduction in locomotion speed during lethargus. By contrast, *aptf-*

*1(-)* mutant animals almost never showed quiescence bouts (Fig 3I), but nevertheless

locomotion speed was decreased by 20% during the lethargus phase (Fig 7). Consistent

with the behavioral activity reduction, there was a significant reduction of command

interneuron activity during lethargus also in *aptf-1(-)* mutant animals (Fig 7 and S12 Fig).

To further characterize the neuronal changes upstream of RIS-mediated sleep induction,

we imaged the activity of RIM during lethargus in *aptf-1(-)* mutants. In wild-type

animals, RIM regularly showed activation transients before lethargus but did not show

many transients during lethargus. RIM showed not only a change in transient frequency

across the lethargus cycle but also a reduction in baseline calcium activity. In *aptf-1(-)*

mutant worms, RIM continued showing calcium transients during lethargus, indicating

that RIS inhibits calcium transients in RIM during sleep bouts. However, reduction of

baseline calcium activity was preserved in *aptf-1(-)*, indicating that RIM activity is

dampened during lethargus independently of RIS at the level of baseline calcium activity.

Together these experiments indicate that a dampening of behavioral and neural baseline

activity occurs during lethargus that is independent of RIS. This neuronal baseline and

behavioral dampening itself appears not to be sufficient to constitute normal sleep bouts,

but could hypothetically lead to an activity change and decreased mutual inhibition in

command interneurons thus promoting sleep induction [55,70].

**Fig 7**

**The dampening of neural and behavioral baseline activity levels during lethargus is**

**independent of RIS function**

Reduction of command interneuron activity levels during lethargus occurs in wild-type

worms and *aptf-1(gk794)* mutants. In the wild-type condition activity levels are reduced

to -0.16 ± 0.02. In the mutant condition activity levels are reduced -0.08 ± 0.02. $**p <$

0.01, Wilcoxon Signed Rank Test. (S1 Data, Sheet 7A and 7B)

**An arousing stimulus inhibits RIS through RIM**

Arousal plays a major role in inhibiting sleep but the circuits that mediate the effect of

arousing stimuli on RIS inhibition are not well understood. We hence studied the circuit

by which stimulation of a nociceptor, the ASH neurons, leads to a reverse escape

response and inhibition of RIS [71]. We optogenetically stimulated ASH using ReaChR

and green light and followed RIS and RIM activities. ASH activation led to a strong

activation of the RIM neuron and triggered a backwards response as previously described

[35,71]. Simultaneously, RIS was inhibited (Fig 8A). RIM can inhibit PVC through

reverse interneurons which it synchronizes [49,72]. Furthermore, strong RIM activation

can inhibit RIS directly. To test whether ASH indirectly inhibits RIS through RIM, we

ablated RIM genetically by expression of *egl-1* from the *tdc-1* promoter [35,38] and

repeated the optogenetic stimulation of ASH. In RIM-ablated L4 animals, activation of

ASH caused the opposite effect on RIS activity. Instead of inhibiting RIS, ASH activated

RIS, while it still increased behavioral activity (Fig 8B). Consistent with our calcium

imaging data, ASH stimulation after RIM ablation predominantly caused a forward

locomotion response (Fig 8C). There are two ways ASH might inhibit RIS through RIM.

One possibility is that arousal strongly activates reverse interneurons, thus inhibiting

forward PVC neurons and RIS during stimulation. Consistent with this idea, gentle tail

touch increased RIS activity more strongly when RIM was ablated (S13 Fig). Another

option is that RIM inhibits RIS directly through tyramine and FLP-18. Both circuits

might play together (Fig 8D). These results delineate a circuit model for how sensory

stimulation can control RIS activation.

**Fig 8**

**Arousing stimulation inhibits RIS and sleep through RIM**

A) ASH depolarization in wild-type worms leads to RIS inhibition and RIM

761            activation, sleep suppression and mobilization. $*p < 0.05$, $**p < 0.01$, $***p <$

0.001, Wilcoxon Signed Rank Test for GCaMP and speed, Fisher's Exact Test for

sleep fraction. (S1 Data, Sheet 8A, B)

B) ASH depolarization in RIM-ablated worms leads to weaker sleep suppression,

mobilization, and RIS activation. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, Wilcoxon

Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction.

(S1 Data, Sheet 8A, B)

C) The response direction following ASH activation in wild-type worms is

predominantly reverse, while in RIM-ablated worms it is predominantly forward.

This suggests that the forward direction is connected to RIS activation. $***p <$

$0.001$, Fisher's Exact Test. (S1 Data, Sheet 8C)

D) A circuit model for RIS regulation through arousal by ASH. Activating synaptic

input is shown as green arrows, inhibitory synaptic input is shown as red arrows, and

gap junctions are indicated as black connections. RIM could serve as a synchronizer

of AVE and AVA to regulate PVC and therefore RIS inhibition. Additionally, RIM

could inhibit RIS directly.

## Discussion

### A wake-active circuit that controls locomotion also promotes sleep

Optogenetic activation and inhibition showed how the activity of presynaptic neurons

affects RIS depolarization during developmental sleep. Several presynaptic neurons can

activate RIS. RIM appears to be a potent direct inhibitor when activated strongly, but can

also act as an activator of RIS. Loss-of-function experiments showed that the command

circuit controls activation of RIS, with PVC presenting a key activator of RIS. PVC has

long been known to mediate the forward escape response by transmitting information

from posterior sensory neurons to activate AVB pre-motor neurons to trigger forward

locomotion [46,49,50]. Consistent with promoting the forward escape response,

optogenetic activation of PVC leads to an increase in forward movement [50,73] (Fig

2D). Reverse movement, in turn, is mediated by AVA, AVE, and AVD command

premotor interneurons, which activate reverse motor neurons. Forward PVC and reverse

AVA/AVE/AVD command interneurons are presynaptic to and mutually inhibit each
other, which ensures discrete forward and reverse locomotion states analogous to a flip-
flop switch [49,55,74].

Our finding that PVC and RIM neurons present key activators of RIS that act in concert
suggests a model for how RIS is controlled and provides a potential mechanism for
linking sleep induction to decreasing arousal and for homeostatically maintaining a series
of sleep bouts. According to this model, during conditions of high arousal, such as during
development outside of lethargus, larvae are constantly awake. The command interneuron
circuit cycles between forward and reverse states, leading to the activation of forward or
reverse motor programs, respectively [49,74,75]. PVC activation has been associated
with the activity of forward states and RIM has mostly been associated with the activity
of reverse states. As neither activation of only the PVC nor the RIM neurons appears to
be sufficient for sleep induction, RIS should not be activated sufficiently to induce sleep
during either forward or reverse states. At the transition between forward and reverse
states, locomotion pauses can occur. It has been shown that in adult worms, RIS shows
activation transients in the nerve ring during locomotion pauses. These calcium transients
appear to be much smaller compared with activation transients during sleep bouts that
extend to the cell soma. Locomotion pausing is reduced after RIS ablation, suggesting
that weak RIS activation promotes pausing [76].

Lethargus induces a modest dampening of neuronal baseline activity that is independent
of RIS and that includes the RIM neurons. The RIM neurons become less excitable,
which should reduce their inhibitory effects on RIS and instead favor their activating
effects. PVC becomes resistant to inhibition and more potent in its capacity to activate
RIS. We hypothesize that these shifts in the properties of the interneurons of the
locomotion circuit favor the activation of the RIS neuron. RIS activation appears to
require concerted activation from PVC and RIM neurons (a process that is perhaps aided
by other locomotion interneurons). Both PVC and RIM appear to depolarize prior to RIS
activation and both types of neurons contribute to RIS depolarization. This suggests that
RIS might be activated when both PVC and RIM exert activating effects. Such an

overlapping activating effect of PVC and RIM on RIS would most likely occur at the
transition from forward to reverse locomotion states, where there could be an overlap of
both forward and reverse neuronal activities. This would suggest that both locomotion
stop and sleep bouts might be induced by locomotion control interneurons at the
transition between forward and reverse locomotion states. The difference between
locomotion stop and a sleep bout would be that in the former RIS would only be
modestly activated, whereas in the latter RIS would be strongly activated (Fig 9).
Consistent with this model, sleep bouts are typically induced at the end of long forward
movements, whereas the exit from the sleep bout is often through a reverse movement
[70,75,77]. Arousal promotes reverse command interneurons activity and strong RIM
activation that can inhibit RIS. Locomotion control and periods of behavioral activity and
rest are already present in animals that do not have a nervous system. It has thus been
hypothesized that sleep and sleep-active neurons evolved from systems controlling
locomotion activity and rest [8]. The finding that a sleep-active neuron can also act as a
locomotion pause neuron [76] and the discovery presented here that the locomotion
circuit controls the depolarization of a sleep-active neuron suggests that sleep might have
evolved from locomotion-controlling neurons and that thus locomotion quiescence and
sleep could be regarded to be homologous behaviors.

**Fig 9**
**A circuit model for RIS activation through locomotion interneurons**
A)    Activating synaptic input is shown as green arrows, inhibitory synaptic input is
shown as red arrows, and gap junctions are indicated as black connections.
Outside of lethargus, the nervous system cycles between forward and reverse
states. RIS is not activated sufficiently to cause a sleep bout, neither during the
forward state during which PVC is active nor during the reversal state during
which RIM is active. The locomotion circuit activates RIS briefly to cause a
locomotion pause at the transition from forward to reverse movement.
Speculatively, the circuit that controls RIS during sleep also controls RIS during
locomotion pauses.

B)       During lethargus motion bouts, the nervous system still cycles between forward

855           and reverse states. Baseline activity and excitability in RIM are reduced, PVC

856           becomes resistant to inhibition and more potent to activate RIS. These changes in

857           locomotor interneurons shift the balance to favor strong RIS activation and

858           induction of a sleep bout, a process which may involve simultaneous activation

859           from multiple neurons including RIM and PVC. Such an overlap activation of RIS

860           by otherwise mutually exclusive neurons could occur at the transition from

861           forward to reverse locomotion states. Perhaps, RIS activation and sleep could

862           occur similarly at the transition from reverse to forward locomotion states.

Our model suggests that the sleep switch is tripartite and includes not only wake-active

wake-promoting neurons and inhibitory sleep-active sleep-promoting neurons, but also

wake-active sleep-promoting neurons as mediators of switch flipping. This sleep switch

acts as an amplifier that can translate a modest reduction of arousal into a massive

shutdown of behavioral activity during sleep. Dampening of neural activity and altered

properties of wake-active sleep-promoting locomotion neurons independently of sleep-

active neurons could be interpreted as a neural equivalent of sleepiness that leads to an

increased propensity to activate sleep-active neurons and to induce sleep bouts.

Mutations that increase arousal and suppress sleep increase the activity of reversal

neurons, whereas conditions that decrease arousal decrease the activity of the reversal

neurons and therefore increase the amount of sleep [67,68,74]. Also, the ablation of

reverse command interneurons such as AVE reduces reversals and leads to ectopic

quiescence, and increases sleep [46,52](and this study). According to our model,

increasing arousal should increase the activity of RIM and other reverse command

interneurons and thus should inhibit RIS. Conversely, reducing arousal could promote

weaker RIM activation and PVC activation that should shift the equilibrium to stronger

RIS activation.

What causes the termination of sleep bouts? The RIS neuron might not be able to sustain

prolonged activity leading to the spontaneous cessation of a sleep bout. The RIS

activation transient and thus sleep bout can be blunted prematurely by a sensory or
optogenetic arousing stimulus [25,27,70,78]. Arousing stimulation, for instance by
activating the nociceptive sensory neurons, triggers a reverse escape response through
backwards command and RIM interneurons [35,72,75,79]. Strong optogenetic RIM
depolarization inhibits RIS and stimulation of the nociceptive ASH neurons causes
inhibition of RIS that depends on RIM, suggesting that a main physiological role of
strong RIM activation is to inhibit sleep upon arousing stimulation, perhaps by
synchronizing the reverse interneurons [72]. RIM activation can inhibit sleep also in
response to acute food deprivation [80,81]. Thus, RIM might present not only an
activator of RIS but also an arousal module that can be activated upon sensing various
external conditions that signal the need to suppress sleep.

RIS inactivation leads to disinhibition of arousal and brain activity starting anew the
cycle of locomotion interneuron activity and locomotion behavior. Depending on the
arousal levels, the locomotion circuit causes RIS reactivation and thus a return to sleep
either immediately or after a delay. The timing at which the rebound activation occurs
can be controlled by the level of arousal, with strong arousal leading to longer wake
periods before the return to sleep, whereas milder stimulations cause the immediate
return to sleep [23]. Consistent with this circuit model of recurrent RIS activation, RIS
activity oscillates, resulting in the typical pattern of sleep bouts that are interrupted by
activity bouts [22]. This circuit design allows homeostatic sleep maintenance of a series
of consecutive sleep bouts with sensory stimulation restarting the cycle of RIS activation
thus prompting an acutely increased RIS activation causing the return to sleep (Fig 6A)
[23,70]. Our model predicts that RIS calcium transient strength is a function of prior
behavioral activity. Consistent with this view, RIS calcium transients are stronger at the
beginning and end of lethargus, when motion bouts are high but less pronounced in the
middle of lethargus when motion bouts are less pronounced (Fig 1A) [22,23]. Thus, the
tripartite flip-flop circuit design allows an adaptation of RIS activity to the strength
required to induce sleep bouts at a given behavioral activity level.

Here we have identified a circuit controlling sleep-active neuron depolarization in *C. elegans*. This work built on the neural connectome and was facilitated by the small size and invariance of the nervous system as well as the transparency of the organism. While the *C. elegans* sleep circuit clearly is built from less cells than the mammalian sleep circuit [8,82,83], there are many conceptual similarities. For instance, in both *C. elegans* and humans, sleep is controlled by inhibitory sleep-active sleep-promoting neurons that depolarize at sleep onset to actively induce sleep by inhibiting wake circuits. A main difference is that humans have many brain centers each consisting of thousands of sleep-active neurons [12]. The single RIS neuron is the major inhibitory sleep-active neuron required for sleep induction in *C. elegans* [25]. Work in mammals revealed the general principles of wake-active wake-promoting neurons and sleep-active sleep-promoting neurons as well as their mutual inhibition. While this information explains the flip-flop nature of sleep and wake states, there is no satisfactory understanding of what flips the sleep switch, i.e. how wakefulness is detected when the system is set to sleep, prompting the activation of inhibitory sleep-active neurons [6]. Our model for the operation of the *C. elegans* sleep circuit indicates that flipping of the sleep switch can be understood if wake-active sleep-promoting neurons are added to the switch model. In this tripartite flip-flop sleep switch model, the sleep-active sleep-promoting center is activated by wake-active neurons. This activation should, however, only occur when the system is set to sleep, a state that could present a neural correlate of sleepiness.

Sleep is reversible by stimulation and hyperarousal is the major cause for insomnia in humans [3,84,85]. Homeostatic sleep maintenance is an essential feature of sleep and is found from worms to humans [19-21,23]. R2 ring neurons in *Drosophila* present an integrator of wake time causing subsequently increased depolarization of ExFl1 sleep-inducing neurons, thus forming a chronic sleep homeostat [86,87]. In vertebrates, serotonergic raphe neurons are active during wakefulness, and can reduce behavioral activity and increase sleep pressure [88]. Our model of a tripartite flip-flop circuit suggests that wake-active sleep-promoting neurons are an essential part of an acute sleep homeostat that translates acute brain activity into increased sleep neuron activity when the system is set to sleep. Wake-active sleep-promoting neurons measure systemic

activity, i.e. they become active together with a global brain activity increase, and can
then activate inhibitory sleep-active neurons. Thus, the interplay of sleep-active sleep-
promoting and wake-active sleep-promoting neurons form an oscillator that periodically
sends out sleep-inducing pulses. Macroscopically, sleep in mammals exists as cortical
oscillations of global down states, known as slow waves [89]. Micro arousals trigger
cortical up states that are followed by cortical down states, known as k-complexes [19-
21]. Both slow wave activity as well as k-complexes could be hypothetically generated
by wake-active sleep-promoting neurons.
954

955 **Materials and Methods**
956

**Worm maintenance and strains**
*C. elegans* worms were grown on Nematode Growth Medium (NGM) plates seeded with
*E. coli* OP50 and were kept at 15-25°C [90]. Crossed strains were genotyped through
Duplex PCR genotyping of single worms [91]. The primer sequences that were used for
Duplex PCR can be found below. To confirm the presence of transgenes after crossings,
fluorescent markers were used. All strains and primers that were used in this study can be
found in the supporting information (S1 Text and S1 Table).
964

965 **Strain generation**
966

DNA constructs were cloned with the three fragments Gateway System (Invitrogen,
Carlsbad, CA) into pCG150 to generate new strains [92]. The *ArchT*, the *ReaChR* and the
*egl-1* genes were expression-optimized for *C. elegans* [93]. The *tdc-1::egl-1* transgene
specifically expresses the apoptosis inducing protein EGL-1 in RIM and RIC. Hence,
RIM and RIC are genetically ablated in worms carrying this transgene. The ablation is
probably incomplete in L1 worms. The *nmr-1::egl-1* transgene leads to the expression of
*egl-1* in all command interneurons causing their genetic ablation. Similar to the *tdc-*
*1::egl-1* transgene, ablation might be incomplete in L1 worms. In both lines *egl-1* was co-
expressed with *mKate2*, which was used to verify the genetic ablations. Transgenic
strains were generated by microparticle bombardment or by microinjection. For

977 microparticle bombardment *unc-119(ed3)* was used. The rescue of the *unc* phenotype was

978 hence used as a selection marker [94,95]. The transgenes were backcrossed twice against

979 N2 wild-type worms to remove the *unc-119(ed3)* background. Extrachromosomal arrays

980 were generated by DNA microinjection. DNA was injected in wild-type, mutant or

981 transgenic worms. For injection, DNA was prepared as follows: construct 30-100 ng/µl,

982 co-injection marker 5-50 ng/µl, pCG150 up to a concentration of 100 ng/µl if required.

983 Positive transformants were selected according to the presence of co-injection markers. A

984 table of all plasmids and a list of all constructs that were generated for this study can be

985 found in the supporting information (S2 Table and S2 Text).

987 **Generation of gene modifications using CRISPR**

989 The following alleles were designed by us in silico and were generated by SunyBiotech.

990 Correctness of the alleles was verified by using Sanger sequencing.

992 **PHX816:** *flp-11(syb816 [SL2::mKate2::linker(GSGSG)::tetanustoxin_LC]) X.*

994 The coding sequences of tetanus toxin light chain and mKate2 were codon-optimized and

995 intronized as described previously and were synthesized [93]. The final sequence can be

996 found in the supporting information (S3 Text).

998 **Imaging**

1000 Cameras and software

1001

1002 All imaging experiments were conducted using either an iXon EMCCD (512x512 pixels),

1003 an iXon Ultra EMCCD (1024x1024 pixels), a Photometrics Prime 95B back-illuminated

1004 sCMOS camera (1200x1200 pixels) or a Nikon DS Qi2 (4908x3264 pixels). For the iXon

1005 cameras the EM Gain was set between 100-200. The exposure times used were between

1006 5-30ms. Andor IQ 2 and 3 and NIS Elements 5 were used for image acquisition.

1007

1008    Illumination systems

1009

A standard 100W halogen lamp together with an infrared filter (Semrock Brightline HC

785/62) were used for Differential Interference Contrast (DIC) microscopy or bright-field

imaging. For calcium imaging and blue light stimulation an LED illumination (CoolLED)

with a 490nm LED and standard GFP filter set (EGFP, Chroma) were used. Optogenetic

stimulations and RFP imaging were performed with an LED illumination (CoolLED)

with a 585nm LED and standard TexasRed filter set (Chroma).

1016

1017    **Agarose Microchamber Imaging**

1018

Long-term imaging experiments were conducted in agarose microchambers as previously

described [96,97]. To summarize, a PDMS mold was used to cast box-shaped

indentations in a hydrogel, which consisted of 3 or 5% agarose dissolved in S-Basal [98].

Two different sizes were used. We imaged L1 larvae in 190 x 190 x 15 μm

microchambers and L4 larvae were imaged in 370 x 370 x 25 μm microchambers.

Depending on the developmental state of the worm that was imaged, either pretzel stage

eggs or L3 larvae were picked into the chambers with OP50 bacteria. Before imaging

worms were kept at either 20°C or 25°C.

1027

For time lapse calcium imaging experiments, L1 worms were filmed every 5s (Figs 1A,

3H-J, 4B-D, 5A, S8C-F and S11A Figs), every 8s (Fig 7, S9A, S9C and S10 Figs) or

every 10s (Figs 3F and 4A) with differential interference contrast (DIC) or bright-field

imaging and widefield fluorescence. The DIC and bright-field light source was left on

continuously, filtered through an infrared filter and was blocked by a shutter during

fluorescence image acquisition. LED illumination was triggered by the EMCCD camera

using the TTL exposure output of the camera. An objective with 20x magnification, an

LED with 480 nm (light intensity was between 0.15-2 mW/mm$^2$), and EM gain of 100-

200 was used. With the 20x objective and a 0.7 lens 4 worms could be imaged

simultaneously in one field. 1-4 fields could be filmed in parallel in one experiment.

These image sequences gave measurable neuronal calcium transients and clear DIC or

bright-field images to identify pumping or non-pumping phases.

1040

1041    During the continuous experiments in Figs 3F, 4B-D and 5A only DIC or bright-field

1042    images were taken.

1043

1044    **AVA inhibition experiment**

1045

1046    NGM plates were prepared with histamine (HA, Sigma, 10mM) as previously described

1047    [54]. Young adult worms expressing a histamine chloride channel in AVA and control

1048    worms were picked onto NGM HA plates the night before the experiments. The next

1049    morning eggs together with *E. coli* bacteria from the NGM HA plates were picked into

1050    microfluidic chambers and DIC imaged as previously described [96,97].

1051

1052    **Optogenetic experiments**

1053

Optogenetic experiments were either conducted in agarose-microchambers as described

above or the worms were immobilized. For immobilization experiments the agarose was

solved in S-Basal. We used three methods of immobilization for optogenetic

experiments.

1) Immobilization on a 3% agarose pad with 25mM Levamisole (Sigma Aldrich) (S6

Fig).

2) Immobilization on a 10% agarose pad with 0.1µm Polybead microspheres

(Polyscience) [99] (Figs 1B/ SDQL, 1C/ SDQL, 3B-D, S1A/SDQL, S5A/ SDQL,

S4 and S8A Figs).

3) Immobilization on a 10% agarose pad with 0.1µm Polybead microspheres

(Polyscience) [99] and 25mM Levamisole (Figs 1B/PVC, 1C/PVC, 2A, 2C, 3B-

D, 4F-G, S1A/PVC, S5A/PVC, S7A, S7C-D, S8B and S8H Figs).

1066

1067 Worms were imaged within 30min of immobilization. A 100x oil objective was used for

1068 illumination and imaging in most experiments. For images in S6A-B Fig a 1.5 lens was

1069 added (S6A and S6B Figs). The imaging in Fig 2A was done using a 40x objective.

1070

1071 ReaChR for neuronal depolarization or ArchT for hyperpolarization was utilized. For

1072 optogenetic stimulation, a 585nm LED and a standard TexasRed (Chroma) filter set were

1073 used.

1074

For optogenetic experiments with L1 larvae, either L4 stage worms or young adult worms

were pre-picked onto NGM plates with all-trans-retinal (ATR, Sigma Aldrich) and grown

at 20 or 25°C. During the two days after exposure to ATR, pretzel stage eggs or L1

worms were taken from this plate for optogenetic experiments. For optogenetic

experiments with L4 larvae an agar chunk containing a mixed population of growing

worms was added to NGM plates containing ATR. Worms for optogenetic experiments

were taken from this plate within the next two days.

1082

Calcium imaging was conducted with an interval of 3s and with an exposure time of 5-

200 ms. A standard optogenetic protocol included calcium imaging during a baseline.

This was followed by a stimulation time, in which the worms were optogenetically

stimulated. The 585 nm light exposure was continuous except for brief interruptions

during the time calcium imaging was conducted. After the optogenetic stimulation,

calcium images were acquired during a recovery period.

1089

In mobile worms this standard protocol was preceded by 20 DIC frames that were taken

every 500ms to determine if the worm was pumping. The overall protocol was repeated

every 15-30min. L1 mobile worms were imaged with a 20x objective and a 0.7 lens.

Mobile L4 worms were imaged with either a 10x objective (Figs 8A-C) or a 20x

objective (Fig 1B/CEP, 1C/URY, S1A/CEP, S5A/URY and S3 Figs). Fixed worms were

usually imaged between 1-4 trials. A delay preceded the standard protocol to allow the

worm to recover from immobilization and between trials. To specifically manipulate

PVC and SDQL in Figs 1B/PVC, 1B/SDQL, 1C/PVC, 1C/SDQL, 2D, S1A/PVC,

S1A/SDQL, S4, S5A/PVC, S5A/SDQL and S7C-D Figs the stimulating illumination was

restricted to the neuronal areas. This was achieved by reducing the size of the field

aperture of the fluorescence illumination. To activate a specific neuron, it was moved into

the illuminated area by using an automated stage. To image RIS, this neuron was moved

into the illuminated area by the automated stage, while the optogenetic light stimulus was

switched off and imaging light was switched on. The details for optogenetic experiments

can be found in in the supporting information (S3 Table).

1105

1106     **Behavioral imaging while PVC activation**

1107

Worms were prepared on retinal plates and picked into microchambers as described

before. A 20x objective was utilized for imaging. The entire chamber was imaged

through bright-field imaging. For tail specific illumination, the LED blend was adjusted

to illuminate a circular area with a radius of 58μm. The 580nm LED was manually turned

on after 1min of imaging and off after 4min of imaging. A Prior XY stage was manually

operated to keep specifically the tail of the worm in the by the LED illuminated area

during stimulation. Worms were imaged with a frame rate of 8 Hz. Only every eighth

image was used for analysis.

1116

1117     **Activity measurements of command interneurons**

1118

GCaMP3.3 was expressed in command interneurons using the *glr-1* promoter [100]. L1

larvae were placed in microfluidic chambers and were imaged using a time-lapse

protocol. 1 DIC and 1 GFP image was taken every 8 seconds using a 20x objective and a

0.7 lens. The 490nm intensity for GFP imaging was set to 0.15 mW/mm$^2$. Intensity

values of all command interneurons located in the head of worms were extracted

manually and analyzed as one entity.

1125

1126     **Pan-neuronal activity measurements**

1127

GCaMP6s and RFP were pan-neuronally expressed under the *rab-3* promoter [101]. As

in the activity measurements of command interneurons, L1 lethargus was imaged in

microfluidic devices. For the optogenetic experiment (Fig 6C), each 30min, first 20 DIC

were taken to determine lethargus. This was followed by GFP images that were taken all

5.8s for 9min. The 490nm intensity was set to 0.07 mW/mm$^2$. In the blue light

stimulation experiment, additional RFP images were taken. A costume written MATLAB

code detected the mean intensity of all head neurons in each GFP and RFP frame. The

head neurons were hence analyzed as one entity.

1136

1137    **Blue-light stimulation experiments**

1138

L1 worms were placed in microfluidic-chambers for blue light stimulation experiments.

The protocol was repeated every 15min. First, 20 DIC pictures were taken every 500ms

to determine whether the worm was pumping or not. Next, baseline GCaMP was imaged

for 3min, the stimulation phase then lasted 18s and a recovery phase was imaged for

3min. The 490nm intensity for calcium imaging was 0.07mW/mm$^2$. The 490nm intensity

for stimulation was set to 1.01 mW/mm$^2$ with a 20x objective. The same LED was used

for calcium imaging and stimulation. The intensity levels were controlled with Andor

IQ2 software.

1147

1148    The RFP signal of the pan-neuronal strain was imaged in addition to the GCaMP signal

1149    during the protocol every 3s with 585nm LED illumination, which was set to

1150    0.17mW/mm$^2$.

1151

1152    **Mechanical stimulation using dish tapping**

1153

The mechanical tapping set up was described previously[67,102]. L1 larvae were imaged

in microfluidic-chambers using a 20x objective and a 0.7 lens. Microfluidic-chambers

were put in a specialized dish. The dish was tapped by a piston driven by an

electromagnet. The piston and the electromagnet were held in a homemade aluminum

frame as described previously [102](model used was Kuhnke, product number H2246).

The voltage used for stimulus application was 5V, the tapping stimulus was applied
between image acquisition using TTL triggering to avoid blurring. Imaging was
controlled with Andor IQ2 software. The imaging protocol was repeated every 15min.
First, 20 DIC pictures were taken with a frequency of 2 pictures per second to determine
the status of worms. Throughout all following steps GCaMP measurements were taken
every 3 seconds. The 490nm intensity for calcium imaging was $0.15mW/mm^2$. Baseline
GCaMP was measured over 3min. Following the tap GCaMP was measured for 3min.
This experiment was initially planned to be combined with optogenetic stimulation,
hence a weak green light stimulus was applied, starting 15 seconds before and ending 45
seconds after the tapping stimulation. The part of the experiment during which a green
light stimulus was applied was selected for presentation in this study. Green light
(585nm) for stimulation was set to $0.17mW/mm^2$. As we did not see any noticeable
changes upon applying green light, we presume that it does not strongly affect the
experiment.

1173

1174    **Simultaneous Calcium Imaging of RIS and PVC**

1175

In order to simultaneously image RIS and PVC, L1 lethargus worms were transferred
from a growing plate using a platinum wire worm pick and were fixed on 10% agarose
pads with 0.1µm Polybead microspheres (Polyscience) [99] and 25mM Levamisole. The
worms were then imaged through a 40x oil objective with an image taken every 3s for
30min with 490nm light of 1.35mW/mm to image GCaMP. Fluorescence intensities for
PVC and RIS were cropped by using a region of interest. A custom written MATLAB
script then detected all RIS peaks. For this, the GCaMP data was first smoothed over 30
values through the inbuilt function *smooth,* which is a $1^{st}$ degree polynomial local
regression model. Through the in-built MATLAB function *islocalmax*, and a minimum
prominence value of 0.2, the locations of RIS peaks were detected. The data of RIS as
well as PVC GCaMP intensity was aligned to the detected RIS peak location.

1187

1188    **Spinning Disc Confocal Microscopy**

1189

L4 worms were fixed with Levamisole. Spinning disc imaging was done with an Andor

Revolution disc system using a 488nm (0.34 mW/mm$^2$) and a 565nm (0.34 mW/mm$^2$)

laser and a Yokogawa (Japan) CSU-X1 spinning disc head. Worms were imaged through

a 100x oil objective. In S6A-B Fig an additional 1.5 lens was used. Z-stacks with z-

planes 0.5µm apart spanning a total distance of 10µm were taken and a maximum

intensity projection calculated in ImageJ.

1196

1197    **Tail touch experiment**

1198

L4 worms were grown and filmed on NGM plates with OP50 bacteria at 20°C. An

eyelash was used to gently touch the tail of the worms during L4 lethargus. The time

from tail touch until the worms were immobile again was measured with a timer. If

worms did not mobilize upon tail touching the time was counted as zero. For GCaMP

intensities worms were imaged before and after tail touch each second for a total of 30s.

They were illuminated with a Leica EL6000 LED.

1205

1206    **Image Analysis**

1207

Image sequences for analysis were selected either based on lethargus or molting

timepoints. Lethargus was determined through DIC or bright-field images as the non-

pumping phase before molting. Time points were classified to be in or outside of

lethargus. Typically, the entire lethargus time and two hours before lethargus were

analyzed. Worms, which were immobilized during the measurements were classified

according to their pumping behavior on NGM plates directly before imaging. Two

parameters were extracted from the image sequences.

1215

1) Calcium signals were extracted automatically or manually with custom-written

MATLAB codes. These codes extracted defined regions of each image and detected

intensity and position data. Extracted regions were chosen slightly bigger than the sizes

of measured neurons. From these extracted regions a certain percentage of highest

intensity pixel was taken as signal. The remaining pixels were taken as background. From

the signal the background was subtracted. For the pan-neuronal and interneuron activity

measurements, the signal in the head was treated as one large neuron and analyzed in the

same way as single neurons. All head neurons expressed under the *rab-3* promoter were

included in the pan-neuronal GCaMP measurements.

1225

For all stimulation experiments, optogenetic and blue light stimulation experiments, the

baseline measurement of each time point was utilized for signal normalization and ΔF/F

generation, except for Fig 6C. In Fig 6C, a mean of all baseline intensities for all wake

time points for each worm was calculated. The mean was then utilized for normalization

for all time points for each worm to better show the different RIS activities during wake

and sleep. The pan-neuronal signal in Fig 6B was normalized over the measured RFP

signal to retrieve ΔR/R. For the transient alignments in Fig 3G, peaks and corresponding

speeds were extracted through a custom-written MATLAB script and aligned as time

point zero.

1235

1236 2) The speeds of the worms were calculated from the positions of the tracked neuron,

1237 except for experiments, in which no GCaMP intensity was measured. To analyze these

1238 experiments frame subtraction of DIC or bright-field images was done with a custom-

1239 written MATLAB routine instead.

1240

1241 **Baseline extraction**

1242

In S12A and C Fig baseline of RIM GCaMP data was extracted by excluding the 95-100

percentile range for wild type and by excluding the 75-100 percentile range for *aptf-*

*1(gk794)* mutants. The baseline was smoothed through a second-degree polynomial local

regression model and with weighted linear least squares. Zero weight was assigned to

data points six means outside the absolute deviation. The amount of data points used for

smoothing was three percent.

1249

1250 **Sleep bout analysis**

1251

Sleep bouts were extracted from selected parts of the time-lapse movies. Dependent on
the experiment, a specific period of the movie sequence was selected and processed:

1) the lethargus period (Figs 3F-J, 4A, 5A, 7, S9A, S9C, S10E-I and S12 Figs)

2) the period from two hours before lethargus up to the end of lethargus (Fig 1A)

3) either three (Fig 4B-D and S8C-F Fig) or four hours (S10A-D Fig) before
shedding of the cuticle

To extract sleep bouts, speeds and subtraction values were first smoothed. In Figs 1A,
3G-J, 4A-D, 5A, S8C-F, S10 and S12 Figs speeds were smoothed through a 1$^{st}$ degree
polynomial local regression model over 20 time points. Other experiments were
smoothed through a second-degree polynomial local regression model and with weighted
linear least squares. Zero weight was assigned to data points six means outside the
absolute deviation. Data was smoothed either over three percent of all data (Figs 3H-J, 7
and S9C-F Fig) or over 40 data points (S9A, S9C, S12A and S12C Figs). This was
achieved with the *smooth* function in MATLAB. Smoothed speeds were normalized
between 0 and 1, with 0 representing the lowest and 1 the highest smoothed speed value
of each worm. In order to be scored as a sleep bout, the normalized speed had to be under
a defined percentage threshold of the normalized speed for a minimum time. The exact
speed and time thresholds were adjusted empirically to represent the worms' behavior
[103]. In Fig 3G-I and S8C-F Fig worms had to have a speed below 5% of their
maximum smoothed speeds for at least 2min in order to be counted as sleeping. For all
other experiments the speed threshold was 10% and the time threshold was 2min. The
2min time threshold was implemented to exclude short pauses of the worm that may not
represent sleep bouts. It was determined empirically. The sleep bout analysis was carried
out with a custom-written MATLAB script.

1276

For stimulation experiments, the baseline and recovery time measurements were too short
to include a minimum time threshold in the sleep bout analysis. Hence, immobility was
used as a proxy for sleep. A mean of the wake speeds was calculated for each worm.
Depending on the strain used, the worms were counted as sleeping when they were below
a threshold of 5 to 30% of its mean wake speed. In most experiments a worm was
counted as sleeping when its speed was below 10% of the calculated mean of the wake

speeds. In S11B Fig the threshold was adjusted to 5%, in S8G Fig to 20%, in S7B Fig to 25%, in Fig 6C, S1C, S2A-C, S3, S5B, S9D-E and S11I Figs to 30% and in S11C-E Fig to 50% to account for a different locomotor behavior of the worms. RIS signals and speeds of wild type and mutants were aligned to sleep bout onset for comparison in Figs 3H, 4A, S8F, S9A, S10D and S10H Figs. RIM signals and speeds were aligned to sleep bout onset in Fig 4A. For GCaMP normalization ten data points before sleep bout onset were taken as baseline in order to calculate $\Delta F/F$. In S11A Fig motion bouts were assigned whenever there was no detected sleep bout.

**RIS peak alignment**

For RIS wide peak detection (Fig 3G-H), first, the normalized GCaMP data was smoothed over 60 time points with the in-built MATLAB function *smooth*. Wide peaks were then detected with the in-built MATLAB function *findpeaks* and a minimum peak prominence threshold of 0.15. GCaMP intensities, speeds and sleep fractions were then aligned to the detected peak maxima. Analysis for narrow peaks was conducted similarly, only two aspects were changed (S8C-F Fig). To find narrow peaks, smoothing was limited to only 5 time points and a minimum peak prominence threshold was set to 0.2.

**Detection of direction of movement**

The direction of movement was analyzed with a costume written MATLAB script. This MATLAB script took two points, the nose and the pharynx, to calculate the direction. For two consecutive images the distance of the nose in the first image to the pharynx in the first image was compared to the distance of the nose in the second image to the pharynx of the first image. If the distance increased, the worm was counted as moving forward, if it decreased, it counted as moving reverse. If the worm was below a threshold of 2µm/s it was counted as sleeping in experiments Figs 2D and 4E. The position of nose and pharynx were detected manually (Figs 2D and 4E). For correction of the stage movement while manually tracking PVC (Fig 2D), the position of a corner of the stage was used.

1314    **Fitting**

1315

1316    The data in Fig 6D was fitted to an asymptote and the data in Fig 6E was fitted to a

1317    BoxLucasFit1 with Origin software. The data in S11A Fig was fitted to a logistic

1318    regression using Origin software. Exact functions and $R^2$ values can be found in the

1319    respective Figs.

1320

**Statistics**

Sample sizes were determined empirically based on previous studies. If possible,

experiments were carried out with internal controls. If this was not possible, control and

experimental condition were alternated. Researchers were not blinded to the genotype for

data analysis, as data analysis was performed by automated routines. Sample exclusion is

described in the respective methods section. To compare GCaMP intensities and speeds

of one sample group at different time points the Wilcoxon Signed Rank Test was utilized.

The Fisher's Exact Test was used to compare the sleep fractions of one sample group at

different time points. The entirety of the baseline was compared to the entirety of the

stimulation period unless otherwise stated through significance bars. Data from different

strains was either compared with the Kolmogorov-Smirnov-Test or the Welch Test. The

p-values can be taken from the respective Figure descriptions. In the graph depicted is the

mean +/- SEM unless otherwise stated. The box in the box plots represents the inter

quartile range with the median. The whiskers show the 10-90-percentile range. The

individual data points are furthermore plotted on top of the box.

**Acknowledgments**

We are grateful to Jonathan Packer and Robert Waterson for suggesting the use of the *zk673.11* promoter. We thank Cori Bargmann, Andrew M. Leifer, Andres Mariqc, Shai Shaham, Yun Zhang, Manuel Zimmer and the Caenorhabdis Genetics Center for strains. The strain PHX816 was generated by SunyBiotech. We are grateful to Kaveh Ashrafi, Ithai Rabinowitch, and Yoshinori Tanizawa for plasmids.

**References**

1.      Irwin MR. Why sleep is important for health: a psychoneuroimmunology perspective. Annu Rev Psychol. 2015;66:143-72. Epub 2014/07/26. doi: 10.1146/annurev-psych-010213-115205. PubMed PMID: 25061767; PubMed Central PMCID: PMCPMC4961463.
2.      McEwen BS, Karatsoreos IN. Sleep Deprivation and Circadian Disruption: Stress, Allostasis, and Allostatic Load. Sleep medicine clinics. 2015;10(1):1-10. Epub 2015/06/10. doi: 10.1016/j.jsmc.2014.11.007. PubMed PMID: 26055668.
3.      Drake CL, Roehrs T, Roth T. Insomnia causes, consequences, and therapeutics: an overview. Depression and anxiety. 2003;18(4):163-76. Epub 2003/12/09. doi: 10.1002/da.10151. PubMed PMID: 14661186.
4.      Knutson KL, Spiegel K, Penev P, Van Cauter E. The metabolic consequences of sleep deprivation. Sleep medicine reviews. 2007;11(3):163-78. Epub 2007/04/20. doi: 10.1016/j.smrv.2007.01.002. PubMed PMID: 17442599; PubMed Central PMCID: PMCPMC1991337.
5.      Van Cauter E, Spiegel K, Tasali E, Leproult R. Metabolic consequences of sleep and sleep loss. Sleep Med. 2008;9 Suppl 1:S23-8. Epub 2008/12/17. doi: 10.1016/S1389-9457(08)70013-3. PubMed PMID: 18929315; PubMed Central PMCID: PMCPMC4444051.
6.      Saper CB, Fuller PM, Pedersen NP, Lu J, Scammell TE. Sleep state switching. Neuron. 2010;68(6):1023-42. Epub 2010/12/22. doi: 10.1016/j.neuron.2010.11.032. PubMed PMID: 21172606; PubMed Central PMCID: PMC3026325.
7.      Campbell SS, Tobler I. Animal sleep: a review of sleep duration across phylogeny. Neurosci Biobehav Rev. 1984;8(3):269-300. Epub 1984/01/01. PubMed PMID: 6504414.
8.      Bringmann H. Sleep-Active Neurons: Conserved Motors of Sleep. Genetics. 2018;208(4):1279-89. Epub 2018/04/06. doi: 10.1534/genetics.117.300521. PubMed PMID: 29618588; PubMed Central PMCID: PMC5887131.
9.      Lu J, Sherman D, Devor M, Saper CB. A putative flip-flop switch for control of REM sleep. Nature. 2006;441(7093):589-94. Epub 2006/05/12. doi: 10.1038/nature04767. PubMed PMID: 16688184.

10.    Morairty S, Rainnie D, McCarley R, Greene R. Disinhibition of ventrolateral
preoptic area sleep-active neurons by adenosine: a new mechanism for sleep promotion.
Neuroscience. 2004;123(2):451-7. Epub 2003/12/31. PubMed PMID: 14698752.
11.    Saper CB, Scammell TE, Lu J. Hypothalamic regulation of sleep and circadian
rhythms. Nature. 2005;437(7063):1257-63. Epub 2005/10/28. doi: 10.1038/nature04284.
PubMed PMID: 16251950.
12.    Weber F, Dan Y. Circuit-based interrogation of sleep control. Nature.
2016;538(7623):51-9. Epub 2016/10/07. doi: 10.1038/nature19773. PubMed PMID:
27708309.
13.    Zhang Z, Zhong P, Hu F, Barger Z, Ren Y, Ding X, et al. An Excitatory Circuit in
the Perioculomotor Midbrain for Non-REM Sleep Control. Cell. 2019;177(5):1293-307
e16. Epub 2019/04/30. doi: 10.1016/j.cell.2019.03.041. PubMed PMID: 31031008.
14.    Borbely AA. A two process model of sleep regulation. Human neurobiology.
1982;1(3):195-204. Epub 1982/01/01. PubMed PMID: 7185792.
15.    Frank MG. Astroglial regulation of sleep homeostasis. Curr Opin Neurobiol.
2013;23(5):812-8. Epub 2013/03/23. doi: 10.1016/j.conb.2013.02.009. PubMed PMID:
23518138.
16.    Krueger JM, Frank MG, Wisor JP, Roy S. Sleep function: Toward elucidating an
enigma. Sleep medicine reviews. 2016;28:46-54. Epub 2015/10/09. doi:
10.1016/j.smrv.2015.08.005. PubMed PMID: 26447948; PubMed Central PMCID:
PMC4769986.
17.    Donlea JM, Pimentel D, Miesenbock G. Neuronal machinery of sleep
homeostasis in Drosophila. Neuron. 2014;81(4):860-72. Epub 2014/02/25. doi:
10.1016/j.neuron.2013.12.013. PubMed PMID: 24559676; PubMed Central PMCID:
PMC3969244.
18.    Pimentel D, Donlea JM, Talbot CB, Song SM, Thurston AJ, Miesenbock G.
Operation of a homeostatic sleep switch. Nature. 2016;536(7616):333-7. Epub
2016/08/04. doi: 10.1038/nature19055. PubMed PMID: 27487216; PubMed Central
PMCID: PMC4998959.
19.    Halasz P. Hierarchy of micro-arousals and the microstructure of sleep.
Neurophysiologie clinique = Clinical neurophysiology. 1998;28(6):461-75. Epub
1999/01/23. PubMed PMID: 9894227.
20.    Amzica F, Steriade M. Cellular substrates and laminar profile of sleep K-
complex. Neuroscience. 1998;82(3):671-86. Epub 1998/03/04. PubMed PMID: 9483527.
21.    Cash SS, Halgren E, Dehghani N, Rossetti AO, Thesen T, Wang C, et al. The
human K-complex represents an isolated cortical down-state. Science.
2009;324(5930):1084-7. Epub 2009/05/23. doi: 10.1126/science.1169626. PubMed
PMID: 19461004; PubMed Central PMCID: PMC3715654.
22.    Iwanir S, Tramm N, Nagy S, Wright C, Ish D, Biron D. The microarchitecture of
C. elegans behavior during lethargus: homeostatic bout dynamics, a typical body posture,
and regulation by a central neuron. sleep. 2013;36(3):385-95. Epub 2013/03/02. doi:
10.5665/sleep.2456. PubMed PMID: 23449971; PubMed Central PMCID: PMC3571756.
23.    Nagy S, Tramm N, Sanders J, Iwanir S, Shirley IA, Levine E, et al. Homeostasis
in C. elegans sleep is characterized by two behaviorally and genetically distinct
mechanisms. eLife. 2014;3:e04380. Epub 2014/12/05. doi: 10.7554/eLife.04380.
PubMed PMID: 25474127; PubMed Central PMCID: PMC4273442.

24.     Joiner WJ. Unraveling the Evolutionary Determinants of Sleep. Curr Biol.
2016;26(20):R1073-R87. Epub 2016/10/26. doi: 10.1016/j.cub.2016.08.068. PubMed
PMID: 27780049; PubMed Central PMCID: PMC5120870.
25.     Turek M, Lewandrowski I, Bringmann H. An AP2 transcription factor is required
for a sleep-active neuron to induce sleep-like quiescence in C. elegans. Curr Biol.
2013;23(22):2215-23. Epub 2013/11/05. doi: 10.1016/j.cub.2013.09.028. PubMed PMID:
24184105.
26.     Bennett HL, Khoruzhik Y, Hayden D, Huang H, Sanders J, Walsh MB, et al.
Normal sleep bouts are not essential for C. elegans survival and FoxO is important for
compensatory changes in sleep. BMC neuroscience. 2018;19(1):10. Epub 2018/03/11.
doi: 10.1186/s12868-018-0408-1. PubMed PMID: 29523076.
27.     Wu Y, Masurat F, Preis J, Bringmann H. Sleep Counteracts Aging Phenotypes to
Survive Starvation-Induced Developmental Arrest in C. elegans. Curr Biol.
2018;28(22):3610-24 e8. Epub 2018/11/13. doi: 10.1016/j.cub.2018.10.009. PubMed
PMID: 30416057; PubMed Central PMCID: PMCPMC6264389.
28.     White JG, Southgate E, Thomson JN, Brenner S. The structure of the nervous
system of the nematode Caenorhabditis elegans. Philos Trans R Soc Lond B Biol Sci.
1986;314(1165):1-340. Epub 1986/11/12. PubMed PMID: 22462104.
29.     Raizen DM, Zimmerman JE, Maycock MH, Ta UD, You YJ, Sundaram MV, et
al. Lethargus is a Caenorhabditis elegans sleep-like state. Nature. 2008;451(7178):569-
72. Epub 2008/01/11. doi: nature06535 [pii]10.1038/nature06535. PubMed PMID:
18185515.
30.     Trojanowski NF, Raizen DM. Call it Worm Sleep. Trends in neurosciences.
2016;39(2):54-62. Epub 2016/01/10. doi: 10.1016/j.tins.2015.12.005. PubMed PMID:
26747654; PubMed Central PMCID: PMC4738085.
31.     Kayser MS, Biron D. Sleep and Development in Genetically Tractable Model
Organisms. Genetics. 2016;203(1):21-33. Epub 2016/05/18. doi:
10.1534/genetics.116.189589. PubMed PMID: 27183564; PubMed Central PMCID:
PMC4858775.
32.     Cook SJ, Jarrell TA, Brittin CA, Wang Y, Bloniarz AE, Yakovlev MA, et al.
Whole-animal connectomes of both Caenorhabditis elegans sexes. Nature.
2019;571(7763):63-71. Epub 2019/07/05. doi: 10.1038/s41586-019-1352-7. PubMed
PMID: 31270481.
33.     Sulston JE, Horvitz HR. Post-embryonic cell lineages of the nematode,
Caenorhabditis elegans. Dev Biol. 1977;56(1):110-56. Epub 1977/03/01. doi: 0012-
1606(77)90158-0 [pii]. PubMed PMID: 838129.
34.     Cassada RC, Russell RL. The dauerlarva, a post-embryonic developmental variant
of the nematode Caenorhabditis elegans. Dev Biol. 1975;46(2):326-42. Epub 1975/10/01.
doi: 0012-1606(75)90109-8 [pii]. PubMed PMID: 1183723.
35.     Guo ZV, Hart AC, Ramanathan S. Optical interrogation of neural circuits in
Caenorhabditis elegans. Nat Methods. 2009;6(12):891-6. Epub 2009/11/10. doi:
nmeth.1397 [pii]
10.1038/nmeth.1397. PubMed PMID: 19898486.
36.     Piggott BJ, Liu J, Feng Z, Wescott SA, Xu XZ. The neural circuits and synaptic
mechanisms underlying motor initiation in C. elegans. Cell. 2011;147(4):922-33. Epub

2011/11/15. doi: 10.1016/j.cell.2011.08.053. PubMed PMID: 22078887; PubMed Central
PMCID: PMC3233480.
37.     Taylor SR, Santpere G, Reilly M, Glenwinkel L, Poff A, McWhirter R, et al.
Expression profiling of the mature <em>C. elegans</em> nervous system by single-cell
RNA-Sequencing. bioRxiv. 2019:737577. doi: 10.1101/737577.
38.     Alkema MJ, Hunter-Ensor M, Ringstad N, Horvitz HR. Tyramine Functions
independently of octopamine in the Caenorhabditis elegans nervous system. Neuron.
2005;46(2):247-60. Epub 2005/04/26. doi: S0896-6273(05)00167-4 [pii]
10.1016/j.neuron.2005.02.024. PubMed PMID: 15848803.
39.     Rogers C, Reale V, Kim K, Chatwin H, Li C, Evans P, et al. Inhibition of
Caenorhabditis elegans social feeding by FMRFamide-related peptide activation of NPR-
1. Nat Neurosci. 2003;6(11):1178-85. Epub 2003/10/14. doi: 10.1038/nn1140. PubMed
PMID: 14555955.
40.     Serrano-Saiz E, Poole RJ, Felton T, Zhang F, De La Cruz ED, Hobert O. Modular
control of glutamatergic neuronal identity in C. elegans by distinct homeodomain
proteins. Cell. 2013;155(3):659-73. Epub 2013/11/19. doi: 10.1016/j.cell.2013.09.052.
PubMed PMID: 24243022; PubMed Central PMCID: PMCPMC3855022.
41.     Lee RY, Sawin ER, Chalfie M, Horvitz HR, Avery L. EAT-4, a homolog of a
mammalian sodium-dependent inorganic phosphate cotransporter, is necessary for
glutamatergic neurotransmission in caenorhabditis elegans. J Neurosci. 1999;19(1):159-
67. Epub 1998/12/31. PubMed PMID: 9870947; PubMed Central PMCID:
PMCPMC3759158.
42.     Zimmer M, Gray JM, Pokala N, Chang AJ, Karow DS, Marletta MA, et al.
Neurons detect increases and decreases in oxygen levels using distinct guanylate
cyclases. Neuron. 2009;61(6):865-79. Epub 2009/03/28. doi:
10.1016/j.neuron.2009.02.013. PubMed PMID: 19323996; PubMed Central PMCID:
PMCPMC2760494.
43.     Sawin ER, Ranganathan R, Horvitz HR. C. elegans locomotory rate is modulated
by the environment through a dopaminergic pathway and by experience through a
serotonergic pathway. Neuron. 2000;26(3):619-31. Epub 2000/07/15. doi: S0896-
6273(00)81199-X [pii]. PubMed PMID: 10896158.
44.     Han X, Chow BY, Zhou H, Klapoetke NC, Chuong A, Rajimehr R, et al. A high-
light sensitivity optical neural silencer: development and application to optogenetic
control of non-human primate cortex. Frontiers in systems neuroscience. 2011;5:18. Epub
2011/08/04. doi: 10.3389/fnsys.2011.00018. PubMed PMID: 21811444; PubMed Central
PMCID: PMC3082132.
45.     Okazaki A, Takagi S. An optogenetic application of proton pump ArchT to C.
elegans cells. Neuroscience research. 2013;75(1):29-34. Epub 2012/10/10. doi:
10.1016/j.neures.2012.09.002. PubMed PMID: 23044183.
46.     Li W, Kang L, Piggott BJ, Feng Z, Xu XZ. The neural circuits and sensory
channels mediating harsh touch sensation in Caenorhabditis elegans. Nat Commun.
2011;2:315. Epub 2011/05/19. doi: 10.1038/ncomms1308. PubMed PMID: 21587232;
PubMed Central PMCID: PMC3098610.
47.     Cao J, Packer JS, Ramani V, Cusanovich DA, Huynh C, Daza R, et al.
Comprehensive single-cell transcriptional profiling of a multicellular organism. Science.

2017;357(6352):661-7. Epub 2017/08/19. doi: 10.1126/science.aam8940. PubMed
PMID: 28818938; PubMed Central PMCID: PMC5894354.
48.    Packer JS, Zhu Q, Huynh C, Sivaramakrishnan P, Preston E, Dueck H, et al. A
lineage-resolved molecular atlas of C. elegans embryogenesis at single-cell resolution.
Science. 2019. Epub 2019/09/07. doi: 10.1126/science.aax1971. PubMed PMID:
31488706.
49.    Chalfie M, Sulston JE, White JG, Southgate E, Thomson JN, Brenner S. The
neural circuit for touch sensitivity in Caenorhabditis elegans. J Neurosci. 1985;5(4):956-
64. Epub 1985/04/01. PubMed PMID: 3981252.
50.    Husson SJ, Costa WS, Wabnig S, Stirman JN, Watson JD, Spencer WC, et al.
Optogenetic analysis of a nociceptor neuron and network reveals ion channels acting
downstream of primary sensors. Curr Biol. 2012;22(9):743-52. Epub 2012/04/10. doi:
10.1016/j.cub.2012.02.066. PubMed PMID: 22483941; PubMed Central PMCID:
PMC3350619.
51.    Turek M, Besseling J, Spies JP, Konig S, Bringmann H. Sleep-active neuron
specification and sleep induction require FLP-11 neuropeptides to systemically induce
sleep. eLife. 2016;5. Epub 2016/03/08. doi: 10.7554/eLife.12499. PubMed PMID:
26949257; PubMed Central PMCID: PMC4805538.
52.    Katz M, Corson F, Iwanir S, Biron D, Shaham S. Glia Modulate a Neuronal
Circuit for Locomotion Suppression during Sleep in C. elegans. Cell reports.
2018;22(10):2575-83. Epub 2018/03/08. doi: 10.1016/j.celrep.2018.02.036. PubMed
PMID: 29514087; PubMed Central PMCID: PMCPMC5870883.
53.    Chalfie M, Hart AC, Rankin CH, Goodman MB. Assaying mechanosensation.
WormBook. 2014. Epub 2014/08/06. doi: 10.1895/wormbook.1.172.1. PubMed PMID:
25093996; PubMed Central PMCID: PMC4448936.
54.    Pokala N, Liu Q, Gordus A, Bargmann CI. Inducible and titratable silencing of
Caenorhabditis elegans neurons in vivo with histamine-gated chloride channels. Proc
Natl Acad Sci U S A. 2014;111(7):2770-5. Epub 2014/02/20. doi:
10.1073/pnas.1400615111. PubMed PMID: 24550306; PubMed Central PMCID:
PMCPMC3932931.
55.    Zheng Y, Brockie PJ, Mellem JE, Madsen DM, Maricq AV. Neuronal control of
locomotion in C. elegans is modified by a dominant mutation in the GLR-1 ionotropic
glutamate receptor. Neuron. 1999;24(2):347-61. Epub 1999/11/26. PubMed PMID:
10571229.
56.    Brockie PJ, Mellem JE, Hills T, Madsen DM, Maricq AV. The C. elegans
glutamate receptor subunit NMR-1 is required for slow NMDA-activated currents that
regulate reversal frequency during locomotion. Neuron. 2001;31(4):617-30. Epub
2001/09/08. PubMed PMID: 11545720.
57.    Nguyen JP, Shipley FB, Linder AN, Plummer GS, Liu M, Setru SU, et al. Whole-
brain calcium imaging with cellular resolution in freely behaving Caenorhabditis elegans.
Proc Natl Acad Sci U S A. 2016;113(8):E1074-81. Epub 2015/12/30. doi:
10.1073/pnas.1507110112. PubMed PMID: 26712014; PubMed Central PMCID:
PMCPMC4776509.
58.    Mahn M, Gibor L, Patil P, Cohen-Kashi Malina K, Oring S, Printz Y, et al. High-
efficiency optogenetic silencing with soma-targeted anion-conducting channelrhodopsins.

Nat Commun. 2018;9(1):4125. Epub 2018/10/10. doi: 10.1038/s41467-018-06511-8.
PubMed PMID: 30297821; PubMed Central PMCID: PMCPMC6175909.
59.    Richmond JE, Davis WS, Jorgensen EM. UNC-13 is required for synaptic vesicle
fusion in C. elegans. Nat Neurosci. 1999;2(11):959-64. Epub 1999/10/20. doi:
10.1038/14755. PubMed PMID: 10526333; PubMed Central PMCID:
PMCPMC2585767.
60.    Davis MW, Morton JJ, Carroll D, Jorgensen EM. Gene activation using FLP
recombinase in C. elegans. PLoS genetics. 2008;4(3):e1000028. Epub 2008/03/29. doi:
10.1371/journal.pgen.1000028. PubMed PMID: 18369447; PubMed Central PMCID:
PMCPMC2265415.
61.    Nagy S, Wright C, Tramm N, Labello N, Burov S, Biron D. A longitudinal study
of Caenorhabditis elegans larvae reveals a novel locomotion switch, regulated by Galphas
signaling. eLife. 2013;2:e00782. Epub 2013/07/11. doi: 10.7554/eLife.00782. PubMed
PMID: 23840929; PubMed Central PMCID: PMC3699835.
62.    Belfer SJ, Chuang HS, Freedman BL, Yuan J, Norton M, Bau HH, et al.
Caenorhabditis-in-drop array for monitoring C. elegans quiescent behavior. sleep.
2013;36(5):689-98G. Epub 2013/05/02. doi: 10.5665/sleep.2628. PubMed PMID:
23633751; PubMed Central PMCID: PMC3624823.
63.    Cirelli C, Bushey D, Hill S, Huber R, Kreber R, Ganetzky B, et al. Reduced sleep
in Drosophila Shaker mutants. Nature. 2005;434(7037):1087-92. Epub 2005/04/29. doi:
nature03486 [pii]10.1038/nature03486. PubMed PMID: 15858564.
64.    Koh K, Joiner WJ, Wu MN, Yue Z, Smith CJ, Sehgal A. Identification of
SLEEPLESS, a sleep-promoting factor. Science. 2008;321(5887):372-6. Epub
2008/07/19. doi: 10.1126/science.1155942. PubMed PMID: 18635795; PubMed Central
PMCID: PMC2771549.
65.    Kume K, Kume S, Park SK, Hirsh J, Jackson FR. Dopamine is a regulator of
arousal in the fruit fly. J Neurosci. 2005;25(32):7377-84. Epub 2005/08/12. doi:
10.1523/JNEUROSCI.2048-05.2005. PubMed PMID: 16093388.
66.    Singh K, Ju JY, Walsh MB, DiIorio MA, Hart AC. Deep Conservation of Genes
Required for Both Drosphila melanogaster and Caenorhabditis elegans Sleep Includes a
Role for Dopaminergic Signaling. sleep. 2014;37(9). PubMed PMID: 25142568.
67.    Schwarz J, Bringmann H. Reduced sleep-like quiescence in both hyperactive and
hypoactive mutants of the Galphaq Gene egl-30 during lethargus in Caenorhabditis
elegans. PLoS One. 2013;8(9):e75853. Epub 2013/09/28. doi:
10.1371/journal.pone.0075853. PubMed PMID: 24073282; PubMed Central PMCID:
PMC3779211.
68.    Huang H, Hayden DJ, Zhu CT, Bennett HL, Venkatachalam V, Skuja LL, et al.
Gap Junctions and NCA Cation Channels Are Critical for Developmentally Timed Sleep
and Arousal in Caenorhabditis elegans. Genetics. 2018;210(4):1369-81. Epub
2018/10/17. doi: 10.1534/genetics.118.301551. PubMed PMID: 30323068; PubMed
Central PMCID: PMCPMC6283151.
69.    Pirri JK, Alkema MJ. The neuroethology of C. elegans escape. Curr Opin
Neurobiol. 2012;22(2):187-93. Epub 2012/01/10. doi: 10.1016/j.conb.2011.12.007.
PubMed PMID: 22226513; PubMed Central PMCID: PMCPMC3437330.

70.    Nichols ALA, Eichler T, Latham R, Zimmer M. A global brain state underlies C.
elegans sleep behavior. Science. 2017;356(6344). Epub 2017/06/24. doi:
10.1126/science.aam6851. PubMed PMID: 28642382.
71.    Kaplan JM, Horvitz HR. A dual mechanosensory and chemosensory neuron in
Caenorhabditis elegans. Proc Natl Acad Sci U S A. 1993;90(6):2227-31. Epub
1993/03/15. PubMed PMID: 8460126.
72.    Gordus A, Pokala N, Levy S, Flavell SW, Bargmann CI. Feedback from network
states generates variability in a probabilistic olfactory circuit. Cell. 2015;161(2):215-27.
Epub 2015/03/17. doi: 10.1016/j.cell.2015.02.018. PubMed PMID: 25772698; PubMed
Central PMCID: PMC4821011.
73.    Stirman JN, Crane MM, Husson SJ, Wabnig S, Schultheis C, Gottschalk A, et al.
Real-time multimodal optical control of neurons and muscles in freely behaving
Caenorhabditis elegans. Nat Methods. 2011;8(2):153-8. Epub 2011/01/18. doi:
10.1038/nmeth.1555. PubMed PMID: 21240278; PubMed Central PMCID:
PMC3189501.
74.    Kawano T, Po MD, Gao S, Leung G, Ryu WS, Zhen M. An imbalancing act: gap
junctions reduce the backward motor circuit activity to bias C. elegans for forward
locomotion. Neuron. 2011;72(4):572-86. Epub 2011/11/22. doi:
10.1016/j.neuron.2011.09.005. PubMed PMID: 22099460.
75.    Kato S, Kaplan HS, Schrodel T, Skora S, Lindsay TH, Yemini E, et al. Global
brain dynamics embed the motor command sequence of Caenorhabditis elegans. Cell.
2015;163(3):656-69. Epub 2015/10/20. doi: 10.1016/j.cell.2015.09.034. PubMed PMID:
26478179.
76.    Steuer Costa W, Van der Auwera P, Glock C, Liewald JF, Bach M, Schuler C, et
al. A GABAergic and peptidergic sleep neuron as a locomotion stop neuron with
compartmentalized Ca2+ dynamics. Nat Commun. 2019;10(1):4095. Epub 2019/09/12.
doi: 10.1038/s41467-019-12098-5. PubMed PMID: 31506439.
77.    Skora S, Mende F, Zimmer M. Energy Scarcity Promotes a Brain-wide Sleep
State Modulated by Insulin Signaling in C. elegans. Cell reports. 2018;22(4):953-66.
Epub 2018/02/02. doi: 10.1016/j.celrep.2017.12.091. PubMed PMID: 29386137.
78.    Cho JY, Sternberg PW. Multilevel modulation of a sensory motor circuit during
C. elegans sleep and arousal. Cell. 2014;156(1-2):249-60. Epub 2014/01/21. doi:
10.1016/j.cell.2013.11.036. PubMed PMID: 24439380; PubMed Central PMCID:
PMC3962823.
79.    Schrodel T, Prevedel R, Aumayr K, Zimmer M, Vaziri A. Brain-wide 3D imaging
of neuronal activity in Caenorhabditis elegans with sculpted light. Nat Methods.
2013;10(10):1013-20. Epub 2013/09/10. doi: 10.1038/nmeth.2637. PubMed PMID:
24013820.
80.    Gallagher T, Kim J, Oldenbroek M, Kerr R, You YJ. ASI regulates satiety
quiescence in C. elegans. J Neurosci. 2013;33(23):9716-24. Epub 2013/06/07. doi:
10.1523/JNEUROSCI.4493-12.2013. PubMed PMID: 23739968; PubMed Central
PMCID: PMCPMC3726252.
81.    Goetting DL, Soto R, Van Buskirk C. Food-Dependent Plasticity in
Caenorhabditis elegans Stress-Induced Sleep Is Mediated by TOR-FOXA and TGF-beta
Signaling. Genetics. 2018. Epub 2018/06/22. doi: 10.1534/genetics.118.301204. PubMed
PMID: 29925566.

82. Gaus SE, Strecker RE, Tate BA, Parker RA, Saper CB. Ventrolateral preoptic
nucleus contains sleep-active, galaninergic neurons in multiple mammalian species.
Neuroscience. 2002;115(1):285-94. Epub 2002/10/29. PubMed PMID: 12401341.
83. Sherin JE, Shiromani PJ, McCarley RW, Saper CB. Activation of ventrolateral
preoptic neurons during sleep. Science. 1996;271(5246):216-9. Epub 1996/01/12.
PubMed PMID: 8539624.
84. Chrousos G, Vgontzas AN, Kritikou I. HPA Axis and Sleep. In: De Groot LJ,
Chrousos G, Dungan K, Feingold KR, Grossman A, Hershman JM, et al., editors.
Endotext. South Dartmouth (MA): Endotext.org; 2016.
85. Panossian LA, Avidan AY. Review of sleep disorders. The Medical clinics of
North America. 2009;93(2):407-25, ix. Epub 2009/03/11. doi:
10.1016/j.mcna.2008.09.001. PubMed PMID: 19272516.
86. Liu S, Liu Q, Tabuchi M, Wu MN. Sleep Drive Is Encoded by Neural Plastic
Changes in a Dedicated Circuit. Cell. 2016;165(6):1347-60. Epub 2016/05/24. doi:
10.1016/j.cell.2016.04.013. PubMed PMID: 27212237; PubMed Central PMCID:
PMC4892967.
87. Donlea JM, Pimentel D, Talbot CB, Kempf A, Omoto JJ, Hartenstein V, et al.
Recurrent Circuitry for Balancing Sleep Need and Sleep. Neuron. 2018;97(2):378-89 e4.
Epub 2018/01/09. doi: 10.1016/j.neuron.2017.12.016. PubMed PMID: 29307711;
PubMed Central PMCID: PMCPMC5779612.
88. Oikonomou G, Altermatt M, Zhang RW, Coughlin GM, Montz C, Gradinaru V, et
al. The Serotonergic Raphe Promote Sleep in Zebrafish and Mice. Neuron.
2019;103(4):686-701 e8. Epub 2019/06/30. doi: 10.1016/j.neuron.2019.05.038. PubMed
PMID: 31248729; PubMed Central PMCID: PMCPMC6706304.
89. Dijk DJ. Regulation and functional correlates of slow wave sleep. Journal of
clinical sleep medicine : JCSM : official publication of the American Academy of Sleep
Medicine. 2009;5(2 Suppl):S6-15. Epub 2009/12/17. PubMed PMID: 19998869; PubMed
Central PMCID: PMC2824213.
90. Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974;77(1):71-94.
Epub 1974/05/01. PubMed PMID: 4366476.
91. Ahringer A. Reverse genetics. WormBook, ed The C elegans Research
Community. 2006. doi: doi/10.1895/wormbook.1.47.1.
92. Merritt C, Seydoux G. Transgenic solutions for the germline. WormBook.
2010:1-21. Epub 2010/02/20. doi: 10.1895/wormbook.1.148.1. PubMed PMID:
20169625.
93. Redemann S, Schloissnig S, Ernst S, Pozniakowsky A, Ayloo S, Hyman AA, et
al. Codon adaptation-based control of protein expression in C. elegans. Nat Methods.
2011;8(3):250-2. Epub 2011/02/01. doi: nmeth.1565 [pii]10.1038/nmeth.1565. PubMed
PMID: 21278743.
94. Praitis V, Casey E, Collar D, Austin J. Creation of low-copy integrated transgenic
lines in Caenorhabditis elegans. Genetics. 2001;157(3):1217-26. Epub 2001/03/10.
PubMed PMID: 11238406; PubMed Central PMCID: PMC1461581.
95. Wilm T, Demel P, Koop HU, Schnabel H, Schnabel R. Ballistic transformation of
Caenorhabditis elegans. Gene. 1999;229(1-2):31-5. Epub 1999/03/30. PubMed PMID:
10095101.

96.     Bringmann H. Agarose hydrogel microcompartments for imaging sleep- and
wake-like behavior and nervous system development in Caenorhabditis elegans larvae. J
Neurosci Methods. 2011;201(1):78-88. Epub 2011/08/02. doi: S0165-0270(11)00418-3
[pii]10.1016/j.jneumeth.2011.07.013. PubMed PMID: 21801751.
97.     Turek M, Besseling J, Bringmann H. Agarose Microchambers for Long-term
Calcium Imaging of Caenorhabditis elegans. Journal of visualized experiments : JoVE.
2015;(100):e52742. Epub 2015/07/02. doi: 10.3791/52742. PubMed PMID: 26132740;
PubMed Central PMCID: PMC4544933.
98.     Stiernagle T. Maintenance of C. elegans. WormBook. 2006:1-11. Epub
2007/12/01. doi: 10.1895/wormbook.1.101.1. PubMed PMID: 18050451.
99.     Kim E, Sun L, Gabel CV, Fang-Yen C. Long-term imaging of Caenorhabditis
elegans using nanoparticle-mediated immobilization. PLoS One. 2013;8(1):e53419. Epub
2013/01/10. doi: 10.1371/journal.pone.0053419. PubMed PMID: 23301069; PubMed
Central PMCID: PMCPMC3536676.
100.    Hart AC, Sims S, Kaplan JM. Synaptic code for sensory modalities revealed by C.
elegans GLR-1 glutamate receptor. Nature. 1995;378(6552):82-5. Epub 1995/11/02. doi:
10.1038/378082a0. PubMed PMID: 7477294.
101.    Nguyen JP, Linder AN, Plummer GS, Shaevitz JW, Leifer AM. Automatically
tracking neurons in a moving and deforming brain. PLoS computational biology.
2017;13(5):e1005517. Epub 2017/05/26. doi: 10.1371/journal.pcbi.1005517. PubMed
PMID: 28545068; PubMed Central PMCID: PMC5436637.
102.    Schwarz J, Lewandrowski I, Bringmann H. Reduced activity of a sensory neuron
during a sleep-like state in Caenorhabditis elegans. Curr Biol. 2011;21(24):R983-4. Epub
2011/12/24. doi: S0960-9822(11)01207-3[pii]10.1016/j.cub.2011.10.046. PubMed
PMID: 22192827.
103.    Nagy S, Raizen DM, Biron D. Measurements of behavioral quiescence in
Caenorhabditis elegans. Methods. 2014. Epub 2014/03/20. doi:
10.1016/j.ymeth.2014.03.009. PubMed PMID: 24642199.

1720
1721

1722 **Supporting Information**

1723

**S1 Fig**

**Weak optogenetic RIM depolarization using the *gcy-13* promoter can induce RIS**

**activation or inhibition**

A)     Control experiments. Optogenetic depolarization of RIS presynaptic neurons

without the addition of ATR. For statistical calculations baseline neural activities (0 -

0.95min) were compared to neural activity levels during the stimulation period (1 -

1.95min). $*p < 0.05$, $**p < 0.01$, Wilcoxon Signed Rank Test for GCaMP. (S2 Data,

Sheet S1A)

B)      Optogenetic RIC depolarization induced an RIS activity increase outside of and

during lethargus. An average of all responsive trials is shown in this figure. Trials were

classified as responsive or non-responsive. In responsive trials a RIS activity increase

correlated with the onset of the stimulation period. In non-responsive trials no change in

RIS activity levels could be seen. n represents the number of animals tested, r represents

the number of trials. For statistical analysis RIS baseline activity levels (0 - 0.95min)

were compared to activity levels during (1 – 1.95min) and after (2 - 2.95min) the

stimulation. *p < 0.05, **p < 0.01, ***p < 0.001, Wilcoxon Signed Rank Test for

GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S1B)

C)      Depolarization of RIM using ReaChR expressed under the *gcy-13* promoter had

no net effect on RIS function. Neural baseline activity levels (0 - 0.95min) were

compared to neuronal levels during the stimulation (1 - 1.95min) and after the stimulation

(2 - 2.95min). *p < 0.05, **p < 0.01, ***p < 0.001, Wilcoxon Signed Rank Test for

GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S1C-E)

D)      RIM optogenetic depolarization using ReaChR expressed under the *gcy-13*

promoter induced either RIS activation or inhibition. Single trials were classified as

activating, if an activity increase in RIS correlated with onsets of optogenetic stimulation

periods. Trials were classified as inhibitory, if an activity decrease in RIS correlated with

onsets of optogenetic stimulation periods. n represents the number of animals tested, r

represents the number of trials. For statistical testing baseline neural activities (0 -

0.95min) were compared to neural activity levels during the stimulation period (1 -

1.55min). *p < 0.05, **p < 0.01, ***p < 0.001, Wilcoxon Signed Rank Test for GCaMP

and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S1C-E)

E)      Percentage of RIS activation and inhibition following optogenetic RIM activation

in different lethargus phases. Lethargus of each individual worm was split into 3 phases

of comparable size (lethargus onset, middle of lethargus and lethargus end). In each

interval, for all worms tested the amount of trials showing an RIS activation or RIS

inhibition were compared to the total amount of trials in this interval. (S2 Data, Sheet

S1C-E)

1761

 **S2 Fig**

 **RIM inhibition of RIS requires tyramine and FLP-18**

 Optogenetic RIM manipulations in these experiments were all performed with ReaChR

 expressed from the *tdc-1* promoter.

 A)      Optogenetic RIM depolarization in *flp-18(db99)* single mutants. Outside of

 lethargus, RIS inactivation caused by RIM optogenetic depolarization was reduced to

 37% of wild-type inhibition levels. During lethargus in *flp-18(db99)* mutants animals

 inhibition levels were only 25% of wild-type level. Neuronal activity levels before (0

 0.95min), during (1 - 1.95min), and after (2.5 – 2.95min) optogenetic RIM depolarization

 were compared. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, Wilcoxon Signed Rank Test for

 GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S2A)

 B)      Optogenetic RIM depolarization in *tdc-1(n3420)* single mutants. Outside of

 lethargus optogenetic RIM depolarization in *tdc-1(n3420)* single mutants no longer

 induced changes in RIS activity levels. During lethargus inhibition levels during the

 stimulation period only reached 40% of wild-type levels. Neuronal activity levels before

 (0 - 0.95min), during (1 - 1.95min) and after (2.5 – 2.95min) optogenetic RIM

 depolarization were compared. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, Wilcoxon Signed

 Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet

 S2B)

 C)      Optogenetic RIM depolarization in *flp-18(db99)* and *tdc-1(n3420)* double mutants

 had no effect on RIS function. Neuronal activity levels before (0 - 0.95min), during (1 -

 1.95min) and after (2.5 – 2.95min) optogenetic RIM depolarization were compared. *$p <$

 $0.05$, **$p < 0.01$, ***$p < 0.001$, Wilcoxon Signed Rank Test for GCaMP and speed,

 Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S2C)

 D)      Quantification of inhibition strength. RIS activity levels during optogenetic RIM

 depolarization in *flp-18(db99)*, *tdc-1(n3420)* and *flp-18(db99)*, *tdc-1(n3420)* double

 mutants were compared to wild-type levels. Wild-type data is depicted in Fig 1B, RIM

 panel. Inhibition strength was calculated by subtracting RIS activity levels during (1 –

 1.95min) the stimulation from activity levels before the stimulation (0 – 0.95min).

Samples were tested for normal distribution using the Shapiro-Wilk test. Wild type and

mutants were compared with a Welch Test. \*\*\*p < 0.001. (S2 Data, Sheet S2D-E)

E)      Quantification of RIS activity levels following RIM optogenetic depolarization.

Activity levels in *flp-18(db99)*, *tdc-1(n3420)* and *flp-18(db99)*; *tdc-1(n3420)* double

mutants were compared to wild-type levels. Wild-type data is depicted in Fig 1B in the

RIM panel. For statistical calculations RIS activity levels before the stimulation (0 –

0.95min) were subtracted from activity levels after the stimulation (2.5 – 2.95min).

Samples were tested for a normal distribution using the Saphiro-Wilk test. To compare

genotypes a Welch Test was performed for all conditions, except for the comparison of

activity levels between wild type and *tdc-1(n3420)* single mutants during lethargus. The

*tdc-1(n3420)* data was not normally distributed, and hence a Kolmogorov-Smirnov-Test

was used. \*\*\*p < 0.001. (S2 Data, Sheet S2D-E)

1804

 **S3 Fig**

 **RIM activation of RIS requires glutamatergic signaling**

A)      RIM optogenetic depolarization using ReAChR expressed under the *gcy-13*

promoter induced robust RIS activation in L4 larvae. In the L4 larvae, RIS activation by

RIM optogenetic depolarization was more robust compared with the same experiment in

L1 larvae. No trial selection was required. For statistical analysis RIS baseline activity

levels (0 - 0.95min) were compared to activity levels during (1 – 1.95min) and after (2 -

2.95min) the stimulation. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, Wilcoxon Signed Rank

Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S3A)

B)      The activating input of RIM optogenetic depolarization on RIS was almost

completely abolished in *eat-4(ky5)* mutants. For statistical analysis RIS baseline activity

levels (0-0.95min) were compared to activity levels during (1 – 1.95min) and after (2 -

2.95min) the stimulation. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, Wilcoxon Signed Rank

Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S3B)

1819

**S4 Fig**

**Activation of RIS by PVC is strongly enhanced during lethargus**

Optogenetic PVC depolarization in L2 larvae led to RIS activation outside of and during

lethargus. RIS activation during lethargus was strongly enhanced. Plotted data represent

the average over all experimental trials. Neural activity levels before the stimulation (0 -

0.95min) were compared to activity levels during the stimulation (1 - 1.95min). *p <

0.05, **p < 0.01, Wilcoxon Signed Rank Test. (S2 Data, Sheet S4)

1827

**S5 Fig**

**Optogenetic hyperpolarization experiments**

A)      Control experiments. Optogenetic hyperpolarization of RIS presynaptic neurons

without the addition of ATR. For statistical calculations baseline neural activities (0 -

0.95min) were compared to neural activity levels during the stimulation period (1 -

1.95min). *$p < 0.05$, **$p < 0.01$, Wilcoxon Signed Rank Test for GCaMP. (S2 Data,

Sheet S5A)

B)      Hyperpolarization of RIM using ArchT expressed under the *gcy-13* promoter had

no net effect on RIS function. Neural baseline activity levels (0 - 0.95min) were

compared to neuronal levels during the stimulation (1 - 1.95min) and after the stimulation

(2 - 2.95min). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, Wilcoxon Signed Rank Test for

GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S5B, C)

C)      RIM optogenetic hyperpolarization using ArchT expressed under the *gcy-13*

promoter caused a decrease in RIS activity levels in selected trials. Single trials were

classified as activating, if an activity increase in RIS occurred at the onset of the

optogenetic stimulation period. Trials were classified as inhibitory, if an activity decrease

in RIS occurred at the onset of the optogenetic stimulation period. n represents the

number of animals tested, r represents the number of trials. For statistical calculations

neural baseline activity levels (0 - 0.95min) were compared to levels during the

stimulation period (1 - 1.75min). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, Wilcoxon Signed

Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet

S5B, C)

D)      Simultaneous optogenetic hyperpolarization of CEP and URY neurons does not

induce changes in RIS activity levels. For statistical testing, baseline neural activities (2 -

2.95min) were compared to neural activity levels during the stimulation period (3 –

3.95min) and after the stimulation (6 – 6.95min). **$p < 0.01$, ***$p < 0.001$, Wilcoxon

Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data,

Sheet S5D)

1856

1857 **S6 Fig**

1858 *zk673.11* **is expressed in PVC, RID and cholinergic motor neurons**

1859 A-B) Expression of *nmr-1* and *zk673.11* only overlaps in PVC in the tail.

1860 C-D) Expression of *nmr-1* and *zk673.11* does not overlap in head neurons.

1861

**S7 Fig**

**PVC has multiple functions**

A)      PVC excitability remained unchanged during lethargus. Experiments were

performed in immobilized L1 larvae to ensure PVC-specific green light illumination. A

long baseline of 10 minutes was used to achieve stable baseline conditions. Activity

levels of PVC during optogenetic depolarization were indistinguishable outside and

during lethargus. PVC displayed a negative rebound transient after optogenetic

depolarization. However, there was no difference in the amount of negative rebound

outside and during lethargus. (S2 Data, Sheet S7A)

B)      RIS showed a rebound after mechanical stimulation. This rebound was stronger in

worms during lethargus and only during lethargus the RIS rebound was accompanied by

a strongly increased immobilization of worms. $**p < 0.01$, $***p < 0.001$, Wilcoxon

Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data,

Sheet S7B)

C-D) Effects of PVC stimulation on AVB activity. L1 larvae were immobilized for

optogenetic experiments to ensure cell-specific stimulation of PVC. AVB activated upon

optogenetic PVC depolarization with the same response strength during and outside of

lethargus. AVB displayed an oscillatory activity pattern in 44% of all trials in worms

outside of lethargus. AVB activity oscillated in 70% of all trials during lethargus. $*p <$

$0.05$, $**p < 0.01$, Wilcoxon Signed Rank Test for GCaMP. (S2 Data, Sheet S7C-D)

1882

**S8 Fig**

**Effects of Optogenetic RIS activation and inhibition on PVC and RIM activity**

A) RIS depolarizes during optogenetic activation in fixed animals. As controls, experiments were performed in the absence of ATR. ***p < 0.001, Wilcoxon Signed Rank Test. (S2 Data, Sheet S8A)

B) RIS hyperpolarization led to a weak PVC depolarization outside and during lethargus. For statistical calculations neural activities before the stimulation period (0 - 1min) were compared to activity levels during the stimulation period (1 - 2min). *p < 0.05, **p < 0.01, compared before and during stimulation, Wilcoxon Signed Rank Test. (S2 Data, Sheet S8B)

C-D) Sample trace of RIS activity and worm locomotion behavior 3 hours before shedding of the cuticle of *aptf-1(gk794)* and *flp-11(tm2705)* mutants. (S2 Data, Sheet S8C and S8D)

E-F) *flp-11(tm2705)* mutants have a significantly increased number of short RIS peaks that do not correlate with sleep. E) **p < 0.01, ***p < 0.001, Welch Test. F) **p < 0.01, Kolmogorov-Smirnov-Test. (S1 Data, Sheet 3G-I)

G) Optogenetic RIS depolarization has no effect on RIM activity outside of and during lethargus. Neuronal activity levels before (0 - 0.95min) and during (1 -1.95min) the stimulation period was compared. *p < 0.05, **p < 0.01, ***p < 0.001, Wilcoxon Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S8G)

H) Optogenetic RIS hyperpolarization induced increased RIM activity both outside of and during lethargus. Measurements were performed in immobilized L1 larvae to reduce measurement noise. Activity levels during baseline measurements (0 – 0.95min) were compared to levels during optogenetic RIS manipulation (1 - 1.95min). *p < 0.05, Wilcoxon Signed Rank Test for GCaMP. (S2 Data, Sheet S8H)

**S9 Fig**

**Command interneurons are required for RIS activation and sleep induction**

A) RIS activation in sleep bouts was strongly reduced in command interneuron-ablated worms. Samples were tested for normal distribution using the Saphiro-Wilk test. $*p < 0.05$, Welch Test. (S2 Data, Sheet S9A-C)

B) Command interneuron-ablated worms moved much slower than wild-type worms. Command interneurons were genetically ablated by expressing ICE from the *nmr-1* promoter. Samples were tested for normal distribution using the Saphiro-Wilk test. $***p < 0.001$, Welch Test for the wake condition and Kolmogorov-Smirnov-Test for the sleep condition. (S2 Data, Sheet S9A-C)

C) Sample traces of RIS activity levels and worm locomotion behaviors outside of and during lethargus in command interneuron-ablated worms and wild-type worms. In command interneuron-ablated worms quiescence bouts occurred only around the middle of the lethargus period. (S2 Data, Sheet S9A-C)

D-E) Mosaic analysis of worms expressing an extrachromosomal array of *nmr-1::ArchT*. Worm were selected that expressed the transgene only in head neurons (D) or head neurons and PVC (E). Neuronal activity levels before (2 - 2.95min) and during (3 - 3.95min) the stimulation period was compared. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, Wilcoxon Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S9D and S9E)

**S10 Fig**

**Glutamatergic signaling is required for sleep induction**

A-D) Sleep bout analysis of *eat-4(ky5)* mutant larvae. *eat-4(ky5)* animals lacked

significant RIS activation at sleep bout onset. Consistent with this finding, mutant worms

displayed a strong reduction in quiescence during lethargus. Samples were tested for a

normal distribution using the Saphiro-Wilk test. $**p < 0.01$, $***p < 0.001$, Welch Test

for comparisons of sleep bout lengths, sleep bout frequencies and sleep fractions.

Wilcoxon Signed Rank Test for quantifications of RIS activity levels in sleep bouts. (S2

Data, Sheet S10A-D)

E-I) Sleep bout analysis of *nmr-1(ak4)* mutant animals. RIS activity levels in sleep bouts

were slightly reduced in the mutant. *nmr-1(ak4)* mutants did not show a reduced amount

of quiescence during lethargus. Samples were tested for a normal distribution using the

Saphiro-Wilk test. $*p < 0.05$, Welch Test for comparisons of sleep bout frequencies,

sleep fractions, maximum RIS activity levels in sleep bouts and RIS activity levels at the

end of sleep bouts. Kolmogorov-Smirnov-Test for the comparison of sleep bout lengths.

(S2 Data, Sheet S10E-I)

1945

**S11 Fig**

**RIS rebound activation following optogenetic hyperpolarization requires synaptic transmission**

A)      RIS GCaMP transient intensities in wild-type worms are correlated with the length of the preceding motion bout. The longer the preceding motion bout, the stronger the RIS activation. (S2 Data, Sheet S11A)

B)      RIM was inhibited during and post hyperpolarization. *p < 0.05, **p < 0.01, ***p < 0.001, Wilcoxon Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S11B)

C)      RIS was optogenetically hyperpolarized with stimuli lasting for 48 seconds (C), 5 minutes (D) or 10 minutes (E). Worms not showing a rebound activation transient were excluded from the analysis, which was no worm for 48 seconds, 1 out of 7 worms for 5 minutes and 1 out of 13 worms for 10 minutes stimulation experiments. Data from these plots were used to generate a dose-response curve of optogenetic RIS hyperpolarization (Fig 6D-E). *p < 0.05, **p < 0.01, ***p < 0.001, Wilcoxon Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S1 Data, Sheet 6D,E)

F-H)    Following optogenetic hyperpolarization, RIS displayed strong rebound activation during lethargus (F). Rebound activation was abolished in a strain that is deficient for neurotransmission specifically in RIS (*flp-11::TetX*). (G) Rebound activation was abolished also by a mutation which impaired global synaptic transmission (*unc-13(s69)*) (H) *p < 0.05, **p < 0.01, Wilcoxon Signed Rank Test. (S2 Data, Sheet S11F-H)

I)      RIS showed a negative rebound following its own optogenetic depolarization. The strength of the negative rebound transient depended on the lethargus status of the worm. Worms during lethargus displayed a three times stronger negative rebound compared to worms outside of lethargus. *p < 0.05, **p < 0.01, ***p < 0.001, Wilcoxon Signed Rank Test for GCaMP and speed, Fisher's Exact Test for sleep fraction. (S2 Data, Sheet S11I)

**S12 Fig**

**RIM baseline activity levels are dampened during lethargus independently of RIS**

A)      Sample traces of RIM transient frequencies, RIM baseline activities and worm locomotion behaviors outside of and during lethargus in wild-type worms and *aptf-1(gk794)* mutants. (S2 Data, Sheet S12)

B)      Wild-type worms, but not *aptf-1(gk794)* mutant worms display changes in RIM transient frequencies across lethargus. Transient frequencies were assessed manually. To be counted as a transient, RIM activity levels had to be at least twice as high as baseline activity levels. ***$p < 0.001$ Kolmogorov-Smirnov-Test for wild-type condition, Welch Test for mutant condition. (S2 Data, Sheet S12)

C)      The reduction of RIM baseline activity levels during lethargus is preserved in *aptf-1(gk794)* mutants. **$p < 0.01$, Wilcoxon Signed Rank Test. (S2 Data, Sheet S12)

1984 **S13 Fig**

1985 **Assaying gentle tail touch reveals an inhibitory role of RIM on RIS**

1986 A) RIM ablation increases the reinstating of immobility following gentle tail touch during

1987 lethargus. *p < 0.05. Kolmogorov-Smirnov-Test. (S2 Data, Sheet S13A)

1988 B) RIM ablation increases RIS activation in response to gentle tail touch. **p < 0.01.

1989 Kolmogorov-Smirnov-Test. (S2 Data, Sheet S13B)

1990

1991 **S1 Text**

1992 A list of strains which were used during this study

1993

1994 **S2 Text**

1995 A list of generated constructs during this study

1996

1997 **S3 Text**

1998 Sequence of the strain PHX816, which was generated during this study

1999

2000 **S1 Table**

2001 List of primers, which were used during this study

2002

2003 **S2 Table**

2004 List of plasmids, which were used during this study

2005

2006 **S3 Table**

2007 Experimental details of all optogenetic experiments conducted during this study

2008

2009 **S1 Data**

2010 Raw data for all experiments from the main figures (Figs 1-9)

2011

2012 **S2 Data**

2013 Raw data for the experiments from the supporting figures (S1-S13 Figs)

**A)** outside of lethargus

forward locomotion stop? reverse

**B)** during lethargus

forward sleep bout reverse

→ synaptic activation

⊣ synaptic inhibition

■—■ gap juction