Translatomic database of cortical astroglia across male and female mouse development

reveals two distinct developmental phenotypes

Gareth M. Rurak[1], Stephanie Simard[1], François Charih[4], Amanda Van Geel[1], John

Stead[1], Barbara Woodside[1,3], James R. Green[4], Gianfilippo Coppola*[2], Natalina

Salmaso*[1,2]

Running Title:

Affiliations: 1-Department of Neuroscience, Carleton University, Ottawa, Ontario;

2-Child Study Center, Yale University; 3- Concordia University, Montreal, Quebec

4-Department of Systems and Computer Engineering, Carleton University, Ottawa,

Ontario

*co-corresponding authors

Authors for Correspondence:

Natalina Salmaso

Carleton University, 1125 Colonel By Drive, Ottawa, Ontario, K1S 5B6, Canada

Tel: 613-520 2600

Email: natalina.salmaso@carleton.ca

Gianfilippo Coppola

Yale University, 333 Cedar st, New Haven, CT 06511, USA

Email: gianfilippo.coppola@yale.edu

Abstract

Astroglial cells are emerging as key players in the development and homeostatic maintenance of neurons and neuronal networks. Astroglial cell functions are critical to neuronal migration and maturation, myelination, and synapse dynamics, however little is known about astroglial phenotypic changes over development. Furthermore, astroglial cells express steroid hormone receptors and show rapid responses to hormonal manipulations, however, despite important sex differences in telencephalic regions such as the cortex and hippocampus few studies have examined sex differences in astroglial cells in development and outside of the hypothalamus and amygdala. To phenotype cortical astroglial cells across postnatal development while considering potential sex differences, we used translating ribosome affinity purification together with RNA sequencing (TRAPseq) and immunohistochemistry to phenotype the entire astroglial translatome in males and females at key developmental time points: P1, P4, P7, P14, P35 and in adulthood. Overall, we found two distinct astroglial phenotypes between early (P1-P7) and late development (P14-Adult). We also noted that although astroglia show few basal sex differences in adulthood, they show significant sex differences in developmental gene expression patterns, with peak sex differences observed at P7. At least part of the sex differences observed at P7 appear to be due to males reaching a more mature astroglial phenotype earlier than females. Together, these data clearly delineate and phenotype astroglia across development and identify sex differences in astroglial developmental programs. Importantly, these developmental sex differences could have an impact on the construction and maintenance of neuronal networks and potential developmental windows of vulnerability to neurologic and psychiatric disease.

Acknowledgements

We would like to thank our funding providers, the Carleton University Discovery

Grant to N.S., Natural Sciences and Engineering Research Council of Canada Discovery

Grant to N.S., the Canadian Foundation for Innovation to N.S., and the Canadian

Research Chairs to N.S..

We thank Dr. Julianna Tomlinson, Dr. Michael Schlossmacher and his laboratory

members at the University of Ottawa for providing us with infrastructural support and

more. Finally, we would like to thank Dr. Alfonso Abizaid for his invaluable input into the

planning and preparation of this research project.

Introduction

The cerebral cortex is a complex network of layered neuronal cells that uses a
balance of excitatory and inhibitory signals to communicate both within the cerebral cortex
itself and to lower brain regions[1]. It is the most recently evolved component of the
mammalian central nervous system and is uniquely involved in higher order cognitive
functions[2] that are central to cognitive and emotional processing. The complex layered
structure of the cerebral cortex emerges during embryonic development in an "inside-out"
pattern consisting of radial glial cell (RGC)-assisted neuroblast migration to lower layers
(i.e. VI) before the upper layers are formed [3,4]. The appropriate migration of developing
neuroblasts is an integral part of the maturation of the cerebral cortex. Early insults or
perturbations during this critical period of development may disrupt the gross organization
of excitatory and inhibitory networks leading to neurodevelopmental disorders including
autism spectrum disorders (ASD), schizophrenia, and mood disorders such as bipolar
disorder[5,6]. Importantly, even subtle modifications in the number or functionality of cortical
synapses may lead to serious behavioural consequences and have been related to a
number of psychiatric diseases[7].

In human populations the prevalence of many, if not all, neurodevelopmental
disorders and mood disorders related to cortical function is dependent on sex [8] .Thus
understanding sex differences in the organization and function of the cerebral cortex is
critical to understanding the neurodevelopmental underpinnings of psychiatric disease.
There is a great deal of literature that suggests that gonadal hormones can directly
modulate disease states associated with schizophrenia, anxiety and mood disorders[9,10].
Similarly, in rodent models, basal changes in behaviours such as in the forced swim task
have been demonstrated across the estrous cycle[11]. Together, these data support an
important role for ovarian hormones and in particular, estrogen, in mediating those
behaviours related to higher order cognitive functioning in situ. Effects on behaviour and
physiology that depend on current levels of hormones such as those described above are
referred to as activational. The sex steroid milieu of embryonic and neonatal life, however,
can also have organizational effects. These are changes induced in this early stage of
development that persist even when the hormonal state producing them has dissipated.

Testosterone secreted from the testes of XY-male embryos acting either directly or via
aromatisation to estradiol is the primary organizing influence in the sexual differentiation
of the brain[12,13]; The role of estradiol in mediating organizational sex differences through
neuronal estrogen receptor (ERs) signaling has been extensively studied in the
hypothalamus and amygdala. For example, in the medial pre-optic area (MPOA),
estrogen receptor signalling induces prostaglandin-mediated synapse formation through
microglial and astroglial-mediated functions[13]. Neuroanatomical and functional sex
differences are also well defined in various neuronal populations of the hypothalamus[14],
amygdala[15], and hippocampus[16]. The cerebral cortex, however, has received less
attention than other telencephalic regions[17].

While it is possible that sex differences in cortical network functioning may be
completely mediated by direct hormonal modulation, sex differences in the prevalence of
anxiety disorders[9] and ASD[18] are apparent prior to puberty, and pro-dromal behaviours
associated with psychosis also show sex differences early in development[19], suggesting
that organizational sex differences (presumably in cortical networks) may be involved in
differences in risk or vulnerability for psychiatric disease.

Astroglial cells have recently emerged as key mediators in brain development and
the etiology of neurodevelopmental and mood disorders has been related to astroglial-
mediated functions such as neuroblast migration and synapse formation, maturation, and
elimination[20,21]. Cortical astroglial cells are a heterogenous cell population traditionally
categorised into three main subtypes easily distinguished by morphology, although recent
work has suggested that many more subtypes exist when astroglial cells are categorised
through single cell RNAseq or other gene and protein expression methods[22,23]. The first
subtype of astroglial cells to emerge in development are RGCs[24]. In addition to providing
scaffolding for migrating neuroblasts, astroglial cells exhibit stem-like potential in early
development and following injury that allows them to give rise to other neuronal cells
before maturing and populating the cortex as fibrous or protoplasmic astroglial cells[25,26].
As astroglia mature, their stem cell potential diminishes, as marked by an increase in
glutamate synthetase and decreased expression of Sox2 and vimentin[27]. The two main
mature morphological subtypes in the cortex are characterized as protoplasmic; canonical

astroglia with large domains of many fine processes, and fibrous; astroglial cells with
many long fibrous extensions[28]. These mature astroglial cell populations are responsible
for the construction and maintenance of both excitatory[29] and inhibitory[30] synapses in the
developing neocortex, modulating neuron signaling and maturation[31], and the elimination
of synapses during cortical maturation[21]. Work from the Barres laboratory characterised
astroglial cells in brain development using next generation sequencing techniques,
however these studies did not examine the contributions of biological sex, or early
developmental periods[32]. Astroglial cells express ERα[33], ERβ[34], and G-protein coupled
ERs[35], and although the full extent of ER functionality in astroglial cells is not
understood[36], astroglia show rapid responses to hormones across various states and
manipulations[37].

Because astroglial cells contribute to cortical development through neuroblast
migration, neuronal network integration, and synapse formation, maturation and
elimination, and because astroglia show rapid responses to hormones, we hypothesized
that cortical astroglial cells may 1) show phenotypic sex differences both overall and
across development and 2) execute developmental functions in a sexually dimorphic way.
To address this, we employed a transgenic mouse model expressing enhanced green
fluorescent protein (eGFP) fused to the L10 ribosome under the pan-astroglial cell
promoter AldH-L1 and used immunohistochemistry and TRAPseq to phenotype the
astroglial translatome and protein expression. We examined time points pertaining to
critical periods of postnatal neocortical development.
141
142

Results

Our initial characterization of neocortical astroglial cells set to measure total

numbers of neocortical astroglial cells (astroglial cell number) across development in both

males and females. We used unbiased stereology to quantify eGFP+ cells (eGFP was

expressed under the control of the pan-astroglial promoter, AldH1l1). No sex differences

in total astroglial number were observed ($F_{1,41}$=0.716, p=0.402, Hη=0.017) (Figure 1A),

however there was a main effect of age on this measure ($F_{5,41}$=58.313, p<0.001,

Hη=0.874), whereby the number of astroglial cells plateaued in both males or females

after P14 (Figure 1A). Additionally, neocortical volume did not exhibit any significant

sexual dimorphism ($F_{1,41}$=0.009, p=0.923, Hη=0) (Figure 1B). Not surprisingly, we

observed a significant increase of brain volume with age (($F_{5,41}$=135.181, p<0.001,

Hη=0.943) (Figure 1B).

Astroglial cells show a great deal of morphological heterogeneity that are typically

associated with functionality (Sofroniew & Vinters, 2010). One difference in morphology

that is seen during cortical development is an early abundance of the radial phenotype

that is replaced by protoplasmic astroglia over development. We assessed gross

morphological changes in development (from radial to non-radial) across age to ascertain

whether sex differences exist. eGFP+ cells were characterized as radial or non-radial

based on the presence of a leading, radial process in the neocortex. No significant sex

effects were found ($F_{1,41}$=0.004, p=0.953, Hη=0), however there was a significant effect

of age ($F_{5,41}$=89.291, p<0.001, Hη=0.916) (Figure 1C). Nearly all (~95% in both males

and females) astroglial cells in the neocortex at P1 exhibited radial morphology (Figure

1C).

To generate a more complete characterization of neocortical astroglia over

development and in relation to sex, we employed TRAPseq to assess the astroglial-

specific translatome in males and females at each time point. Gene expression levels (as

RPKM) for all expressed genes can be found in our searchable database at http://cu-

bic.ca/stardb (see Methods). To validate our TRAPseq methodology, we used q-RT-PCR

to compare the unbound fragment to its bound counterpart on levels of two astroglial

genes (*GFAP and Glul*) and two neuronal genes (*Rbfox3 and Pvalb*). As expected, we

found a large increase in astroglial gene expression in the bound fragment and a large

reduction in neuronal-specific gene expression, suggesting a successful enrichment of

the astroglial translatome (Supplemental Figure 3). In order to further characterise

astroglial-specific markers by age and sex, we plotted RPKM data for each of the top 20

genes expressed only in postnatal neocortical astroglia from both the Cahoy dataset[32],

28 genes from the Zhang dataset[38] and 48 genes from the Rowitch dataset that identified

layer-specific astroglial cell gene expression (Bayaktar et al. 2020) that were non-

overlapping (Supplemental Figures 4, 5 & 6). Interestingly, expression of most astroglia-

specific genes from the Zhang and Cahoy datasets (27/48) increases across

development, for example *CYP4F14*, a Vitamin E hydroxylase and *SLC39A12*, a zinc

transporter involved in the pathophysiology of schizophrenia[39] both have low expression

levels early on and rise to adult levels at P14 (Supplemental Figure 4). Only one astroglial

gene decreased in expression over development, *Pla2g3,* a gene involved in oxidative

stress that has been linked to Alzheimer's disease[40].

Supplemental Figure 5 depicts the 14/48 astroglial genes that show stable

expression over development and finally, Supplemental Figure 6 depicts those genes

(8/48) that show consistent low expression over the entire developmental period

assessed. Interestingly, we did not find any apparent sex differences in these astroglial-

specific genes, although a few appear as DEGs at specific time points, for example, the

cytokine Il33 is a DEG at P7 (lower in females- see below). Supplemental figure 7 shows

astroglial cell gene expression for Layers 2-4 and deep layers 5-6, supplemental figure 8

shows upper-layer biased astroglial cell gene expression across development, and

supplemental figure supplementary figure 9 shows gene expression for genes identified

with the AST2, AST3 subtypes.

Astroglial cell show age related differences in translatome

Next, we examined the overall sample-to-sample relationship by hierarchical

clustering to identify patterns throughout development and across sex. Clustering shows

a pattern of organization that is primarily affected by age, with P1, P4 and P7 (early) and

P14, P35 and adult mice (late) showing clustered expression patterns that differ from the

early time points (Figure 2A). We manually sorted samples by sex and age and although

some organization by sex is noted in Figure 2B, this is not as clearly demarcated or as

extensive as that observed by age. Indeed, when we collapsed gene expression levels

across all time points, only 16 genes were differentially expressed between female and

males (Supplementary Data File 2) and a subset of these sex differences in gene

expression is for genes are located on the sex chromosomes, suggesting little to no sex

differences in neocortical astroglia across all time points overall. However, when we look

at sex differences over time, we do note sex differences in developmental patterns. It is

clear that there are two major neocortical astroglial phenotypes across age: an early (P1-

P7) and late (P14-adult) phenotype. However, male samples show an "intermediate"

cluster and begin to show the late or mature astroglial phenotype at P7, whereas females

show this phenotype at P14 (Figure 2B).

To validate the early and late phenotype that we found using TRAPseq, we

compared TRAP extracted RNA on six genes differentially expressed between the early

and late timepoints, 3 associated with synapse/network maturation (*Gja1, Lynx1, Grin2c*)

and three steroid hormone receptors (*Ar, Pgr, Esr1*). Each of these showed a similar

pattern of results between TRAPseq and qPCR (Supplemental Figure 10), which is

highlighted by the significant positive correlations shown in Supplemental Figure 11. In

addition, we further validated the early and late astroglial cell phenotype, by assessing

neocortical astroglial cells protein expression using immunohistochemistry of common

astroglial markers at similar timepoints. Using the Aldhl1-L10 eGFP+ mice, we quantified

expression of the astroglia-specific cytoskeletal filament protein glial acidic fibrillary

protein, (GFAP). GFAP expression was not sexually dimorphic across development

($F_{1,41}=1.259$, $p=0.268$, $H\eta=0.029$), however, there was a main effect of age ($F_{5,41}=32.271$,

$p<0.001$, $H\eta=0.793$) (Figure 3A). The number of GFAP+ cells reached peak levels at P14

and remained stable into P35 and adulthood (Figure 3A). Prior to P14, no more than 15%

of eGFP+ cells co-expressed GFAP, from P14-onwards approximately 25% of eGFP+

cells expressed GFAP, consistent with previous work that has demonstrated that a large

proportion of cortical astroglia do not express GFAP under basal conditions (Simard et

al., 2018). Next, we quantified the number of cells that expressed vimentin, another

intermediate filament protein in astroglial cells, that is generally associated with a stem-
like radial sub-type of astroglial cells. A significant interaction of sex by age was found
($F_{5,41}$=3.871, p=0.006, Hη=0.321) (Figure 3B,D), such that females showed significantly
more vimentin+ cells compared with males at P7 (t=3.41, p=0.011) and males showed
significantly more vimentin+ cells in adulthood (t=3.895, p=0.005) (Figure 3B). Because
vimentin expression is associated with astroglial cell neurogenic and stem cell potential
and because we observed sex differences in vimentin expression, we quantified the
proportion of actively dividing eGFP+ cells across the postnatal period. Using a marker of
active s-phase cell division, Ki67, we quantified the percent of Ki67+eGFP+ cells. Ki67
was present in less than 2% of the astroglial cell population throughout the postnatal
period in both males and females (Figure 3C). There were no significant sex differences
across development ($F_{1,41}$=0.862, p=0.358, Hη=0.021), however there was a main effect
of age ($F_{5,41}$=26.799, p<0.001, Hη=0.766). Importantly, the findings here that the number
of astroglial cells expressing vimentin show sex and age differences, and proportion of
astroglia expressing GFAP and Ki67 show distinct early and late expression closely
resemble the expression patterns revealed by TRAPseq analysis.

The developmental pattern of the astroglial translatome is sexually dimorphic
To further characterize the *"early"* and *"late"* phenotype observed in the astroglial cell
translatome, we examined the number and proportion of up- and downregulated genes
in the early versus late developmental phenotypes, both overall and by sex (Figure 4A).
The early versus late phenotype in males showed 8983 DEGs whereas the female early
versus late phenotype had 9688 DEGs (full list of DEGs in Supplementary Data File 1).
Interestingly, the proportion of up- or downregulated genes was similar, with males and
females showing 62% and 55% of DEGs in the early versus late phenotype being
upregulated, respectively. When we collapsed the DEGs across sex to generate a general
early vs late phenotype for astroglial cells, the number of DEGs reaches 12131 with 58%
of those genes being upregulated in the late phenotype (Figure 4A). This suggests that
the difference in the early and late developmental phenotype is not merely reflective of
turning off developmental genes but rather is a combination of downregulating genes

associated with early development and turning on genes associated with later neocortical
development and basal functioning in early adulthood. Some key DEGs stand out as they
have not previously been functionally described in astroglial cells. In particular, *Trim67*,
an important and evolutionarily preserved guidance cue for neurite outgrowth[41], is high in
early development and, as expected, decreases with age. Additionally, *Ntng1*, codes for
Netrin G1, another guidance cue that is paramount to developing neuroblasts, is higher
during the "late" developmental phenotype in astroglial cells. Differential *Ntng1*
expression appears when examining both the overall and the sex-specific early versus
late astroglial cell phenotype. *Ntng1* has been implicated in the etiology of familial
schizophrenia[42] and *Ntng1* mRNA is in lower cortical samples from individuals with
schizophrenia and autism[43].

We compared the proportion of overlap between the female-specific and male-
specific DEGs in the early vs late phenotype. Of the 11415 unique DEGs, approximately
64% overlap between males and females, however about 21% of total DEGs are specific
to females and only about 15% are specific to males (Figure 4B). When we further
characterize the sex-specific canonical pathways (CPs) and gene ontology (GO) by gene
set over-representation, a proportion of the top five female-specific CPs are related to cell
cycle/mitosis, marked by CPs in ribosome function. In comparison, male-specific CPs are
related to oxidative phosphorylation and small molecule transport. With respect to GO, a
large proportion of the top five female-specific and male-specific GOs are related to
intracellular parts and organelles. This suggests that there are distinct developmental
patterns in neocortical astroglial cells that emerge between the sexes. To broaden the
understanding of sex differences, we identified DEGs of females versus males at each
time point (Figure 4C, Supplementary Data File 2). At P1, there are only 20 DEGs,
however there are 183 DEGs at P7 and 100 DEGs at P14 (Figure 4C; full list of DEGs in
Supplementary Data File 2). Analysis of DEGs at P1 calls one CP related to histone
demethylation. When we analyze DEGs between timepoints and sexes, we observe a
peak in the number of DEGs when comparing P7 vs 14 in both females and males (Figure
4D and Supplementary Data File 3). Within this peak (P7 vs. P14), 1658 DEGs are
identified in males and 3685 DEGs are identified females. These sex differences between

P7-P14 are particularly striking given that these are the developmental time points, where sex hormones are largely dormant. This suggests that these sex differences in astroglial gene expression are part of a developmental program induced by perinatal exposure to hormones involved in sexual differentiation and/or directly induced by sex-linked chromosomes. Remarkably 39% of DEGs at P7 and 79% of DEGs at P14 are downregulated in females. Within these downregulated genes are typical markers of cell activity, the immediate early genes *Fos* and *Arc,* suggesting decreased basal activity in female astroglial cells at these timepoints (Supplementary Data File 3, Tab a). Analysis of DEGs at P7 show that three of the top five CPs are related to synapse development and transmission while the remaining two are related to intracellular signalling pathways. Similar analysis of DEGs at P14 show that the majority of the top five CPs are related to cell cycle. In addition, the top five GO terms from DEGs at P7 are related to myelination and synaptic components and at P14, the top five GO terms are related to cell cycle.

A more complete view of sex differences is captured by differential network dynamics. We used co-expression network analysis (see Online Methods), to infer, separately a co-expression network in females and males throughout development, and identified 10 "female" (F) and 12 "male" (M) co-expression modules, plus one unassigned gene module, grey (Supplementary Data File 4 & Supplementary Data File 5, Tab1). In females, the F3, F6, F7 and F9 modules are enriched in genes upregulated in late vs. early, and the F1, F2, F4 and F10 modules are enriched in genes downregulated in late vs. early (see Supplementary Data File 4,Tab4). The F3, F6 and F9 (upregulated) modules are enriched in astroglial genes. The F2 and F4 modules (downregulated) are enriched in cell cycle terms, and the F10 module (downregulated) is enriched in genes involved in proteasome activity and cell cycle. In the males' network, the M2, M3, M4, M8 and M12 modules are enriched in genes upregulated in late vs. early, and the M1, M7 and M10 modules are enriched in genes downregulated in late vs. early in males (see Supplementary Data File 5, Tab 4). Similar to what was observed in the female network, some of the modules enriched in upregulated genes in the late vs. early phenotype in males (M2, M3, M6 and M8), are enriched in astrocyte genes. It is likely that, in both networks, the astrocyte signature for the modules enriched in upregulated genes reflects

a shift in a mature astroglial phenotype during development. Finally, the M10 module

(downregulated) is enriched in cell cycle terms, and the M1 module (downregulated) is

enriched in genes involved in mRNA processing. The cell cycle signature for some of the

downregulated modules likely reflects a reduction of proliferative potential of the cells over

time, as would be expected over development.

We conducted an eigengene analysis of the network modules to evaluate how

modules changed over development. Overall, we found consistent results across males

and females whereby modules enriched with astroglial genes were increased over

development, modules enriched with genes involved in cell cycle regulation were

downregulated over development, modules involved in translation (ribosomes) were also

downregulated and finally modules enriched in genes related to mitochondria and

mitochondrial function were increased with age, further lending support to the "early" and

"late" astroglial cell phenotypes (Supplementary Data File 5, Tab 4 and Figure 5 A-C).

A closer look at the temporal dynamics highlights some interesting features, with

the female network characterized, mostly, by monotonic, sigmoid like temporal profiles,

and the male network characterized by a larger variety of patterns (subset shown in Figure

5 A-C). In females, the sigmoid profiles are either decreasing with age (F1, F2, F4 and

F10) or increasing with age (F3 and F9). We also find two modules (F6 and F7),

increasing with age, with a possible peak at P7 and P14, respectively. In the sigmoid

profiles, we also identified a 'slow' subgroup (F1, F2 and F9), characterized by a slow

change between P1 and P7 and more activity around P14; and a 'fast' subgroup (F3, F4,

F5, F7 and F10), characterized by a fast change around P7 and lower activity from P14

onward. The temporal dynamic pattern analysis is consistent with the "early" and "late"

hierarchical clustering. Unlike the female network, most of the modules in the male

network show a relatively faster temporal dynamics at P1 to P4-P7. Noticeable are also

the pronounced early peaks in the M5 and M6 modules, the late peaks in the M4, M8 and

M12 modules and a pronounced deep at P7 in the M10 module. Functional analysis

shows that the 'early peaks modules', M5 and M6, are enriched in neurotransmitter

secretion and synaptic transmission terms, respectively; the 'late peaks' modules, M4,

M8 and M12, are enriched in transmembrane transport, signaling receptor

activity/astrocyte, and transmembrane transport terms, respectively. The M10 module is

enriched in mRNA processing, and the M1 in neuron terms.

We tested the modules for enrichment in Estrogen and Testosterone related gene

sets (i.e. ESR1 and ESR2 gene targets and genes/proteins interacting with the

chemical Testosterone (Rouillard et al., 2016) and found significant overlap between the

M2 module in males with the ESR1 targets. The male M2 module expression increases

rapidly, from P1 to P14, where it levels off. This may be consistent with the active process

of masculinization of the brain early in development. No significant overlap is found with

any of the modules in the female set.

Network dynamics highlight differential astrocyte subtype differentiation

We tested the male and female modules for enrichment in gene marker for

astrocyte sub-types (Bayraktar et al., 2020; Nature Neuroscience; Batiuk et al., 2020;

Nature Communications) to investigate sexual dimorphism in astrocyte subtype

differentiation, and describe them following the developmental timeline (see

Supplementary Data File 4, and Supplementary Data File 5, Tab5).

The AST4 (neural stem or progenitor cells) markers overlap with modules F2, F6

and F10 in females, and M1, M8 and M11 in males. The overlap with modules decreasing

with time is expected and in line with the stem cell/progenitor phenotype of AST4, while

the overlap with F6 (females) and M8 (males) is more surprising, given that both modules

first decrease, then increase from P7 (see Figure 5B). Interestingly, vimentin expression

is one of the major markers of the AST4 subtype (Batiuk et al., 2020; Nature

Communications) (in Supplementary Data File 2), therefore our observation of sex

differences in vimentin expression over time (Figure 3B) is consistent with sexual

dimorphism in these particular network dynamics.

The AST5 (intermediate progenitors) markers overlap with F3 and F9 in females

and M1, M2 and M6 in males (see Figure 5C). All modules show increased expression

over time at a faster rate than modules overlapping with the mature astrocytes, and

differently in males than in females. Interestingly, in males, we find overlap with module,

M1, a module decreasing over time, and lack of overlap with a similar module in females

(see Figure 5C). Top scoring terms for this module are mRNA processing, Spliceosome

and Proteasome. Down-regulation of this module could be associated with cell cycle and

the associated dismantling and ciliogenesis of astroglial cilia (Sterpka and Chen, 2018).

Alternatively, the proteasome gene down-regulation could also be consistent with the

GFAP increase over time (Middeldorp et al., 2009). The faster temporal dynamics and

the male specific module point to sexual dimorphisms in astroglial differentiation

dynamics that become explicit at the intermediate progenitor stage; these warrant further

investigation in future studies.

The AST1, AST2 and AST3 (mature astrocytes) markers overlap with modules

showing increasing expression with time both in females (F9 and F3) and in males (M2,

M3 and M12). Notably, the F9 module in females (AST1, AST2) has a slower rise than

F3 (AST2, AST3), suggesting a delay in AST3 with respect to AST1 (see Figure 5A). In

males, M2 is enriched in all the mature cell types, probably capturing common underlying

processes; the M12 (AST1, AST2) module starts increasing late, at P14, M3 has a slower

increase with respect to M2, again suggesting a delay in AST3 differentiation with respect

to AST1 and AST2. The overlap with modules increasing expression in time is consistent

with the mature phenotype of these astroglial classes. Overall, some female modules

show little change between P1 and P4, and a steeper increase from P4 to P7, while male

modules tend to change from P1, suggesting a delay in females. Also, we find two shared

modules in males, across two transitions, M1 in AST4 to AST5, and, M2, in AST5 to

AST1/AST2/AST3; conversely, females do not have any shared modules in the first

transition, AST4 to AST5, and have two shared modules, F3 and F9, both in the second

transition AST5 to AST1/AST2/AST3.

Differential Transcriptional Regulation Dynamics Drives Sexually Dimorphic

Astrocyte Differentiation Programs

The different developmental trajectories and the faster temporal dynamics in

males supports the idea of differential regulation of astrocytic differentiation ?between

males and females, already present at P1. We then investigated the regulation of network

modules to identify determinants of network dynamics and subtype differentiation. We

first identified a number of putative transcription factors regulating the male and female
network modules and found an average of 320 and 340 significant TFs per module,
respectively, for a total of 856 and 858 TFs (see Methods), with 2 male-specific (HELT,
KAT7) and 4 female-specific (CARF, CPEB1, MTF1, PHF1 TFs. Interestingly, HELT and
CPEB1 target AST4 genes markers involved in different processes (HELT: neural crest
differentiation, cell fate specification; CPEB1: cell cycle related; cell projection;
cytoskeleton) (see Table GF4 and GF5 Tab6).
We then used the expression of TFs target genes as a reporter of activity, similarly
to the Viper algorithm (Alvarez et al., 2016), and estimated TF differential activity between
females and males at each time point, and between time points, using the camera
algorithm (Wu and Smyth, 2012) (see Methods).
We first identified the subsets of differentially active TFs (DATFs) between P1 and
P4 in males and females, as putative determinants of early transcriptional dynamics,
resulting, respectively, in a set of 839 and 510 DATFs, respectively. Within this subset,
we identified DATFs at P1 between males and females, as putative TFs driving the
difference in early transcriptional dynamics, resulting, respectively, in a set of 775 and
119 DATFs, suggesting a stronger activity in transcriptional regulation in males vs
females. Next, we focused on the subset of DATFs targeting gene markers of individual
astrocyte sub-types, to focus on cell-type specific differential regulation. We found 551
DATFs in males and 33 in females, with targets co-expressed in the modules M1 (57),
M11 (494) and F2 (24), F10 (9), and enriched marker genes for AST4. At P1, the already
consistent number of differentially active TFs might reflect the differential priming by
testosterone and/or aromatization of estradiol in males of astrocyte progenitor cells, which
would reach peak levels at this time. Interestingly, ESR1 is a DATFs controlling targets in
M11, suggesting that some of the sex differences in the developmental trajectories of
cortical astroglia, may be directly initiated by the perinatal exposure to estradiol critical to
male sexual differentiation.
We then aggregated the TF-target genes and built a regulome (i.e. a regulatory
network represented as a directed graph) for the males (13536 genes and 404928 edges)
and females (13756 genes and 449097edges) using known regulatory relationships, and

identified the putative TFs, potentially upstream of the other TFs/genes, i.e. the source nodes of the graph (see Methods). The source nodes are the nodes in a directed graph, with a non-zero out-degree and a zero in-degree. We found ESR1, ZFP3, MZF1, PRRX1, SOX7, MSX2, ZGPAT, TCF24, and MYRFL in the male network and BARX2, FOXR2, and MYRFL in the female network. Interestingly, while ESR1 is present in both, it is a source node only in the male network, supporting a central role of ESR1 in AST differentiation in males and of network rewiring early in masculinization.

Early masculinization of the brain occurs largely as a result of a perinatal surge in testosterone that is aromatised into estradiol in the brain of males, therefore we further examined P1 as a critical time point in setting up differential developmental patterns in the astroglia of males and females. Extensive work from the McCarthy laboratory has demonstrated that prostaglandins, and specifically, glial release of cyclooxengase 2 (COX-2) is a key regulator of estrogen-mediated architectural sex differences in the hypothalamus and the amygdala[44,45]. The gene for COX2, *Ptgs2*, is a sex-specific DEG in our data set, in the overall females versus males analysis, in the early female versus male phenotype, and at P1 female versus male comparison (Supplementary Data File 1 & Supplementary Data File 2) Furthermore, Cox2 is listed as an ESR1 target in our dataset and it is well- known to be downstream of ESR1 signalling (Stossi et al., 2004). To validate COX2 expression over development, we counted the number of cortical COX2+ cells in astroglia (co-localised with eGFP) and saw a similar pattern of expression to that observed in our TRAPseq, with a significant sex difference in astroglial Cox2 protein expression at P1 ($F_{1,41}$=12.517 p=0.017, $\mathrm{H}\eta$=0.715)(Figure 6A). COX2-mediated masculinization of the hypothalamus occurs in response to estradiol, which is aromatised from the surge in testosterone in males that occurs perinatally in rodents (Figure 6B). This perinatal surge in testosterone has been linked to sexual differentiation of the hypothalamus, amygdala, and mediates sexually dimorphic juvenile and adult behaviours[46-48]. To validate our findings and to test if astroglial COX2 protein levels are similarly regulated by estrogen in the cortex, we compared vehicle injected males and females with females that were masculinized by injecting estradiol benzoate at P0 and then sacrificed all groups 36 hours later. The overall one-way ANOVA showed a trend for

significance ($F_{3,20}$=2.781, p=0.068, Hη=0.294), however because our previous
experiment showed a difference between males and females, we conducted a planned
comparison of these groups. Females at P1 show fewer COX2+ cells in the frontal cortex
compared to males ($F_{1,9}$=7.827, p=0.021, Hη=0.465) and females exposed to estrogen
($F_{1,9}$=5.759, p=0.04, Hη=0.390) (Figure 6C). This suggests that estrogen exposure at
P0 is sufficient to masculinize COX2 expression the cortex.
479
(Sex differences in) Astrocyte differentiation and relevance to disorders
We investigated the relevance of the male and female co-expression network to human
disease, by testing the network modules for overlap with disease ontologies (DOs) (see
Methods). We focused on the subset of the modules in overlap with astrocyte subtype
markers, and report only results for brain-specific disorders (see Supplementary Data File
4 & Supplementary Data File 5, Tab7). In females, top scoring DO terms are intellectual
disability and mental deficiency (Module F2); intellectual disability, bipolar disorder,
epilepsy, schizophrenia, cognitive delay, autistic disorder, Alzheimer's disease (Module
F3); Unverricht-Lundborg syndrome, lysosomal storage diseases, amyloidosis,
progressive myoclonic epilepsies (Module F6); epilepsy seizures, Alzheimer's disease,
schizophrenia, intellectual disability, major depressive disorder, neurodegenerative
disorders, autistic disorder, Parkinson's disease (Module F9). In males, top scoring DO
terms are intellectual disability (Module M1); mitochondrial diseases, Alzheimer's
disease, neurodegenerative disorders, epileptic encephalopathy, seizures, bipolar
disorder, epilepsy, schizophrenia, autistic disorder, Parkinson's disease (Module M2);
peripheral neuropathy (Module M3); mitochondrial diseases, Leigh disease (Module M6);
intellectual disability (Module M8); microcephaly, ciliopathies, intellectual disability
(Module M11); epilepsy, bipolar disorder, mental depression, unipolar depression,
depressive disorder, Alzheimer's disease, schizophrenia, Intellectual disability, autistic
disorder (Module M12). Notably, many of the diseases are in overlap between males and
females, and there does not appear to be disease specificity per se as a wide variety of
neurological, psychiatric, and developmental disorders are called. Importantly, however,
each of these diseases have been linked to astroglial dysfunction either primarily or in a

reactive capacity and therefore the wide variety of diseases called likely speaks to the

general involvement of astroglia in all neural processes (Molofsky et al., 2012). Modules

associated to differentiated astrocyte subtypes, i.e. increasing with age, are enriched in

neurodegenerative disorders terms. Interestingly, these same modules also show

enrichment in autism-associated genes, and only modules F9 and M2 show enrichment

in Parkinson's disease genes, unlike Alzheimer's and schizophrenia. Perhaps reflecting

regional involvement in disease pathology, with Parkinson's origins to mid-brain-striatal

functions rather than cortical origins. Module F6, associated with astrocyte progenitor

cells, is enriched in epilepsy related genes including a number of GABA transporters and

GABA receptor subunits (*gabra3, gabrg2, gabrb2, slc6a1*), unlike its M8 counterpart.

Interestingly, both astroglial cells and GABA subunits are central to epilepsy pathology

(Coltour and Steinhäuser, 2015; Fritschy, 2009), and sex differences in gaba subunits

have been linked to sex differences in epilepsy (Li et al., 2007) however, whether

astroglial expression of these subunits is related to developmental sex differences in

epilepsy susceptibility is unknown. Finally, the "early peak" module M6 associated with

intermediate progenitor cells, is enriched in mitochondrial diseases terms. This male-

specific expression is consistent with mitochondrial diseases having a higher incidence

in males, thought to be due to uniparental or maternal inheritance (Beekman et al., 2014).

Discussion

In the current study, we generated a translatome database of cortical astroglial

cells across development in both male and female mice. We show two clear astroglial

phenotypes across development, an early (before P14) and late (after P14) phenotype.

These early and late gene expression profiles reflect astroglia associated with early

developmental functions related to gliogenesis and synaptogenesis and astroglia

associated with functions involved in maturation, synapse elimination and maintaining a

stable, homeostatic environment in adulthood (Figure 7D).

Surprisingly, cortical astroglial cells showed very little sexual dimorphism in either

morphology or gene expression particularly in adulthood when very few DEGs were not

sex-chromosome associated genes. Importantly, to assess basal differences between

males and females in a conservative way, we chose to examine females during the

metestrus phase, when ovarian hormones are at their lowest and therefore more similar

to levels observed in males. As such, it is likely that greater sex differences in the

astroglial translatome would be observed during other estrous cycle phases, such as in

proestrus, when estrogen peaks. Indeed, we have previously demonstrated changes in

cortical astroglial expression of GFAP and fibroblast growth factor 2 (FGF2), an astroglial

mitogen and growth factor across the estrous cycle[49], and a number of studies have

shown functional, morphological and protein expression changes in astroglia in response

to sex hormones (Verkhratsky and Nedergaard, 2018; Azcoitia et al., 2010) Together

these results would suggest that astroglial cells show phenotypic changes in response to

ovarian hormones (or activational effects of hormones), but at least in the neocortex,

astroglia do not show basal sexual dimorphism (organizational effects of hormones).

Interestingly, the sex differences in DEGs observed at P7,and P14, were not

related to sex-specific canonical pathways or functions per se, but rather, were related to

general developmental phenomena such as cell cycle, and neuronal, dendritic and

synaptic development, suggesting that astroglia modulate neuronal networks in a sexually

dimorphic way. This could, in turn, potentially lead to the formation of sexually dimorphic

neuronal networks, and presumably, sexually dimorphic behavioural expression later in

life. Given the critical role of the neocortex in psychiatric disease and complex human

behaviours[50], one might imagine that changes in neuronal network development may be
induced by estrogen-modulated changes in astroglia. In addition, these may be
particularly susceptible to developmental perturbations that modulate hormone
expression levels including exposure to stress hormones and environmental endocrine
disruptors. Moreover, because hormone levels vary between individuals under normal
conditions, this allows for a potential range of sexual differentiation of cortical networks,
which would mirror behavioural sex differences observed across species, that are
typically expressed on a continuum.

In contrast to the sex differences in gene expression seen in P7-P35, we see few
DEGs between astroglia in male and female mice in the early postnatal period and in
adulthood. This, in addition to the sex differences in CP of early versus late astroglial
phenotype and the apparent early peak of some networks in male mice, suggests that
cortical astroglial cells employ different developmental strategies or timing in
developmental trajectories to reach a similar adult phenotype, with males accelerating
faster than females in early development. The idea that distinct developmental trajectories
between the sexes might lead to similar functional results is similar to the concept of
compensation put forward by Woolley[51], which was based on findings of estradiol
potentiating glutamatergic synapses, albeit on a much more general level. Earlier
DeVries[52] postulated that sex differences in gene expression patterns over development
may actually prevent sex differences in specific regions or ensure that neuronal systems
develop and are functional in the presence of cues such as sex hormones that are crucial
for stimulating sex differences in target regions. It is possible that the sex differences in
gene expression profiles between astroglia ensure that these systems develop "equally"
in both sexes to achieve similar functionality in adulthood in the presence of sex
chromosomes and hormones that are necessary to differentiate regions critical to sexually
dimorphic behaviours, such as the hypothalamus.

The question remains as to what induces sex differences in cortical astroglia gene
expression patterns over development. Three potential scenarios emerge: 1) sex
chromosome-linked genes directly induce changes in gene expression patterns in
astroglia. 2) The perinatal testosterone surge initiates cascades of gene expression

patterns in astroglia that modulate cell maturation and network development. 3) Astroglial
cells have a similar start and end point in our dataset and it is possible that sex differences
in astroglia gene expression represents an active process to compensate for sex
differences in developing neuronal imputs to the cortex that arise from dramatic changes
in neuronal number, arbours, and synapses on neurons that project from the
hypothalamus to the neocortex Importantly, these scenarios need not be mutually
exclusive and may be active in parallel systems.

Scenario 1. Kdm5d appears as one of the few DEGs between females and males
at P1 in our dataset, female expression levels are lower than males. *Kdm5d* is a Y-linked
gene that encodes for demethylation enzymes and is required and sufficient to induce
sexual differentiation of gene expression[53]. Indeed, changes in gene methylation may
have profound effects that persist throughout time and cascades of gene expression may
or may not be initiated, altering the responsiveness of the cell, and thus the network as a
whole. Moreover, even in the absence of the testes determining *Sry* gene, the Y-
chromosome has been shown to exert differential effects on developing peripheral and
nervous tissue (reviewed in[54]), lending support to the idea that sex chromosome linked
genes induce later sex differences in astroglia across development.

Scenario 2. Similarly, our results suggest that the perinatal testosterone surge, and
consequentially, estrogen, is involved in the induction of developmental sex differences
in astroglia. Our weighted network analysis shows enrichment of genes in several
modules associated with responses to sex hormones. We show sex differences in the
prostaglandin system (including Cox2) and that neonatal estrogen administration induces
changes in cortical COX2 expression. Similar changes in astroglia have been
demonstrated in the hypothalamus and are sufficient to produce observable sex
differences in neuronal networks and associated behaviour[13]. Thus, the endogenous
expression of estrogen early in development may facilitate a cascade of cellular
maturation that in-turn will alter subsequent genetic programs. As such, the neuronal
network will facilitate different programs as a result of the staggered pattern of maturation
and communication.

Finally, *scenario 3* suggests that sex differences in the hypothalamus are
necessary for a functional behavioural outcome that is overt and essential for sexual
reproduction, first suggested by de Vries[52]. However, there are few overt measures that
can be linked to cortical functioning that exhibit similar sexual dimorphisms, although sex
differences in risk and resilience for psychiatric disease associated with cortical
functioning are well-supported. For example, schizophrenia and ASD are more prominent
in the male population[8] and both show changes in developmental pathophysiology that
occur both during the prenatal and postnatal periods[55,56]. There may be a relationship
between astroglial neurogenic potential and the terminal phases of neuronal migration
and construction and maturation of functional cortical networks that buffers females
against developmental anomalies which, if left unchecked, may lead to schizophrenia or
ASD morbidity.
In-line with this hypothesis, there are a number of models where estrogen
treatment improves recovery or where females are protected in developmental injury
models. Chronic postnatal hypoxia in rodents is used as a model of premature birth and
induces loss of brain volume, particularly in the cortex and hippocampus, and leads to
motor and cognitive impairments later in life[57,58]. The loss of cortical volume in the hypoxic
model is less pronounced in females[57] and interestingly, estradiol treatment at the time of
the injury improves white matter damage recovery[59]. It is also known that chronic
postnatal hypoxia increases astroglial stem cell capacity[25,60]. This process may be
enhanced in females allowing augmented recovery from the aversive effects of hypoxia,
but may also speak to the nature of enhanced early postnatal plasticity in females
compared to males. It is possible that females experience an enhanced or shifted
neurogenic period early postnatally, when we have observed an increase in vimentin+
astroglia, that may translate to increased neurogenic potential of astroglial cells in the
cortex. Similarly, while overt network or cellular sex differences may not modulate sex
differences in risk for psychiatric disease, sex differences in developmental programs may
leave one sex more vulnerable to perturbations during that time. Future studies will be
needed to determine the functional significance of sex differences in developmental
programs. Nevertheless, our data show a clear developmental pattern in cortical astroglia

that differs between males and females, while no apparent basal sex differences in

astroglia exist.

Online Methods

Animals

54 male and female C57/Bl6-AldHl1-L10-GFP transgenic mice generated from our

breeding colony maintained at the Carleton University animal facility within the University

of Ottawa. Mice were genotyped as in (Simard et al., 2018) and sex was determined by

the presence or absence of tissues in the urogenital tract and confirmed by genotyping

for the *Zfy* gene. All animals were group-housed until sacrifice in standard (27cm x 21cm

x 14cm), fully transparent polypropylene cages with chew block, bedding, house and *ad*

libitum access to standard lab chow (2014 Teklad Global 14% protein®) and water.

Animals were raised in the standard environment with no outside manipulation except for

standard care and to monitor estrous cycle stage (when applicable). The mice were

maintained on a 12-hour light/dark cycle in a temperature controlled (21 degrees) facility.

All animal use procedures have been approved by the Carleton University Committee for

Animal Care, according to the guidelines set by the Canadian Council for the Use and

Care of Animals in Research. Mice were overdosed and perfused for

immunohistochemistry or rapidly decapitated for TRAPseq processing each on postnatal

day 1* (P1) (36 hours postnatal), P4, P7, P14, P35 and adulthood (defined as 7-9 weeks

of age) at the same time of day. Developmental time points and subject numbers are

summarized in Supplementary Figure 1.

Organizational hormones study- 5 pregnant dams were ordered from Charles

River and arrived at the Carleton University animal facility at the University of Ottawa at

gestational day 15/16 and given 3 days to acclimatize to the environment. Dams were

monitored intermittently overnight starting on gestational day 19 using red light. 31 male

and female pups were randomly assigned to control or treatment groups approximately 1

hour following birth and sacrificed 35 hours later. The majority of groups in all experiments

were formed with subjects from 3-5 separate litters to control for litter effects.

Estrous Cycle Monitoring

All adult group female animals were monitored daily after P35 to identify stage of

estrous cycle using a saline lubricated swab inserted into the opening of the vagina to

collect cells from the vaginal wall. Samples were smeared on a glass microscope slide

and examined under a 10x objective using a light microscope (VistaVision™). To
understand basal organizational sex differences in astroglia, we used female mice in the
phase of the estrous cycle with the lowest circulating ovarian hormones that has the least
effect on cortical astroglial morphology and protein expression: metestrus[61]. After the
stage of estrous cycle (Metestrus, Diestrus, Proestrus and Estrus) had been established
for at least two full cycles, females were sacrificed during the light cycle of the metestrus
stage.

Organizational hormones-injections

Estradiol benzoate (Chem Cruze, sc-205314) was dissolved in peanut oil
(0.8g/mL) as per Hisasue et al., 2010 to allow for an injection of 20µg in a 25µL injection
volume. Letrozole (Sigma-Aldrich, LG545-10MG) was dissolved in 0.3% hydroxypropyl
cellulose PBS (Aldrich, 435007-5G) as per[62], to allow for an injection of 250mg/kg in a
25µL injection volume, sufficient to block all testosterone conversion based on average
pup weight of 5g. Estradiol benzoate and letrozole solutions were prepared using glass
materials and stored in a glass amber bottle to prevent absorption and disruption from
plastics or ultraviolet exposure.

Animal Sacrifice

Rapid Decapitation for immunohistochemistry and qPCR- Animals in the P1

immunohistochemical group were rapidly decapitated and brains immediately placed in a

4% paraformaldehyde (Fisher Scientific) (PFA) solution at 4 ℃ for 24h after which the

brains were switched to 30% sucrose at 4 ℃ for 24h. Following this period, brains were

flash frozen until slicing. For the organizational hormone experiment, pups underwent

rapid decapitation 36h following treatment. Brains were harvested and split along the

central fissure. One hemisphere was flash frozen at -80°C for analysis in qPCR and the

other hemisphere was fixed as above.

Cardiac Perfusion for Immunohistochemical Analysis- All mice were sacrificed for

tissue collection at the same time of day during the light cycle. All mice in the

immunohistochemical group underwent cardiac perfusion aside from those in group P1

(see above). Animals in P4, P7, P14, P35 and adult groups were given an overdose of

44mg/kg sodium pentobarbital (CDMV Canada), followed by intra-cardiac perfusion upon

all spinal reflex cessation. Animals from the P4 and P7 groups underwent manual cardiac

perfusion with 1mL of saline solution before receiving 1mL of 4% PFA via syringe. Mice

in the P14, P35, and adult groups blood were flushed using 10mL of saline through the

left atrium followed by 20mL of 4% PFA to fix the tissue. Brains were extracted and placed

into vials containing 4% PFA and stored at 4 ℃. Following a 24h period, brains were

transferred to a 30% sucrose in PBS (Fisher Scientific) solution and placed at 4 ℃*.*

Tissue sectioning

Following sucrose treatment, all brains were then flash frozen at -80 ℃ until

sectioning on a Leica (Leica™ CM1900) cryostat (30µm thick). 15 sets of sagittal sister

sections were adhered to electrostatic slides (Fisher Scientific™) in rotating order, each

slide therefore contained a full representation of the brain for stereological analysis as per

previous studies[60,63].

Immunohistochemistry

All immunohistochemical processes took place at room temperature (~21 ℃) as

per our previous studies[25,60,64]. One representative slide was taken from every subject

and all subjects were processed simultaneously. Brain tissue from all groups were

prepared for immunohistochemistry using a 10% horse serum (Gibco™) PBS-T (0.3%

TritonX (Fisher Scientific)) pre-block solution for 1 hour before being incubated with the

respective primary antibody solution (see Supplementary Figure 1). Primary antibodies

were diluted in 10% horse serum PBS-T (0.3% TritonX). Following approximately twenty-

four hours of primary antibody incubation, slices were washed in a 1x PBS solution 3

times to remove unbound antibodies before being incubated with the species-appropriate

fluorescein conjugated secondary antibody for visualization. Secondary antibodies were

diluted in 10% horse serum PBS-T (0.3% TritonX) and incubated for 2 hours before slides

were submerged in a 1x PBS wash 3 times to remove unbound antibodies. Slides were

then coated with a nuclear stain, DAPI with hard setting mounting medium (VectorLabs),

to fix glass cover slips and allowed to set before analysis. See Supplementary Figure 1

for antibodies and dilutions used.

Stereological Analysis

Unbiased estimates of cell numbers were obtained through use of Zeiss

AxioImager M2 with ApoTome motorized fluorescent microscope (Carl Zeiss, Thornwood,

NY, USA) in conjunction with a motorized stage and a computer running on Windows 7

using the program StereoInvestigator™ (MicroBrightfield, Colchester, VT, USA). Serial

sagittal sections of the right hemisphere obtained through cryosectioning at 30µm on 15

sister sections were used for stereological analysis. Contours encompassing the whole

right hemisphere neocortex on each section were drawn as boundaries in

StereoInvestigator™. Cells were counted for expression of individual and/or co-

expressed proteins using the optical fractionator probe at 40x. Sampling grids were

optimized for cortical contours to include at minimum 3 sampling sites per contour to allow

for a systematic and unbiased method to estimate cell density and cell quantification for

right hemisphere of the cortex regardless of cell shape, size, orientation, spatial

distribution, or post-mortem brain shrinkage[65]. Sampling boxes are randomly placed by

StereoInvestigator™ within the sampling frame measuring 150µm x 150µm x 30µm with

3 of 6 exclusion borders. Total number of cells per count are reported via

StereoInvestigator™ output. For analysis of astroglial morphology, StereoInvestigator™

(MBF, Vermont™) software on a Zeiss Observer with Apotome (Zeiss™) was employed.

We identified and distinguished between radial and non-radial astroglial cells, counting
these using unbiased sampling via the StereoInvestigator (MBF, Vermont ™) at 40x.
Representative confocal mosaic images used for photomicrographs were taken using
ZEN software (Zeiss™) with the Airyscan 800 microscope (Zeiss™).

TRAPseq

Translating ribosome affinity purification (TRAP)-sequencing was performed as
per[66]. Briefly, eGFP-positive animals were rapidly decapitated and the brains removed
and placed into a 1000x activated cyclohexamide (CHX) (100µg/ml, Sigma, dissolved in
methanol, American Bioanalytical) dissection buffer at which point the cortex, removing
any remaining white matter was dissected and flash-frozen in CHX dissection buffer at -
80°C. When appropriate, brains were thawed on ice in homogenization buffer containing
protease inhibitors, 1000x activated CHX, and RNAse inhibitors. Tissue samples were
mechanically homogenized and then incubated on a rotisserie with anti-GFP (HtzGFP_04
(clone19F7) and HtzGFP_02 (clone 19C8): Memorial Sloan-Kettering Monoclonal
Antibody Facility) coated biotinylated magnetic beads (Streptavidin MyOne T1
Dynabeads; Invitrogen # 65601) coated with Protein L (Fisher # PI-29997) at 4 °C
overnight. Following incubation with magnetic beads, the samples were collected on a
magnet (DynaMag-2; Invitrogen #123-21D) and the unbound fragment was collected and
frozen at -80 °C. Bound mRNA fragments washed in polysome buffer underwent RNA
extraction with DNAse treatment using Absolutely RNA Nanoprep Kit (Stratagene
#400753).

RNA samples were sent to the Yale K.E.C.K facility for sample QC as per Simard
et al. 2018: RNA quality was determined using a nanodrop and RNA integrity was
determined by running an Agilent Bioanalyzer gel (RINs >8). mRNA was purified from
approximately 500ng of total RNA with oligo-dT beads and sheared by incubation at 94C.
Following first-strand synthesis with random primers, second strand synthesis was
performed with dUTP for generating strand-specific sequencing libraries. The cDNA
library was then end-repaired and A-tailed, adapters were ligated and second-strand
digestion was performed by Uricil-DNA-Glycosylase. Indexed libraries that met
appropriate cut-offs for both were quantified by qRT-PCR using a commercially available

kit: Kapa Library Quant Kit (Illumina) (KAPA Biosystems # KK4854-07960298001) and

insert size distribution was determined with the LabChip GX or Agilent Bioanalyzer. Only

samples with a yield of ≥0.5 ng/ul were used for sequencing.

Sample concentrations were normalized to 10 nM and loaded onto Illumina Rapid

or High-output flow cells at a concentration that yields 130-250 million passing filter

clusters per lane. Samples were sequenced using 75bp paired-end sequencing on an

Illumina HiSeq 2500 according to Illumina protocols. The 6bp index was read during an

additional sequencing read that automatically follows the completion of read 1. Data

generated during sequencing runs were simultaneously transferred to the YCGA high-

performance computing cluster. A positive control (prepared bacteriophage Phi X library)

provided by Illumina is spiked into every lane at a concentration of 0.3% to monitor

sequencing quality in real time.

Quantitative Real-Time PCR

To validate the enrichment of astroglial cell genes in our bound,

immunoprecipitated fragment, the astroglial specific genes GFAP and Glul expression

levels were compared between input and TRAP-immunoprecipitated fragments. In

addition, we evaluated RbFox3 and Pvalb gene expression assays to validate a

concurrent decrease in neuronal gene expression. 20ng of RNA (concentration

determined using NanoDrop (Thermofisher Scientific)) was used to produce cDNA using

SuperScript IV First Strand Sythesis System (ThermoFisher Scientific). Best-coverage

TAQMAN assays (Life Technologies) were run as per manufacturer's instructions and

analyzed using the Applied Biosystems 7500 Real-Time PCR system and software.

To validate the TRAP-seq results we used RNA from all bound (*n*=55),

immunoprecipitated fragments to produce cDNA using SuperScript III First-Strand

Synthesis SuperMix (Thermo Fisher Scientific; catalog #11752050). We then performed

a preamplification reaction (14 cycles) using TaqMan PreAmp Master Mix (Thermo Fisher

Scientific; catalog #4384267), with a pooled-assay mix generated according to user

guidelines. Finally, best-coverage TaqMan Gene Expression Assays (see Supp. Figure

3) were run as per the manufacturer's instructions using the Bio Rad CFX Connect Real

Time system and software.

Statistical Analysis

All immunohistochemical and qPCR data were analyzed for using a 2 (male vs.

female) x 6 (P1 vs. P4 vs. P7 vs. P14 vs. P35 vs. Adult) between-subject factorial

analyses of variance (ANOVA) design using IBM SPSS Statistics (Version 20.0). Because

the treatment differed in males and females, for the organizational hormones study we

employed a one-way ANOVA to determine overall significance. Post-hoc analysis using

Bonferroni correction of pairwise comparisons for non-orthogonal comparisons were

conducted when p<0.05. We re-ran all of the statistics including litter composition (male

vs. female) as a covariate and found no significant effects of litter on any of the variables

measured (data not shown).

RNAseq data analysis. Sequencing reads were mapped to the mouse genome

(GRCm38.p5) and the Ensembl (releas-84) transcriptome annotation, using HISAT2

(Pertea et al., 2016). The mapping rates ranged from 74% to 95%. This resulted in a

number of mapped pairs from ~19.5 million up to 40 million (see Supplementary Table 6

for more details). The resulting SAM files were converted to BAM formant using

SAMtools[67]. The gene expression levels as counts were estimated using featureCounts[68]

with the following options: -t exon; -g gene_id; -F "GTF"; -B; -s 0; -p; -a. Counts from

technical replicates were added. Raw read counts were then filtered requiring more than

1 CPM (counts per million) in at least 25% of the samples and 16208 survived the filter

and were considered for further analysis. We next used the sva Bioconductor package

(Leek et al., 2012) to identify and correct for latent variables. Hierarchical clustering (using

the genes with a sd in expression level above the 95%) pre- and post sva correction show

the sample-to-sample relationship and relative clustering (see Supplemental Figure 2).

FastQC v0.10.1 and RNA-SeQC (v1.1.8) were used for QC (see Supplemental Table

QC). The edgeR function rpkm was used to estimate gene expression levels as RPKM.

Differential expression analysis. Differential expressed genes were then inferred

using the edgeR pipeline[69], using the trended dispersion to estimate the biological

variance and the GLM capability of the package. Nominal p-values from differential

expression analysis were FDR corrected, and an FDR cut-off of 0.05 was used for all the
tests.

Functional enrichment analysis. ConsensusPathDB[70] was used to test
differentially expressed genes for overrepresentation in Gene Ontologies and Canonical
Pathways.

Weighted gene co-expression network analysis. We used Weighted Gene Co-
expression Network Analysis (WGCNA)[71] for co-expression network analysis using gene
expression estimates (as log2(RPKM+1)), separately, from the 25 male samples and the
26 female samples, after cqn and sva correction. We estimated male and female co-
expression networks and modules using the function blockwiseModule, using the bicorr
as correlation estimate, a "signed" network type and a minModuleSize=50. The power
cut-off was set to 16 for both sets. The analysis produced a male network of 13 modules
and a female network of 15 modules, including the grey module of unassigned genes
(See WGCNA Summary Data).

Transcription Factor Analysis. We used the ChEA3 online suite (Keenan et al.,
2019) to test each network module for enrichment in Transcription Factors (TFs),
retaining only significant calls at FDR<0.05. We considered all the available databases
within ChEA3. This resulted in duplicated significant TFs, which were combined,
considering the union of all the reported target genes. We estimate the activity for each
TF using the camera gene set enrichment analysis algorithm (Wu and Smyth, 2012).
considering the P4 vs P1 difference. FDR corrected for multiple comparisons and used
FDR<0.05 as cut-off. Any TF estimated as non-significantly active was not retained for
further analysis. We then built a directed network from the remaining TFs, using the
known TF-target interaction and the WGCNA adjacency as edge weights.

871   References

872   1   Molyneaux, B. J., Arlotta, P., Menezes, J. R. L. & Macklis, J. D. Neuronal subtype
873   specification in the cerebral cortex. *Nature Reviews Neuroscience* **8**, 427-437,
874   doi:10.1038/nrn2151 (2007).
875   2   Barton, R. A. & Harvey, P. H. Mosaic evolution of brain structure in mammals. *Nature*
876   **405**, 1055-1058, doi:10.1038/35016580 (2000).
877   3   Berry, M. & Rogers, A. W. The migration of neuroblasts in the developing cerebral cortex.
878   *Journal of anatomy* **99**, 691-709 (1965).
879   4   Rakic, P. Mode of cell migration to the superficial layers of fetal monkey neocortex. *The*
880   *Journal of Comparative Neurology* **145**, 61-83, doi:10.1002/cne.901450105 (1972).
881   5   Strakowski, S. M. *et al.* The functional neuroanatomy of bipolar disorder: a consensus
882   model. *Bipolar Disorders* **14**, 313-325, doi:10.1111/j.1399-5618.2012.01022.x (2012).
883   6   Lin, C.-H. & Lane, H.-Y. Early Identification and Intervention of Schizophrenia: Insight
884   From Hypotheses of Glutamate Dysfunction and Oxidative Stress. *Frontiers in psychiatry*
885   **10**, 93, doi:10.3389/fpsyt.2019.00093 (2019).
886   7   Bayés, À. *et al.* Characterization of the proteome, diseases and evolution of the human
887   postsynaptic density. *Nature Neuroscience* **14**, 19-21, doi:10.1038/nn.2719 (2011).
888   8   Bao, A.-M. & Swaab, D. F. Sexual differentiation of the human brain: Relation to gender
889   identity, sexual orientation and neuropsychiatric disorders. *Frontiers in*
890   *neuroendocrinology* **32**, 214-226, doi:10.1016/j.yfrne.2011.02.007 (2011).
891   9   Donner, N. C. & Lowry, C. A. Sex differences in anxiety and emotional behavior. *Pflugers*
892   *Archiv : European journal of physiology* **465**, 601-626, doi:10.1007/s00424-013-1271-7
893   (2013).
894   10   Jaric, I., Rocks, D., Cham, H., Herchek, A. & Kundakovic, M. Sex and Estrous Cycle
895   Effects on Anxiety- and Depression-Related Phenotypes in a Two-Hit Developmental
896   Stress Model. *Frontiers in molecular neuroscience* **12**, 74, doi:10.3389/fnmol.2019.00074
897   (2019).
898   11   Kokras, N. *et al.* Forced swim test: What about females? *Neuropharmacology* **99**, 408-421,
899   doi:10.1016/J.NEUROPHARM.2015.03.016 (2015).
900   12   McEwen, B. S., Lieberburg, I., Chaptal, C. & Krey, L. C. Aromatization: Important for
901   sexual differentiation of the neonatal rat brain. *Hormones and Behavior* **9**, 249-263,
902   doi:10.1016/0018-506X(77)90060-5 (1977).
903   13   Wright, C. L., Schwarz, J. S., Dean, S. L. & McCarthy, M. M. Cellular mechanisms of
904   estradiol-mediated sexual differentiation of the brain. *Trends in Endocrinology &*
905   *Metabolism* **21**, 553-561, doi:10.1016/j.tem.2010.05.004 (2010).
906   14   GORSKI, R. A. & WAGNER, J. W. Gonadal Activity and Sexual Differentiation of the
907   Hypothalamus. *Endocrinology* **76**, 226-239, doi:10.1210/endo-76-2-226 (1965).
908   15   Johnson, R. T., Breedlove, S. M. & Jordan, C. L. Androgen receptors mediate
909   masculinization of astrocytes in the rat posterodorsal medial amygdala during puberty. *The*
910   *Journal of comparative neurology* **521**, 2298-2309, doi:10.1002/cne.23286 (2013).
911   16   Woolley, C. S., Gould, E., Frankfurt, M. & McEwen, B. S. Naturally occurring fluctuation
912   in dendritic spine density on adult hippocampal pyramidal neurons. *Journal of*
913   *Neuroscience* **10** (1990).

17   Bowers, J. M., Waddell, J. & Mccarthy, M. M. A developmental sex difference in
hippocampal neurogenesis is mediated by endogenous oestradiol. *Biology of Sex*
Differences **1**, 8, doi:10.1186/2042-6410-1-8 (2010).
18   Frazier, T. W., Georgiades, S., Bishop, S. L. & Hardan, A. Y. Behavioral and Cognitive
Characteristics of Females and Males With Autism in the Simons Simplex Collection.
Journal of the American Academy of Child & Adolescent Psychiatry **53**, 329-340.e323,
doi:10.1016/J.JAAC.2013.12.004 (2014).
19   Parellada, M., Gomez-Vallejo, S., Burdeus, M. & Arango, C. Developmental Differences
Between Schizophrenia and Bipolar Disorder. *Schizophrenia Bulletin* **43**, 1176-1189,
doi:10.1093/schbul/sbx126 (2017).
20   Sloan, S. A. & Barres, B. A. Mechanisms of astrocyte development and their contributions
to neurodevelopmental disorders. *Current opinion in neurobiology* **27**, 75-81,
doi:10.1016/j.conb.2014.03.005 (2014).
21   Clarke, L. E. & Barres, B. A. Emerging roles of astrocytes in neural circuit development.
Nature Reviews Neuroscience **14**, 311-321, doi:10.1038/nrn3484 (2013).
22   Loo, L. *et al.* Single-cell transcriptomic analysis of mouse neocortical development. *Nature*
communications **10**, 134-134, doi:10.1038/s41467-018-08079-9 (2019).
23   Bayraktar, O. A. *et al.* Single-cell in situ transcriptomic map of astrocyte cortical layer
diversity. *bioRxiv*, 432104, doi:10.1101/432104 (2018).
24   Anthony, T. E., Klein, C., Fishell, G. & Heintz, N. Radial Glia Serve as Neuronal
Progenitors in All Regions of the Central Nervous System. *Neuron* **41**, 881-890,
doi:10.1016/S0896-6273(04)00140-0 (2004).
25   Bi, B. *et al.* Cortical glial fibrillary acidic protein-positive cells generate neurons after
perinatal hypoxic injury. *The Journal of neuroscience : the official journal of the Society*
for Neuroscience **31**, 9205-9221, doi:10.1523/JNEUROSCI.0518-11.2011 (2011).
26   Ganat, Y. M. *et al.* Early Postnatal Astroglial Cells Produce Multilineage Precursors and
Neural Stem Cells In Vivo. *Journal of Neuroscience* **26**, 8609-8621,
doi:10.1523/JNEUROSCI.2532-06.2006 (2006).
27   Salmaso, N., Tomasi, S. & Vaccarino, F. M. Neurogenesis and maturation in neonatal brain
injury. *Clinics in perinatology* **41**, 229-239, doi:10.1016/j.clp.2013.10.007 (2014).
28   Sofroniew, M. V. & Vinters, H. V. Astrocytes: biology and pathology. *Acta*
neuropathologica **119**, 7-35, doi:10.1007/s00401-009-0619-8 (2010).
29   Chever, O., Pannasch, U., Ezan, P. & Rouach, N. Astroglial connexin 43 sustains
glutamatergic synaptic efficacy. *Philosophical Transactions of the Royal Society of*
London B: Biological Sciences **369** (2014).
30   Diniz, L. P. *et al.* Astrocyte transforming growth factor beta 1 promotes inhibitory synapse
formation via CaM kinase II signaling. *Glia* **62**, 1917-1931, doi:10.1002/glia.22713
(2014).
31   Di Garbo, A., Barbi, M., Chillemi, S., Alloisio, S. & Nobile, M. Calcium signalling in
astrocytes and modulation of neural activity. *Bio Systems* **89**, 74-83,
doi:10.1016/j.biosystems.2006.05.013 (2007).
32   Cahoy, J. D. *et al.* A Transcriptome Database for Astrocytes, Neurons, and
Oligodendrocytes: A New Resource for Understanding Brain Development and Function.
The Journal of Neuroscience **28**, 264 LP-278, doi:10.1523/JNEUROSCI.4178-07.2008
(2008).

33   Milner, T. A. *et al.* Ultrastructural evidence that hippocampal alpha estrogen receptors are
located at extranuclear sites. *Journal of Comparative Neurology* **429**, 355-371,
doi:10.1002/1096-9861(20010115)429:3<355::AID-CNE1>3.0.CO;2-#
10.1002/1096-9861(20010115)429:33.0.CO;2-# (2001).
34   Platania, P. *et al.* Differential Expression of Estrogen Receptors Alpha and Beta in the
Spinal Cord during Postnatal Development: Localization in Glial Cells.
Neuroendocrinology **77**, 334-340, doi:10.1159/000070899 (2003).
35   Lee, E. *et al.* GPR30 regulates glutamate transporter GLT-1 expression in rat primary
astrocytes. *The Journal of biological chemistry* **287**, 26817-26828,
doi:10.1074/jbc.M112.341867 (2012).
36   Piechota, M. *et al.* Transcriptional signatures of steroid hormones in the striatal neurons
and astrocytes. *BMC Neuroscience* **18**, 37, doi:10.1186/s12868-017-0352-5 (2017).
37   Acaz-Fonseca, E., Avila-Rodriguez, M., Garcia-Segura, L. M. & Barreto, G. E. Regulation
of astroglia by gonadal steroid hormones under physiological and pathological conditions.
Progress in Neurobiology **144**, 5-26, doi:10.1016/J.PNEUROBIO.2016.06.002 (2016).
38   Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons,
and vascular cells of the cerebral cortex. *The Journal of neuroscience : the official journal*
of the Society for Neuroscience **34**, 11929-11947, doi:10.1523/JNEUROSCI.1860-14.2014
(2014).
39   Scarr, E. *et al.* Increased cortical expression of the zinc transporter SLC39A12 suggests a
breakdown in zinc cellular homeostasis as part of the pathophysiology of schizophrenia.
npj Schizophrenia **2**, 16002, doi:10.1038/npjschz.2016.2 (2016).
40   Martinez-Garcia, A. *et al.* PLA2G3, a gene involved in oxidative stress induced death, is
associated with Alzheimer's disease. *Journal of Alzheimer's disease : JAD* **22**, 1181-1187,
doi:10.3233/jad-2010-101348 (2010).
41   Yaguchi, H. *et al.* TRIM67 protein negatively regulates Ras activity through degradation
of 80K-H and induces neuritogenesis. *The Journal of biological chemistry* **287**, 12050-
12059, doi:10.1074/jbc.M111.307678 (2012).
42   Aoki-Suzuki, M. *et al.* A family-based association study and gene expression analyses of
netrin-G1 and -G2 genes in schizophrenia. *Biological psychiatry* **57**, 382-393,
doi:10.1016/j.biopsych.2004.11.022 (2005).
43   Eastwood, S. L. & Harrison, P. J. Decreased mRNA expression of netrin-G1 and netrin-
G2 in the temporal lobe in schizophrenia and bipolar disorder. *Neuropsychopharmacology*
: official publication of the American College of Neuropsychopharmacology **33**, 933-945,
doi:10.1038/sj.npp.1301457 (2008).
44   Bowers, J. M., Waddell, J. & McCarthy, M. M. A developmental sex difference in
hippocampal neurogenesis is mediated by endogenous oestradiol. *Biology of Sex*
Differences **1**, 8-8, doi:10.1186/2042-6410-1-8 (2010).
45   Lenz, K. M. & McCarthy, M. M. Organized for sex - steroid hormones and the developing
hypothalamus. *The European journal of neuroscience* **32**, 2096-2104, doi:10.1111/j.1460-
9568.2010.07511.x (2010).
46   Lenz, K. M. & McCarthy, M. M. Organized for sex - steroid hormones and the developing
hypothalamus. *The European journal of neuroscience* **32**, 2096-2104, doi:10.1111/j.1460-
9568.2010.07511.x (2010).

	47	Clarkson, J. & Herbison, A. E. Hypothalamic control of the male neonatal testosterone surge. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **371**, 20150115, doi:10.1098/rstb.2015.0115 (2016).
	48	Swaab, D. F. Sexual differentiation of the brain and behavior. *Best Practice & Research Clinical Endocrinology & Metabolism* **21**, 431-444, doi:10.1016/j.beem.2007.04.003 (2007).
	49	Salmaso, N. & Woodside, B. Fluctuations in astrocytic basic fibroblast growth factor in the cingulate cortex of cycling, ovariectomized and postpartum animals. *Neuroscience* **154**, 932-939, doi:10.1016/J.NEUROSCIENCE.2008.03.063 (2008).
	50	Arnsten, A. F. T. Ameliorating prefrontal cortical dysfunction in mental illness: inhibition of phosphotidyl inositol-protein kinase C signaling. *Psychopharmacology* **202**, 445-455, doi:10.1007/s00213-008-1274-9 (2009).
	51	Oberlander, J. G. & Woolley, C. S. 17β-Estradiol Acutely Potentiates Glutamatergic Synaptic Transmission in the Hippocampus through Distinct Mechanisms in Males and Females. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **36**, 2677-2690, doi:10.1523/JNEUROSCI.4437-15.2016 (2016).
	52	De Vries, G. J. Minireview: Sex Differences in Adult and Developing Brains: Compensation, Compensation, Compensation. *Endocrinology* **145**, 1063-1068, doi:10.1210/en.2003-1504 (2004).
	53	Mizukami, H. *et al.* KDM5D-mediated H3K4 demethylation is required for sexually dimorphic gene expression in mouse embryonic fibroblasts. *The Journal of Biochemistry* **165**, 335-342, doi:10.1093/jb/mvy106 (2018).
	54	Arnold, A. P. *et al.* Minireview: Sex Chromosomes and Brain Sexual Differentiation. *Endocrinology* **145**, 1057-1062, doi:10.1210/en.2003-1491 (2004).
	55	Galvez-Contreras, A. Y., Campos-Ordonez, T., Gonzalez-Castaneda, R. E. & Gonzalez-Perez, O. Alterations of Growth Factors in Autism and Attention-Deficit/Hyperactivity Disorder. *Frontiers in psychiatry* **8**, 126, doi:10.3389/fpsyt.2017.00126 (2017).
	56	Matute, C., Melone, M., Vallejo-Illarramendi, A. & Conti, F. Increased expression of the astrocytic glutamate transporter GLT-1 in the prefrontal cortex of schizophrenics. *Glia* **49**, 451-455, doi:10.1002/glia.20119 (2005).
	57	Mayoral, S. R., Omar, G. & Penn, A. A. Sex differences in a hypoxia model of preterm brain damage. *Pediatric research* **66**, 248-253, doi:10.1203/PDR.0b013e3181b1bc34 (2009).
	58	van der Kooij, M. A. *et al.* Mild neonatal hypoxia–ischemia induces long-term motor- and cognitive impairments in mice. *Brain, Behavior, and Immunity* **24**, 850-856, doi:https://doi.org/10.1016/j.bbi.2009.09.003 (2010).
	59	Gerstner, B. *et al.* 17beta-estradiol protects against hypoxic/ischemic white matter damage in the neonatal rat brain. *Journal of neuroscience research* **87**, 2078-2086, doi:10.1002/jnr.22023 (2009).
	60	Salmaso, N. *et al.* Environmental enrichment increases the GFAP+ stem cell pool and reverses hypoxia-induced cognitive deficits in juvenile mice. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **32**, 8930-8939, doi:10.1523/JNEUROSCI.1398-12.2012 (2012).
	61	Gillies, G. E. & McArthur, S. Estrogen actions in the brain and the basis for differential action in men and women: a case for sex-specific medicines. *Pharmacological reviews* **62**, 155-198, doi:10.1124/pr.109.002071 (2010).

62 Bowen, R. S., Ferguson, D. P. & Lightfoot, J. T. Effects of Aromatase Inhibition on the
Physical Activity Levels of Male Mice. *Journal of steroids & hormonal science* **1**, 1-7,
doi:10.4172/2157-7536.S1-001 (2011).
63 Komitova, M. *et al.* Hypoxia-induced developmental delays of inhibitory interneurons are
reversed by environmental enrichment in the postnatal mouse forebrain. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience **33**, 13375-13387,
doi:10.1523/JNEUROSCI.5286-12.2013 (2013).
64 Salmaso, N. *et al.* Contribution of maternal oxygenic state to the effects of chronic
postnatal hypoxia on mouse body and brain development. *Neuroscience letters* **604**, 12-17,
doi:10.1016/j.neulet.2015.07.033 (2015).
65 Schmitz, C. & Hof, P. R. Design-based stereology in neuroscience. *Neuroscience* **130**, 813-
831, doi:10.1016/j.neuroscience.2004.08.050 (2005).
66 Simard, S. *et al.* Profiling changes in cortical astroglial cells following chronic stress.
Neuropsychopharmacology : official publication of the American College of
Neuropsychopharmacology **43**, 1961-1971, doi:10.1038/s41386-018-0105-x (2018).
67 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford,*
England) **25**, 2078-2079, doi:10.1093/bioinformatics/btp352 (2009).
68 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics (Oxford, England)* **30**, 923-
930, doi:10.1093/bioinformatics/btt656 (2014).
69 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics (Oxford,*
England) **26**, 139-140, doi:10.1093/bioinformatics/btp616 (2010).
70 Kamburov, A., Stelzl, U., Lehrach, H. & Herwig, R. The ConsensusPathDB interaction
database: 2013 update. *Nucleic acids research* **41**, D793-800, doi:10.1093/nar/gks1055
(2013).
71 Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**, 559-559, doi:10.1186/1471-2105-9-559 (2008).

1077

1078

 Figure Captions

 Figure 1. A- Total number of eGFP+ cells in the neocortex. B- Estimated neocortical
 volume. C- Percentage of radial-like eGFP+ cells. * indicates a significant difference
 from all other time points. Bars denote group mean and error bars displayed are ± SEM.

 Figure 2. A- Clustered heat map of individual sample gene expression levels from
 TRAPseq data. B- Represents a heat map of samples manually sorted by sex and age.

 Figure 3. A- Total number of cells expressing GFAP. B- Total number of vimentin+ cells.
 C- Percent GFP+ co-expressing Ki67. D- Representative mosaic of confocal
 pictomicrographs at 10x, whole cortex sagittal section (left), zoom on the molecular
 layer (right). * indicates a significant difference from all other time points; # indicates a
 significant difference between male and female at that time point. Bars denote group
 mean and error bars displayed are ± SEM.

 Figure 4. A- Number of DEGs between early and late phenotype in males and females.
 B- Venn diagram displaying overlap in DEGs between females and males. C- Bar plot
 showing the number of up- and downregulated DEGs between males and females at
 each time point (females=reference group). D- Bar plot showing the number of DEGs
 between time points.

 Figure 5- Plot of modules Eigen-value vs time. Represented is the subset of the male
 and female modules enriched in astrocyte subtype markers. . A- modules enriched in
 AST1, AST2, AST3 subtypes; top row: modules M2, M3, M12; bottom row: F3, F9. B-
 modules enriched in AST4 subtypes; top row: modules M8, M11, M1; bottom row: F6,
 F2, F10. C- modules enriched in AST5 subtypes; top row: modules M2, M6, M1; bottom
 row: F3, F9.

 Figure 6. A- Density of COX2+ GFP+ astroglial cells in the cortex across examined
 timepoints. Bars represent group means, error bars are displayed as ± SEM.
 Representative mosaic reconstruction of Cox2, eGFP staining in a male and female
 mouse within the PFC region of the cortex at P1 (top). A 63X+oil image highlighting a
 Cox2+, eGFP+ astroglia (white arrow) and a Cox2-, eGFP+ astroglia (pink arrow) in the
 PFC (bottom). B- top panel- Graphical illustration of arbitrary levels circulating gonadal
 hormones (testosterone, solid lines; estrogen, broken lines) in males (blue) and females
 (red) overlay with ages examined in the study. B- lower panel- important cortical
 neurulation events represented by colour saturation. Darker saturation indicates when a
 process is more active, lower saturation indicates when a process is less active.
 (Adapted from: Ellis 2004; Gillies and McArthur, 2010; Anderson, 2003; Estes and
 McAllistter, 2016). C- Density of of COX2+ cells in the cortex at P1 in males and
 females treated with estradiol or letrozole. Bars represent group means, error bars are
 displayed as ± SEM. * indicates significance between treatment and vehicle.

Figure 1

Figure 2

A

B

Figure 3

Figure 4

A

B

Early vs. Late

Males — 1727 — 7256 — 2432 — Females

C **D**

Figure 5

A-subtypes 1, 2 & 3

B- subtype 4

C- subtype 5

Figure 6

A

Supplemental Figure 1

Animal numbers for immunohistochemistry

Age/Sex	P1*	P4	P7	P14	P35	Adult
M	4	5	3	4	4	5
F	4	4	4	6	5	5
Total	8	9	7	10	9	10

Animal numbers for TRAPseq

Age/Sex	P1*	P4	P7	P14	P35	Adult
M	4	4	4	5	3	5
F	4	4	8	5	3	4
Total	8	8	12	10	6	9

Animal numbers for organizational hormones study

Sex/Group	Vehicle	Estradiol/Letrozole
Male	6	7
Female	5	6

Antibodies used for immunohistochemistry

Antibody	Supplier	Species	Dilution
anti-GFP	Abcam (ab19370)	Chicken	1:1000
anti-GFAP	Abcam (ab134436)	Chicken	1:500
anti-GFAP	Abcam (ab10062)	Mouse	1:1000
anti-GFAP	Sigma (AB5804)	Rabbit	1:500
anti-GS	Abcam (ab73593)	Mouse	1:500
anti-Ki67	Invitrogen (PA1-21520)	Rabbit	1:500
anti-Vimentin	Invitrogen (PA5-27231)	Rabbit	1:500
Donkey anti-chicken	Alexafluor 488		1:1000
Donkey anti-mouse	Alexafluor 488		1:1000
Donkey anti-rabbit	Alexafluor 546		1:500
Donkey anti-mouse	Alexafluor 546		1:500

Supplemental Figure 2

A- Pre-correction

B- Post-correction

Top- Hierarchical clustering (using the genes with a sd in expression level above the 95%) pre- and post sva correction show the sample-to-sample relationship and relative clustering

Bottom- Principal component analysis plots. Represented are the sample distribution along the first principal component (PC1) and then second principal component (PC2). Shows in brackets os the percentage of the variance captured by PC1 and PC2. Circles: female samples; squares: male samples. Symbols are color coded by age.

Supplemental Figure 3

Supplemental Figure 4

Developmental Changes

Supplemental Figure 5

Stable through Development

Supplemental Figure 6

Low Expression

Supplemental Figure 7

Supplemental Figure 8

Upper Biased but Pan Astro

Supplemental Figure 9

TaqMan Gene Expression Assays	Gene Symbol	Dye	Assay ID
Glutamine Synthetase	Glul	FAM	Mm00725701_s1
Glial Fibrillary Acidic Protein	GFAP	FAM	Mm01253033_m1
RNA Binding Fox-1 Homolog 3	Rbfox3	FAM	Mm01248771_m1
Parvalbumin	Pvalb	FAM	Mm00443100_m1
Ly6/Neurotoxin 1	Lynx1	FAM	Mm01204957_g1
Estrogen Receptor 1	ESR1	FAM	Mm00433149_m1
Androgen Receptor	AR	FAM	Mm00442688_m1
Glutamate Ionotropic Receptor NMDA Type Subunit 2C	Grin2c	FAM	Mm00439180_m1
Progesterone Receptor	PGR	FAM	Mm00435628_m1
G protein-coupled Estrogen Receptor 1	GPER1	FAM	Mm02620446_s1
Actin Beta	Actb PL	VIC	Mm00607939_s1
Actin Beta	Actb PL	VIC	Mm02619580_g1

Supplemental Figure 10

Q-RT-PCR validation of gene expression levels between early (P1-P7) and late timepoints (P14-Adult). Expressed as fold change

Supplemental Figure 11

	1	2	3	4	5	6	7	8	9	10	11	12
1. Gja1 Fold Change	1											
2. Gja1 RPKM	**0.56****	1										
3. Lynx1 Fold Change	0.48**	0.76**	1									
4. Lynx1 RPKM	0.46**	0.87**	**0.84***	1								
5. Esr1 Fold Change	-0.12	-0.02	0.08	0.08	1							
6. Esr1 RPKM	-0.01	-0.07	-0.15	-0.04	**0.39****	1						
7. Ar Fold Change	0.38**	0.51**	0.76**	0.53**	0.29**	-0.13	1					
8. Ar RPKM	0.47**	0.78**	0.91**	0.89**	0.46	-0.18	**0.75****	1				
9. Grin2c Fold Change	0.68**	0.68**	0.50**	0.63**	-0.28*	-0.07	0.18	0.56**	1			
10. Grin2c RPKM	0.53**	0.88**	0.77**	0.89**	-0.03	-0.14	0.48**	0.80**	**0.76****	1		
11. Pgr Fold Change	0.42**	0.53**	0.78**	0.57**	0.24	-0.08	0.96**	0.77**	0.22	0.51**	1	
12. Pgr RPKM	0.43**	0.76**	0.77**	0.83**	0.09	0.11	0.57**	0.84**	0.49**	0.70**	**0.61****	1

* Correlation significant where $p < 0.05$ (2-tailed)
** Correlation significant where $p < 0.01$ (2-tailed)

Correlation Table. Pearson correlations between Q-RT-PCR (fold change) validation of gene expression levels in TRAP samples and RPKM levels from TRAP-RNASeq.

Supplemental Figure 12

Supplemental Figure 12. Contingency table of the male modules (rows) and female modules (columns). The module color, and the total number of genes in the module label both the rows and the columns. Numbers within the table represent the counts of genes in the intersection of the corresponding row and column module. The table is color-coded by the $-\log_{10}$ of the Bonferroni corrected Fisher exact test p value (legend on then right).