Characterizing of novel magnetotactic bacteria using a combination of magnetic

column separation (MTB-CoSe) and *mamK*-specific primers

Veronika V. Koziaeva[1], Lolita M. Alekseeva[1,2], Maria M. Uzun[1,2], Pedro Leão[3],

Marina V. Sukhacheva[1], Ekaterina O. Patutina[1], Tatyana V. Kolganova[1], Denis S.

Grouzdev[1*]

[1] Institute of Bioengineering, Research Center of Biotechnology of the Russian

Academy of Sciences, Moscow, 119071, Russia

[2] Lomonosov Moscow State University, Moscow, 199991, Russia,

[3] Instituto de Microbiologia Professor Paulo de Góes, Universidade Federal do Rio de

Janeiro, Rio de Janeiro, RJ, 21941-902, Brazil

*Corresponding author

Denis S. Grouzdev. Moscow, Prospect 60 Letiya Oktyabrya 7 bld 1, 117312, Russia.

+74991351240, denisgrouzdev@gmail.com

Running title

Novel approaches for MTB investigation

ABSTRACT

Magnetotactic bacteria (MTB) belong to different taxonomic groups according to 16S

rRNA or whole-genome phylogeny. Magnetotactic representatives of the class

Alphaproteobacteria and the order *Magnetococcales* are the most frequently isolated

MTB in environmental samples. This bias is due in part to limitations of currently

available methods to isolate MTB. Here we describe a new approach for isolation of MTB cells that does not depend on cell motility and will allow collecting bacteria both south- and north-seeking movement. We also designed a specific primer system for the gene encoding the MamK protein that effectively detects diverse MTB phylogenetic groups in any sample type. The combination of these two approaches allowed the identification of a novel MTB belonging to the family *Syntrophaceae* of the class *Deltaproteobacteria*. Moreover, we found that *Nitrospirae* bacteria predominated in the MTB fraction of a sample taken from Lake Beloe Bordukovskoe near Moscow, Russia. We describe the novel dominant *Nitrospirae* bacterium 'Candidatus Magnetomonas plexicatena' and propose its taxonomic name.

IMPORTANCE

Among magnetotactic bacteria (MTB), the members of phyla *Proteobacteria*, *Nitrospirae* and 'Ca. Omnitrophica' have been studied extensively using the existing approaches. However, in recent years, analyses of the metagenomic databases have revealed the presence of MTB in phylogenetic groups, which had not been previously detected using standard approaches. This finding indicates that the biodiversity of MTB is much broader than is currently known. The difficulty of identifying MTB based on comparative analysis of 16S rRNA genes lies in the existence of closely related species of non-magnetotactic bacteria. Moreover, there is an absence of 16S rRNA MTB sequences from such taxonomic groups as 'Latescibacteria' and *Planctomycetes*. In addition, the standard methods of separating MTB can benefit bacteria with high motility. Developing novel strategies for investigation offers great

47 promise towards identifying MTB groups. We have proposed new approach to

48 separate MTB cells from environmental samples and have also proposed a specific

49 primer system for the MTB identification.

INTRODUCTION

Prokaryotes having directed active movement that is guided by geomagnetism are

collectively called magnetotactic bacteria (MTB) (1). The term MTB has no

taxonomic meaning such that it representatives are physiologically, morphologically

and phylogenetically different and share only the ability to synthesize special

organelles called magnetosomes. Magnetosomes consist of nanosized magnetite

(Fe_3O_4) (2) or greigite (Fe_3S_4) (3–5) crystals surrounded by a lipid bilayer membrane

having proteins specific to the organelle (6, 7). Magnetosomes frequently assemble

into chains inside the cell (8). MTB evolved the ability to conduct a special type of

movement called magnetotaxis, which is based on orientation relative to magnetic

field lines (9, 10).

MTB are found among the phyla *Proteobacteria*, *Nitrospirae*, *Planctomycetes*, the

candidate phylum *'Omnitrophica'* and the candidate phylum *'Latescibacteria'* (11–

15). Despite the high diversity of MTB found in environmental samples, they are

difficult to isolate in axenic culture. Therefore, culture-independent techniques are an

indispensable approach to study these bacteria.

Unlike other uncultivated bacteria, MTB can be isolated from environmental samples

based on their magnetotatic activity. Isolation methods include primary separation

using a magnet (16), as well as various magnetic trap techniques, such as 'race-track'

(17) or 'MTB-trap' (18). Although such magnetic enrichment approaches are

efficient for isolation of environmental MTB, members of the *Alphaproteobacteria*

and *'Etaproteobacteria'* classes tend to predominate when these separation methods

are used (14, 18–23). Since they are in higher concentration on the environment, are

usually faster (24, 25) and more resistant to higher oxygen concentration than other

MTB (26). The time used in standard concentration methods (20–30 minutes) make it

hard the separation of MTB that have slow swimming ability (e.g., *Magnetovibrio*

blakemorei MV-1) (27). Longer concentration time leads to the emergence of non-

magnetic pollutants, which makes the enrichment of slowly swimming cells difficult.

The magnetic field intensity, the distance to the magnet and chemotaxis can also

negatively affect enrichment efficiency of MTB. Due to these and other

characteristics, a narrow group of bacteria is preferentially collected using standard

methods, which gives a biased representation of MTB diversity in the microbial

community (26).

In some cases, nonmagnetic pollutants may interfere with the phylogenetic definition

of MTB. This interference is because the ability to synthesize magnetosomes is not a

taxonomic descriptor, and both MTB and non-MTB may belong to the same

taxonomic group. For example, the magnetotactic *Desulfovibrio magneticus* RS-1 is

closely related to *D. carbinolicus* and *D. burkinensis* (16S rRNA sequence similarity

of 99.5% and 99.4%, respectively), which do not form magnetosomes (28).

Furthermore, the frequency of MTB in the microbial community is low (~1–3%), as

shown in studies using next generation sequencing (NGS) techniques (29, 30).

Therefore, identification of MTB using 16S rRNA sequences alone is difficult, since

there are no universal identification markers for MTB. To date, MTB-specific

primers are available only for the identification of individuals from the

Magnetospirillum genus (31) and freshwater coccus from *Magnetococcales* order

(32). The absence of MTB reference strains in different taxonomic groups also

complicates the description of MTB diversity based on 16S rRNA sequence analysis.

For example, putative magnetotosome genes were found in contigs from the draft

genome of bacteria belonging to *'Latescibacteria'* and *Planctomycetes* phyla, but no

16S rRNA sequences were associated with these contigs (13, 14). As a result, MTB

having high levels of 16S rRNA similarity to non-MTB and MTB that are present

with low abundance in the community can remain undetected.

Development of improved techniques for detection and separation of MTB is thus

needed for isolation of novel MTB that are less frequent and/or swim slowly.

In this study, we propose a novel approach for MTB isolation called **MTB Co**lumn

Separation (MTB-CoSe) that overcomes the problems associated with bias

separation. Additionally, we developed a universal primer system for identifying

MTB in an environment that addresses the limitations of using 16S rRNA profiling

for MTB populations. We chose as an universal marker gene for MTB the gene

encoding the actin-like protein MamK, which drives the ordered arrangement of the

magnetosome chain in MTB cells (33).

RESULTS

Primer system design

All known MTB possess the genomic magnetosome island (MAI), comprising clusters of genes that regulate magnetosome mineralization (34). MTB share a common set of genes in the *mamAB* operon, which has been recognized as being indispensable for biomineralization. Currently, there are nine highly conserved primary genes in magnetite- and greigite-producing MTB: *mamA*, *mamB*, *mamM*, *mamQ*, *mamO*, *mamI*, *mamP*, *mamK* and *mamE* (12). For primer system design, we analyzed all nine genes across all known MTB (Table S1). The analysis of conserved regions allowed selection of genes that are amenable to the design of universal primer systems. The *mamM* and *mamB* genes, as well as *mamE* and *mamO*, are homologues, and thus are difficult to differentiate. Therefore, they were excluded from the analysis. Furthermore, *mamA*, *mamP* and *mamQ* genes have conserved sites only within a narrow taxonomic group, which precludes their use in a universal identification system. The *mamI* gene was highly conserved, but has a short nucleotide sequence. Thus, the *mamK* gene was chosen as the most optimal molecular marker for MTB identification. Based on analysis of conserved regions of *mamK*, several primer system variations were constructed and tested. The best results were demonstrated for nested PCR using two pairs of degenerate primers (mamK_79F-mamK_577R and mamK_86F-mamK_521R; Fig. S1). Upon testing this primer system on MTB, a product with the desired length of ~440 bp was obtained. Testing of non-MTB *M. aberrantis* and *E. coli* as negative controls yielded no product. Sequencing of the PCR products obtained with the primer set showed that the *mamK* gene fragment had the correct sequence. In testing the primer system on

DNA from the total microbial community, a product of the correct length was also detected (Fig. 1a).

Together these results showed that the method of magnetotactic bacteria identification based on *mamK* as a molecular marker is applicable not only to axenic cultures but also to total DNA from whole communities.

MTB-CoSe approach

Magnetic columns were used as a new method to separate MTB from enrironmental samples. This method minimizes oxygen sensitivity and motility bias seen in classical methods of MTB magnetic separation. In addition, it performs north- and south-seeking MTB isolation simultaneously.

Initially, testing was performed on an axenic culture of *M. magneticum* AMB-1 mixed with *E. coli* DH10B at a 1:3 ratio. The cell suspension was loaded into the column and rinsed with PBS buffer. The presence of *E. coli* cells was checked in the flow-through and after elution of the magnetic fraction. Separation of the cell suspension was successful in that *E. coli* was found in the flow-through and AMB-1 was retained in the column. We then tested this method on a bottom-sediment sample from Lake Beloye Bordukovskoe, which is characterized by low pH levels, a reduced quantity of anions and cations and a high organic carbon content (Table S2). Immediately after MTB-CoSe, a light microscopy examination revealed the presence of $1.12 \pm 0.06 \times 10^6$ MTB in 20 ml of the total sample, indicating that up to 10^8 MTB cells can be obtained from one liter of sediment. Vibrios were the most common MTB morphotype, and ovoid cells, which accounted for about 30% of the magnetic

fraction, were also detected. Nonmagnetic cells were also present in small quantities.

The isolated MTB cells were divided into two parts. One part was used to isolate

DNA for metagenome sequencing and creating libraries of the *mamK* and 16S rRNA

gene sequences. The second part was fixed for FISH/TEM studies.

Diversity analysis by *mamK* and 16S rRNA clone libraries

Clone libraries from the *mamK* and 16S rRNA gene sequences consisted of 346 and

449 sequences, respectively, after the removal of chimeras and low-quality reads.

Based on the MamK sequences, a phylogenetic analysis was conducted that yielded

six OTUs. The tree topology was compared with the tree based on the 16S rRNA

sequence, and identity with the nearest microorganisms was obtained to identify the

phylogenetic position of the detected OTUs (Fig. 2). Based on the MamK sequences,

the OTUs LBB_01, LBB_02 and LBB_03 were the most well-represented and

formed a separate branch on the phylogenetic tree together with the MamK sequences

of well-known strains of the Nitrospirae phylum. OTU LBB_01 was dominant (87%

of the clone library) and formed a separate clade. The closest to LBB_01, with 61.8%

sequence identity, was MamK sequence from *'Candidatus* Magnetobacterium

bavaricum*'* TM-1. The dominant group LBB_01 was also present on the

phylogenetic tree based on 16S rRNA sequences and forms a separate branch within

the phylum *Nitrospirae*. The identity with *'Ca*. Magnetobacterium bavaricum*'* TM-1,

'Ca. Magnetominusculus xianensis*'* and *'Ca.* Magnetoovum mohavensis*'* 16S rRNA

sequences were 92.2%, 89.9% and 89.0%, respectively. The second OTU, LBB_02,

clustered with the MamK sequence of *'Ca*. Magnetominusculus xianensis*'* HCH-1

with 85.4% homology. OTU LBB_02 was present on the tree based on 16S rRNA

and also clustered with *‘Ca*. Magnetominusculus xianensis*’* HCH-1. The sequence

similarity was 99.2%, which indicated that they belong to the same species. The third

most-frequent OTU (7.2%) was LBB_03. On the phylogenetic tree based on MamK,

LBB_03 formed a separate clade with Nitrospirae bacterium MYbin3 with 87.5%

sequence similarity. OTU LBB_03 was also present on the tree of 16S rRNA

sequences; it formed a branch together with cultivated members of the genus

Thermodesulfovibrio of the family *Nitrospiraceae* and clustered with C_I49, *‘Ca*.

Thermomagnetovibrio*’* HSMV-1 and MYbin3 sequences. The sequence identity with

these three species was 95.8%, 91.2% and 92.8%, respectively. The level of

similarity with 16S rRNA sequence from *‘Ca*. Magnetobacterium bavaricum*’* TM-1

was 87.9%, which indicates that they may belong to different families. The sequence

similarity of the 16S rRNA sequence of OTUs LBB_01 and LBB_02 was 84–86%

with the closest cultivated strain *Thermodesulfovibrio hydrogeniphilus* of the family

Nitrospiraceae, which indicates that they belong to a separate family. In accordance

with the taxonomy proposed by Parks et al. (35), the level of similarity and branching

order suggest their affiliation with the family *‘Ca*. Magnetobacteriaceae*’*. OTU

LBB_03 and *‘Ca* Thermomagnetovibrio*’* HSMV-1 belong to the same family as

strains of the genus *Thermodesulfovibrio*, but form separate genera.

The minor groups were OTUs LBB_05 and LBB_06, which clustered with the

MamK sequences from MTB belonging to *Magnetococcales* and *Rhodospirillales*

order; they represented 0.3% and 0.9% of the *mamK* clone library, respectively. OTU

LBB_05 clustered with MamK belonging to Alphaproteobacterium WMHbin7 and

clustered with the MamK sequence obtained in the fosmid library Fos002 described

by Jogler et al. (18). The level of similarity with MamK of WMHbin7 and MamK

from Fos002 was 100%. The level of similarity with MamK of the nearest cultured

organism, *Magnetospirillum magneticum*, was 74.6%. OTU LBB_05 was found in

the 16S rRNA library and had a high level of similarity with the magnetotactic vibrio

LM-1 (98.5%) and the bacterium WMHbin7 (98.3%). Due to the high level of

similarity in both the MamK and 16S rRNA sequences, it is possible that OTU

LBB_05, bacterium LM-1, WMHbin7 and the bacterium from the fosmid library

Fos002 are the same species.

OTU LBB_06 formed a separate branch with the MamK of HA3dbin1, belonging to

the order *Magnetococcales*. The level of similarity between sequences was 98.4%.

The level of similarity with the MamK of the cultivated coccus *'Magnetofaba*

australis' IT-1 was 66.4%. OTU LBB_06 was also present on the 16S rRNA

phylogenetic tree and had 98.9% similarity to *Magnetococcus* sp. clone OTU2.

One interesting finding was the presence of OTU LBB_04, which had no MamK that

was closely related to any known MTB. The level of similarity to the closest MamK

of Nitrospirae bacterium MYbin3 was 69.2%, with coverage of 45%. Thus, the

phylogenetic affiliation of OTU LBB_04 required a more thorough analysis. After

searching for the nearest homologues for MamK of OTU LBB_04, a protein with a

high sequence identity (97.0%) was found in the IMG database. This *mamK* gene was

found in metagenome 3300021602, which was obtained by Neufeld et al. (36) from a

boreal shield lake in Ontario. After binning of the metagenome, a bin

(MAG_21602_syn32) containing the target *mamK* sequence was obtained (Table S3).

The 16S rRNA sequence of the received bin was also identified (acc. №

Ga0194060_100555694). Analysis of that sequence on GTDB showed that it belongs

to the order '*Syntrophales*' and, according to NCBI, the order *Syntrophobacterales*.

MTB of this taxonomic group have not been previously detected. On the 16S rRNA

phylogenetic tree, OTU LBB_04 formed a separate branch with representatives of the

Syntrophaceae family. The 16S rRNA sequence derived from the

MAG_21602_syn32 was also included in this cluster and had 96.8% identity with the

16S rRNA gene of OTU LBB_04. The level of similarity with the closest validly

described strains, *Syntrophus aciditrophicus* and *Smitella propionica*, was 93.5% and

91.2%, respectively, which indicates that OTU LBB_04 belongs to a novel genus

within the family *Syntrophaceae*.

In addition to the findings described above, there were sequences on the 16S rRNA

tree belonging to the genera *Sphingomonas*, *Halomonas*, *Rhodococcus*, and

Pseudomonas, constituting 12.2%, 10.4%, 4.8% and 2.4% of the entire library,

respectively. Members of these genera can produce metal particles (37), meaning that

they can be attracted by a magnet alongside MTB. Thus, for all OTUs identified on

the MamK tree, the corresponding OTU on the 16S rRNA tree was determined.

Based on those 16S rRNA gene sequences, probes were designed to identify the

morphology of the matching MTB by FISH-TEM.

FISH-TEM evidence for new MTB groups

To correlate the phylogenetic data with the MTB morphotype, FISH-TEM was

performed. The LBB_01 probe hybridized only with vibrioid-shaped bacteria (Fig.

3a-c). TEM images of the same area used for FISH analysis (Fig. 3d) revealed that

this group of MTB presented a thick chain of magnetosomes organized along the

long axis of the bacterial cell body (Fig. 3e). The observed magnetosomes were

anisotropic (Fig. 3f) and presented [111] as the elongation axis. In addition, the fast

Fourier transform (FFT) pattern of the magnetosome crystalline structure

corresponded to the iron oxide magnetite (Fig. 3g). The LBB_03 probe correlated to

another vibrioid cell presenting anisotropic magnetosomes in a single chain (Fig. 4a-

e), whereas the LBB_02 probe was associated with a small, ovoid MTB with two

magnetosome chains of anisotropic crystals (Fig. 4f-j). The probe LBB_04 identified

a rod-shaped MTB with disorganized magnetosomes located close to the center of the

bacterial cell body (Fig. 4k-o).

Cell morphology of LBB_01

Further analysis of cells compatible with those identified using the LBB_01 probe

confirmed that these cells have a vibrioid shape and a 2.0 ± 0.4 μm and 0.5 ± 0.1 μm

length and width, respectively (n=32). Each cell contained 33 ± 9 anisotropic

magnetite magnetosomes (n=32) organized as described above (Fig. 3e). The

magnetosome tips were not always sequentially oriented and were sometimes pointed

in opposite directions (Fig. 3f), as was previously seen for other MTBs with

anisotropic magnetosomes (38). Immature crystals were found in different chain

segments. A detailed analysis of the magnetosome size showed that they have a mean

length and width of 108 ± 21.1 nm and 45 ± 8.1 nm, respectively, and a mean shape

factor (width/length) of 0.45 (n = 1061; Fig. S2).

General information and phylogeny of the LBB_01 genome

Using metagenomic data, we assembled the genome of the dominant group LBB_01 into 1154 contigs that included 2.3 Mbp with N_{50} equal to 3183. The completeness of the genome assembly was 65.15% with 1.87% contamination. The average G + C content was 42.1 mol%.

To determine the phylogenetic position of the LBB_01 genome we obtained, we built a phylogenomic tree (Fig. 5). *'Ca*. Magnetobacterium bavaricum*'* TM-1 and *'Ca*. Magnetoovum chiemensis*'* CS-04 were not included in the analysis because they failed the quality check due to a high percentage of contamination (34% and 17%, respectively). The genomes quality scores were determined as completeness − 5x contamination and had a threshold equal to 50. Bacterial genomes for which the quality score was less than 50 were not selected for analysis, according to Parks et al. (35). These two species had quality scores of -78.41 and 3.11, respectively, and thus were not analyzed further. As a result, the LBB_01 genome and the genomes of members of the genus *Thermodesulfovibrio* formed a separate branch from *Nitrospira moscoviensis* on the phylogenomic tree. Inside this branch, the LBB_01 genome, together with other MTB, clustered separately from the genus *Thermodesulfovibrio*. In addition, based on the 16S rRNA sequence, LBB_01 formed a branch separate from the genera *'Ca*. Magnetobacterium*'* and *'Ca*. Magnetominusculus*'*. Thus, based on the phylogenomic analysis, the strain LBB_01 represented a novel species and belonged to a novel genus within the family *'Ca*. Magnetobacteriaceae*'*.

Magnetosome genes in the LBB_01 genome

Analysis of the LBB_01 genome revealed genes for 17 magnetosome proteins that

were identified in several contigs (Fig. S3a; Table S4). Among them, several proteins

coded by the main *mamAB* operon, such as MamA, MamE, MamI, MamK, MamM

and MamO-Cter, were found. MamK sequence was identical to OTU LBB 01

sequence retrieved from a clonal library. Homology analyses based on amino acid

sequences of those Mam proteins showed that the magnetosome proteins of the

strains belonging to the *Nitrospirae* phylum (Table S4) share the highest identity with

them. A phylogenetic analysis based on the concatenated amino acid sequences of

MamAIKMP proteins showed that LBB_01 clustered together with the Mam proteins

of *'Ca* Magnetobacterium*'* and *'Ca* Magnetominusculus'. However, upon analysis of

conserved Mam proteins, strain LBB_01 formed a separate branch on the

phylogenomic tree (Fig. S3b). A comparison of the topology of the phylogenomic

tree of core proteins and the tree of concatenated proteins MamAIKMP indicated a

vertical inheritance of strain LBB_01 MAI.

In addition to proteins encoded by the main operon, 10 *mad* related to the

magnetotactic *Deltaproteobacteria* and *Nitrospirae* were also found (39). Three of

them formed a small genomic cluster comprising *mad24*, *mad25* and *mad26*. *Mad28*

was present in two copies. *Man1*, a gene found only in the genome of MTB

belonging to the *Nitrospirae* phylum, was also identified in the LBB_01 genome at a

site adjacent to *mamK* (40). This organization of magnetosome genes resembled the

island of *'Ca* Magnetobacterium casensis*'* MYR-1. The specific operons *mamGFDC*,

mamXY and *mms6* related to *Alphaproteobacteria* and *Magnetococcales* were not

found.

DISCUSSION

The amount of information concerning MTB has been increasing significantly. Among the growing body of genomic data, new taxonomic groups of MTB have been found, which indicates a wider diversity of MTB than was previously known (13, 14). Developing new approaches to study MTB in natural samples will expand our knowledge of the diversity and distribution of MTB in different environments. In this study, we proposed a new approach to study MTB diversity that allows collection of cells without relying on motility or taxis characteristics and uses a universal marker gene. A large number of MTB cells can be collected directly from the bottom sediment, thus bypassing the collection stage that requires a magnet. We demonstrated that a large fraction of the gathered cells can be used to obtain DNA suitable for sequencing the metagenome, whereas the separated cells could be subjected to single-cell sorting for analysis of minority groups. In addition, due to the preservation of viability, the cells can be used to produce enriched cultures.

To identify certain groups of bacteria in microbial communities, one effective method is to identify genes specific to bacterial groups (41). Currently, there are many primer systems used to study bacteria that share some functional trait (42–46). For instance, genes encoding two reaction center proteins, PufLM (47, 48) and the Fenna-Matthews-Olson-protein (FmoA) (49) are used to study bacterial photosynthesis, whereas ammonia monooxygenase (*amoCAB*) genes (50) are mostly applied to investigate ammonia-oxidizing archaea. However, there is no universal primer system for all MTB, even though these bacteria have magnetosome mineralization

genes that could serve as primer targets. Here we developed a MTB-specific primer

system and tested it on the magnetic fraction collected from Lake Beloye

Bordukovskoe. We found that *mamK* gene sequences can be used to identify and

distinguish different taxonomic groups of MTB. Moreover, *mamK* can be used as a

marker gene to investigate the diversity of MTB.

By combining novel enrichment and detection approaches, we found many

representatives of the phylum *Nitrospirae* in oligotrophic Lake Beloye

Bordukovskoe. In previous studies, *Magnetococcales* and *Rhodospirillales* were

reported to be the dominant group of MTB in freshwater environments. However,

using both 16S rRNA and *mamK* gene analyses in our approach, we showed that

these two types of MTB were instead minority groups. Our results are consistent with

an earlier study that showed a significant predominance of MTB belonging to the

Nitrospirae phylum when the DNA was isolated directly from the sediment without

prior separation of the MTB. (29). Our finding is also supported by evidence that

Nitrospirae is the dominant MTB phylum in the freshwater lakes Miyun and

Chiemsee in China and Germany, respectively (51). MTB of the orders

Magnetococcales and *Rhodospirillales* may also be overrepresented due to the high

number of reference sequences among the *Alphaproteobacteria* and

'Etaproteobacteria' classes. As such, studies of MTB diversity typically focused on

bacteria in these two classes. Studying the diversity of MTB based on a functional

gene will expand our understanding of MTB diversity in various environments.

Using both 16S rRNA and *mamK* gene analyses, we identified *mamK* and

corresponding 16S rRNA genes that belonged to a previously unknown taxonomic

group of MTB. The 16S rRNA analysis classified the bin MAG_21602_syn32 related to OTU LBB_04 into the class *Deltaproteobacteria*. Furthermore, FISH/TEM analysis confirmed that the OTU LBB_04 of the *Syntrophaceae* family belongs to a MTB and showed that the *Syntrophaceae* magnetosomes are bullet-shaped.

Using MTB-CoSe, we could collect suitable numbers of cells for DNA extraction and subsequent metagenome sequencing. We obtained the metagenome of the dominant group of the phylum *Nitrospirae*, which currently contains only one family, *Nitrospiraceae*, comprising 3 genera: *Leptospirillum*, *Nitrospira* and *Thermodesulfovibrio*. The branching order of the phylogenomic tree and the 15% difference in 16S rRNA sequences with the closest cultivated species, *Thermodesulfovibrio hydrogeniphilus*, suggest that strain LBB_01 belongs to a separate family. The 8% sequence difference from bacteria of the candidate genus '*Ca.* Magnetobacterium' indicated that strain LBB_01 likely represents a novel genus and species within the candidate family '*Ca.* Magnetobacteriaceae', for which we propose the name '*Ca.* Magnetomonas plexicatena'. The family *'Ca.* Magnetobacteriacea'* currently includes several candidate genera to which only MTB belong: *'Ca.* Magnetobacterium, '*Ca*. Magnetoovum*', '*Ca*. Magnetominusculus*' and the proposed genus *'Ca.* Magnetomonas*'*. Members of these genera differ in their morphology: '*Ca.* Magnetobacterium bavaricum', '*Ca.* Mb. Casensis' and the bacteria of Lake Miyun '*Ca.* Mb. bavaricum*'-like are rod-shaped and synthesize multiple chains of bullet-shaped magnetosomes (51). Meanwhile, '*Ca.* Magnetoovum chiemensis' CS-04 (52) and '*Ca.* Mo. Mohavensis' LO-1 are large ovoid and synthesize only 100–200 magnetosomes per cell (53). For bacteria of the genera '*Ca.*

Magnetominusculus', the morphology of '*Ca.* Mm. xianensis' HCH-1, was

previously unknown (54), but our analysis showed that these bacteria are small ovoid

and synthesize two bundles of bullet-shaped magnetosomes. The newly described

genus '*Ca.* Magnetomonas' has a vibrioid shape and synthesizes one bundle of

bullet-shaped magnetosomes.

CONCLUDING REMARKS

This study presents novel approaches to investigating magnetotactic bacteria that

allowed us, for the first time, to identify MTB of the family *Syntrophaceae*. Our

methods using *mamK* as a marker gene and column separation allowed the

identification of low-abundance MTB in different microbial communities. Our

approach could be used to characterize magnetotactic representatives from

'*Latescibacteria*' and *Planctomycetes* phyla, for which the magnetosome morphology

has not yet been determined.

TAXONOMIC PROPOSALS

Phylogenetic analysis of the LBB_01 genome classified it as a novel genus. We

propose Latin names for the novel candidate genus and species:

Candidatus Magnetomonas

Magnetomonas (Ma.gne.to.mo'nas. Gr. n. *magnes*, *-etos* a magnet; N.L. pref.

magneto- pertaining to a magnet; N L. fem. n. *monas* unit, monad; N.L. fem. n.

Magnetomonas a magnetic monad)

Candidatus Magnetomonas plexicatena

Magnetomonas plexicatena (ple.xi.ca.te'na. L. past part. *plexus* interwoven; L. fem.

n. *catena* chain; N.L. fem. n. *plexicatena* an interwoven chain)

MATERIALS AND METHODS

Sampling and physicochemical analysis

Water and sediment samples were taken from the freshwater lake Beloye

Bordukovskoe, Shatura District, Russia (55°37'56"N, 39°44'38"E). Samples having

1:2 sediment:water were incubated in the dark at room temperature for several

months. Water from the microcosm was analyzed according to 41 parameters at the

National Research Nuclear University MEPhI water quality control testing

laboratory. The elemental composition of organic carbon (C%) in the sediment was

measured using a Flash 1112 elemental analyzer coupled with a Thermo-Finnigan

Delta V Plus isotope mass spectrometer (Thermo Fisher Scientific, USA) at the Core

Facility of IPEE A.N. Severtsov RAS.

***mamK* primer system design and selection of amplification conditions**

We searched for nucleotide sequences of magnetosome genomic island (MAI) genes

in GenBank and IMG databases. Nucleotide sequence alignment procedures were

performed using the ClustalW algorithm (55). Primer design was carried out using

the program Oligo6 (Molecular Biology Insights, USA).

The selection of PCR conditions and testing of the ability of the designed primers to

identify the target product were performed using end-point PCR. The reaction buffer

(25 µl) had the following composition: 5 µl Mas^{CFE}TaqMIX-2025 buffer (Dialat

LTD, Russia), 0.5 pmol/µl each primer, 0.04% BSA and 10–50 ng of a DNA

template. Amplification was performed on a Mastercycler gradient instrument

(Eppendorf, Germany). The temperature-time profile of the reaction was: initiation, 3

min at 95 °C; 4 cycles of 30 s at 95 °C, 40 s at 58 °C, and 1 min at 72 °C; 36 cycles

of 30 s at 95 °C, 40 s at 52 °C, 1 min at 72 °C; and a final cycle of 7 min at 72 °C. A

fragment of the *mamK* gene was detected using nested PCR using two pairs of

degenerate primers: external (mamK_79F and mamK_577R) and internal

(mamK_86F and mamK_521R) (Table S5).

The efficiency, specificity and versatility of the primer system was tested on axenic

cultures of the genus *Magnetospirillum* (*M. moscoviense* BB-1, *M. kuznetsovii* LBB-

42 and *M. caucaseum* SO-1) and on a sample from the River Uda containing the

magnetotactic coccus UR-1 of the order *Magnetococcales* only. DNA from the

nonmagnetic bacteria *Escherichia coli* DH10B and *M. aberrantis* SpK served as

negative controls. *M. aberrantis* is closely related to MTB and cannot form

magnetosomes (56). *E. coli* carries the gene *mreB*, which codes for a rod-shape

determining protein and has high sequence similarity to *mamK*. Amplified 16S rRNA

sequences were used as controls for the DNA template. The resulting PCR products

were sequenced to control nonspecific annealing of the primers. Amplification of the

mamK gene was also verified in DNA isolated from whole microbial communities

from the Soda Lake Kiran and Moskva River, which contain different taxonomic

groups of MTB (23).

Magnetotactic bacteria column separation (MTB-CoSe)

The proposed method involves use of MACS® columns (Miltenyi Biotec, Germany)

(Fig. 6). Since large sediment particles can occupy space in the column and in turn

increase the washing time, we introduced additional steps. To desorb cells from large

clay particles, PBS buffer (final concentration 1.25X) and glass microbeads (2 g;

150–200 μm diameter; Sigma-Aldrich, USA) were added to a 50 ml tube containing

20 ml of the samples (sediment + water). Additional filtration and centrifugation

steps allowed removal of large soil particles and concentrated the bacterial cells. The

mixture was then placed on a ThermoShakerTS-100 (Biosan, Latvia) for 15 min at

100 rpm before the homogenate was transferred to a Bunsen flask and filtered using a

paper filter under vacuum pressure to remove large soil particles. The filtrate was

transferred to 50 ml tubes and centrifuged (Eppendorf Centrifuge 5804R) for 10–15

minutes at 8,000 rpm. The majority of the supernatant was discarded, and 2 ml was

retained for cell resuspension. Next, a miniMACS column was washed with 1.25X

PBS buffer and placed on a magnet, and the cell mixture was applied to the column.

MTB cells that adsorbed onto the column were washed 4–5 times with 2 ml 1.25X

PBS buffer until there was a complete absence of cells in the washing liquid. Next,

the column was removed from the magnet, and the MTB cells were eluted twice with

100 μl 1.25X PBS buffer into a clean 1.5 ml tube. For samples having high cell

concentrations, the eluate was reapplied onto the column, which was rinsed to

completely remove nonmagnetic cells. The presence of MTB cells and nonmagnetic

bacteria was assessed by microscopy of 5 μl aliquots of eluted liquid.

This method is suitable for separating MTB from a large sample volume in a

relatively short time. After completing all the manipulations, the magnetic fraction

had low amounts of nonmagnetic pollutants.

Light and electron microscopy

The bacterial morphology after MTB-CoSe was examined using an Eclipse E200

(Nikon, Japan) light microscope. For conventional transmission electron microscopy

(TEM), magnetically enriched cells were deposited on Formvar-coated 300-mesh

copper grids, washed with distilled water and imaged using a Morgagni (FEI, USA)

TEM operated at 80 kV. For high-resolution TEM (HRTEM), the same grids

prepared for conventional TEM were imaged using a Tecnai G2 F20 FEG (FEI,

USA) operated at 200 kV and equipped with a 4k×4k Gatan UltraScan 1000 CCD

camera. Measurements and fast Fourier transform from the HRTEM images were

obtained using Digital Micrograph software (Gatan, USA).

Amplification, cloning and sequencing fragments of genes encoding 16S rRNA

and *mamK*

Genomic DNA was extracted using a DNeasy PowerSoil kit (Qiagen, Netherlands)

according to the manufacturer's instructions. Amplification of the 16S rRNA

sequences was performed using the universal primers 27F and 1492R (57). The PCR

products were purified from a 0.7% agarose gel using the Wizard SV Gel and PCR

Clean-Up System kit (Promega, United States) according to the manufacturer's

recommendations. Cloning was carried out with the pGEM-T Easy Vector System I

(Promega, USA). *E. coli* DH10B competent cells were transformed using a

multiporator (Eppendorf, Germany). The target insert of the *mamK* gene was

sequenced using the primer M13F, and 16S rRNA was sequenced using the primers

341F, 530F, 1114F and 519R (58). Sequencing was performed using the Sanger

method on an ABI3730 DNA Analyzer sequencer (Applied Biosystems, USA) with

the BigDye® Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, USA), as

recommended by the manufacturer.

Phylogenetic analysis

Nucleotide and amino acid sequences were aligned using MAFFT (59). The GTDB-

Tk v. 0.1.3 toolkit was used to find 120 single-copy bacterial marker genes and

construct multiple alignments of concatenated single-copy gene sequences (35).

Concatenated *mamAIKMP* sequences were obtained using Gblocks (60). Obtained

sequences of the MamK and 16S rRNA were grouped in OTUs using identity

threshold 97%. Phylogenetic analysis was performed using the IQ-TREE program

(61) with selection of an evolutionary model using ModelFinder (62) and estimation

of branch supports using UFBoot2 (63). The MamK and 16S rRNA sequences for the

identified OTUs were deposited in GenBank under the accession numbers

MK636828-33 and MK63285-90, respectively.

Fluorescence in situ hybridization/transmission electron microscopy

(FISH/TEM)

For fluorescence *in situ* hybridization (FISH), collected MTB cells were fixed in 3%

paraformaldehyde for 1.5 hours. For the association of the retrieved *rrs* gene

sequences with different MTB morphotypes, a drop of sample was added to center-

marked or index copper grids covered with Formvar for electron microscopy. A thin

layer of carbon was sputtered onto the grids (Balzers CED-030/Baltec) atop each

sample to provide stabilization. FISH was performed on each grid using conditions

and buffers described in Pernthaler et al. (2001) (64) and 30% formamide as the

hybridization buffer. After this procedure, the grids were stained with 0.1 µg/ml 4,6-

diamidino-2-phenylindole (DAPI) for 10 minutes, placed between a glass slide and

cover glass, and observed with a Zeiss AxioImager microscope equipped with an

AxioCam Mrc (Zeiss, Germany).

Probe sets used for FISH were designed based on the *rrs* sequences retrieved from

sequencing of the magnetically enriched samples (Table S5). In addition, a mix of

EUB388I, EUB388II and EUB388III bacterial universal probes labelled with Alexa

488 was used as a control (65).

After performing FISH, the same grids were placed on a Morgagni (FEI, USA)

transmission electron microscope (TEM) operated at 80 kV, and images were taken

in the same region where hybridization with the specific probe occurred.

Genome sequencing

Libraries were constructed with NEBNext DNA library prep reagent set for Illumina,

per the kit protocol. Sequencing was performed using the Illumina HiSeq 1500

platform with single-end 220-bp reads. The dried DNA was dissolved in 50 µl MQ.

Raw reads were quality-checked with FastQC v. 0.11.7

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/), and low-quality reads

were trimmed using Trimmomatic v. 0.36 (66). The quality-filtered reads were

assembled *de novo* with SPAdes v. 3.12.0 using the default settings (67). Annotations

of the scaffolds were carried out using the NCBI Prokaryotic Genome Annotation

Pipeline (68). Results for this whole-genome shotgun project were deposited in

DDBJ/ENA/GenBank under the accession number SNNQ00000000. To evaluate

genome statistics, an automatic assembly quality evaluation tool (QUAST) was used

(69).

Metagenome binning and analysis

The assembled metagenome of Lake Beloe Bordukovskoe and the metagenome

3300021602 obtained in the IMG database were binned using three different tools

(MaxBin2 (70), MyCC (71), and Busy Bee Web (72)) prior to dereplication and

refinement with the DAS Tool (73), which performs a consensus binning to produce

the final bin set. Genome contamination was removed using an approach based on

taxonomic assignments in RefineM v. 0.0.24 (74). Completeness and contamination

rates were assessed using CheckM v. 1.0.12 (75) with the 'lineage wf' command and

default settings. Magnetosome island genes were found using local BLAST

compared with reference sequences of magnetotactic bacteria.

ACKNOWLEDGMENTS

We thank Professor Josh D. Neufeld for permission to use metagenomic data

(3300021602). We thank Professor Aharon Oren for his help with Latin names. We

thank Dr. Svetlana Zhenilo for microscopy assistance and Dr. Vasil Gaisin for

comments that improved an earlier version of this manuscript. We thank Jefferson

Cypriano, CENABIO and Unimicro for helping with accsess to the TEM used in this

work.

FUNDING INFORMATION

This study was funded by RFBR as research project No. 18-34-01005 and the

Ministry of Science and Higher Education of the Russian Federation. This study was

performed using scientific equipment at the Core Research Facility 'Bioengineering'

(Research Center of Biotechnology RAS). The work conducted by the U.S.

Department of Energy Joint Genome Institute, a DOE Office of Science User

Facility, was supported by the Office of Science of the U.S. Department of Energy

under Contract No. DE-AC02-05CH11231.

REFERENCES

1. Blakemore R. 1975. Magnetotactic bacteria. Science (80-) 190:377–379.

2. Frankel RB, Blakemore RP, Wolfe RS. 1979. Magnetite in Freshwater

Magnetotactic Bacteria. Science (80-) 1355–1356.

3. Farina M, Esquivelt DMS, Lins De Barrost HGP. 1990. Magnetic iron-sulphur

crystals from a magnetotactic microorganism. Nature 343:256–258.

4. Mann S, Sparks NHC, Frankelt RB, Bazylinski DA. 1990. Biomineralization of

ferrimagnetic greigite (Fe3S4) and iron pyrite (FeS2) in a magnetotactic

bacterium. Nature 343:258–261.

5.   Heywood BR, Mann S. 1990. Controlled biosynthesis of greigite (Fe3S4) in

magnetotactic bacteria. Naturwissenschaften 77:536–538.

6.   Balkwill DL, Maratea D, Blakemore RP. 1980. Ultrastructure of a

magnetotactic spirillum. J Bacteriol 141:1399–1408.

7.   Gorby YA, Beveridge TJ, Blakemore RP. 1988. Characterization of the

bacterial magnetosome membrane. J Bacteriol 170:834–841.

8.   Frankel RB. 1984. Magnetic guidance of organisms. Annu Rev Biophys Bioeng

13:85–103.

9.   Frankel RB, Bazylinski DA, Johnson MS, Taylor BL. 1997. Magneto-aerotaxis

in marine coccoid bacteria. Biophys J 73:994–1000.

10.   Faivre D, Schuler D. 2008. Magnetotactic Bacteria and Magnetosomes. Chem

Rev 108:4875–4898.

11.   Kolinko S, Jogler C, Katzmann E, Wanner G, Peplies J, Schüler D. 2012.

Single-cell analysis reveals a novel uncultivated magnetotactic bacterium

within the candidate division OP3. Environ Microbiol 14:1709–1721.

12.   Lefèvre CT, Bazylinski DA. 2013. Ecology, diversity, and evolution of

magnetotactic bacteria. Microbiol Mol Biol Rev 77:497–526.

13.   Lin W, Pan Y. 2015. A putative greigite-type magnetosome gene cluster from

the candidate phylum *Latescibacteria*. Environ Microbiol Rep 7:237–242.

14.   Lin W, Pan Y, Bazylinski DA. 2017. Diversity and ecology of and

biomineralization by magnetotactic bacteria. Environ Microbiol Rep.

15. Dziuba M, Koziaeva V, Grouzdev D, Burganskaya E, Baslerov R, Kolganova

T, Chernyadyev A, Osipov G, Andrianova E, Gorlenko V, Kuznetsov B. 2016.

Magnetospirillum caucaseum sp. nov., *Magnetospirillum marisnigri* sp. nov.

and *Magnetospirillum moscoviense* sp. nov., freshwater magnetotactic bacteria

isolated from three distinct geographical locations in European Russia. Int J

Syst Evol Microbiol 66:2069–2077.

16. Bazylinski D, Lefèvre C, Schüler D. 2013. Magnetotactic bacteria, p. 453–494.

In The Prokaryotes: Prokaryotic Physiology and Biochemistry. Berlin.

17. Wolfe RS, Thauer RK, Pfennig N. 1987. A "capillary racetrack" method for

isolation of magnetotactic bacteria. FEMS Microbiol Lett 45:31–35.

18. Jogler C, Lin W, Meyerdierks A, Kube M, Katzmann E, Flies C, Pan Y, Amann

R, Reinhardt R, Schuler D. 2009. Toward cloning of the magnetotactic

metagenome: Identification of magnetosome island gene clusters in

uncultivated magnetotactic bacteria from different aquatic sediments. Appl

Environ Microbiol 75:3972–3979.

19. Pan H, Zhu K, Song T, Yu-zhang K, Lefèvre C, Xing S, Liu M, Zhao S, Xiao

T, Wu L. 2008. Characterization of a homogeneous taxonomic group of marine

magnetotactic cocci within a low tide zone in the China Sea. Environ Microbiol

10:1158–1164.

20. Flies CB, Jonkers HM, De Beer D, Bosselmann K, Böttcher ME, Schüler D.

2005. Diversity and vertical distribution of magnetotactic bacteria along

chemical gradients in freshwater microcosms. FEMS Microbiol Ecol 52:185–

195.

21.   Lin W, Wang Y, Li B, Pan Y. 2012. A biogeographic distribution of

magnetotactic bacteria influenced by salinity. ISME J 6:475–479.

22.   Flies CB, Peplies J, Schüler D. 2005. Combined approach for characterization

of uncultivated magnetotactic bacteria from various aquatic environments. Appl

Env Microbiol 71:2723–2731.

23.   Kozyaeva V V., Grouzdev DS, Dziuba M V., Kolganova T V., Kuznetsov BB.

2017. Diversity of magnetotactic bacteria of the Moskva River. Microbiology

86:106–112.

24.   Zhang S Da, Petersen N, Zhang WJ, Cargou S, Ruan J, Murat D, Santini CL,

Song T, Kato T, Notareschi P, Li Y, Namba K, Gué AM, Wu LF. 2014.

Swimming behaviour and magnetotaxis function of the marine bacterium strain

MO-1. Environ Microbiol Rep 6:14–20.

25.   Araujo ACV, Morillo V, Cypriano J, Teixeira LCRS, Leão P, Lyra S, Almeida

LG de, Bazylinski DA, Vasconcellos ATR de, Abreu F, Lins U. 2016.

Combined genomic and structural analyses of a cultured magnetotactic

bacterium reveals its niche adaptation to a dynamic environment. BMC

Genomics 17:363–375.

26.   Lin W, Tian L, Li J, Pan Y. 2008. Does capillary racetrack-based enrichment

reflect the diversity of uncultivated magnetotactic cocci in environmental

samples? FEMS Microbiol Lett 279:202–206.

27.   Bazylinski DA, Williams TJ, Lefevre CT, Trubitsyn D, Fang J, Beveridge TJ,

Moskowitz BM, Ward B, Schubbe S, Dubbels BL, Simpson B. 2013.

Magnetovibrio blakemorei gen. nov., sp. nov., a magnetotactic bacterium

(*Alphaproteobacteria*: *Rhodospirillaceae*) isolated from a salt marsh. Int J Syst

Evol Microbiol 63:1824–1833.

28.   Sakaguchi T, Arakaki A, Matsunaga T. 2002. *Desulfovibrio magneticus* sp.

nov., a novel sulfate-reducing bacterium that produces intracellular single-

domain-sized magnetite particles. Int J Syst Evol Microbiol 52:215–221.

29.   Xu C, Zhang W, Pan H, Du H, Xiao T. 2018. Distribution and diversity of

magnetotactic bacteria in sediments of the Yellow Sea continental shelf. J Soils

Sediments 2634–2646.

30.   Liu J, Zhang W, Li X, Li X, Chen X, Li J-H, Teng Z, Xu C, Santini C-L, Zhao

L, Zhao Y, Zhang H, Zhang W-J, Xu K, Li C, Pan Y, Xiao T, Pan H, Wu L-F.

2017. Bacterial community structure and novel species of magnetotactic

bacteria in sediments from a seamount in the Mariana volcanic arc. Sci Rep

7:17964.

31.   Lefèvre CT, Schmidt ML, Viloria N, Trubitsyn D, Schüler D, Bazylinski DA.

2012. Insight into the evolution of magnetotaxis in *Magnetospirillum* spp.,

based on *mam* gene phylogeny. Appl Environ Microbiol 78:7238–7248.

32.   Lin W, Pan Y. 2009. Specific primers for the detection of freshwater

alphaproteobacterial magnetotactic cocci. Int Microbiol 12:237–242.

33.   Komeili A, Li Z, Newman DK, Jensen GJ. 2006. Magnetosomes are cell

membrane invaginations organized by the actin-like protein MamK. Science

(80-) 311:242–245.

34.   Bazylinski DA, Frankel RB. 2004. Magnetosome formation in prokaryotes. Nat

Rev Microbiol 2:217–230.

35. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz P. 2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. Nat Biotechnol 36:996–1004.

36. Schiff SL, Tsuji JM, Wu L, Venkiteswaran JJ, Molot LA, Elgood RJ, Paterson MJ, Neufeld JD. 2017. Millions of boreal shield lakes can be used to probe Archaean Ocean biogeochemistry. Sci Rep 7:46708.

37. Krumov N, Perner-Nochta I, Oder S, Gotcheva V, Angelov A, Posten C. 2009. Production of inorganic nanoparticles by microorganisms. Chem Eng Technol 32:1026–1035.

38. Leão P, Chen Y-R, Abreu F, Wang M, Zhang W-J, Zhou K, Xiao T, Wu L-F, Lins U. 2017. Ultrastructure of ellipsoidal magnetotactic multicellular prokaryotes depicts their complex assemblage and cellular polarity in the context of magnetotaxis. Environ Microbiol 19:2151–2163.

39. Lefèvre CT, Trubitsyn D, Abreu F, Kolinko S, Jogler C, de Almeida LGP, de Vasconcelos ATR, Kube M, Reinhardt R, Lins U, Pignol D, Schüler D, Bazylinski DA, Ginet N. 2013. Comparative genomic analysis of magnetotactic bacteria from the *Deltaproteobacteria* provides new insights into magnetite and greigite magnetosome genes required for magnetotaxis. Environ Microbiol 15:2712–2735.

40. Lin W, Deng A, Wang Z, Li Y, Wen T, Wu L-F, Wu M, Pan Y. 2014. Genomic insights into the uncultured genus "*Candidatus* Magnetobacterium" in the phylum *Nitrospirae*. ISME J 8:2463–77.

41. Imhoff JF. 2016. New dimensions in microbial ecology — functional genes in

studies to unravel the biodiversity and role of functional microbial groups in the

environment. Microorganisms 1–41.

42.　Petri R, Imhoff JF. 2020. The relationship of nitrate reducing bacteria on the

basis of *narH* gene sequences and comparison of *narH* and 16S rONA based

phylogeny. Syst Appl Microbiol 23:47–57.

43.　Meyer B, Kuever J. 2007. Molecular analysis of the diversity of sulfate-

reducing and sulfur-oxidizing prokaryotes in the environment , using *aprA* as

functional marker gene. Appl Environ Microbiol 73:7664–7679.

44.　Petri R, Podgorsek L, Imho JF. 2001. Phylogeny and distribution of the *sox B*

gene among thiosulfate-oxidizing bacteria. FEMS Microbiol Lett 197:171–178.

45.　Junier P, Kim O, Junier T. 2009. Community analysis of betaproteobacterial

ammonia-oxidizing bacteria using the *amoCAB* operon. Appl Microbiol

Biotechnol 175–188.

46.　Vetriani C, Voordeckers JW, Crespo-medina M, Brien CEO, Giovannelli D,

Lutz RA. 2014. Deep-sea hydrothermal vent *Epsilonproteobacteria* encode a

conserved and widespread nitrate reduction pathway (Nap). ISME J 1510–

1521.

47.　Tank M, Thiel V, Imhoff JF. 2009. Phylogenetic relationship of phototrophic

purple sulfur bacteria according to *pufL* and *pufM* genes. Int Microbiol 175–

185.

48.　Gaisin VA, Kalashnikov AM, Sukhacheva M V., Namsaraev ZB, Barhutova

DD, Gorlenko VM, Kuznetsov BB. 2015. Filamentous anoxygenic phototrophic

bacteria from cyanobacterial mats of Alla hot springs (Barguzin Valley,

Russia). Extremophiles 19:1067–1076.

49. Thiel V, Tank M, Neulinger SC, Gehrmann L, Dorador C, Imhoff JF. 2010.

Unique communities of anoxygenic phototrophic bacteria in saline lakes of

Salar de Atacama (Chile): evidence for a new phylogenetic lineage of

phototrophic *Gammaproteobacteria* from *pufLM* gene analyses. FEMS

Microbiol Ecol 74:510–522.

50. Francis CA, Roberts KJ, Beman JM, Santoro AE, Oakley BB. 2005. Ubiquity

and diversity of ammonia-oxidizing archaea in water columns and sediments of

the ocean. Proc Natl Acad Sci 102:14683–14688.

51. Lin W, Jogler C, Schüler D, Pan Y. 2011. Metagenomic analysis reveals

unexpected subgenomic diversity of magnetotactic bacteria within the phylum

Nitrospirae. Appl Environ Microbiol 77:323–326.

52. Li J, Menguy N, Gatel C, Boureau V, Snoeck E, Patriarche G, Leroy E, Pan Y,

Menguy N. 2015. Crystal growth of bullet-shaped magnetite in magnetotactic

bacteria of the *Nitrospirae* phylum. J R Soc Interface 12(103):20141288.

53. Lefèvre CT, Pósfai M, Abreu F, Lins U, Frankel RB, Bazylinski DA. 2011.

Morphological features of elongated-anisotropic magnetosome crystals in

magnetotactic bacteria of the *Nitrospirae* phylum and the *Deltaproteobacteria*

class. Earth Planet Sci Lett 312:194–200.

54. Lin W, Paterson GA, Zhu Q, Wang Y, Kopylova E, Li Y, Knight R, Bazylinski

DA, Zhu R, Kirschvink JL, Pan Y. 2017. Origin of microbial biomineralization

and magnetotaxis during the Archean. Proc Natl Acad Sci 114:2171–2176.

55. Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W : improving the

sensitivity of progressive multiple sequence alignment through sequence

weighting , position-specific gap penalties and weight matrix choice. Nucleic

Acids Res 22:4673–4680.

56. Gorlenko VM, Dzyuba M V., Maleeva AN, Panteleeva AN, Kolganova T V.,

Kuznetsov BB. 2011. *Magnetospirillum aberrantis* sp. nov., a new freshwater

bacterium with magnetic inclusions. Microbiology 80:692–702.

57. Lane DJ. 1991. 16S/23S rRNA sequencing In: Stackebrandt E, Goodfellow M

(eds) Nucleic acid techniques in bacterial systematics. Wiley, New York 115–

175.

58. Sambrook J, Fritsch EF, Maniatis T. 1989. Molecular cloning: a laboratory

manual. Cold Spring Harbor Laboratory Press, 2nd ed. New York.

59. Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software

version 7: improvements in performance and usability. Mol Biol Evol 30:772–

80.

60. Castresana J. 2000. Selection of conserved blocks from multiple alignments for

their use in phylogenetic analysis. Mol Biol Evol 17:540–552.

61. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2014. IQ-TREE: a fast

and effective stochastic algorithm for estimating maximum-likelihood

phylogenies. Mol Biol Evol 32:268–274.

62. Wong TKF, Jermiin LS, Minh BQ, Kalyaanamoorthy S, von Haeseler A. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat

Methods 14:587–589.

63. Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2017.

UFBoot2: improving the ultrafast bootstrap approximation. Mol Biol Evol

35:518–522.

64. Pernthaler J, Glöckner F-O, Schönhuber W, Amann R. 2001. Fluorescence in

situ hybridization (FISH) with rRNA-targeted oligonucleotide probes. Methods

Microbiol 30:207–226.

65. Daims H, Brühl A, Amann R, Schleifer K-H, Wagner M. 1999. The domain-

specific probe EUB338 is insufficient for the detection of all Bacteria:

development and evaluation of a more comprehensive probe set. Syst Appl

Microbiol 22:434–444.

66. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for

Illumina sequence data. Bioinformatics 30:2114–2120.

67. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS,

Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin A V., Sirotkin A V.,

Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: A New

Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J

Comput Biol 19:455–477.

68. Tatusova T, Dicuccio M, Badretdin A, Chetvernin V, Nawrocki P, Zaslavsky L,

Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic

genome annotation pipeline. Nucleic Acids Res 44:6614–6624.

69. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: Quality

assessment tool for genome assemblies. Bioinformatics 29:1072–1075.

70. Wu Y-W, Tang Y-H, Tringe SG, Simmons BA, Singer SW. 2014. MaxBin: an

automated binning method to recover individual genomes from metagenomes

using an expectation-maximization algorithm. Microbiome 2:26.

71. Lin H-H, Liao Y-C. 2016. Accurate binning of metagenomic contigs via

automated clustering sequences using information of genomic signatures and

marker genes. Sci Rep 6:24175.

72. Laczny CC, Kiefer C, Galata V, Fehlmann T, Backes C, Keller A. 2017.

BusyBee Web : metagenomic data analysis by bootstrapped supervised binning

and annotation. Nucleic Acids Res 45:W171–W179.

73. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, Banfield

JF. 2018. Recovery of genomes from metagenomes via a dereplication,

aggregation and scoring strategy. Nat Microbiol 3:836-843.

74. Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN,

Hugenholtz P, Tyson GW. 2017. Recovery of nearly 8,000 metagenome-

assembled genomes substantially expands the tree of life. Nat Microbiol

2:1533-1542.

75. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015.

CheckM : assessing the quality of microbial genomes recovered from isolates ,

single cells , and metagenomes. Genome Res 25:1043–1055.

Figure legends

Figure 1. Evaluation of nested PCR for MTB analyses. **a)** PCR amplicons obtained

with the primers mamK_86F-mamK_521R after nested PCR on MTB and non-MTB

isolates and the entire microbial community; **b)** PCR amplicons with16S rRNA used

as a control DNA template. Lane 1: control sample without DNA; Lane 2: *E. coli*

DH10B; Lane 3: *Magnetospirillum aberranis* SpK; Lane 4: *M. moscoviense* BB-1*;*

Lane 5: *M. kuznetsovii* LBB-42; Lane 6: *M. caucaseum* SO-1; Lane 7:

Magnetococcales bacterium UR-1; Lane 8: Moskva River; Lane 9: Soda Lake Kiran;

Lane M: DNA ladder GeneRuler 100 bp Plus.

Figure 2. Maximum-likelihood phylogenetic trees based on **a)** magnetosome

associated protein MamK (285 amino acid sites) reconstructed with evolutionary

model LG+I+G4; and **b)** 16S rRNA sequences (1,399 nucleotide sites) reconstructed

using the evolutionary model GTR+F+I+G4.

Figure 3. FISH-TEM of the MTB '*Ca*. Magnetomonas plexicatena' LBB_01 and

HRTEM of its magnetosomes. **a)** Phase contrast image of magnetically enriched

environmental sample deposited on Formvar- coated TEM grids; **b)** Bacteria

observed following hybridization with a EUB probe; **c)** Image after hybridization

with a probe specific for *Ca*. Magnetomonas plexicatena species. The arrow

highlights a small coccus that was not hybridized with the specific probe; **d)** TEM

image of the same region marked as a box in panel **a**; **e)** Higher magnification of the

highlighted area in **d** showing a chain of anisotropic magnetosomes organized along

the long axis of the cell; **f)** Higher magnetic activity of the area inside the square in

figure **e**. Magnetosomes in this cell had occasional rough edges and a bullet shape; **g)**

HRTEM of the magnetosome indicated by an asterisk in **f**. FFT of the magnetosome

crystalline structure support the possibility that these magnetite magnetosomes align

along an [111] elongation axis.

Figure 4. FISH-TEM image of MTB from environmental samples. The circles and

arrows indicate cells that did and did not, respectively, hybridize with the specific

probe tested. **a**), **f**) and **k**) Phase contrast image of magnetically enriched

environmental samples on Formvar-coated TEM grids; **b**), **g**) and **l**) Bacteria present

in the same region as imaged in **a**, **f** and **k**, respectively, stained with DAPI; **c**), **h**)

and **m**) Image of the same area as shown in **b**, **g** and **l**, respectively, after

hybridization of bacterial cells with the EUB probe; **d**) Image of the same area

captured in image **c** after hybridization with the LBB_03 probe; **e**) TEM image of the

same cell that showed hybridization with the LBB_03 probe in **d**. The cell has a rod

shape and anisotropic magnetosomes; **i**) Image of the same area captured in **h** after

cell hybridization with probe LBB_02; **j**) TEM image of the MTB that showed

hybridization with probe LBB_02 in image **i**. The MTB is a small coccus having two

chains of anisotropic magnetosomes; **n**) TEM image of the same area represented in

image **m** after hybridization with probe LBB_04. This cell had many disorganized

anisotropic crystals in the cytoplasm.

Figure 5. Maximum-likelihood phylogenetic tree derived from concatenated core

protein sequences. Phylogenetic analysis was performed with a LG+F+I+G4 model

based on 34,747 amino acid positions. The scale bar represents amino acid

substitutions per site.

Figure 6. MTB-CoSe separation procedure that allowed collection of magnetotactic

components directly from sediment samples.

A

Nitrospirae

Delta-

Alpha-

'Eta'-

B

A **Ph**
B **EUB**
C **LBB_01**

10μm

D 5 μm

E 500 nm

F 50 nm
*

G
111 110

ZA [1 1̄ 0]

10 nm