Title: Dual SMAD inhibition and Wnt inhibition enhances the differentiation of induced pluripotent stem cells into Retinal Ganglion cells (iPSC-RGCs)

Venkata R. M. Chavali*[1], Naqi Haider[1], Sonika Rathi[1], Teja Alapati[1], Jie He, Kamaljot Gill[2], Sergei Nikonov[2], Roman Nikonov[2], Thu T. Duong[2], Devin S. McDougald[2], Joan O' Brien[1] and Jason A. Mills[2]*

Affiliation:

[1]Scheie Eye Institute, University of Pennsylvania Perelman School of Medicine, Pennsylvania, PA, 19104, USA; [2]F.M. Kirby Center for Molecular Ophthalmology and Center for Advanced Retinal and Ocular Therapeutics (CAROT),

*Corresponding authors: Venkata R. M. Chavali, Scheie Eye Institute, University of Pennsylvania, 133A Anatomy Chemistry Building, 3620 Hamilton Walk, Philadelphia, Pennsylvania, USA, 19104. Ph: 215-898-8320, Email: vchavali@pennmedicine.upenn.edu and
Jason A. Mills, F.M. Kirby Center for Molecular Ophthalmology and Center for Advanced Retinal and Ocular Therapeutics (CAROT), Scheie Eye Institute, University of Pennsylvania, Perelman School of Medicine, 309B Stellar Chance Laboratories, Philadelphia, Pennsylvania, USA, 19104. Telephone: 215-746-3482. Email: millsja@pennmedicine.upenn.edu.

Running Title:
Keywords: Pluripotent stem cells, retinal ganglion cells, differentiation

Abstract:
Glaucoma is a group of progressive optic neuropathies that share common biological and clinical characteristics including irreversible changes to the optic nerve and visual field loss caused by death of retinal ganglion cells (RGCs). The loss of RGCs manifests as characteristic cupping or optic nerve degeneration, resulting in peripheral vision loss in patients with Glaucoma. Several initial studies carried out RGC differentiation from induced pluripotent stem cells (iPSCs) or embryonic stem cells (ESCs), by modulating classical RGC signaling pathways to mimic in vivo retinal development. Recent studies used small molecules and peptide modulators at specific intervals to fine tune RGC differentiation, and also increase the yields to produce purified RGCs. In this study, we present a chemically defined and functionally novel *in vitro* methodology for cultivating unprecedented yields of RPC populations from iPSCs, that are then directed toward the RGC lineage. Using this method, we differentiated control iPSC lines into RGCs using in in a stepwise manner using small molecules and peptide modulator treatment to inhibit BMP and TGF-β (SMAD), and canonical Wnt pathways yielding a robust population of iPSC-RGCs.

Introduction:

Glaucoma encompasses a heterogenous group of optic neuropathies, tied together by their exhibition of permanent damage to the optic nerve.[1] This degeneration is due to the death of retinal ganglion cells (RGCs), with subsequent peripheral vision loss. [2]Primary open-angle glaucoma (POAG), the most common form of glaucoma, is characterized by chronic and progressive optic nerve degeneration and corresponding visual field deficits in the presence of an open and normal iridocorneal chamber angle.[3] This disease is the leading cause of irreversible blindness worldwide [4], with an estimated 11.1 million expected to become blind from POAG by 2020.[5] It is also the leading cause of vision loss among African Americans.[6] Despite the prevalence of POAG, its pathogenesis remains poorly understood. Elevated intraocular pressure (IOP) is a major risk factor [7], but degeneration of the optic nerve and loss of RGCs may still occur in those with normal eye pressure.[6]   The complexity of glaucoma certainly makes it possible, if not probable, that RGCs become inherently susceptible to this disease process.

RGCs are found in the ganglion cell layer of the retina and serve as the projection neurons of the retina, utilizing long axons to effectively connect the eye to the brain. They transmit both image-forming and non-image-forming visual information, processed by retinal cells such as photoreceptors and horizontal cells, to higher visual centers in the lateral geniculate body through the optic nerve.[8] Currently, no promising treatment exists for glaucoma or RGC degeneration. Although shown to be effective in animal models of glaucoma, neuroprotective approaches have not proven practical in human settings.[9, 10] As mature mammalian RGCs are a terminally differentiated lineage, they do not regenerate after succumbing to disease[11], consequently leading to the irreparable blindness. Understandably, there is a great desire for an application devised to rejuvenate or replace injured RGCs. Hence, it is with good reason that translating

human stem cell technology to a viable regenerative therapy for use in degenerative conditions has become a worldwide priority.

Human pluripotent stem cells (hPSCs) offer a tantalizing value in medicine - they can be guided to differentiate into practically any type of cell within an organism. Owing to a variety of techniques that are established to derive every key retinal cell lineage, blinding disorders of the retina can be particularly well-modeled with hPSCs.[12, 13] Though the field was previously reliant on embryonic stem cells (ESCs), advances in stem cell research led to the discovery that overexpressing genes associated with "stemness" in somatic cells can reprogram them into induced pluripotent stem cells (iPSCs).[14, 15] The advent of iPSC manipulation has opened up opportunities for implementing this technology to cure diseases stemming from retinal cells.[16] Taking hPSCs and differentiating them toward RGC lineage commitment may generate enough healthy cells to compensate for the degenerative RGCs in glaucoma. As a result, stem-cell based therapy holds promise as a method to restore vision in retinal degeneration; however, success of these treatment strategies hinges on de novo synthesis of RGCs with stable phenotypes from hPSCs.[17, 18]

Several methods for differentiating ESCs and iPSCs into RGCs have been reported. Of the retinal lineages, RGC differentiation is perhaps the most complicated, historically involving three-dimensional (3D) culture of hPSCs.[19, 20] This enables the primordial eye to self-organize into 3D cell aggregates such as embryoid bodies (EBs), followed by optic vesicles (OVs) and later, optic cups, all of which are analogous to structures developed in vivo.[20, 21] The retinal progenitor cells (RPCs) within eventually mature into RGCs, characteristically the first neuronal subtype to develop, as well as other retinal cell types.[22, 23] While most of the lineages of the retina do arise in a properly organized manner, therein lies the problem with the earlier RGC generation protocols: instead of predominantly RGCs, majority of neural retinal cell types are present.[24-26] This heterogeneity and inefficient yield in culture restricts selection of specifically RGCs without an enrichment step, even if the 3D platform phase were to precede an adherent culture.[24, 27] To better specify the RGC cell lineage and ideally obviate 3D culture, latter studies modulated RGC signaling pathways, including Hedgehog and Notch, using various ligands and other factors.[26, 28] While this did enhance efficiency, there still resulted a suboptimal RGC yield and ambiguous picture of which pathways to best modulate, due to the ill-defined addition and administration of components. This shortcoming has been improved upon by administration of small molecules and peptide modulators at appropriate intervals in a calculated manner to more precisely modulate cellular signals and pathways such as bone morphogenetic protein (BMP), Wnt, IGF1, and TGF-β.[29] Canonical Wnt, BMP, and nodal are the three signaling pathways classically inhibited in vitro to spur pluripotent stem cells toward the neural retinal fate.[30-32] Once RPCs form the neural retina, the basic helix-loop-helix transcription factor (TF) ATOH7 plays a central role in RGC commitment and expression regulation of RGC-specific markers BRN3+ and ISL1.[27, 33, 34] Regardless of the protocol, adequate differentiation relies on emulating the pathways present in generating these cell lineages in vivo.

Developing a therapeutic approach employing stem cell technology requires an efficient, reproducible, and safe differentiation protocol for RGCs. To this end, we detail a chemically defined and functionally novel in vitro technique for cultivating unprecedented yields of RPC populations from iPSCs that are then directed toward the RGC lineage. True to their in vivo development, we directed unaffected control lines of iPSCs to differentiate in a stepwise manner using a temporal induction method centered on small molecule and peptide modulator treatment to inhibit BMP and TGF-β (SMAD), and canonical Wnt yielding a robust population of iPSC-RGCs after Day 30. Analysis at the molecular and physiologic levels using flow cytometry, immunolabeling, gene expression, and electrophysiology yielded results in accordance with the RGC lineage and allowed specific subtype identification.[26, 28, 35]

Methods:

Human induced pluripotent stem cell culture

Undifferentiated IPS cells (Control 1 (CHOPWT8). Control 2 (CHOPWT9), and Control 3 (CHOPWT10) were derived and characterized as previously published showing a complete analysis of iPSC characteristics. [36-38] All human samples were collected in accordance with institutional review board (IRB) requirements at the Children's Hospital of Philadelphia with informed consent. IPSc cells were maintained in IPSC medium ((Dulbecco's modified essential medium/ Ham's F12 nutrient media; DMEM/F12 (50:50; corning)) containing 1 x glutatmax (glut, invitrogen), 1 x penicillin/streptomycin (PS, invitrogen), 15% Knockout serum replacement (KSR; invitrogen), 1 x non-essential amino acids (NEAA; invitrogen), 0.1mM β-Mercaptoethanol (2-ME; invitrogen), and 10 ng/mL of basic fibroblast growth factor (bFGF; R&D Systems) on 0.1% gelatin coated dishes with irradiated mouse embryonic fibroblast (iMEFs).

Retinal Progenitor Cells (RPCs) generation

iPSCs were cultured on 0.1% gelatinized plates containing iMEFs in 37°C 5% O_2 and 5% CO_2 conditions. Cells were maintained until approximately 70-75% confluence, then feeder cells were depleted and approximately 1.5 - 1.8 x 10^6 iPSCs were seeded one well of 6-well (9.8 cm^2) tissue culture dish (Corning) coated with 1:100 diluted growth factor reduced (GFR) Matrigel in iPSC media containing Thiazovivin (2 mM, Tocris). iPSCs were maintained in iPSC: MEF-conditioned medium (80:20) + 20 ng/mL of bFGF. iPSCs were maintained in 37C 5% O_2 and 5% CO_2 until reaching 100% confluence then transferred to in 37°C, 5% CO_2 overnight prior to induction. All induction media changes were performed on a daily basis. On the first day, iPSC cells were changed to RPC induction media 1 (RPC1) (DMEM/F12 (50:50), 1 x P/S, 1 x Glut, 2% B27 supplement (w/o vitamin A), 1% N2 supplement containing 2 μM XAV939 (X) (Wnt inhibitor; SelleckChem, 10 mM SB431542 (SB) (TGF-Beta inhibitor; SelleckChem), 100 nM LDN193189 (L) (BMP inhibitor; Stemgent), 10 mM nicotinamide (Sigma-Aldrich), 10 ng/mL IGF1 (R&D Systems), cultures were fed for 4 days. On day 5, cultures were fed with retinal progenitor media 2 (RPC2) (DMEM/F12 (50:50), 1 x P/S, 1 x Glut, 2 x B27 supplement (w/o vitamin A), 1 x N2 supplement containing 2 μM XAV939 (Wnt inhibitor,

SelleckChem, 10 mM SB431542 (TGF-Beta inhibitor, SelleckChem), 100 nM LDN193189 (BMP inhibitor, Stemgent), 10 ng/mL IGF1 (R&D Systems), and 10 ng/mL bFGF (R&D Systems) with media changes daily for 12 days.

Retinal ganglion cell (RGC) differentiation conditions.

On day 16 of differentiation, four conditions were evaluated for RGC generation. The base media included DMEM/F12 (50:50),1 x P/S, 1 x Glut, 2 x B27 supplement (w/o vitamin A) and 1 x N2 supplement. For experimental conditions 1) LSBX, 2) LSB, 3) LX, and 4) LSB-CHIR (CHIR, Wnt agonist) were evaluated. Cultures were treated for 5 days prior to RGC patterning using SHH (100 ng/mL; R&D systems) or SAG (100 nM; Tocris), SHH was replaced with SAG after establishing equal competence, FGF8 (100 ng/mL, R&D systems), DAPT (3-10 mM Tocris), Y27632 (10 mM, Tocris), Follistatin 300 (100 ng/mL; R&D systems), cyclopamine (0.5 mM; Tocris) BDNF (50 ng/mL; R&D systems), NT4 (5 ng/mL; R&D systems), CNTF (5 ng/mL; R&D systems), forskolin (25 mM; SelleckChem), and cAMP (400 mM; Sigma-Aldrich) for RGC patterning. see figure 1A and 2A for kinetics. Cells were replated at a density of 1.0×10^5 cells/well of 6-well using the crosshatching technique as previously described (Espuny-Camacho et al. 2013) in RPCs base media containing Follistatin 300, Cyclo, DAPT, and 10 mM Rock inhibitor (Y27632, Tocris) on day 23 and culture media changed 24h post-plating with RPC2 media containing Follistatin 300 and DAPT. After maturation, medium was exchanged every 3-4 days with neural basal media supplemented with 2% B27 without vitamin-A and 1 x Glutamax.

Flow cytometry analysis of RGCs

Culture of RGCs were collected using TrypLE (Invitrogen) and run through a 100 ☐m filter to ensure single cell suspension. A portion of cells were incubated with antibodies anti-CD90-PE-cy7 (Thy1; Biolegend 1:100) and anti-SSEA4-APC (Biolegend 1:100) in 1 x PBS supplemented with 0.5% bovine serum albumin (BSA) +0.1% Na Azide (FACS buffer) and incubated at RT for 30 minutes on ice as previously described.[41] We analyzed stained cells AccuriC6 on and data analyzed using FlowJo Version 10.0.8 software (TreeStar).

Immunohistochemistry

Immunohistochemistry analysis was performed as previously published (Duong et al. 2018). For immunolocalization of antigens, RGCs on coverslips were fixed and thoroughly washed with phosphate-buffered saline, cryoprotected in 30% sucrose and embedded in OCT tissue freezing medium before sectioning. For characterizing RGCs immunohistochemistry was performed using SOX2, PAX6, RAX, Tuj1, Thy1/CD90 antibodies and dilution as described earlier.[38-41] Sections were mounted in mounting media containing DAPI (4',6-diamidino-2-phenylindole), observed under an Olympus FV1000 Confocal microscope, and images were captured with the use of appropriate filters and lasers.

Voltage recordings

The circular cover glass coated with 1:100 matrigel containing adherent cells was transferred to the recording chamber filled with iNS maintenance medium. The camber was placed on the microscope stage (Olympus BX-61 upright microscope) and perfused with the oxygenated Ames's solution (Sigma-Aldrich) resulting in gradual replacement of the growth medium with the Ames medium. The temperature of the solution inside the chamber was gradually increased to 37°C prior to the beginning of the imaging and recording session using Warner Instruments TC 344B temperature controller. Confocal images of the cells were acquired with Olympus Fluoview 1000 MPE system and cells demonstrating strong GFP expression were selected for patch-clamp recording. Whole cell configuration was achieved in the voltage-clamp mode at -60 mV holding potential. After achieving whole-cell configuration cells were either maintained in the current clamp mode at zero holding current and depolarized with calibrated current steps, or in the voltage-clamp mode at -60 mV holding potential from which level they were depolarized with calibrated voltage steps. Warner Instruments PC 505B amplifier and Molecular Devices Digidata 1440 digitizer under the control of the Clampex software were used for controlling and recording of the membrane voltages and currents. Patch pipettes were pulled from 1.2/0.69 mm OD/ID borosilicate glass on Sutter Instruments P-87 puller. Pipette filling solution included (in mM) 110 K-gluconate, 12 NaCl, 10 HEPES, 1 EGTA. Pipette were mounted on Sutter Instruments MPC-200 micromanipulators, MTI CCD 72 camera system was used to provide video control over pipette and cell positioning in the chamber.

Quantitative Real-time PCR (qRT-PCR)

Total RNA was extracted using Purelink RNA isolation kit (invitrogen), with DNAse treatment. For each sample, 1 ug of total RNA was reverse transcribed using the high-capacity cDNA reverse transcription kit using random primers (ThermoFisher). Amplified material was detected using Taqman probes (target-FAM and housekeeping-TET) and AmpliTaq Gold Master mix (ThermoFisher) on a 7500 Fast Real-time PCR system (Applied Biosystems). All results were normalized to a B2M control (housekeeping) and are from 3 technical replicated of 3 independent biological samples for each time-point and experimental condition.

Statistical Analysis
Quantitative data were obtained from three independent experiments per cell line in triplicate. Statistical analysis was performed with Student T-test in Prism. *p-values of <0.05 were considered statistically significant.

RESULTS:

Combined inhibition of WNT, BMP and TGF-β and nicotinamide contribute to efficient generation of early retinal progenitor cells.

The hiPSC retinal differentiation methods differ between two- and three-dimensional culture conditions, and the use of proteins, and small molecules to recapitulate processes responsible for vertebrate retina development [28, 42]. These methods require a mid-differentiation enrichment step that involved the establishment of retinal rosettes followed by manual isolation and maturation to generate retinal subtypes. We have developed and characterized a modified two-stage protocol that differentiates hiPSCs into an enriched population of retinal progenitor cell (RPC) cultures followed by targeted patterning to retinal ganglion cells (RGCs) [28]. In order to accomplish this, hiPSCs were grown to confluence and subsequently treated with a retinal induction media containing, DMEM/F12 plus N2 and B27, XAV939 (WNT inhibitor), SB431542, (TGF-β inhibitor), LDN193189, (BMP inhibitor), nicotinamide, and IGF1 for 4 day (Fig. 1A). The inhibition of Wnt and BMP signaling has been documented to enhanced the expression of eye field transcription factors (EFTFs) during retinal differentiations of hPSCs [29]. We observed that addition of TGF-beta inhibition induced greater EFTFs expression during early retinal patterning. Nicotinamide was added to the differentiation media (d0-4) to promote the expression of early eye field markers LHX2 and RAX, as previously published [43]. Nicotinamide has been shown to promote cell expansion and adaptation to a radial/rosette morphology.[44] Patterning factors such as insulin like growth factor-1 (IGF-1) and basic growth factor-2 (bFGF2) aid in the specification of eye field identity to differentiating retinal progenitors.[29] From day 5-10, nicotinamide was removed and bFGF was added to retinal induction media. Analysis at day 7, showed an enriched population of SOX2, RAX and PAX6 positive cells (Fig. 1B). The expression of early retinal progenitor markers, LHX2 and RAX, were identified in over 97% of day 7 cultures (Fig.1 C) indicating and efficient generation of RPCs. Quantification of EFTFs, LHX2, RAX, PAX6, SIX6, SIX3, and VSX2, showed significant expression relative to pluripotent stem cells in all three individuals.

Extended exposures of BMP, WNT, and TGF-β inhibition leads to efficient generation of retinal ganglion cell neurons.

The efficient induction of retinal progenitors is largely dependent on the inhibition of pathways responsible for patterning of the diencephalon; however, how modification on of these inhibitory pathways play a role on retinal ganglion cells generation is unclear. Here, we tested whether pharmacological inhibition of WNT, TGF-β, and BMP can improve RGC induction and subsequently enable the controlled generation of mature iPSC-RGC neurons. We evaluated four different treatment modalities to determine pathways that enhance or restrict RGC generation from hiPSCs. Among the conditions tested: 1) BMP (L), TGB-B (SB), and Wnt inhibition (X) (LSBX), 2) LSB, 3) LX, and 4) LSB-CHIR (CHIR, Wnt agonist) (Fig. 1A). The progression of the differentiation proceeded through three phases; uniform RPC cultures (DIV7), rosette forming cultures (D16), then late stages neuronal cultures (D35) (Fig 2B). At day 21, cells were treated with RGC induction conditions which included the activation of sonic hedgehog, by augmenting with SHH (100 ug/mL). Comparative studies showed the use of SAG (100nM, smoothened agonist) could replace SHH with no difference in RGC induction changes. Fibroblast growth factor 8 (FGF8) and notch inhibitor (DAPT) were added along with SHH or SAG. At day 23, we SHH, TGF-β, and notch signaling were inhibited

to promote RGC maturation using cyclopamine, follistatin, and DAPT. RGC cultures expanded using a cross-hatching technique known separate cells into small clusters in RPC base media supplemented with rho-kinase inhibitor (Y-27632), follistatin, and DAPT to improve survival of RGCs during the expansion [45]. Neurotrophins, BDNF, neurotrophin 4 (NT4) and CNTF, were added to the media to promote the development and survival of cells with neuronal fate.[46] Forskolin and Y-27632 were included as a potent promoter of RGC neurite growth, survival and lineage commitment.[42] These conditions were established by combining previously described conditions used by Sluch et al., 2015) and Teotia et al., 2017 for hiPSC-RGCs.

After 35 days in vitro, we dissociated cultures and performed flow cytometry analysis of the single cell suspension obtained from three independent iPSC lines, and data showed that using the LSBX condition resulted in the highest proportion of RGC, with a range between 34-45% Thy-1 (CD90) positive iPSC-RGCs in our study (Fig. 2C-E). The LX, CHIR, and LSB conditions resulted in 26.7% (±3.2), 22.5% (±4.7), and 27.2% (±6.2), respectively. We observed consistent results across individuals and experiments (Fig. 2D). All the RGCs generated displayed small soma and elongated axonal processes, with Immunocytochemical analysis showing the expression of Thy-1/Tuj1, Brn3a and SNCG (Fig. 2A, 5).

Gene expression profiles during differentiation of iPSC to RPCs and RGCs:

During eye formation in vertebrates, cell intrinsic and extrinsic signals and/or transcription factors control the differentiation and fate determination of retinal cells. We evaluated the gene expression profile of three eye specific transcription factors (RAX, PAX6 and Six3) called eye field transcription factors (EFTFs) that are known to play a role in the anterior neural plate. The expression of Rx (encoded by RAX gene) is maximum in the RPC inhibited with L and SB when compared to the other conditions at DIV23. The detection of RAX expression at DIV35 RGCs was minimal suggesting a commitment to a more differentiated retinal fate in at the consequence of retinal progenitor cell expansion. PAX6 is expressed in the cornea, lens, ciliary body, and retina through development and plays a role in determining their cell fate. The expression of PAX6 gene transcript is expressed in all experimental conditions in RPCs and RGCs, however PAX6 is predominantly expressed at DIV 23 and 35 in the CHIR condition, which stimulated the expression of canonical Wnt signaling. This indicates that a prolonged stimulation of RPCs with Wnt restricts the differentiation potential and maintains more cells as multipotent progenitors.

Among the other gene expression transcripts analyzed, we observed an increase in SIX3 gene expression in both LX and LSBX conditions DIV23 days when compared to RPCs stimulated with LSB or CHIR. The expression of Six3 decreased in RGCs at DIV35 indicating that neurospecification is reaching completion at this stage.

The SOX11 transcript is heavily expressed in developing retina during embryonic stages [47]. It is required for the maintenance of hedgehog signaling and is critical for axonal growth, extension and driving adult neurogenesis [48-50]. The expression of SOX11 increases nearly two-fold in LSBX and LX condition when compared to LSB and

CHIR in RPCs at DIV23. The expression of SOX11 is significantly reduced in RGCs at DIV35. Gli3 has a dual function as a transcriptional activator and a repressor of the sonic hedgehog (Shh) pathway. GLI1 is a simple transcriptional activator encoded by a target gene of Shh signaling. We observed increased expression of Gli3 in RPCs at DIV23 when compared to RGCs at DIV35. Expression of both Gli3 and Gli1 genes show that sonic hedgehog signaling is important for development of RPCs.

Direction selective RGC (DS-RGC), that respond to motion of light in different directions can be identified by expression of specific molecular markers, such as CART, CDH6, and FSTL4, among other genes. [51-54] CDH6 mediates axon-target matching and promotes wiring specificity that does not lead to image formation. These data provide some of the first evidence that a specific classical cadherin can promote wiring specificity in the mammalian visual system. Cadherin mediated cell-cell adhesion ensures precise connectivity of neurons in the eye to target nuclei in the brain. Increased expression of CDH6 was observed in LSBX condition in DIV35 RGCs demonstrating their maturation towards increased specificity for axonal wiring between RGCs. The CARTPT transcript (cocaine- and amphetamine-regulated transcript) is expressed by a major subtype of RGCs, ooDSGCs. In our study, the expression of *CARTPT* is seen in RGCs matured only in LSB, CHIR and LSBX conditions at DIV35 indicating that these conditions develop specific subtypes of mature RGCs that are known to be markers for ON-OFF direction-selective RGCs.[55] FSTL4 is a gene expressed in ON DS-RGCs and is colocalized with BRN3B in few RGCs [56]. We have seen increased expression of FSTL4 gene transcript in CHIR induced RGCs when compared with other conditions at DIV35 indicating that development of various subtypes of mature RGCs in our culture conditions. We did not see any expression of CARTPT or FSTL4 transcripts in RPCs at DIV23.

The RGCs at DIV35 expressed Brn3a transcript predominantly in LSBX growth condition when compared to other conditions. Interestingly we did see ATOH7 expression early in RPC differentiation at DIV23 and lower expression in DIV35 RGCs. We detected low expression of CRX in DIV35 RGC cultures across all culture conditions implying that the differentiated RGCs using these conditions restrict the photoreceptor precursor cell population. We observed low expression of RCVRN (expressed by photoreceptors) and MITF in our cultures at DIV23 and 35 indicating that our cultures are differentiated predominantly towards RGC fate, with minimal retinal pigmented epithelium cell identity. We observed the development of few interneurons in mature RPCs (amacrine or horizontal cells) as evidenced by expression of CALB2 especially in RGCs with Wnt activation at DIV35.

Functional Analysis of iPSC-Retinal Ganglion Cells (iPSC-RGCs)

Confocal imaging of iPSC-RGC cultures at DIV35 cell show the presence of the GFP-expressing neurons with neurite projections, and an increase in the density and complexity of the projections over the period of one month (Fig. 5). Two out of the six cells tested demonstrated action potential firing in the current clamp mode in response to the depolarizing currents from patch pipettes. The responses of one of these cells

(*cell a*) are illustrated in Fig. 6. Although cells are not expected to fire action potentials under ideal voltage-clamp conditions, we believe that imperfect clamping in distal processes can lead to such firing we observed as illustrated on the Fig. 6 for the same *cell a* from Fig. 5. Interestingly, all but one cell (6 out of 6 and 4 out of 5 DIV35 old cells) demonstrated reliable firing when depolarized with voltage steps in the voltage-clamp mode. On Fig. 7 *cell b* illustrates firing of the older cells and *cell c* illustrates firing of the younger cells in response to the depolarizing voltage steps. As expected, later stage iPSC-RGCs cultures (DXX) produced higher frequency, sustained firing and generated larger spikes when compared to early born iPSC-RGCs.

Cells firing under current-clamp conditions had reasonably strong resting potential around -50 mV while all cells that fired under voltage-clamp but not current-clamp conditions had rather weak resting potentials around and above -30 mV. Maintaining these cells at zero-current level before and between depolarizing current steps in the current-clamp mode keeps membrane potential at its weak resting level, likely resulting in the weaker sodium/potassium gradient precluding cells from firing upon depolarization. Alternatively, in the voltage clamp mode membrane potential was maintained at -60 mV before and between depolarizing voltage steps likely resulting in recharging of ion gradients which enabled action potential firing in response to the depolarization. This may partly be the reason why some cells fired under voltage-clamp but not current-clamp conditions.

Discussion:

In the current study, we employed a two-step/stage differentiation to induce RGC differentiation from iPSCs by modification of existing methodologies.[28, 29, 56] The first stage involves differentiation of iPSCs to RPCs. The RPCs were matured in a stage specific manner using small molecules to modulate sonic hedgehog pathway, Wnt pathway and Notch signaling to produce abundant RGCs reliability, that stain positive for RGC markers and emulate action potential.

Historically generation of RGCs from hiPSCs used matrigel as an adherent support to culture cells obtained from Embryoid bodies (EB) or ESCs. A cocktail of factors supplemented with FGF2 and proneural supplements was later used to differentiation.[57] RGCs resulting from such differentiations were characterized for their expression of Brn3b, Neurofilament-200. Matrigel was successfully replaced as a matrix for RGC differentiations using several additive factors like Noggin, DKK-1, IGF-1 and FGF in follow-up protocols. [58] Teotia *et al*, 2017 successfully recapitulated the "Lamba protocol" by differentiating hiPSCs initially into neural rosettes and then to functional RGCs by recapitulating retinal developmental pathways using chemical regulators.[28] Generation of a CRISPR engineered iPSC line with a fluorescent RFP tag reporter in the Brn3B locus, greatly assisted in evaluation of pathways necessary for RGC differentiation and characterization. [59]

We developed a quick and efficient RGC generation protocol involve embryoid bodies or need retinal rosette production or enrichment to initiate RGC differentiation. In our

method, the entire pluripotent iPSC culture is differentiated into RPCs in 21 days using a chemically defined medium. We employ crosshatching technique to generate clumps of cells that undergo stage specific differentiation to produce functionally mature RGCs by DIV28. This is a must faster timeline considering other published methods to date. [24, 28, 59-61]. We used the a modified chemically defined media composition standardized by Teotia *et al*, 2017 [28] to take advantage of 15-day RGC differentiation and paired it with our novel RPC generation protocol to generate RGCs making our method unique. The RPCs thus generated showed immunoreactivity to RAX and LHX2 in the retinal precursor lines for over 97% of cells in iPSC culture indicating that our protocol committed cells towards RPC lineage. Using our method, we reliability differentiated six control iPSC lines and multiple clones from those lines (data not shown) to generate mature RGCs that stain positive for Thy1/Tuj1, Brn3a and SNCG markers. Using different chemical conditions, we were able to produce Thy1 positive cells upto/ in over 58% of the differentiated RGCs. We also generated other retinal cell types in addition to mature RGCs that express pan-retinal markers and appear to be amacrine and/or bipolar cells. We also did not find any gene transcripts produced by any cells that relates to RPE and photoreceptors.

Several published methods to generate RGCs from iPSC and ESCs in 2D cultures used stepwise differentiation protocols generated different types of RGCs based on morphological and functional characteristics [56]. The RGC subtypes were characterized by their distinct single-cell RNA sequencing profiles, morphological features and distinct expression profiles. [56] Based on the nature and type of physiological responses, several RGC subtypes like ON-, OFF- and alpha-RGCs were observed. Evaluation of the gene expression profile of transcripts relevant several RGC markers at DIV18 and DIV28 time point of RGC differentiation confirming the validity of our method. Futher, gene expression profiles revealed that in addition to the known RGC subtypes that express ATOH7, Brn3a, SNCG, we produced several specialized RGCs like direction selective RGCs (DS-RGCs) that demonstrate gene expression profiles for *CART*, *CDH6*, and *FSTL4*, among other genes.

A strategy to bypass embryoid body (EB) formation by adding exogenous factors and in presence of proneural supplements like B27 and N2 was developed by Reichman *et al.*, 2014.[62] The long-term cultures of retinal organoids (RO) when supplemented with B27 allowed the differentiation of all retinal cell types including RGCs.[63] The RGC survival in these ROs derived retinal cultures decreased in vitro as the culture time was extended as floating cultures were a hinderance to the normal RGC growth and induced apoptosis. Adherent cultures obtained by enzymatic dissociation of ROs greatly improved the survival of these RGC [64] Several protocols following this study, differentiated iPSCs to neural rosettes and embryoid bodies that were further differentiated into RGCs using chemically defined media. [28, 56, 64, 65] These methods have several shortcomings for large scale growth and culture of RGCs as, they not only introduce bias during manual selection of desired RPCs, but also introduce variability in the total numbers or RGCs produced resulting from RPC numbers used for differentiation. Hence to reduce variability and increase the efficiency of the RGC generation, we differentiate the entire iPSC culture vessel to RPCs, and use a

crosshatching technique that does not need manual selection/screening to produce abundant RPC clumps that can be readily differentiated into RGCs.

Methods to isolate and purify RGCs have improved significantly with protocols using magnetic-activated cell sorting (MACS) with magnetic beads coupled to CD90 antibody, fluorescent cell sorting (FACSs), or immunopanning demonstrated enrichment of Thy1 positive RGCs cells.[65, 66] Using CD90 Thy1 magnetic beads, we were able to purify our Thy1 positive RGCs to over 90% to generate enriched cultures with defined characteristics.

The maintenance of later stage RGC cultures were cultured initially in NeuralBasal Media containing 1 x N2, and 1 x B27; however, this resulted in significant overgrowth of proliferative retinal progenitors cells. The removal of N2 supplement reduced the number of dividing cells in our cultures, providing a culture with predominantly neurons. We have also used 1 x CultureOne supplement within the RGC maturation media, which also reduced the number of non-RGC cell growth in day 45 in culture and beyond. After day 35 in culture, the mature RGCs produce appropriate morphological and physiological features. We observed extensive neurite outgrowth, and migration of RGCs to form cell clusters with elongated axonal projections forming a compartmentalized space by pushing away other cell aggregates may be by secreting unknown paracrine secretions, which warrants further investigation (Supplementary Fig).[26] Furthermore, the RGCs differentiated using the LSBX condition exhibited a electrphysiological function, with the ability to conduct sodium and potassium through voltage-dependent channels and fire action potentials. When comparing action potentials obtained from RGCs at day 35, day 70 and day 100, we observed that RGCs were able to fire continuously producing larger spikes with higher frequency, as they mature and age in culture when compared to younger cells.

We were unable to estimate the proportions of different RGC subtypes generated using our protocol. A systematic analysis of RGCs by single-cell RNA sequencing at different maturation time points is currently underway in our laboratory to identify RGC subtype markers and any maturation markers in the RGCs. Our validated methodology can reliably harness iPSC technology as a renewable source of RPCs to efficiently produce highly pure populations of RGCs for in vitro studies of glaucoma and potential therapeutic modalities for incurable RGC-related diseases.

Acknowledgements:
Supported in part by the National Eye Institute of the National Institutes of Health under Award Number R21EY028273-01A1 (VRMC), R21EY027936-01A1 (JAM), and 1RO1EY023557-01 (JMO). BrightFocus Foundation grant (VRMC), Research to Prevent Blindness Unrestricted Grant Funds to Scheie Eye Institute (VRMC, JAM, JMO), F.M. Kirby Foundation, and The Paul and Evanina Bell Mackall Foundation Trust for their support The Center for Advanced Retinal and Ocular Therapeutics (CAROT), The Foundation for Fighting Blindness (FFB), Research to Prevent Blindness (RPB), the National Institute of Health vision core grant P30EY01583-26, National Institutes of

Health, Department of Health and Human Services, under eyeGENETM and contract Nos. HHSN260220700001C and HHSN263201200001C.

Author contributions:
VRMC proposed the project and developed the concept with JAM. VRMC, NH, SN, TTD, SR, JAM conceived the method. VRMC, NH, SR, SN, RN, KG, JH, TTD, JAM conducted the experiments. DM generated the SNCG-GFP construct. VRMC, JAM, TTD, SR, NH, and SN performed the data analysis. TA, VRMC and JAM wrote the manuscript. VRMC, JAM, TTD, SR, NH, SN, RN, JMO, JH and TA revised the manuscript. VRMC and JAM supervised this work. All authors read and approved the final manuscript.

Additional information:

Competing financial interests: The authors declare no competing financial interests.

References:

1.   Janssen SF, Gorgels TG, Ramdas WD, Klaver CC, van Duijn CM, Jansonius NM, Bergen AA: **The vast complexity of primary open angle glaucoma: disease genes, risks, molecular mechanisms and pathobiology**. *Progress in retinal and eye research* 2013, **37**:31-67.
2.   Tham YC, Li X, Wong TY, Quigley HA, Aung T, Cheng CY: **Global prevalence of glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-analysis**. *Ophthalmology* 2014, **121**(11):2081-2090.
3.   Weinreb RN, Leung CK, Crowston JG, Medeiros FA, Friedman DS, Wiggs JL, Martin KR: **Primary open-angle glaucoma**. *Nat Rev Dis Primers* 2016, **2**:16067.
4.   Kapetanakis VV, Chan MP, Foster PJ, Cook DG, Owen CG, Rudnicka AR: **Global variations and time trends in the prevalence of primary open angle glaucoma (POAG): a systematic review and meta-analysis**. *Br J Ophthalmol* 2016, **100**(1):86-93.
5.   Quigley HA, Broman AT: **The number of people with glaucoma worldwide in 2010 and 2020**. *Br J Ophthalmol* 2006, **90**(3):262-267.
6.   Kwon YH, Fingert JH, Kuehn MH, Alward WL: **Primary open-angle glaucoma**. *N Engl J Med* 2009, **360**(11):1113-1124.
7.   Liu Y, Allingham RR: **Major review: Molecular genetics of primary open-angle glaucoma**. *Exp Eye Res* 2017, **160**:62-84.
8.   Yu DY, Cringle SJ, Balaratnasingam C, Morgan WH, Yu PK, Su EN: **Retinal ganglion cells: Energetics, compartmentation, axonal transport, cytoskeletons and vulnerability**. *Prog Retin Eye Res* 2013, **36**:217-246.
9.   Dahlmann-Noor AH, Vijay S, Limb GA, Khaw PT: **Strategies for optic nerve rescue and regeneration in glaucoma and other optic neuropathies**. *Drug Discov Today* 2010, **15**(7-8):287-299.

10. Danesh-Meyer HV, Levin LA: **Neuroprotection: extrapolating from neurologic diseases to the eye**. *Am J Ophthalmol* 2009, **148**(2):186-191 e182.
11. Sluch VM, Zack DJ: **Stem cells, retinal ganglion cells and glaucoma**. *Dev Ophthalmol* 2014, **53**:111-121.
12. Meyer JS, Howden SE, Wallace KA, Verhoeven AD, Wright LS, Capowski EE, Pinilla I, Martin JM, Tian S, Stewart R *et al*: **Optic vesicle-like structures derived from human pluripotent stem cells facilitate a customized approach to retinal disease treatment**. *Stem Cells* 2011, **29**(8):1206-1218.
13. Singh R, Shen W, Kuai D, Martin JM, Guo X, Smith MA, Perez ET, Phillips MJ, Simonett JM, Wallace KA *et al*: **iPS cell modeling of Best disease: insights into the pathophysiology of an inherited macular degeneration**. *Hum Mol Genet* 2013, **22**(3):593-607.
14. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S: **Induction of pluripotent stem cells from adult human fibroblasts by defined factors**. *Cell* 2007, **131**(5):861-872.
15. Takahashi K, Yamanaka S: **Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors**. *Cell* 2006, **126**(4):663-676.
16. Chamling X, Sluch VM, Zack DJ: **The Potential of Human Stem Cells for the Study and Treatment of Glaucoma**. *Invest Ophthalmol Vis Sci* 2016, **57**(5):ORSFi1-6.
17. Tang Z, Zhang Y, Wang Y, Zhang D, Shen B, Luo M, Gu P: **Progress of stem/progenitor cell-based therapy for retinal degeneration**. *J Transl Med* 2017, **15**(1):99.
18. Wright LS, Phillips MJ, Pinilla I, Hei D, Gamm DM: **Induced pluripotent stem cells as custom therapeutics for retinal repair: progress and rationale**. *Exp Eye Res* 2014, **123**:161-172.
19. Eiraku M, Takata N, Ishibashi H, Kawada M, Sakakura E, Okuda S, Sekiguchi K, Adachi T, Sasai Y: **Self-organizing optic-cup morphogenesis in three-dimensional culture**. *Nature* 2011, **472**(7341):51-56.
20. Nakano T, Ando S, Takata N, Kawada M, Muguruma K, Sekiguchi K, Saito K, Yonemura S, Eiraku M, Sasai Y: **Self-formation of optic cups and storable stratified neural retina from human ESCs**. *Cell Stem Cell* 2012, **10**(6):771-785.
21. Gill KP, Hewitt AW, Davidson KC, Pebay A, Wong RC: **Methods of Retinal Ganglion Cell Differentiation From Pluripotent Stem Cells**. *Transl Vis Sci Technol* 2014, **3**(4):7.
22. Chuang JH, Yarmishyn AA, Hwang DK, Hsu CC, Wang ML, Yang YP, Chien KH, Chiou SH, Peng CH, Chen SJ: **Expression profiling of cell-intrinsic regulators in the process of differentiation of human iPSCs into retinal lineages**. *Stem Cell Res Ther* 2018, **9**(1):140.
23. Marquardt T, Gruss P: **Generating neuronal diversity in the retina: one for nearly all**. *Trends Neurosci* 2002, **25**(1):32-38.
24. Tanaka T, Yokoi T, Tamalu F, Watanabe S, Nishina S, Azuma N: **Generation of retinal ganglion cells with functional axons from human induced pluripotent stem cells**. *Sci Rep* 2015, **5**:8344.

25.   Maekawa Y, Onishi A, Matsushita K, Koide N, Mandai M, Suzuma K, Kitaoka T, Kuwahara A, Ozone C, Nakano T *et al*: **Optimized Culture System to Induce Neurite Outgrowth From Retinal Ganglion Cells in Three-Dimensional Retinal Aggregates Differentiated From Mouse and Human Embryonic Stem Cells**. *Curr Eye Res* 2016, **41**(4):558-568.

26.   Ohlemacher SK, Sridhar A, Xiao Y, Hochstetler AE, Sarfarazi M, Cummins TR, Meyer JS: **Stepwise Differentiation of Retinal Ganglion Cells from Human Pluripotent Stem Cells Enables Analysis of Glaucomatous Neurodegeneration**. *Stem Cells* 2016, **34**(6):1553-1562.

27.   Deng F, Chen M, Liu Y, Hu H, Xiong Y, Xu C, Liu Y, Li K, Zhuang J, Ge J: **Stage-specific differentiation of iPSCs toward retinal ganglion cell lineage**. *Mol Vis* 2016, **22**:536-547.

28.   Teotia P, Chopra DA, Dravid SM, Van Hook MJ, Qiu F, Morrison J, Rizzino A, Ahmad I: **Generation of Functional Human Retinal Ganglion Cells with Target Specificity from Pluripotent Stem Cells by Chemically Defined Recapitulation of Developmental Mechanism**. *Stem Cells* 2017, **35**(3):572-585.

29.   Lee J, Choi SH, Kim YB, Jun I, Sung JJ, Lee DR, Kim YI, Cho MS, Byeon SH, Kim DS *et al*: **Defined Conditions for Differentiation of Functional Retinal Ganglion Cells From Human Pluripotent Stem Cells**. *Invest Ophthalmol Vis Sci* 2018, **59**(8):3531-3542.

30.   Zhou L, Wang W, Liu Y, Fernandez de Castro J, Ezashi T, Telugu BP, Roberts RM, Kaplan HJ, Dean DC: **Differentiation of induced pluripotent stem cells of swine into rod photoreceptors and their integration into the retina**. *Stem Cells* 2011, **29**(6):972-980.

31.   Cavodeassi F, Carreira-Barbosa F, Young RM, Concha ML, Allende ML, Houart C, Tada M, Wilson SW: **Early stages of zebrafish eye formation require the coordinated activity of Wnt11, Fz5, and the Wnt/beta-catenin pathway**. *Neuron* 2005, **47**(1):43-56.

32.   Hirami Y, Osakada F, Takahashi K, Okita K, Yamanaka S, Ikeda H, Yoshimura N, Takahashi M: **Generation of retinal cells from mouse and human induced pluripotent stem cells**. *Neurosci Lett* 2009, **458**(3):126-131.

33.   Yang Z, Ding K, Pan L, Deng M, Gan L: **Math5 determines the competence state of retinal ganglion cell progenitors**. *Dev Biol* 2003, **264**(1):240-254.

34.   Sun Y, Dykes IM, Liang X, Eng SR, Evans SM, Turner EE: **A central role for Islet1 in sensory neuron development linking sensory and spinal gene regulatory programs**. *Nat Neurosci* 2008, **11**(11):1283-1293.

35.   Lipton SA, Tauck DL: **Voltage-dependent conductances of solitary ganglion cells dissociated from the rat retina**. *J Physiol* 1987, **385**:361-391.

36.   Sullivan SK, Mills JA, Koukouritaki SB, Vo KK, Lyde RB, Paluru P, Zhao G, Zhai L, Sullivan LM, Wang Y *et al*: **High-level transgene expression in induced pluripotent stem cell-derived megakaryocytes: correction of Glanzmann thrombasthenia**. *Blood* 2014, **123**(5):753-757.

37.   Duong TT, Vasireddy V, Ramachandran P, Herrera PS, Leo L, Merkel C, Bennett J, Mills JA: **Use of induced pluripotent stem cell models to probe the**

pathogenesis of Choroideremia and to develop a potential treatment. *Stem Cell Res* 2018, **27**:140-150.

38. Maguire JA, Gagne A, Mills JA, Gadue P, French DL: **Generation of human control iPS cell line CHOPWT9 from healthy adult peripheral blood mononuclear cells**. *Stem Cell Res* 2016, **16**(1):14-16.

39. Lyssenko NN, Haider N, Picataggi A, Cipollari E, Jiao W, Phillips MC, Rader DJ, Chavali VRM: **Directional ABCA1-mediated cholesterol efflux and apoB-lipoprotein secretion in the retinal pigment epithelium**. *J Lipid Res* 2018, **59**(10):1927-1939.

40. Chavali VR, Khan NW, Cukras CA, Bartsch DU, Jablonski MM, Ayyagari R: **A CTRP5 gene S163R mutation knock-in mouse model for late-onset retinal degeneration**. *Hum Mol Genet* 2011, **20**(10):2000-2014.

41. Chavali VR, Sommer JR, Petters RM, Ayyagari R: **Identification of a promoter for the human C1Q-tumor necrosis factor-related protein-5 gene associated with late-onset retinal degeneration**. *Invest Ophthalmol Vis Sci* 2010, **51**(11):5499-5507.

42. Sluch VM, Davis CH, Ranganathan V, Kerr JM, Krick K, Martin R, Berlinicke CA, Marsh-Armstrong N, Diamond JS, Mao HQ *et al*: **Differentiation of human ESCs to retinal ganglion cells using a CRISPR engineered reporter cell line**. *Sci Rep* 2015, **5**:16595.

43. Meng Y, Ren Z, Xu F, Zhou X, Song C, Wang VY, Liu W, Lu L, Thomson JA, Chen G: **Nicotinamide Promotes Cell Survival and Differentiation as Kinase Inhibitor in Human Pluripotent Stem Cells**. *Stem Cell Reports* 2018, **11**(6):1347-1356.

44. Buchholz DE, Pennington BO, Croze RH, Hinman CR, Coffey PJ, Clegg DO: **Rapid and efficient directed differentiation of human pluripotent stem cells into retinal pigmented epithelium**. *Stem Cells Transl Med* 2013, **2**(5):384-393.

45. Espuny-Camacho I, Michelsen KA, Gall D, Linaro D, Hasche A, Bonnefont J, Bali C, Orduz D, Bilheu A, Herpoel A *et al*: **Pyramidal neurons derived from human pluripotent stem cells integrate efficiently into mouse brain circuits in vivo**. *Neuron* 2013, **77**(3):440-456.

46. Proenca CC, Song M, Lee FS: **Differential effects of BDNF and neurotrophin 4 (NT4) on endocytic sorting of TrkB receptors**. *J Neurochem* 2016, **138**(3):397-406.

47. Jiang Y, Ding Q, Xie X, Libby RT, Lefebvre V, Gan L: **Transcription factors SOX4 and SOX11 function redundantly to regulate the development of mouse retinal ganglion cells**. *J Biol Chem* 2013, **288**(25):18429-18438.

48. Bergsland M, Werme M, Malewicz M, Perlmann T, Muhr J: **The establishment of neuronal properties is controlled by Sox4 and Sox11**. *Genes Dev* 2006, **20**(24):3475-3486.

49. Wen W, Pillai-Kastoori L, Wilson SG, Morris AC: **Sox4 regulates choroid fissure closure by limiting Hedgehog signaling during ocular morphogenesis**. *Dev Biol* 2015, **399**(1):139-153.

50. Pillai-Kastoori L, Wen W, Wilson SG, Strachan E, Lo-Castro A, Fichera M, Musumeci SA, Lehmann OJ, Morris AC: **Sox11 is required to maintain proper**

levels of Hedgehog signaling during vertebrate ocular morphogenesis.
PLoS Genet 2014, **10**(7):e1004491.

51. De la Huerta I, Kim IJ, Voinescu PE, Sanes JR: **Direction-selective retinal ganglion cells arise from molecularly specified multipotential progenitors**. *Proc Natl Acad Sci U S A* 2012, **109**(43):17663-17668.

52. Kay JN, De la Huerta I, Kim IJ, Zhang Y, Yamagata M, Chu MW, Meister M, Sanes JR: **Retinal ganglion cells with distinct directional preferences differ in molecular identity, structure, and central projections**. *J Neurosci* 2011, **31**(21):7753-7762.

53. Huberman AD, Manu M, Koch SM, Susman MW, Lutz AB, Ullian EM, Baccus SA, Barres BA: **Architecture and activity-mediated refinement of axonal projections from a mosaic of genetically identified retinal ganglion cells**. *Neuron* 2008, **59**(3):425-438.

54. Yonehara K, Shintani T, Suzuki R, Sakuta H, Takeuchi Y, Nakamura-Yonehara K, Noda M: **Expression of SPIG1 reveals development of a retinal ganglion cell subtype projecting to the medial terminal nucleus in the mouse**. *PLoS One* 2008, **3**(2):e1533.

55. Trakhtenberg EF, Pho N, Holton KM, Chittenden TW, Goldberg JL, Dong L: **Cell types differ in global coordination of splicing and proportion of highly expressed genes**. *Sci Rep* 2016, **6**:32249.

56. Langer KB, Ohlemacher SK, Phillips MJ, Fligor CM, Jiang P, Gamm DM, Meyer JS: **Retinal Ganglion Cell Diversity and Subtype Specification from Human Pluripotent Stem Cells**. *Stem Cell Reports* 2018, **10**(4):1282-1293.

57. Lamba DA, McUsic A, Hirata RK, Wang PR, Russell D, Reh TA: **Generation, purification and transplantation of photoreceptors derived from human induced pluripotent stem cells**. *PLoS One* 2010, **5**(1):e8763.

58. Singh RK, Mallela RK, Cornuet PK, Reifler AN, Chervenak AP, West MD, Wong KY, Nasonkin IO: **Characterization of Three-Dimensional Retinal Tissue Derived from Human Embryonic Stem Cells in Adherent Monolayer Cultures**. *Stem Cells Dev* 2015, **24**(23):2778-2795.

59. Sluch VM, Chamling X, Liu MM, Berlinicke CA, Cheng J, Mitchell KL, Welsbie DS, Zack DJ: **Enhanced Stem Cell Differentiation and Immunopurification of Genome Engineered Human Retinal Ganglion Cells**. *Stem Cells Transl Med* 2017, **6**(11):1972-1986.

60. Meyer JS, Shearer RL, Capowski EE, Wright LS, Wallace KA, McMillan EL, Zhang SC, Gamm DM: **Modeling early retinal development with human embryonic and induced pluripotent stem cells**. *Proc Natl Acad Sci U S A* 2009, **106**(39):16698-16703.

61. Riazifar H, Jia Y, Chen J, Lynch G, Huang T: **Chemically induced specification of retinal ganglion cells from human embryonic and induced pluripotent stem cells**. *Stem Cells Transl Med* 2014, **3**(4):424-432.

62. Reichman S, Terray A, Slembrouck A, Nanteau C, Orieux G, Habeler W, Nandrot EF, Sahel JA, Monville C, Goureau O: **From confluent human iPS cells to self-forming neural retina and retinal pigmented epithelium**. *Proc Natl Acad Sci U S A* 2014, **111**(23):8518-8523.

63.   Reichman S, Slembrouck A, Gagliardi G, Chaffiol A, Terray A, Nanteau C, Potey A, Belle M, Rabesandratana O, Duebel J *et al*: **Generation of Storable Retinal Organoids and Retinal Pigmented Epithelium from Adherent Human iPS Cells in Xeno-Free and Feeder-Free Conditions**. *Stem Cells* 2017, **35**(5):1176-1188.
64.   Rabesandratana O, Goureau O, Orieux G: **Pluripotent Stem Cell-Based Approaches to Explore and Treat Optic Neuropathies**. *Front Neurosci* 2018, **12**:651.
65.   Kobayashi W, Onishi A, Tu HY, Takihara Y, Matsumura M, Tsujimoto K, Inatani M, Nakazawa T, Takahashi M: **Culture Systems of Dissociated Mouse and Human Pluripotent Stem Cell-Derived Retinal Ganglion Cells Purified by Two-Step Immunopanning**. *Invest Ophthalmol Vis Sci* 2018, **59**(2):776-787.
66.   Lamba DA, Karl MO, Ware CB, Reh TA: **Efficient generation of retinal progenitor cells from human embryonic stem cells**. *Proc Natl Acad Sci U S A* 2006, **103**(34):12769-12774.

Figure Legends:

Figure 1: Differentiation of iPSCs into neural retina and retinal ganglion cells (RGCs). A) Schematic showing iPSCs differentiation in neural retinal differentiation media (DMEM/F12 + 2% B27 + 1% N2 + 10 ng/mL IGF1), including inhibitors of BMP, WNT and TGF- beta for 15 days, bFGF2 was included in media from day 4-14. At 16 days *in vitro*, cells were treated with RGC induction and maturation media. Differentiation using four different conditions, LX, LSB, CHIR and LSBX are shown. B) Characterization of RPCs using immunocytochemistry with SOX2, RAX and PAX6 of early retinal progenitor markers. C) Percentage of RPCs expressing retinal progenitor markers (RAX and LHX2) by flow cytometry D) Gene expression profiles showing RPC markers in three different iPSC lines, showing a high enrichment. Mean ± SEM represented from at least 3 independent experiments per individual.

Figure 2: Efficient generation of iPSC-RGCs require inhibition of TGF-beta, Wnt, and BMP signaling. A) Phase contract images of different stages of iPSC-RGC differentiation at DIV7, DIV16 and DIV35 is shown. B) Differentiation of retinal progenitor cells (RPCs) to retinal ganglion cells (RGCs) using defined inhibition or activation of developmental pathways, C) Percentage of Thy-1 (CD90) positive iPSC-RGCs in our studies in different differentiation conditions, D) Proportions of CD90 positive RGC cells across four differentiation conditions is shown. Each symbol represents an individual experiment. E) Percentage of CD90 positive cells obtained across three control iPSC lines across each differentiation condition. Mean ± SEM represented from at least 3 independent experiments per individual.

Figure 3: Transcriptional Profiling of iPSC-RPCs. Quantitative-RT-PCR profiles showing relative expression of retinal ganglion cell and retinal subtypes

(photoreceptors, amacrine, horizontal, and RPE) at Day 23 and Day 35 stages of development from iPSCs to RGCs.

Figure 4: Functional Characteristics of Induced Retinal Ganglion Cells. A) RGCs were transduced with AAV7m8-SNCG-eGFP and immunofluorescence RGC markers (BRN3a, BRN3b, THY1, TUBB3). B) Density of RGCs expressing GFP and other axonal processes increases between DIV35 (4B, cell c) and maturation D72 (4B, cells a and b). Blue arrows point to cells selected for patch clamp recordings and that fired action potential upon depolarization. C) Action potential firing in response to depolarizing voltage step recorded from cell a and cell b (D72) and from cell c (DIV35). D) This trace demonstrates action potential firing in response to the two different depolarizing steps of current recorded from *cell a* from Fig. 4B. We also observe a spontaneous spike firing at the start of the trace. Same scale bar applies to all images.

A

Days in vitro (DIV)

DIV7 (RPC) → DIV16 → DIV35

B

DIV 21 — 23 — 25 — 27 — 35

SHH or SAG	Follistatin	Follistatin	BDNF	Forskolin	DAPT
FGF8	Cyclo	DAPT	NT4	cAMP	
DAPT	DAPT		CNTF	Y-27632	

Long-term RGC cultures

Neural Basal (NB) + B27

C

Days in vitro (DIV35)

LSB   CHIR   LSBX   LX

Autofluorescence ↑

Thy1-APC →

D

% CD90+ cells (Thy1)

Individual per treatment

- LSB
- CHIR
- LSBX
- LX

E

% CD90+ (Thy-1)

- Control 1
- Control 2
- Control 3

LSB   CHIR   LSBX   LX

A

| pSNCG-GFP | CD90/BRN3A | BRN3B/TUBB3 | CD90/TUBB3 |

50 µm

50 µm

B

cell a

cell b

cell c

50 µm

C

cell a

−30 mV   HP=−60 mV

−200

cell b

−30 mV   HP=−60 mV

−200

cell c

−30 mV   HP=−60 mV

−50

−100

1 s

D

0 pA   10 pA   20 pA   holding current

−20

−40

−60

Membrane voltage (mV)

1 s