1   **Edgy conservation: Canadian plants at-risk are overwhelmingly range-edge populations**

2   **and under-studied**

4   S. Klemet-N'Guessan[1][*][†], R. Jackiw[2][*], C.G. Eckert[2][*], A. L. Hargreaves[1][*][†][†]

5   1. Department of Biology, McGill University, 1205 Dr Penfield Ave, Montreal, Quebec,

6   Canada, H3A 1B1

7   2. Department of Biology, Queen's University, 116 Barrie St., Kingston, ON, Canada, K7L

8   3N6

*SKN and RJ made equal contributions, as did CGE and ALH as senior authors.

†Current address: Department of Biology, Trent University, 2140 East Bank Drive,

Peterborough, Ontario, Canada, K9L 1Z8.

††Corresponding Author: anna.hargreaves@mcgill.ca; 514 398 7401

ABSTRACT

As biodiversity declines toward the poles, high-latitude countries will contain the poleward range

edge of many species, potentially focusing national conservation toward range-edge populations

whose global conservation value remains contentious. Using the >200 vascular plants assessed

for protection in Canada, we ask whether national species-conservation rankings are biased

toward range-edge populations and supported by adequate research. Of 192 plant taxa deemed

at-risk in Canada, 77% were only found in Canada at the northernmost 20% or less of their

range. Higher threat categories had more peripheral taxa, and the mismatch between national and

global threat rankings was greater for peripheral vs. non-peripheral taxa. Almost half (43%) of

Canadian at-risk plants had not been studied in the peer-reviewed, conservation-relevant

literature, 57% had not been studied in Canada, and peripheral populations received even less

research effort than non-peripheral taxa. Only 5% of 7-9 conservation-relevant studies assessed

at-risk populations in the context of their geographic range—information that is critical to

establishing their relative conservation value. Thus, flora conservation in Canada is largely the

conservation of edge populations, yet edge populations themselves and the geographic context

that makes them unique are understudied, a research gap we must close to improve evidence-

based conservation.

Keywords: species distributions, range limits, wildlife conservation, cold-edge populations,

endangered species, botanical diversity

INTRODUCTION

The latitudinal gradient in species diversity poses an interesting conservation dilemma for high-latitude countries. As species drop out toward higher latitudes, large polar countries—particularly in the northern hemisphere where landmass is greater—are likely to contain the poleward range edge of many species. All else being equal, species that occur in a jurisdiction only at the edge of their range (hereafter 'peripheral species') will occupy less area and so have fewer individuals than non-peripheral species. This potentially makes peripheral species more likely to be locally rare and deemed 'at risk' even if they have robust populations in their range core (Lesica and Allendorf 1995; Glass et al. 2017). Northern jurisdictions must therefore decide how much of their conservation resources to spend protecting peripheral versus more endemic species (Bunnell et al. 2004). This allocation is tricky as the relative conservation value of peripheral populations remains controversial (e.g. Lesica and Allendorf 1995; Bunnell et al. 2004; Glass et al. 2017)

On one hand, peripheral populations may be less important to species' overall persistence and harder to conserve than populations closer to the range core (Lesica and Allendorf 1995). In the absence of dispersal barriers, range edges are thought to arise from declines in the abundance and quality of habitat, such that peripheral populations are predicted to be less fit, smaller, and more isolated than central populations (Brown et al. 1996; Eckert et al. 2008). Small populations are more prone to genetic drift, which can lead to the fixation of deleterious mutations and interfere with selection of favourable mutations, thereby eroding population fitness and adaptive potential (Brown et al. 1996). This is particularly true of poleward populations that arose from post-glacial range expansions, as expansion additionally reduces genetic diversity and fitness via successive founder effects and 'surfing' of deleterious alleles (Waters et al. 2013; Pironon et al.

2017). Small, isolated populations are also more prone to extinction from demographic and

environmental stochasticity, and it has often proven challenging to maintain or increase

population sizes at species range edges (Bunnell et al. 2004). For these reasons, some biologists

and policy makers have argued against allocating limited conservation resources to edge

populations (Hunter and Hutchinson 1994; Bunnell et al. 2004).

Despite their demographic vulnerability, peripheral populations may harbour distinctive

genetic diversity important for preserving species' evolutionary potential (Hunter and

Hutchinson 1994; Hampe and Petit 2005) and hence capacity for response to rapid

environmental change. If peripheral populations occupy different environments than core

populations (Lesica and Allendorf 1995), they will experience novel selection (Mayr 1954;

Chevin and Lande 2011). Transplant experiments reveal that peripheral populations can be

locally adapted to range-edge conditions (Sexton et al. 2011; Hargreaves and Eckert 2019),

contributing uniquely to intra-species diversity. Dispersal ability is also strongly selected on at

range edges (Phillips et al. 2010; Williams et al. 2016), and both adaptation to local habitat and

increased dispersal ability can prime peripheral populations for colonizing habitat beyond the

species current range (Darling et al. 2008; Hargreaves and Eckert 2014; Hargreaves et al. 2015).

Poleward populations are increasingly valuable in this regard, as they are geographically poised

to initiate poleward range shifts under climate warming (Gibson et al. 2009).

As the above controversy suggests, range position alone may be insufficient to determine

a population's conservation value. Edge populations are not always smaller or more isolated than

range-centre populations, nor does range position predict genetic diversity, genetic quality, or

demographic stability consistently enough to guide conservation policy (Eckert et al. 2008;

Pironon et al. 2017). Empirical evidence for local adaptation in peripheral populations is

decidedly mixed, though still poorly tested (Hargreaves et al. 2014). Finally, even 'super edge

populations' with high performance at many sites in many years may be hard to identify without

large experiments, as ecologically important adaptations can be masked in benign years or by

poor maternal provisioning or inbreeding (Sexton et al. 2011; Hargreaves and Eckert 2019).

Assessing the conservation value of specific edge populations therefore requires empirical

research, yet earlier syntheses suggest peripheral populations are under-studied compared to core

populations (Eckert et al. 2008; Sexton et al. 2009).

Canada is an excellent case for studying peripheral species' representation in

conservation policy and research in high-latitude countries. Canada is the world's second largest

country, spanning almost 10 million km^2 and more than 41° of latitude—as much latitude as

separates Canada from the Equator. Canada's biodiversity is clustered at its southern border

(Coristine et al. 2018), and many species only occur in Canada at the northern tip of a much

larger range (Fig. 1). Canadian conservation legislation seems especially likely to protect such

peripheral species, as both the initial conservation assessment body (Committee on the Status of

Endangered Wildlife in Canada; COSEWIC) and federal Species at Risk Act (SARA) largely

ignore populations outside Canada when determining national conservation priority. Previous

estimates suggest that ~75% of Canada's at-risk species only occur in Canada at their northern

range edge (Yakimowski and Eckert 2007; Gibson et al. 2009). Yet in 2012, three species were

denied federal protection because the government deemed they were 'only' peripheral

populations (Fraser 2000; SARA 2012), logic that could jeopardize the protection of many

species in Canada.

Here we explore the relationships between peripherality (whether a taxon occurs in a

jurisdiction only at its geographic range edge), conservation priority, and research effort. We

consider vascular plants as they are of high conservation value globally but often receive

disproportionately little conservation funding (Raven 1987; Schemske et al. 1994); are relatively

diverse in Canada (e.g. 3600 species compared to ~150 mammals); and make up the largest

proportion (~23%) of taxa assessed by COSEWIC. Using 214 plant taxa with a published

COSEWIC assessment and range map, we designated each as peripheral (northernmost 20% or

less of its range in Canada) or non-peripheral (>20% of its range in Canada). We then tested (*i*)

whether the proportion of peripheral taxa increases from low to high COSEWIC threat rankings

(not-at-risk, special concern, threatened, endangered) and (*ii*) whether Canadian and international

threat rankings differ more for peripheral vs. non-peripheral taxa using standardized NatureServe

rankings. We surveyed the research on at-risk taxa to (*iii)* test whether the conservation-relevant

research effort was evenly spread between peripheral and non-peripheral taxa and (*iv)* assess

how range-wide studies of at-risk taxa could inform conservation.

METHODS

Canadian conservation designation

We considered the 214 vascular plant taxa assessed by COSEWIC as of July 2017 for which a

published COSEWIC assessment was available (one not-at-risk species, *Hackelia ciliata,* was

excluded due to lack of published assessment). Eligibility for protection under Canada's Species

at Risk Act (SARA) is determined by the COSEWIC Vascular Plant subcommittee using

quantitative criteria established by the International Union for the Conservation of Nature

(IUCN). Initial COSEWIC assessments consider only populations in Canada (Gärdenfors 2001;

COSEWIC 2017), but can be adjusted by secondary assessments that consider the likelihood of

genetic replenishment from core populations, local adaptations in Canadian populations,

Canadian habitat's importance to the taxon's persistence, and the condition of populations

outside Canada (IUCN 2018). COSEWIC assessments use the 'best biological information on a

species', including scientific studies, community knowledge, and Aboriginal traditional

knowledge (SARA 2010). The final decision to nationally protect a taxon under SARA rests with

the federal government after considering the socioeconomic implications (SARA 2010); we use

COSEWIC rather than SARA designations throughout as they more closely reflect biology.

Taxa characterization

For each of the 214 taxa (including species, subspecies, varieties, and populations), we recorded

its COSEWIC designation. 192 taxa were designated at-risk, i.e. of special concern (may become

threatened or endangered due to biological constraints and other threats: 21%), threatened (likely

to become endangered: 22%), or endangered (facing imminent extirpation or extinction: 47%).

The remaining assessed taxa were designated 'not-at-risk' (do not face an imminent extirpation

or extinction risk given the current circumstances: 10%). For four species with separate listings

for two geographically distinct populations (*Eleocharis geniculata, Psilocarphus brevissimus*,

Smilax rotundifolia, *Solidago speciosa*), we included both listings as separate data points. For

each taxon, we determined its taxonomic group (angiosperm, gymnosperm, or pteridophyte

(ferns and allies)), lifespan (annual/biennial, perennial, or mixed), growth form (herb/graminoid,

woody shrub, or tree), habitat (aquatic or terrestrial), and general location in Canada. This

information was generally in COSEWIC reports, otherwise we used SARA and plants USDA

databases (SARA Act 2011; USDA 2018).

(i) Proportion of peripheral taxa from lowest to highest COSEWIC threat ranking

To assess whether peripherality is associated with COSEWIC threat ranking, we designated each

taxon as either 'peripheral' (only the northernmost 20% or less of their distribution in Canada;

Fig. 1), or 'non-peripheral'. *Micranthes spicata* had ≤20% of its eastern range edge in the Yukon

with the rest in Alaska; it was deemed non-peripheral. Peripheral designation was determined

visually from range maps in COSEWIC reports (214 taxa), or from the USDA database (USDA

2018) if the COSEWIC report lacked a map (*Pedicularis furbishiae*). For four taxa close to the

20% threshold, we checked visual estimates by digitizing COSEWIC maps and calculating the

proportion range area in Canada (P. Caissy & A.L. Hargreaves, unpublished data).

Analyses were done in R 3.5.1 (R Development Core Team 2018). We tested whether the

proportion of peripheral taxa differed among the four COSEWIC threat rankings using a 4×2

Fisher's exact test (*fisher.test*). As it did, we tested which ranks differed from each other using a

pairwise post-hoc comparison holding the overall $\alpha = 0.05$ (*pairwiseNominalIndependence*,

rcompanion package version 2.1.7 (Mangiafico 2018)). To test whether the pattern of

peripherality vs. threat ranking differed among habitats, life spans, or growth forms, we ran one

binomial generalized linear model (GLM) per grouping, with peripherality as a binomial

response (yes/no) and threat rank and group as interacting categorical predictors (peripheral ~

threat_rank $\times$ group, *glm*, logit link function, Fig. S1). Models of life span excluded 5 taxa with

'mixed' life spans as there were no 'not-at-risk' taxa in this category. Group levels are described

above (*Taxa characterization*). Significance of GLM predictors was assessed using type III tests

(*Anova*, car package version 2.1.4 (Fox and Weisberg 2018)). Finally, we tested whether

peripheral taxa were more endangered overall by converting the four COSEWIC categories to

numeric ranks (1-4) and testing whether this differed between peripheral and non-peripheral taxa

using a non-parametric two-sample Wilcoxon test (*wilcox.test*).

(ii) Canadian vs. global NatureServe threat rankings

We compared Canadian vs. global threat rankings from NatureServe (i.e. 'rounded global status';

NatureServe 2018); NatureServe ranks use consistent criteria so national and global rankings are

directly comparable. Ranks range from 1 (most threatened) to 5 (least threatened, NatureServe

2018). While NatureServe also considers taxa below the species level, Canadian or global ranks

were missing for seven taxa (three at-risk and four not-at-risk), so $n = 207$ taxa (full list of taxa's

COSEWIC, SARA and NatureServe ranks in Dryad data). We calculated the 'rank discrepancy'

for each taxon as 'Global rank – Canadian rank'. Values are numeric but only range from 0 (i.e.

no difference) to 4 (i.e. taxon ranked as 5 (least threatened) globally and 1 (most threatened) in

Canada), so are non-parametric. We tested whether rank discrepancy differed between peripheral

and non-peripheral taxa using a two-sample Wilcoxon test.

Although Nature Conservancy and COSEWIC ranks use different numbers of categories

(five and four, respectively), both are derived from IUCN criteria. To test whether the two bodies

ranked taxa consistently, we converted COSEWIC ranks to numeric ranks from 4 (not-at-risk) to

1 (endangered); as NatureServe did not give any species a 5 (least threatened) in Canada, both

NatureServe Canadian ranks and numeric COSEWIC ranks varied from 1 to 4. We used a paired

Wilcoxon test to assess whether COSEWIC and NatureServe ranks for Canadian populations

differed overall (the five taxa with two distinct COSEWIC designations were each counted

twice). As NatureServe and COSEWIC ranks did differ significantly (Fig. S2a), we tested

whether the difference between NatureServe – COSEWIC Canadian ranks was greater for

peripheral taxa (unpaired Wilcoxon test; Fig. S2b), and whether it altered conclusions re. the

'Global rank – Canadian rank' (paragraph above) by redoing that analysis replacing NatureServe

Canadian ranks with COSEWIC ranks (Fig. S3).

(iii) Research effort on plant taxa considered at-risk in Canada

To assess the research effort on plants at-risk in Canada, we searched ISI Web of Science for

studies on each at-risk taxon using its Latin name, common name, and synonyms listed in its

COSEWIC assessment. While COSEWIC evaluated some specific subspecies, varieties or

populations, we found few studies for these smaller designations so considered all taxa at the

species level (n = 189 at-risk species). We narrowed results to ecological or evolutionary studies

by including a second search term *"*ecolog* OR evolution* OR population* OR demograph*

OR genetic* OR conservation* OR fitness"*. We discarded studies that did not present data on the

taxon of interest (e.g. only mentioned it in key words), yielding 2940 studies.

We categorized the resulting studies on their conservation and geographic relevance. We

categorized conservation relevance as: 1—no data on natural populations or that could be used

by COSEWIC; 2—data from natural populations but not relevant to COSEWIC (e.g. how much

a plant species contributed to a herbivore's diet); or 3—data of potential use to COSEWIC and

conservation (e.g. performance, local adaptation or genetic diversity of natural populations);

ultimately we only considered this third category in analyses (below). We also classified whether

studies sampled wild Canadian populations, wild populations in the USA, both (providing a

wider geographic context for the at-risk Canadian populations), or neither (sampled populations

outside Canada/USA or no specific population). Studies that investigated more than one at-risk

taxon were counted for each taxon included.

We then assessed the distribution of research effort potentially relevant to conservation

(category 3). We first tested whether species that were peripheral or not peripheral in Canada

(binary predictor) differed in the probability that they had been studied anywhere in their range

(yes/no response; binomial GLM), or differed in the number of studies per species from

anywhere in their range (over-dispersed count response, negative binomial GLM, *glm.nb*, MASS

package version 7.3.45 (Venables and Ripley 2013)). We then reran both analyses considering

only studies that included Canadian populations. For negative binomial models, we assessed the

significance of peripherality by comparing the model with peripherality as a predictor (#studies ~

peripheralYN) to the null model (#studies ~ 1) using a likelihood ratio test with significance

evaluated using the Chi squared distribution (*anova*, df = 1).

(iv) Conservation insights from comparing edge vs core populations

Studies that sample populations from both Canada and the USA should reveal the most about the

relative conservation value of peripheral populations in Canada. We therefore read each such

study and noted examples where the results yielded insights relevant to conservation that could

not have been gleaned from smaller-scale sampling.

RESULTS

Of 192 at-risk plant taxa in Canada, 77% were peripheral, with only the northernmost 20% or

less of their distribution in Canada. Many occurred only in Ontario or British Columbia (68% of

148 peripheral taxa vs. 30% of 44 non-peripheral taxa; Fig. 1). Thus, these provinces are

disproportionately responsible for conserving at-risk plants, and peripheral species in particular.

(i) Proportion of peripheral taxa from lowest to highest COSEWIC threat ranking

The percentage of plant taxa that only occur peripherally in Canada increased with increasing

endangerment ($\chi^2_{df=3}$= 17.7, P = 0.0005). The percentage of peripheral taxa was significantly

higher in the Endangered vs. Special concern rank (post-hoc $P = 0.0007$), and the differences

between Endangered and Not at risk and between Threatened and Special concern were almost

significant (post-hoc $P = 0.055$; Fig. 2). The pattern of peripherality vs. threat ranking was

consistent between aquatic vs. terrestrial habitats, herbaceous/graminoid vs. woody growth

forms, and annual/biennial vs. perennial life spans (Fig. S1). Overall, peripheral taxa were

significantly more endangered than non-peripheral taxa (Wilcoxon test $W = 2827$, $P < 0.001$).

(*ii*) *Canadian vs. global NatureServe threat rankings*

For the 207 taxa in our dataset with both Canadian and global NatureServe ranks (157 peripheral,

50 not peripheral), the disparity between Canadian and global ranks was greater for taxa that are

peripheral in Canada compared to taxa that are not (Wilcoxon test, $W = 6756$, $P < 0.001$; Fig. 3).

Of the 203 plant taxa that NatureServe considered at-risk (ranks 1 to 3) in Canada, 67% were

considered secure (ranks 4 or 5) across their global range; most of these at-risk-nationally but

secure-globally taxa are peripheral in Canada (125 peripheral, 14 non-peripheral; Fig. 3).

NatureServe tended to consider taxa more threatened than COSEWIC, but this difference

reflected higher threat rankings for non-peripheral taxa (Fig. S2); the greater discrepancy

between national and global threat ranks for peripheral taxa (Fig. 3) remains if one compares

NatureServe global ranks and COSEWIC ranks directly (Fig. S3).

(*iii*) *Research effort on plant species considered at-risk in Canada*

We found 709 conservation-relevant studies on plant species considered at-risk (by COSEWIC)

in Canada. Almost half (43%) of Canada's 189 at-risk plant species had not been studied in peer-

reviewed work that could inform conservation (Fig. 4a). Though this does not preclude the

existence of studies in the non-refereed literature or journals not indexed on Web of Science, it

suggests that the 'best biological information' is sparse for these taxa. Species peripheral in

Canada and species not peripheral in Canada did not differ in the likelihood that they had been

studied somewhere in their range (43% vs 45%; binomial GLM: peripheral_designation $\chi^2_{df=1}$ =

0.10, P = 0.75) or in the number of studies overall (negative binomial GLM:

peripheral_designation $\chi^2_{df=1}$ = 0.89, P = 0.34; Fig. 4a).

Of the 709 conservation-relevant studies, only 188 included Canadian populations; more

than half (57%) of plant species at-risk in Canada had not been studied in Canada (Fig. 4b).

Compared to non-peripheral species, species that are peripheral in Canada were as likely to have

been studied in Canada (57% vs 55%; binomial GLM: peripheral_designation $\chi^2_{df=1}$ = 0.16, P =

0.69) but had significantly fewer studies per species that included Canadian populations

(negative binomial GLM: peripheral_designation $\chi^2_{df=1}$ = 7.27, P = 0.0070; Fig. 4b).

(iv) Conservation insights from comparing edge vs core populations

Only 42 (5%) of 709 studies on plants at-risk in Canada studied Canadian populations in the

context of their geographic range (i.e. included both Canadian and USA populations). These

studies covered 21 (11%) of 189 at-risk species, and almost half were on one of three species:

Whitebark pine (*Pinus albicaulis*; 8 studies), Butternut (*Juglans cinerea*, 7 studies), and Juniper

sedge (*Carex juniperorum*, 3 studies). Studies that took a range-wide perspective provided

unique insights into the conservation of peripheral populations (Table 1). These include whether

edge populations differ from core populations genetically or demographically, or in their traits or

habitat. For instance, populations of Deerberry decreased in size and frequency toward the

species northern range edge in Canada, but nevertheless were as productive and genetically

diverse as core populations, and showed evidence of local adaptation and high dispersal ability

(Yakimowski and Eckert 2007, 2008). Thus the demographic and genetic value of these

populations was not predicted by their peripherality, size, or spatial isolation.

DISCUSSION

Our results show that conservation of plants in Canada is fundamentally the conservation of

range-edge populations. Three quarters of nationally at-risk plant taxa only occur in Canada at

the northernmost 20% or less of their North American range, in line with earlier estimates for all

at-risk taxa combined (Gibson et al. 2009). While many not-at-risk plants are also peripheral in

Canada (Fig. 2), peripheral taxa had higher threat rankings overall and a greater discrepancy

between their Canadian vs. global threat ranking (Fig. 3). Higher national threat ranks for

peripheral taxa could reflect real increased risk per population, since human activity is also

highest in southern Canada (Coristine and Kerr 2011), or the precautionary principle since

Canada has little control over the fate of populations outside its borders. Either way, northern-

edge populations are geographically poised to initiate northward range shifts under climate

warming, so may be critical for Canada's future biodiversity (Gibson et al. 2009). Moreover,

northward range shifts will bring new species to Canada's southern border, requiring an

informed policy on conserving peripheral populations. Unfortunately, conservation of peripheral

taxa has been debated in the absence of much relevant scientific evidence.

While peripheral populations are prioritized for conservation in Canada, they do not seem

prioritized for peer-reviewed research that could inform their conservation. Based on our Web of

Science survey, more than half of the plant species with at-risk populations in Canada had not

been studied in Canada in a way that could guide their conservation, and peripheral plants had

significantly fewer studies that included Canadian populations than non-peripheral plants (Fig.

4). Even the best-studied at-risk peripheral plant (*Cirsium pitcheri*) had only 8 studies that

included Canadian populations. While this could reflect difficulty in obtaining permits or

adequate sample sizes, taxonomic bias is likely. For example, one bird species that is both

peripheral and at-risk in Canada had almost 50 studies that included Canadian populations

(Marbeled murrelet; Web of Science search May 2019). Thus peripheral plants appear

systematically under-studied in Canada.

Conservation-relevant studies that include both Canadian and US populations illustrate

the value of studying peripheral populations directly and in a broad geographical context.

However, most such studies have assessed neutral genetic diversity and population structure

(Table 1). Conservation would particularly benefit from studies that assess characteristics

important for long-term persistence and range expansion, such as habitat preferences, population

demography and dispersal ability (Schemske et al. 1994). Future genetic work could move

beyond neutral variation to evaluating the adaptive diversity likely to be important in responding

to environmental change (Shaw and Etterson 2012), and local adaptation through which range-

edge populations may contribute uniquely to species' biodiversity (Yeaman et al. 2016). Whether

researchers will close these knowledge gaps depends partially on how government agencies

incentivise (i.e. fund) and remove barriers to (i.e. permit) research on at-risk peripheral

populations. Unfortunately, the "peripherality issue" is not currently highlighted in federal

programs that fund species-at-risk research in Canada (e.g. Government of Canada 2019).

We hope that exposing the lack of peer-reviewed study inspires future work on at-risk

edge populations, but recognize that amassing this work will take time, and that some of the most

informative types of study, e.g. large reciprocal transplants, will be impossible with endangered

taxa. We therefore suggest that we have an excellent and potentially under-used body of research

that could inform Canadian conservation: the already extensive theory and empirical research on

species range edges (Sexton et al. 2009; Pironon et al. 2017). While this research clearly shows

that edge populations can vary significantly from one another in demography (Sagarin et al.

2006) and adaptation (Hargreaves and Eckert 2019), it also reveals broad scale patterns that can

be predictive, e.g. that poleward range edges are often dispersal limited whereas high-elevation

edge populations are often population sinks (Halbritter et al. 2013; Hargreaves et al. 2014). For

countries like Canada whose biodiversity is disproportionately comprised of range-edge

populations, leveraging the species-distribution literature could help inform conservation,

helping meet our commitments to protecting current and future biodiversity.

ACKNOWLEDGEMENTS

We thank Christie Whelan (Department of Fisheries and Oceans), Gary Allen (Parks Canada),

Mark Major (Parks Canada), and Joanna Freeland (Trent University) for providing insight into

Canadian conservation policy and practise, and Pascale Caissy for making the map in Figure 1.

Funding was provided by Natural Sciences and Engineering Research Council of Canada

(Discovery grants to ALH and CGE), and by William E. Bembridge Science Undergraduate

Research Award to SKN.

AUTHOR CONTRIBUTIONS

CGE designed and supervised the original study, which RJ conducted as an undergraduate

project (2014). SKN updated the database in 2017, ALH did the statistics, and SKN and ALH

co-wrote the current manuscript with edits from CGE and RJ.

Table 1. Studies with wide geographic coverage can shed light on the conservation needs and

value of peripheral populations. Key findings taken from studies of plant taxa deemed at-risk in

Canada that sampled both Canadian and USA populations. All taxa are peripheral in Canada or

threatened throughout their range.

Response		
Finding for edge populations	Species	Reference
Genetic uniqueness		
Genetically differentiated from core populations (neutral variation)	Branched bartonia	Ciotir et al. 2013
	Green dragon	Boles et al. 1999
	Spalding's Campion	Lesica et al. 2016
Greatest population differentiation	Butternut	Hoban et al. 2010
Second highest cold tolerance	Whitebark pine	Bower and Aitken 2006
Genetic diversity		
Not lower neutral genetic diversity than core populations	Heartleaf plantain	Mymudes and Les 1993
	Deerberry	Yakimowski and Eckert 2008
	Green dragon	Boles et al. 1999
	Golden paintbrush	Godt et al. 2005
	Cucumber tree	Budd et al. 2015
	Whitebark pine	Liu et al. 2016
Population demography		
Not smaller than core populations	Golden paintbrush	Godt et al. 2005
Smaller & more isolated than core populations	Deerberry	Yakimowski and Eckert 2007
Performance & ecology		
Similar habitat to core populations	Small whorled pogonia orchid	Mehrhoff 1989
Lowest seed viability & distinct dormancy patterns	Green dragon	Yang et al. 1999

Higher seed mass & equal sexual reproduction / productivity as core populations	Deerberry	Yakimowski and Eckert 2007

Peripheral: Branched bartonia = *Bartonia paniculata ssp. paniculata*; Cucumber tree = *Magnolia*
acuminata; Deerberry = *Vaccinium stamineum*; Golden paintbrush = *Castilleja levisecta*; Green
dragon = *Arisaema dracontium*; North American ginseng = *Panax quinquefolius*; Small whorled
pogonia orchid = *Isotria medeleoides*; Heartleaf plantain = *Plantago cordata*; Spalding's
campion = *Silene spaldingii*

Not peripheral: Butternut = *Juglans cinerea*; Whitebark pine = *Pinus albicaulis*
360

Figure 1. Two examples of 'peripheral' species that only occur in Canada at the

northernmost 20% or less of their range. Left is Dense spike-primrose (*Epilobium*

densiflorum, Onagraceae), right is Colicroot (*Aletris farinosa,* Nartheciaceae), both endangered

in Canada. Green polygons show each species' Canadian distribution, dark grey polygons show

its distribution outside Canada. As was the case for 70% of plant taxa considered at risk in

Canada, these species occur in the Canadian biodiversity hotspots of southern British Columbia

(left) or southern Ontario (right). Distribution maps were modified from COSEWIC reports

(COSEWIC 2005, 2015) by Pascale Caissy. Photo credits: Debra L Cook (left); Victoria

McPhail (right).

Figure 2. Higher threat rankings include more peripheral taxa. The percentage of peripheral

taxa (those with ≤ 20% of their northernmost range in Canada) varied among COSEWIC threat r

ankings (total n = 214 plant taxa). Threat ranks are from least threatened (Not at risk) to most thr

eatened (Endangered). The number of taxa per rank is shown at the base of each bar. Contrasting

letters indicate significant differences among categories; the differences between Endangered an

d Not at risk and between Threatened and Special Concern were almost significant (P = 0.055).

381
382

Figure 3. Discrepancy between Canadian and global threat ranks is greater for peripheral

species. Diagonal lines indicate Canadian populations have the same threat ranking as the taxon

globally. Taxa that are peripheral in Canada (left, *n* = 157 taxa) have a greater mismatch between

their Canadian and global threat ranks (more taxa listed as threatened in Canada but secure

globally) than taxa with >20% of their range in Canada (right, *n* = 50 taxa). Point shape indicates

lifespan; colour indicates growth form and habitat (brown = woody shrub or tree, green = non-

woody terrestrial plant, blue = aquatic plant).

Figure 4. Distribution of conservation-relevant research on plant species at-risk in

Canada. (A) Top panels show distribution of all 702 ecological or evolutionary studies that

could inform conservation; (B) bottom panels show 188 studies from (A) that included data from

wild Canadian populations. Left panels = peripheral species (≤ 20% of their range in Canada);

right panels = non-peripheral species >20% of their range in Canada). Black bars indicate species

for which we found no peer-reviewed studies.

REFERENCES

Boles, R.L., Lovett-Doust, J., and Lovett-Doust, L. 2000. Population genetic structure in green dragon (*Arisaema dracontium*, Araceae). Can. J. Bot. **77**(10): 1401-1410. doi:10.1139/b99-089.

Bower, A.D., and Aitken, S.N. 2006. Geographic and seasonal variation in cold hardiness of whitebark pine. Can. J. For. Res. **36**(7): 1842-1850. doi:10.1139/x06-067.

Brown, J.H., Stevens, G.C., and Kaufman, D.M. 1996. The geographic range: size, shape, boundaries, and internal structure. Annu. Rev. Ecol. Syst. **27**: 597-623. doi:10.1146/annurev.ecolsys.27.1.597.

Budd, C., Zimmer, E., and Freeland, J.R. 2015. Conservation genetics of *Magnolia acuminata*, an endangered species in Canada: Can genetic diversity be maintained in fragmented, peripheral populations? Conserv. Genet. **16**(6): 1359-1373. doi:10.1007/s10592-015-0746-9.

Bunnell, F.L., Campbell, R.W., and Squires, K.A. 2004. Conservation priorities for peripheral species: the example of British Columbia. Can. J. For. Res.-Rev. Can. Rech. For. **34**(11): 2240-2247. doi:10.1139/x04-102.

Chevin, L.M., and Lande, R. 2011. Adaptation to marginal habitats by evolution of increased phenotypic plasticity. J. Evol. Biol. **24**(7): 1462-1476. doi:10.1111/j.1420-9101.2011.02279.x.

Ciotir, C., Yesson, C., and Freeland, J. 2013. The evolutionary history and conservation value of disjunct *Bartonia paniculata subsp. paniculata* (Branched Bartonia) populations in Canada. Botany **91**(9): 605-613. doi:10.1139/cjb-2013-0063.

Coristine, L.E., and Kerr, J.T. 2011. Habitat loss, climate change, and emerging conservation challenges in Canada. Can. J. Zool. **89**(5): 435-451. doi:10.1139/z11-023.

Coristine, L.E., Jacob, A.L., Schuster, R., Otto, S.P., Baron, N.E., Bennett, N.J., Bittick, S.J., Dey, C., Favaro, B., and Ford, A. 2018. Informing Canada's commitment to biodiversity conservation: A science-based framework to help guide protected areas designation through Target 1 and beyond. Facets(3): 531-562. doi:10.1139/facets-2017-0102.

COSEWIC. 2005. COSEWIC assessment and status report on the Dense-Spike primrose

Epilobium densiflorum in Canada. Committee on the Status of Endangered Wildlife in

Canada, Ottawa. p. vi + 26.

COSEWIC. 2015. COSEWIC assessment and status report on the Colicroot *Aletris farinosa* in

Canada. Committee on the Status of Endangered Wildlife in Canada, Ottawa. p. xiii + 39.

COSEWIC. 2017. COSEWIC wildlife species assessment: candidate wildlife species. Available

from https://www.canada.ca/en/environment-climate-change/services/committee-status-

endangered-wildlife/wildlife-species-assessment-process-categories-

guidelines/candidate.html [accessed 17 November 2018].

Darling, E., Samis, K.E., and Eckert, C.G. 2008. Increased seed dispersal potential towards

geographic range limits in a Pacific coast dune plant. New Phytol. **178**(2): 424-435.

doi:10.1111/j.1469-8137.2007.02349.x.

Eckert, C., Samis, K., and Lougheed, S. 2008. Genetic variation across species' geographical

ranges: the central–marginal hypothesis and beyond. Mol. Ecol. **17**(5): 1170-1188.

doi:10.1111/j.1365-294X.2007.03659.x.

Fox, J., and Weisberg, S. 2018. An R companion to applied regression. Sage Publications.

Fraser, D.F. Species at the edge: the case for listing of "peripheral" species. *In* Proceedings of a

Conference on the Biology and Management of species and Habitats at Risk. Kamloops,

British Columbia, 15–19 February 1999 2000. *Edited by* L. Darling. British Columbia

Ministry of Environment, Lands and Parks, Victoria, B.C. pp. 49-53.

Gärdenfors, U. 2001. Classifying threatened species at national versus global levels. Trends

Ecol. Evol. **16**(9): 511-516. doi:10.1016/S0169-5347(01)02214-5.

Gibson, S.Y., Van der Marel, R.C., and Starzomski, B.M. 2009. Climate change and

conservation of leading-edge peripheral populations. Conserv. Biol. **23**(6): 1369-1373.

doi:10.1111/j.1523-1739.2009.01375.x.

Glass, W.R., Corkum, L.D., and Mandrak, N.E. 2017. Living on the edge: Traits of freshwater

fish species at risk in Canada. Aquat. Conserv.: Mar. Freshw. Ecosyst. **27**(5): 938-945.

doi:10.1002/aqc.2781.

Godt, M.J.W., Caplow, F., and Hamrick, J. 2005. Allozyme diversity in the federally threatened

golden paintbrush, *Castilleja levisecta* (Scrophulariaceae). Conserv. Genet. **6**(1): 87-99.

doi:10.1007/s10592-004-7746-5.

Government of Canada. 2019. Canada Nature Fund. Available from

https://www.canada.ca/en/environment-climate-change/services/nature-legacy/fund.html

[accessed 16 June 2019].

Halbritter, A.H., Alexander, J.M., Edwards, P.J., and Billeter, R. 2013. How comparable are

species distributions along elevational and latitudinal climate gradients? **22**: 1228-1237.

Hampe, A., and Petit, R.J. 2005. Conserving biodiversity under climate change: the rear edge

matters. Ecol. Lett. **8**(5): 461-467. doi:10.1111/j.1461-0248.2005.00739.x.

Hargreaves, A.L., and Eckert, C.G. 2014. Evolution of dispersal and mating systems along

geographic gradients: implications for shifting ranges. Funct. Ecol. **28**(1): 5-21.

doi:10.1111/1365-2435.12170.

Hargreaves, A.L., and Eckert, C.G. 2019. Local adaptation primes cold- edge populations for

range expansion but not warming- induced range shifts. Ecol. Lett. **22**(1): 78-88.

doi:10.1111/ele.13169.

Hargreaves, A.L., Samis, K.E., and Eckert, C.G. 2014. Are species' range limits simply niche

limits writ large? A review of transplant experiments beyond the range. Am. Nat. **183**(2):

157-173. doi:10.1086/674525.

Hargreaves, A.L., Bailey, S.F., and Laird, R.A. 2015. Fitness declines towards range limits and

local adaptation to climate affect dispersal evolution during climate- induced range

shifts. J. Evol. Biol. **28**(8): 1489-1501. doi:10.1111/jeb.12669.

Hoban, S.M., Borkowski, D.S., Brosi, S.L., McCleary, T.S., Thompson, L.M., McLachlan, J.S.,

Pereira, M.A., Schlarbaum, S.E., and Romero-Severson, J. 2010. Range- wide

distribution of genetic diversity in the North American tree *Juglans cinerea*: a product of

range shifts, not ecological marginality or recent population decline. Mol. Ecol. **19**(22):

4876-4891. doi:10.1111/j.1365-294X.2010.04834.x.

Hunter, M.L., and Hutchinson, A. 1994. The virtues and shortcomings of parochialism:

conserving species that are locally rare, but globally common. Conserv. Biol. **8**(4): 1163-

1165. doi:10.1046/j.1523-1739.1994.08041163.x.

IUCN. 2018. The IUCN Red List of Threatened Species. Version 2018-2. Available from
http://www.iucnredlist.org [accessed 17 November 2018].

Lesica, P., and Allendorf, F.W. 1995. When are peripheral populations valuable for
conservation? Conserv. Biol. **9**(4): 753-760. doi:10.1046/j.1523-1739.1995.09040753.x.

Liu, J.-J., Sniezko, R., Murray, M., Wang, N., Chen, H., Zamany, A., Sturrock, R.N., Savin, D.,
and Kegley, A. 2016. Genetic diversity and population structure of whitebark pine (*Pinus*
albicaulis Engelm.) in western North America. PloS ONE **11**(12): e0167986.
doi:10.1371/journal.pone.0167986.

Mangiafico, S. 2018. rcompanion: Functions to Support Extension Education Program
Evaluation. R package version 2.0.3.

Mayr, E. 1954. Change of genetic environment and evolution. Allen & Unwin, London. pp. 157-
180.

Mehrhoff, L.A. 1989. Reproductive vigor and environmental factors in populations of an
endangered North American orchid, *Isotria medeoloides* (Pursh) Rafinesque. Biol.
Conserv. **47**(4): 281-296. doi:10.1002/j.1537-2197.1993.tb13809.x.

Mymudes, M.S., and Les, D.H. 1993. Morphological and genetic variability in *Plantago cordata*
(Plantaginaceae), a threatened aquatic plant. Am. J. Bot. **80**(3): 351-359.
doi:10.1016/0006-3207(89)90071-2.

NatureServe. 2018. NatureServe Explorer: NatureServe Conservation Status. Available from
http://explorer.natureserve.org/ranking.htm [accessed 17 November 2018].

Phillips, B., Brown, G., and Shine, R. 2010. Evolutionarily accelerated invasions: the rate of
dispersal evolves upwards during the range advance of cane toads. J. Evol. Biol. **23**(12):
2595-2601. doi:10.1111/j.1420-9101.2010.02118.x.

Pironon, S., Papuga, G., Villellas, J., Angert, A.L., García, M.B., and Thompson, J.D. 2017.
Geographic variation in genetic and demographic performance: new insights from an old
biogeographical paradigm. Biol. Rev. **92**(4): 1877-1909. doi:10.1111/brv.12313.

R Development Core Team. 2018. A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0.

Raven, P.H. Scope of the plant conservation problem world-wide. *In* Proceedings of an
International Conference on botanic gardens and the world conservation strategy. Las

Palmas de Gran Canaria, 26-30 November 1985 1987. *Edited by* D. Bramwell. London:

Academic Press.

Sagarin, R.D., Gaines, S.D., and Gaylord, B. 2006. Moving beyond assumptions to understand

abundance distributions across the ranges of species. Trends Ecol. Evol. **21**(9): 524-530.

doi:10.1016/j.tree.2006.06.008.

Schemske, D.W., Husband, B.C., Ruckelshaus, M.H., Goodwillie, C., Parker, I.M., and Bishop,

J.G. 1994. Evaluating approaches to the conservation of rare and endangered plants.

Ecology **75**(3): 584-606. doi:10.2307/1941718.

Sexton, J.P., Strauss, S.Y., and Rice, K.J. 2011. Gene flow increases fitness at the warm edge of

a species' range. Proc. Natl. Acad. Sci. **108**(28): 11704-11709.

doi:10.1073/pnas.1100404108.

Sexton, J.P., McIntyre, P.J., Angert, A.L., and Rice, K.J. 2009. Evolution and ecology of species

range limits. Annu. Rev. Ecol. Syst. **40**: 415-436.

doi:10.1146/annurev.ecolsys.110308.120317

Shaw, R.G., and Etterson, J.R. 2012. Rapid climate change and the rate of adaptation: insight

from experimental quantitative genetics. New Phytol. **195**(4): 752-765.

doi:10.1111/j.1469-8137.2012.04230.x.

Species At Risk Act. 2010. Bill C-5, An Act respecting the protection of wildlife species at risk

in Canada. *Edited by* S.I. Act. Canada Gazette Part II, Ottawa.

Species At Risk Act. 2011. SARA Schedule 1: Species Index. Available from https://wildlife-

species.canada.ca/species-risk-registry/sar/index/default_e.cfm [accessed 3 December

2018].

Species At Risk Act. 2012. List of Wildlife Species at Risk (Decisions Not to Add Certain

Species) Order *Edited by* S.I. Act. Canada Gazette Part II, Ottawa.

USDA. 2018. Plants database.

Venables, W.N., and Ripley, B.D. 2013. Modern applied statistics with S-PLUS. Springer

Science & Business Media.

Waters, J.M., Fraser, C.I., and Hewitt, G.M. 2013. Founder takes all: density-dependent

processes structure biodiversity. Trends Ecol. Evol. **28**(2): 78-85.

doi:10.1016/j.tree.2012.08.024.

Williams, J.L., Kendall, B.E., and Levine, J.M. 2016. Rapid evolution accelerates plant

population spread in fragmented experimental landscapes. Science **353**(6298): 482-485.

doi:10.1126/science.aaf6268.

Yakimowski, S.B., and Eckert, C.G. 2007. Threatened peripheral populations in context:

geographical variation in population frequency and size and sexual reproduction in a

clonal woody shrub. Conserv. Biol. **21**(3): 811-822. doi:10.1111/j.1523-

1739.2007.00684.x.

Yakimowski, S.B., and Eckert, C.G. 2008. Populations do not become less genetically diverse or

more differentiated towards the northern limit of the geographical range in clonal

Vaccinium stamineum (Ericaceae). New Phytol. **180**(2): 534-544. doi:10.1111/j.1469-

8137.2008.02582.x.

Yang, J., Lovett- Doust, J., and Lovett- Doust, L. 1999. Seed germination patterns in green

dragon (*Arisaema dracontium*, Araceae). Am. J. Bot. **86**(8): 1160-1167.

doi:10.2307/2656980.

Yeaman, S., Hodgins, K.A., Lotterhos, K.E., Suren, H., Nadeau, S., Degner, J.C., Nurkowski,

K.A., Smets, P., Wang, T., and Gray, L.K. 2016. Convergent local adaptation to climate

in distantly related conifers. **353**(6306): 1431-1433.

Edgy conservation: Canadian at-risk plants are overwhelmingly range-edge populations and under-studied

Figure S1. The relation between peripherality and threat status does not differ between habitats, lifespans, or growth forms. Each panel was assessed with a binomial GLM: proportion peripheral taxa ~ threat_rank × group, where group was habitat, lifespan, or growth form. The interaction term was non-significant in full models (threat × habitat: $\chi^2_{df=3} = 0.99$, $P = 0.80$; threat × lifespan: $\chi^2_{df=3} = 4.60$, $P = 0.20$; threat × growth form: $\chi^2_{df=3} = 2.92$, $P = 0.41$), so was dropped from final models. In all final models (peripheral ~ threat_rank + group), the proportion of peripheral taxa did not differ among groups (habitat: $\chi^2_{df=1} = 1.68$, $P = 0.19$; lifespan: $\chi^2_{df=1} = 0.51$, $P = 0.47$; growth form: $\chi^2_{df=1} = 0.20$, $P = 0.65$), but differed significantly with COSEWIC status (more peripheral taxa in the endangered vs. special concern rank, as per main results Fig. 2). Threat categories are listed from least to most threatened, a taxon is considered peripheral if it occurs in Canada at the northernmost 20% or less of its range, and numbers at bottom give the number of taxa per bar (total n is lower for (b) as 5 taxa with a mixed lifespan (annual/perennial or biennial/perennial) were excluded).

Figure S2. NatureServe ranks Canadian plant populations as more nationally-imperiled than does COSEWIC. NatureServe ranks range from 1 (most threatened) to 5 (least threatened), but none of the plant taxa assessed by COSEWIC were given a rank of 5. COSEWIC categorical ranks were converted to numeric ranks from 1= endangered to 4 = not at risk. (a) Even assuming equivalent scales (i.e. NatureServe 4 equivalent to COSEWIC 4), NatureServe ranks of Canadian populations differed from COSEWIC ranks (paired Wilcoxon test $W = 1078$, $P = 0.0006$; n = 207 taxa) in that NatureServe tended to rank taxa as more nationally threatened than COSEWIC. (b) The difference between NatureServe and COSEWIC rankings in (a) was driven primarily by non-peripheral taxa, which showed a bigger discrepancy (unpaired Wilcoxon test $W = 4788$, $P = 0.008$).

Figure S3. Peripheral taxa have a greater discrepancy between Canadian and global threat ranks whether one uses NatureServe Canadian ranks (a) or COSEWIC ranks (b). Threat rankings are described in Fig. S2 (1=most threatened). (a) Same data as shown in Fig. 3: the difference in NatureServe Global – NatureServe Canadian threat ranks is greater for peripheral vs non-peripheral species. (b) The difference between NatureServe Global – COSEWIC Canadian ranks is also greater for peripheral vs. non-peripheral species (Wilcoxon test, W = 7072, P < 0.001).