High conservation priority of range-edge plant populations not matched by habitat

protection or research effort

Running head: Conserving range-edge plant populations

P. Caissy[1*†], S. Klemet-N'Guessan[1*††], R. Jackiw[2], C.G. Eckert[2], A. L. Hargreaves[1†††]

*joint first-authors.

1. Department of Biology, McGill University, 1205 Dr Penfield Ave, Montreal QC, Canada,

H3A 1B1

2. Department of Biology, Queen's University, 116 Barrie St., Kingston ON, Canada, K7L 3N6

†Present address: Département de Biologie, Université Laval, 1045 Avenue de la Médecine,

local 3058, Québec, QC, Canada, G1V 0A6.

††Present address: Department of Biology, Trent University, 2140 East Bank Drive,

Peterborough, ON, Canada, K9L 1Z8.

†††Corresponding Author: anna.hargreaves@mcgill.ca; 514 398 7401

Keywords: species distributions, range limits, wildlife conservation, cold-edge populations,

endangered species, botanical diversity

ABSTRACT

High-latitude countries tend to contain the polar range-edge of many species that are nationally

rare but globally common. This can focus national conservation efforts toward range-edge

populations, whose conservation needs and value are disputed. Using plants in Canada as a case

study, we ask whether national species-conservation rankings prioritize range-edge populations,

and whether conservation priority is matched by habitat protection and research effort. We found

that >75% of federally-protected plants only occur in Canada peripherally, at the northernmost

20% or less of their total range, and that the most imperilled taxa had the smallest percentage of

their range in Canada (endangered plants: median=1.0%). Occurring peripherally in Canada was

associated with higher threat even after accounting for range area, potentially because range-

edge taxa experienced 85% higher human population densities in their Canadian range than non-

peripheral taxa. High conservation priority was not matched by habitat protection, as more

imperilled and more peripheral taxa had smaller fractions of their Canadian range in protected

areas. Finally, peer-reviewed research on plants at-risk in Canada was low. Only 42% of plants

considered at-risk in Canada had been studied in Canada, and only 11% of species with large

distributions outside Canada had been studied in the context of their wider geographic range—

information that is critical to establishing their relative conservation value. Our results illustrate

that plant conservation in Canada is fundamentally linked to conserving range-edge populations,

yet edge populations themselves are understudied, a research gap we must close to improve

evidence-based conservation.

INTRODUCTION

The conservation value of populations at the edge of a species' distribution is contentious. Species range edges often coincide with declining suitability and abundance of habitat; about three quarters of transplant experiments find declines in performance beyond a species' range (Hargreaves et al. 2014; Lee-Yaw et al. 2016). Edge populations in poor-quality or isolated habitat are, therefore, predicted to be small (Hengeveld and Haeck 1982; Brown et al. 1996), eroding their genetic quality through genetic drift (Ellstrand and Elam 2003). Small population size and low genetic quality can make edge populations less important to species' persistence and harder to conserve (Bunnell et al. 2004). However, even dramatically reduced performance beyond the range does not necessarily mean that habitat at the range edge is low quality (Hargreaves and Eckert 2019), and while some edge populations are smaller, less genetically diverse, or more inbred than core populations (Sexton et al. 2011), these patterns are far from universal (Villellas et al. 2013; Pironon et al. 2015, 2017; de Medeiros et al. 2018).

Indeed, range-edge populations may be particularly valuable for species' long-term success either genetically or geographically. For widespread species, range edges often include extreme or unusual habitats (Thakur et al. 2018); if edge populations are locally adapted, they may contribute uniquely to species' overall genetic diversity (Bunnell et al. 2004; Sexton et al. 2011). For species whose ranges have expanded and contracted with glacial cycles, populations at the equatorial range edge may harbour disproportionate genetic diversity that could be critical for long-term persistence (Hampe and Petit 2005). Finally, while many taxa are shifting to higher elevations and latitudes in response to climate warming (Chen et al. 2011; Freeman et al. 2018), warming is expected to outpace dispersal ability for 11% of species globally (Thomas et al.

2004). Cool-edge populations are geographically poised to initiate range shifts (Gibson et al.

2009), and may have evolved higher dispersal abilities due to past expansions or demographic

instability (Phillips et al. 2010; Hargreaves et al. 2015), making them invaluable for successful

range shifts.

The conservation value of range-edge populations is especially germane when conservation

policy prioritizes (or deprioritizes) them implicitly or explicitly. Many political jurisdictions rank

species by local conservation need, e.g., national, state, or provincial Red Lists. As political

borders rarely follow biogeographic boundaries, jurisdictions may contain range-edge

populations of species widely distributed outside their borders (hereafter 'peripheral taxa'; Hunter

and Hutchinson 1994). For example, most US states contain >20 peripheral reptiles or

amphibians, but many explicitly deprioritize peripheral species in conservation listings,

sometimes jeopardizing species' overall persistence (Steen and Barrett 2015). Conversely,

peripheral taxa may be implicitly prioritized; all else being equal they will occupy less area in a

jurisdiction, making them more likely to be locally rare and deemed 'at-risk' (Lesica and

Allendorf 1995; Glass et al. 2017). In jurisdictions with local endemism, local conservation

rankings can show considerable mismatch from global conservation priorities. For example,

>75% of Finland's rare beetles (Komonen 2007) and >70% of Canada's at-risk flora and fauna

(Gibson et al. 2009; Cameron and Hargreaves 2020) have wide distributions south of these

countries' borders.

For jurisdictions with many peripheral populations, resolving their conservation value is highly

relevant to effective conservation policy, but may require direct study of edge populations

themselves. Range position does not consistently predict population size, genetic diversity, or

demographic stability (Eckert et al. 2008; Pironon et al. 2017), and the extent of local adaptation

to range-edge conditions can be hard to identify without large experiments (Sexton et al. 2011;

Hargreaves and Eckert 2019; Anderson & Wadgymar 2020). However, earlier syntheses suggest

range edges are under-studied compared to core populations (Eckert et al. 2008; Sexton et al.

2009), potentially impeding evidence-based conservation.

An excellent case study for exploring 'peripherality' (the extent to which a species occurs in a

jurisdiction at the edge of its range) in conservation is Canada. Canada is the world's second

largest country, spanning almost 10 million km^2 and >41° of latitude—as much latitude as

separates Canada from the Equator—and contains the northernmost potential land for species in

the Americas. Canada's biodiversity is clustered at its southern border (Coristine et al. 2018), as

is Canada's human population, potentially increasing threats to range-edge taxa and creating

conflicts between conservation and economic development. Previous estimates suggest ~75% of

terrestrial taxa designated nationally at-risk only occur in Canada at their northern range edge

(Yakimowski and Eckert 2007; Gibson et al. 2009), but this has not been formally quantified.

Finally, risk-assessments are publicly available for all taxa assessed for federal protection (by the

Committee on the Status of Endangered Wildlife in Canada, COSEWIC) and generally include a

detailed range map.

We test the relationships between peripherality and: range area in Canada; conservation priority

in Canada; conservation risk (human population density) and habitat protection (protected areas);

and peer-reviewed conservation research effort. We use vascular plants as they are relatively

diverse (3600 species in Canada compared to ~150 mammals), are of high conservation value

globally, and often receive disproportionately little conservation funding (Raven 1987;

Schemske et al. 1994). Using COSEWIC risk assessments and range maps (Fig. 1),

complemented by NatureServe risk rankings, human population censes, and literature searches,

we address five key questions. *Q1)* Do Canadian and global threat rankings differ more for taxa

that are peripheral in Canada vs. those that are not, as expected if peripheral taxa have smaller

ranges or populations in Canada or face greater threats (see below)? *Q2)* Do more nationally-

imperilled taxa have smaller ranges in Canada or smaller proportions of their total range in

Canada (i.e. more peripheral), as expected if small ranges and increased peripherality are

associated with conservation risk? *Q3*) Do taxa that occur more peripherally in Canada

experience higher human population density in their Canadian range, potentially explaining an

association between peripherality and conservation risk? *Q4*) Do more-imperilled or more

peripheral taxa have a greater proportion of their range in protected areas, potentially indicating

active conservation effort? *Q5*) Is conservation research effort evenly spread between range-edge

taxa (<20% of their total range in Canada) and non-edge taxa, and how might range-wide studies

inform conservation?

Figure 1. Most plant taxa assessed for protection in Canada only occur in Canada at their northern range edge. Maps illustrate the Canadian distributions, digitized from COSEWIC risk assessments, of 182 of the 220 extant vascular plant taxa assessed from 1977 to 2018, grouped by their COSEWIC-designated risk status (the remaining 38 taxa did not have digitizeable maps but estimates of peripherality are similar; *Table 1*). If taxa were assessed more than once we use the most recent assessment. Most plant taxa in Canada have not been assessed by COSEWIC, so the *Not at Risk* category does not represent all non-imperilled plants but those that were thought to be potentially at risk and later deemed secure. Thick black line shows the Canada-USA border, coloured dots introduce the colour scheme used in subsequent figures.

METHODS

Threat status

Flora eligibility for protection under Canada's Species at Risk Act (SARA) is determined by the

COSEWIC Vascular Plant subcommittee using quantitative criteria established by the

International Union for the Conservation of Nature (IUCN). COSEWIC was created in 1977, and

SARA was passed in 2002 (SARA 2002). There are three steps to a COSEWIC recommendation.

1) COSEWIC prioritizes which taxa to assess. In this phase COSEWIC prioritizes taxa likely to

become (globally) extinct (SARA 15.1b). 2) If a taxon is prioritized for assessment, COSEWIC

commissions and then reviews and approves an assessment report using the 'best biological

information' available, including scientific, community, and Aboriginal traditional knowledge

(SARA 2002). 3) COSEWIC uses the report to recommend a status: Special concern (may

become threatened or endangered due to biological constraints and other threats); Threatened

(likely to become endangered if threats not mitigated); Endangered (facing imminent extirpation

or extinction), Extirpated, Extinct, or Data deficient (we do not consider Extirpated, Extinct, or

Data deficient taxa as they have no current Canadian range or reliable range map); or Not at risk

(no imminent risk of extirpation or extinction). COSEWIC determines a status considering only

Canadian populations, then reviews whether adjustments are warranted given the likelihood of

demographic rescue from populations outside Canada (Environment and Climate Change Canada

2017). The final decision to protect taxa under SARA rests with the federal government after

considering the socioeconomic implications of COSEWIC's recommendation (SARA 2002). We

therefore use COSEWIC rather than SARA designations as they more closely reflect biology.

As of August 2018, COSEWIC assessment reports were available for 220 plant populations (1

report each, most available on line but some obtained from COSEWIC directly; sample size

details in Table 1). If the taxon had been reassessed we used the most recent assessment. A few

taxa had reports for two geographically distinct populations ('designatable units' in COSEWIC

terminology). If both populations had the same threat status we combined them, otherwise we

excluded the taxon as it was unclear how to calculate a corresponding global range per

population (Table 1). We excluded reports without range maps or descriptions from which we

could estimate the proportion of each range in Canada, yielding 209 taxa including species and

subspecies (Table 1). We recorded each taxon's COSEWIC threat status: Special concern,

Threatened, Endangered, or Not at risk; the first three categories are considered 'at-risk'.

COSEWIC only assesses taxa that might be imperilled. The 'Not at risk' category does not

reflect all plants that are secure in Canada, but a group of taxa deemed potentially at risk

(therefore worth assessing) but ultimately secure (e.g., not in sufficient decline for listing,

reassessment revealed other populations, taxon not a true taxonomic unit).

Table 1. Sample sizes, with data used for each Question in bold. At-risk taxa are those
considered Special concern, Threatened, or Endangered, and can have any proportion of their
range in Canada. Peripheral taxa include at-risk and not-at-risk taxa. 'At-risk' and 'Peripheral'
columns give the number of populations or taxa and the percentage of the total (i.e. divided by
column 2). Analyses for Q2–4 use a continuous measure of peripherality only available for
species with digitized range maps. Analyses for Q1 and Q5 define peripherality categorically and
so include a larger data set: main analyses use a 20% cut off (species with <20% of their global
range in Canada = peripheral), and include taxa with digitized maps and taxa for whom we could
estimate peripherality (Y/N) from range descriptions. Using only taxa with digitized maps and a
10% cut-off yielded consistent results (Supp Info).

| | | | Extant plants in Canada | | |
| | | | Peripheral | | |
	Total	At-risk (COSEWIC)	20% cut-off	10% cut-off	Question
(a) populations with a COSEWIC status	220	197 (90%)	–	–	–
(b) taxa with COSEWIC report and range map[1]	209	**189** (90%)	165 (79%)[2]	–	Q5
(c) taxa in (b) with NatureServe ranks for world and Canada	**202**	186 (92%)	159 (79%)[2]	–	Q1
(d) taxa in (b) with digitizeable range map	**182**	166 (91%)	141 (77%)[3]	127 (70%)[3]	Q2, Q3, Q4

1. From (a), (b) excludes 2 taxa with no available COSEWIC report (*Hackelia ciliata*,
Symphyotrichum sericeum), 2 taxa whose COSEWIC report has no map (*Carex nebrascensis,*
Pedicularis furbishia), and 6 populations from 3 taxa where 2 designatable units (populations)
had different COSEWIC statuses (*Eleocharis geniculata, Psilocarphus brevissimus, Solidago*
speciosa). *Smilax rotundifolia* had 2 populations assessed with the same COSEWIC status;
these have been merged into 1 taxon.
2. For 27 taxa whose distributions could not be digitized (b minus d), we estimated whether they
were peripheral in Canada (20% cut-off) based on undigitizeable maps and range descriptions.
We did not do this for the 10% cut-off as it was harder to determine reliably.
3. 27 taxa maps could not be digitized as they did not show the complete range (19), or were in a
projection that could not be digitized (4) or were otherwise unclear (4).
195

For each taxon with a COSEWIC range map, we compiled Canadian and global threat rankings

from NatureServe (i.e. 'rounded global status'; NatureServe 2018). NatureServe ranks use

consistent criteria so national and global rankings are directly comparable (a detailed comparison

with COSEWIC statuses is in Supp Info Fig. S1). Ranks range from 1 (most threatened) to 5

(least threatened; NatureServe 2018). NatureServe Canadian or global ranks were missing for

seven species, so n = 202 taxa (Table 1). We calculated the 'rank disparity' for each taxon as

'Global rank – Canadian rank'.

Range maps & area

Of the 209 COSEWIC range maps, we were able to digitize 182 using the geographic

information software *Quantum GIS 2.18* (QGIS Development Team 2018; Fig 1). We

georeferenced maps using ≥10 dispersed coordinates of obvious landmarks, such as country,

province, county or water-body boundaries. All maps underwent a thin plane spline

transformation to allow local deformation, and to standardize map projections to the World

Geodetic System 1984 projection (WGS 84, EPSG 4326), a standard projection for worldwide

geographic datasets and convenient when working with latitude and longitude coordinates (Tim

Elrick, McGill Geographic Information Centre administrator; personal communication).

COSEWIC provided maps as polygons (160 taxa) or points occurrences (29 taxa). For polygon

maps, we traced digital polygons by hand, and for point occurrence maps, we generated a convex

polygon including all points (generally equivalent to COSEWIC's 'extent of occupancy'). If we

were unable to generate a convex polygon (<4 points on the map; n = 2 taxa), we generated a 1

km buffer around each point creating distinct polygons. We could not digitize maps for 27 taxa

whose maps were incomplete or imprecise (as noted by COSEWIC or if mapped as presence

/absence by province) or used a projection that did not allow for proper digitization (e.g.,

circumpolar projection; Table 1).

For each taxon with a range map, we estimated the proportion of its range in Canada. For 182

taxa with a digitized map, we calculated *a)* global range area (km^2), *b)* Canadian range area

(km^2), *c)* proportion range in Canada ($b \div a$) as a quantitative measure of peripherality. For area

calculations, we projected individual shapefiles in QGIS, using the World Mollweide equal-area

projection (WGS 84, EPSG 54 009) since some taxa had large ranges encompassing lower

latitudes such as Mexico (Usery and Seong 2000). Range area was extracted with the *gArea*

function in the "rgeos" package (version 0.4-1; Bivand and Rundel 2018) in R version 3.2.2. (R

Core Team 2017).

For all 209 taxa with COSEWIC range maps, we also assigned a categorical designation of

peripherality to simplify analyses comparing national vs. global threat rank and research effort

(Questions 1 & 5). We designated taxa as peripheral if they had <20% of their total range in

Canada, otherwise not. Any threshold is arbitrary; we chose 20% as all 27 taxa whose map could

not be digitized could still be unambiguously categorized as peripheral or not (i.e. clearly had

<20% or >20% of their range in Canada) from maps and range descriptions provided in

COSEWIC assessments and recovery plans. To test the sensitivity of results we ran two

supplementary analyses. First, we re-ran analyses using only taxa with digitized range maps

(Table 1d). Second, we used a more conservative cut-off of 10% (Table 1d), the cut-off used by

COSEWIC to define their lowest level of Canadian responsibility (Environment and Climate

Change Canada 2017).

Covariates

For taxa with digitized range maps (*n* = 182), we calculated two covariates. First, we estimated

the human population density in each taxon's Canadian range using the dissemination blocks

from the 2016 Canadian census (Statistics Canada 2016). Dissemination blocks are the smallest

geographic units used by Statistics Canada (equivalent to a city block bounded by intersecting

streets, with block size determined by road density), for which inhabitants/block data are

available (Statistics Canada 2011). Using *QGIS 2.18* (QGIS Development Team 2018), we

overlaid each taxon's Canadian range polygon on the dissemination block map (McKie 2016). In

R, we then summed the inhabitants across all blocks and partial blocks within each species

range, and then divided this sum by the taxon's Canadian range area to estimate human

population density.

Second, we calculated the officially protected area within each taxon's Canadian range using two

protected area databases. The Canadian Council on Ecological Areas database includes all levels

of protected area (e.g., municipal, provincial, federal) in all Canadian provinces and territories

except the province of Quebec (CCEA 2016). Quebec's 'Ministère du Développement Durable

Environnement et Lutte contre les Changements Climatiques' database maps all protected areas

in Quebec (MDEL 2016). We projected the protected area shapefiles to the World Mollweide

equal-area projection (WGS4 EPSG 54 009), then overlaid them on each species' Canadian

range polygon to generate a geographic file of protected habitat. We then calculated the area

(km^2) of protected habitat in each range polygon in R.

Literature search for peer-reviewed studies on at-risk plants

To assess the peer-reviewed research effort on plants deemed at-risk in Canada, we searched

Web of Science for studies on each at-risk taxon with a COSEWIC range map (Table 1b) using

its scientific name, English common name, and synonyms listed in its COSEWIC assessment (up

to August 2017). We searched all taxa at the species level since few studies were available at

lower taxonomic designations (*n* = 209 at-risk species). We narrowed results to ecological or

evolutionary studies by including the search term *"*ecolog* OR evolution* OR population* OR*

demograph* OR genetic* OR conservation* OR fitness". We discarded studies that did not

present data on the taxon of interest (e.g., only mentioned it in key words), yielding >2900

studies.

We assessed the conservation and geographic relevance of the resulting studies. Studies were

deemed conservation relevant if they contained data on natural populations that would be

potential use to COSEWIC (e.g., population censuses, performance, life-history, local

adaptation, genetic diversity). Studies that contained no data on natural plant populations or data

that was not relevant to their conservation (e.g., how much a plant species contributed to a

herbivore's diet) were not considered further. We further classified whether each relevant study

sampled wild Canadian populations, wild populations in the USA, both (providing a wider

geographic context for at-risk Canadian populations), or neither (sampled populations outside

Canada/USA or no specific population). Studies that investigated more than one at-risk taxon (32

studies) were counted for each taxon included.

Analyses

Q1) Do Canadian & global NatureServe ranks differ more for peripheral vs. non-peripheral

taxa?

Taxa cannot be less threatened (i.e. more secure) nationally than they are globally, so we are not

testing whether a disparity exists or the direction of the disparity, but whether the disparity is

bigger for peripheral vs. non-peripheral taxa. Threat-rank disparity (NatureServe Global rank –

NatureServe Canadian rank) is numeric but only ranges from 0 (i.e. no difference) to 4 (i.e. taxon

ranked as 5 (least threatened) globally and 1 (most threatened) in Canada), requiring non-

parametric analyses. We tested whether threat-rank disparity was greater for peripheral vs. non-

peripheral taxa using a non-parametric two-sample Wilcoxon test. We used categorical

peripherality (peripheral if <20% of range in Canada, otherwise not; Table 1c) to match the (lack

of) resolution in the response variable. As detailed above, we tested whether our results were

robust to using a) only taxa with digitized maps, and b) defining peripheral as ≥10% of range in

Canada (Supp Info).

Q2) Do more imperilled taxa have smaller ranges or range proportions in Canada?

We ran one generalized linear model (GLM) for each of three response variables: global range

area, Canadian range area, and percentage of range in Canada. Each model considered

COSEWIC status as a categorical predictor (model structure: response ~ status). We predicted

that global range area would not differ among COSEWIC ranks, but that Canadian range area

and percentage range in Canada would decline with increasing threat. We used negative binomial

error distributions for over-dispersed count data (range area), and quasi-binomial distributions

for over-dispersed proportional data (range %). Here and for all GLMs, we assessed predictor

significance by comparing models with and without the predictor of interest using a likelihood

ratio test compared to a Chi-squared distribution (*anova* function in R). When COSEWIC status

was significant, we assessed which statuses differed using least squared mean contrasts with a

Tukey correction to maintain alpha = 0.05 (package *lsmeans*, version 2.30-0; Lenth 2016). To

test whether peripherality was associated with increased imperilment even after accounting for

range area, we translated COSEWIC status to integers (1 = Not at risk, 4 = Endangered). We

used this 'numeric status' as the response in a quasiPoisson GLM, with Canadian range area and

proportion of range in Canada as predictors (numeric COSEWIC status ~ Canadian range area +

proportion range in Canada).

COSEWIC procedures changed slightly once SARA was passed in 2002. At the suggestion of

people familiar with COSEWIC, we reran all analyses for Q2 including a categorical covariate

for whether taxa were assessed before or after 2002, to see whether results differed (Supp Info).

Q3) Do more peripheral and/or imperilled taxa have more people in their Canadian range?

We tested whether human population density (response) varied with the proportion of global

range in Canada (proportional predictor) and among COSEWIC statuses (categorical predictor),

as expected if human population density and peripheral taxa co-occur close to Canada's southern

border (human density ~ status + proportion range in Canada). We used a negative binomial

GLM and assessed significance as for *Q2*.

Q4) Do more peripheral and/or imperilled taxa have more of their Canadian range protected?

We tested whether the proportion of a taxon's Canadian range that overlapped with a protected

area (overdispersed proportional response) varied among COSEWIC statuses (categorical

predictor) and with peripherality (proportional predictor) using a quasibinomial GLM: proportion

Canadian range protected ~ status + proportion range in Canada. Predictor significance was

determined as in *Q2*, and we extracted the fit lines and 95% confidence intervals for the effect of

peripherality using the *visreg* R package (version 2.4.1, Breheny and Burchett 2017).

Q5) Conservation research effort and insights from range-wide studies on at-risk plants

For the 189 plant species at-risk in Canada with quantifiable ranges (Table 1b), we tested

whether peripheral vs. non-peripheral taxa differed in whether or how often they had been

studied in the conservation-relevant literature, both in across their entire range and in Canada

specifically. All four GLMs used categorical peripherality (Y if <20% of global range in Canada)

as a predictor. Response variables were: 1) whether the taxa had been studied anywhere in its

range (binomial response/GLM); 2) the number of studies from anywhere in the range (negative

binomial response/GLM); 3) whether the taxa had been studied in Canada (binomial GLM); 4)

the number of studies that included Canadian populations (negative binomial GLM). The effect

of being peripheral was evaluated using likelihood ratio tests, as in *Q2*.

For taxa that have a large fraction of their range outside Canada, studies that sample populations

from both Canada and the USA should reveal the most about the relative conservation value of

peripheral populations in Canada. We therefore reduced the data above (189 taxa) to species with

less than half their range in Canada, selected the conservation-relevant studies on these taxa that

included Canadian populations, and read each study carefully to note examples that yield insights

relevant to conservation that could not have been gleaned from smaller-scale sampling.

RESULTS

Of 189 plant taxa considered at-risk in Canada (Table 1b), 152 taxa (80%) occurred in Canada in

less than 20% of their range, of which 151 were at their northern range edge (*Micranthes spicata*

has the eastern tip of its range in Canada, with the rest in Alaska, Fig. 1 top right). Many at-risk

plants occurred only in southern Ontario or British Columbia (68% of 152 peripheral at-risk taxa,

19% of 37 non-peripheral taxa; Fig. 1). Thus, these provinces are disproportionately responsible

for conserving at-risk and range-edge plants.

Q1) Do Canadian & global NatureServe ranks differ more for peripheral vs. non-peripheral

taxa?

As predicted, the disparity between NatureServe's Canadian and global ranks was greater for

taxa that are peripheral in Canada compared to taxa that are not (Wilcoxon test, $W = 5878$, $P <$

0.0001, $n = 202$; Fig. 2). Of the 198 plant taxa that NatureServe considered at-risk (ranks 1 to 3)

in Canada, 67% were considered secure (ranks 4 or 5) across their global range; most of these

nationally-at-risk but globally-secure taxa are peripheral in Canada (123 peripheral, 10 non-

peripheral; Fig. 2). Results were consistent using COSEWIC vs. NatureServe Canadian ranks

(Fig. S1), or using only taxa with digitized maps and a 10% vs 20% threshold for peripherality

(Table S1).

371
Figure 2. Disparity between Canadian and global threat ranks is greater for peripheral species. (Results for *Question 1*). Diagonal lines indicate Canadian populations have the same threat ranking as the species does globally (values above the line are not possible; points are jittered for visualization). Taxa with <20% of their range in Canada (left, *n* = 159) have a greater mismatch between their Canadian and global threat ranks (more taxa listed as threatened in Canada but secure globally) than taxa with >20% of their range in Canada (right, *n* = 43). Point shape indicates lifespan; colour indicates growth form and habitat (brown = woody shrub or tree, green = non-woody terrestrial plant, blue = aquatic plant). Sample sizes details in Table 1c.

Q2) Do more imperilled taxa have smaller ranges or range percentages in Canada?

Plants considered at-risk by COSEWIC (status = Special Concern, Threatened, or Endangered) generally had large global distributions (median = 390,521 km^2), with much smaller ranges in Canada (median = 4,598 km^2) that were clustered toward Canada's southern border (Fig. 1). At-risk taxa had a median of 1.8% of their global range in Canada. As predicted, global range size did not differ among Canadian status categories (χ^2_{df3} = 7.5, *P* = 0.057; Fig. 3a). However, taxa assessed and deemed at-risk had significantly smaller Canadian ranges than taxa assessed and deemed Not-at-risk taxa (χ^2_{df3} = 40.7, *P* < 0.0001; Fig. 3b), and the most imperilled taxa had the

smallest percentages of their range in Canada (χ^2_{df3} = 15.0, P = 0.0002; Fig. 3c). Results did not

differ between taxa assessed before or after SARA was passed (Table S2).

The extent to which taxa occurred in Canada at the edge of their range was associated with

increased conservation threat, even after accounting for absolute range area in Canada. Numeric

COSEWIC status increased (i.e. more imperilled) as range size in Canada decreased (χ^2_{df1} = 8.8,

P = 0.003) and as range percentage in Canada decreased (χ^2_{df1} = 16.9, P < 0.001; Fig. S3).

398
399
Figure 3. Global vs. Canadian range size for plants assessed for protection in Canada.
(Results for *Question 2*). Differing letters indicate significant differences among COSEWIC
threat statuses within panels. The *Not at Risk* status does not represent all non-threatened plants
but those that were thought to be potentially at risk but later deemed secure. The lower, middle
and upper horizontal lines in each boxplot indicate the 25th percentile, the median and the 75th
percentile, respectively. Whiskers extend 1.58 interquartile range / square root *n* from the median
or to the extreme points, whichever is less. Coloured points show raw data (horizontal jitter to
facilitate visualization of overlapping points); sample sizes in Table 1d.
408

Q3) Do more peripheral and/or imperilled taxa have more people in their Canadian range?

Human population density differed with threat status, but not with peripherality. More imperilled

plants had significantly higher human population densities in their Canadian ranges compared to

taxa assessed at lower threat ranks ($\chi^2_{df3} = 36.7$, $P < 0.0001$; Fig. 4). After accounting for

differences among threat ranks, taxa that are more peripheral in Canada did not have higher

human densities within their Canadian range ($\chi^2_{df1} = 1.7$, $P = 0.20$).

Figure 4. Human population density varied among COSEWIC threat statuses. (Results for
Question 3). Boxplots are as in Fig. 3. Sample size details in Table 1d.

Q4) Do more peripheral and/or imperilled taxa have more of their Canadian range protected?

Plant taxa considered at-risk in Canada had only 3.7% (median) of their Canadian range

protected, and only 8 taxa had >50% of their Canadian range protected (Fig. 5). Habitat

protection varied among threat statuses (COSEWIC status: $\chi^2_{df3} = 15.1$, $P = 0.0017$), but contrary

to predictions the taxa the are the most imperilled in Canada tended to have the lowest fraction of

their Canadian range protected; Fig. 5a). Habitat protection also varied with peripherality

(proportion range in Canada: $\chi^2_{df1} = 37.0$, $P < 0.001$). As predicted, the more peripherally a taxon

occurred in Canada, the smaller the fraction of its Canadian range was protected (Fig. 5b).

Figure 5. Habitat protection varies with threat status and peripherality. (Results for
Question 4). (A) Taxa more imperilled in Canada (threatened, endangered) have the smallest
proportion of their Canadian range protected. Boxplot formatting and *n* as in Figs. 2&3. (B) Taxa
with less of their global range in Canada (more peripheral) have smaller proportions of their
Canadian range protected. Line, shading, and points show fit, 95% confidence intervals, and
residuals extracted from the quasi binomial GLM: proportion Canadian range protected ~
COSEWIC status + proportion range in Canada.

Q5) Conservation research effort and insights from range-wide studies on at-risk plants

Our literature searches yielded 657 peer-reviewed, conservation-relevant studies on the 189 plant

species that are at-risk in Canada and whose geographic distribution we can assess (Table 1b).

Almost half (44%) of the 189 species had not been studied in peer-reviewed work that could

inform conservation (Fig. 6a). Though this does not preclude the existence of studies in the non-

refereed literature or journals not indexed on Web of Science, it suggests that the 'best biological

information' is sparse for many taxa. Compared to species with more than 20% of their range in

Canada, species that only occur peripherally in Canada did not differ in the likelihood that they

had been studied ($\chi^2_{df=1}$ = 1.73, P = 0.19) or in the number of studies ($\chi^2_{df=1}$ = 0.21, P = 0.88;

Fig. 6a). Comparisons remained non-significant if we used only taxa with digitized range maps

or 10% peripheral criterion (Table S3).

Of 657 conservation-relevant studies, only 187 included Canadian populations. Less than half

(42%) of plant species at-risk in Canada had been studied in Canada (Fig. 6b). Peripheral and

non-peripheral species did not differ in the likelihood that they had been studied in Canada

($\chi^2_{df=1}$ = 0.02, P = 0.90), nor in the number of studies that included Canadian populations ($\chi^2_{df=1}$

= 2.72, P = 0.099; Fig. 6b). However, if one considers only taxa with a digitized range map or a

stricter definition of peripheral (<10% range in Canada), peripheral species had fewer studies

that included Canadian populations than non-peripheral species (Table S3).

Figure 6. Distribution of conservation-relevant research on plants at-risk in Canada. (A) distribution of all 657 studies that could inform conservation; (B) 187 studies from (A) that included data from wild Canadian populations. Left panels = peripheral species (< 20% of range in Canada); right panels = non-peripheral species. Black bars show species with no peer-reviewed studies.

Most (162 of 189) plant species at risk have less than half their range in Canada, such that

understanding Canadian populations in the context of their wider geographic range could inform

conservation. But only 6% (34 of 536) of studies on these species included both Canadian and

USA populations, and these 34 studies covered only 20 (11%) of the 162 species. Studies that

performed range-wide sampling provided unique insights into conserving peripheral populations

(Table 2). These include whether edge populations differ from core populations genetically,

demographically, or in key traits or habitat affinity. For instance, populations of Deerberry

(*Vaccinium stamineum*) decreased in size and frequency toward the species northern range edge

in Canada, but were nevertheless as productive and genetically diverse as core populations, and

showed evidence of local adaptation and high dispersal ability (Yakimowski and Eckert 2007,

2008). Thus the demographic and genetic value of these populations was not predicted by their

peripherality, size, or spatial isolation.

Table 2. Studies with wide geographic coverage can shed light on the conservation needs and
value of peripheral populations. Key findings are taken from studies of plant taxa deemed at-risk
in Canada that sampled both Canadian and USA populations. All taxa are peripheral in Canada.

Response		
Finding for edge populations	Species	Reference
Genetic uniqueness		
Genetically differentiated from core populations (neutral variation)	Branched bartonia	Ciotir et al. 2013
	Green dragon	Boles et al. 2000
	Spalding's Campion	Lesica et al. 2016
Greatest population differentiation	Butternut	Hoban et al. 2010
Genetic diversity		
Not lower neutral genetic diversity than core populations	Heartleaf plantain	Mymudes and Les 1993
	Deerberry	Yakimowski and Eckert 2008
	Green dragon	Boles et al. 2000
	Golden paintbrush	Godt et al. 2005
	Cucumber tree	Budd et al. 2015
Population demography		
Not smaller than core populations	Golden paintbrush	Godt et al. 2005
Smaller & more isolated than core populations	Deerberry	Yakimowski and Eckert 2007
Performance & ecology		
Similar habitat to core populations	Small whorled pogonia orchid	Mehrhoff 1989
Lowest seed viability & distinct dormancy patterns	Green dragon	Yang et al. 1999
Higher seed mass & equal sexual reproduction / productivity as core populations	Deerberry	Yakimowski and Eckert 2007

Branched bartonia = *Bartonia paniculata ssp. paniculata*; Butternut = *Juglans cinerea;*
Cucumber tree = *Magnolia acuminata*; Deerberry = *Vaccinium stamineum*; Golden paintbrush =
Castilleja levisecta; Green dragon = *Arisaema dracontium*; North American ginseng = *Panax*
quinquefolius; Small whorled pogonia orchid = *Isotria medeleoides*; Heartleaf plantain =
Plantago cordata; Spalding's campion = *Silene spaldingii*.

DISCUSSION

Our results show that conservation of plants in Canada is fundamentally the conservation of

range-edge populations. Three quarters of nationally at-risk plant taxa only occur in Canada at

the northernmost 20% or less of their global range, in line with earlier estimates for all at-risk

taxa combined (Gibson et al. 2009). We do not have range maps for the thousands of plants in

Canada that have not been assessed by COSEWIC, so cannot directly test whether at-risk plants

are more peripheral than average. However, taxa with <20% of their range in Canada had a

greater disparity between their Canadian vs. global threat ranking than taxa with more of their

range in Canada (Fig. 2), and the most imperilled taxa were significantly more peripheral

(smaller proportion of their range in Canada) than less imperilled taxa (Fig. 2C), suggesting a

real relationship between occurring as range-edge populations and being nationally at-risk.

Range-edge taxa could be more nationally threatened because they have smaller ranges and

therefore fewer individuals in Canada, or because their Canadian populations are

disproportionately threatened. Our results suggest that both are true. Smaller range area in

Canada was associated with higher COSEWIC threat status (Fig. 3B), but peripherality was

associated with higher threat even after accounting for Canadian range area (Fig. S3).

Endangered taxa are both the most imperilled and most peripheral group (Fig. 3C) and had

significantly higher human population densities in their Canadian range (Fig. 4), although

peripherality was not associated with human density after accounting for threat rank. We did not

test effects of human activity not associated with high population density (e.g., agriculture), but

overall human activity is also highest in southern Canada (Coristine and Kerr 2011), where most-

at-risk and almost all peripheral taxa occur (Fig. 1). Thus, higher national threat ranks for

peripheral taxa probably reflect real increased risk per population.

Stewarding Canada's future biodiversity clearly requires an informed policy on conserving

peripheral populations. Not only are most at-risk flora range-edge populations, but these

populations are geographically poised to initiate northward range shifts under climate warming

(Gibson et al. 2009), and so will make up more of Canada's flora in the future. Unfortunately,

conservation of peripheral taxa has been debated in the absence of much relevant scientific

evidence. Less than half the plant species with at-risk populations in Canada have been studied in

Canada in a way that could guide their conservation. While this could reflect difficulty in

obtaining permits or adequate sample sizes, taxonomic bias is likely. For example, one bird

species that is both peripheral and at-risk in Canada had almost 50 studies that included

Canadian populations (Marbeled murrelet; Web of Science search May 2019), far more than any

peripheral at-risk plant species (Fig. 6B).

The few conservation-relevant studies that include both Canadian and US populations illustrate

the value of studying peripheral populations directly and in a broad geographic context.

However, most of these studies have assessed neutral genetic diversity and population structure

(Table 2). Conservation would particularly benefit from studies of characteristics important for

long-term persistence and range expansion, such as habitat preferences, population demography

and dispersal ability (Schemske et al. 1994). Future genetic work could move beyond neutral

variation to evaluating the adaptive diversity likely to be important in responding to

environmental change (Shaw and Etterson 2012), and local adaptation through which range-edge

populations may contribute uniquely to species' biodiversity (Yeaman et al. 2016). Whether

researchers will close these knowledge gaps depends partially on how government agencies

incentivise (i.e. fund) and remove barriers to (i.e. permit) research on at-risk peripheral

populations. Unfortunately, the "peripherality issue" is not currently highlighted in federal

programs that fund species-at-risk research in Canada (e.g. Government of Canada 2019).

We hope that exposing this research need inspires future work on at-risk edge populations, but

recognize that amassing this work will take time, and that some of the most informative types of

study, e.g., large reciprocal transplants, will be impossible with endangered taxa. In the

meantime, we have a potentially under-used body of research that could inform Canadian

conservation: the already extensive theory and empirical research on species range edges (Sexton

et al. 2009; Pironon et al. 2017). While this research clearly shows that edge populations can

vary significantly from one another in demography (Sagarin et al. 2006) and degree of adaptation

(Hargreaves and Eckert 2019), it also reveals broad scale patterns that can be predictive, e.g., that

poleward range edges are often dispersal limited whereas high-elevation edge populations are

often demographic sinks (Halbritter et al. 2013; Hargreaves et al. 2014), and suggests novel

conservation strategies, e.g., increasing gene-flow among isolated range-edge populations to

spread broadly beneficial alleles (Sexton et al. 2011; Hargreaves and Eckert 2019). For countries

like Canada whose biodiversity is disproportionately comprised of range-edge populations,

leveraging this literature could help meet commitments to protecting current and future

biodiversity.

ACKNOWLEDGEMENTS

We thank Christie Whelan (Dept Fisheries and Oceans), Gary Allen & Mark Major (Parks

Canada), Joanna Freeland (Trent University), Jeanette Whitton (UBC), and Karen Samis

(Queen's U) for providing insight into Canadian conservation policy and practise. We also thank

Tim Elrick (McGill University) for his help with geographical data. Funding was provided by

Natural Sciences and Engineering Research Council of Canada (Discovery grants to ALH and

CGE), by E. Gordon Edwards and Vivian Astroff Science Undergraduate Research Awards to

PC, and by William E. Bembridge Science Undergraduate Research Award to SKN.

COMPETING INTERESTS STATEMENT

Authors have no competing interests to declare.

REFERENCES

Anderson, J., and Wadgymar, S. 2020. Climate change disrupts local adaptation and favours
upslope migration. Ecol. Lett. **23**(1): 181-192.
Bivand, R., and Rundel, C. 2018. rgeos: Interface to Geometry Engine - Open Source ('GEOS').
Available from https://cran.r-project.org/package=rgeos.
Boles, R.L., Lovett-Doust, J., and Lovett-Doust, L. 2000. Population genetic structure in green
dragon (Arisaema dracontium, Araceae). Can. J. Bot. **77**(10): 1401–1410.
doi:10.1139/b99-089.
Breheny, P., and Burchett, W. 2017. Visualization of Regression Models Using visreg.
Brito, D., Ambal, R.G., Brooks, T., Silva, N.D., Foster, M., Hao, W., Hilton-Taylor, C., Paglia,
A., Rodríguez, J.P., and Rodríguez, J.V. 2010. How similar are national red lists and the
IUCN Red List? Biol. Conserv. **143**(5): 1154–1158. doi:10.1016/j.biocon.2010.02.015.
Brown, J.H., Stevens, G.C., and Kaufman, D.M. 1996. The geographic range: size, shape, and
internal structure. Annu. Rev. Ecol. Syst. **27**: 597–623.
doi:10.1146/annurev.ecolsys.27.1.597.
Budd, C., Zimmer, E., and Freeland, J.R. 2015. Conservation genetics of Magnolia acuminata, an
endangered species in Canada: Can genetic diversity be maintained in fragmented,
peripheral populations? Conserv. Genet. **16**(6): 1359–1373. doi:10.1007/s10592-015-
0746-9.
Bunnell, F.L., Campbell, R.W., and Squires, K.A. 2004. Conservation priorities for peripheral
species: the example of British Columbia. Can. J. For. Res. **34**(11): 2240–2247.
doi:10.1139/x04-102.
Cameron, V., and Hargreaves, A.L. 2020. Spatial distribution and hotspots of mammals in
Canada. bioRxiv: 2020.01.29.925461. doi:10.1101/2020.01.29.925461.
CCEA. 2016. Conservation Areas Reporting and Tracking System Procedures Manual and
Database Schema Implementation version - 2016. : 43.
Chen, I.-C., Hill, J.K., Ohlemüller, R., Roy, D.B., and Thomas, C.D. 2011. Rapid range shifts of
species associated with high levels of climate warming. Science **333**(6045): 1024–6.
doi:10.1126/science.1206432.
Ciotir, C., Yesson, C., and Freeland, J. 2013. The evolutionary history and conservation value of
disjunct Bartonia paniculata subsp paniculata (Branched Bartonia) populations in Canada.
Botany **91**(9): 605–613. doi:10.1139/cjb-2013-0063.
Coristine, L.E., Jacob, A.L., Schuster, R., Ford, A., Woodley, S., Shiffman, D.S., Orihel, D.,
Baron, N.E., Bennett, N.J., Venter, O., Bittick, S.J., Favaro, B., Dey, C., Otto, S.P.,
Polfus, J.L., Nowlan, L., and Palen, W.J. 2018. Informing Canada's commitment to
biodiversity conservation: A science-based framework to help guide protected areas
designation through Target 1 and beyond. Facets **3**(1): 531–562. doi:10.1139/facets-
2017-0102.
Coristine, L.E., and Kerr, J.T. 2011. Habitat loss, climate change, and emerging conservation
challenges in Canada. Can. J. Zool. **89**(5): 435–451. doi:10.1139/z11-023.
Eckert, C.G., Samis, K.E., and Lougheed, S.C. 2008. Genetic variation across species'
geographical ranges: the central–marginal hypothesis and beyond. Mol. Ecol. **17**(5):
1170–1188. doi:10.1111/j.1365-294X.2007.03659.x.

Ellstrand, N.C., and Elam, D.R. 2003. Population Genetic Consequences of Small Population
Size: Implications for Plant Conservation. Annu. Rev. Ecol. Syst. **24**(1): 217–242.
doi:10.1146/annurev.es.24.110193.001245.
Environment and Climate Change Canada. 2017. COSEWIC wildlife species assessment:
candidate wildlife species. Available from www.canada.ca/en/environment-climate-
change/services/committee-status-endangered-wildlife/wildlife-species-assessment-
process-categories-guidelines/candidate.html [accessed 18 November 2019].
Freeman, B.G., Lee-Yaw, J.A., Sunday, J.M., and Hargreaves, A.L. 2018. Expanding, shifting
and shrinking: The impact of global warming on species' elevational distributions. Glob.
Ecol. Biogeogr. **27**(11): 1268–1276. doi:10.1111/geb.12774.
Gibson, S.Y., Van der Marel, R.C., and Starzomski, B.M. 2009. Climate change and
conservation of leading-edge peripheral populations. Conserv. Biol. **23**(6): 1369–1373.
doi:10.1111/j.1523-1739.2009.01375.x.
Glass, W.R., Corkum, L.D., and Mandrak, N.E. 2017. Living on the edge: Traits of freshwater
fish species at risk in Canada. Aquat. Conserv. Mar. Freshw. Ecosyst. **27**(5): 938–945.
doi:10.1002/aqc.2781.
Godt, M.J.W., Caplow, F., and Hamrick, J.L. 2005. Allozyme diversity in the federally
threatened golden paintbrush, Castilleja levisecta (Scrophulariaceae). Conserv. Genet.
6(1): 87–99. doi:10.1007/s10592-004-7746-5.
Government of Canada. 2019. Canada Nature Fund. Available from
https://www.canada.ca/en/environment-climate-change/services/nature-legacy/fund.html
[accessed 16 June 2019].
Halbritter, A.H., Alexander, J.M., Edwards, P.J., and Billeter, R. 2013. How comparable are
species distributions along elevational and latitudinal climate gradients? Glob. Ecol.
Biogeogr. **22**(11): 1228–1237. doi:10.1111/geb.12066.
Hampe, A., and Petit, R.J. 2005. Conserving biodiversity under climate change: The rear edge
matters. Ecol. Lett. **8**(5): 461–467. doi:10.1111/j.1461-0248.2005.00739.x.
Hargreaves, A.L., Bailey, S.F., and Laird, R.A. 2015. Fitness declines towards range limits and
local adaptation to climate affect dispersal evolution during climate-induced range shifts.
J. Evol. Biol. **28**(8): 1489–1501. doi:10.1111/jeb.12669.
Hargreaves, A.L., and Eckert, C.G. 2019. Local adaptation primes cold-edge populations for
range expansion but not warming-induced range shifts. Ecol. Lett. **22**(1): 78–88.
doi:10.1111/ele.13169.
Hargreaves, A.L., Samis, K.E., and Eckert, C.G. 2014. Are Species' Range Limits Simply Niche
Limits Writ Large? A Review of Transplant Experiments beyond the Range. Am. Nat.
183(2): 157–173. The University of Chicago PressThe American Society of Naturalists.
doi:10.1086/674525.
Hengeveld, R., and Haeck, J. 1982. The distribution of abundance. J. Biogeogr. **9**(4): 303–316.
Hoban, S.M., Borkowski, D.S., Brosi, S.L., McCleary, T.S., Thompson, L.M., McLachlan, J.S.,
Pereira, M.A., Schlarbaum, S.E., and Romero- Severson, J. 2010. Range-wide
distribution of genetic diversity in the North American tree Juglans cinerea: a product of
range shifts, not ecological marginality or recent population decline. Mol. Ecol. **19**(22):
4876–4891. doi:10.1111/j.1365-294X.2010.04834.x.

Hunter, M.I., and Hutchinson, A. 1994. The Virtues and Shortcomings of Parochialism:
Conserving Species That Are Locally Rare, but Globally Common. Conserv. Biol. **8**(4):
1163–1165.
Komonen, A. 2007. Are we conserving peripheral populations? An analysis of range structure of
longhorn beetles (Coleoptera: Cerambycidae) in Finland. J. Insect Conserv. **11**(3): 281–
285. doi:10.1007/s10841-006-9043-8.
Lee-Yaw, J.A., Kharouba, H.M., Bontrager, M., Mahony, C., Csergő, A.M., Noreen, A.M.E., Li,
Q., Schuster, R., and Angert, A.L. 2016. A synthesis of transplant experiments and
ecological niche models suggests that range limits are often niche limits. Ecol. Lett.
19(6): 710–722. doi:10.1111/ele.12604.
Lenth, R.V. 2016. Least-Squares Means: The R Package lsmeans. doi:10.18637/jss.v069.i01.
Lesica, P., Adams, B., and Smith, C.T. 2016. Can physiographic regions substitute for
genetically-determined conservation units? A case study with the threatened plant, Silene
spaldingii. Conserv. Genet. **17**(5): 1041–1054. doi:10.1007/s10592-016-0842-5.
Lesica, P., and Allendorf, F.W. 1995. When Are Peripheral Populations Valuable for
Conservation? Conserv. Biol. **9**(4): 753–760.
McKie, D. 2016. David McKie | Your home for data journalism. Available from
http://www.davidmckie.com/ [accessed 22 December 2018].
MDEL. 2016. Registre des aires protégées au Québec.
de Medeiros, C.M., Hernández-Lambraño, R.E., Ribeiro, K.A.F., and Sánchez Agudo, J.Á. 2018.
Living on the edge: do central and marginal populations of plants differ in habitat
suitability? Plant Ecol. **219**(9): 1029–1043. doi:10.1007/s11258-018-0855-x.
Mehrhoff, L.A. 1989. Reproductive vigor and environmental factors in populations of an
endangered North American orchid, Isotria medeoloides (Pursh) rafinesque. Biol.
Conserv. **47**(4): 281–296. doi:10.1016/0006-3207(89)90071-2.
Mymudes, M.S., and Les, D.H. 1993. Morphological and Genetic Variability in Plantago cordata
(Plantaginaceae), a Threatened Aquatic Plant. Am. J. Bot. **80**(3): 351–359.
doi:10.2307/2445359.
NatureServe. 2018. NatureServe Explorer: NatureServe Conservation Status. Available from
http://explorer.natureserve.org/ranking.htm [accessed 17 November 2019].
Phillips, B.L., Brown, G.P., and Shine, R. 2010. Evolutionarily accelerated invasions: The rate of
dispersal evolves upwards during the range advance of cane toads. J. Evol. Biol. **23**(12):
2595–2601. doi:10.1111/j.1420-9101.2010.02118.x.
Pironon, S., Papuga, G., Villellas, J., Angert, A.L., García, M.B., and Thompson, J.D. 2017.
Geographic variation in genetic and demographic performance: new insights from an old
biogeographical paradigm. Biol. Rev. **92**(4): 1877–1909. doi:10.1111/brv.12313.
Pironon, S., Villellas, J., Morris, W.F., Doak, D.F., and García, M.B. 2015. Do geographic,
climatic or historical ranges differentiate the performance of central versus peripheral
populations? Glob. Ecol. Biogeogr. **24**(6): 611–620. doi:10.1111/geb.12263.
QGIS Development Team. 2018. QGIS Geographic Information System. Available from
http://qgis.osgeo.org.
R Core Team. 2017. R: A language and environment for statistical computing. Vienna. Available
from https://www.r-project.org/.

692 Raven, P. 1987. Scope of the plant conservation problem world-wide in Bramwell D, editor. *In*
693    Proceedings of an International Conference on botanic gardens and the world
694    conservation strategy. London: Academic Press., Las Palmas de Gran Canaria.
695 Sagarin, R.D., Gaines, S.D., and Gaylord, B. 2006. Moving beyond assumptions to understand
696    abundance distributions across the ranges of species. Trends Ecol. Evol. **21**(9): 524–530.
697    doi:10.1016/j.tree.2006.06.008.
698 SARA. 2002. Species at Risk Act.
699 Schemske, D.W., Husband, B.C., Ruckelshaus, M.H., Goodwillie, C., Parker, I.M., and Bishop,
700    J.G. 1994. Evaluating approaches to the conservation of rare and endangered plants.
701    Ecology **75**(3): 584–606.
702 Sexton, J.P., McIntyre, P.J., Angert, A.L., and Rice, K.J. 2009. Evolution and Ecology of Species
703    Range Limits. Annu. Rev. Ecol. Evol. Syst. **40**(1): 415–436.
704    doi:10.1146/annurev.ecolsys.110308.120317.
705 Sexton, J.P., Strauss, S.Y., and Rice, K.J. 2011. Gene flow increases fitness at the warm edge of
706    a species' range. Proc. Natl. Acad. Sci. **108**(28): 11704–11709.
707    doi:10.1073/pnas.1100404108.
708 Shaw, R.G., and Etterson, J.R. 2012. Rapid climate change and the rate of adaptation: insight
709    from experimental quantitative genetics. New Phytol. **195**(4): 752–765.
710    doi:10.1111/j.1469-8137.2012.04230.x.
711 Statistics Canada. 2011. Census Dictionary.
712 Statistics Canada. 2016. Geographic Attribute File, Reference Guide.
713 Steen, D.A., and Barrett, K. 2015. Should states in the USA value species at the edge of their
714    geographic range? J. Wildl. Manag. **79**(6): 872–876. doi:10.1002/jwmg.897.
715 Thakur, M., Schättin, E.W., and McShea, W.J. 2018. Globally common, locally rare: revisiting
716    disregarded genetic diversity for conservation planning of widespread species. Biodivers.
717    Conserv. **27**(11): 1–5. doi:10.1007/s10531-018-1579-x.
718 Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C.,
719    Erasmus, B.F.N., de Siqueira, M.F., Grainger, A., Hannah, L., Hughes, L., Huntley, B.,
720    van Jaarsveld, A.S., Midgley, G.F., Miles, L., Ortega-Huerta, M.A., Townsend Peterson,
721    A., Phillips, O.L., and Williams, S.E. 2004. Extinction risk from climate change. Nature
722    **427**(6970): 145–148. doi:10.1038/nature02121.
723 Usery, E.L., and Seong, J.-C. 2000. A Comparison of Equal-Area Map Projections for Regional
724    and Global Raster Data. PhD Thesis. Available from http://carto-
725    research.er.usgs.gov/projection/pdf/nmdrs.usery.prn.pdf.
726 Villellas, J., Morris, W.F., and Garcia, M.B. 2013. Variation in stochastic demography between
727    and within central and peripheral regions in a widespread short-lived herb. Ecology
728    **94**(6): 1378–1388. doi:10.1890/12-1163.1.
729 Yakimowski, S.B., and Eckert, C.G. 2007. Threatened peripheral populations in context:
730    Geographical variation in population frequency and size and sexual reproduction in a
731    clonal woody shrub. Conserv. Biol. **21**(3): 811–822. doi:10.1111/j.1523-
732    1739.2007.00684.x.
733 Yakimowski, S.B., and Eckert, C.G. 2008. Populations do not become less genetically diverse or
734    more differentiated towards the northern limit of the geographical range in clonal
735    Vaccinium stamineum (Ericaceae). New Phytol. **180**(2): 534–544. doi:10.1111/j.1469-
736    8137.2008.02582.x.

737   Yang, J., Lovett- Doust, J., and Lovett- Doust, L. 1999. Seed germination patterns in green
738   dragon (Arisaema dracontium, Araceae). Am. J. Bot. **86**(8): 1160–1167.
739   doi:10.2307/2656980.
740   Yeaman, S., Hodgins, K.A., Lotterhos, K.E., Suren, H., Nadeau, S., Degner, J.C., Nurkowski,
741   K.A., Smets, P., Wang, T., Gray, L.K., Liepe, K.J., Hamann, A., Holliday, J.A.,
742   Whitlock, M.C., Rieseberg, L.H., and Aitken, S.N. 2016. Convergent local adaptation to
743   climate in distantly related conifers. Science **353**(6306): 1431–1433.
744   doi:10.1126/science.aaf7812.
745

FIGURES AND TABLES

SUPPLEMENTARY INFORMATION

QUESTION 1

Comparability of NatureServe and COSEWIC ranks

Although Nature Conservancy and COSEWIC ranks use different numbers of categories (five

and four, respectively), both are derived from IUCN criteria. To test whether the two bodies

ranked taxa consistently, we converted COSEWIC ranks to numeric ranks from 4 (not-at-risk) to

1 (endangered); as NatureServe did not give any species a 5 (least threatened) in Canada, both

NatureServe Canadian ranks and numeric COSEWIC ranks varied from 1 to 4. We used a paired

Wilcoxon test to assess whether COSEWIC and NatureServe ranks for Canadian populations

differed overall. NatureServe and COSEWIC ranks did differ significantly, as NatureServe

tended to consider taxa more nationally threatened than COSEWIC (Fig. S1).

766   **Figure S1. (**a) NatureServe tended to consider taxa more threatened than COSEWIC, i.e. more
767   points fall below the diagonal line reference line than above (1 = most threatened; Wilcoxon
768   paired rank test = 924.5, P = 0.0003). (b) This difference reflected higher threat rankings for
769   non-peripheral taxa (Wilcoxon unpaired rank test = 4139, P = 0.016). Centre lines and boxes
770   show the median, 25th and 75th quartiles (R Core Team 2017).
771

However, the systematic difference between NatureServe Canadian ranks and COSEWIC ranks

did not alter conclusions for Q1. If we used COSEWIC ranks instead of NatureServe Canadian

ranks to calculate the discrepancy between global and national conservation priority, peripheral

taxa still have greater discrepancy than non-peripheral taxa (Fig. S2).

Figure S2. The disparity between national and global threat ranks is greater for species
peripheral in Canada, whether one assesses Canadian threat using NatureServe Canadian ranks
(shown in (a) – same data as depicted in Fig 2) or COSEWIC ranks (b: Wilcoxon unpaired test
on whether the disparity between global and Canadian threat ranks differs between peripheral
and non-peripheral taxa: 5877.5, P < 0.0001). Diagonal lines indicate Canadian populations have
the same threat ranking as the taxon globally. Taxa that are peripheral in Canada (left) have a
greater mismatch between their Canadian and global threat ranks (more taxa listed as threatened
in Canada but secure globally) than taxa with >20% of their range in Canada (right).

Changing definition of peripheral

Our main analyses used 202 taxa (Table 1c) and paired Wilcoxon tests to test disparity between

NatureServe Canadian and global rankings among taxa deemed peripheral in Canada (<20% of

global range in Canada, determined from digitized maps or maps and range descriptions) or non-

peripheral. We tested the sensitivity to our definition of 'peripheral' by rerunning analyses using

only taxa with a digitized range map and quantitative assessment of peripherality, and using a

stricter threshold for 'peripheral' of 10% of range in Canada. Conclusions did not change (Table

S1).

Table S1. Reanalysis of Question 1 with different definitions of peripheral. Top row gives

results reported in the main paper.

Definition of peripheral	n taxa total (peripheral)	Wilcoxon W, P
<20% of range in Canada, determined from digitized maps or non-digitized maps & range descriptions	202 (159)	5878, $P < 0.0001$
<20% of range in Canada, determined from digitized maps only	175 (135)	4686, $P < 0.0001$
≤10% of range in Canada, determined from digitized maps only	175 (123)	684, $P < 0.0001$

QUESTION 2

Figure S3. Being more peripheral (having a smaller % of global range) in Canada is associated
with being more nationally threatened even after accounting for absolute range area in Canada
(*Question 2*). QuassiPoisson GLM: numeric COSEWIC status ~ Canadian range area + Canadian
range fraction. Canadian range area is numeric, but is displayed categorically to help visualize
patterns.
807

A colleague who served on COSEWIC's vascular plants subcommittee pointed out that the way

COSEWIC prioritizes and assesses taxa changed once the Species at Risk Act was passed in

2002, potentially reducing prioritization and ranking of peripheral populations. We tested this by

re-running models in Question 2 with a categorical predictor denoting whether species had been

assessed by COSEWIC before 2002. Taxa assessed before or after SARA did not vary in their

global range area, Canadian range area, or fraction of their global range in Canada (Table S2).

Taxa were designated as slightly more at risk after SARA passed (least squared mean numeric

COSEWIC rank = 3.2, where 1=Not at risk and 4=Endangered), than before (mean = 2.5)

potentially indicating more effective prioritization. The effect of range area or percent on

COSEWIC status did not differ between taxa assessed before or after SARA (model with vs.

without interactions: $\chi^2_{df=2}$, $P = 0.50$).

Table S2. Reanalyzing models in Question 2 accounting for whether taxa were assessed before

or after SARA passed in 2002.

Response	Predictors in model	Predictor significance $\chi^2_{df=1}$, P
Global range area	COSEWIC threat rank	7.50, $P = 0.057$
	Before/After SARA	0.01, $P = 0.91$
Canadian range area	COSEWIC threat rank	21.1, $P < 0.001$
	Before/After SARA	3.3, $P = 0.070$
% range in Canada	COSEWIC threat rank	13.8, $P = 0.003$
	Before/After SARA	1.5, $P = 0.16$
COSEWIC threat rank (numeric)	Canadian range area	11.7, $P < 0.0001$
	Percent range in Canada	4.7, $P < 0.031$
	Before/After SARA	17.2, $P < 0.0001$

 QUESTION 5

Changing definition of peripheral

Our main analyses for Q5 used 189 at-risk taxa for which we could quantify or estimate

peripherality, defined as <20% of their global range in Canada (Table 1b). To test whether

conclusions were sensitive to this definition of peripheral, we reran analyses using only at-risk

taxa with a digitized range map and quantitative assessment of peripherality ($n = 166$), and again

using a stricter threshold for 'peripheral' ($\leq$10% of range in Canada). The effect of being

peripheral generally remained non-significant, except for the number of studies that included

Canadian populations (Table S3).

Table S3. Reanalysis of Question 5 with different definitions of peripheral. Note that Q5
considers only at-risk taxa. Cells give $\chi^2_{df=1}$ test statistics for effect of peripheral (yes or no) and
associated *P* values, df = 1 in all cases. Leftmost results column gives results reported in main
manuscript, using all taxa for which peripherality could be quantified or estimated (Table 1b).
837

GLM (error distribution)	Threshold for peripheral (% global range in Canada) Taxa considered N species (peripheral spp)		
	<20% COSEWIC map n = 189 (159)	<20% Digitized map n = 166 (132)	$\leq$10% Digitized map n = 166 (120)
Likelihood studied anywhere (qbinomial)	$\chi^2 = 1.73$ $P = 0.19$	$\chi^2 = 1.71$ $P = 0.19$	$\chi^2 = 0.02$ $P = 0.88$
Studies per species (negative binomial)	$\chi^2 = 0.21$ $P = 0.88$	$\chi^2 = 0.26$ $P = 0.61$	$\chi^2 = 1.10$ $P = 0.29$
Likelihood studied in Canada (qbinomial)	$\chi^2 = 0.02$ $P = 0.90$	$\chi^2 = 0.12$ $P = 0.90$	$\chi^2 = 1.32$ $P = 0.25$
Studies per species in Canada (negative binomial)	$\chi^2 = 2.72$ $P = 0.099$	$\chi^2 = 3.66$ $P = 0.056$	$\chi^2 = 6.97$ $P = 0.008$