Quantifying within-host diversity of H5N1 influenza viruses in humans and

poultry in Cambodia

Louise H. Moncla[1]*, Trevor Bedford[1,2], Philippe Dussart[3], Srey Viseth Horm[3], Sareth Rith[3],

Philippe Buchy[4], Erik A Karlsson[3], Lifeng Li[5,6], Yongmei Liu[5,6], Huachen Zhu[5,6], Yi Guan[5,6],

Thomas C. Friedrich[7,8], Paul F. Horwood[3,9]*

Author affiliations

1. Fred Hutchinson Cancer Research Center, Seattle, Washington, United States.

2. University of Washington, Seattle, Washington, United States.

3. Virology Unit, Institut Pasteur du Cambodge, Institut Pasteur International Network, Phnom

Penh, Cambodia.

4. GlaxoSmithKline, Vaccines R&D, Singapore, Singapore.

5. Joint Influenza Research Centre (SUMC/HKU), Shantou University Medical College,

Shantou, People's Republic of China.

6. State Key Laboratory of Emerging Infectious Diseases/Centre of Influenza Research, School

of Public Health, The University of Hong Kong, Hong Kong, SAR, People's Republic of China.

7. Department of Pathobiological Sciences, University of Wisconsin School of Veterinary

Medicine, Madison, WI, United States.

8. Wisconsin National Primate Research Center, Madison, WI, United States.

9. College of Public Health, Medical and Veterinary Sciences, James Cook University,

Townsville, Australia.

* correspondence: lhmoncla@gmail.com and paul.horwood@jcu.edu.au

Abstract

Avian influenza viruses (AIVs) periodically cross species barriers and infect humans. The

likelihood that an AIV will evolve mammalian transmissibility depends on acquiring and selecting

mutations during spillover, but data from natural infection is limited. We analyze deep

sequencing data from infected humans and domestic ducks in Cambodia to examine how H5N1

viruses evolve during spillover. Overall, viral populations in both species are predominated by

low-frequency (<10%) variation shaped by purifying selection and genetic drift, and half of the

variants detected within-host are never detected on the H5N1 virus phylogeny. However, we do

detect a subset of mutations linked to human receptor binding and replication (PB2 E627K, HA

A150V, and HA Q238L) that arose in multiple, independent humans. PB2 E627K and HA A150V

were also enriched along phylogenetic branches leading to human infections, suggesting that

they are likely human-adaptive. Our data show that H5N1 viruses generate putative human-

adapting mutations during natural spillover infection, many of which are detected at >5%

frequency within-host. However, short infection times, genetic drift, and purifying selection likely

restrict their ability to evolve extensively during a single infection. Applying evolutionary methods

to sequence data, we reveal a detailed view of H5N1 virus adaptive potential, and develop a

foundation for studying host-adaptation in other zoonotic viruses.

Author summary

H5N1 avian influenza viruses can cross species barriers and cause severe disease in humans.

H5N1 viruses currently cannot replicate and transmit efficiently among humans, but animal

infection studies and modeling experiments have suggested that human adaptation may require

only a few mutations. However, data from natural spillover infection has been limited, posing a

challenge for risk assessment. Here, we analyze a unique dataset of deep sequence data from

H5N1 virus-infected humans and domestic ducks in Cambodia. We find that well-known

markers of human receptor binding and replication arise in multiple, independent humans. We

also find that 3 mutations detected within-host are enriched along phylogenetic branches

leading to human infections, suggesting that they are likely human-adapting. However, we also

show that within-host evolution in both humans and ducks are shaped heavily by purifying

selection and genetic drift, and that a large fraction of within-host variation is never detected on

the H5N1 phylogeny. Taken together, our data show that H5N1 viruses do generate human-

adapting mutations during natural infection. However, short infection times, purifying selection,

and genetic drift may severely limit how much H5N1 viruses can evolve during the course of a

single infection.

Introduction

Influenza virus cross-species transmission poses a continual threat to human health. Since

emerging in 1997, H5N1 avian influenza viruses (AIVs) have caused 860 confirmed infections

and 454 deaths in humans[1]. H5N1 viruses naturally circulate in aquatic birds, but some

lineages have integrated into poultry populations. H5N1 viruses are now endemic in domestic

birds in some countries[2–4], and concern remains that continued human infection may one day

facilitate human adaptation.

The likelihood that an AIV will adapt to replicate and transmit among humans depends on

generating and selecting human-adaptive mutations during spillover. Influenza viruses have

high mutation rates[5–8], short generation times[9], and large populations, and rapidly generate

diversity within-host. Laboratory studies using animal models[10–12] show that only 3-5 amino

acid substitutions may be required to render H5N1 viruses mammalian-transmissible[10–12],

and that viral variants present at frequencies as low as 5% may be transmitted by respiratory

droplets[13]. Subsequent modeling studies suggest that within-host dynamics are conducive to

generating human-transmissible viruses, but that these viruses may remain at frequencies too

low for transmission[14,15]. Although these studies offer critical insight for H5N1 virus risk

assessment, it is unclear whether they adequately describe how cross-species transmission

proceeds in nature.

H5N1 virus outbreaks offer rare opportunities to study natural cross-species transmission, but

data are limited. One study of H5N1 virus-infected humans in Vietnam identified mutations

affecting receptor binding, polymerase activity, and interferon antagonism; however, they

remained at low frequencies throughout infection[16]. Recent characterization of H5N1 virus-

infected humans in Indonesia identified novel mutations within-host that enhance polymerase

activity in human cells[17]. Unfortunately, neither of these studies include data from naturally

infected poultry, which would provide a critical comparison for assessing whether infected

humans exhibit signs of adaptive evolution. A small number of studies have examined within-

host diversity in experimentally infected poultry[18–20], but these may not recapitulate the

dynamics of natural infection.

As part of ongoing diagnostic and surveillance effort, the Institut Pasteur du Cambodge collects

and confirms samples from AIV-infected poultry during routine market surveillance, and from

human cases and poultry during AIV outbreaks. Since H5N1 was first detected in Cambodia in

2004, 56 human cases and 58 poultry outbreaks have been confirmed and many more have

gone undetected[21]. Here we analyze previously generated deep sequence data[22] from 8

infected humans and 5 infected domestic ducks collected in Cambodia between 2010 and 2014.

We find that viral populations in both species are dominated by low-frequency variation shaped

by population expansion, purifying selection, and genetic drift. We identify a handful of

mutations in humans linked to improved mammalian replication and transmissibility, two of

which were detected in multiple samples, suggesting that adaptive mutations arise during

natural spillover infection. Although most within-host mutations are not linked to human

infections on the H5N1 virus phylogeny, three mutations identified within-host are enriched on

phylogenetic branches leading to human infections. Our data suggest that known adaptive

mutations do occur in natural H5N1 virus infection, but that a short duration of infection,

randomness, and purifying selection may together limit the evolutionary capacity of these

viruses to evolve extensively during any individual spillover event.

Methods

Viral sample collection

The Institute Pasteur in Cambodia is a World Health Organization H5 Reference Laboratory

(H5RL) and has a mandate to assist the Cambodian Ministry of Health and the Ministry of

Agriculture, Forestry, and Fisheries in conducting investigations into human cases and poultry

outbreaks of H5N1 virus, respectively. Surveillance for human cases of H5N1 virus infection is

conducted through influenza-like-illness, severe acute respiratory illness, and event-based

surveillance in a network of hospitals throughout the country [23]. Poultry outbreaks of H5N1

virus are detected through passive surveillance following reports from farmers and villagers of

livestock illness or death. The H5RL conducts confirmation of H5N1 virus detection and further

characterization (genetic and antigenic) of H5N1 virus strains.

Human subjects and IRB approval

The Cambodian influenza surveillance system is a public health activity managed by the

Ministry of Health in Cambodia and has a standing authorization from the National Ethics

Committee for Human Research. The deep sequence analysis of H5N1 influenza virus from

human samples was approved for this study by the National Ethics Committee for Human

Research (#266NECHR).

RNA isolation and RT-qPCR

RNA was extracted from swab samples using the QIAmp Viral RNA Mini Kit (Qiagen, Valencia,

CA, USA), following manufacturer's guidelines and eluted in buffer AVE. Extracts were tested

for influenza A virus (M-gene)[24] and subtypes H5 (primer sets H5a and H5b), N1, H7, and H9

by using quantitative RT-PCR (qRT-PCR) using assays sourced from the International Reagent

Resource (https://www.internationalreagentresource.org/Home.aspx), as previously

outlined[25]. Only samples with high viral load (≥10^3 copies/µl of extracted viral RNA in buffer

AVE), as assessed by RT-qPCR, were selected for sequence analysis. All samples were

sequenced directly from the original specimen, without passaging in cell culture or eggs.

Information on the samples included in the present analyses are presented in **Table 1**.

cDNA generation and PCR

cDNA was generated using Superscript IV Reverse Transcriptase (Invitrogen, Carlsbad, CA,

USA) and custom influenza primers targeting the conserved ends for whole genome

amplification[26]. The following primers were pooled together in a 1.5 : 0.5 : 2.0 : 1.0 ratio: Uni-

1.5: ACGCGTGATCAGCAAAAGCAGG, Uni-0.5: ACGCGTGATCAGCGAAAGCAGG, Uni-2.0:

ACGCGTGATCAGTAGAAACAAGG, and Uni-1.0: AGCAAAAGCAGG. 1 µl of this primer pool

was added to 1 µl of 10 mM dNTP mix (Invitrogen) and 11 µl of RNA. Contents were briefly

mixed and heated for 5 minutes at 65°C, followed by immediate incubation on ice for at least 1

minute. Next, a second mastermix was made with 4 µl of 5X Superscript IV Buffer, 1 µl of 100

mM DTT, 1 µl of RNaseOut Recombinant RNase Inhibitor, and 1 µl of SuperScript IV Reverse

Transcriptase (200 U/µl) (Invitrogen). 7 µl of mastermix was added to each sample, for a total

volume of 20 µl. This mixture was briefly mixed, incubated at 55°C for 20 minutes, then

inactivated by incubating at 80°C for 10 minutes. Whole genomic amplification of the influenza

virus was conducted using Ex Taq™ Hot Start Version (TaKaRa). Forward primers were Uni-1.5

and Uni-0.5 mixed in a ratio of 3:2, and reverse primer was Uni-2.0. The temperature cycle

parameters were 98°C for 2 min, and then 5 cycles (98°C for 30 seconds, 45°C for 30 seconds,

and 72°C for 3 minutes), followed by 25 cycles (98°C for 30 seconds, 55°C for 30 seconds, and

72°C for 3 minutes).

152 Library preparation and sequencing

For each sample, amplicons were quantified using the Qubit^{TM} dsDNA BR Assay Kit

(Invitrogen), pooled in equimolar concentrations, and fragmented using the NEBNext dsDNA

Fragmentase (New England BioLabs, Ipswich, MA). DNA fragments with the size of 350-700 bp

were separated on an agarose gel during electrophoresis and purified for input into the

NEBNext Ultra DNA Library Prep Kit for Illumina® (New England BioLabs). Prepared libraries

were quantified using KAPA Library Quantification Kits for Illumina® platforms (KAPA

Biosystems) and pooled in equimolar concentrations to a final concentration of 4 nM, and run

using an MiSeq Reagent Kit v2 (Illumina, San Diego, CA) for 500 cycles (2 x 250 bp).

Demultiplexed files were output in FASTQ format.

162 Processing of raw sequence data, mapping, and variant calling

Human reads were removed from raw FASTQ files by mapping to the human reference genome

GRCH38 with bowtie2[27] version 2.3.2 (http://bowtie-bio.sourceforge.net/bowtie2/index.shtml).

Reads that did not map to human genome were output to separate FASTQ files and used for all

subsequent analyses. Illumina data was analyzed using the pipeline described in detail at

https://github.com/lmoncla/illumina_pipeline. Briefly, raw FASTQ files were trimmed using

Trimmomatic[28] (http://www.usadellab.org/cms/?page=trimmomatic), trimming in sliding

windows of 5 base pairs and requiring a minimum Q-score of 30. Reads that were trimmed to a

length of <100 base pairs were discarded. Trimming was performed with the following

command: java -jar Trimmomatic-0.36/trimmomatic-0.36.jar SE input.fastq output.fastq

SLIDINGWINDOW:5:30 MINLEN:100. Trimmed reads were mapped to consensus sequences

previously derived[22] using bowtie2[27] version 2.3.2 (http://bowtie-

bio.sourceforge.net/bowtie2/index.shtml), using the following command: bowtie2 -x

reference_sequence.fasta -U read1.trimmed.fastq,read2.trimmed.fastq -S output.sam --local.

Duplicate reads were removed with Picard (http://broadinstitute.github.io/picard/) with: java -jar

picard.jar MarkDuplicates I=input.sam O=output.sam REMOVE_DUPLICATES=true. Mapped

reads were imported into Geneious (https://www.geneious.com/) for visual inspection and

consensus calling. Consensus sequences were called by reporting the majority base at each

site. For nucleotide sites with <100x coverage, a consensus base was not reported, and was

instead reported as an "N". To avoid issues with mapping to an improper reference sequence,

we then remapped each sample's trimmed FASTQ files to its own consensus sequence. These

bam files were again manually inspected in Geneious, and a final consensus sequence was

called. We were able to generate full-genome data for all samples except for

A/Cambodia/X0128304/2013, for which we were lacked data for PB1. These BAM files were

then exported and converted to mpileup files with samtools[29]

(http://samtools.sourceforge.net/), and within-host variants were called using VarScan[30,31]

(http://varscan.sourceforge.net/). For a variant to be reported, we required the variant site to be

sequenced to a depth of at least 100x with a minimum, mean PHRED quality score of 30, and

for the variant to be detected in both forward and reverse reads at a frequency of at least 1%.

We called variants using the following command: java -jar VarScan.v2.3.9.jar mpileup2snp

input.pileup --min-coverage 100 --min-avg-qual 30 --min-var-freq 0.01 --strand-filter 1 --output-

vcf 1 > output.vcf. VCF files were parsed and annotated with coding region changes using

custom software available here (https://github.com/blab/h5n1-

cambodia/blob/master/scripts/H5N1_vcf_parser.py). All amino acid changes for HA are reported

and plotted using native H5 numbering, including the signal peptide, which is 16 amino acids in

length. For ease of comparison, some amino acid changes are also reported with mature H5

peptide numbering in the manuscript when indicated.

Phylogenetic reconstruction

We downloaded all currently available H5N1 virus genomes from the EpiFlu Database of the

Global Initiative for Sharing All Influenza Data[32,33] (GISAID, https://www.gisaid.org/) and all

currently available full H5N1 virus genomes from the Influenza Research Database (IRD,

http://www.fludb.org)[34] and added consensus genomes from our 5 duck samples and 8

human samples. Sequences and metadata were cleaned and organized using fauna

(https://github.com/nextstrain/fauna), a database system part of the Nextstrain platform.

Sequences were then processed using Nextstrain's augur software[35]

(https://github.com/nextstrain/augur). Sequences were filtered by length to remove short

sequences using the following length filters: PB2: 2100 bp, PB1: 2100 bp, PA: 2000 bp, HA:

1600 bp, NP: 1400 bp, NA: 1270 bp, MP: 900 bp, and NS: 800 bp. We excluded sequences with

sample collection dates prior to 1996, and those for which the host was annotated as laboratory

derived, ferret, or unknown. We also excluded sequences for which the country or geographic

region was unknown. Sequences for each gene were aligned using MAFFT[36], and then

trimmed to the reference sequence. We chose the A/Goose/Guangdong/1/96(H5N1) genome

(GenBank accession numbers: AF144300-AF144307) as the reference genome.

IQTREE[37,38] was then used to infer a maximum likelihood phylogeny, and TreeTime[39] was

used to infer a molecular clock and temporally-resolved phylogeny. Tips which fell outside of 4

standard deviations away from the inferred molecular clock were removed. Finally,

TreeTime[39] was used to infer ancestral sequence states at internal nodes and the geographic

migration history across the phylogeny. We inferred migration among 9 defined geographic

regions, China, Southeast Asia, South Asia, Japan and Korea, West Asia, Africa, Europe, South

America, and North America, as shown by color in **Fig. 1** and **Fig. S2**. Our final trees are

available at https://github.com/blab/h5n1-cambodia/tree/master/data/tree-jsons, and include the

following number of sequences: PB2: 4063, PB1: 3867, PA: 4082, HA: 6431, NP: 4070, NA:

5357, MP: 3940, NS: 3678. Plotting was performed using baltic

(https://github.com/evogytis/baltic).

Tajima's _D_ calculation

Tajima's _D_ was calculated with the following equation:

$$D = \frac{d}{\sqrt{V(d)}} = \frac{\pi - \frac{S}{a_1}}{\sqrt{e_1 S + e_2 S(S-1)}}$$

where:

$e_1 = \dfrac{c_1}{a_1}$ $e_2 = \dfrac{c_2}{a_1^2 + a_2}$

$c_1 = b_1 - \dfrac{1}{a_1}$ $c_2 = b_2 - \dfrac{n+2}{a_1^2 + a_2} + \dfrac{a_2}{a_1^2}$

$b_1 = \dfrac{n+1}{3(n-1)}$ $b_2 = \dfrac{2(n^2 + n + 3)}{9n(n-1)}$

$a_1 = \sum_{i=1}^{n-1} \dfrac{1}{i}$ $a_2 = \sum_{i=1}^{n-1} \dfrac{1}{i^2}$

$\pi = \pi_N$ or π_S as calculated below in "Diversity (π) calculation", and S is the number of

segregating sites, i.e., the number of within-host single nucleotide variants called for a given

sample and coding region. Within-host variants were called as described above, requiring a

minimum coverage of 100x, a minimum frequency of 1%, a minimal base quality score of Q30,

and detection on both forward and reverse reads. For each sample, we treated synonymous

variants and nonsynonymous variants separately, calculating D for nonsynonymous variation as

the difference between π_N and S_N, and D for synonymous variation as the difference between

π_S and S_S. For n, we used the average coverage across the coding region. Values shown in

Fig. 2c represent mean D when values were combined across all human or duck samples. To

calculate the 95% confidence interval, we performed a bootstrap. We resampled our D values

with replacement, 10,000 times, and calculated the mean of the resampled values in each

iteration. We then calculated the 2.5% and 97.5% percentile of these bootstrapped means and

report this as the 95% confidence interval.

Diversity (π) calculation

Within-host variants were called as described above, requiring a minimum coverage of 100x, a

minimum frequency of 1%, a minimal base quality score of Q30, and detection on both forward

and reverse reads. Variants were annotated as nonsynonymous or synonymous. For each

sample and coding region, we computed the average number of pairwise nonsynonymous

pairwise differences per nonsynonymous site (π_N) and the average number of pairwise

synonymous differences per synonymous site and (π_S) with SNPGenie[40,41]

(https://github.com/chasewnelson/SNPGenie). We used the same set of within-host variants as

reported throughout the manuscript (minimum frequency of 1%) for these diversity calculations.

In both **Fig. 3** and **Table 2**, we present the mean π_N (dark colors) or π_S (light colors) when

values were combined across all humans (red bars) or ducks (blue bars). To calculate the

standard error of these estimates, we performed a bootstrap. We resampled our diversity values

with replacement, 10,000 times, and calculated the mean of the resampled values in each

iteration. We then calculated the standard deviation among our sampled means, and report this

as the standard error. Error bars in **Fig. 3** reflect this calculated standard error.

Comparison to functional sites

We used the Sequence Feature Variant Types tool from the Influenza Research

Database[34] to download all currently available annotations for H5 hemagglutinins, N1

neuraminidases, and all subtypes for the remaining gene segments. We then annotated each

within-host SNV identified in our dataset that fell within an annotated region or site. The

complete results of this annotation are available in **Table S1**. We next filtered our annotated

SNVs to include only those located in sites involved in "host-specific" functions or interactions,

i.e., those that are distinct between human and avian hosts. We defined host-specific

functions/interactions as receptor binding, interaction with host cellular machinery, nuclear

import and export, immune antagonism, 5' cap binding, temperature sensitivity, and

glycosylation. We also included sites that have been phenotypically identified as determinants of

transmissibility and virulence. Sites that participate in binding interactions with other viral

subunits or vRNP, conserved active site domains, drug resistance mutations, and epitope sites

were not categorized as host-specific for this analysis. We annotated both synonymous and

nonsynonymous mutations in our dataset, but only highlight nonsynonymous changes in **Fig. 4**

and **Table 3**.

Shared sites permutation test

To test whether human or duck samples shared more polymorphisms than expected by chance,

we performed a permutation test. We first counted the number of variable amino acid sites, n, in

which an SNV altered the coded amino acid, across coding regions and samples. For example,

if two SNVs occurred in the same codon site, we counted this as 1 variable amino acid site.

Next, for each gene and sample, we calculated the number of amino acid sites that were

covered with sufficient sequencing depth that a mutation could have been called using our SNV

calling criteria. To do this, we calculated the length in amino acids of each coding region, L, that

was covered by at least 100 reads. Non-coding regions were not included. For each coding

region and sample, we then simulated the effect of having n variable amino acid sites placed

randomly along the coding region between sites 1 to L, and recorded the site where the

polymorphism was placed. After simulating this for each gene and sample, we counted the

number of sites that were shared between at least 2 human or at least 2 duck samples. This

process was repeated 100,000 times. The number of shared polymorphisms at each iteration

was used to generate a null distribution, as shown in **Fig. 5b**. We calculated p-values as the

number of iterations for which there were at least as many shared sites as observed in our

actual data, divided by 100,000. For the simulations displayed in **Fig. 5c** and **Fig. 5d**, we

wanted to simulate the effect of genomic constraint, meaning that only some fraction of the

genome could tolerate mutation. For these analyses, simulations were done exactly the same,

except that the number of sites at which a mutation could occur was reduced to 70% (**Fig. 5c**)

or 60% (**Fig. 5d**). Code for performing the shared sites permutation test is freely available at

https://github.com/blab/h5n1-cambodia/blob/master/figures/figure-5b-shared-sites-permutation-

test.ipynb.

Reconstruction of host transitions along the phylogeny

We used the phylogenetic trees in **Fig. S2** to infer host transitions along each gene's phylogeny.

As described above, we used TreeTime[39] to reconstruct ancestral nucleotide states at each

internal node and infer amino acid mutations along each branch along these phylogenetic trees.

We then classified host transition mutations along branches that lead to human or avian tips as

follows (**Fig. 6a**). For each branch in the phylogeny, we enumerated all tips descending from

that branch. If all descendent tips were human, we considered this a monophyletic human

clade. If the current branch's ancestral node also led to only human descendants, we labelled

the current branch a "to-human" branch. If a branch leading to a monophyletic human clade had

an ancestral node that included avian and human descendants, then we considered the current

branch an "avian-to-human" branch, and also labelled it as "to-human". All other branches were

considered "to-avian" branches. We did not explicitly allow for human-to-avian branches in this

analysis. Because avian sampling is poor relative to human sampling, and because H5N1 virus

circulation is thought to be maintained by transmission in birds, we chose to only label branches

explicitly leading to human infections as to-human branches. We also reasoned that for

instances in which a human tip appears to be ancestral to an avian clade, this more likely

results from poor avian sampling than from true human-to-avian transmission. Using these

criteria, we then gathered the inferred amino acid mutations that occurred along each branch in

the phylogeny, and counted the number of times they were associated with each type of host

transition. We then queried each SNV detected within-host in our dataset, in both human and

duck samples, to determine the number of host transitions that they occurred on in the

phylogeny, as shown in **Fig. 6b**. To test whether individual mutations were enriched along

branches leading to human infections, we performed Fisher's exact tests comparing the number

of to-avian and to-human transitions along which the mutation was detected vs. the overall

number of to-avian and to-human transitions that were observed along the tree. Mutations that

showed statistically significant enrichment are annotated in **Fig. 6b**.

General availability of analysis software and data

All code used to analyze data and generate figures for this manuscript are publicly available at

https://github.com/blab/h5n1-cambodia. Raw FASTQ files with human reads removed are

available under SRA accession number PRJNA547644, and accessions SRX5984186-

SRX5984198. All reported variant calls and phylogenetic trees are available at

https://github.com/blab/h5n1-cambodia/tree/master/data.

334   Results

Sample selection and dataset information

We analyzed full-genome sequence data from primary, influenza virus-confirmed samples from

infected humans and domestic ducks from Cambodia (**Table 1**). Four domestic duck samples

(pooled organs) were collected as part of poultry outbreak investigations, while one was

collected during live bird market surveillance (pooled throat and cloacal swab)[4]. All human

samples (throat swabs) were collected via event-based surveillance upon admittance to various

hospitals throughout Cambodia[22]. Because of limited sample availability and long storage

times, generating duplicate sequence data for each sample was not possible. We therefore

focused on samples whose viral RNA copy numbers after viral RNA extraction were $\geq 10^3$

copies/µl of buffer as assessed by RT-qPCR (**Table 1**), and whose mean coverage depth

exceeded 100x (**Fig. S1**). We analyzed full genome data for 7 human and 5 duck samples, and

near complete genome data for A/Cambodia/X0128304/2013, for which we lack data from the

PB1 gene.

H5 viruses circulating in Cambodia were exclusively clade 1.1.2[4] until 2013, when a novel

reassortant virus emerged[42]. This reassortant virus expressed a hemagglutinin (HA) and

neuraminidase (NA) from clade 1.1.2, with internal genes from clade 2.3.2.1a[22]. All 2013/2014

samples in our dataset come from this outbreak, while samples collected prior to 2013 are clade

1.1.2 (**Table 1**, **Fig. 1**, and **Fig. S2**). All HA sequences (with the exception of

A/duck/Cambodia/Y0224304/2014, which expresses a divergent HA) derive from the same

lineage that has been circulating in southeast Asia for years (**Fig. 1**). For the internal gene

segments, samples collected between 2010-2012 and samples collected between 2013-2014

fall into distinct parts of the tree, each nested within the diversity of other southeast Asian

viruses (**Fig. S2**). The 2013 reassortant viruses share 4 amino acid substitutions in HA, S123P,

S133A, S155N, and K266R[22] (H5, mature peptide numbering). S133A and S155N have been

linked to improved α-2,6 linked sialic acid binding, independently and in combination with

S123P[43–45]. All samples encode a polybasic cleavage site in HA (XRRKRR) between amino

acids 325-330 (H5, mature peptide numbering), a virulence determinant for H5N1 AIVs[46,47],

and a 20 amino acid deletion in NA. This NA deletion is a well-documented host range

determinant[48–51].

Duck samples are not immediately ancestral to the human samples in our dataset, and they

therefore are unlikely to represent transmission pairs. We therefore treat these samples as

examples of within-host diversity in naturally infected humans and ducks, rather than direct

transmission pairs. With this caveat, we aimed to use this subset of 8 human and 5 duck

samples to determine whether positive selection would promote adaptation in humans. Positive

selection increases the frequency of beneficial variants, and is often identified by tracking

mutations' frequencies over time. While multiple time points were not available in our dataset, all

human samples were collected 5-12 days after reported symptom onset[22]. Animal infection

studies have observed drastic changes in within-host variant frequencies in 3-7 days[11,13],

suggesting that 5-12 days post symptom onset may provide sufficient time for transmitted

diversity to be altered within-host. We reasoned that while we expect positive selection to

promote the emergence of human-adapting mutation in humans, H5N1 viruses should be well-

adapted for replication in ducks, which are a natural host species. We therefore hypothesize to

observe the following patterns: (1) During replication in humans, positive selection should

increase the frequencies of human-adaptive mutations, resulting in elevated rates of

nonsynonymous diversity and a higher proportion of high-frequency variants. In contrast,

viruses in ducks should be fit for replication and be shaped by purifying selection, leading to an

excess of synonymous variation and an excess of low-frequency variants. (2) Viruses in

humans should harbor mutations phenotypically linked to mammalian adaptation. (3) If selection

is strong at a particular site, then viruses in humans should exhibit evidence for convergent

evolution, i.e., the same mutation arising across multiple samples. (4) If human-adaptive

variants arising within humans are present on the H5N1 phylogeny, then they should be more

likely to occur on branches leading to human infections than on branches leading to bird

infections.

Within-host diversity in humans and ducks is dominated by low-frequency variation

We called within-host variants across the genome that were present in ≥1% of sequencing

reads and occurred at a site with a minimum read depth of 100x and a minimum PHRED quality

score of 30 (see Methods for details). All coding region changes are reported using native H5

numbering, including the signal peptide, unless otherwise noted. Most single nucleotide variants

(SNVs) were present at low frequencies (**Fig. 2a**). We identified a total of 206 SNVs in humans

(111 nonsynonymous, 91 synonymous, 4 missense) and 40 in ducks (16 nonsynonymous, 23

synonymous, 1 missense). Human samples had more SNVs than duck samples on average

(mean SNVs per sample: humans = 26 ± 19, ducks = 8 ± 3, p = 2.79 x 10^{-17}, Fisher's exact test),

although the number of SNVs per sample was variable among samples in both species (**Fig.**

S3).

To determine whether humans had more high-frequency variants than ducks, we generated a

site frequency spectrum (**Fig. 2b**). Purifying selection removes new variants from the

population, generating an excess of low-frequency variants, while positive selection promotes

accumulation of high-frequency polymorphisms. Exponential population expansion also leads to

an excess of low-frequency variation. In both humans and ducks, over 80% of variants (both

synonymous and nonsynonymous) were present in <10% of the population, and the distribution

of SNV frequencies were strikingly similar (**Fig. 2b**). In both host species, there is an excess of

low-frequency variation compared to the expectation under a neutral model (no population size

changes or selection), and a deficiency of intermediate and high-frequency variants (**Fig. 2b**,

grey dots and connecting line). Overall, the frequencies of SNVs in humans and ducks were not

statistically different (p=0.11, Mann Whitney U test), and mean SNV frequencies were similar

(mean SNV frequency in human samples = 5.8%, mean in duck samples = 6.6%).

To determine whether the excess of low-frequency variation we observed was better explained

by purifying selection or demography, we summarized the frequency spectrum by calculating

Tajima's D (**Fig. 2c**). Tajima's D measures the difference between the average number of

pairwise differences between a set of sequences (π) with the number of variable sites (S). π is

weighted by variant frequencies, and will be largest when the population has a large number of

high-frequency variants, while S is simply a count of the number of variable sites, and is not

impacted by variant frequencies. Both population expansion and purifying selection should lead

to an excess of low-frequency variation and negative Tajima's D. However, while population

expansion should impact nonsynonymous and synonymous sites equally, purifying selection

should have a greater effect on nonsynonymous variants. If the excess of low-frequency

variation we observed was driven solely by demographic factors, then we expect synonymous

and nonsynonymous sites to have similar Tajima's D values, while purifying selection should

lead to more negative Tajima's D values at nonsynonymous sites. When calculated across the

full genome, Tajima's *D* was similar between humans and ducks, and was comparable when

calculated for synonymous and nonsynonymous sites. Taken together, these data suggest that

in both humans and ducks, viral populations are dominated by low-frequency variation.

Furthermore, this excess of low-frequency variation can be explained by within-host population

expansion.

Purifying selection and genetic drift shape within-host diversity

Comparing nonsynonymous (π_N) and synonymous (π_S) polymorphism in a population is another

common measure for selection that is robust to differences in sequencing coverage depth[52].

An excess of synonymous polymorphism ($\pi_N/\pi_S < 1$) indicates purifying selection, an excess of

nonsynonymous variation ($\pi_N/\pi_S > 1$) suggests positive selection, and approximately equal

rates ($\pi_N/\pi_S \sim 1$) suggest that genetic drift is the predominant force shaping diversity. We

calculated the average number of nonsynonymous and synonymous pairwise differences

between DNA sequences, and normalized these values to the number of synonymous and

nonsynonymous sites. In both species, most genes exhibited $\pi_N < \pi_S$, although there was

substantial variation among samples (**Table 2** and **Fig. 3**). The difference between π_s and π_N

was generally not statistically significant (**Table 2**), with the exception of human M2 ($\pi_N =$

0.00017, $\pi_S = 0$, p = 0.042, paired t-test) and PB1 ($\pi_N = 0.000083$, $\pi_S = 0.00038$, p = 0.049,

paired t-test), which exhibited weak evidence of purifying selection. When diversity estimates

across all genes were combined, both species exhibited $\pi_N/\pi_S < 1$ (**Fig. 3**) (human $\pi_N/\pi_S = 0.36$,

p = 0.0059, unpaired t-test; duck $\pi_N/\pi_S = 0.21$, p = 0.038, unpaired t-test). Genome-wide

diversity was not correlated with days post symptom onset (**Fig. S4a**). Taken together, these

data suggest that H5N1 within-host populations in both humans and ducks are broadly shaped

by weak purifying selection and genetic drift. We do not find evidence for widespread positive

selection in any individual coding region.

SNVs are identified in humans at functionally relevant sites

Influenza phenotypes can be drastically altered by single amino acid changes. We took advantage of the Influenza Research Database[29] Sequence Feature Variant Types tool, a catalogue of amino acids that are critical to protein structure and function, and that have been experimentally linked to functional alteration. We downloaded all available annotations for H5 HAs, N1 NAs, and all subtypes for the remaining proteins, and annotated each mutation in our dataset that fell within an annotated region (**Table S1**). We then filtered these annotated amino acids to include only those located in sites involved in host-specific functions (see Methods for details).

Of the 218 unique, polymorphic amino acid sites in our dataset (including both human and duck samples), we identified 34 nonsynonymous mutations at sites involved in viral replication, receptor binding, virulence, and interaction with host cell machinery (**Fig. 4**). Some sites are explicitly linked to H5N1 virus mammalian adaptation (**Table 3**). PB2 E627K was detected as a minor variant in A/Cambodia/W0112303/2012, and in A/Cambodia/V0417301/2011 at consensus. A lysine at position 627 is a conserved marker of human adaptation[51,53] that enhances H5N1 replication in mammals[11,12,51,54]. A/Cambodia/W0112303/2012 also encoded PB2 D701N at consensus. Curiously, this patient also harbored the reversion mutation, N701D, at low-frequency within-host. An asparagine (N) at PB2 701 enhances viral replication and transmission in mammals[55,56], while an aspartate (D) is commonly identified in birds. We cannot distinguish whether the founding virus harbored an asparagine or aspartate, so our data are consistent with two possibilities: transmission of a virus harboring asparagine and within-host generation of aspartate; or, transmission of a virus with asparate followed by within-host selection but incomplete fixation of asparagine. All other human and avian samples in our dataset encoded the "avian-like" amino acids, glutamate at PB2 627, and aspartate at PB2 701. None of the adaptive polymerase mutations that were recently identified by Welkers et al.[17] in

H5N1 virus-infected humans in Indonesia were present in our samples, nor were any of the

human-adaptive mutations identified in a recent deep mutational scan of PB2[57].

We also identified HA mutations linked to human receptor binding. Two human samples

encoded an HA A150V mutation (134 in mature, H5 peptide numbering, **Fig. 4**). A valine at HA

150 improves α-2,6 linked sialic acid binding in H5N1 viruses[58,59], and was also identified in

H5N1 virus-infected humans in Vietnam[16]. Finally, HA Q238L was detected in

A/Cambodia/V0417301/2011 and A/Cambodia/V0401301/2011. HA 238L (222 in mature, H5

peptide numbering) was shown in H5N1 virus transmission studies to confer a switch from α-2,3

to α-2,6 linked sialic acid binding[11] and mediate transmission[11,12]. An HA Q238R mutation

was identified in A/Cambodia/X0125302/2013, although nothing is known regarding an arganine

(R) at this site.

Mutations annotated as host-specific were not detected at higher frequencies than non-host-

specific mutations (mean frequency for host-specific mutations = 8.2% ± 8.8%, mean frequency

for non-host-specific mutations = 5.2% ± 4.7%, p-value = 0.084, unpaired t-test). Additionally,

the proportion of mutations that were host-specific was not higher in samples from longer

infections (p-value = 0.72, Fisher's exact test, **Fig. S4b**). All 8 human samples harbored at least

1 mutant in a host-specific site. Critically though, the functional impacts of influenza virus

mutations strongly depend on sequence context[60], and we did not phenotypically test these

mutations. We caution that confirming functional impacts for these mutations would require

further study. Still, our data show that putative human-adapting mutations are generated during

natural spillover. Our results also highlight that even mutations that have been predicted to be

strongly beneficial (e.g., PB2 627K and HA 238L) may remain at low frequencies in vivo.

Shared diversity is limited

Each human H5N1 infection is thought to represent a unique avian spillover event. If selection is

strong at a given site in the genome, then mutations may arise at that site independently across

multiple patients. We identified 13 amino acid sites in our dataset that were polymorphic in at

least 2 samples, 4 of which were detected in both species (PB1 371, PA 307, HA 265 and NP

201). Of the 34 unique polymorphic amino acid sites in ducks, 3 sites were shared by at least 2

duck samples; of the 188 unique polymorphic amino acid sites in humans, 9 were shared by at

least 2 human samples (**Fig. 5a**). Two of these shared sites, HA 150 and HA 238, are linked to

human-adapting phenotypes **(Table 3)**. To determine whether the number of shared sites we

observe is more or less than expected by chance, we performed a permutation test. For each

species, we simulated datasets with the same number of sequences and amino acid

polymorphisms as our actual dataset, but assigned each polymorphism to a random amino acid

site. For each iteration, we then counted the number of polymorphic sites that were shared by

≥2 samples. We ran this simulation for 100,000 iterations for each species, and used the

number of shared sites per iteration to generate a null distribution (**Fig. 5b**, colored bars).

Comparison to the observed number of shared sites (3 and 9, dashed lines in **Fig. 5b**),

confirmed that humans share slightly more polymorphisms than expected by chance (p =

0.046), while ducks share significantly more (p = 0.00006).

Viral genomes are highly constrained [61], which could account for the convergence we

observe. Experimental measurements of the distribution of fitness effects in influenza A virus

have estimated that approximately 30% of genome mutations are lethal [61], while estimates

from other RNA viruses suggest that lethal percentage ranges from 20-40% [62]. We repeated

our simulations to restrict the number of amino acid sites that could tolerate a mutation to 70%

or 60%, representing a lethal fraction of 30% or 40%. When 70% of the coding region was

permitted to mutate, ~23% of simulations resulted in ≥9 shared sites in humans (p = 0.23)(**Fig.**

5c), and when 60% of the genome was permitted to mutate, ~39% of simulations resulted in ≥9

shared sites (p = 0.39)(**Fig. 5d**). In contrast, the probability of observing 3 shared sites among

duck samples remained low regardless of genome constraint (70% of genome tolerates

mutation: p = 0.00014; 60% of genome tolerates mutation: p = 0.00028), suggesting a

significant, although low, level of convergence (**Fig. 5c, d**). Taken together, our results suggest

that duck samples share significantly more variants than expected by chance. In humans,

despite the presence of shared polymorphisms with known human-adaptive phenotypes, the

degree of convergence we observe is no more than expected given genome constraint.

Within-host SNVs are not enriched on spillover branches

If within-host mutations are human-adapting, then those mutations should be enriched among

H5N1 viruses that have caused human infections in the past. To test this hypothesis, we

inferred full genome phylogenies using all available full-genome H5N1 viruses from the

EpiFlu[32,33] and IRD[34] databases (**Fig. 1** and **Fig. S2)**, reconstructed ancestral nucleotide

states at each internal node, and inferred amino acid mutations along each branch. We then

classified host transition mutations along branches that led to human or avian tips (**Fig. 6a)**. If a

branch fell within a clade that included only human tips, that branch was labelled as a "to-

human" transition. If a branch led to a human-only clade but its ancestral branch included avian

descendants, this was inferred to be an avian-to-human transition, and was also labelled as "to-

human". All other transitions were labelled "to-avian" (**Fig. 6a,** see Methods for more details).

We then curated the mutations that occurred on each type of host transition, and compared

these counts to the mutations identified within-host in our dataset.

Of the 120 nonsynonymous within-host SNVs we identified in our dataset, 60 (50%) were not

detected on the phylogeny at all. This suggests that many of the mutations generated within-

host are purged from the H5N1 viral population over time. Additionally, because humans are

generally dead-end hosts for H5N1 viruses, even human-adapting variants arising within-host

are likely to be lost due to lack of onward transmission. Of the within-host mutations that were

detected on the phylogeny, most occurred on branches leading to avian infections (**Fig 6b,** blue

bars). However, there were a few exceptions (**Fig 6b,** red bars). Across the phylogeny, we

enumerated a total of 31,939 to-avian transitions, and 2,787 to-human transitions, so that we

expect a 11.46:1 ratio of to-avian transitions relative to to-human transitions. In contrast, PB2

E627K was heavily enriched among human infections, detected on 15 to-avian transitions and

36 to-human transitions ($p = 4.21 \times 10^{-28}$, Fisher's exact test). HA A150V was detected in only

one to-avian transition, but in 8 to-human transitions ($p = 1.46 \times 10^{-8}$, Fisher's exact test), and

HA N198S was detected on 4 to-avian transitions and 3 to-human transitions ($p = 0.014$,

Fisher's exact test). Although nothing is known regarding a serine at HA 198, a lysine at that site

can confer α-2,6-linked sialic acid binding[43,63]. Taken together, these data suggest that the

majority of mutations detected within-host are not associated with human spillover. However,

they agree with selection for human-adapting phenotypes at a small subset of sites (PB2

E627K, HA A150V, HA N198S).

Discussion

Our study utilizes a unique dataset of to quantify H5N1 virus diversity in natural spillover

infections. We establish a set of hypotheses to interrogate whether H5N1 viruses adapt to

humans during natural spillover, and find support for two of them. We detect putative human-

adapting mutations (PB2 E627K, HA A150V, and HA Q238L) during human infection, two of

which arose multiple times (supporting hypothesis 2). PB2 E627K and HA A150V are enriched

along phylogenetic branches leading to human infections, supporting their potential role in

human adaptation (supporting hypothesis 4). However, we also find that population growth,

genetic drift, and weak purifying selection broadly shape viral diversity in both hosts (rejecting

hypothesis 1), and that convergent evolution in human viruses can be explained by genomic

constraint (rejecting hypothesis 3). Together, our data show that during spillover, H5N1 viruses

have the capacity to generate well-known markers of mammalian adaptation in multiple,

independent hosts. However, none of these markers reached high-frequencies within-host. We

speculate that during spillover, short infection times, genetic drift, demography, and purifying

selection may together limit the capacity of H5N1 viruses to evolve extensively during a single

human infection.

Although data from spillovers are limited, our results align with data from Vietnam[16] and

Indonesia[17]. Welkers et al.[17] identified markers of mammalian replication (PB2 627K) and

transmission (HA 220K) in humans, but found that adaptive markers were not widespread.

Welkers et al. also characterized new mutations that improved human replication, suggesting

that there are yet undiscovered pathways for adaptation. Imai et al.[16] characterized SNVs in

H5N1-infected humans that altered viral replication, receptor binding, and interferon

antagonism, but these mutations stayed at low frequencies. Imai et al. also showed that most

within-host variants elicited neutral or deleterious effects on protein function in humans, aligning

with the purifying selection we detect within-host, and the absence of ~50% of within-host

variants in the phylogeny. These findings also agree with predictions by Russell et al.[14], who

hypothesized that H5N1 viruses would generate human-adapting mutations during infection, but

that these mutations would remain at low frequencies and fail to be transmitted.

One unexpected result is that mutations hypothesized to be strongly beneficial, like PB2 627K

and HA 238L, remained low-frequency during infection. These mutations could have arisen late

in infection or been linked to deleterious mutations. Additionally, epistasis is crucial to influenza

virus evolution, and mutations that promote human adaptation in one background may not be

well-tolerated in others. PB2 E627K is widespread among clade 2.2.1 H5N1 viruses, but only

sparsely detected in other H5N1 clades. Soh et al.[57] recently uncovered strongly human-

adapting PB2 mutations that are rare in nature, likely because they are inaccessible via single

site mutations. Genetic background plays a vital role in determining how AIVs evolve, and may

at least partially explain our findings. Importantly, our study involves a small number of samples

from a single geographic location, and two H5N1 virus clades. Continued characterization of

H5N1 virus spillover in other clades is necessary to define whether our observations are

generalizable across H5N1 virus outbreaks.

An important caveat of our study is that the human and duck samples described likely do not

represent transmission pairs. Although the samples analyzed in this study descend from the

same HA lineage (with the exception of A/duck/Cambodia/Y0224304/2014), the duck samples

are not phylogenetically ancestral to the human samples in this dataset (**Fig. 1** and **Fig. S2**),

and most likely were not the source of the human infections. We therefore caution that each

sample in this dataset merely represents an example of within-host diversity in a naturally

infected host, rather than a before and after snapshot of individual cross-species transmission

events.

Assessing zoonotic risk is critical but challenging. By quantifying patterns of within-host

diversity, identifying mutations at adaptive sites, measuring convergent evolution, and

comparing within-host diversity to long-term evolution, we can assemble a nuanced

understanding of AIV evolution. These methods provide a foundation for understanding cross-

species transmission that can readily be applied to other avian influenza virus datasets, as well

as newly emerging zoonotic viruses.

References

1. Organization WH. Cumulative number of confirmed human cases for avian influenza
A(H5N1) reported to WHO, 2003-2018.

2. Chen H, Smith GJD, Li KS, Wang J, Fan XH, Rayner JM, et al. Establishment of multiple
sublineages of H5N1 influenza virus in Asia: Implications for pandemic control. Proceedings
of the National Academy of Sciences. 2006; doi:10.1073/pnas.0511120103

3. Nguyen DT, Jang Y, Nguyen TD, Jones J, Shepard SS, Yang H, et al. Shifting Clade
Distribution, Reassortment, and Emergence of New Subtypes of Highly Pathogenic Avian
Influenza A(H5) Viruses Collected from Vietnamese Poultry from 2012 to 2015. J Virol.
2017; doi:10.1128/JVI.01708-16

4. Horm SV, Tarantola A, Rith S, Ly S, Gambaretti J, Duong V, et al. Intense circulation of
A/H5N1 and other avian influenza viruses in Cambodian live-bird markets with serological
evidence of sub-clinical human infections. Emerg Microbes Infect. 2016;
doi:10.1038/emi.2016.69

5. Nobusawa E, Sato K. Comparison of the Mutation Rates of Human Influenza A and B
Viruses. J Virol. 2006;80: 3675–3678.

6. Parvin JD, Moscona A, Pan WT, Leider JM, Palese P. Measurement of the mutation rates
of animal viruses: influenza A virus and poliovirus type 1. J Virol. 1986;59: 377–383.

7. Pauly MD, Procario MC, Lauring AS. A novel twelve class fluctuation test reveals higher
than expected mutation rates for influenza A viruses. Elife. 2017;6. doi:10.7554/eLife.26437

8. Suárez P, Valcárcel J, Ortín J. Heterogeneity of the mutation rates of influenza A viruses:
isolation of mutator mutants. J Virol. 1992;66: 2491–2494.

9. Baccam P, Beauchemin C, Macken CA, Hayden FG, Perelson AS. Kinetics of influenza A
virus infection in humans. J Virol. 2006;80: 7590–7599.

10. Imai M, Watanabe T, Hatta M, Das SC, Ozawa M, Shinya K, et al. Experimental adaptation
of an influenza H5 HA confers respiratory droplet transmission to a reassortant H5
HA/H1N1 virus in ferrets. Nature. 2012;486: 420–428.

11.   Linster M, van Boheemen S, de Graaf M, Schrauwen EJA, Lexmond P, Mänz B, et al.
Identification, Characterization, and Natural Selection of Mutations Driving Airborne
Transmission of A/H5N1 Virus. Cell. 2014;157: 329–339.

12.   Herfst S, Schrauwen EJA, Linster M, Chutinimitkul S, de Wit E, Munster VJ, et al. Airborne
transmission of influenza A/H5N1 virus between ferrets. Science. 2012;336: 1534–1541.

13.   Wilker PR, Dinis JM, Starrett G, Imai M, Hatta M, Nelson CW, et al. Selection on
haemagglutinin imposes a bottleneck during mammalian transmission of reassortant H5N1
influenza viruses. Nat Commun. 2013;4: 2636.

14.   Russell CA, Fonville JM, Brown AEX, Burke DF, Smith DL, James SL, et al. The Potential
for Respiratory Droplet–Transmissible A/H5N1 Influenza Virus to Evolve in a Mammalian
Host. Science. 2012;336: 1541–1547.

15.   Sigal D, Reid JNS, Wahl LM. Effects of Transmission Bottlenecks on the Diversity of
Influenza A Virus. Genetics. 2018;210: 1075–1088.

16.   Imai H, Dinis JM, Zhong G, Moncla LH, Lopes TJS, McBride R, et al. Diversity of Influenza
A(H5N1) Viruses in Infected Humans, Northern Vietnam, 2004–2010. Emerg Infect Dis.
2018;24: 1128–1238.

17.   Welkers MRA, Pawestri HA, Fonville JM, Sampurno OD, Pater M, Holwerda M, et al.
Genetic diversity and host adaptation of avian H5N1 influenza viruses during human
infection. Emerg Microbes Infect. 2019;8: 262–271.

18.   Milani A, Fusaro A, Zamperin G, Bonfante F, Mancin M, Mastrorilli E, et al. Viral population
diversity in vaccinated poultry host infected with H5N1 highly pathogenic avian influenza
virus. Int J Infect Dis. 2016;53: 104.

19.   Iqbal M, Xiao H, Baillie G, Warry A, Essen SC, Londt B, et al. Within-host variation of avian
influenza viruses. Philos Trans R Soc Lond B Biol Sci. 2009;364: 2739–2747.

20.   Gutiérrez RA, Viari A, Godelle B, Buchy P. Biased mutational pattern and quasispecies
hypothesis in H5N1 virus. Infect Genet Evol. 2013;15: 69–76.

21.   Suttie A, Karlsson EA, Deng Y-M, Hurt AC, Greenhill AR, Barr IG, et al. Avian influenza in
the Greater Mekong Subregion, 2003-2018. Infect Genet Evol. 2019;74: 103920.

22.   Rith S, Davis CT, Duong V, Sar B, Horm SV, Chin S, et al. Identification of Molecular
Markers Associated with Alteration of Receptor-Binding Specificity in a Novel Genotype of
Highly Pathogenic Avian Influenza A(H5N1) Viruses Detected in Cambodia in 2013. J Virol.
2014;88: 13897–13909.

23.   Horwood PF, Karlsson EA, Horm SV, Ly S, Heng S, Chin S, et al. Circulation and
characterization of seasonal influenza viruses in Cambodia, 2012-2015. Influenza Other
Respi Viruses. 2019;13: 465–476.

24.   Zhu H, Wang J, Wang P, Song W, Zheng Z, Chen R, et al. Substitution of lysine at 627
position in PB2 protein does not change virulence of the 2009 pandemic H1N1 virus in
mice. Virology. 2010;401: 1–5.

25.   Horwood PF, Horm SV, Suttie A, Thet S, Y P, Rith S, et al. Co-circulation of Influenza A H5,

H7, and H9 Viruses and Co-infected Poultry in Live Bird Markets, Cambodia. Emerg Infect
Dis. 2018;24: 352–355.

26. Zhou B, Donnelly ME, Scholes DT, St George K, Hatta M, Kawaoka Y, et al. Single-
reaction genomic amplification accelerates sequencing and vaccine production for classical
and Swine origin human influenza a viruses. J Virol. 2009;83: 10309–10313.

27. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9: 357–359.

28. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30: 2114–2120.

29. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.

30. Koboldt DC, Chen K, Wylie T, Larson DE, McLellan MD, Mardis ER, et al. VarScan: variant
detection in massively parallel sequencing of individual and pooled samples.
Bioinformatics. 2009;25: 2283–2285.

31. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2: somatic
mutation and copy number alteration discovery in cancer by exome sequencing. Genome
Res. 2012;22: 568–576.

32. Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative contribution
to global health. Global Challenges. 2017;1: 33–46.

33. Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data – from vision to
reality. Eurosurveillance. 2017;22: 30494.

34. Zhang Y, Aevermann BD, Anderson TK, Burke DF, Dauphin G, Gu Z, et al. Influenza
Research Database: An integrated bioinformatics resource for influenza virus research.
Nucleic Acids Res. 2017;45: D466–D474.

35. Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, et al. Nextstrain: real-
time tracking of pathogen evolution. Kelso J, editor. Bioinformatics. 2018;34: 4121–4123.

36. Katoh K, Misawa K, Kuma KK-I, Miyata T. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002;30: 3059–
3066.

37. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol.
2015;32: 268–274.

38. Chernomor O, von Haeseler A, Minh BQ. Terrace Aware Data Structure for Phylogenomic
Inference from Supermatrices. Syst Biol. 2016;65: 997–1008.

39. Sagulenko P, Puller V, Neher RA. TreeTime: Maximum-likelihood phylodynamic analysis.
Virus Evolution. 2018;4. doi:10.1093/ve/vex042

40. Nelson CW, Moncla LH, Hughes AL. SNPGenie: estimating evolutionary parameters to
detect natural selection using pooled next-generation sequencing data. Bioinformatics.

2015;31: 3709–3711.

41. Nei M, Gojobori T. Simple methods for estimating the numbers of synonymous and
739   nonsynonymous nucleotide substitutions. Mol Biol Evol. 1986;3: 418–426.

740   42. Sorn S, Sok T, Ly S, Rith S, Tung N, Viari A, et al. Dynamic of H5N1 virus in Cambodia and
741   emergence of a novel endemic sub-clade. Infect Genet Evol. 2013;15: 87–94.

742   43. Yamada S, Suzuki Y, Suzuki T, Le MQ, Nidom CA, Sakai-Tagawa Y, et al. Haemagglutinin
743   mutations responsible for the binding of H5N1 influenza A viruses to human-type receptors.
744   Nature. 2006;444: 378–382.

745   44. Yang Z-Y, Wei C-J, Kong W-P, Wu L, Xu L, Smith DF, et al. Immunization by avian H5
746   influenza hemagglutinin mutants with altered receptor binding specificity. Science.
747   2007;317: 825–828.

748   45. Wang M, Tscherne DM, McCullough C, Caffrey M, García-Sastre A, Rong L. Residue Y161
749   of influenza virus hemagglutinin is involved in viral recognition of sialylated complexes from
750   different hosts. J Virol. 2012;86: 4455–4462.

751   46. Suguitan AL, Matsuoka Y, Lau Y-F, Santos CP, Vogel L, Cheng LI, et al. The multibasic
752   cleavage site of the hemagglutinin of highly pathogenic A/Vietnam/1203/2004 (H5N1) avian
753   influenza virus acts as a virulence factor in a host-specific manner in mammals. J Virol.
754   2012;86: 2706–2714.

755   47. Schrauwen EJA, Herfst S, Leijten LM, van Run P, Bestebroer TM, Linster M, et al. The
756   multibasic cleavage site in H5N1 virus is critical for systemic spread along the olfactory and
757   hematogenous routes in ferrets. J Virol. 2012;86: 3975–3984.

758   48. Zhou H, Yu Z, Hu Y, Tu J, Zou W, Peng Y, et al. The Special Neuraminidase Stalk-Motif
759   Responsible for Increased Virulence and Pathogenesis of H5N1 Influenza A Virus. Martin
760   DP, editor. PLoS One. 2009;4: e6277.

761   49. Zhou H, Jin M, Chen H, Huag Q, Yu Z. Genome-sequenee Analysis of the Pathogenic
762   H5N1 Avian Influenza A Virus Isolated in China in 2004. Virus Genes. 2006;32: 85–95.

763   50. Matsuoka Y, Swayne DE, Thomas C, Rameix-Welti M-A, Naffakh N, Warnes C, et al.
764   Neuraminidase Stalk Length and Additional Glycosylation of the Hemagglutinin Influence
765   the Virulence of Influenza H5N1 Viruses for Mice. J Virol. 2009;83: 4704–4708.

766   51. Hatta M, Gao P, Halfmann P, Kawaoka Y. Molecular basis for high virulence of Hong Kong
767   H5N1 influenza A viruses. Science. 2001;293: 1840–1842.

768   52. Zhao L, Illingworth CJR. Measurements of intrahost viral diversity require an unbiased
769   diversity metric. Virus Evol. 2019;5: vey041.

770   53. Subbarao EK, Kawaoka Y, Murphy BR. Rescue of an influenza A virus wild-type PB2 gene
771   and a mutant derivative bearing a site-specific temperature-sensitive and attenuating
772   mutation. J Virol. 1993;67: 7223–7228.

773   54. Le QM, Sakai-Tagawa Y, Ozawa M, Ito M, Kawaoka Y. Selection of H5N1 influenza virus
774   PB2 during replication in humans. J Virol. 2009;83: 5278–5281.

55. Gabriel G, Dauber B, Wolff T, Planz O, Klenk H-D, Stech J. The viral polymerase mediates
adaptation of an avian influenza virus to a mammalian host. Proc Natl Acad Sci U S A.
2005;102: 18590–18595.

56. Steel J, Lowen AC, Mubareka S, Palese P. Transmission of Influenza Virus in a
Mammalian Host Is Increased by PB2 Amino Acids 627K or 627E/701N. Baric RS, editor.
PLoS Pathog. 2009;5: e1000252.

57. Soh YQS, Moncla LH, Eguia R, Bedford T, Bloom JD. Comprehensive mapping of
adaptation of the avian influenza polymerase protein PB2 to humans. Elife. 2019;8.
doi:10.7554/eLife.45079

58. Auewarakul P, Suptawiwat O, Kongchanagul A, Sangma C, Suzuki Y, Ungchusak K, et al.
An avian influenza H5N1 virus that binds to a human-type receptor. J Virol. 2007;81: 9950–
9955.

59. Naughtin M, Dyason JC, Mardy S, Sorn S, von Itzstein M, Buchy P. Neuraminidase inhibitor
sensitivity and receptor-binding specificity of Cambodian clade 1 highly pathogenic H5N1
influenza virus. Antimicrob Agents Chemother. 2011;55: 2004–2010.

60. Lyons DM, Lauring AS. Mutation and epistasis in influenza virus evolution [Internet].
Viruses 2018. doi:10.3390/v10080407

61. Visher E, Whitefield SE, McCrone JT, Fitzsimmons W, Lauring AS. The Mutational
Robustness of Influenza A Virus. PLoS Pathog. 2016;12: e1005856.

62. Sanjuán R. Mutational fitness effects in RNA and single-stranded DNA viruses: common
patterns revealed by site-directed mutagenesis studies. Philos Trans R Soc Lond B Biol
Sci. 2010;365: 1975–1982.

63. Watanabe Y, Ibrahim MS, Ellakany HF, Kawashita N, Mizuike R, Hiramatsu H, et al.
Acquisition of Human-Type Receptor Binding Specificity by New H5N1 Influenza Virus
Sublineages during Their Emergence in Birds in Egypt. Fouchier RAM, editor. PLoS
Pathog. 2011;7: e1002068.

64. Guilligay D, Tarendeau F, Resa-Infante P, Coloma R, Crepin T, Sehr P, et al. The structural
basis for cap binding by influenza virus polymerase subunit PB2. Nat Struct Mol Biol.
2008;15: 500–506.

65. Nerome R, Hiromoto Y, Fukushima T, Nerome K, Lim W, Yamazaki Y, et al. Evolutionary
characterization of the six internal genes of H5N1 human influenza A virus. J Gen Virol.
2000;81: 1293–1303.

66. Xu L, Bao L, Zhou J, Wang D, Deng W, Lv Q, et al. Genomic Polymorphism of the
Pandemic A (H1N1) Influenza Viruses Correlates with Viral Replication, Virulence, and
Pathogenicity In Vitro and In Vivo. Digard P, editor. PLoS One. 2011;6: e20698.

67. Bussey KA, Desmet EA, Mattiacio JL, Hamilton A, Bradel-Tretheway B, Bussey HE, et al.
PA Residues in the 2009 H1N1 Pandemic Influenza Virus Enhance Avian Influenza Virus
Polymerase Activity in Mammalian Cells. J Virol. 2011;85: 7020–7028.

68. Hiromoto Y, Saito T, Lindstrom S, Nerome K. Characterization of Low Virulent Strains of

Highly Pathogenic A/Hong Kong/156/97 (H5N1) Virus in Mice after Passage in
Embryonated Hens' Eggs. Virology. 2000;272: 429–437.

69. Webster RG, Govorkova EA, Kaverin NV, Varich NL, Gitelman AK, Lipatov AS, et al.
Structure of antigenic sites on the haemagglutinin molecule of H5 avian influenza virus and
phenotypic variation of escape mutants. J Gen Virol. 2002;83: 2497–2505.

70. Yen H-L, Aldridge JR, Boon ACM, Ilyushina NA, Salomon R, Hulse-Post DJ, et al. Changes
in H5N1 influenza virus hemagglutinin receptor binding domain affect systemic spread.
Proc Natl Acad Sci U S A. 2009;106: 286–291.

71. Stevens J, Blixt O, Chen L-M, Donis RO, Paulson JC, Wilson IA. Recent Avian H5N1
Viruses Exhibit Increased Propensity for Acquiring Human Receptor Specificity. J Mol Biol.
2008;381: 1382–1394.

72. Wang W, Lu B, Zhou H, Suguitan AL, Cheng X, Subbarao K, et al. Glycosylation at 158N of
the hemagglutinin protein and receptor binding specificity synergistically affect the
antigenicity and immunogenicity of a live attenuated H5N1 A/Vietnam/1203/2004 vaccine
virus in ferrets. J Virol. 2010;84: 6570–6577.

73. Chutinimitkul S, van Riel D, Munster VJ, van den Brand JMA, Rimmelzwaan GF, Kuiken T,
et al. In vitro assessment of attachment pattern and replication efficiency of H5N1 influenza
A viruses with altered receptor specificity. J Virol. 2010;84: 6825–6833.

74. Stevens J, Blixt O, Tumpey TM, Taubenberger JK, Paulson JC, Wilson IA. Structure and
receptor specificity of the hemagglutinin from an H5N1 influenza virus. Science. 2006;312:
404–410.

75. Maines TR, Chen L-M, Van Hoeven N, Tumpey TM, Blixt O, Belser JA, et al. Effect of
receptor binding domain mutations on receptor binding and transmissibility of avian
influenza H5N1 viruses. Virology. 2011;413: 139–147.

76. Chen L-M, Blixt O, Stevens J, Lipatov AS, Davis CT, Collins BE, et al. In vitro evolution of
H5N1 avian influenza virus toward human-type receptor specificity. Virology. 2012;422:
105–113.

77. Weber F, Kochs G, Gruber S, Haller O. A Classical Bipartite Nuclear Localization Signal on
Thogoto and Influenza A Virus Nucleoproteins. Virology. 1998;250: 9–18.

78. Grantham ML, Wu W-H, Lalime EN, Lorenzo ME, Klein SL, Pekosz A. Palmitoylation of the
influenza A virus M2 protein is not required for virus replication in vitro but contributes to
virus virulence. J Virol. 2009;83: 8655–8661.

79. Holsinger LJ, Shaughnessy MA, Micko A, Pinto LH, Lamb RA. Analysis of the
posttranslational modifications of the influenza virus M2 protein. J Virol. 1995;69: 1219–
1225.

80. Li Y, Yamakita Y, Krug RM. Regulation of a nuclear export signal by an adjacent inhibitory
sequence: The effector domain of the influenza virus NS1 protein. Proceedings of the
National Academy of Sciences. 1998;95: 4864–4869.

81. Hale BG, Barclay WS, Randall RE, Russell RJ. Structure of an avian influenza A virus NS1

protein effector domain. Virology. 2008;378: 1–5.

82. Imai H, Shinya K, Takano R, Kiso M, Muramoto Y, Sakabe S, et al. The HA and NS Genes
of Human H5N1 Influenza A Virus Contribute to High Virulence in Ferrets. Basler CF,
editor. PLoS Pathog. 2010;6: e1001106.

Acknowledgments

We would like to thank Katherine Xue for her careful reading and comments on the manuscript.

Figure legends

Figure 1: Phylogenetic placement of H5N1 samples from Cambodia

All currently available H5N1 sequences were downloaded from the Influenza Research

Database and the Global Initiative on Sharing All Influenza Data and used to generate full

genome phylogenies using Nextstrain's augur pipeline as shown in the trees on the left.

Phylogenies for the full genome are shown in Figure S2. Colors represent the geographic region

in which the sample was collected (for tips) or the inferred geographic location (for internal

nodes). The x-axis position indicates the date of sample collection (for tips) or the inferred time

to the most recent common ancestor (for internal nodes). In the full phylogeny (left), H5N1

viruses from Cambodia selected for within-host analysis are indicated by tan circles with black

outlines. The subtrees containing the Cambodian samples selected for within-host analysis are

shown to the right and are indicated with grey, dashed arrows. In these trees, human tips are

marked with a tan circle with a black outline, while duck tips are denoted with a tan square with

a black outline. All samples from our within-host dataset are labelled in the subtrees with their

strain name. Internal genes from samples collected prior to 2013 belong to clade 1.1.2, while

internal genes from samples collected in 2013 or later belong to clade 2.3.2.1a. All HA and NA

sequences in this dataset, besides A/duck/Cambodia/Y0224304/2014, belong to clade 1.1.2.

Figure 2: Within-host diversity in humans and ducks is dominated by low-frequency

variation

(a) Within-host polymorphisms present in at least 1% of sequencing reads were called in all

human (red) and duck (blue) samples. Each dot represents one unique single nucleotide variant

(SNV), the x-axis represents the nucleotide site of the SNV, and the y-axis represents its

frequency within-host. (b) For each sample in our dataset, we calculated the proportion of its

synonymous (light blue and light red) and nonsynonymous (dark blue and dark red) within-host

variants present at frequencies of 1-10%, 10-20%, 20-30%, 30-40%, and 40-50%. We then took

the mean across all human (red) or duck (blue) samples. Bars represent the mean proportion of

variants present in a particular frequency bin and error bars represent standard error. Grey dots

and connecting lines represent the expected proportion of variants in each bin under a neutral

model. (c) We calculated Tajima's D across the full genomes of humans and ducks, separately

for synonymous and nonsynonymous sites. Values represent the mean Tajima's D across all

humans or ducks, and values in parentheses represent the 95% confidence interval.

Figure 3: Purifying selection and genetic drift shape within-host diversity

For each sample and gene, we computed the average number of pairwise nonsynonymous

differences per nonsynonymous site (π_N) and the average number of pairwise synonymous

differences per synonymous site (π_S). We then calculated the mean for each gene and species.

Each bar represents the mean and error bars represent the standard error calculated by

performing 10,000 bootstrap resamplings. Human values are shown in red and duck values are

shown in blue.

Figure 4: Mutations are present at functionally relevant sites

We queried each amino acid changing mutation identified in our dataset against all known

annotations present in the Influenza Research Database Sequence Feature Variant Types tool.

Each mutation is colored according to its function. Shape represents whether the mutation was

identified in a human (circle) or duck (square) sample. Mutations shown here were detected in

at least 1 human or duck sample. Filled in shapes represent nonsynonymous changes and open

shapes represent synonymous mutations. Grey, transparent dots represent mutations for which

no host-related function was known. Each nonsynonymous colored mutation, its frequency, and

its phenotypic effect is shown in **Table 3**, and a full list of all mutations and their annotations are

available in **Table S1**.

Figure 5: Ducks share more polymorphisms than expected by chance

(a) All amino acid sites that were polymorphic in at least 2 samples are shown. This includes

sites at which each sample had a polymorphism at the same site, but encoded different variant

amino acids. There are 3 amino acid sites that are shared by at least 2 duck samples, and 9

polymorphic sites shared by at least 2 human samples. 3 synonymous changes are detected in

both human and duck samples (PB1 371, PA 397, and NP 201). Frequency is shown on the y-

axis. (b) To test whether the level of sharing we observed was more or less than expected by

chance, we performed a permutation test. The x-axis represents the number of sites shared by

at least 2 ducks (blue) or at least 2 humans (red), and the bar height represents the number of

simulations in which that number of shared sites occurred. Actual observed number of shared

sites (3 and 9) are shown with a dashed line. (c) The same permutation test as shown in (b),

except that only 70% of amino acid sites were permitted to mutate. (d) The same permutation

test as shown in (b), except that only 60% of amino acid sites were permitted to mutate.

Figure 6: A small subset of within-host variants are enriched on spillover branches

(a) A schematic for how we classified host transitions along the phylogeny. Branches within

monophyletic human clades were labelled "to-human" (red branches). Branches leading to a

monophyletic human clade, whose parent node had avian children were also labelled as "to-

human" (half red, half blue branches), and all other branches were labelled "to-avian" (blue

branches). (b) Each amino acid-changing SNV we detected within-host in either ducks (left) or

humans (right) that was present in the H5N1 phylogeny is displayed. Each bar represents an

amino acid mutation, and its height represents the number of to-avian (blue) or to-human (red)

transitions in which this mutation was present along the H5N1 phylogeny. Significance was

assessed with a Fisher's exact test. * indicates $p < 0.05$, **** indicates $p < 0.0001$.

Supporting information legends

Figure S1: Genome coverage

The mean coverage depth at each nucleotide site (x-axis) for each gene across our 8 human

and 5 duck samples is shown. Solid black lines represent the mean coverage across samples,

and the grey shaded area represents the standard deviation of coverage depth across samples.

Figure S2: Full genome phylogenetic placement of H5N1 virus samples from Cambodia

All currently available H5N1 virus sequences were downloaded from the Influenza Research

Database and the Global Initiative on Sharing All Influenza Data and used to generate full

genome phylogenies using Nextstrain's augur pipeline. Colors represent the geographic region

in which the sample was collected (for tips) or the inferred geographic location (for internal

nodes). The x-axis position indicates the date of sample collection (for tips) or the inferred time

to the most recent common ancestor (for internal nodes). In the full phylogenies (left), H5N1

viruses from Cambodia selected for within-host analysis are indicated by tan circles with black

outlines. The subtrees containing the Cambodian samples selected for within-host analysis are

shown to the right in the order that they appear in the full tree. In these trees, human tips are

marked with a tan circle with a black outline, while duck tips are denoted with a tan square with

a black outline. Both human and duck tips are labelled with their strain names. Internal genes

from samples collected prior to 2013 belong to clade 1.1.2, while internal genes from samples

collected in 2013 or later belong to clade 2.3.2.1a. All HA and NA sequences in this dataset,

besides A/duck/Cambodia/Y0224304/2014, belong to clade 1.1.2.

Figure S3: All within-host variants detected in our dataset

All within-host variants detected in our study are shown. Each row represents one sample and

each column represents one gene. The x-axis shows the nucleotide site and the y-axis shows

the frequency that the variant was detected within-host. Filled circles represent nonsynonymous

changes, while open circles represent synonymous changes. Blue dots represent variants

identified within duck samples, while red dots represent variants identified in human samples.

Blank plots indicate that no variants were identified in that sample and gene.

Figure S4: Neither diversity nor host-specific mutations increase over time
969
(a) For each human sample, the full genome nucleotide diversity (π_N or π_S) is plotted vs. the

days post-symptom onset. Dark red dots represent the mean, full-genome nonsynonymous

diversity for a given sample (π_N), and light red dots represent the mean, full-genome

synonymous diversity for that same sample (π_S). Neither nonsynonymous nor synonymous

diversity are correlated with days post symptom onset (nonsynonymous: r^2 = -0.17, p = 0.69;

synonymous: r^2 = -0.22, p = -0.61). (b) To compare whether the number of putative host-

adapting mutations increased over time in humans, we compared the number of host-specific

and non-host specific mutations in humans sampled either in "early infection" (5-8 days post

symptom onset), or in "late infection" (9-12 days post symptom onset). We divided the data into

these categories by splitting on the mean days post symptom onset for human samples, which

was 8 days. We then compared the proportion of host-specific variants during early and late

infections with a Fisher's exact test. The proportion of variants that are host-specific is not

different in early vs. late infections (p = 0.72).

Table S1: All within-host SNVs with annotations

Every SNV identified in humans and ducks within-host are displayed with their frequency,

coding region change, and functional annotation. All annotations for H5 HAs, N1 NAs, and all

subtypes for all other genes were downloaded from the Influenza Research Database

Sequence Feature Variant Types tool. Each SNV was then annotated as shown in the

"description" column. These descriptions are paraphrased from annotations presented in the

Influenza Research Database. We then manually curated annotated mutations to determine

whether they were involved in "host-specific" functions or not, as shown in the "host-specific?"

column. We defined host-specific functions/interactions as receptor binding, interaction with host

cellular machinery, nuclear import and export, immune antagonism, 5' cap binding, temperature

sensitivity, and glycosylation. We also included sites that have been phenotypically identified as

determinants of transmissibility and virulence. Sites that participate in binding interactions with

other viral subunits or vRNP, conserved active site domains, drug resistance mutations, and

epitope sites were not categorized as host-specific for this analysis. We annotated both

synonymous and nonsynonymous mutations in our dataset.

1000

1001

1002

1003

1004

Tables

Table 1: Sample information

Sample ID	Host	Sample type	Collection	Date	Days post-symptom onset	vRNA copies/µl (after vRNA extraction)	Clade
A/duck/Cambodia/PV027D1/2010	Domestic duck	Pooled organs	Poultry outbreak investigation	April 2010	NA	5.45×10^6	1.1.2
A/duck/Cambodia/083D1/2011	Domestic duck	Pooled organs	Poultry outbreak investigation	September 2011	NA	3.74×10^7	1.1.2
A/duck/Cambodia/381W11M4/2013	Domestic duck	Pooled throat and cloacal swab	Live bird market surveillance	March 2013	NA	7.37×10^5	1.1.2/2.3.2.1a reassortant
A/duck/Cambodia/Y0224301/2014	Domestic duck	Pooled organs	Poultry outbreak investigation	February 2014	NA	2.0×10^5	1.1.2/2.3.2.1a reassortant
A/duck/Cambodia/Y0224304/2014	Domestic duck	Pooled organs	Poultry outbreak investigation	February 2014	NA	5.0×10^6	1.1.2/2.3.2.1a reassortant
A/Cambodia/V0401301/2011	Human (10F, died)	Throat swab	Event-based surveillance	April 2011	9	5.02×10^3	1.1.2

A/Cambodia/V0417301/2011	Human (5F, died)	Throat swab	Event-based surveillance	April 2011	5	8.98×10^4	1.1.2
A/Cambodia/W0112303/2012	Human (2M, died)	Throat swab	Event-based surveillance	January 2012	7	2.05×10^3	1.1.2
A/Cambodia/X0125302/2013	Human (1F, died)	Throat swab	Event-based surveillance	January 2013	12	6.84×10^4	1.1.2/2.3.2.1a reassortant
A/Cambodia/X0128304/2013	Human (9F, died)	Throat swab	Event-based surveillance	January 2013	8	5.09×10^3	1.1.2/2.3.2.1a reassortant
A/Cambodia/X0207301/2013	Human (5F, died)	Throat swab	Event-based surveillance	February 2013	12	1.73×10^5	1.1.2/2.3.2.1a reassortant
A/Cambodia/X0219301/2013	Human (2M, died)	Throat swab	Event-based surveillance	February 2013	12	1.66×10^3	1.1.2/2.3.2.1a reassortant
A/Cambodia/X1030304/2013	Human (2F, died)	Throat swab	Event-based surveillance	October 2013	8	1.08×10^4	1.1.2/2.3.2.1a reassortant

Table 2: Mean π_N and π_S values per gene

Gene	Species	Mean π_N	Mean π_S	π_N/π_S	p-value
PB2	Human	0.00015	0.00023	0.65	0.50
PB2	Duck	0.00	0.00031	0.00	0.27
PB1	**Human**	**0.000083**	**0.00038**	**0.22**	**0.049**
PB1	Duck	0.000009	0.000066	0.14	0.31
PA	Human	0.00012	0.00044	0.27	0.083
PA	Duck	0.000037	0.00016	0.23	0.094
HA	Human	0.00044	0.00035	1.26	0.61
HA	Duck	0.000054	0.00025	0.22	0.40
NP	Human	0.000050	0.00050	0.10	0.12
NP	Duck	0.00011	0.00028	0.39	0.49
NA	Human	0.000078	0.0005	0.16	0.064
NA	Duck	0.000056	0.00023	0.24	0.27
M1	Human	0.00010	0.00063	0.14	0.23
M1	Duck	0.000068	0.00	NA	0.18
M2	**Human**	**0.00017**	**0.00**	**NA**	**0.042**
M2	Duck	0.00	0.00	NA	NA
NS1	Human	0.000014	0.00056	0.03	0.20
NS1	Duck	0.000036	0.00	NA	0.37

NEP	Human	0.000064	0.00	NA	0.18
NEP	Duck	0.000030	0.00013	0.23	0.37
Full genome	**Human**	**0.000139**	**0.000381**	**0.36**	**0.0059**
Full genome	**Duck**	**0.000039**	**0.00018**	**0.22**	**0.038**

For each gene and sample, we computed nonsynonymous (π_N), and and synonymous (π_S) diversity as the average number of pairwise differences between a set of DNA sequences. Values of 0.00 indicate that there were no SNPs identified in that gene for that host species and mutation type. We then combined values from each sample to generate a diversity estimate for each gene and host species. Significance was assessed by a paired t-test testing the null hypothesis that $\pi_N = \pi_S$. Bold values of p < 0.05.

Table 3: Mutations identified at functionally relevant sites

Sample	Gene	Nt site	Ref base	Variant base	Coding region change	Freq.	Description	Type
A/Cambodia/X0128304/2013	PB2	1069	A	T	N348Y	6.15%	Putative m7GTP cap binding site[64].	replication
A/Cambodia/V0401301/2011	PB2	1202	A	C	N392H	3.61%	Putative m7GTP cap binding site[64].	replication
A/Cambodia/W0112303/2012	PB2	1891	G	A	E627K	6.63%	A Lys at 627 enhances mammalian replication[51,53].	replication
A/Cambodia/X0125302/2013	PB2	2022	G	A	V667I	2.99%	An Ile at 667 was associated with human-infecting H5N1 virus strains[65].	replication
A/Cambodia/W0112303/2012	PB2	2113	A	G	N701D	16.49%	An Asn at 701 enhances mammalian replication[55,56].	replication
A/Cambodia/X0125302/2013	PB2	2163	A	G	S714G	9.59%	An Arg at 714 enhances mammalian replication[55].	replication
A/Cambodia/X1030304/2013	PB1	631	A	G	R211G	2.34%	Nuclear localization motif.	interaction with host machinery

A/Cambodia/X0125302/2013	PB1	1078	A	G	K353R	2.94%	An Arg at 353 is associated with higher replication and pathogenicity of an H1N1 pandemic strain[66].	replication
A/Cambodia/X0125302/2013	PB1	1716	A	T	T566S	5.20%	An Ala at 566 is associated with higher replication and pathogenicity of an H1N1 pandemic virus[66].	replication
A/Cambodia/X0219301/2013	PA	265	A	G	T85A	2.84%	An Ile at 85 enhances polymerase activity of pandemic H1N1 in mammalian cells[67].	replication
A/Cambodia/X0128304/2013	PA	1868	A	G	K615R	2.47%	An Asn at PA 615 has been associated with adaptation of avian influenza polymerases to humans[55].	replication
A/Cambodia/X0207301/2013	PA	1903	A	G	S631G	1.79%	A Ser at 631 enhances virulence of H5N1 viruses in mice[68].	virulence
A/Cambodia/X0128304/2013	HA	299	A	G	E91G	6.33%	A Lys at 91 enhances α-2,6 binding[43]. (H5 mature: 75)	receptor binding
A/Cambodia/V0417301/2011	HA	425	A	G	E142G	3.20%	Putative glycosylation site[69]. (H5 mature: 126)	virulence
A/Cambodia/V0401301/2011	HA	449	C	T	A150V	20.24%	A Val at 150 confers enhanced α-2,6 sialic acid binding in H5N1 viruses[58,59]. (H5 mature: 134)	receptor binding

A/Cambodia/X0125302/2013	HA	449	C	T	A150V	15.09%	A Val at 150 confers enhanced α-2,6 sialic acid binding in H5N1 viruses[58,59]. (H5 mature: 134)	receptor binding
A/Cambodia/X0128304/2013	HA	542	A	C	K172T	11.50%	Part of putative glycosylation motif that improves α-2,6 binding[70–72]. (H5 mature: 156)	receptor binding
A/Cambodia/V0401301/2011	HA	517	T	C	Y173H	5.04%	Residue involved in sialic acid recognition[45]. (H5 mature: 157)	receptor binding
A/Cambodia/V0401301/2011	HA	593	A	G	N198S	3.32%	A Lys at 198 confers α-2,6 sialic acid binding [43,73](H5 mature: 182)	receptor binding
A/Cambodia/X0128304/2013	HA	703	A	G	T226A	28.91%	An Ile at 226 enhanced α-2,6 sialic acid binding[63]. (H5 mature: 210)	receptor binding
A/Cambodia/V0401301/2011	HA	713	A	T	Q238L	2.80%	A Leu at 238 confers a switch from α-2,3 to α-2,6 sialic acid binding and is a determinant of mammalian transmission[11,12,73–76]. (H5 mature: 222)	receptor binding
A/Cambodia/V0417301/2011	HA	713	A	T	Q238L	8.45%	A Leu at 238 confers a switch from α-2,3 to α-2,6 sialic acid binding and is a determinant of mammalian transmission[11,12,73–76]. (H5 mature: 222)	receptor binding
A/Cambodia/X0125302/2013	HA	713	A	G	Q238R	40.30%	A Leu at 238 confers a switch from α-2,3 to α-2,6 sialic acid binding and is a determinant of mammalian transmission[11,12,73–76]. (H5 mature: 222)	receptor binding

A/duck/Cambodia/Y0224304/2014	NP	674	C	T	T215I	3.69%	Nuclear targeting motif[77].	interaction with host machinery
A/Cambodia/X1030304/2013	M2	861	G	A	C50Y	2.03%	A Cys at position 50 is a palmitoylation site that enhances virulence[78,79].	virulence
A/Cambodia/X0128304/2013	NS1	502	C	T	P159L	2.8%	Part of the NS1 nuclear export signal mask[80].	interaction with host machinery
A/duck/Cambodia/Y0224301/2014	NS1	646	T	C	L207P	2.22%	NS1 flexible tail, which interacts with host machinery[81].	interaction with host machinery
A/duck/Cambodia/Y0224301/2014	NS1	654	C	T	P210S	2.55%	NS1 flexible tail, which interacts with host machinery[81].	interaction with host machinery
A/Cambodia/X0207301/2013	NEP	609	A	G	E47G	4.59%	This site was implicated in enhanced virulence of H5N1 viruses in ferrets[82].	virulence

All nonsynonymous mutations that were identified in sites with putative links to host-specific phenotypes are shown. We identify a handful of amino acid mutations that have been explicitly linked to mammalian adaptation of avian influenza viruses. For HA mutations, all mutations use native H5 numbering, including the signal peptide. For ease of comparison, the corresponding amino acid number in mature, H5 peptide numbering is also provided in parentheses in the description column. Full annotations for all mutations in our data are shown in **Table S1**.

Figure 1: Phylogenetic placement of H5N1 samples from Cambodia
All currently available H5N1 sequences were downloaded from the Influenza Research Database and the Global Initiative on Sharing All Influenza Data and used to generate full genome phylogenies using Nextstrain's augur pipeline as shown in the trees on the left. Phylogenies for the full genome are shown in Figure S2. Colors represent the geographic region in which the sample was collected (for tips) or the inferred geographic location (for internal nodes). The x-axis position indicates the date of sample collection (for tips) or the inferred time to the most recent common ancestor (for internal nodes). In the full phylogeny (left), H5N1 viruses from Cambodia selected for within-host analysis are indicated by tan circles with black outlines. The subtrees containing the Cambodian samples selected for within-host analysis are shown to the right and are indicated with grey, dashed arrows. In these trees, human tips are marked with a circles, while duck tips are denoted with squares, and are labelled with their strain name. All HA and NA sequences in this dataset, besides A/duck/Cambodia/Y0224304/2014, belong to clade 1.1.2. Internal genes from samples collected prior to 2013 belong to clade 1.1.2, while internal genes from samples collected in 2013 or later belong to clade 2.3.2.1a.

a.

b.

c.

Figure 2: Within-host diversity in humans and ducks is dominated by low-frequency variation
(a) Within-host polymorphisms present in at least 1% of sequencing reads were called in all human (red) and duck (blue) samples. Each dot represents one unique single nucleotide variant (SNV), the x-axis represents the nucleotide site of the SNV, and the y-axis represents its frequency within-host. **(b)** For each sample in our dataset, we calculated the proportion of its synonymous (light blue and light red) and nonsynonymous (dark blue and dark red) within-host variants present at frequencies of 1-10%, 10-20%, 20-30%, 30-40%, and 40-50%. We then took the mean across all human (red) or duck (blue) samples. Bars represent the mean proportion of variants present in a particular frequency bin and error bars represent standard error. Grey dots and connecting lines represent the expected proportion of variants in each bin under a neutral model. **(c)** We calculated Tajima's *D* across the full genomes of humans and ducks, separately for synonymous and nonsynonymous sites. Values represent the mean Tajima's *D* across all humans or ducks, and values in parentheses represent the 95% confidence interval.

Figure 3: Purifying selection and genetic drift shape within-host diversity

For each sample and gene, we computed the average number of pairwise nonsynonymous differences per nonsynonymous site (π_N) and the average number of pairwise synonymous differences per synonymous site π_S). We then calculated the mean for each gene and species. Each bar represents the mean and error bars represent the standard error calculated by performing 10,000 bootstrap resamplings. Human values are shown in red and duck values are shown in blue.

Figure 4: Mutations are present at functionally relevant sites

We queried each amino acid changing mutation identified in our dataset against all known annotations present in the Influenza Research Database Sequence Feature Variant Types tool. Each mutation is colored according to its function. Shape represents whether the mutation was identified in a human (circle) or duck (square) sample. Mutations shown here were detected in at least 1 human or duck sample. Filled in shapes represent nonsynonymous changes and open shapes represent synonymous mutations. Grey, transparent dots represent mutations for which no host-related function was known. Each nonsynonymous colored mutation, its frequency, and its phenotypic effect is shown in Table 3, and a full list of all mutations and their annotations are available in Table S1.

Figure 5: Ducks share more polymorphisms than expected by chance
(a) All amino acid sites that were polymorphic in at least 2 samples are shown. This includes sites at which each sample had a polymorphism at the same site, but encoded different variant amino acids. There are 3 amino acid sites that are shared by at least 2 duck samples, and 9 polymorphic sites shared by at least 2 human samples. 3 synonymous changes are detected in both human and duck samples (PB1 371, PA 397, and NP 201). Frequency is shown on the y-axis. **(b)** To test whether the level of sharing we observed was more or less than expected by chance, we performed a permutation test. The x-axis represents the number of sites shared by at least 2 ducks (blue) or at least 2 humans (red), and the bar height represents the number of simulations in which that number of shared sites occurred. Actual observed number of shared sites (3 and 9) are shown with a dashed line. **(c)** The same permutation test as shown in **(b)**, except that only 70% of amino acid sites were permitted to mutate. **(d)** The same permutation test as shown in **(b)**, except that only 60% of amino acid sites were permitted to mutate.

a.

b.

Figure 6: A small subset of within-host variants are enriched on spillover branches
(a) A schematic for how we classified host transitions along the phylogeny. Branches within monophyletic human clades were labelled "to-human" (red branches). Branches leading to a monophyletic human clade, whose parent node had avian children were also labelled as "to-human" (half red, half blue branches), and all other branches were labelled "to-avian" (blue branches). **(b)** Each amino acid-changing SNV we detected within-host in either ducks (left) or humans (right) that was present in the H5N1 phylogeny is displayed. Each bar represents an amino acid mutation, and its height represents the number of to-avian (blue) or to-human (red) transitions in which this mutation was present along the H5N1 phylogeny. Significance was assessed with a Fisher's exact test. * indicates p < 0.05, **** indicates p < 0.0001.