Neuromodulatory selection of motor neuron recruitment patterns in

a visuomotor behavior increases speed

Authors

Urvashi Jha[1,2], Vatsala Thirumalai[1, *]

Highlights

• Zebrafish larvae generate swims of increased speed during optomotor

response when D1-like receptors are activated.

• D1-like receptor activation increases the extent of tail bending during forward

swims and turns resulting in increased swim speed.

• Neuromodulation via D1-like receptors increases motor drive by enhancing

excitability of 'slow' motor neurons. In addition, D1-like receptor activation

recruits quiescent 'fast' motor neurons to increase swim speed.

• This demonstrates neuromodulatory selection of motor neurons belonging to

different 'speed' modules to alter swimming behavior.

[1] National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore, Karnataka-560065; India

[2] SASTRA Deemed University, School of Chemical and Biotechnology, Thanjavur, Tamil Nadu-613401; India

* Correspondence: vatsala@ncbs.res.in

Summary

Animals generate locomotion at different speeds to suit their behavioral needs. Spinal circuits generate locomotion at these varying speeds by sequential activation of different spinal interneurons and motor neurons. Larval zebrafish can generate slow swims for prey capture and exploration by activation of secondary motor neurons and much faster and vigorous swims during escapes and struggles via the additional activation of primary motor neurons. Neuromodulators are known to alter motor output of spinal circuits yet their precise role in speed regulation is not understood well. Here, in the context of optomotor response (OMR), an innate, evoked locomotor behavior, we show that dopamine (DA) provides an additional layer to regulation of swim speed in larval zebrafish. Activation of D1-like receptors increases swim speed during OMR in free-swimming larvae. By analysing tail bend kinematics in head-restrained larvae, we show that the increase in speed is actuated by larger tail bends. Whole cell patch clamp recordings from motor neurons reveal that during OMR, typically only secondary motor neurons are active while primary motor neurons are quiescent. Activation of D1-like receptors increases motor drive from secondary motor neurons by decreasing spike threshold and latency. In addition, D1-like receptor activation enhances excitability and recruits quiescent primary motor neurons. Our findings provide an example of neuromodulatory reconfiguration of spinal motor neuron speed modules such that members are selectively recruited and motor drive is increased to effect changes in locomotor speed.

Keywords

Graphical abstract

Introduction

Survival for most animals depends on their ability to adapt speed, gait and direction of movement quickly in response to sensory stimuli. In vertebrates, these processes are controlled by the concerted activity of distributed circuits in the brain and the spinal cord. In particular, the regulation of speed by spinal locomotor circuits has received much attention recently. The ability to genetically label specific neuronal populations and to target them for imaging or electrophysiology has allowed us to identify recruitment patterns of interneurons and motor neurons at different speeds. Studies in mice and zebrafish have identified subtypes of spinal interneurons and motor neurons that are selectively recruited at different speeds of locomotion [1–8]. These interneurons and motor neurons make selective synaptic connections among them and are thought to constitute 'slow' or 'fast' network modules, that are sequentially recruited as the locomotor speed increases [9,10]. Thus, behaviors such as escapes, which involve fast swimming, are generated by the fast module, and spontaneous swimming behaviors, occurring at much lower swim frequencies are controlled by the slow module [11].

Although this modular view of spinal network operation is widely acknowledged [12,13], we also know that the spinal network is the target of a variety of neuromodulators [14–17]. Neuromodulators have been shown to fuse independent circuits and to construct 'de novo' circuits from cells belonging to disparate circuits [18,19]. We wished to understand if swim speed regulation in zebrafish larvae is subject to neuromodulation and if yes, whether the rules for modular recruitment are altered by neuromodulators.

Dopamine has long been known to be a key neuromodulator of motor circuits [17,20] and is critical for activation, maturation and modulation of locomotor patterns

across invertebrates and vertebrates [21–26]. In zebrafish, dopaminergic input from

the diencephalon to the spinal cord develops early [27] and is known to regulate

developmental maturation of swim motor patterns [24]. Nevertheless, the role of

spinal dopamine in regulating speed of locomotion has thus far not been investigated

in any vertebrate.

We chose to study speed regulation and its dopaminergic modulation, during

the optomotor response (OMR), an innate behaviour evoked by whole-field visual

motion, to which zebrafish larvae respond by turning towards and moving in the

direction of perceived motion. By adjusting their locomotor speed to match the speed

of the perceived motion, larvae try to maintain a stable position with respect to their

surroundings [28,29]. However, the spinal mechanisms by which speed is controlled

during OMR have thus far not been investigated.

We used a combination of behavioral monitoring in free-swimming larvae,

kinematic characterization in head-restrained larvae and whole-cell patch clamping

in immobilized larvae to ask how speed is controlled during OMR and how it is

modulated at the neuronal level. Here, we show that dopamine acting via D1-like

receptors (D1-like-R) increases the speed of swimming during OMR by increasing

the extent of tail bending. This in turn is caused by two factors: (1) increased firing of

action potentials in motor neurons of the slow module, typically recruited during OMR

and (2) novel recruitment of motor neurons belonging to the fast module, not typically

active during OMR. Thus, dopamine selectively activates and recruits neurons from

disparate spinal modules to increase speed. Our results suggest that activity and

recruitment pattern of neurons in these circuit modules are not only defined by their

cellular   properties   and   synaptic   connectivity.   Neuromodulatory   inputs   can

dramatically alter the dynamics of these modules by recruiting neurons that would

have otherwise not participated under a given context.

Results

D1-like-R activation increases swim speed during OMR.

OMR was evoked in freely swimming zebrafish larvae between 6-7 days post

fertilization (dpf) by presenting square-wave gratings moving in a clockwise direction

from below (Figure 1A , Video S1). Zebrafish larvae respond by following the

direction of the stimulus and matching their swim speed with that of the moving

grating [29]. Larval centroid was tracked to extract kinematic parameters. Zebrafish

larvae swim intermittently in a beat-and-glide pattern where brief periods of discrete

tail oscillations (bout) are followed by a period of no tail movement [30]. We further

segmented tracked data to identify individual swim bouts (Figure 1B).

We compared average swim speed (including both bouts and inter-bout

quiescent periods) during OMR before and after bath application of 20 µM D1-like-R

agonist, SKF-38393. Larvae swam with higher average swim speed post D1-like-R

activation (Figure 1C, D, Video S1). We then asked if D1-like-R activation increases

speeds achieved during bouts and compared average bout speed before and after

bath application of D1-like-R agonist. Average bout speed increased (Figure 1E)

resulting in greater distance travelled per bout (Figure 1F) post application of D1-like-

R agonist.

To further confirm that the increase in swim speed is a result of D1-like-R

activation and not an artefact of repeated stimulus presentation, we also compared

these parameters before and after bath application of 20 µM D1-like-R antagonist,

SCH-23390. Average swim speed decreased post application of D1-like-R

antagonist (Figure 1G, H). Consistently, a decrease in average bout speed (Figure

1I) and average distance travelled per bout (Figure 1J) were observed post

application of D1-like-R antagonist.

D1-like-R signaling is associated with increased spontaneous locomotor

activity in larval zebrafish [24,31,32]. Similar to spontaneous swimming, we find D1-

like-R signaling to have a stimulatory effect on the motor output during OMR.

Importantly, these results indicate that locomotory speed during an innate and

reflexive behavior is subject to neuromodulation via D1-like-R activation.

D1-like-R signaling controls swim speed by modulating tail beat amplitude.

We sought to further understand which kinematic variables are altered by D1-like-R

signaling to bring about the increase in bout speed we observe. For characterization

of kinematic variables, we evoked OMR in a head-restrained preparation which

allows for a more accurate and detailed tracking of tail kinematics.

In response to caudal-to-rostral moving gratings, partially restrained larvae

reliably performed forward OMR characterized by mostly symmetrical, alternating

left-right tail beats (Figure 2A, Video S2). Tail-tip position was tracked to extract tail

beat amplitudes, tail beat frequency and duration of bouts (Figure 2B). As DA has

been shown to modulate swim initiation in larval zebrafish [33], we also quantified

the number of swim bouts initiated.

Activation of D1-like-R increased mean tail beat amplitude and tail beat

frequency of forward swims (Figure 2C-E, Video S2). However, no difference in

duration of bouts or number of bouts initiated was observed post application of 100

μM D1-like-R agonist, SKF-38393 (Figure 2F, G). In contrast, application of 100 μM

D1-like-R antagonist, SCH-23390, decreased mean tail beat amplitude of forward

locomotion (Figure 2H, I). Tail beat frequency, bout duration and number of swim

bouts initiated did not change significantly (Figure 2J-L).

Modulation of tail beat amplitude is a key factor contributing to control of

swim speed [34,35]. Thus, by modulating tail beat amplitude D1-like-R activation

results in faster swim bouts during forward OMR.

Activation of D1-like-R results in larger tail bend angle during turns.

Free swimming OMR behavioural assay required larvae to swim both in a straight

trajectory as well as make routine turns (Video S1). Since we observed an increase

in both average swim speed and average bout speed after D1-like-R activation

during free-swimming OMR (Figure 1), we surmised that D1-like-R activation will

likely affect tail bend kinematics during routine turns also. To investigate the role of

D1-like-R signaling in modulation of turning behavior, we presented leftward and

rightward moving gratings to head-restrained larvae (Figure 3A). Larvae responded

to the stimuli with asymmetric tail deflections in the direction of grating movement

(Figure 3A, Video S3). Tail-tip position was tracked and maximum tail bend angle

was measured before and after application of 100 µM D1-like-R agonist and

antagonist (Figure 3B).

Larvae performed turns with larger tail bend angles post application of D1-

like-R agonist (Figure 3C, E, Video S3). In contrast, application of the D1-like-R

antagonist showed a decrease in the magnitude of the tail bend angle (Figure 3D, F).

Taken together, these results indicate that D1-like-R activation modulates the

extent of tail bends during both forward locomotion and turns resulting in greater

displacement.

Activation of D1-like-R intensifies drive from 'slow' motor neurons

D1-like-R mediated increase in swim speed could result from an increase in the firing of active motor neurons and/or by the recruitment of new cells to the active pool [1]. Zebrafish axial motor neurons can be broadly divided into two classes: primary and secondary [36]. Early born, dorsally located primary motor neurons innervate fast muscle fibres and are active during fast swimming, struggle and escape responses swimming whereas later born, ventral secondary motor neurons are active during slower swimming [1,37]. Besides anatomical position and axonal innervation pattern, the two classes of motor neurons can also be clearly distinguished based on their cellular properties and response to injected current [38] (Table S1, Figure S1). As secondary motor neurons are active during low-frequency swimming such as those observed in OMR , we first asked if they are modulated by D1-like-R. We obtained whole-cell patch clamp recordings from dorsal secondary motor neurons while presenting moving gratings from below to evoke forward OMR (Figure 4A). 7/9 recorded dorsal secondary motor neurons fired bursts of action potentials in response to moving grating presentation (Figure 4B). An increase in the number of action potentials generated during moving gratings presentation was observed post application of 20 µM D1-like-R agonist (Figure 4C, G). Interestingly, we also observed motor neurons otherwise quiescent during visual stimulus presentation (2/9), fire action potentials post application of D1-like-R agonist (Figure 4D, E). D1-like-R agonist mediated effects were antagonized in 4/4 secondary motor neurons post subsequent application of D1-like-R antagonist (Figure 4F, G). Number of action potentials spontaneously generated during stationary grating presentation did not differ significantly before and after application of D1-like-R agonist (Figure S2A).

Each motor neuron burst has been shown to correspond to motor bursts in

fictive swims and is a measure of swim frequency [2,38]. Further analysis of bursts

showed an increase in average number of spikes per burst without a significant

increase in the number of bursts post D1-like-R activation (Figure 4H, I). Voltage

oscillations underlying these bursts reflect synaptic drive to these motor neurons

(Figure 4J; [38]). Average frequency of these oscillations did not differ significantly

post application of D1-like-R agonist (Figure 4K). This suggests that upstream

rhythm generating circuits are unlikely to be modulated by D1-like-R signaling.

These results indicate that activation of D1-like-R signaling increases swim

speed by enhancing drive from secondary motor neurons. This in turn is mediated by

recruitment of quiescent neurons and increased firing per swim cycle of already

active neurons. These changes point to D1-like-R mediated increase in the

excitability of secondary motor neurons, although an increase in synaptic drive

during OMR cannot be ruled out.

D1-like-R activation reduces spike latency and action potential threshold of

secondary motor neurons.

To test if D1-like-R signaling directly affects the excitability of secondary motor

neurons, or if it engages synaptic drive operating during OMR, we asked if

secondary motor neurons show changes in firing behavior in response to direct

current injection when D1-like R agonist is applied. We analyzed firing responses of

secondary motor neurons to a series of depolarizing current steps (25 pA-400pA, 25

pA step size) before and after application of 20 μM D1-like-R agonist (Figure 5A).

D1-like-R agonist significantly reduced first spike latency (Figure 5B, C). The

decrease in spike latency was observed for all current steps and was less

pronounced when injected currents were high (Figure 5D).

D1-like-R activation also reduced the voltage threshold for the first action

potential generated (Figure 5 B, E).

We observed no significant effect of D1-like-R activation on rheobase, input

resistance and instantaneous firing rate of first two spikes (Figure 5 F-H).   Resting

membrane potential, spike afterhyperpolarization (AHP) and spike-width did not

change post application of D1-like-R agonist (Figure S3).

These results show that D1-like-R activation increases secondary

motoneuronal excitability by reducing spike latency and action potential threshold.

Quiescent primary motor neurons are recruited during OMR by D1-like-R

activation.

As primary motor neurons are active only during fast and vigorous swimming [1,37],

we next examined if D1-like-R signaling can modulate primary motor neuron's

excitability as well to bring about faster locomotion. We obtained whole-cell patch

clamp recordings from primary motor neurons while presenting drifting gratings from

below to evoke OMR (Figure 6A). Primary motor neurons usually showed sub-

threshold responses (Figure 6C) or were quiescent (Figure 6F) during visual stimulus

presentation. Bath application of 20 µM D1-like-R agonist increased excitability of

primary motor neurons. 6/12 primary motor neurons recorded showed recruitment

during moving grating presentation post application of D1-like-R agonist (Figure

6D,G). Subsequent application of D1-like-R antagonist on 4 motor neurons reduced

the number of action potentials generated in all these cells (Figure 6E, H). Primary

motor neurons are further classified as caudal primary (CaP), middle primary (MiP)

and rostral primary (RoP) based on the rostrocaudal position of their somata. Of the six primary motor neurons that were recruited, 3 were identified as CaP, 2 as RoP and 1 as MiP. In contrast, number of action potentials spontaneously generated during stationary grating presentation did not differ significantly before and after application of D1-like-R agonist (Figure S2B).

Frequency of voltage oscillations underlying bursts (Figure 6I,J) were found to be similar to swimming frequency at which primary motor neuron have been shown to be recruited [1]. These results show that D1-like-R signaling recruits novel neurons from the 'fast' motor pool to increase the speed of locomotion during OMR.

D1-like-R activation increases excitability of primary motor neurons.

Next, we investigated the mechanism by which D1-like-R activation recruits primary motor neurons during OMR. Responses to a series of depolarizing current steps were obtained from primary motor neurons. Bath applications of 20 µM D1-like-R agonist were associated with an increase in the number of evoked action potentials (Figure 7A). D1-like-R activation modulated several membrane properties to increase the excitability of primary motor neurons. Spike analysis revealed a decrease in first spike latency (Figure 7B, C). Significant reduction in spike latencies were observed for all current steps injected (Figure 7D).

D1-like-R activation also reduced threshold for action potential generation (Figure 7B, E) and rheobase (Figure 7F). However, a decrease in input resistance was observed post applications of D1-like-R agonist (Figure 7G). Modulation of these membrane properties significantly increased the instantaneous firing rate of first two spikes (Figure 7H), suggesting that with D1-like-R activation, the same synaptic drive to primary motor neurons could still elicit stronger firing. Together, our

analysis of secondary and primary motor neuron firing properties in response to

grating stimuli and current injections reveal a stimulatory effect of D1-like-R

signaling. Activation of D1-like-R increases the number of action potentials fired per

burst by decreasing spike latency and spike threshold. The additional spikes fired

per burst will act to increase the extent of tail bends, thus increasing the

displacement generated in the same amount of time. While not ruling out actions of

D1-like-R elsewhere in the motor hierarchy, these effects at the level of motor

neurons provide a parsimonious mechanism for the speed regulation observed

during OMR.

Discussion

Neuromodulatory selection of novel neurons

Zebrafish larvae generate slow swims when presented with slowly drifting gratings.

During this behavior they activate secondary motor neurons belonging to the 'slow'

module, while primary motor neurons belonging to the 'fast' module are quiescent.

However, when D1-like receptors are activated, primary motor neurons are recruited

and secondary motor neurons fire more action potentials, resulting in larger tail

bends and therefore greater displacement and faster swims. Our results

demonstrate that even under conditions where the sensory input is constant,

neuromodulators can remap sensorimotor transformations, such that the behavioral

output is altered. In addition, our experiments underline the fact that the pattern of

recruitment of neurons during behaviors is heavily dependent on the

neuromodulatory context. Recent advances in microscopy and the development of

novel neuronal activity sensors have allowed us to map neuronal circuits that are

active during a variety of behaviors. Our study serves as a reminder that such maps

are a function of the neuromodulatory status of the animal.

Dopaminergic modulation of motor neuronal excitability

In zebrafish, spinal dopaminergic projections arise solely from the A11 cluster, also

known as dopaminergic diencephalospinal neurons (DDNs) [27,33]. Dopaminergic

neuromodulation underlies developmental maturation of swim motor patterns [23]

and in particular, the DDNs have been shown to mediate the switch from long to

short swim bouts [24]. DDNs are more likely to fire bursts of action potentials during

swim bouts and fire tonically during periods of rest [33]. Since burst firing could

potentially lead to higher calcium influx into presynaptic terminals, more dopamine is

likely to be released by bursting DDNs onto their spinal targets during periods of

active locomotion. These findings lend support to our view that endogenous

dopamine is capable of altering locomotor speeds *in vivo* in the freely moving

animal.

Tyrosine hydroxylase (TH) immunoreactive fibers in the spinal cord make

contact with both primary and secondary motor neurons [39] implying direct

modulation of motoneuronal excitability by A11 dopaminergic neurons. We show that

activation of D1-like receptors enhances the excitability of primary and secondary

motor neurons, most likely by acting directly on them. Importantly, we show that both

spike latency and threshold are decreased, suggestive of direct dopaminergic

modulation of the transient A-type potassium conductance (I_A). Indeed, dopaminergic

reduction of I_A leading to a decrease in first spike latency has been reported in

invertebrate [40] and vertebrate [41] motor neurons. In addition, in primary motor

neurons neuronal gain is increased, suggesting modulatory actions on voltage

dependent conductances [42]. These results complement previous studies on

mechanistic bases of speed control. The 'size principle' states that neurons are

sequentially recruited as speed increases such that the smaller, high input resistance

secondary motor neurons get recruited at slow speeds and the larger, low input

resistance primary motor neurons are recruited at high speeds. To match this

recruitment order, gradations in endogenous rhythmicity have also been observed

[38]. In addition, at higher speeds, primary motor neurons preferentially receive

enhanced excitation from pre-motor interneurons to enable their recruitment [43]. In

contrast, dopaminergic neuromodulation does not seem to tap into any of these

previously identified mechanisms of speed regulation. Instead, the neuromodulatory

regulation of speed that we identified appears to target intrinsic spike generation

mechanisms directly to effect recruitment. Although we show specific modulation of

intrinsic properties of motoneurons, modulation of spinal and supra-spinal pre-motor

circuits cannot be ruled out.

Regulation of speed during OMR by spinal interneurons

The recruitment patterns of spinal pre-motor interneurons during OMR has thus far

not been documented. Nevertheless, recruitment patterns across a range of

locomotor speeds in other contexts have been reported in both larval and adult

zebrafish. In adult zebrafish, premotor V2a interneurons connect preferentially to

slow, intermediate or fast motor neurons forming distinct speed modules that are

sequentially recruited as speed increases [8–10]. However, in larval zebrafish,

premotor interneurons active at slow speeds are silenced as swim speed increases,

and distinct sets of interneurons are recruited from quiescence at faster speeds [2].

Such a phenomenon has also been reported for speed-related gait changes during limbed locomotion in mice [3–5,44]. In our studies, the dopaminergic modulation of speed during OMR could be augmented by enhanced suppression of slow-speed interneurons and excitation of fast-speed interneurons. Selective suppression of slow-speed interneurons [45] or activation of fast-speed interneurons [46] are known to play roles in switching swim speeds. Further experiments targeting these interneuronal types should throw light on whether they are targets of dopaminergic neuromodulation.

Supraspinal circuits governing the OMR behavior

OMR relies on circuits distributed across the nervous system. Retinal input, including responses from direction-selective RGCs, arrives in arborization fields (AF) 5 and 6, which are innervated by the dendrites of pre-tectal neurons. Pretectal neurons show direction-selective tuning, respond to binocular optic flow and send long range projections to the cerebellum and the hindbrain reticular formation [47,48], in turn leading to the generation of the appropriate swim command in the spinal cord. Several hindbrain neurons are activated when the fish perform OMR [50]. Among these, neurons in the nucleus of the medial longitudinal fasciculus (nMLF) are active during forward OMR and the ventromedial spinal projection neurons (vSPNs) are active during turns [49,51]. The nMLF makes direct synaptic contacts with primary and secondary motor neurons [52], regulates swim speed during OMR [29] and receives dense innervation from TH-positive fibers [53].

Recently, neurons in the posterior tuberal nucleus (PTN), a sub-group of the A11 dopaminergic cluster, were identified to show large calcium transients in response to forward moving gratings in larval zebrafish [54]. The activation of PTN dopaminergic

neurons was delayed by many seconds after optic flow started and lasted for several

seconds after it was turned off, suggesting a slow build-up of the modulatory drive.

Further, activity in the PTN neurons was suppressed following motor bouts implying

a feedback mechanism that limits the dopaminergic modulation. Projections from

PTN are restricted to the hindbrain and could potentially be the source of TH-fibers

projecting to nMLF. In lamprey, dopamine released from the posterior tuberculum

activates reticulospinal cells to increase locomotor output through a D1 receptor-

dependent mechanism [55]. Taken together, our results, in the context of the above

studies, suggest that during OMR, dopaminergic neurons in the A11 cluster act to

increase swim speed via their concomitant modulatory actions at the spinal and

supraspinal levels.

Acknowledgements

The authors would like to thank the following sources of funding support: Wellcome

Trust-DBT India Alliance Intermediate and Senior fellowships (VT), Department of

Biotechnology (VT), Science and Engineering Research Board, Department of

Science and Technology (VT), Department of Atomic Energy (VT), and CSIR-UGC

fellowship (UJ). The authors would also like to thank Sriram Narayanan, Pushkar

Paranjpe and Dilawar Singh for technical assistance, and T.P. Jagadeesh for

maintaining the fish facility.

Author Contributions

Designed experiments (UJ and VT); Performed experiments (UJ); Analyzed data (UJ

and VT); Wrote manuscript (UJ and VT).

Declaration of Interests

The authors declare that they have no competing interests. The funding agencies did

not have any role in the design or outcomes of this study.

Figure legends

Figure 1. D1-like-R activation increases swim speed during OMR in free-

swimming zebrafish larvae.

(A) Schematic of experimental set up. OMR was evoked in freely swimming

zebrafish larvae (6-7 dpf) by presenting radial gratings moving in a clockwise

direction on a screen below the arena. Zebrafish larvae respond to the stimulus by

swimming in the direction of moving grating. Videos were acquired from above and

larval centroid was tracked. **(B)** *Left,* instantaneous speed of a representative larva

during a trial. *Right,* zoomed view of region marked with the black rectangle showing

characteristic intermittent swimming pattern. Brief period of activity represents

individual swim bouts (highlighted by line on top) and is followed by a period of no

tail movement. **(C)** Overlaid tracked centroid for a trial duration of 60 seconds

showing trajectory of a representative larva before (Control) and after bath

application of 20 µM D1-like-R agonist, SKF-38393. Scale bar represents 10 mm.

(D-F) Paired plots for average speed, average bout speed and average bout

distance respectively, before (black) and after (blue) application of D1-like-R agonist.

(G) Overlaid tracked centroid of a representative larva for a trial duration of 60

seconds before (Control) and after bath application of 20 µM D1-like-R antagonist,

SCH-23390. Scale bar represents 10 mm. **(H-J)** Paired plots for average speed,

average bout speed and average bout distance respectively, before (black) and after

(red) application of D1-like-R antagonist. $N_{SKF-38393}$=14 larvae, $N_{SCH-23390}$=9 larvae,

Wilcoxon signed rank test; ** p<0.001. See also Video S1.

Figure 2. D1-like-R signaling modulates tail beat amplitude during forward

OMR in head restrained larvae.

(A) Schematic representation of experimental set-up. Z-projection of a swim bout

showing characteristic alternating left-right tail beats evoked in a head-restrained

larval zebrafish (top) in response to caudal-to-rostral moving gratings (bottom). Tail-

tip position was tracked for measurement of kinematic variables. (**B**) *Left,*

representative tracked tail-position for a trial. *Right,* zoomed view of highlighted bout

showing kinematic variables used for behavioral quantification. **(C)** Overlaid tracked

tail-tip position for a representative trial before (top) and after (bottom) application of

100 μM D1-like-R agonist, SKF-38393. Summary data of **(D)** average tail beat

amplitude, **(E)** average tail beat frequency, **(F)** average bout duration and **(G)**

number of bouts initiated before (black) and after (blue) bath application of D1-like-R

agonist. **(H)** Overlaid tracked tail-tip position for a representative trial before (top) and

after (bottom) application of 100 μM D1-like -R antagonist, SCH-23390. Summary

data of **(I)** average tail beat amplitude, **(J)** average tail beat frequency, **(K)** average

bout duration and **(L)** number of bouts initiated before (black) and after (red) bath

application of D1-like-R antagonist. Scale bar represents 1mm. $N_{SKF-38393}$=37 larvae,

$N_{SCH-23390}$=18 larvae, Wilcoxon signed rank test; *** p<0.00001, * p<0.05, ns: not

significant. See also video S2.

Figure 3. D1-like-R activation increases tail bend angle during turns.

(A) Schematic representation of experimental set-up showing a turn characterized by asymmetric tail bend about the midline (top) evoked using sideward moving vertical bars (bottom) in a head-restrained zebrafish larva. Tail-tip position was tracked for measurement of tail bend angle. **(B)** *Left,* tail bend angle for a representative trial. Maximum tail bend angle for each turn is highlighted with a solid circle. *Right,* snapshot of a turn showing references used for calculation of tail bend angle. **(C)** Overlaid tracked tail-tip position during a trial before (left) and after (right) bath application of 100 µM D1-like-R agonist, SKF-38393. **(D)** Overlaid tracked tail-tip position during a trial before (left) and after (right) bath application of 100 µM D1-like-R antagonist, SCH-23390. **(E-F)** Summary data of average tail bend angle before and after application of D1-like-R agonist and antagonist. Scale bar represents 1mm. $N_{SKF-38393}$=29 larvae, $N_{SCH-23390}$=26 larvae, Wilcoxon signed-rank test; **p<0.001, *p<0.05. See also video S3.

Figure 4. D1-like-R activation enhances drive from secondary motor neurons during OMR.

(A) Schematic representation of experimental set-up. Whole-cell patch clamp recordings from secondary motor neurons were obtained while presenting 10s of stationary gratings alternating with 10s of forward moving gratings. **(B)** *Left,* recording from a representative secondary motor neuron during a trial. Region under small black line is shown in a zoomed view on the *right.* Shaded areas represent timing of moving gratings presentation. **(C)** *Left,* recording from the same neuron as in **(B)** post application of 20 µM D1-like-R agonist, SKF-38393. Region under the

small blue line is shown expanded on the *right*. **(D)** Representative recording from a secondary motor neuron that was quiescent during moving gratings presentation before any drug application (black). **(E)** Recording from the same secondary motor neuron as in **(D)** after subsequent application of D1-like-R agonist (blue) and **(F)** D1-like-R antagonist (red). **(G)** Paired plot of number of action potentials fired during moving grating presentation before (black), after application of D1-like-R agonist (blue) and post subsequent application of D1-like-R antagonist (red). **(H)** Paired plot of average number of spikes per burst before (black) and after application of D1-like-R agonist (blue) **(I**) Paired plot of number of bursts during moving gratings presentation before (black) and after (blue) application of D1-like-R agonist, SKF-38393. **(J)** *Top,* representative recordings from a secondary motor neuron during moving grating representation before (black) and after (blue) application of D1-like-R agonist. *Bottom,* zoomed view of region marked with black rectangle showing voltage oscillations (highlighted in pink). **(K)** Average frequency of voltage oscillations before (black) and after (blue) application of D1-like-R agonist. N=9 cells from 9 larvae, Wilcoxon signed-rank test; *p<0.05, ns: not significant. See also Table S1, Figure S1, Figure S2.

Figure 5. D1-like-R activation reduces spike latency and action potential threshold of secondary motor neurons.

(A) Schematic of experimental set up. Region marked by the red rectangle is shown below. *Right,* Responses of a secondary motor neuron to a series of depolarizing current steps were compared before (Control) and after application of 20 µM D1-like-R agonist, SKF-38393. **(B)** Representative traces before (black) and after (blue) application of D1-like-R agonist. Vertical dotted lines indicate spike latency and

horizontal dotted lines mark threshold for action potential generation before (black) and after (blue) D1-like-R agonist application. **(C)** Summary data for first spike latency. Wilcoxon signed-rank test; **p<0.01 **(D**) Summary data of spike latency for each step of depolarizing current injected. Mixed-Effects model; p<0.0001 **(E)** Summary data for threshold of action potential generation. Wilcoxon signed-rank test; *p<0.05. **(F)** Summary data for rheobase. Paired ttest, p=0.59 **(G)** Summary data for input resistance. Wilcoxon signed-rank test; p= 0.07 **(H)** Instantaneous firing frequency of the first two spikes measured in response to series of current steps before (black) and after (blue) application of D1-like-R agonist. Mixed-Effects model; p= 0.822. ns: not significant. N=9 secondary motor neurons recorded from 9 larvae. Error bars represent SEM. See also Figure S3.

Figure 6. Quiescent primary motor neurons are recruited during OMR by D1-like-R activation. (A) Schematic representation of experimental set-up. Whole-cell patch clamp recordings from primary motor neurons were obtained while presenting 10 s of stationary gratings alternating with 10 s of forward moving gratings. Motor neurons were targeted in the region marked by the red rectangle. Zoomed view of the same region is shown on the right. **(B)** Number of action potentials fired by primary motor neurons during forward moving grating presentation before (black), after bath application of 20 µM D1-like-R agonist (blue) followed by bath application of 20 µM D1-like-R antagonist (red) **(C-E)** Representative recordings from a rostral primary motor neuron (RoP) before any drug application (black), after application of D1-like-R agonist (blue) followed by D1-like-R antagonist (red) application. **(F-H)** Recordings from a middle primary motor neuron (MiP) before any drug application (black), after application of D1-like-R agonist (blue) followed by D1-like-R antagonist

(red) application. Shaded areas represent timing of moving gratings presentation. **(I)**

Left, recording from a caudal primary motor neuron (CaP) during forward moving

grating presentation post application of D1-like-R agonist. *Right,* zoomed view of the

highlighted region showing rhythmic voltage oscillations (marked in pink). **(J)**

Distribution of frequency of rhythmic oscillations from all the primary motor neurons

that were recruited post application of D1-like-R agonist. N=12 primary motor

neurons recorded from 12 larvae, Wilcoxon signed-rank test; *p<0.05. See also

Table S1, Figure S1, Figure S2.

Figure 7. D1-like-R activation increases excitability of primary motor neurons.

(A) Schematic of experimental set up. Region marked by the red rectangle is shown

below. Responses of a representative primary motor neuron to a series of

depolarizing current steps were compared before (Control) and after application of

20 µM D1-like-R agonist, SKF-38393. **(B)** Representative traces before (black) and

after (blue) application of D1-like-R agonist. Vertical dotted lines indicate spike

latency, horizontal dotted lines mark threshold for action potential generation and

lines on top indicate first inter-spike interval (F_{inst}) before (black) and after (blue) D1-

like-R agonist application. **(C)** Summary data for first spike latency. Wilcoxon

signed-rank test; *p<0.05 **(D**) Summary data of spike latency for each step of

depolarizing current injected. Mixed-Effects model; *p<0.05 **(E)** Summary data for

threshold of action potential generation. Wilcoxon signed-rank test; ***p<0.001. **(F)**

Summary data for rheobase. Wilcoxon signed-rank test; *p<0.05. **(G)** Summary data

for input resistance. Wilcoxon signed-rank test; *p<0.05. **(H)** Instantaneous firing

frequency of the first two spikes measured in response to series of current steps

before (black) and after (blue) application of D1-like-R agonist. Mixed-Effects model;

*p<0.05. N=12 primary motor neurons recorded from 12 larvae. Error bars represent

SEM. See also Figure S4.

Methods

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All experiments were approved by the institutional animal ethics committee and the

institutional biosafety committee. Experiments were performed on Indian wild type

zebrafish (*Danio rerio)* larvae between 6 and 7 days post fertilization (dpf) at room

temperature (25-28 °C)*.* Larvae have not undergone sex specification at these

stages. Larvae were maintained in 14:10 light-dark cycle at 28 °C in E3 medium

(composition in mM: 5 NaCl, 0.17 KCl, 0.33 $CaCl_2$, and 0.33 $MgSO_4$, pH 7.8).

METHOD DETAILS

Free swimming behavior assay

Zebrafish larvae swam freely in a circular arena 55 mm in diameter. A block of

agarose in the centre of the arena restricted swimming in a narrower lane 20 mm in

width. Videos were acquired at 120 fps using a CMOS camera (FL3-U3-13-S2M-

CS, Point Grey, Richmond, USA) with IR illumination provided from below. A long

pass filter (715 nm colored glass filter, Thorlabs, New Jersey, USA) was placed in

front of the camera to remove the visual stimulus. Radially spinning square wave

gratings were displayed on a screen underneath the arena (Figure 1A). Stimulus

presentation was performed using Processing [56]. Acquisition of videos and

tracking of larval centroid was performed using Bonsai [57].

Average speed was calculated as total distance travelled in trial duration of 60

s. Bout start and end indices were identified from a plot of instantaneous speed vs

time (Figure 1B). Distance travelled within a bout was calculated as bout distance

and was divided by duration of respective bout to calculate bout speed. Average of

these parameters across all bouts per larva was then calculated to give average bout

distance and average bout speed.

Head-restrained behavior assay

Larvae were embedded in 1.5% low gelling agarose (Sigma-Aldrich; Missouri, USA).

E3 medium was added after the agarose had congealed. Agarose around the tail

leaving pectoral fins free was removed for observing tail movements. Optomotor

response was evoked by presenting black and white gratings moving with spatial

period 10 mm and temporal period 10 mm/s. Visual stimulus was projected on a

screen below the larva using a DLP projector (Acer C120; Acer Inc., New Taipei City,

Taiwan) after reflection by a cold mirror (FM203; Thorlabs, New Jersey, USA). The

arena was IR illuminated from below. Videos were acquired at 500 fps using a high-

speed camera (Phantom Miro eX4; Vision research, New Jersey, USA) fitted with a

zoom lens (AF Nikkor 60 mm f/2.8D, Nikon, Tokyo, Japan) and infrared pass filter

(768 nm, Pixelteq, Florida, USA). Each trial consisted of linearly drifting grating

presentation for 12s. Three trials before and four trials after the drug application were

performed.

Data were analyzed using custom written scripts in ImageJ, IPython [58] and

MATLAB (Mathworks, Natick, USA). Videos were processed in Fiji [59] to obtain

larval mid-line. A* search algorithm was used to track tail-tip position using custom

written macros in ImageJ. Tail-tip position was used for extracting kinematic

parameters. Analysis of kinematic variables was performed offline. Struggles were identified as vigorous body movements with large tail bends and were removed from subsequent analyses. Tracking data was segmented into swim bouts (Figure 2B, left). Amplitude was defined as the maximum (peak-to-trough) tail tip displacement in a bout. Time difference between movement onset and offset of a bout was defined as bout duration (Figure 2B, right). These variables were extracted for individual bouts and average values across all bouts per larva were calculated. Tail beat frequency was defined as the time taken to perform a complete oscillation and was calculated using fast fourier transform. Tail bend angle was calculated using head and tail-tip coordinate. Maximum tail bend angle for each turn was extracted (Figure 3B) and average maximum tail bend angle per larva was calculated.

Electrophysiology

Whole-cell patch clamp recordings from motor neurons were obtained while presenting caudal-to-rostral moving gratings on a screen below the stage. Larvae were anesthetized in 0.01% MS-222 (Sigma-Aldrich; Missouri, USA) and transferred to a recording chamber. The larvae were pinned down through notochord using fine tungsten wire (California Fine Wire, CA, USA). The MS222 was then replaced by external solution (composition in mM: 134 NaCl, 2.9 KCl, 1.2 $MgCl_2$, 10 HEPES, 10 glucose, 2.1 $CaCl_2$, 0.01 D-tubocurarin; pH 7.8; 290 mOsm) and skin along the tail was carefully removed using forceps (Fine Science Tools, Foster City, USA). Muscles in a hemi-segment (between the 10^{th} and 13^{th} myotomes) were carefully removed to expose the spinal cord. Cells were targeted at 4X magnification with a 60X/1.0 numerical aperture (NA) water-immersion objective (LUMPlanFL) on a

compound microscope (BX61WI, Olympus; Tokyo, Japan) with differential

interference optics (DIC). Patch pipettes were made using thick walled borosilicate

capillaries (1.5 mm OD;0.86 mm ID; Warner Instruments, Hamden, CT, United

States) pulled to 1-1.5 μm tip diameter using a Flaming brown P-97 pipette puller

(Sutter Instruments, Novato, CA, United States). Pipettes were backfilled with K-

gluconate internal solution (composition in mM: 115 K gluconate, 15 KCl, 2 $MgCl_2$,

10 HEPES, 10 EGTA, 4 ATP Disodium salt; pH 7.2; 290 mOsm) and typically had

resistances between 8 and 10 MΩ. Sulforhodamine was added to the patch internal

solution for visualization of cell-morphology and axon tracts.

Motor neurons were targeted blind and classified as primary or secondary

post hoc based on their cellular properties (Table S1) and responses to current

injections. Primary motor neurons fired tonically whereas the dorsal secondary motor

neurons showed chattering response to current injections (Figure S1). Six trials of

visual stimulation were presented, with each trial consisting of 10s of stationary

gratings alternating with 10s of forward moving gratings. Visual stimuli were

displayed on an LCD screen (Waveshare, Hongkong) driven by Raspberry Pi (Model

B RASP-PI-3, Raspberry Pi Foundation, Cambridge, UK). Visual stimulus

presentation and electrophysiological recordings were triggered simultaneously and

acquired using Multiclamp 700B amplifier, Digidata 1440A digitizer and pCLAMP

software (Molecular Devices, San Jose, USA). Corrections for bridge balance were

applied in current clamp mode. Membrane potentials are reported without correction

for liquid junction potential which was measured to be +8 mV for the external and

internal solutions mentioned above. The data were low-pass filtered at 2kHz using a

Bessel filter and sampled at 20kHz at a gain of 1.

Data were analyzed using Clampfit (Molecular Devices, San Jose, USA) and

MATLAB (Mathworks, Natick, USA). Clampfit was used to identify spikes and bursts.

To calculate the frequency of oscillations underlying bursts, recordings for trials

where forward grating was presented, was first filtered to remove spikes and low

frequency periodic depolarization. Frequency with maximum power post fast fourier

transform of the filtered recording was used to estimate oscillation frequency for a

trial.

To determine intrinsic firing pattern of motor neurons, 200 ms long

depolarizing current pulses in the range (25-400 pA, step size 25 pA) were applied in

absence of visual stimulus presentation. First spike latency was defined as the time

between onset of the current step and the peak of the first spike. Spike threshold

was defined for the first spike at rheobase as the membrane potential where the

speed of voltage change reaches 5mV/ms. Rheobase was defined as the minimum

current required to elicit an action potential. Input resistance was calculated by

injecting 200 ms long hyperpolarizing pulses (-100pA to -25 pA).

Drugs

Drug concentrations for each experiment was determined in pilot experiments (data

not shown). Drugs were dissolved in E3 for behavioral experiments and in external

solution for electrophysiological experiments.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data were tested for normality using one sample Konglomorov-Smirnov test. Paired

ttest or Wilcoxon signed rank test were performed for comparisons between the two

groups in MATLAB. Mixed effects model using nmle package [60] in R was used to

test the effect of drug on latency and firing frequency for different values of current

injected. Significance level was defined as 0.05. Details on statistical test, p values

and sample size can be found in figure legend.

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, Peptides, and Recombinant Proteins		
SKF-38393 hydrochloride	SIGMA-ALDRICH	D047; CAS:62717-42-2
SCH-23390 hydrochloride	SIGMA-ALDRICH	D054; CAS:125941-87-9
Adenosine 5'-triphosphate disodium salt	SIGMA-ALDRICH	A2383; CAS:34369-07-8
Potassium D- gluconate	SIGMA-ALDRICH	G4500; CAS:299-27-4
Sulforhodamine B	SIGMA-ALDRICH	230162; CAS:3530-42-1
Tubocurarine chloride pentahydrate	SIGMA-ALDRICH	93750; CAS:6989-98-6
EGTA	SIGMA-ALDRICH	03777; CAS:67-42-5
Agarose, low gelling temperature	SIGMA-ALDRICH	A9414; CAS:39346-81-1
Ethyl 3-aminobenzoate methanesulfonate (MS-222)	SIGMA-ALDRICH	E10521; CAS:888-86-2
Magnesium chloride	Fisher Scientific	M33500; CAS:7791-18-6
Glucose	Fisher Scientific	D16500; CAS:50-99-7
HEPES	HIMEDIA	RM380; CAS:7365-45-9
Sodium chloride	HIMEDIA	GRM853; CAS:7647-14-5
Potassium chloride	Qualigens	Product#13305
Software and Algorithms		
Matlab	Mathworks, Natick, USA	RRID:SCR_001622
Processing	[56]	https://processing.org
Bonsai	[57]	http://www.open-ephys.org/bonsai
Ipython	[58]	https://ipython.org; RRID:SCR_001815
Fiji	[59]	http://fiji.sc; RRID:SCR_002285
nlme package in R	[60]	https://CRAN.R-project.org/package=nlme.

References

1.   McLean, D.L., Fan, J., Higashijima, S., Hale, M.E., and Fetcho, J.R. (2007). A

topographic map of recruitment in spinal cord. Nature *446*, 71. Available at:

https://doi.org/10.1038/nature05588.

2.   McLean, D.L., Masino, M.A., Koh, I.Y.Y., Lindquist, W.B., and Fetcho, J.R.

(2008). Continuous shifts in the active set of spinal interneurons during

changes in locomotor speed. Nat. Neurosci. *11*, 1419. Available at:

https://doi.org/10.1038/nn.2225.

3.   Bellardita, C., and Kiehn, O. (2015). Phenotypic Characterization of Speed-

Associated Gait Changes in Mice Reveals Modular Organization of Locomotor

Networks. Curr. Biol. *25*, 1426–1436. Available at:

https://doi.org/10.1016/j.cub.2015.04.005.

4.   Gosgnach, S., Lanuza, G.M., Butt, S.J.B., Saueressig, H., Zhang, Y.,

Velasquez, T., Riethmacher, D., Callaway, E.M., Kiehn, O., and Goulding, M.

(2006). V1 spinal neurons regulate the speed of vertebrate locomotor outputs.

Nature *440*, 215–219. Available at: https://doi.org/10.1038/nature04545.

5.   Zhong, G., Sharma, K., and Harris-Warrick, R.M. (2011). Frequency-

dependent recruitment of V2a interneurons during fictive locomotion in the

mouse spinal cord. Nat. Commun. *2*, 274. Available at:

https://doi.org/10.1038/ncomms1276.

6.   Gabriel, J.P., Ausborn, J., Ampatzis, K., Mahmood, R., Eklöf-Ljunggren, E.,

and El Manira, A. (2010). Principles governing recruitment of motoneurons

during swimming in zebrafish. Nat. Neurosci. *14*, 93. Available at:

https://doi.org/10.1038/nn.2704.

7.   Ampatzis, K., Song, J., Ausborn, J., and El Manira, A. (2013). Pattern of

Innervation and Recruitment of Different Classes of Motoneurons in Adult

Zebrafish. J. Neurosci. *33*, 10875 LP – 10886. Available at:

http://www.jneurosci.org/content/33/26/10875.abstract.

8. Ausborn, J., Mahmood, R., and El Manira, A. (2012). Decoding the rules of

recruitment of excitatory interneurons in the adult zebrafish locomotor network.

Proc. Natl. Acad. Sci. *109*, E3631 LP-E3639. Available at:

http://www.pnas.org/content/109/52/E3631.abstract.

9. Ampatzis, K., Song, J., Ausborn, J., and El Manira, A. (2014). Separate

Microcircuit Modules of Distinct V2a Interneurons and Motoneurons Control

the Speed of Locomotion. Neuron *83*, 934–943. Available at:

https://www.sciencedirect.com/science/article/pii/S0896627314006291.

10. Song, J., Dahlberg, E., and El Manira, A. (2018). V2a interneuron diversity

tailors spinal circuit organization to control the vigor of locomotor movements.

Nat. Commun. *9*, 3370. Available at: https://doi.org/10.1038/s41467-018-

05827-9.

11. Berg, E.M., Björnfors, E.R., Pallucchi, I., Picton, L.D., and El Manira, A. (2018).

Principles Governing Locomotion in Vertebrates: Lessons From Zebrafish   .

Front. Neural Circuits   *12*, 73. Available at:

https://www.frontiersin.org/article/10.3389/fncir.2018.00073.

12. Kiehn, O. (2016). Decoding the organization of spinal circuits that control

locomotion. Nat. Rev. Neurosci. *17*, 224–238. Available at:

https://www.ncbi.nlm.nih.gov/pubmed/26935168.

13. Arber, S. (2017). Organization and function of neuronal circuits controlling

movement. EMBO Mol. Med. *9*, 281–284. Available at:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5331227/.

14. Marder, E., and Thirumalai, V. (2002). Cellular, synaptic and network effects of

neuromodulation. Neural Networks *15*, 479–493. Available at:

http://www.sciencedirect.com/science/article/pii/S0893608002000436.

15.   Marder, E. (2012). Neuromodulation of Neuronal Circuits: Back to the Future.

Neuron *76*, 1–11. Available at:

https://www.sciencedirect.com/science/article/pii/S0896627312008173.

16.   Harris-Warrick, R.M. (2011). Neuromodulation and flexibility in Central Pattern

Generator networks. Curr. Opin. Neurobiol. *21*, 685–692. Available at:

https://www.ncbi.nlm.nih.gov/pubmed/21646013.

17.   Miles, G.B., and Sillar, K.T. (2011). Neuromodulation of Vertebrate Locomotor

Control Networks. Physiology *26*, 393–411. Available at:

https://doi.org/10.1152/physiol.00013.2011.

18.   Dickinson, P.S., Mecsas, C., and Marder, E. (1990). Neuropeptide fusion of

two motor-pattern generator circuits. Nature *344*, 155–158. Available at:

https://doi.org/10.1038/344155a0.

19.   Lieske, S.P., Thoby-Brisson, M., Telgkamp, P., and Ramirez, J.M. (2000).

Reconfiguration of the neural network controlling multiple breathing patterns:

eupnea, sighs and gasps. Nat. Neurosci. *3*, 600–607. Available at:

https://doi.org/10.1038/75776.

20.   Sharples, S.A., Koblinger, K., Humphreys, J.M., and Whelan, P.J. (2014).

Dopamine: a parallel pathway for the modulation of spinal locomotor networks.

Front. Neural Circuits *8*, 55. Available at:

https://www.ncbi.nlm.nih.gov/pubmed/24982614.

21.   Kiehn, O., and Kjaerulff, O. (1996). Spatiotemporal characteristics of 5-HT and

dopamine-induced rhythmic hindlimb activity in the in vitro neonatal rat. J.

Neurophysiol. *75*, 1472–82. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/8727391.

22.   Barriere, G., Mellen, N., Cazalets, J.-R.J.-R., Barrière, G., Mellen, N., and

Cazalets, J.-R.J.-R. (2004). Neuromodulation of the locomotor network by

dopamine in the isolated spinal cord of newborn rat. Eur. J. Neurosci. *19,*

1325–1335. Available at: https://doi.org/10.1111/j.1460-9568.2004.03210.x.

23.   Thirumalai, V., and Cline, H.T. (2008). Endogenous Dopamine Suppresses

Initiation of Swimming in Prefeeding Zebrafish Larvae. J. Neurophysiol. *100,*

1635–1648. Available at:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2544474/.

24.   Lambert, A.M., Bonkowsky, J.L., and Masino, M.A. (2012). The conserved

dopaminergic diencephalospinal tract mediates vertebrate locomotor

development in zebrafish larvae. J. Neurosci. *32*, 13488–13500. Available at:

https://www.ncbi.nlm.nih.gov/pubmed/23015438.

25.   Puhl, J.G., and Mesce, K.A. (2008). Dopamine Activates the Motor Pattern for

Crawling in the Medicinal Leech. J. Neurosci. *28*, 4192 LP – 4200. Available

at: http://www.ncbi.nlm.nih.gov/pubmed/18417698.

26.   Whelan, P., Bonnot, A., and O'Donovan, M.J. (2000). Properties of Rhythmic

Activity Generated by the Isolated Spinal Cord of the Neonatal Mouse. J.

Neurophysiol. *84*, 2821–2833. Available at:

https://doi.org/10.1152/jn.2000.84.6.2821.

27.   Tay, T.L., Ronneberger, O., Ryu, S., Nitschke, R., and Driever, W. (2011).

Comprehensive catecholaminergic projectome analysis reveals single-neuron

integration of zebrafish ascending and descending dopaminergic systems. Nat.

Commun. *2*, 171. Available at: http://dx.doi.org/10.1038/ncomms1171.

28.   Portugues, R., and Engert, F. (2009). The neural basis of visual behaviors in

the larval zebrafish. Curr. Opin. Neurobiol. *19*, 644–647. Available at:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4524571/.

29.   Severi, K.E., Portugues, R., Marques, J.C., O'Malley, D.M., Orger, M.B., and

Engert, F. (2014). Neural control and modulation of swimming speed in the

larval zebrafish. Neuron *83*, 692–707. Available at:

https://www.ncbi.nlm.nih.gov/pubmed/25066084.

30.   Buss, R.R., and Drapeau, P. (2001). Synaptic Drive to Motoneurons During

Fictive Swimming in the Developing Zebrafish. J. Neurophysiol. *86*, 197–210.

Available at: https://doi.org/10.1152/jn.2001.86.1.197.

31.   Irons, T.D., Kelly, P.E., Hunter, D.L., MacPhail, R.C., and Padilla, S. (2013).

Acute administration of dopaminergic drugs has differential effects on

locomotion in larval zebrafish. Pharmacol. Biochem. Behav. *103*, 792–813.

Available at:

http://www.sciencedirect.com/science/article/pii/S0091305712003346.

32.   Souza, B.R., Romano-Silva, M.A., and Tropepe, V. (2011). Dopamine D2

Receptor Activity Modulates Akt Signaling and Alters GABAergic Neuron

Development and Motor Behavior in Zebrafish Larvae. J. Neurosci. *31*, 5512–

5525. Available at: http://www.jneurosci.org/content/31/14/5512.long.

33.   Jay, M., De Faveri, F., and McDearmid, J.R. (2015). Firing Dynamics and

Modulatory Actions of Supraspinal Dopaminergic Neurons during Zebrafish

Locomotor Behavior. Curr. Biol. *25*, 435–444. Available at:

https://doi.org/10.1016/j.cub.2014.12.033.

34.   Green, M.H., Ho, R.K., and Hale, M.E. (2011). Movement and function of the

pectoral fins of the larval zebrafish (Danio rerio) during slow swimming. J. Exp.

Biol. *214*, 3111 LP – 3123. Available at:

http://jeb.biologists.org/content/214/18/3111.abstract.

35.   Bainbridge, R. (1958). The Speed of Swimming of Fish as Related to Size and

to the Frequency and Amplitude of the Tail Beat. J. Exp. Biol. *35*, 109–133.

Available at: http://jeb.biologists.org/content/jexbio/35/1/109.full.pdf.

36.   Myers, P.Z. (1985). Spinal motoneurons of the larval zebrafish. J. Comp.

Neurol. *236*, 555–561. Available at:

http://doi.wiley.com/10.1002/cne.902360411.

37.   Liu, D.W., and Westerfield, M. (1988). Function of identified motoneurones and

co-ordination of primary and secondary motor systems during zebra fish

swimming. J. Physiol. *403*, 73–89. Available at:

https://doi.org/10.1113/jphysiol.1988.sp017239.

38.   Menelaou, E., and McLean, D.L. (2012). A Gradient in Endogenous

Rhythmicity and Oscillatory Drive Matches Recruitment Order in an Axial

Motor Pool. J. Neurosci. *32*, 10925 LP – 10939. Available at:

http://www.jneurosci.org/content/32/32/10925.abstract.

39.   McLean, D.L., and Fetcho, J.R. (2004). Relationship of tyrosine hydroxylase

and serotonin immunoreactivity to sensorimotor circuitry in larval zebrafish. J.

Comp. Neurol. *480*, 57–71. Available at: https://doi.org/10.1002/cne.20281.

40.   Harris-Warrick, R.M., Coniglio, L.M., Levini, R.M., Gueron, S., and

Guckenheimer, J. (1995). Dopamine modulation of two subthreshold currents

produces phase shifts in activity of an identified motoneuron. J. Neurophysiol.

74, 1404–1420. Available at: https://doi.org/10.1152/jn.1995.74.4.1404.

41.   Han, P., Nakanishi, S.T., Tran, M.A., and Whelan, P.J. (2007). Dopaminergic

Modulation of Spinal Neuronal Excitability. J. Neurosci. *27*, 13192–13204.

Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.1279-

07.2007.

42.   Kispersky, T.J., Caplan, J.S., and Marder, E. (2012). Increase in Sodium

Conductance Decreases Firing Rate and Gain in Model Neurons. J. Neurosci.

32, 10995 LP – 11004. Available at:

http://www.jneurosci.org/content/32/32/10995.abstract.

43.   Kishore, S., Bagnall, M.W., and McLean, D.L. (2014). Systematic Shifts in the

Balance of Excitation and Inhibition Coordinate the Activity of Axial Motor

Pools at Different Speeds of Locomotion. J. Neurosci. *34*, 14046 LP – 14054.

Available at: http://www.jneurosci.org/content/34/42/14046.abstract.

44.   Talpalar, A.E., Bouvier, J., Borgius, L., Fortin, G., Pierani, A., and Kiehn, O.

(2013). Dual-mode operation of neuronal networks involved in left–right

alternation. Nature *500*, 85–88. Available at:

http://www.nature.com/articles/nature12286 [Accessed June 11, 2019].

45.   Kimura, Y., and Higashijima, S. (2019). Regulation of locomotor speed and

selection of active sets of neurons by V1 neurons. Nat. Commun. *10*, 2268.

Available at: https://doi.org/10.1038/s41467-019-09871-x.

46.   Knafo, S., Fidelin, K., Prendergast, A., Tseng, P.-E.B., Parrin, A., Dickey, C.,

Böhm, U.L., Figueiredo, S.N., Thouvenin, O., Pascal-Moussellard, H., *et al.*

(2017). Mechanosensory neurons control the timing of spinal microcircuit

selection during locomotion. Elife *6*, e25260. Available at:

https://www.ncbi.nlm.nih.gov/pubmed/28623664.

47.   Naumann, E.A., Fitzgerald, J.E., Dunn, T.W., Rihel, J., Sompolinsky, H., and

Engert, F. (2016). From Whole-Brain Data to Functional Circuit Models: The

Zebrafish Optomotor Response. Cell *167*, 947-960.e20. Available at:

http://www.sciencedirect.com/science/article/pii/S0092867416314027.

48.   Kramer, A., Wu, Y., Baier, H., and Kubo, F. (2019). Neuronal Architecture of a

851   Visual Center that Processes Optic Flow. Neuron. Available at:

852   http://www.sciencedirect.com/science/article/pii/S0896627319303757.

853   49.   Orger, M.B., Kampff, A.R., Severi, K.E., Bollmann, J.H., and Engert, F. (2008).

854   Control of visually guided behavior by distinct populations of spinal projection

855   neurons. Nat. Neurosci. *11*, 327. Available at: https://doi.org/10.1038/nn2048.

856   50.   Severi, K.E., Böhm, U.L., and Wyart, C. (2018). Investigation of hindbrain

857   activity during active locomotion reveals inhibitory neurons involved in

858   sensorimotor processing. Sci. Rep. *8*, 13615. Available at:

859   https://doi.org/10.1038/s41598-018-31968-4.

860   51.   Huang, K.-H., Ahrens, M.B., Dunn, T.W., and Engert, F. (2013). Spinal

861   Projection Neurons Control Turning Behaviors in Zebrafish. Curr. Biol. *23*,

862   1566–1573. Available at: https://doi.org/10.1016/j.cub.2013.06.044.

863   52.   Wang, W.-C., and McLean, D.L. (2014). Selective responses to tonic

864   descending commands by temporal summation in a spinal motor pool. Neuron

865   *83*, 708–721. Available at: https://www.ncbi.nlm.nih.gov/pubmed/25066087.

866   53.   McLean, D.L., and Fetcho, J.R. (2004). Ontogeny and innervation patterns of

867   dopaminergic, noradrenergic, and serotonergic neurons in larval zebrafish. J.

868   Comp. Neurol. *480*, 38–56. Available at:

869   https://www.ncbi.nlm.nih.gov/pubmed/15515022.

870   54.   Reinig, S., Driever, W., and Arrenberg, A.B. (2017). The Descending

871   Diencephalic Dopamine System Is Tuned to Sensory Stimuli. Curr. Biol. *27*,

872   318–333. Available at:

873   https://www.sciencedirect.com/science/article/pii/S0960982216314464?via%3

874   Dihub.

55.   Ryczko, D., Grätsch, S., Schläger, L., Keuyalian, A., Boukhatem, Z., Garcia,

C., Auclair, F., Büschges, A., and Dubuc, R. (2017). Nigral Glutamatergic

Neurons Control the Speed of Locomotion. J. Neurosci. *37*, 9759 LP – 9770.

Available at: http://www.ncbi.nlm.nih.gov/pubmed/28924005.

56.   Reas, C., and Fry, B. (2006). Processing: programming for the media arts. AI

Soc. *20*, 526–538. Available at: https://doi.org/10.1007/s00146-006-0050-9.

57.   Lopes, G., Bonacchi, N., Frazão, J., Neto, J.P., Atallah, B. V, Soares, S.,

Moreira, L., Matias, S., Itskov, P.M., Correia, P.A., *et al.* (2015). Bonsai: an

event-based framework for processing and controlling data streams. Front.

Neuroinform. *9*, 7. Available at:

https://www.frontiersin.org/article/10.3389/fninf.2015.00007.

58.   Pérez, F., and Granger, B.E. (2007). IPython: A System for Interactive

Scientific Computing. Comput. Sci. Eng. *9*, 21–29. Available at:

https://aip.scitation.org/doi/abs/10.1109/MCSE.2007.53.

59.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M.,

Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., *et al.* (2012).

Fiji: An open-source platform for biological-image analysis. Nat. Methods *9*,

676–682. Available at: http://www.nature.com/articles/nmeth.2019.

60.   Pinheiro J, Bates D, DebRoy S, Sarkar D, R.C.T. (2019). nlme: Linear and

Nonlinear Mixed Effects Models. Available at: https://cran.r-

project.org/package=nlme.

A

B

duration

Amplitude

Tail tip position (mm)

Time (s)

C

D
Amplitude (mm)
***

E
Tail beat frequency (Hz)
*

F
Bout duration (ms)
ns

G
Bout number
ns

H

I
Amplitude (mm)
*

J
Tail beat frequency (Hz)
ns

K
Bout duration (ms)
ns

L
Bout number
ns

• Control   • Agonist   • Antagonist

A

B

Max. tail bend angle

C Control · D1-like-R agonist

D Control · D1-like-R antagonist

E **

F *

- Control · Agonist · Antagonist

A

Control

D1-like-R agonist

Primary motor neuron

10 mV

100 ms

275 pA

B

F$_{inst.}$

Threshold

Latency

C *

First spike latency (ms)

D *

Spike latency (ms)

Current injected (pA)

E ***

AP threshold (mV)

F *

Rheobase (pA)

G *

R$_{in}$(MΩ)

H *

Firing frequency (Hz)

Current injected (pA)

— Control — D1-like-R agonist