Behavioral training of marmosets and electrophysiological recording from the cerebellum

Ehsan Sedaghat-Nejad[1], David J. Herzfeld[1,3] *, Paul Hage[1] *,
Kaveh Karbasi[1], Tara Palin[1], Xiaoqin Wang[2], and Reza Shadmehr[1]

[1]Laboratory for Computational Motor Control
[2]Laboratory for Auditory Neurophysiology
Department of Biomedical Engineering
Johns Hopkins School of Medicine

[3] current address: Dept. of Neurobiology, Duke University School of Medicine, Durham, NC.
* Indicates equal contribution

Correspondence: Ehsan Sedaghat-Nejad or Reza Shadmehr, Traylor Building Room 416, Johns Hopkins School of Medicine, 720 Rutland Ave., Baltimore, MD 21205. Email: e.sedaghatnejad@gmail.com or shadmehr@jhu.edu. Phone: 410-614-3424.

Acknowledgements: The work was supported by grants from the NIH (5-R01-NS078311), the Office of Naval Research (N00014-15-1-2312), and the National Science Foundation (CNS-1714623). We are grateful to Tahl Holtzman from Cambridge Neurotech, who made important suggestions regarding choice of microdrive, electrodes, and silicone gel for protection of the dura. ESN collected some of the behavioral data on subject B and all of the behavioral and neurophysiological data on subject M. He performed surgery on subjects M and R, analyzed the behavioral and neurophysiological data, and made figures. DJH designed the survival surgery, wrote the software for behavioral training, and collected behavioral data from subject B. He performed surgery on subject B. PH designed the image guided procedures for construction of the head-post, designed the alignment ruler, and made figures. KK wrote some of the software, collected some of the behavioral data on subject M, and analyzed the neurophysiological data. TP performed image co-registration and made figures. XW provided the animals as well as technical help with surgery and feedback for electrophysiology designs. RS collected some of the neurophysiological data and wrote the manuscript. We are very grateful to Seth Koehler, Xiao-Ping Liu, and Wang Lab technicians in helping with surgeries and animal care.

Abstract
The common marmoset (*Callithrix Jacchus*) is a promising new model for study of neurophysiological
basis of behavior in primates. Like other primates, it relies on saccadic eye movements to monitor and
explore its environment. Previous reports have demonstrated some success in training marmosets to
produce goal-directed actions in the laboratory. However, the number of trials per session has been
relatively small, thus limiting the utility of marmosets as a model for behavioral and neurophysiological
studies. Here, we report the results of a series of new behavioral training and neurophysiological
protocols aimed at increasing the number of trials per session while recording from the cerebellum. To
improve the training efficacy, we designed a precisely calibrated food regulation regime that motivated
the subjects to perform saccade tasks, resulting in about a thousand reward-driven trials on a daily
basis. We then developed a multi-channel recording system that used imaging to target a desired region
of the cerebellum, allowing for simultaneous isolation of multiple Purkinje cells in the vermis. In this
report, we describe (1) the design and surgical implantation of a CT guided, subject specific head-post,
(2) the design of a CT and MRI guided alignment tool for trajectory guidance of electrodes mounted on
an absolute encoder microdrive, (3) development of a protocol for behavioral training of subjects, and
(4) simultaneous recordings from pairs of Purkinje cells during a saccade task.
17

New and Noteworthy
Marmosets present the opportunity to investigate genetically based neurological disease in primates; in
particular, diseases that affect social behaviors, sound production, and eye movements. All of these
behaviors depend on the integrity of the cerebellum. Here, we present training methods that motivate
the subjects, allowing for improved performance during recordings. We also present
electrophysiological techniques that precisely target the subject's cerebellum, allowing for simultaneous
isolation of multiple Purkinje cells.

Introduction

The common marmoset (*Callithrix jacchus*) has gained considerable attention for neurophysiological investigation in recent years because of its potential for transgenic manipulation (Miller et al., 2016). Marmosets are small, New World primates with no known lethal zoonotic diseases that are transmittable to humans (Wakabayashi et al., 2018). Behaviorally, they share important attributes with humans: they are social and live in family units, they are vocal and rely on species-specific sound production for their communication, and they are visual and use saccadic eye movements to explore their environment. Notably, for a primate they have a particularly short gestation period (~ 5 months) and regularly give birth to twins or triplets. Thus, they have a high breeding efficiency with potential for germline transmission of genetically modified models.

Indeed, marmosets research is benefiting from transgenic (Sasaki et al., 2009), gene-editing (Kishi et al., 2014), and optogenetic tools that target specific neurons (MacDougall et al., 2016). Thus, marmosets have the potential to become a model system for investigating cognitive and social behaviors (Miller et al., 2016;Takahashi et al., 2017;Mustoe et al., 2015), auditory (Wang, 2018) and visual perception (Solomon and Rosa, 2014), as well as neural control of vocalization (Roy et al., 2011;Eliades and Wang, 2013;Eliades and Miller, 2017) and eye movements (Mitchell et al., 2014;Johnston et al., 2018).

Many aspects of social behavior, vocalization, and eye movements depend on the integrity of the cerebellum. For example, children who suffer from autism spectrum disorder (ASD), a developmental disorder that leads to impairments in social and communication skills, exhibit anatomic abnormalities in their cerebellum, including reduced number of Purkinje cells (Whitney et al., 2008), particularly along the vermis (Murakami et al., 1989;Hashimoto et al., 1995;Courchesne et al., 2001;Scott et al., 2009). Notably, in children with ASD, we found that damage to the cerebellum is prevalent in lobule VI and parts of lobule VIII (Marko et al., 2015), regions that in the macaque are critical for control of saccadic eye movements (Takagi et al., 1998;Barash et al., 1999). Thus, marmosets may present a particularly good opportunity to investigate the role of the cerebellum in eye movements, social communication, and neurological diseases such as ASD.

However, despite these attractive features, there is a critical concern: can marmosets be employed for studying the neural basis of goal-direct behavior in a laboratory setting? This concern is due to the fact that in the current literature, the number of trials that these animals can perform within a session appears to be quite limited. For example, one recent study that trained marmosets to perform saccades found that on average, they performed 80-100 trials per session (Johnston et al., 2018), while another study reported 300-800 trials per session (Mitchell et al., 2014). Some of these numbers are on the lower bound of what is needed to identify task-related neurons, and may be insufficient to measure neurophysiological changes during learning or decision-making. As a result, it is unclear whether marmosets can serve as a model for neurophysiological studies of operant conditioned behavior.

Over the past four years, our team has been building a new marmoset laboratory, aiming to develop behavioral and neurophysiological protocols that allow for electrophysiological recording from the cerebellum of head-fixed animals. From a behavioral perspective, our goal was to ask whether the subjects could be motivated to produce about a thousand rewarded trials on a daily basis in the head-fixed configuration. From an electrophysiological perspective, our aim was to produce precise targeting

of a region of the cerebellum, thus providing the means to simultaneously isolate multiple Purkinje cells
using high-density electrodes.
To solve the training problem, we designed a carefully calibrated food regulation regime that
motivated the subjects to perform a saccade task. To solve the electrophysiological problem, we
developed a high-precision, acute multi-channel recording system that resulted in simultaneous
recording from multiple Purkinje cells within the cerebellar vermis.
In this report, we provide an account of our experience, both in terms of procedures that failed
and those that succeeded, going from initial behavioral training through neurophysiological recording.
We describe (1) the design and surgical implantation of a CT guided subject-specific head-post, (2)
development of a CT and MRI guided alignment mechanism for trajectory guidance of electrodes
mounted on an absolute encoder microdrive, (3) design of a protocol for training of subjects in a reward
driven saccade task, and (4) isolation of Purkinje cells via tetrodes, heptodes, and high-density silicon
probes during the task.
14
Methods
Subjects
The surgical procedures were carried out on three marmosets: two female (subject B, 420g, 11 years
old; subject M, 350g, 4 years old) and one male (subject R, 420g, 4 years old). The behavioral procedures
were tested on subjects B and M. All three subjects were born and raised in a breeding colony that one
of the authors (XW) has maintained at the Johns Hopkins School of Medicine since 1996. The
experimental procedures were evaluated and approved by Johns Hopkins University Animal Care and
Use Committee in compliance with the guidelines of the United States National Institutes of Health.
23
Design of head-post and chamber
Recording from Purkinje cells of the cerebellum imposes particular demands on stability: one needs to
not only isolate a neuron, but also establish identity of that neuron via presence of both simple and
complex spikes, which requires movable electrodes and longer periods of recordings. In order to
stabilize the head during behavioral sessions, the current technique is to build an acrylic based thick
"helmet" atop the skull, which then serves as a base that holds two or more metal posts (Lu et al.,
2001a). The glue-inserted metal posts are attached to bars that are fixed to a primate chair, thereby
holding the head in a fixed position. A more recent method is a halo-like device that holds the skull
inside a ring (Johnston et al., 2018). Both approaches have been used for recordings from the cerebral
cortex. However, reaching the cerebellum requires open access to the posterior parts of the skull,
making the halo approach less desirable. Furthermore, we aimed to produce an approach that reduced
disruption of the underlying skin and muscles, resulting in a design that allowed for skin coverage of the
skull and integrity of the temporalis muscles. This led us to develop a titanium based head-post and
chamber design that eliminated use of the acrylic based helmet, and maximized the open area in the
posterior part of the skull (Fig. 1).
To prepare for head-fixation, we designed a head-post based on the specific geometry of the
surface of each subject's skull and manufactured it using 3D printed titanium. We began with a
preoperative CT (Fig. 1A), and used it to build a 3D model of the skull (Fig. 1B). This was done using 3D
Slicer, an open-source image analysis and visualization software (Fedorov et al., 2012). The 3D skull

model was then imported into a CAD environment (SolidWorks and Autodesk Fusion 360), which informed the design of an X-shaped titanium head-post (Fig. 1B and Fig. 1C), a titanium base recording chamber (Fig. 1D), and a plastic protective chamber cap. The head-post and base chamber designs were optimized to remain light weight and within the anatomical constraints of the frontal eye ridge and the lateral ridges on the skull, allowing us to minimize damage to the temporalis muscle attachment. The head-post and base chamber were 3D printed with laser melted, grade 2 titanium (6AI-4V), producing a lightweight (3g), biocompatible, corrosion resistant structure. The protective cap that covered the chamber was 3D printed with PLA plastic filament (0.5g).

Surgery

Implanting the head-post and chamber was conducted by a surgical team led by lab members, and aided by veterinary technicians and veterinarians. The animal was anesthetized with Alfaxalone (12mg/Kg), then intubated with #2.0 uncuffed endotracheal (ET) tube, and administered isoflurane (1-1.25%, adjusted as needed) to maintain the anesthetic plane, as well as atropine and dexamethasone via intramuscular (IM) injections. The subject was then head-fixed by inserting bilateral ear bars into the osseous ear canal, which were fastened to a stereotaxic frame, while a bite bar was slid into place with a forehead clamp attachment to ensure that the head was positioned appropriately and securely with respect to the chest. 2% Lidocaine and 1:100000 epinephrine was then injected into the subcutaneous space beneath the scalp, and the operative region was treated with alternating cycles of diluted chlorhexidine and alcohol and at the very end sprayed with betadine. Next, using a surgical marker, we drew an outline for a midline incision on the scalp, extending 2mm posterior to the supraorbital fat pad and extending to the occipital ridge. We then made the incision using a scalpel and the skin was separated from the underlying fascia as well as the fascia from the underlying muscle. From here, the surface of the skull was prepared by removing residual tissue using curettes and 3% hydrogen peroxide solution, while the head-post was positioned for implantation. The temporalis muscle was pushed backed from its bony insertions on the spots that intersected with the head-post.

To attach the head-post, phosphoric acid etchant was applied for 15-20 seconds and then rinsed. Next, a thin layer of Optibond Solo Plus (Kerr Corp.) was applied to the surface of the skull, covering the entire area to be occupied by the head-post, cranial ground screw, and base chamber. After application, the Optibond was cured using a UV gun. The screw locations were then marked and drilled using a 2.5mm diameter drill bit (#54), with a stopper limiting the hole depth to 1.1mm. NX3 dual-cure dental cement (Kerr Corp.) was applied to the center of the bottom of the head-post, placed on the skull, and then cured using UV light.

Titanium screws (2.4x4mm, 0.04g, or 2.4x5mm, 0.05g) were then used to affix the legs of the head-post onto the skull. A cranial ground screw (titanium, 2.4x8mm, 0.08g) was screwed into the prepared hole (Fig. 1B). We confirmed that screw depth into the skull did not exceed 1.1mm. Next, NX3 cement was applied around screws inside head-post screw holes, and around the ground screw to ensure structural integrity of the implanted screws.

NX3 cement was also applied to the bottom edge of the base chamber and the base chamber was placed between the rear legs of the head-post (Fig. 1B) so that it was approximately located above the visual cortex and directed straight down. NX3 cement underwent a UV light curing. A final layer of

NX3 cement was applied over the whole structure and was smoothed using Parafilm tape, and then get cured with UV light.

The total weight of the NX3 cement, the installed apparatus, and the screws was less than 8.5g. With the implantation complete, the temporalis muscle and fascia were replaced to their original positions. Finally, the two skin flaps were re-attached together via sutures using a 0/4 single filament thread. As a result, the surgical procedure produced an animal in which the skin covered almost the entire skull, except for small regions dedicated to the frontal ground screw, the main bar of the head-post, and the base chamber.

At this point anesthesia was discontinued and the ET tube was removed. The animal was monitored as consciousness and motor function were regained. A second dose of Dexamethasone (0.25mg/Kg) was administered via IM injection 12 hours after the first dose. Recovery was monitored with daily postoperative check-ups for 13 days, including inspection of the surgery site for infection, checking for integrity of the sutures, recording of vital signs, feeding, and the administering of drugs.

Establishing a chamber-based coordinate system for the cerebellum

Because of the small size of the skull and our chamber design, we found it difficult if not impossible to attach a micromanipulator to the head-post. Given that the micromanipulator was external to the head-post, we were concerned that we may not be able to accurately target and reproducibly approach a desired location in the cerebellum. This led us to design an electrode guide system that we integrated into the chamber design (Fig. 2). Our problem was that during surgery, the base chamber was placed by hand on the skull, and therefore its location with respect to the cerebellum was unknown. Furthermore, post-surgical CT or MRI imaging of the implanted animal was possible but problematic because the titanium pieces generated substantial artifacts, making it difficult to precisely localize the base chamber.

To solve this problem, we designed a CT visible reference ruler (Fig. 2) that we inserted into the base chamber. The reference ruler consisted of 7 layers of a 1.5mm marked grid system, 3D printed with PLA plastic filament (Fig. 2B). With the reference ruler inserted into the chamber, we performed a post-operative CT scan (Fig. 2C). This produced a clear image of the ruler, but with artifacts from the titanium. We then co-registered pre-operative CT (Fig. 1A) and post-operative CT (Fig. 2C) using bony areas and co-registered pre-operative CT (Fig. 1A) and the pre-operative MRI (Fig. 2A). Fixing pre-operative CT between 2 registrations produced a co-registered image of all three images. The result was a full representation of the subject's head, including the reference ruler, the chamber, and the brain (Fig. 2D).

We used the ruler, chamber, and skull geometry to plan electrode trajectories in order to arrive at 20 distinct points of interest in the cerebellum. In this design, all 20 trajectories traveled through a single 1.5mm diameter burr hole. To calculate each trajectory, we placed 21 points at various positions on the reference ruler (red marks, Fig. 2D), 1 point on the surface of the skull for a craniotomy burr hole (burr hole point, Fig. 2D), and 20 points at various locations along lobules V, VI, and VII of the vermis (cerebellar marks, Fig. 2D). The point representing the burr hole was placed at 1.1mm lateral to the midline in order to avoid the superior sagittal sinus.

The various points were then imported into a CAD environment, along with a model of the brain that we generated with 3DSlicer from the MRI images (Fig. 2E). We aligned the imported points to their respective targets, guided by the co-registered image. The process relied on the alignment of all reference axis ruler points from the co-registered image (Fig. 2D) to the reference ruler within the model

environment (Fig. 2E). This automatically aligned the burr hole and points on the cerebellum to the skull and brain models, creating an accurate representation of the distances and geometry of the desired cerebellar points, the desired burr hole, and base chamber.

Using the coordinates of the various points we created a set of 20 trajectories that originated from the desired recording locations in the cerebellum, converged through the single burr hole, and traveled out past the base chamber (Fig. 2F). Each line in 3D space represented a trajectory that an electrode would travel to reach a given cerebellar destination.

Design of an electrode guidance tool

Based on the electrode trajectories that we had defined in the post-op model (Fig. 2F), we designed a tool that would be attached to the base chamber and provide alignment of the electrode to the specific trajectory that traveled through the burr hole and arrived at the desired cerebellar point of interest (Fig. 3A). This alignment tool featured 20 cylindrical tracks, each a cylinder of 1/32" in diameter. All cylinders converged onto the single burr hole, from which point the trajectories diverged to arrive at the various cerebellar destinations. The 20 cylinders (guide cylinders, Fig. 3B) were configured in a 4X5 grid which corresponded to the four targeted cerebellar lobules (VIIa, VIa-c, VIe, and Vd), spanning from 0 to 2.5mm along the midline-lateral axis. The cylinders in the alignment tool provided a physical representation of the electrode trajectories above the burr hole. The tool was 3D printed using an Objet260 printer with VeroClear plastic filament.

Alignment of the electrode

To advance the electrode, we chose a piezoelectric, high precision microdrive (0.5 micron resolution) with an integrated absolute encoder (M3-LA-3.4-15 Linear smart stage, New Scale Technologies). The microdrive held the electrode and advanced it along a single axis defined by the chosen cylinder of the electrode alignment tool. Thus, the next problem was to position the microdrive in 3D space so that its single direction of motion was precisely aligned with the desired cylinder in the electrode alignment tool.

To align the axis of the microdrive with the desired cylinder, we began by attaching the microdrive to a mechanical stereotaxic micromanipulator (SM-11-HT, Narishige, Japan). With the alignment tool installed in the base chamber, we inserted a 1/32" rod (outer diameter) inside the desired cylinder of the alignment tool (Fig. 4A), and then attached a 16G tube (0.033 inch inner diameter) to the microdrive using the same electrode holder that would also hold the electrode. The tube was fixed to the microdrive, serving as a model for the axis of travel of the electrode.

We then used the stereotaxic micromanipulator to maneuver the microdrive, thereby positioning the tube held by the electrode holder so that it precisely traveled along the axis defined by the rod. Alignment of the tube (representing the electrode) with the rod (representing the guide) was confirmed under a microscope by advancing the microdrive and demonstrating that this motion inserted the rod into the tube. Once this alignment was confirmed, the stereotactic manipulator that held the microdrive was locked into place. We then removed the tube from the microdrive and replaced it with the recording electrode (Fig. 4B). All further motion of the electrode was now under the control of the microdrive, limited to a single axis of travel that was along the desired trajectory.

Craniotomy

Our design required that the 20 trajectories specified by the alignment tool converge on a single point on the skull and then travel within the brain to reach the cerebellum. The next step was to localize this point on the skull and drill a burr hole. This was achieved by using the alignment tool to mark the skull and localize the burr hole.

The subject was head-fixed and the general craniotomy area, the base chamber, and the alignment tool were sterilized using alcohol, rinsed with saline, and then suctioned dry. The alignment tool was installed on the base chamber and a 22G blunt needle tip was dipped into ink and then inserted into one of the cylinders (the choice was arbitrary as all cylinders converged to the same burr hole location). The ink tipped needle was advanced until contact was made with the bone and a mark was left on the surface.

Using the procedures for aligning an electrode, we aligned a sterilized drill bit so that its trajectory of travel precisely matched the trajectory specified by a cylinder on the alignment tool. We attached a DC motor to the bit, and then attached the motor to the stereotaxic micromanipulator. At this point the alignment tool was removed, exposing the marked skull. From our co-registered pre-op and post-op CT we estimated that the total thickness of the skull and dental cement was between 900um-1100um. Under a microscope, the Z-axis of the stereotaxic micromanipulator was advanced to a near touch of the bone surface. The skull thickness estimate was used to guide the advancement of the drill bit, which concluded with observation of wet bone chips and/or liquid.

Sealing the burr hole with a transparent silicone gel

Our design included a single burr hole, thus requiring us to continue using a single hole for many weeks during daily electrode penetrations. To accomplish this, we experimented with a silicone gel to seal the exposed dura at the burr hole.

Silicone gels can help preserve the integrity of the intra-cranial space after a craniotomy and in prolonging the life and functionality of burr holes. We wanted to use a gel that was non-toxic, transparent, elastic, easy to apply, and had resealing capabilities, allowing multiple penetrations by electrodes over an extended period of time.

We experimented with a commercially available soft silicone gel (DOWSIL 3–4680, Dow Corning, DuraGel, Cambridge Neurotech). This gel is a polydimethylsiloxane (PDMS) based silicone and is primarily used for hydrophobic encapsulation of electronic microchips. However, due to its biocompatible and antibacterial properties, it has recently been used as a dura substitute/cover for chronic recording in mice (Jackson and Muthuswamy, 2008;Jiang et al., 2017). These previous reports had tested biocompatibility, cytotoxicity, and sealing capability of the gel in mice, suggesting that it is a safe product, effective in sealing the dura from air contact, thereby minimizing possibility of infection. The published reports suggested that the gel had a number of attractive features: it was biocompatible, reduced possibility of inflammation, reduced cerebro-spinal fluid leakage, reduced humidity loss from the craniotomy, and remained soft for a period of days to weeks, thereby allowing microelectrode penetration with minimal force.

Immediately after drilling the burr hole and cleaning the remaining bone chips we covered it with the silicone gel (DuraGel, Cambridge Neurotech). Once the gel was cured (about 30 minutes), its mechanical properties allowed the electrodes to penetrate it for weeks. Moreover, the viscosity of the

gel allowed us to place it inside of the burr hole, thereby covering the dura completely, sealing it from air. The gel is transparent, which allowed us to monitor the dura beneath the gel.

Data acquisition

We recorded from the cerebellum using three types of electrodes: quartz insulated 4 fiber (tetrode) or 7 fiber (heptode) metal core (platinum/tungsten 95/05) electrodes (Thomas Recording), and 64 contact high density silicon probes (Cambridge Neurotech). None of these electrodes could penetrate the marmoset dura. Therefore, we performed a micro-duratomy using a 30G needle, which was installed on the stereotaxic micromanipulator frame and advanced to the surface of dura until a puncture was made. Once the puncture was made, the various electrodes could travel through the dura.

We connected each electrode to a 32 or 64 channel headstage amplifier and digitizer (Intan Technologies, USA), and then connected the headstage to a communication system (RHD2000, Intan Technologies, USA), as shown in Fig. 4. In addition, we connected to the RHD2000 system digital outputs from our custom behavioral software and analog output from an optical sensor mounted on the TV screen.

We used OpenEphys (Siegle et al., 2017), an open-source extracellular electrophysiology data acquisition software, for interfacing with the RHD2000 system and recording of signals. The signals were post processed and analyzed using MATLAB (Mathworks, USA) and Python.

Behavioral training

The next problem was to train the animals to perform goal-direct movements and remain motivated for about a thousand trials. In our task, following fixation of a center target, a primary target appeared at a random location. As the animal made a saccade to this primary target, that target was erased, and a secondary target was presented. The animal was rewarded if following the primary saccade it made a saccade to the secondary target and maintained fixation.

We implemented a training protocol consisting of multiple phases of reward based conditioning to familiarize the subjects with the experimental conditions and the demands of the tasks. Immediately following surgical recovery, the animal was placed on a food regulated diet. For 5 days a week, this diet consisted of 15g of lab diet power, 10g of apple/mango sauce, mixed in 30g of water. This resulted in net 40mL of deliverable food, which we administered via a syringe pump during the task. During the weekend, the food consisted of 30g for the first day and 25g for the second day of a solid lab diet.

Weight was monitored daily to ensure the health of the animal: weight was maintained within 85-100% of the average weight before the food regulation regime. If the weight fell below 85%, food regulation was stopped and the animal was fed in the colony until weight recovered to at least 90%. The animal was progressively acclimated to being held by a handler, entering its carrier independently, and sitting in its task chair. The experiment room was maintained at 74-84^{o}F.

Saccade training began with a frequency of 5 days/week . Visual targets were presented on a TV screen (Curved MSI 32" 144hz - model AG32CQ) while binocular eye movements were tracked using an EyeLink-1000 eye tracking system (SR Research, USA). We began with fixation training, followed by fixation with target-jump to induce a primary saccade. As the subject became accustomed to these paradigms, we introduced a secondary, corrective saccade. The amplitude of the primary target was 5 deg. The amplitude of the secondary target was 2 deg. At this point, we shifted towards the goal of

maximizing the number of successful trials that the subject was able to achieve, and cross referenced

this performance with her weight post food regulation. Successful trials resulted in food being dispensed

at a rate of 0.020mL/trial.

Data analysis

To detect simple spikes, we began with a high-pass filtered version of the signal (300 Hz), and then

subtracted at each time point the median of the signal across all contacts, a method known as common

average referencing. We then used a threshold based techniques to hand-sort the data. Here, we

consider only the largest spike recorded by each contact. To detect complex spikes, we relied on

frequency-domain analysis that tentatively identified these spikes via their power spectrum properties.

To confirm that the complex and simple spikes originated from the same cell, we compared the

conditional probability $\Pr\big(S(t)\big|C(0)\big)$ with $\Pr\big(S(t)\big|S(0)\big)$. These probabilities describe spike-triggered

histograms. For example, $\Pr\big(S(t)\big|C(0)\big)$ is the probability that a simple spike occurred at time *t*, given

that a complex spike was generated at time zero. $\Pr\big(S(t)\big|S(0)\big)$ is the probability that a simple spike

occurred at time *t*, given that a simple spike was generated at time zero. Simple spikes that originate

from a single cell produce a refractory period. Thus, $\Pr\big(S(t)\big|S(0)\big)$ should exhibit a low probability

period of roughly 10ms in duration centered at time zero. On the other hand, a complex spike

suppresses production of future simple spikes, but not those that occurred before. Thus, $\Pr\big(S(t)\big|C(0)\big)$

should be asymmetric, with a long period of low simple spike probability following time point zero.

Multi-contact electrodes allow for analysis of simultaneously recorded neurons. However,

spiking activity in one neuron can easily influence the data recorded by two nearby contacts, thus giving

an illusion that the two contacts are picking up two distinct neurons. To guard against this, after we

sorted the data in each contact, we waveform triggered the data recorded by contact A by the spikes

recorded in contact B. This identified the waveform of the neuron recorded by contact B on the

waveform recorded in contact A. We compared this cross-contact triggered waveform with the within

contact triggered waveform generated by the spikes recorded by contact A. The cross-contact triggered

waveform should produce a different cluster of spikes in A than the main neuron isolated by A. If there

were spikes that occurred within 1ms of each other in contacts A and B, we used these coincident-spike

events to trigger the waveform in A. The spikes in A that were identified to be coincident with B should

look approximately the same as the non-coincident spikes in A.

To quantify coordination between activities of two Purkinje cells, we computed conditional

probabilities. For contacts 1 and 2, we computed $\Pr\big(S1(t)\big|C2(0)\big)$, testing whether complex spikes in

contact 2 produced any changes in the simple spikes on contact 1. If the data were sorted properly, this

conditional probability should be essentially flat. To visualize coordination between the neurons, we

next plotted the spike-triggered waveform of voltages recorded by contact 1, triggered by the simple

spikes in contact 2. Finally, we computed $\Pr\big(S1(t)\big|S2(0)\big)$. This measure quantified whether the

occurrence of a simple spike in contact 2 altered the probability of simple spikes in contact 1.

Results

The methods that we described above allowed us to train subjects to produce around a thousand trials per session during cerebellar recordings. However, we arrived at these methods following a learning process that involved a number of unsuccessful attempts. Here, we describe our experience in terms of both types of results.

Behavioral results

For subject B, we experimented with saccade training without the head-post using a loose fitting head-restraining system (days -416 to 0, training without head-post, Fig. 5A). After a number of attempts we abandoned this approach because we were unable to produce robust calibration.

Following head-post surgery, we began behavioral training of subject B using the traditional approach (Mitchell et al., 2014;Johnston et al., 2018) of feeding the animals in the colony and supplementing that food with reward during task performance in the head-fixed condition (training without food regulation, Fig. 5A). We found that this was insufficient to motivate this subject.

We next instituted a different approach: eliminating home-cage feeding on experiment days, and instead providing task-based food rewards during the head-fixation period (food regulation, Fig. 5A and Fig. 5B). We ensured that regardless of performance, the animals always received their required allotment of food (at least 25 ml of food per day) and weight was maintained at around 90% of the pre-food regulation regime. However, this food was delivered only while the animal was head-fixed. This produced improved performance in subject B (days 911-982, Fig. 5A). We then fully tested this approach in subject M (Fig. 5B).

Following recovery from surgery, subject M was food restricted and was fed only in the head-fixed condition. Within 2 months subject M was able to perform around 1000 rewarded trials per session, and then maintained this performance for over a year in various tasks. During neurophysiological experiments, subject M produced 600-1500 rewarded trials per session (neurophysiology trials, running average data presented in Fig. 5B). Notably, during a year of food-restricted training, subject M remained healthy, as suggested by her stable weight and lack of complications.

An important component of our training was that the animals received a small amount of food per successful trial: 0.02 mL per trial. At completion of a successful trial, the computer generated a sound, and food delivery was signaled by the sound of the food pump. However, the low reward rate encouraged the animals to withhold licking until the food accumulated across multiple trials. In effect, the number of successful trials increased because the animals learned to perform the task in short blocks of rewarded trials.

Simultaneous recordings from pairs of Purkinje cells

We explored lobules V, VI and VII of the cerebellum using tetrodes, heptodes, and high density silicon arrays. We thought that the cranial ground screw would be required to provide a ground for our electrophysiological recordings. However, with experience we learned that the ground screw was unnecessary: the head-post by itself acted as a reliable reference for the electrical signals during neurophysiological recordings.

As the electrode approached the tentorium, there was a noticeable increase in background
activity, indicative of proximity to Purkinje cells and their high baseline discharge rate. When the
electrode passed through the tentorium, there was a distinct "pop", which was then followed by a
gradual increase in spiking activity.
Identification of a Purkinje cell required presence of both simple and complex spikes. Examples
of putative complex and simple spikes, recorded by a silicon array, a tetrode, and a heptode are
presented in Fig. 6A. Complex spikes have a stereotypical positive wave that has slow dynamics,
resulting in greater power in the lower frequencies (<300 Hz range) than the waveform of simple spikes
(Warnaar et al., 2015). Furthermore, complex spikes are relatively rare events, presenting an inter-spike
interval that is roughly two orders of magnitude more than simple spikes.
A useful real-time tool for identification of complex spikes is a low-pass filter (300 Hz), which
under good recording conditions can provide a tentative label for the complex spikes. Fig. 6B shows an
example of recording data from a heptode that simultaneously isolated three neurons: two putative
Purkinje cells, and a third unidentified neuron. In this figure, the voltages for each of the three
simultaneously recorded neurons are represented as black traces, and the low-pass filtered version of
the same signal is shown in red. The complex spikes are tentatively identified by the sharp increase in
the low-frequency power of the signal.
An important feature of a Purkinje cell is that production of a complex spike suppresses the
generation of simple spikes for a period of 10ms or longer. Thus, the shared cellular origin of complex
and simple spikes can be verified by using the complex spikes to trigger the voltage waveform. The
upper trace of Fig. 6C provides an example of this. We quantified this pattern via the condition
probability $\Pr\left(S(t)\middle|C(0)\right)$, which measures the probability of a simple spike at time t, given that a
complex spike occurred at time 0. This measure illustrates that the simple spikes occurred with equal
frequency before the complex spike, but were then completely suppressed for around 10ms following
the complex spike.
A second feature of a Purkinje cell is that the simple spikes have a refractory period. This feature
can be verified by using the simple spikes to trigger the voltage waveform. The lower trace of Fig. 6C
provides an example of this. We quantified this pattern via the conditional probability $\Pr\left(S(t)\middle|S(0)\right)$,
which measures the probability of a simple spike at time t, given that a simple spike occurred at time 0.
This measure illustrates that the simple spikes exhibited a 5ms refractory period.
Two examples of simultaneously recorded neurons from lobule VII during the saccade task are
shown in Fig. 7. In Fig. 7A, the heptode isolated three neurons, two of which were Purkinje cells (high-
pass filtered signals are shown in black, the low-pass filtered signal is shown in red). In this trial, the
animal was presented with a target at 5° on the horizontal axis and made a saccade at 220ms latency
(bottom row, Fig. 7A). The saccade landed about 0.5° short of the target, and was followed by a
secondary saccade. The Purkinje cell on contact 6 (2nd row, Fig. 7A) produced a complex spike before
the primary saccade. A second Purkinje cell, recorded in contact 5 (top row, Fig. 7A), did not produce a
complex spike before the onset of the primary saccade, but rather produced a complex spike before the
onset of the secondary saccade. A third neuron was isolated by contact 7, but this neuron was not a
Purkinje cell as it lacked complex spikes.

The waveforms for the complex and simple spikes in this recording session for each contact are shown in the left column of Fig. 7B. The waveforms illustrate the slower dynamics of the complex spikes, thus providing the basis for why a low pass filter is generally useful in their identification.

The middle column of Fig. 7B displays the probability densities of the inter-spike intervals for each type of spike. The number of simple spikes in a given period of time was roughly two orders of magnitude larger than the number of complex spikes.

The right column of Fig. 7B illustrates the within-cell timing properties of the spikes via conditional probabilities. $\Pr\big(S(t)\big|C(0)\big)$ demonstrates that following a complex spike at time 0, there was a silent period of ~10ms during which simple spikes were absent. Thus, production of a complex spike briefly suppressed production of simple spikes, demonstrating that the complex and simple spikes originated from the same Purkinje cell. $\Pr\big(S(t)\big|S(0)\big)$ demonstrates a 5ms refractory period, which is critical for demonstrating that the simple spikes were correctly attributed to a single Purkinje cell, and not misattributed because of cross-contaminated spikes from a neighboring contact. Finally, a comparison of $\Pr\big(S(t)\big|C(0)\big)$ with $\Pr\big(S(t)\big|S(0)\big)$ demonstrates that the complex-spike induced suppression of simple spikes was longer than the typical simple spike refractory period, another indication that the source of complex and simple spikes was a single Purkinje cells.

Another example of a recording session is shown in Fig. 7C. In this trial, the target was presented at 5^o along the vertical axis, and the animal made a saccade at a latency of 220ms. At saccade onset, the primary target was erased and a new target at 2^o to the right of the primary target was displayed. Thus, at the conclusion of the vertical saccade the target was not on the fovea, encouraging the animal to make a corrective horizontal saccade. This is an example of cross-axis adaptation (Xu-Wilson et al., 2009). A Purkinje cell was isolated on contact 1, while a second Purkinje cell was isolated on contact 7.

Fig. 7D illustrates the within contact timing properties of the complex and simple spikes. Like the data in Fig. 7B, the conditional probabilities demonstrate that simple spikes exhibited a 5m refractory period, and following a complex spike, there was suppression of simple spikes for a period of 10ms.

Spike timing properties of simultaneously recorded Purkinje cells
The ability to record simultaneously from multiple Purkinje cells allows one to ask whether nearby cells coordinate their activity, a function that may be fundamental in driving neurons in the cerebellar nucleus (Person and Raman, 2012). However, with multiple contacts there is a danger that some of the spikes that are recorded in contact A and attributed to Purkinje cell 1, are in fact generated by nearby Purkinje cell 2, which in turn is also being recorded by contact B. This cross-contact influence (cell 2 affecting recordings in both contacts A and B) would result in spurious coordination. One way to guard against this misattribution is to compare the spike-triggered waveforms.

Fig. 8A (left column) illustrates the waveform of simple spikes recorded by contact 6, during the same session as that shown in Fig. 7A. The middle column illustrates the waveform recorded by contact 6 but triggered by the simple spikes in contact 5. The data suggest that on average, the voltages recorded by contact 6 reflect in only a minor way the spikes being recorded by contact 5. Occasionally, the two P-cells produced a simple spike within 1ms of each other. The waveform in contact 6 as triggered by only these synchronous events (right column of Fig. 8A) looks very similar to the waveform

of the usual spikes (left column of Fig. 8A). Thus, these features suggest that any coordinated activity
present in the data of these two contacts would be because two independent cells fired simple spikes at
approximately the same time.
A useful technique to visualize the coordination between two Purkinje cells is by triggering the
voltages recorded by one contact via the simple spikes generated by a different Purkinje cell on a
separate contact. The resulting spike-triggered histogram of waveforms recorded by contact 6, triggered
by spikes in contact 5, is shown in Fig. 8B. The term $\Pr\big(S6(t)\big|S5(0)\big)$ measures the probability of simple
spikes on contact 6 at time t, given that a simple spike occurred on contact 5 at time 0. The conditional
probability exhibits a peak at around 0ms, suggesting a measure of coordination. This coordination could
have arisen because both Purkinje cells were saccade related, or driven by presentation of a visual
stimulus. It is also possible that these two cells received similar inputs (Heck et al., 2007), or that activity
in one affected activity in the other (Han et al., 2018).
The top plot of Fig. 8C illustrates the spike timing properties of these pairs of Purkinje cells. The
conditional probability $\Pr\big(S5(t)\big|C6(0)\big)$ quantifies the effect of complex spikes in contact 6 at time 0 on
the simple spikes in contact 5 at time t. It demonstrates that the complex spike in contact 6 did not
suppress the simple spikes in contact 5. Similarly, the conditional probability $\Pr\big(S6(t)\big|C5(0)\big)$
demonstrates that the complex spike in contact 5 did not suppress the simple spikes in contact 6. This
provides further evidence that the two nearby contacts picked up two distinct Purkinje cells.
The conditional probability $\Pr\big(C5(t)\big|C6(0)\big)$, displayed in the right plot of Fig. 8C, quantifies the
relationship between complex spikes in contacts 5 and 6. This plot illustrates that the complex spikes in
these two Purkinje cells were not well coordinated (little or no co-occurrence at 0ms latency).
In summary, these data from a pair of simultaneously recorded Purkinje cells, cells that were
likely less than 50μm apart, suggested that the cells were not served by the same inferior olive neuron.
However, the simple spikes exhibited a measure of coordination, suggesting that they may share
common inputs, or that the electrical activity in one cell influenced the activity in the other (Han et al.,
2018).
27
Silicone gel coating of the burr hole
We sealed the burr hole with a gel that we hoped would provide long-term protection to the dura, while
allowing for daily penetrations. We experimented with DuraGel, a silicone based polymer that is
primarily used for hydrophobic encapsulation of electronic microchips, but recently tested in mice as a
dura substitute and cover during chronic recordings (Jackson and Muthuswamy, 2008;Jiang et al., 2017).
Our experience with this product has been positive. Fig. 9 provides images of the burr hole after
drilling and following application of the gel. The gel remained in place for two weeks as we performed
daily recordings. At two weeks we removed the gel to examine the dura under a microscope, and then
took samples and performed tissue culture. This was repeated at intervals of two weeks. In all cases the
results suggested a healthy dura without any evidence of infection. Thus, DuraGel appears compatible
with the marmoset dura, providing long-term protection while allowing for daily electrode penetration.
39

Discussion

While the marmoset presents a number of attractive features as a primate model for the study of neural control of social behaviors, verbal communication, and saccadic eye movements, there is concern regarding whether the animals can be motivated to produce a sufficient number of trials. Previous reports on head-restrained marmosets had demonstrated operant conditioned behaviors in auditory tasks (Osmanski and Wang, 2011;Osmanski et al., 2013;Remington et al., 2012) as well as eye movement tasks (Mitchell et al., 2014;Johnston et al., 2018). However, performance has been rather limited: around 100 trials per session in the auditory task (Osmanski and Wang, 2011;Osmanski et al., 2013;Remington et al., 2012), 80-100 trials per session in the saccade task (Johnston et al., 2018), and 300-800 trials per session in the eye movement task (Mitchell et al., 2014). Here, we aimed to improve on these results, and found that through a behavioral training protocol, it was possible to motivate the animals to consistently perform around 1000 trials per session.

In addition, in order to investigate the neural basis of motor learning, we aimed to build tools that allowed for neurophysiological investigation of the cerebellum. We began with a CT+MRI based geometric model of the subject's skull, and used that data to compute the curvature of the base of a titanium head-post and chamber. Using the imaging data, we designed an alignment tool that had cylinders that guided an electrode through the burr hole via an absolute encoder microdrive. This resulted in stable recordings from multiple Purkinje cells during the saccade task.

Behavioral training

In previous reports, marmosets typically underwent a moderate food regulation regime and then were presented with the opportunity to acquire a form of liquid reward such as diluted apple sauce. In our approach, the subjects were gradually trained to receive their main source of nutrition as a consequence of rewarded behavior. Although our experience is limited to only two subjects, the results are encouraging: we were able to maintain the subjects' weight within roughly 90% of their pre-food regulation regime, keep them in excellent health, and motivate them to perform the task.

In previous reports, the amount of reward delivered per trial was relatively large: 0.1-0.2 mL per trial (Osmanski and Wang, 2011;Osmanski et al., 2013;Remington et al., 2012), 0.07 mL per trial (Johnston et al., 2018), or 0.05-0.06 mL per trial (Mitchell et al., 2014). Here, we trained the subjects to expect a lower reward rate, 0.02 mL per trial. The sound of the successful trial and the engagement of the food pump indicated food delivery, but the low reward rate encouraged the subjects to wait two or three correct trials before harvesting the accumulated food. This low reward rate may have been an important factor in motivating the subjects to consistently produce a relatively large number of trials per session.

Head-post design

Our head-post design departed significantly from epoxy-based helmet design that was employed in the past two decades in the study of the marmoset auditory cortex (Lu et al., 2001b;Wang et al., 2005;Eliades and Wang, 2008;Roy and Wang, 2012). Our design was made possible because of recent advances in 3D printing of titanium, allowing the use of imaging to build a subject-specific head-post.

The X-shaped design of our head-post was in contrast to the halo design recently demonstrated by the Everling group (Johnston et al., 2018) in the marmoset. Each design has its own advantages. The

halo design maximizes the open surface of the skull, thus allowing for access to many regions of the brain, which is ideal for simultaneous recording from multiple cortical areas. In contrast, our X-design limits the access to only the posterior regions, but has the advantage that the animal's skin can cover the entire head-post base, leaving only the chamber and the head-post pole exposed. Thus, our design allowed the skull to remain largely covered (for about a year following surgery, at which point the skin receded).

In our experience, the X-shaped head-post provided excellent stability: on occasion, during neurophysiological recording the animal would use its arms or legs to push against the chair, which then resulted in loss of the cell. However, this loss was temporary: often the electrode recovered the cell within a few seconds after the animal returned to its calm state.

The need to record from Purkinje cells during sensorimotor learning
One of the tenets of motor control is the idea that in order to move accurately, the brain relies on internal models that predict sensory consequences of motor commands. Evidence for this idea comes from human behavioral experiments (Maschke et al., 2004;Smith and Shadmehr, 2005;Izawa et al., 2012;Golla et al., 2008;Rabe et al., 2009;Donchin et al., 2012;Roth et al., 2013) and animal lesion studies (Buttner et al., 1994;Goffart et al., 2003;Barash et al., 1999;Machado et al., 2015), suggesting that a critical structure for forming internal models is the cerebellum.

However, it has been difficult to decipher how the cerebellum learns internal models: while in some tasks like smooth pursuit the activity of Purkinje cells is a simple function of eye velocity (Krauzlis, 2000), for most other tasks such as saccadic eye movements (Helmchen and Buttner, 1995;Thier et al., 2000), wrist movements (Ishikawa et al., 2014;Tanaka et al., 2019), and arm movements (Roitman et al., 2005;Roitman et al., 2009;Hewitt et al., 2011;Hewitt et al., 2015), it has been difficult to relate Purkinje cell activity to the ongoing motion. That is, the neural code with which the cerebellum represents internal models has been difficult to decipher.

Because learning involves a change in the neural encoding, this poor understanding of the representation of internal models has made it difficult to relate the rich behavioral changes that have been observed during motor learning, such as multiple timescales of memory (Smith et al., 2006;Kording et al., 2007), spontaneous recovery of memory (Ethier et al., 2008), and savings (Pekny et al., 2011), with the neural mechanisms of learning in the cerebellum. Our long-term goal in building this marmoset lab is to search for the neural basis of learning internal models in the cerebellum. However, the first step is to understand the neural code with which Purkinje cells make predictions.

Recently, using data in the macaque we proposed a new theory: the Purkinje cells may be organized in micro-clusters (Herzfeld et al., 2015;Herzfeld et al., 2018), wherein each micro-cluster is composed of Purkinje cells that share a common preference for error. The preference for error is expressed through the complex spike tuning of the Purkinje cell (Soetedjo and Fuchs, 2006;Soetedjo et al., 2008). We found that if Purkinje cells were organized in this way, the language of internal models as expressed by Purkinje cells became decipherable: simple spikes as a population predicted the real-time motion of the eye during a saccade (Herzfeld et al., 2015). Thus, this theory suggested a potential solution to the encoding problem: to investigate internal models, organize the simple spikes of a population of Purkinje cells based on the functional properties of each cell's complex spikes.

If the neural representation of internal model relies on a population coding of Purkinje cells, the challenge is to simultaneously record simple and complex spikes from multiple Purkinje cells in the awake, behaving animals. However, this has proven to be difficult. Array electrodes have simultaneously isolated multiple Purkinje cells in rodents (Welsh et al., 1995;Sugihara et al., 2007;Blenkinsop and Lang, 2011;Tang et al., 2016;Tang et al., 2019) and decerebrated cats (Ebner and Bloedel, 1981), but this has often been in the anesthetized animal (however, see (Han et al., 2018)). In the awake behaving subject, calcium imaging has been used to track complex spikes in multiple Purkinje cells (Najafi et al., 2014;Kostadinov et al., 2019), but this technique does not have the temporal resolution yet to track simple spikes. In primates, simultaneous recordings from Purkinje cells are quite rare. The report by Medina and Lisberger (2007) is the only example we are aware of in which pairs of Purkinje cells were recorded in the primate during goal-directed behavior.

Simultaneous recording from multiple Purkinje cells in the marmoset cerebellum
Although the marmoset is roughly the same weight as a rat, its cerebellum is about twice the size (Fujita et al., 2010). Similar to the rat, stripes of different aldolase C expression intensities separate the marmoset cerebellum into longitudinal compartments. However, olivo-cortical and cortico-nuclear projections indicate that the marmoset cerebellum has several compartments that are not present in rodents (compartments in the flocculus, nodulus, and the most lateral hemispheres) (Fujita et al., 2010). In comparison to the macaque, marmoset cerebellum has fewer folia, but a relatively larger vermis. Because the vermis is likely the region critical for saccadic eye movements, as well as control of the tongue during vocalization, the marmoset vermis may be particularly well suited for study of internal models in motor control.

We found that with modern electrodes it was possible to isolate multiple Purkinje cells from the vermis during goal-direct saccades. We employed tetrodes, heptodes, and silicon arrays, finding that each had its own advantages. Tetrodes and heptodes had a 3D geometry that gave them the potential to record from multiple Purkinje cells within the same folium, whereas silicon arrays provided the possibility of recording from Purkinje cells across folia.

We also found that the computational tools used for identifying Purkinje cells in other animals worked well in the marmoset. Like other animals, the complex spikes in the marmoset Purkinje cells had a waveform that exhibited slow dynamics, making it possible to use frequency based filtering techniques to tentatively distinguish complex spikes from simple spikes. Furthermore, following a complex spike, there was a 10-15ms period of simple spike suppression, helping to confirm that a single Purkinje cell produced both spikes.

Populations of Purkinje cells can coordinate their spiking, and this coordination may play a critical role in regulating activity of neurons in the deep cerebellar nucleus (Person and Raman, 2012). Perhaps the earliest demonstration of simple spike coordination was by Bell and Grimm (1969), who made microelectrode recordings in anesthetized cats, finding that Purkinje cells that were located <70um apart often fired nearly simultaneously. Later work found further evidence in support of this idea (Ebner and Bloedel, 1981;De Zeeuw et al., 1997;Shin and De, 2006;Heck et al., 2007;Wise et al., 2010). For example, Heck et al. (2007) demonstrated that in rats, P-cells within a few hundred microns fired synchronously during the period of reach to grasp of a food pellet, but not during the period after grasp completion.

Here we observed some evidence for sub-millisecond coordination in simple spikes of nearby Purkinje cells during performance of a saccade task. A recent study in mice (Han et al., 2018) also found that firing of many neighboring Purkinje cells exhibited sub-millisecond coordination. That study noted that when chemical synapses were blocked in brain slices, some synchrony persisted. They suggested that sub-millisecond coordination of neighboring Purkinje cells was not via shared synaptic inputs, but via electrical activity that in one Purkinje cell resulted in opening of sodium channels in the neighboring Purkinje cell. The question of whether Purkinje cells in the primate produce coordinated activity during goal-directed movements remains to be explored.

In summary, in order to better understand the neural basis of internal models, we have built a new laboratory that focuses on marmosets. Here, we reported on our attempts to improve marmoset behavioral training and electrophysiological recording methods. We found that through behavioral shaping, it was possible to motivate the animals to perform around 1000 trials in a head-fixed saccade task. We also found that with the aid of imaging, it was possible to build a system for simultaneous isolation of multiple Purkinje cells in the cerebellum.

1
2
Fig. 1. Design of the head-post and base chamber. **A**. Preoperative CT image that we used to build a
geometric model of the surface of the skull. **B**. Model of the head-post and chamber fitted to the CT-
based model of the animal's skull. The screws, head-post, and chamber are titanium. We used 2.4X8mm
screw for the cranial ground (0.08g), and 2.4x4mm screws for securing the legs of the head-post (0.04g).
C. The head-post (2.39g), designed to precisely fit the curved geometry of the subject. **D**. Base chamber
(0.48g).
9

2
Fig. 2. Development of a chamber-based geometric model to define electrode trajectories and specify
location of the burr hole. **A**. Pre-operative MRI image used for identifying the desired regions of interest
in the cerebellum. **B**. Reference axis ruler (1.104g) that was inserted into the base chamber before post-
op CT imaging. **C**. Post-operative CT image after the surgical installation of the head-post, base chamber,
and reference ruler. While the reference ruler is clearly visible, the titanium head-post and chamber
have produced significant artifacts. **D**. Co-registered pre-operative CT, post-operative CT, and pre-
operative MRI. Markers identify points on the reference axis and points in the cerebellum. A mark
identifies the burr hole location. **E**. A 3D model that has co-registered the skull, brain, chamber, and
reference ruler. **F**. Using the 3D model, we drew trajectories that began at points of interest in the
cerebellum, converged on a single 1.5mm diameter burr hole on the skull, and then diverged beyond
the chamber as cylinders within the reference ruler. These trajectories represented desired electrode
paths.
15
16

Fig. 3. Design of the electrode trajectory guidance tool. **A**. The alignment tool was designed to allow precise placement of the electrode along each of the 20 desired trajectories that converged onto various cerebellar destinations. The tool consisted of a 3D printed block (2.03g) that had a single cylinder for each of the desired electrode trajectories. The figure shows a model of the tool with 20 guide cylinders that align with the recording regions of interest and converge through the single craniotomy burr hole. **B**. Guide cylinder trajectories with posterior and sagittal views of the recording range on the cerebellum. The labels identify the cerebellar lobule that the trajectory is aimed toward. **C**. Alignment of the electrode with the desired trajectory. We installed the alignment tool in the base chamber and then placed a rod in the cylinder corresponding to the desired trajectory. The image on the right shows a section view of the alignment tool. We then placed a tube in the electrode holder (the same holder that would later hold the electrode), and attached the electrode holder to the microdrive. The microdrive was then maneuvered by a micromanipulator so that the axis of motion of the microdrive holding the tube was aligned with the rod. **D**. An image of the electrode and the electrode holder aligned to advance along the desired trajectory.

1
2

3
4
5 **Fig. 4.** Behavioral and electrophysiological recording system. AcB: acquisition board. BC: behavioral
6 computer. EpC: electrophysiological computer. EC: eye recording computer.
7
8

1
2
Fig. 5. Behavioral results. **A**. Weight and behavioral data in subject B. Training began without a head-
post, but was unsuccessful because of an inability to produce reliable calibration of eye measurements.
Following surgery, a long period of training without food regulation ensued, but was ultimately
abandoned because of the unwillingness of the animal to work for more than a few hundred trials.
Following food regulation, performance in the task dramatically improved. **B**. Weight and behavioral
data in subject M. Correct trials data represent running average of bin size 2.
9
10

1
2
Fig. 6. Recordings from the marmoset cerebellum. **A**. Complex and simple spikes recorded by a silicon
array, a tetrode, and a heptode. The complex spike is noted by an asterisk. **B**. Example of multiple
neurons simultaneously recorded by a heptode. The black traces are the high-pass filtered voltages and
the red trace is the low-pass filtered version of the same signal (300Hz cutoff). The top and bottom
traces are putative Purkinje cells while the middle trace is an unidentified third neuron. **C**. Spike-
triggered waveforms and histograms. In the upper trace, the voltages recorded by a single contact on a
heptode were triggered by the complex spikes on the same contact. The resulting complex spike-
triggered waveform exhibits a uniform pattern of simple spikes before the complex spike, and then
simple spike suppression. This pattern is quantified in $\Pr\big(S(t)\big|C(0)\big)$, which is the probability of a simple
spike at time *t*, given that a complex spike occurred at time 0. In the lower trace, the voltages were
triggered by the simple spikes. The resulting simple spike-triggered waveform exhibits a symmetric
period of spike suppression, reflecting the Purkinje cell's refractory period. This pattern is quantified in
$\Pr\big(S(t)\big|S(0)\big)$.
16
17

1
2
Fig. 7. Example of heptode recordings during two sessions in the saccade task. **A**. Sample trial,
illustrating activity of a Purkinje cell in contact 5, another Purkinje cell in contact 6, and an unidentified
neuron in contact 7. The black traces are high-pass filtered voltages and the red traces are low-passed
filtered signal (300Hz cutoff). The middle figure shows the geometry of the heptode and the location of
each contact. **B**. Spike waveforms and patterns of spike timing recorded from the session in part A. CS is
complex spike and SS is simple spike. ISI is inter-spike interval. $\Pr\big(S(t)\big|S(0)\big)$ is the probability of a
simple spike at time t, given that a simple spike occurred at time zero. $\Pr\big(S(t)\big|C(0)\big)$ is the probability of
a simple spike at time t, given that a complex spike occurred at time zero. **C**. Sample trial recorded in a
different session, illustrating activity of a Purkinje cell in contact 7, another Purkinje cell in contact 1, and
an unidentified neuron in contact 4. **D**. Spike waveforms and patterns of spike timing recorded from the
session in part B. Bin size is 1ms for conditional probabilities, 25ms for probability of complex spike ISI,
and 2ms for probability of simple spike ISI. Error bars on the spike waveforms are standard deviation.
15

1
2
Fig. 8. Spike timing property of a pair of simultaneously recorded Purkinje cells during the saccade task.
A. A Purkinje cell was isolated by contact 5, and another one by contact 6. The data recorded by contact
6 were triggered by simple spike events in contact 6 (left column), and simple spike events in contact 5
(middle column). The right column is data recorded by contact 6 and triggered if there was a simple
spike event in both contacts 6 and 5 within 1ms of each other. On average, spikes in one contact did not
produce a significant voltage change in another contact, and the shapes of synchronous (within 1ms)
and non-synchronous spikes were nearly identical. **B**. To measure coordination among the Purkinje cells,
we triggered the waveform in contact 6 by the spikes in contact 5. The pattern of coordination is
quantified via the conditional probability $\Pr\left(S6(t)\big|S5(0)\right)$. **C**. Left plot shows the probability of simple
spikes at time t, given that a complex spike occurred at time zero in another channel. There is no
suppression of simple spikes following a complex spike. The plot on the right shows the probability of
co-occurrence of the complex spikes. The two cells did not produce complex spikes that were strongly
coordinated.
16

1
2

1.5mm

Day 0: burr hole   Day 0: dura gel

Day 5   Day 10

3
4   **Fig. 9**. Images from the burr hole before (day 0) and after being sealed by silicone gel.
5
6

1   Reference List

Barash S, Melikyan A, Sivakov A, Zhang M, Glickstein M, Thier P (1999) Saccadic dysmetria and
adaptation after lesions of the cerebellar cortex. J Neurosci 19:10931-10939.

Bell CC, Grimm RJ (1969) Discharge properties of Purkinje cells recorded on single and double
microelectrodes. J Neurophysiol 32:1044-1055.

Blenkinsop TA, Lang EJ (2011) Synaptic action of the olivocerebellar system on cerebellar nuclear spike
activity. J Neurosci 31:14708-14720.

Buttner U, Straube A, Spuler A (1994) Saccadic dysmetria and "intact" smooth pursuit eye movements
after bilateral deep cerebellar nuclei lesions. J Neurol Neurosurg Psychiatry 57:832-834.

Courchesne E, Karns CM, Davis HR, Ziccardi R, Carper RA, Tigue ZD, Chisum HJ, Moses P, Pierce K, Lord C,
Lincoln AJ, Pizzo S, Schreibman L, Haas RH, Akshoomoff NA, Courchesne RY (2001) Unusual brain growth
patterns in early life in patients with autistic disorder: an MRI study. Neurology 57:245-254.

De Zeeuw CI, Koekkoek SK, Wylie DR, Simpson JI (1997) Association between dendritic lamellar bodies
and complex spike synchrony in the olivocerebellar system. J Neurophysiol 77:1747-1758.

Donchin O, Rabe K, Diedrichsen J, Lally N, Schoch B, Gizewski ER, Timmann D (2012) Cerebellar regions
involved in adaptation to force field and visuomotor perturbation. J Neurophysiol 107:134-147.

Ebner TJ, Bloedel JR (1981) Correlation between activity of Purkinje cells and its modification by natural
peripheral stimuli. J Neurophysiol 45:948-961.

Eliades SJ, Miller CT (2017) Marmoset vocal communication: Behavior and neurobiology. Dev Neurobiol
77:286-299.

Eliades SJ, Wang X (2008) Chronic multi-electrode neural recording in free-roaming monkeys. J Neurosci
Methods 172:201-214.

Eliades SJ, Wang X (2013) Comparison of auditory-vocal interactions across multiple types of
vocalizations in marmoset auditory cortex. J Neurophysiol 109:1638-1657.

Ethier V, Zee DS, Shadmehr R (2008) Spontaneous recovery of motor memory during saccade
adaptation. J Neurophysiol 99:2577-2583.

Fedorov A, Beichel R, Kalpathy-Cramer J, Finet J, Fillion-Robin JC, Pujol S, Bauer C, Jennings D, Fennessy
F, Sonka M, Buatti J, Aylward S, Miller JV, Pieper S, Kikinis R (2012) 3D Slicer as an image computing
platform for the Quantitative Imaging Network. Magn Reson Imaging 30:1323-1341.

Fujita H, Oh-Nishi A, Obayashi S, Sugihara I (2010) Organization of the marmoset cerebellum in three-
dimensional space: lobulation, aldolase C compartmentalization and axonal projection. J Comp Neurol
518:1764-1791.

Goffart L, Chen LL, Sparks DL (2003) Saccade dysmetria during functional perturbation of the caudal
fastigial nucleus in the monkey. Ann N Y Acad Sci 1004:220-228.

Golla H, Tziridis K, Haarmeier T, Catz N, Barash S, Thier P (2008) Reduced saccadic resilience and impaired saccadic adaptation due to cerebellar disease. Eur J Neurosci 27:132-144.

Han KS, Guo C, Chen CH, Witter L, Osorno T, Regehr WG (2018) Ephaptic Coupling Promotes Synchronous Firing of Cerebellar Purkinje Cells. Neuron 100:564-578.

Hashimoto T, Tayama M, Murakawa K, Yoshimoto T, Miyazaki M, Harada M, Kuroda Y (1995) Development of the brainstem and cerebellum in autistic patients. J Autism Dev Disord 25:1-18.

Heck DH, Thach WT, Keating JG (2007) On-beam synchrony in the cerebellum as the mechanism for the timing and coordination of movement. Proc Natl Acad Sci U S A 104:7658-7663.

Helmchen C, Buttner U (1995) Saccade-related Purkinje cell activity in the oculomotor vermis during spontaneous eye movements in light and darkness. Exp Brain Res 103:198-208.

Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R (2015) Encoding of action by the Purkinje cells of the cerebellum. Nature 526:439-442.

Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R (2018) Encoding of error and learning to correct that error by the Purkinje cells of the cerebellum. Nat Neurosci 21:736-743.

Hewitt AL, Popa LS, Ebner TJ (2015) Changes in Purkinje cell simple spike encoding of reach kinematics during adaption to a mechanical perturbation. J Neurosci 35:1106-1124.

Hewitt AL, Popa LS, Pasalar S, Hendrix CM, Ebner TJ (2011) Representation of limb kinematics in Purkinje cell simple spike discharge is conserved across multiple tasks. J Neurophysiol 106:2232-2247.

Ishikawa T, Tomatsu S, Tsunoda Y, Lee J, Hoffman DS, Kakei S (2014) Releasing dentate nucleus cells from Purkinje cell inhibition generates output from the cerebrocerebellum. PLoS One 9:e108774.

Izawa J, Criscimagna-Hemminger SE, Shadmehr R (2012) Cerebellar contributions to reach adaptation and learning sensory consequences of action. J Neurosci 32:4230-4239.

Jackson N, Muthuswamy J (2008) Artificial dural sealant that allows multiple penetrations of implantable brain probes. J Neurosci Methods 171:147-152.

Jiang Z, Huxter JR, Bowyer SA, Blockeel AJ, Butler J, Imtiaz SA, Wafford KA, Phillips KG, Tricklebank MD, Marston HM, Rodriguez-Villegas E (2017) TaiNi: Maximizing research output whilst improving animals' welfare in neurophysiology experiments. Sci Rep 7:8086.

Johnston KD, Barker K, Schaeffer L, Schaeffer D, Everling S (2018) Methods for chair restraint and training of the common marmoset on oculomotor tasks. J Neurophysiol 119:1636-1646.

Kishi N, Sato K, Sasaki E, Okano H (2014) Common marmoset as a new model animal for neuroscience research and genome editing technology. Dev Growth Differ 56:53-62.

Kording KP, Tenenbaum JB, Shadmehr R (2007) The dynamics of memory as a consequence of optimal adaptation to a changing body. Nat Neurosci 10:779-786.

Kostadinov D, Beau M, Pozo MB, Hausser M (2019) Predictive and reactive reward signals conveyed by climbing fiber inputs to cerebellar Purkinje cells. Nat Neurosci 22:950-962.

Krauzlis RJ (2000) Population coding of movement dynamics by cerebellar Purkinje cells. Neuroreport 11:1-6.

Lu T, Liang L, Wang X (2001a) Neural representations of temporally asymmetric stimuli in the auditory cortex of awake primates. J Neurophysiol 85:2364-2380.

Lu T, Liang L, Wang X (2001b) Temporal and rate representations of time-varying signals in the auditory cortex of awake primates. Nat Neurosci 4:1131-1138.

MacDougall M, Nummela SU, Coop S, Disney A, Mitchell JF, Miller CT (2016) Optogenetic manipulation of neural circuits in awake marmosets. J Neurophysiol 116:1286-1294.

Machado AS, Darmohray DM, Fayad J, Marques HG, Carey MR (2015) A quantitative framework for whole-body coordination reveals specific deficits in freely walking ataxic mice. Elife 4.

Marko MK, Crocetti D, Hulst T, Donchin O, Shadmehr R, Mostofsky SH (2015) Behavioural and neural basis of anomalous motor learning in children with autism. Brain 138:784-797.

Maschke M, Gomez CM, Ebner TJ, Konczak J (2004) Hereditary cerebellar ataxia progressively impairs force adaptation during goal-directed arm movements. J Neurophysiol 91:230-238.

Medina JF, Lisberger SG (2007) Variation, signal, and noise in cerebellar sensory-motor processing for smooth-pursuit eye movements. J Neurosci 27:6832-6842.

Miller CT, Freiwald WA, Leopold DA, Mitchell JF, Silva AC, Wang X (2016) Marmosets: A Neuroscientific Model of Human Social Behavior. Neuron 90:219-233.

Mitchell JF, Reynolds JH, Miller CT (2014) Active vision in marmosets: a model system for visual neuroscience. J Neurosci 34:1183-1194.

Murakami JW, Courchesne E, Press GA, Yeung-Courchesne R, Hesselink JR (1989) Reduced cerebellar hemisphere size and its relationship to vermal hypoplasia in autism. Arch Neurol 46:689-694.

Mustoe AC, Cavanaugh J, Harnisch AM, Thompson BE, French JA (2015) Do marmosets care to share? Oxytocin treatment reduces prosocial behavior toward strangers. Horm Behav 71:83-90.

Najafi F, Giovannucci A, Wang SS, Medina JF (2014) Sensory-driven enhancement of calcium signals in individual Purkinje cell dendrites of awake mice. Cell Rep 6:792-798.

Osmanski MS, Song X, Wang X (2013) The role of harmonic resolvability in pitch perception in a vocal nonhuman primate, the common marmoset (Callithrix jacchus). J Neurosci 33:9161-9168.

Osmanski MS, Wang X (2011) Measurement of absolute auditory thresholds in the common marmoset (Callithrix jacchus). Hear Res 277:127-133.

Pekny SE, Criscimagna-Hemminger SE, Shadmehr R (2011) Protection and expression of human motor memories. J Neurosci 31:13829-13839.

Person AL, Raman IM (2012) Purkinje neuron synchrony elicits time-locked spiking in the cerebellar nuclei. Nature 481:502-505.

Rabe K, Livne O, Gizewski ER, Aurich V, Beck A, Timmann D, Donchin O (2009) Adaptation to visuomotor rotation and force field perturbation is correlated to different brain areas in patients with cerebellar degeneration. J Neurophysiol 101:1961-1971.

Remington ED, Osmanski MS, Wang X (2012) An operant conditioning method for studying auditory behaviors in marmoset monkeys. PLoS One 7:e47895.

Roitman AV, Pasalar S, Ebner TJ (2009) Single trial coupling of Purkinje cell activity to speed and error signals during circular manual tracking. Exp Brain Res 192:241-251.

Roitman AV, Pasalar S, Johnson MT, Ebner TJ (2005) Position, direction of movement, and speed tuning of cerebellar Purkinje cells during circular manual tracking in monkey. J Neurosci 25:9244-9257.

Roth MJ, Synofzik M, Lindner A (2013) The cerebellum optimizes perceptual predictions about external sensory events. Curr Biol 23:930-935.

Roy S, Miller CT, Gottsch D, Wang X (2011) Vocal control by the common marmoset in the presence of interfering noise. J Exp Biol 214:3619-3629.

Roy S, Wang X (2012) Wireless multi-channel single unit recording in freely moving and vocalizing primates. J Neurosci Methods 203:28-40.

Sasaki E, et al. (2009) Generation of transgenic non-human primates with germline transmission. Nature 459:523-527.

Scott JA, Schumann CM, Goodlin-Jones BL, Amaral DG (2009) A comprehensive volumetric analysis of the cerebellum in children and adolescents with autism spectrum disorder. Autism Res 2:246-257.

Shin SL, De SE (2006) Dynamic synchronization of Purkinje cell simple spikes. J Neurophysiol 96:3485-3491.

Siegle JH, Lopez AC, Patel YA, Abramov K, Ohayon S, Voigts J (2017) Open Ephys: an open-source, plugin-based platform for multichannel electrophysiology. J Neural Eng 14:045003.

Smith MA, Ghazizadeh A, Shadmehr R (2006) Interacting adaptive processes with different timescales underlie short-term motor learning. PLoS Biol 4:e179.

Smith MA, Shadmehr R (2005) Intact ability to learn internal models of arm dynamics in Huntington's disease but not cerebellar degeneration. J Neurophysiol 93:2809-2821.

Soetedjo R, Fuchs AF (2006) Complex spike activity of purkinje cells in the oculomotor vermis during behavioral adaptation of monkey saccades. J Neurosci 26:7741-7755.

Soetedjo R, Kojima Y, Fuchs AF (2008) Complex spike activity in the oculomotor vermis of the cerebellum: a vectorial error signal for saccade motor learning? J Neurophysiol 100:1949-1966.

Solomon SG, Rosa MG (2014) A simpler primate brain: the visual system of the marmoset monkey. Front Neural Circuits 8:96.

Sugihara I, Marshall SP, Lang EJ (2007) Relationship of complex spike synchrony bands and climbing fiber projection determined by reference to aldolase C compartments in crus IIa of the rat cerebellar cortex. J Comp Neurol 501:13-29.

Takagi M, Zee DS, Tamargo RJ (1998) Effects of lesions of the oculomotor vermis on eye movements in primate: saccades. J Neurophysiol 80:1911-1931.

Takahashi DY, Liao DA, Ghazanfar AA (2017) Vocal Learning via Social Reinforcement by Infant Marmoset Monkeys. Curr Biol 27:1844-1852.

Tanaka H, Ishikawa T, Kakei S (2019) Neural Evidence of the Cerebellum as a State Predictor. Cerebellum 18:349-371.

Tang T, Blenkinsop TA, Lang EJ (2019) Complex spike synchrony dependent modulation of rat deep cerebellar nuclear activity. eLife 8.

Tang T, Suh CY, Blenkinsop TA, Lang EJ (2016) Synchrony is Key: Complex Spike Inhibition of the Deep Cerebellar Nuclei. Cerebellum 15:10-13.

Thier P, Dicke PW, Haas R, Barash S (2000) Encoding of movement time by populations of cerebellar Purkinje cells. Nature 405:72-76.

Wakabayashi M, Koketsu D, Kondo H, Sato S, Ohara K, Polyakova Z, Chiken S, Hatanaka N, Nambu A (2018) Development of stereotaxic recording system for awake marmosets (Callithrix jacchus). Neurosci Res 135:37-45.

Wang X (2018) Cortical Coding of Auditory Features. Annu Rev Neurosci 41:527-552.

Wang X, Lu T, Snider RK, Liang L (2005) Sustained firing in auditory cortex evoked by preferred stimuli. Nature 435:341-346.

Warnaar P, Couto J, Negrello M, Junker M, Smilgin A, Ignashchenkova A, Giugliano M, Thier P, De SE (2015) Duration of Purkinje cell complex spikes increases with their firing frequency. Front Cell Neurosci 9:122.

Welsh JP, Lang EJ, Suglhara I, Llinas R (1995) Dynamic organization of motor control within the olivocerebellar system. Nature 374:453-457.

Whitney ER, Kemper TL, Bauman ML, Rosene DL, Blatt GJ (2008) Cerebellar Purkinje cells are reduced in a subpopulation of autistic brains: a stereological experiment using calbindin-D28k. Cerebellum 7:406-416.

1   Wise AK, Cerminara NL, Marple-Horvat DE, Apps R (2010) Mechanisms of synchronous activity in
2   cerebellar Purkinje cells. J Physiol 588:2373-2390.

3   Xu-Wilson M, Chen-Harris H, Zee DS, Shadmehr R (2009) Cerebellar contributions to adaptive control of
4   saccades in humans. J Neurosci 29:12930-12939.
5
6