Restoring ancestral phenotypes by reduction of plasticity is a general

pattern in gene expression during adaptation to different stressors in

Tribolium castaneum

Eva L. Koch[1] and Frédéric Guillaume[1]

[1] Department of Evolutionary Biology and Environmental Studies, University of Zürich, Winterthurerstr. 190, 8057 Zürich, Switzerland

Corresponding author:
Prof. Dr. Frédéric Guillaume
tel: ++41 44 635 6623
frederic.guillaume@ieu.uzh.ch

Abstract

Plasticity and evolution are two processes individuals to respond to environmental change, but how both are related and impact each other is still controversial. We studied plastic and evolutionary responses in gene expression of *Tribolium castaneum* after exposure to new environments that differed from ancestral conditions in temperature, humidity or both. Using experimental evolution with ten replicated lines per condition, we were able to demonstrate adaptation after 20 generations. We measured gene expression in each condition in adapted selection lines and control lines to infer evolutionary and plastic changes. We found more evidence for changes in mean expression (shift in the intercept of reaction norms) in adapted lines than for changes in plasticity (shifts in slopes). Plasticity was mainly preserved and was responsible for a large part of the phenotypic divergence in expression between ancestral and new conditions. However, we found that genes with the largest evolutionary changes in expression also evolved reduced plasticity and often showed expression levels closer to the ancestral stage. Results obtained in the three different conditions were similar suggesting that restoration of ancestral expression levels during adaptation is a general evolutionary pattern. We increased the sample size in the most stressful condition and were then able to detect a positive correlation between proportion of genes with reversion of the ancestral plastic response and mean fitness per selection line.

Introduction

Whenever facing environmental change, populations can adapt to new phenotypic optima by plasticity and evolution. Plasticity is the ability of a single genotype to produce multiple phenotypes as a function of the environment. It is often seen as an immediate response of individuals to changes in their environment. In contrast, evolution requires a change in allele

frequencies within a population. This process occurs over several generations and represents a

more long-term response, which can result in local adaptation. It is still not well understood how

these two processes are related and interact with each other (de Jong, 2005; Forsman, 2015;

Ghalambor, McKay, Carroll, & Reznick, 2007; Price, Qvarnström, & Irwin, 2003; Wund, 2012).

Plasticity does not affect a population's genetic composition and should thus not have long-term

consequences in the following generations. However, it changes the distribution of phenotypes

on which selection can act (de Jong, 2005; Pfennig et al., 2010; Pigliucci, 2005; Price,

Qvarnstrom, & Irwin, 2003). If plasticity perfectly matches the new phenotypic optimum, it

prevents evolution since no genetic changes are required and selection is prevented (Ghalambor

et al., 2007). On the other hand, plasticity is also crucial for a population's persistence and can

reduce the costs of selection (Chevin, Lande, & Mace, 2010; Pavey, Collin, Nosil, & Rogers, 2010).

It prevents extinction and protects populations from bottleneck effects, thereby maintaining a

higher genetic variation on which subsequently selection can act (Fitzpatrick, 2012; Pfennig et

al., 2010; Massimo Pigliucci, 2005). There is both, theoretical (Chevin et al., 2010; Draghi &

Whitlock, 2012; Fierst, 2011) and empirical work (Schaum, Rost, Millar, & Collins, 2013),

demonstrating that more plastic populations exhibit faster evolution. The benefits of plasticity

for persisting in new habitats were also demonstrated in invasive species (Molina-Montenegro,

Peñuelas, Munné-Bosch, & Sardans, 2012; Pichancourt & van Klinken, 2012; Yeh & Price, 2004).

The extent of plasticity can be represented as a reaction norm (Scheiner, 1993), which is the

phenotypic trait value as a function of an environmental variable. Evolution can affect the

reaction norm in two ways: The intercept can be shifted corresponding to a change in the mean

phenotypic value. Alternatively, the slope of the reaction norm, i.e. the plasticity, can be

changed. Thus, plasticity itself can also be subject to evolution given that there is sufficient

genetic variation in reaction norms (Garland & Kelly, 2008; Nussey, Postma, Gienapp, & Visser,

2005).

Dependent on whether plastic responses are adaptive and increase fitness of an individual,

different patterns describing the relationship between ancestral plasticity and evolution are

possible. If plastic responses are adaptive, but not sufficient to reach the phenotypic optimum,

evolution should work in the same direction as the plastic response (referred to as Baldwin effect

(Crispo, 2007) or cogradient variation (Conover, Duffy, & Hice, 2009). In this case, selection may

favour the most plastic individuals, causing evolved populations to exhibit a higher plasticity than

their ancestors (Crispo 2007; Lande 2009). Another possible outcome is genetic assimilation: An

initially environmentally induced phenotype can get fixed by a loss of plasticity and becomes

continuously expressed even in the ancestral environment (Levis & Pfennig, 2016; Pigliucci,

Murren, & Schlichting, 2006). In contrast, if plastic responses are maladaptive, we expect to

observe evolutionary changes opposite to plasticity (countergradient variation ((Conover et al.,

2009) or genetic compensation (Grether 2005)). Maladaptive plasticity was proposed as a

possible mechanism promoting evolution since it moves phenotypes further away from the

optimum and thereby increases the strength of selection on the phenotype (Ghalambor et al.,

2007). Both co-gradient (Barton, Sunnucks, Norgate, Murray, & Kearney, 2014; Conover et al.,

2009) and counter-gradient (Conover et al., 2009; Ghalambor et al., 2015; Laugen, Laurila,

Räsänen, & Merilä, 2003) evolutionary changes have been found, indicating that adaptive and

maladaptive plasticity are common. Reversion of ancestral plasticity occurs more frequently (Ho

& Zhang, 2018), indicating that plastic responses are often not beneficial for long-term

adaptation.

Plastic responses in physiology, behaviour or morphological traits are often initiated by changes

in gene expression (Hodgins-Davis & Townsend, 2009; Wray, 2007). The transcriptome

represents a direct link between genotype and phenotype making it particularly interesting to

study the interplay between plasticity and evolution. Transcription is highly plastic and

modulating expression levels is an important part of an organism's physiological adjustment to

environmental change (Gibson, 2008; McCairns & Bernatchez, 2009). On the other hand, there

are also many studies demonstrating evolutionary divergence in gene expression between locally

adapted populations (Alvarez, Schrey, & Richards, 2015; Guo et al., 2016; Romero, Ruvinsky, &

Gilad, 2012; Townsend, Cavalieri, & Hartl, 2003; Whitehead & Crawford, 2006). Gene expression

may even evolve more rapidly than changes in proteins since mutations affecting the magnitude

of expression are less likely to be deleterious than changes in protein structures (Carroll, 2005;

Wray, 2007). However, it is not clear how fast plasticity in gene expression can change. Some

studies reported changes in plasticity in few genes after adaptation to new conditions (Morris et

al., 2014; Passow et al., 2017; von Heckel, Stephan, & Hutter, 2016), whereas others found only

limited evolution of plasticity (Yampolsky, Glazko, & Fry, 2012) or less than expected (Huang &

Agrawal, 2016).

In our study, we used whole transcriptomes to understand the interplay between plasticity and

evolution at the gene expression level during adaptation to new environments with the model

organism *Tribolium castaneum* (red flour beetle) using an experimental evolution approach.

With RNA-seq, we quantified plastic and evolutionary changes in gene expression and their

contribution to the total phenotypic divergence between populations inhabiting different

environments. More specifically, we were interested to test whether the same genes exhibited

both evolutionary and plastic changes in a new environment and whether evolved changes were

in the same direction as their ancestral plasticity. In doing so, we could test how plasticity

affected evolution. We tested whether plasticity prevented evolution, or whether maladaptive

plasticity increased the strength of selection and promoted evolution. Next, we tested how

evolution affected gene expression reaction norms, i.e. whether adaptation to new conditions

was achieved by changes in slopes or intercepts.

Material and Methods

Animal rearing, experimental evolution

We used the *Tribolium castaneum* Cro1 strain (Milutinović, Stolpe, Peuß, Armitage, & Kurtz,

2013), collected from a wild population in 2010 and adapted to lab standard conditions (32°C,

70% relative humidity (r.h.)) for more than 20 generations. Beetles were kept in 24h darkness on

organic wheat flour mixed with 10% organic baker's yeast. We sterilized flour and yeast by

heating them for 12h at 80°C before use. To test for adaptation to new environmental conditions

we used replicate lines and exposed them to three treatment and control (CT) conditions. The

conditions in the treatments were: Dry (D): 33°C and 30% r. h.; Hot (H): 37°C and 70% r. h.; Hot-

Dry (HD): 37°C and 30% r. h. To generate replicate lines, we used 120 individuals (60 females

and 60 males in the pupal stage) and placed them into a vial containing 80g medium. We

produced six lines per selection regime (treatments plus control), resulting in a total of 24 lines.

For each new generation, we randomly collected 120 pupae and placed them into a new vial.

After seven to ten days, in which the pupae became adults, mated and laid eggs, adult beetles

were removed by sieving the medium. We waited until the next generation (eggs/larvae in the

medium) had reached the pupal stage and again collected 120 pupae per line to establish the

next generation. This is similar to natural selection since individuals, depending on their fitness,

do not contribute equally to the next generation. In generation 15 we produced additional mixed

lines to prevent loss of genetic diversity by gene drift and inbreeding, which might impede

adaptation: We mixed the six replicate lines of each selection regime in equal proportions (20

individuals from each replicate line) four times, resulting in four mixed lines with 120 individuals

each. In total we had 39 lines: six normal and four mixed lines per selection regime (one line in D

became extinct). The transplant experiment to test for adaptation was conducted in generation

22.

Reciprocal transplant and fitness assay

Before testing for adaptation, all lines stayed for two generations in the same condition to

remove potential maternal or epigenetic effects (Supporting information, Figure S1): Beetles of

generation 20 from all selection lines were transferred to control conditions, in which they

stayed for one week to mate and lay eggs. After removal of the adults, we waited until their

offspring had reached the pupal stage and separated males and females. These individuals

(generation 21) developed completely in control conditions. When they reached the adult stage,

we created 13 full-sib families per selection line by transferring one virgin male with one virgin

female of the same selection line in 15mL tubes with 1g of medium. After four days, in which the

beetles could mate and lay eggs, 9g of medium was added to provide food for the developing

offspring and each mating pair was transferred to a new vial. We repeated this three times,

resulting in four vials per mating pair containing medium and eggs. Immediately after removal of

the mating pair, vials of each mating pair were randomly assigned to the four different

conditions, resulting in full-sib families split across all conditions. These beetles were transferred

to the treatments at the egg stage. As soon as offspring in these vials had reached the pupal

stage, males and females (four females and four males per family and condition) were separated

and transferred to 15 mL tubes with 5 g of medium and remained there until they were used for

the fitness assay two weeks later. They developed completely in treatment conditions. We then

assessed their performance in each condition by estimating their fitness to test for adaptation. A

virgin male and a virgin female of the same selection line from the same condition but from

different families were again placed into a 15 mL tube with 1 g medium. After four days, the

mating pair was removed. Males and females were transferred to 1 mL Eppendorf tubes (one

individual per tube), immediately frozen in liquid nitrogen and stored at -80°C to use them for

gene expression measurements. 9 g medium was added to the mating tube. After four weeks (in

CT and H) or five weeks (D and HD), all offspring had reached the adult stage and were counted.

We used the number of adult offspring as an estimate of the fitness of a mating pair.

Statistical analysis

To test whether selection regime significantly influenced number of offspring produced and test

whether 20 generations in the treatments resulted in adaptation, we compared offspring

numbers of selection lines in their native condition to CT-lines transferred to the same condition.

We applied linear mixed models using the R-packages *lme4* (Bates, Mächler, Bolker, & Walker,

2015), and *lmertest* (Kuznetsova, Brockhoff, & Christensen, 2017) and *lsmeans* (Lenth, 2016) to

obtain *p*-values and confidence intervals. We included line and family as random factors,

selection and line type (mixed/normal) and their interaction as fixed effects. To test whether the

selection regime influenced how lines responded to the treatments, we used a linear mixed

model with offspring number in CT and treatment conditions as response variable, condition,

selection regime, line type (normal/mixed) and interactions as fixed effects and line, family and

interaction between line and condition as random effects. A significant interaction between

condition and selection regime indicates a significant effect of the selection regime (evolution)

on the response to the conditions (plastic response).

RNA extraction, library preparation and sequencing

208 female beetles (Table 1) p stored at -80°C were homogenized in Tri-Reagent® (Zymo

Research, California, USA) using an electric bead mill. RNA was extracted with the RNA Mini Prep

kit (Zimo Research, California, USA) following the instructions of the manufacturer. RNA-quality

was checked on a TapeStation (Agilent, Waldbronn, Germany) and concentrations were

measured with aQubit® Fluorometer (Life Technologies, California, USA). Libraries were created

with 500 ng RNA for each individual separately with the LEXOGEN mRNA-Seq Library Kit

following the manual (LEXOGEN GmbH, Vienne, Austria). Library quality was checked on a

TapeStation (Agilent, Waldbronn, Germany) and concentrations were determined by qPCR.

Libraries were diluted to the same molarity and pooled (33-36 libraries per pool). All treatments

and selection regimes were randomized during RNA-extraction, library preparation, and

sequencing. Single-end sequencing was performed in five runs on the Illumina NextSeq 500

(Illumina, Inc, California, USA) using the 75 cycles High Output Kit. After quality control using

FastQC (www.bioinformatics.bbsrc.ac.uk/projects/fastqc) reads (adaptors were trimmed and the

first 10 bases were hard trimmed, minimum average quality Q10, minimum tail quality 10,

minimum read length 20) were mapped against the reference genome

(ftp://ftp.ensemblgenomes.org/pub/release30/metazoa/gtf/tribolium_castaneum/Tribolium_ca

staneum.Tcas3.30.gtf.gz) with STAR v.2.5 (Dobin et al., 2013). We then used FeatureCounts (Liao,

Smyth, & Shi, 2014) to count the number of reads that mapped to each gene in the reference

genome. Mapping as well as read counting was performed within the data analysis framework

SUSHI (Hatakeyama et al., 2016).

Gene expression analysis

Gene expression analysis was done in R (R Core Team, 2017). We used the R package *edgeR*

(Robinson, McCarthy, & Smyth, 2010) for normalizing (method: TMM) expression data to cpm

(counts per million) after filtering lowly expressed genes (minimum of one cpm in at least two

samples). For subsequent differential expression analysis we used the R package *limma* (Law, Chen, Shi, & Smyth, 2014; Ritchie et al., 2015). From the differential expression analysis, we obtained the number of differentially expressed genes (DE genes) within lines between conditions (plastic changes, see Figure S1) or between lines of different origins (CT vs. selection) within conditions (evolutionary changes, see Figure S1). The total phenotypic divergence in gene expression between CT and treatments (i.e. total change TC) is the differential expression (log2-fold change) between CT-lines in CT and selection lines in the treatments (Figure 1A-C and Supporting information Figure S1). The ancestral plasticity (PC_{CT}) is the differential expression of CT-lines between CT and treatment conditions, while the evolved plasticity (PC_{Sel}) is the same difference measured in selection lines. The evolutionary changes are EC_T when measured as differential expression between CT and selection lines in the treatments and EC_{CT} when measured in CT (Figure 1). Finally, differences between plastic responses of CT- and selection lines (the interaction between condition and selection regime) give the evolutionary change in plasticity. To partition TC into changes explained by ancestral plasticity (PC_{CT}) and evolutionary changes (EC_T, see Figure 1A-C), we calculated the relative contribution of each component to the total. We used the log2-fold change of each transcript to evaluate and compare the magnitude of the plastic and evolutionary changes. When computing log2-fold changes, we accounted for non-independence among individuals from the same line by using the *duplicateCorrelation* function (Smyth et al. 2005) and added sequencing runs as batch effect. A gene is classified as differentially expressed (DE) with a FDR ≤ 5% after adjusting for multiple testing (Benjamini and Hochberg 1995).

In a second step, we used the normalized read counts (cpm, TMM-normalized) corrected for batch effects (sequencing runs) using the *removeBatchEffect* function in the *limma* R package (Ritchie et al., 2015). We used these counts to quantify expression levels in in CT and treatments.

This allowed us to conduct an analysis for each selection lines separately as was described by Ho

& Zhang (2018) (details see below).

Evolution of reaction norms

Reaching a new phenotypic optimum can be achieved by changing the mean (intercept of the

reaction norm) or by changing the plasticity of a trait (slope of reaction norm) (Figure 1D,E). We

were interested in assessing which of these patterns was more prevalent in gene expression

evolution. To test for an evolutionary change in plasticity (slope), we first evaluated the similarity

in plasticity between CT- and selection lines by calculating the correlation between PC_{CT} and PC_{Sel}

(Suppurting information Figure S2A) among all genes, independent of the significance. Second,

to test for an evolutionary change in mean expression (intercept), we calculated the correlation

between the evolved differences in CT (EC_{CT}) and in treatment conditions (EC_T). A significant

correlation indicates that the overall mean has changed (Figure S2A). Further, for all genes with

significant EC_T, we quantified how much of this difference could be explained by a shift in the

mean or by a change in plasticity. A shift in the mean corresponds to EC_{CT}, whereas the

remaining difference (EC_T-EC_{CT}) gives the evolutionary changes in plasticity (see also Stoks et al.

2016). Relative contributions of each component were obtained by dividing them by the total

EC_T.

To test for significant effects of the selection regime on differential expression (i.e., on number

of DE genes, log2-fold change, and correlations), we used a permutation test. We randomly

assigned samples and their transcriptomes to either CT selection or treatment (D, H, HD)

selection (number of samples for each selection was not changed) and repeated the DE analysis.

We kept the original assignment to lines and conditions and repeated the DE analysis for each

permuted data set. Observed values (e.g. number of DE genes, correlations) were considered

significant if higher than the most extreme 5% of the distribution calculated from permutations.

Comparing plastic and evolutionary responses

To infer the relationship between ancestral plasticity and evolution, we compared the direction

of PC_{CT} to the direction of EC_T. Evolution may *reinforce* the plastic response when PC_{CT} is in the

same direction as EC_T (Figure 1B). If EC_T is in opposite direction, it *reverses* PC_{CT} (Figure 1A). To

test which of these patterns was more prevalent, we followed Ho and Zhang 2018 (Ho & Zhang,

2018). Expression levels (cpm) of CT-Lines in CT conditions represented the original stage (*Lo*),

CT-Lines in treatment the plastic stage (*Lp*), and selection lines in their respective condition the

adaptive stage (*La*) (Figure 2). For subsequent analysis we used genes with appreciable PC_{CT} (|*Lp*

− *Lo*|> 0.2 *Lo*) and EC_T (|*La* − *Lp*|> 0.2 *Lo*) (Ho & Zhang, 2018) and calculated the proportion of

plastic genes with reinforced (TC > PC_{CT}) or reversed (TC < PC_{CT}) changes.

An alternative classification of plastic responses is to assess whether PC_{CT} brings expression

levels closer to the new optimum (*La).* In this case, the total change TC would be larger than EC_T

(Figure 2C). Such a pattern can occur even if PC_{CT} and EC_T are in opposite directions and classified

as reversion (Figure 2C), but may indicate that ancestral plasticity was beneficial. A PC_{CT} that

moves *Lp* further away from *La* is strongly indicative of maladaptive plasticity.

To confirm that our results were not sensitive to the applied cutoff (20% *Lo*), we repeated the

analysis with a cutoff of 50% *Lo* and without any cutoff (Supporting information, Figure S3.1 A-

H). Recently, it was pointed out (Mallard et al. 2018; Ho and Zhang 2019) that an excess of

reversions relative to reinforcements is expected to be observed due to a statistical artefact that

cannot be completely removed by permutation tests. We therefore applied an additional

parametric bootstrapping test as proposed by Ho and Zhang (Ho and Zhang 2019), to compare

proportion of reversions and reinforcements (Supporting information, Figure S3.1 J-L).

To better understand the relationship between the within-line proportions of reversed genes

and proportions of genes with *La* closer to *Lo* with adaptation, we calculated the Spearman

correlation between the proportions of reversed expression changes (or *La* closer to *Lo*

respectively) and mean offspring number in seven selection lines in HD. We focused on HD

because it was the most extreme environment with the strongest decline in offspring number.

To test for significance, we used permutations: Mean offspring numbers were randomly assigned

to lines and correlation was calculated again. Proportion of permutations with a correlation

coefficient exceeding the observed value gave the respective *P*-value.

Results

Fitness assay showed evolutionary adaptation

We found that selection lines had a higher offspring number in their native condition compared to CT-lines (Dry: $F_{1,14}$ = 9.20, *p* = 0.009; Hot: $F_{1,16}$ = 4.78, *p* = 0.044; Hot-Dry: $F_{1,16}$ = 23.51, *p* = 1.786E-04), confirming that adaptation had occurred (Figure 2). In contrast to treatment conditions, there was no difference in offspring number between CT- and selection lines under CT conditions (CT-lines: 64.89 [61.49, 68.29]; D-lines: 65.76 [61.26, 70.25], H-lines: 67.41 [63.76, 71.06], HD-lines: 65.56 [62.17, 68.94]) (Figure 2). Using three additional mixed models, we compared how lines from different selection regimes responded to treatments. We found significant negative effects for all stress treatments (D: $F_{1,23}$ = 45.37, *p* = 6.85E-07; H: $F_{1,28}$ = 507.68, *p* < 2.200E-16; HD: $F_{1,28}$ = 553.06, *p* < 2.200E-16) (Figure 2). Interaction between selection and treatments, i.e. whether the response to the treatment was different depending on selection regime, was significant for HD-lines ($F_{1,28}$ = 9.39, *p*= 4.754E-03) and for D ($F_{1,23}$ = 4.32, *p* = 0.049), but not for H ($F_{1,28}$ = 0.57, *p* = 0.455). Interaction between selection regime, treatment and line type (normal/mixed) was not significant in any treatment indicating that mixing lines did not have an effect on adaptation. ANOVA tables as well as results of the linear mixed models are in supporting information S4.

Evolution of gene expression

We found many more genes with expression divergence relative to Control (TC) than genes with evolved changes in the treatments (EC_T) or in control conditions (EC_{CT}) (Table 2). Increased temperature caused expression divergence in a larger number of genes than reduced humidity (Table 2). Most DE genes with significant TC also had significant ancestral plasticity PC_{CT} (Dry: 33.9%; Hot: 72.9%, Hot-Dry: 82.2%), but only a minority showed significant evolutionary change

EC_T (Dry: 4.9%; Hot: 1.1%; Hot-Dry: 1.0%, see also Supporting information Figure S3.2). About

half of the DE genes with EC_T in Hot and Hot-Dry then did not diverge significantly from their

ancestral expression levels in Control (TC ~ 0). Overall, TC had a larger log2-fold change

contribution from plastic changes (PC_{CT}) than evolved changes in Hot and Hot-Dry (median

proportions PC_{CT}/TC; Hot: 71.47%; Hot-Dry: 86.30%), but not in Dry (42.90%). Many genes had

PC_{CT} > TC, especially in HD (Figure 3A). In these cases, EC_T was negative and reduced TC.

Evolution of reaction norms

We were then interested to examine whether EC_T were due to a change in mean expression

(shift in the intercept of reaction norms) or to a change in plasticity (different slopes of reaction

norms, see Figure 3). First, we found a significant correlation between EC_{CT} and EC_T, in all three

treatments (permutation tests: $P < 0.0001$) (Table 3, Supporting information Figure S2). This

indicates that for a majority of genes, evolution in the past 20 generations affected their mean

expression levels by shifting the intercept of reaction norms (see Figure 1D). Second, we also

found a highly positive correlation between PC_{CT} and PC_{Sel} (Figure Supporting information S2A),

indicating that plasticity was mainly preserved during adaptation to new conditions. A

permutation test showed that this correlation was not significantly different from correlations

obtained from permuted datasets (Table 3), where samples were randomly assigned to selection

regimes, suggesting that the treatments did not have specific effects on plasticity. When we

quantified the relative contributions of changes in the mean versus changes in plasticity to EC_T,

we found that evolution of the intercept explained more evolutionary divergence than evolution

of the slope of reaction norms, especially in Dry (Figure 3B).

Evolution of plasticity in DE genes

We found only five genes with significantly evolved plasticity in the DE analysis, after correcting

for multiple testing. Although plasticity seemed to be preserved overall, we found evidence that

it evolved among DE genes. The DE analysis revealed a reduced number of plastic genes in selection lines compared to plastic genes in CT-lines (Table 2), although the difference was only significant in HD (Table 3). It further showed that ancestral plasticity PC_{CT} was larger than evolved plasticity PC_{Sel}, in D and HD (Table 3), indicating a reduction of the magnitude of plastic responses after adaptation. We found the same outcome when focusing on genes with significant EC_T, in HD (Mann-Whitney U test: U = 2133, *P* = 2.102E-04; median: CT-lines: 0.25, HD-lines: 0.13), but not in D (U = 144, *P* = 0.584; median: CT-lines: 0.10, D-lines: 0.14) or H (U = 386, *P* = 0.158; median: CT-lines: 0.21, H-lines: 0.12) (Figure 4).

Effect of plasticity on evolutionary responses

We could not find evidence that plasticity in general impeded evolution. In the DE analysis genes with a significant ancestral plastic response were overrepresented among the evolved DE genes in H and HD (H: χ2 = 4.51, df = 1, *P* = 3.38E-02; HD: χ2 = 9.71, df = 1, *P* = 1.84E-03), but not in D (*P* = 0.113 Fisher's exact test). Furthermore, when we analysed each selection line separately and compared their expression levels to CT-lines, the proportion of genes exhibiting substantial PC_{CT} (|*Lp* − *Lo*| > 0.2 *Lo*) were also overrepresented among evolved genes with EC_T (|*La* - *Lp*| > 0.2 *Lo*) (Table S5.1).

Maladaptive plastic changes may be reverted by evolutionary changes when the plastic response overshoots the optimum expression level in a new environment. Among the genes with significant EC_T in the DE analysis, we found that PC_{CT} was generally in opposite direction (Figure 4). Enrichment tests revealed that genes with positive EC_T in the DE analysis were overrepresented among genes with negative PC_{CT} in H and HD (Fisher's exact test D: *P* = 0.076; H: *P* = 0.002; HD: *P* = 9.26e-06), and genes with negative EC_T were overrepresented among genes with positive PC_{CT}, in H and in HD as well (Fisher's exact test D: *P* = 0.073; H: *P* = 0.021; HD: *P* =

0.049)(Figure 4). In almost all genes that showed significant EC_T and PC_{CT}, the responses were in

opposite direction (Dry: 2 out of 2 genes; Hot: 12 out of 13; Hot-Dry: 33 out of 34). We also

found negative correlations between PC_{CT} and EC_T in all treatments, although significant only in H

and HD, but not in D (Supporting information, Figure S3.3).

When we considered mean expression levels per line with $PC_{CT} > 20\%$ *Lo* and $EC_T > 20\%$ *Lo*, we

obtained similar results as in the DE analysis. Reinforcements were present but less frequent

than reversions in all treatments (proportions of plastic responses with reversion/reinforcement:

Dry: 48.5 ± 1.0% / 34.3 ± 0.8 %, $P < 1.151e-07$ (binomial test); Hot: 44.6 ± 0.9% / 28.8 ± 1.2%, $P =$

4.369E-13; Hot-Dry: 43.6 / 27.1 $P < 2.2e-16$; Figure 5A). The reversed genes represented 34.1 ±

0.8% (Dry), 40.0 ± 0.9% (Hot), and 42.96 ± 2.1% (Hot-Dry) of the evolved genes. In contrast, 24.1

± 0.6% (Dry), 25.8 ± 1.1% (Hot), 26.80 ± 1.6% (Hot-Dry) of evolved genes showed reinforcement.

Different cut-off values did not influence these outcomes (Figure S3.1). Parametric bootstrapping

as an additional test (Ho & Zhang, 2019), confirmed our results (Supporting information, Figure

S3.1). We also found that a large proportion of ancestrally plastic genes had *La* closer to *Lo* than

to *Lp* (Dry: 45.4 ± 1.1%; Hot: 41.2 ± 1.2%; Hot-Dry: 40.7 ± 2.5 %)(Figure 5B). Finally, substantial

proportions of genes with EC_T did not show PC_{CT} (Dry: 41.8 ± 0.7%, Hot: 34.6 ± 0.8%, Hot-Dry:

30.3 ± 1.0 %).

In case of genetic assimilation, i.e. loss of plasticity in selection line and continuous expression of

the induced phenotype, evolutionary differences would not be visible in treatment condition.

We calculated the mean expression levels of selection lines in Control and defined genes as

showing assimilation if they met three criteria: no PC_{Sel} (absolute difference between selection

lines in CT and in treatment < 20% *Lo*), $PC_{CT} > 20\%$ *Lo*, and *Lp* ~*La* (absolute difference < 20% *La*).

The proportion of PC_{CT} genes showing genetic assimilation was small in all conditions (Dry: 6.8 %,

Hot: 10.3 %, Hot-Dry: 10.3 %; Supporting information Table S5.2).

Proportions of reversed plastic responses and association with fitness

To gain a better understanding of the adaptive value of the changes of expression levels in the evolved lines, we tested for an association between within-line proportion of reversed or reinforced plastic responses and the average fitness of the lines in the HD treatment. We found that lines with a higher proportion of reversions had a higher average offspring number (correlation: 0.82, *P* = 0.012, Figure 6A) and we found a negative but non-significant correlation between fitness and reinforcements (correlation: -0.43, *P* = 0.85). When we tested for an association between fitness and proportion of ancestrally plastic genes with *La* closer to *Lo*, we also found a positive correlation (correlation: 0.86, *P* = 0.006, Figure 6B). Overall, better adapted lines (higher fitness in HD) showed a higher proportion of reversed ancestral plasticity and these plastic genes were more similar to the original expression levels of CT-lines in CT. Performing the analysis with gene expression data in H and D provided similar correlations, although not significant because of lower sample sizes (Supporting information, Table S5.3).

Discussion

We studied plastic and evolved responses in gene expression of *T. castaneum* in response to three new environmental conditions (Dry, Hot, Hot-Dry). After 20 generations of experimental evolution, we were able to detect adaptation and found significant evolutionary changes in expression levels. Comparing evolutionary changes with ancestral plastic responses showed that a reversion of plasticity was most frequent (> 95 % genes with significant plastic changes in DE analysis; > 40 % of genes with substantial plastic changes, i.e. changes higher than 20% of ancestral levels). The number of genes where ancestral plasticity was reinforced by evolution was significantly smaller (DE analysis: <5%, 27-34 % of genes with substantial plastic changes) and a smaller proportion of genes (7-10 % of substantially plastic genes) showed expression

patterns consistent with genetic assimilation. A high proportion of the originally plastic genes

evolved to expression levels that were closer to control levels than to ancestrally plastic levels.

Although the proportion of non-reversed plastic genes was still high, positive associations

between fitness and proportion of reversions, and compensated plasticity (expression levels

closer to control levels) respectively, suggest that ancestral plasticity was maladaptive for a

majority of responding genes. Although plasticity showed a high degree of preservation in terms

of number of responding genes and direction of the response, we found evidence that selection

lines evolved a reduced plasticity and thus partly compensated the maladaptive ancestral

response. We were further able to show a positive association between the proportion of

reversed plastic responses and adaptation (mean fitness per line) in the most stressful treatment

Hot-Dry.

Maladaptive plasticity

Different patterns describing the relationship between plastic and evolved changes in gene

expression have been documented. It was suggested that plasticity might help populations to

persist after environmental change, or to colonize new habitats by bringing phenotypes closer to

the new optimum. Studies found support for this hypothesis by showing that plastic responses of

non-adapted individuals diminished differences to native populations (Lohman, Stutz, & Bolnick,

2017; Mäkinen, Papakostas, Vøllestad, Leder, & Primmer, 2016). Adaptive plasticity can also be

indicated when plastic and evolutionary responses are in the same direction (Li, Li, Song, Wang,

& Zhang, 2017), or when plasticity is higher in adapted populations (Hasan et al., 2017; McCairns

& Bernatchez, 2009), suggesting that most plastic individuals were favored by selection.

However, there are also examples for the reversed pattern suggesting that plasticity was

maladaptive. In wild populations of *Fundulus heteroclitus*, evolved changes to different

temperatures were opposite to plastic responses of the ancestral population (Dayan et al.,

2015). *Rhagoletis* flies shifting to a new host fruits showed evolutionary responses opposite to

plasticity of non-adapted species (Ragland et al., 2015). Experimental evolution studies found

countergradient evolution in *Drosophila* adapting to different diets (Huang & Agrawal, 2016;

Yampolsky et al., 2012) and in guppies adapting to low predation environments (Ghalambor et

al., 2015). A comparative study (Ho & Zhang, 2018) analyzing data of multiple experimental

evolution suggested that reversions of gene expression changes might be a general pattern

during adaptation.

Our study fits with these previous observations. We found a higher proportion of reversions than

reinforcements in all conditions indicating mostly maladaptive plasticity. An alternative

explanation for the prevalence of reversions without maladaptive plastic responses would be

that CT-lines exhibited a response in the right direction, but overshot an optimum expression

level (Figure 1C). Fine-tuning during long-term adaptation could then lead to a partial reversion

of the plastic response. We took this possibility into account by not only focusing on a reversion

of plasticity, but also testing whether plastic changes brought expression levels closer to the

adapted stage *La* than to original stage *Lo.* If, in contrast, we see that *La* is closer to the original

level (CT lines in CT conditions), it indicates that plastic responses were maladaptive since they

moved expression levels further away from the new optimum and became compensated during

evolution (see Figure 1C). We found that adapted lines showed a high proportion of expression

levels closer to their ancestral level *Lo* than to *Lp*. We could further show that this proportion is

positively associated with higher fitness per selection line. We also found strong positive

associations of within-line proportions of reversions with mean reproductive output, indicating a

possible fitness advantage to reversions.

There was still a large proportion of plastic genes that did not show reversion. They are either

close to the levels of adapted lines, or showed reinforcement. The first case might indicate that

plasticity prevented evolution by matching the new optimum. Reinforced plastic changes could

be examples of adaptive plasticity. However, correlation between proportion of reinforcements

and fitness was not significant and negative. The observed positive correlation between

proportion of reversed plastic genes and mean fitness per line in HD rather suggests that

reversions were favoured during adaptations. Reversions might become more pronounced after

more generations, once evolution had sufficient time to further reverse maladaptive ancestral

plastic responses. Our selection lines still show a strong reduction in offspring number compared

to control levels suggesting further potential to adapt.

Evolution of reaction norms

The ancestral maladaptive plasticity can be compensated by shifts in the intercept or changes in

the slope of reaction norms. Both are not mutually exclusive and can occur together in the same

trait. We were aiming to quantify their relative importance for evolutionary responses in

transcriptomes.

Gene expression studies so far provided mixed results regarding the evolution of plasticity.

Drosophila populations adapted to different temperatures showed local adaptation, but there

was no evidence for evolution of thermal reaction norms of different transcripts and changes

affected mainly expression mean (Clemson, Sgrò, & Telonis-Scott, 2016). Experimental evolution

studying *Drosophila* under variable diets found no significant changes in plasticity (Yampolsky et

al., 2012) or less than expected (Huang & Agrawal, 2016). In contrast, other studies found

differences in temperature responses between tropical and temperate *Drosophila* populations

(Levine, Eckert, & Begun, 2011; von Heckel et al., 2016). Other examples for differences in genes

expression plasticity between adapted and non-adapted populations include temperature

(Morris et al., 2014) and salinity (Gibbons, Metzger, Healy, & Schulte, 2017; McCairns &

Bernatchez, 2009) responses of marine and freshwater sticklebacks, temperature response of

killifish populations from different latitudes, as well as plastic responses to toxic hydrogen

sulphide (H_2S) of fish population from H_2S rich springs versus non-toxic springs (Passow et al.,

2017). There was no consistent pattern regarding the direction in which plasticity evolves: In

some cases adapted population showed an increase in plasticity (Morris et al., 2014), in other

cases plasticity was reduced (Huang & Agrawal, 2016; Ragland et al., 2015; von Heckel et al.,

2016) or reduction and enhancement of plasticity were equally frequent (Gibbons et al., 2017;

Yampolsky et al., 2014). Overall, there is evidence in multiple species that expression plasticity of

some genes can evolve. However, even in some of these studies reporting evolved plasticity

(Dayan et al., 2015; Gibbons et al., 2017; Morris et al., 2014) the number of transcripts with

significant changes in the mean was much higher than transcripts with changed plasticity and

large parts of the plastic responses showed a high degree of preservation.

In accordance with these previous findings we found that changes in the mean contributed more

to the observed expression differences in the treatments than changes in plasticity. A possible

reason might be that genetic variation in mean expression was higher than genetic variation in

plasticity. In addition, we did not select for changes in plasticity directly since the conditions in

the treatments were constant. Selection was therefore on expression levels in the treatment and

only indirectly on plasticity. Plasticity could evolve if mean expression levels were genetically

correlated with plasticity. Although we did not directly test for such correlations, we found

evidence for evolution of plasticity of the DE genes, that is of the genes with significant changes

in expression level between CT and treatment conditions. Although plastic responses showed a

high degree of preservation in terms of affected genes and direction, we found evidence for

evolutionary changes in the magnitude of plastic responses, i.e. the slope of the reaction norm.

Why can plasticity be maladaptive?

New stressors might disturb homeostasis resulting in inappropriate responses and long-term adaptation therefore restores ancestral phenotypes by genetic changes, referred to as genetic compensation (Grether, 2005) or counter-gradient variation (Conover et al., 2009). However, in our study we applied relatively mild stressor treatments, i.e. individuals were able to survive and reproduce. Drought and heat are also stressors, which *T. castaneum* had experienced in the past (Sokoloff, 1972), so there had probably been selection on plastic responses to be beneficial. However, plastic responses might be optimized for a short-term exposure: Allocation of resources from reproduction to protection might increase survival probability and allow individuals to continue reproduction as soon as the stress has disappeared, but this response becomes maladaptive during continuous exposure and should therefore be under negative selection. Expression of stress related genes is in general accompanied by a down-regulation of genes involved in growth and reproduction due to an allocation of resources (Schwenke, Lazzaro, & Wolfner, 2016; Sokolova, 2013). A well-studied example are heat shock proteins (hsp). Hsp are well known for their protective function and to be crucial for survival (Feder & Hoffman, 1999), but it was also shown that their expression comes at a cost (Feder et al., 1998; Sørensen, Kristensen, & Loeschcke, 2003). Accordingly, it was often found that hsp expression in populations adapted to warmer climates is lower compared to non-adapted populations (Fangue, Hofmeister, & Schulte, 2006; Narum & Campbell, 2015; J G Sørensen, Dahlgaard, & Loeschcke, 2001). In general, other protection mechanisms independent of ancestral plasticity may arise during long-term adaptation (e.g. enzymes, which are more stable at high temperature) and make the costly stress response expendable.

An alternative explanation for the reduced plasticity in adapted lines is that the signal responsible for eliciting the plastic responses is based on any kind of damage (e.g. deformations

in macromolecules, membrane lipids, proteins, and DNA) caused by heat or stress in general

(Kültz, 2005). Higher resistance in adapted lines might shift the inducing thresholds, i.e. the

temperature when damages occur and stress response is induced (Sikkink, Reynolds, Ituarte,

Cresko, & Phillips, 2014) above the levels we applied in the treatments.

Interestingly, we found no differences in fitness between lines from different selection regimes

under CT conditions (Figure 2). We could detect some genes with different expression levels in

the treatments and found in general a correlated change in expression levels in control

conditions, but this did not affect offspring number. It indicates a lack of fitness trade-offs,

where alleles providing a fitness advantage in one environment (treatment) are detrimental in

another (CT). Together with the observation that selection lines evolved to bring expression

closer to ancestral CT-levels, it suggests that for many genes the optimal expression level is not

different between conditions. They might be involved in processes important for maintenance

and reproduction. Under stress, limited resources have to be invested into protection, that are

then not available for reproduction (Sokolova, 2013). Long-term adaptation should then work to

restore control levels that are likely to be optimized for highest reproductive output and to

reduce costly stress responses resulting in improved canalization of traits associated with fitness

(Stearns & Kawecki, 2006). Canalization, i.e. robustness against environmental variation, was

found previously in gene expression adaptation (Levine et al., 2011; Shaw et al., 2014; von

Heckel et al., 2016). Genetic differences between control and selection lines that are responsible

for adaptation to the treatments did not have an effect in control conditions and thus represent

cryptic genetic variation (Gibson & Dworkin, 2004). They might either concern genes that are not

expressed in control conditions, or represent changes neutral under control conditions.

Potential caveats

The number of genes with significant plastic changes in the DE analysis was much higher compared to genes showing evolutionary changes. One possible explanation would be that adaptive plasticity prevented evolution. If the plastic responses matched the optimum, no genetic changes in the selection lines are expected to occur. However, when we analysed each line separately and considered a gene as evolved if the mean difference between *La* and *Lp* was more than 20% of the *Lo*, we found approximately the same number of genes with EC_T and PC_{CT} (Supporting information Table S5.1).

In the DE analysis in *limma* we did not analyse each line separately but treated them as biological replicates. Since lines were split across conditions comparisons between conditions, i.e. plastic changes, can be made within lines. They should thus be more precise and statistical power should be higher than comparisons between selection regimes, i.e. evolved changes, that have to be made between lines. Differences between lines from the same selection regime lower the ability to obtain significant evolutionary changes. These differences can arise from genetic drift. Since our population size was relatively small (120 individuals per line) this might have been an important factor. Another explanation is that lines from the same selection regime differed how exactly they improved their fitness in the respective treatment. Since fitness is a highly polygenic trait, the genes contributing to a fitness increase may not be the same in different lines (see Barghi et al., 2019). For the most extreme treatment HD, where we sequenced seven lines, we further found considerable differences in fitness between the lines, suggesting that not all of them were at the same stage of adaptation. It is therefore not surprising that expression levels did not evolve in the same way among lines.

The DE analysis in *limma* requires that a gene shows similar changes in all replicate lines, and is therefore more conservative. If the main interest of a study is to identify promising candidate genes for future more detailed analyses it is the appropriate approach to keep FDR as low as

possible. In contrast, if the focus is more on general patterns, a less stringent analysis using mean

expression levels can give us a more complete picture. Since genetic drift is random it cannot

explain the observed excess of reversions over reinforcements.

Although we demonstrated that gene expression changed during evolution, it is not clear

whether these changes are the cause of an increased fitness in these conditions or whether they

are rather the consequence of adaptation and being less stressed. One disadvantage in studying

whole transcriptomes is that not all responding genes might be of functional importance but are

correlated to other adaptive changes. High intercorrelations within the transcriptome (Ayroles et

al., 2009; McGraw et al., 2011) might lead to correlated responses in many other genes.

Furthermore, observed evolutionary changes might be caused by indirect selection and other

mechanisms, e.g. changes in protein structure of enzymes, were responsible for adaptation of

selection lines. Future studies that manipulate expression and test for correlated changes in

offspring are needed to confirm adaptive value of expression changes.

Conclusions

We found that genes with the strongest plastic responses showed evolutionary changes in

opposite direction suggesting that ancestral plasticity was maladaptive for long-term adaptation.

In the most stressful treatment, selection lines with higher fitness show a higher proportion of

reversions and a higher proportion of originally plastic genes that are closer to ancestral

expression levels. Differences between adapted lines and CT-lines in the treatment were mainly

due to a change in mean expression (i.e. shift in the intercept of reaction norms), while plasticity

was preserved in terms of affected genes and direction of change. However, we found that a

part of the differences in the treatments can be explained by a reduction in the magnitude of

plasticity in adapted lines. Our results add to growing evidence that plasticity and evolution are

often in opposite direction and maladaptive plastic responses might increase strength of

selection. In contrast to previous studies, including fitness data allowed us to give evidence for

adaptation and in the most stressful condition we were able to show an association between

reversion of plasticity and adaptation. Similar results in all three stress treatments indicated that

these findings may represent a general pattern of gene expression adaptation.

Data Accessibility

RNA-seq reads will be available at GEO.

Fitness data will be available at Dryad.

Author Contribution

FG and ELK designed experiment. ELK conducted experiment, laboratory work and analysed the

data. FG and ELK wrote the manuscript.

Acknowledgements

This work was supported by the Swiss National Science Foundation, grants PP00P3_1144846 and

PP00P3_176965 to FG. We thank Sonja Sbilordo, Sara Meier, Cilgia Lippuner, Tim Schoch and Tim

Emmenegger for helping during the fitness assay, Elke Karaus, Martina Berchtold, Encarnación

Lozano for RNA extraction and quality control, Lucy Poveda and Maria Domenica Moccia for

advice in library preparation and sequencing, Lennart Opitz for helping to process the RNA-seq

data.

Figure 1: Possible relations between plastic change PC and evolutionary change EC. Gene expression levels of control lines (white) in the ancestral control condition represent the original stage Lo, expression in the treatment the plastic stage Lp. Expression levels of selection lines (grey) in the treatment give the adapted stage La Arrows indicate the direction of PC (dashed line) and EC (solid line). PC can be in opposite direction to EC (reversion **(A))** or it can be in the same direction (reinforcement **(B))**. However, even if PC and EC are opposite to each other, PC can bring expression levels closer to levels of the adapted lines. In this case, the total change TC, (difference between expression levels of control lines in control conditions and selection lines in treatment) is larger than EC **(C)**. During adaptation, lines could have reached the optimum by either changing mean expression, i.e. shift in the intercept of the reaction norm **(D)** or by changing their plasticity, i.e. the slope of the reaction norm **(E)**. In case of a change in the mean, plastic changes of Control lines PC_{CT} and selection lines PC_{Sel} as well as observed evolutionary change EC_T in treatment and in Control EC_{CT} would be highly correlated because reaction norms (red arrows) remain parallel **(D)**. If observed EC_T in treatment is due to a change in plasticity only, no correlation between EC_T and EC_{CT} should exist, and the correlation between PC_{Sel} and PC_{CT} should be small **(E)**.

Figure 2: Offspring number of Control and selection lines under different conditions. Selection lines could adapt for 20 generations.

645 **Figure 3:** Relative contribution of ancestral plasticity and evolutionary change to total divergence for all genes showing
646 significant total divergence in the DE analysis. Number of genes: Dry: 283, Hot: 1192, Hot-Dry: 2045 **(A)**. Relative
647 contribution of changes in the mean (shift in the intercept of reaction norm) and changes in plasticity (different slopes
648 of reaction norms) to evolutionary differences between control-lines and selection lines in the treatment. Only genes
649 with significant differences in the DE analysis were included. Number of genes: Dry: 18, Hot: 25, Hot-Dry: 55 **(B)**.

Figure 4: Plastic responses in control-lines and adapted selection-lines of genes showing significant evolutionary changes in expression in the treatments Dry **(A)**, Hot **(B)** and Hot-Dry **(C)**. Plastic responses in adapted selection lines are weaker (smaller logFC). Furthermore, plastic changes are mainly opposite to evolved changes. Genes that evolved to lower expression in the treatments (represented in lightgrey) show positive plastic changes in control-lines, i.e. they are up-regulated. Genes that evolved to higher expression levels (in darkgrey) are down-regulated in the non-adapted Control-Lines.

Figure 5: Comparison of plastic and evolutionary changes in gene expression in response to three environmental
conditions. **A:** Proportion of genes (average over selection lines) showing a reversion or reinforcement of ancestral
plasticity during evolution. **B:** Proportion of genes with expression levels after evolution (adapted stage La) closer to
original levels (Lo, i.e. Control-lines in control conditions), or closer to plastic levels (Lp, i.e. expression levels of
Control-lines in treatment). Only genes exhibiting substantial plastic changes ($|Lp-Lo| > 20\%$ of Lo) as well as
evolutionary changes ($La-Lp| > 20\%$ of Lo) were used for analysis.

Figure 6: Relationship between plastic and evolved changes in gene expression in response to hot-dry conditions.
Expression levels of Control-lines in control conditions represent the ancestral stage Lo, Control-Lines in hot-dry
conditions the plastic stage Lp and adapted Hot-Dry-lines in hot-dry conditions the adapted stage La. **A:** Relationship
between the proportion of reversed plastic responses and mean fitness (=offspring number) per Hot-Dry line. **B:**
Relationship between proportion of genes with La closer to Lo and mean fitness. Only genes exhibiting substantial
plastic changes (|Lp-Lo| > 20% of Lo) as well as evolutionary changes (La-Lp| > 20% of Lo) were used for analysis. P-
values of the spearman correlations were obtained by 10,000 permutations. 95 % Confidence intervals are based on a
non-parametric bootstrap test.

683 **Table 1:** Number of sequenced replicate lines and individuals per selection and treatment, which were used for this
684 study. Selection lines could adapt to conditions for 20 generations. Control (CT) conditions: 33°C, 70% relative
685 humidity (r.h.); treatments: Dry (D): 33°C, 30% r.h.; Hot (H): 37°C, 70% r.h.; Hot-Dry (HD): 37°C, 30% r.h.

Selection	Conditions				
	Total	CT	D	H	HD
CT					
Number lines	**7**	7	5	5	7
Number Ind.	**89**	25	20	19	26
D					
Number lines	**4**	4	4	–	–
Number Ind.	**32**	16	16	–	–
H					
Number lines	**5**	5	–	5	–
Number Ind.	**40**	20	–	20	–
HD					
Number lines	**7**	7	–	–	7

	Number Ind.	**56**	28	–	–	28

Table 2: Significantly differently expressed genes. For evolved differences, control-lines and selection-lines are compared within condition (e.g. Dry-lines vs Control-lines in Dry). Plastic response gives the number of genes that changed expression between control and treatment conditions (e.g. Dry-lines in Dry vs Dry-lines in control). Different plasticity gives the number of genes with significant different plastic responses in lines from different selection regimes. Analysis was conducted using the R package limma (Ritchie et al., 2015).

		Dry	**Hot**	**Hot-Dry**
Evolved difference in Control	down	3	0	6
	up	1	9	4
	total	**4**	**9**	**10**
Evolved difference in treatment	down	9	8	22
	up	9	17	33
	total	**18**	**25**	**55**
Plastic response Control-Lines	down	209	1765	2451
	up	156	1714	2200
	total	**365**	**3479**	**4651**
Plastic response Selection-Lines	down	28	1417	1649
	up	21	1381	1470
	total	**49**	**2798**	**3119**
Total change (Selection-lines in treatment vs Control-lines in Control)	down	154	564	1016
	up	129	628	1029
	total	**283**	**1192**	**2045**
different plasticity	total	**0**	**1**	**4**

Table 3: P-values obtained from permutation tests (10,000 permutations). Samples were randomly assigned to either control or treatment selection and differential expression analyses were repeated. Significance was assessed by calculating the proportion of permutations with more extreme values than the observed one. Control (CT) conditions: 33°C, 70% relative humidity r.h. Conditions in treatments: Dry (D): 33°C, 30% r.h.; Hot (H): 37°C, 70% r.h.; Hot-Dry (HD): 37°C, 30% r.h.

	Selection		
	D	**H**	**HD**
CT-Lines have more genes with significant plastic responses compared to adapted selection lines	0.2981	0.3037	**0.0325**
Magnitude of plastic response is higher in CT-Lines	**0.0493**	0.1676	**0.0491**
Individuals from different selection regimes show lower similarity in plastic responses [†]	0.8608	0.3914	0.5217

Number of genes with significant differences in expression levels is higher in treatment than in CT conditions	**0.0095**	**0.0003**	**0.0031**
Differences in expression levels in CT conditions and treatment are correlated	**< 0.0001**	**<0.0001**	**< 0.0001**

†: observed correlation of plastic responses of CT- and selection lines would be lower compared to results of permutations

References

Alvarez, M., Schrey, A. W., & Richards, C. L. (2015). Ten years of transcriptomics in wild populations: What have we learned about their ecology and evolution? *Molecular Ecology*, *24*(4), 710–725. doi:10.1111/mec.13055

Ayroles, J. F., Carbone, M. A., Stone, E. A., Jordan, K. W., Lyman, R. F., Magwire, M. M., … Mackay, T. F. C. (2009). Systems genetics of complex traits in *Drosophila melanogaster*. *Nature Genetics*, *41*(3), 299–307. doi:10.1038/ng.332

Barghi, N., Tobler, R., Nolte, V., Jakšić, A. M., Mallard, F., Otte, K. A., … Schlötterer, C. (2019). *Genetic redundancy fuels polygenic adaptation in Drosophila*. PLoS Biology (Vol. 17). doi:10.1371/journal.pbio.3000128

Barton, M., Sunnucks, P., Norgate, M., Murray, N., & Kearney, M. (2014). Co-gradient variation in growth rate and development time of a broadly distributed butterfly. *PLoS ONE*, *9*(4), 1–8. doi:10.1371/journal.pone.0095258

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using **lme4**. *Journal of Statistical Software*, *67*(1), 1–48. doi:10.18637/jss.v067.i01

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, 289–300.

Carroll, S. B. (2005). Evolution at Two Levels: On Genes and Form. *PLoS Biology*, *3*(7), e245. doi:10.1371/journal.pbio.0030245

Chevin, L. M., Lande, R., & Mace, G. M. (2010). Adaptation, plasticity, and extinction in a changing environment: Towards a predictive theory. *PLoS Biology*, *8*(4). doi:10.1371/journal.pbio.1000357

Clemson, A. S., Sgrò, C. M., & Telonis-Scott, M. (2016). Thermal plasticity in Drosophila melanogaster populations from eastern Australia: quantitative traits to transcripts. *Journal of Evolutionary Biology*, *29*(12), 2447–2463. doi:10.1111/jeb.12969

Conover, D. O., Duffy, T. A., & Hice, L. A. (2009). The covariance between genetic and environmental influences across ecological gradients: Reassessing the evolutionary significance of countergradient and cogradient variation. *Annals of the New York Academy of Sciences*, *1168*(1), 100–129. doi:10.1111/j.1749-6632.2009.04575.x

Crispo, E. (2007). The Baldwin effect and genetic assimilation: Revisiting two mechanisms of evolutionary change mediated by phenotypic plasticity. *Evolution*, *61*(11), 2469–2479. doi:10.1111/j.1558-5646.2007.00203.x

Dayan, D. I., Crawford, D. L., & Oleksiak, M. F. (2015). Phenotypic plasticity in gene expression contributes to divergence of locally adapted populations of *Fundulus heteroclitus*. *Molecular Ecology*, *24*(13), 3345–3359. doi:10.1111/mec.13188

de Jong, G. (2005). Evolution of phenotypic plasticity: patterns of plasticity and the emergence of ecotypes. *New Phytologist*, *166*(2003), 101–118. doi:10.1111/j.1469-8137.2005.01322.x

de Jong, Gerdien. (2005). Evolution of phenotypic plasticity: patterns of plasticity and the emergence of ecotypes. *New Phytologist*, *166*(1), 101–118. doi:10.1111/j.1469-8137.2005.01322.x

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., … Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.

Draghi, J. A., & Whitlock, M. C. (2012). Phenotypic plasticity facilitates mutational variance, genetic variance, and evolvability along the major axis of environmental variation. *Evolution*, *66*(9), 2891–2902. doi:10.1111/j.1558-5646.2012.01649.x

Fangue, N. A., Hofmeister, M., & Schulte, P. M. (2006). Intraspecific variation in thermal tolerance and heat shock protein gene expression in common killifish, *Fundulus heteroclitus*. *The Journal of Experimental Biology*, *209*(Pt 15), 2859–72. doi:10.1242/jeb.02260

Feder, M. E., & Hoffman, G. E. (1999). Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annual Review of Physiology*, *61*, 243–282.

Feder, M. E., & Krebs, R. A. (1998). Natural and Genetic Engineering of the Heat-Shock Protein Hsp70 in Drosophila melanogaster : Consequences for Thermotolerance. *American Zoologist*, *38*(3), 503–517. doi:10.1093/icb/38.3.503

Fierst, J. L. (2011). A history of phenotypic plasticity accelerates adaptation to a new environment. *Journal of Evolutionary Biology*, *24*(9), 1992–2001. doi:10.1111/j.1420-9101.2011.02333.x

Fitzpatrick, B. M. (2012). Underappreciated consequences of phenotypic plasticity for ecological speciation. *International Journal of Ecology*, *2012*, 32–37. doi:10.1155/2012/256017

Forsman, A. (2015). Rethinking phenotypic plasticity and its consequences for individuals, populations and species. *Heredity*, *115*(4), 276–284. doi:10.1038/hdy.2014.92

Garland, T., & Kelly, S. A. (2008). Phenotypic plasticity and experimental evolution. *Journal of Experimental Biology*, *211*(16), 2725–2725. doi:10.1242/jeb.022673

Ghalambor, C. K., Hoke, K. L., Ruell, E. W., Fischer, E. K., Reznick, D. N., & Hughes, K. A. (2015). Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature*, *525*(7569), 372–375. doi:10.1038/nature15256

Ghalambor, C. K., McKay, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology*, *21*(3), 394–407. doi:10.1111/j.1365-2435.2007.01283.x

Gibbons, T. C., Metzger, D. C. H., Healy, T. M., & Schulte, P. M. (2017). Gene expression plasticity in response to salinity acclimation in threespine stickleback ecotypes from different salinity habitats. *Molecular Ecology*, *26*(10), 2711–2725. doi:10.1111/mec.14065

Gibson, G. (2008). The environmental contribution to gene expression profiles. *Nature Reviews Genetics*, *9*(8), 575.

Gibson, G., & Dworkin, I. (2004). Uncovering cryptic genetic variation. *Nature Reviews Genetics*, *5*(9), 681–690. doi:10.1038/nrg1426

Grether, G. F. (2005). Environmental Change, Phenotypic Plasticity, and Genetic Compensation. *The American Naturalist*, *166*(4), E115–E123. doi:10.1086/432023

Guo, J., Liu, R., Huang, L., Zheng, X. M., Liu, P. L., Du, Y. S., … Ge, S. (2016). Widespread and adaptive alterations in genome-wide gene expression associated with ecological divergence of two oryza species. *Molecular Biology and Evolution*, *33*(1), 62–78. doi:10.1093/molbev/msv196

Hasan, M. M., DeFaveri, J., Kuure, S., Dash, S. N., Lehtonen, S., Merilä, J., & McCairns, R. J. S. (2017). Sticklebacks adapted to divergent osmotic environments show differences in plasticity for kidney morphology and candidate gene expression. *The Journal of*

Experimental Biology, *220*(12), 2175–2186. doi:10.1242/jeb.146027

Hatakeyama, M., Opitz, L., Russo, G., Qi, W., Schlapbach, R., & Rehrauer, H. (2016). SUSHI: An
exquisite recipe for fully documented, reproducible and reusable NGS data analysis. *BMC*
Bioinformatics, *17*(1), 1–9. doi:10.1186/s12859-016-1104-8

Ho, W.-C., & Zhang, J. (2018). Evolutionary adaptations to new environments generally reverse
plastic phenotypic changes. *Nature Communications*, *9*(1), 1–11. doi:10.1038/s41467-017-
02724-5

Ho, W.-C., & Zhang, J. (2019). Genetic gene expression changes during environmental
adaptations tend to reverse plastic changes even after the correction for statistical
nonindependence. *Molecular Biology and Evolution*. doi:10.1093/molbev/msz002

Hodgins-Davis, A., & Townsend, J. P. (2009). Evolving gene expression: from G to E to G × E.
Trends in Ecology and Evolution, *24*(12), 649–658. doi:10.1016/j.tree.2009.06.011

Huang, Y., & Agrawal, A. F. (2016). Experimental Evolution of Gene Expression and Plasticity in
Alternative Selective Regimes. *PLoS Genetics*, *12*(9), 1–23.
doi:10.1371/journal.pgen.1006336

Krebs, R. A., & Feder, M. E. (1997). Natural Variation in the Expression of the Heat-Shock Protein
HSP70 in a Population of Drosophila melanogaster and Its Correlation with Tolerance of
Ecologically Relevant Thermal Stress. *Evolution*, *51*(1), 173. doi:10.2307/2410970

Kregel, K. C. (2002). Invited Review: Heat shock proteins: modifying factors in physiological stress
responses and acquired thermotolerance. *Journal of Applied Physiology*, *92*(5), 2177–2186.
doi:10.1152/japplphysiol.01267.2001

Kültz, D. (2005). Molecular and Evolutionary Basis of the Cellular Stress Response. *Annual Review*
of Physiology, *67*(1), 225–257. doi:10.1146/annurev.physiol.67.040403.103635

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear
Mixed Effects Models. *Journal of Statistical Software*, *82*(13), 1–26.
doi:10.18637/jss.v082.i13

Lande, R. (2009). Adaptation to an extraordinary environment by evolution of phenotypic
plasticity and genetic assimilation. *Journal of Evolutionary Biology*, *22*(7), 1435–1446.
doi:10.1111/j.1420-9101.2009.01754.x

Laugen, A. T., Laurila, A., Räsänen, K., & Merilä, J. (2003). Latitudinal countergradient variation in
the common frog (Rana temporaria) development rates – evidence for local adaptation. *J.*
Evol., *16*(5), 996–1005.

Law, C. W., Chen, Y., Shi, W., & Smyth, G. K. (2014). voom: precision weights unlock linear model
analysis tools for RNA-seq read counts. *Genome Biology*, *15*(2), R29. doi:10.1186/gb-2014-
15-2-r29

Lenth, R. V. (2016). Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software*,
69(1), 1–33. doi:10.18637/jss.v069.i01

Levine, M. T., Eckert, M. L., & Begun, D. J. (2011). Whole-genome expression plasticity across
tropical and temperate *Drosophila melanogaster* populations from eastern Australia.
Molecular Biology and Evolution, *28*(1), 249–256. doi:10.1093/molbev/msq197

Levis, N. A., & Pfennig, D. W. (2016). Evaluating ' Plasticity-First ' Evolution in Nature : Key
Criteria and Empirical Approaches. *Trends in Ecology & Evolution*, *31*(7), 563–574.
doi:10.1016/j.tree.2016.03.012

Li, A., Li, L., Song, K., Wang, W., & Zhang, G. (2017). Temperature, energy metabolism, and
adaptive divergence in two oyster subspecies. *Ecology and Evolution*, *7*(16), 6151–6162.
doi:10.1002/ece3.3085

Liao, Y., Smyth, G. K., & Shi, W. (2014). FeatureCounts: An efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–930.
doi:10.1093/bioinformatics/btt656

Loeschcke, V., & Hoffmann, A. a. (2007). Consequences of heat hardening on a field fitness
component in Drosophila depend on environmental temperature. *American Naturalist*,
169(2), 175–183. doi:10.1086/510632
Lohman, B. K., Stutz, W. E., & Bolnick, D. I. (2017). Gene expression stasis and plasticity following
migration into a foreign environment. *Molecular Ecology*, *26*(18), 4657–4670.
doi:10.1111/mec.14234
Mahroof, R., Zhu, K. Y., & Subramanyam, B. (2005). Changes in Expression of Heat Shock Proteins
in Tribolium castaneum (Coleoptera: Tenebrionidae) in Relation to Developmental Stage,
Exposure Time, and Temperature. *Annals of the Entomological Society of America*, *98*(1),
100–107. doi:10.1603/0013-8746(2005)098[0100:CIEOHS]2.0.CO;2
Mäkinen, H., Papakostas, S., Vøllestad, L. A., Leder, E. H., & Primmer, C. R. (2016). Plastic and
evolutionary gene expression responses are correlated in European grayling (*Thymallus*
thymallus) subpopulations adapted to different thermal environments. In *Journal of*
Heredity (Vol. 107, pp. 82–89). doi:10.1093/jhered/esv069
Mallard, F., Jakšić, A. M., & Schlötterer, C. (2018). Contesting the evidence for non-adaptive
plasticity. *Nature*, *555*(7698), E21.
McCairns, R. J. S. S., & Bernatchez, L. (2009). Adaptive divergence between freshwater and
marine sticklebacks: insights into the role of phenotypic plasticity from an integrated
analysis of candidate gene expression. *Evolution*, *64*(4), 1029–1047. doi:10.1111/j.1558-
5646.2009.00886.x
McGraw, E. A., Ye, Y. H., Foley, B., Chenoweth, S. F., Higgie, M., Hine, E., & Blows, M. W. (2011).
High-dimensional variance partitioning reveals the modular genetic basis of adaptive
divergence in gene expression during reproductive character displacement. *Evolution*,
65(11), 3126–3137. doi:10.1111/j.1558-5646.2011.01371.x
Milutinović, B., Stolpe, C., Peuß, R., Armitage, S. A. O., & Kurtz, J. (2013). The Red Flour Beetle as
a Model for Bacterial Oral Infections. *PLoS ONE*, *8*(5). doi:10.1371/journal.pone.0064638
Molina-Montenegro, M. A., Peñuelas, J., Munné-Bosch, S., & Sardans, J. (2012). Higher plasticity
in ecophysiological traits enhances the performance and invasion success of Taraxacum
officinale (dandelion) in alpine environments. *Biological Invasions*, *14*(1), 21–33.
doi:10.1007/s10530-011-0055-2
Morris, M. R. J., Richard, R., Leder, E. H., Barrett, R. D. H., Aubin-Horth, N., & Rogers, S. M.
(2014). Gene expression plasticity evolves in response to colonization of freshwater lakes in
threespine stickleback. *Molecular Ecology*, *23*(13), 3226–3240. doi:10.1111/mec.12820
Narum, S. R., & Campbell, N. R. (2015). Transcriptomic response to heat stress among
ecologically divergent populations of redband trout. *BMC Genomics*, *16*(1), 103.
doi:10.1186/s12864-015-1246-5
Nussey, D. H., Postma, E., Gienapp, P., & Visser, M. E. (2005). Selection on heritable phenotypic
plasticity in a wild bird population. *Science*, *310*(5746), 304–306.
doi:10.1126/science.1117004
Passow, C. N., Henpita, C., Shaw, J. H., Quackenbush, C. R., Warren, W. C., Schartl, M., … Tobler,
M. (2017). The roles of plasticity and evolutionary change in shaping gene expression
variation in natural populations of extremophile fish. *Molecular Ecology*, *26*(22), 6384–
6399. doi:10.1111/mec.14360
Pavey, S. A., Collin, H., Nosil, P., & Rogers, S. M. (2010). The role of gene expression in ecological
speciation. *Annals of the New York Academy of Sciences*, *1206*, 110–129.
doi:10.1111/j.1749-6632.2010.05765.x
Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., & Moczek, A. P.
(2010). Phenotypic plasticity's impacts on diversification and speciation. *Trends in Ecology*
and Evolution, *25*(8), 459–467. doi:10.1016/j.tree.2010.05.006

Pichancourt, J. B., & van Klinken, R. D. (2012). Phenotypic plasticity influences the size, shape and
dynamics of the geographic distribution of an invasive plant. *PLoS ONE*, *7*(2), e32323.
doi:10.1371/journal.pone.0032323
Pigliucci, M., Murren, C. J., & Schlichting, C. D. (2006). Phenotypic plasticity and evolution by
genetic assimilation. *Journal of Experimental Biology*, *209*(12), 2362–2367.
doi:10.1242/jeb.02070
Pigliucci, Massimo. (2005). Evolution of phenotypic plasticity: Where are we going now? *Trends*
in Ecology and Evolution, *20*(9), 481–486. doi:10.1016/j.tree.2005.06.001
Price, T. D., Qvarnstrom, A., & Irwin, D. E. (2003). The role of phenotypic plasticity in driving
genetic evolution. *Proceedings of the Royal Society B: Biological Sciences*, *270*(1523), 1433–
1440. doi:10.1098/rspb.2003.2372
Price, Trevor D., Qvarnström, A., & Irwin, D. E. (2003). The role of phenotypic plasticity in driving
genetic evolution. *Proceedings of the Royal Society B: Biological Sciences*, *270*(1523), 1433–
1440. doi:10.1098/rspb.2003.2372
R Core Team. (2017). R Core Team (2017). R: A language and environment for statistical
computing. *R Foundation for Statistical Computing, Vienna, Austria. URL Http://Www.R-*
Project.Org/.
Ragland, G. J., Almskaar, K., Vertacnik, K. L., Gough, H. M., Feder, J. L., Hahn, D. A., & Schwarz, D.
(2015). Differences in performance and transcriptome-wide gene expression associated
with Rhagoletis (Diptera: Tephritidae) larvae feeding in alternate host fruit environments.
Molecular Ecology, *24*(11), 2759–2776. doi:10.1111/mec.13191
Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic*
Acids Research, *43*(7), e47–e47. doi:10.1093/nar/gkv007
Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics*, *26*(1), 139–
140. doi:10.1093/bioinformatics/btp616
Romero, I. G., Ruvinsky, I., & Gilad, Y. (2012). Comparative studies of gene expression and the
evolution of gene regulation. *Nature Reviews Genetics*, *13*(7), 505–516.
doi:10.1038/nrg3229
Schaum, E., Rost, B., Millar, A. J., & Collins, S. (2013). Variation in plastic responses of a globally
distributed picoplankton species to ocean acidification. *Nature Climate Change*, *3*(3), 298–
302. doi:10.1038/nclimate1774
Scheiner, S. M. (1993). Genetics and Evolution of Phenotypic Plasticity. *Annual Review of Ecology*
and Systematics, *24*(1), 35–68. doi:10.1146/annurev.es.24.110193.000343
Schwenke, R. A., Lazzaro, B. P., & Wolfner, M. F. (2016). Reproduction–Immunity Trade-Offs in
Insects. *Annual Review of Entomology*, *61*(1), 239–256. doi:10.1146/annurev-ento-010715-
023924
Shaw, J. R., Hampton, T. H., King, B. L., Whitehead, A., Galvez, F., Gross, R. H., … Stanton, B. A.
(2014). Natural Selection Canalizes Expression Variation of Environmentally Induced
Plasticity-Enabling Genes. *Molecular Biology and Evolution*, *31*(11), 3002–3015.
doi:10.1093/molbev/msu241
Sikkink, K. L., Reynolds, R. M., Ituarte, C. M., Cresko, W. A., & Phillips, P. C. (2014). Rapid
Evolution of Phenotypic Plasticity and Shifting Thresholds of Genetic Assimilation in the
Nematode Caenorhabditis remanei. *G3: Genes|Genomes|Genetics*, *4*(6), 1103–1112.
doi:10.1534/g3.114.010553
Silbermann, R., & Tatar, M. (2000). Reproductive costs of heat shock protein in transgenic
Drosophila melanogaster. *Evolution*, *54*(6), 2038–2045. doi:10.1111/j.0014-
3820.2000.tb01247.x

Smyth, G. K., Michaud, J., & Scott, H. S. (2005). Use of within-array replicate spots for assessing
differential expression in microarray experiments. *Bioinformatics*, *21*(9), 2067–2075.
doi:10.1093/bioinformatics/bti270
Sokoloff, A. (1972). *The biology of Tribolium with special emphasis on genetic aspects I.*
Clarendon Press and Oxford Univ. Press, Oxford. doi:10.1086/407731
Sokolova, I. M. (2013). Energy-limited tolerance to stress as a conceptual framework to integrate
the effects of multiple stressors. *Integrative and Comparative Biology*, *53*(4), 597–608.
doi:10.1093/icb/ict028
Sørensen, J G, Dahlgaard, J., & Loeschcke, V. (2001). Genetic variation in thermal tolerance
among natural populations of *Drosophila buzzatii*: down regulation of Hsp70 expression
and variation in heat stress resistance traits. *Functional Ecology*, *15*(3), 289–296.
doi:10.1046/j.1365-2435.2001.00525.x
Sørensen, Jesper Givskov, Kristensen, T. N., & Loeschcke, V. (2003). The evolutionary and
ecological role of heat shock proteins. *Ecology Letters*, *6*(11), 1025–1037.
doi:10.1046/j.1461-0248.2003.00528.x
Stearns, S. C., & Kawecki, T. J. (2006). Fitness Sensitivity and the Canalization of Life-History
Traits. *Evolution*, *48*(5), 1438. doi:10.2307/2410238
Stoks, R., Govaert, L., Pauwels, K., Jansen, B., & De Meester, L. (2016). Resurrecting complexity:
The interplay of plasticity and rapid evolution in the multiple trait response to strong
changes in predation pressure in the water flea Daphnia magna. *Ecology Letters*, *19*(2),
180–190. doi:10.1111/ele.12551
Townsend, J. P., Cavalieri, D., & Hartl, D. L. (2003). Population genetic variation in genome-wide
gene expression. *Molecular Biology and Evolution*, *20*(6), 955–963.
doi:10.1093/molbev/msg106
Via, S. (1993). Adaptive Phenotypic Plasticity: Target or By-Product of Selection in a Variable
Environment? *The American Naturalist*, *142*(2), 352–365. doi:10.1086/285542
von Heckel, K., Stephan, W., & Hutter, S. (2016). Canalization of gene expression is a major
signature of regulatory cold adaptation in temperate Drosophila melanogaster. *BMC
Genomics*, *17*(1), 1–14. doi:10.1186/s12864-016-2866-0
Whitehead, A., & Crawford, D. L. (2006). Neutral and adaptive variation in gene expression.
Proceedings of the National Academy of Sciences of the United States of America\r, *103*(14),
5425–5430. doi:10.1073/pnas.0507648103
Wray, G. A. (2007). The evolutionary significance of cis-regulatory mutations. *Nature Reviews
Genetics*, *8*(3), 206–216. doi:10.1038/nrg2063
Wund, M. A. (2012). Assessing the impacts of phenotypic plasticity on evolution. *Integrative and
Comparative Biology*, *52*(1), 5–15. doi:10.1093/icb/ics050
Yampolsky, L. Y., Glazko, G. V., & Fry, J. D. (2012). Evolution of gene expression and expression
plasticity in long-term experimental populations of *Drosophila melanogaster* maintained
under constant and variable ethanol stress. *Molecular Ecology*, *21*(17), 4287–4299.
doi:10.1111/j.1365-294X.2012.05697.x
Yampolsky, L. Y., Zeng, E., Lopez, J., Williams, P. J., Dick, K. B., Colbourne, J. K., & Pfrender, M. E.
(2014). Functional genomics of acclimation and adaptation in response to thermal stress in
Daphnia. *BMC Genomics*, *15*(1), 1–12. doi:10.1186/1471-2164-15-859
Yeh, P. J., & Price, T. D. (2004). Adaptive Phenotypic Plasticity and the Successful Colonization of
a Novel Environment. *The American Naturalist*, *164*(4), 531–542. doi:10.1086/423825
Zhao, L., & Jones, W. A. (2012). Expression of heat shock protein genes in insect stress responses.
Invertebrate Survival Journal, *9*(1), 93–101.
983