Restoring ancestral phenotypes by reduction of plasticity is a general pattern in gene expression during adaptation to different stressors in *Tribolium castaneum*

Eva L. Koch1 and Frédéric Guillaume1

1 Department of Evolutionary Biology and Environmental Studies, University of Zürich,

Winterthurerstr. 190, 8057 Zürich, Switzerland

Abstract

Plasticity and evolution are two processes enabling individuals to respond to environmental change, but how both are related and impact each other is still controversial. We studied plastic and evolutionary responses in gene expression of *Tribolium castaneum* after exposure to new environments that differed from ancestral conditions in temperature, humidity or both. Using experimental evolution with ten replicated lines per condition, we were able to demonstrate adaptation after 20 generations. We measured gene expression in each condition in adapted selection lines and control lines to infer evolutionary and plastic changes. We found more evidence for changes in mean expression (shift in the intercept of reaction norms) in adapted lines than for changes in plasticity (shifts in slopes). Plasticity was mainly preserved and was responsible for a large part of the phenotypic divergence in expression between ancestral and new conditions. However, we found that genes with the largest evolutionary changes in expression also evolved reduced plasticity and often showed expression levels closer to the ancestral stage. Results obtained in the three different conditions were similar suggesting that restoration of ancestral expression levels during adaptation is a general evolutionary pattern. We increased the sample size in the most stressful condition and were then able to detect a positive correlation between proportion of genes with reversion of the ancestral plastic response and mean fitness per selection line.

Introduction

Whenever facing environmental change, populations can adapt to new phenotypic optima by plasticity and evolution. Plasticity is the ability of a single genotype to produce multiple phenotypes as a function of the environment. It is often seen as an immediate response of individuals to changes in their environment. In contrast, evolution requires a change in allele

frequencies within a population. This process occurs over several generations and represents a

more long-term response, which can result in local adaptation. It is still not well understood how

these two processes are related and interact with each other (de Jong, 2005; Forsman, 2015;

Ghalambor, McKay, Carroll, & Reznick, 2007; Price, Qvarnström, & Irwin, 2003; Wund, 2012).

Plasticity does not affect a population's genetic composition and should thus not have long-term

consequences in the following generations. However, it changes the distribution of phenotypes

on which selection can act (de Jong, 2005; Pfennig et al., 2010; Pigliucci, 2005; Price,

Qvarnstrom, & Irwin, 2003). If plasticity perfectly matches the new phenotypic optimum, it

prevents evolution since no genetic changes are required and selection is prevented (Ghalambor

et al., 2007). On the other hand, plasticity is also crucial for a population's persistence and can

reduce the costs of selection (Chevin, Lande, & Mace, 2010; Pavey, Collin, Nosil, & Rogers, 2010).

It prevents extinction and protects populations from bottleneck effects, thereby maintaining a

higher genetic variation on which subsequently selection can act (Fitzpatrick, 2012; Pfennig et

al., 2010; Massimo Pigliucci, 2005). There is both, theoretical (Chevin et al., 2010; Draghi &

Whitlock, 2012; Fierst, 2011) and empirical work (Schaum, Rost, Millar, & Collins, 2013),

demonstrating that more plastic populations exhibit faster evolution. The benefits of plasticity

for persisting in new habitats were also demonstrated in invasive species (Molina-Montenegro,

Peñuelas, Munné-Bosch, & Sardans, 2012; Pichancourt & van Klinken, 2012; Yeh & Price, 2004).

The extent of plasticity can be represented as a reaction norm (Scheiner, 1993), which is the

phenotypic trait value as a function of an environmental variable. Evolution can affect the

reaction norm in two ways: The intercept can be shifted corresponding to a change in the mean

phenotypic value. Alternatively, the slope of the reaction norm, i.e. the plasticity, can be

changed. Thus, plasticity itself can also be subject to evolution given that there is sufficient genetic variation (Garland & Kelly, 2008; Nussey, Postma, Gienapp, & Visser, 2005).

Dependent on whether plastic responses are adaptive and increase fitness of an individual, different patterns describing the relationship between ancestral plasticity and evolution are possible. If plastic responses are adaptive, but not sufficient to reach the phenotypic optimum, evolution should work in the same direction as the plastic response (referred to as Baldwin effect (Crispo, 2007) or cogradient variation (Conover, Duffy, & Hice, 2009). In this case, selection may favour the most plastic individuals, causing evolved populations to exhibit a higher plasticity than their ancestors (Crispo 2007; Lande 2009). Another possible outcome is genetic assimilation: An initially environmentally induced phenotype can get fixed by a loss of plasticity and becomes continuously expressed even in the ancestral environment (Levis & Pfennig, 2016; Pigliucci, Murren, & Schlichting, 2006). In contrast, if plastic responses are maladaptive, we expect to observe evolutionary changes opposite to plasticity (countergradient variation ((Conover et al., 2009) or genetic compensation (Grether 2005)). Maladaptive plasticity was proposed as a possible mechanism promoting evolution since it moves phenotypes further away from the optimum and thereby increases the strength of selection on the phenotype (Ghalambor et al., 2007). Both co-gradient (Barton, Sunnucks, Norgate, Murray, & Kearney, 2014; Conover et al., 2009) and counter-gradient (Conover et al., 2009; Ghalambor et al., 2015; Laugen, Laurila, Räsänen, & Merilä, 2003) evolutionary changes have been found indicating that adaptive and maladaptive plasticity are common. It was even argued that reversion of ancestral plasticity occurs more frequently (Ho & Zhang, 2018) and is often not beneficial for long-term adaptation.

Plastic responses in physiology, behaviour or morphological traits are often initiated by changes in gene expression (Hodgins-Davis & Townsend, 2009; Wray, 2007). The transcriptome

represents a direct link between genotype and phenotype making it particularly interesting to study the interplay between plasticity and evolution. Transcription is highly plastic and modulating expression levels is an important part of an organism's physiological adjustment to environmental change (Gibson, 2008; McCairns & Bernatchez, 2009). On the other hand, there are also many studies demonstrating evolutionary divergence in gene expression between locally adapted populations (Alvarez, Schrey, & Richards, 2015; Guo et al., 2016; Romero, Ruvinsky, & Gilad, 2012; Townsend, Cavalieri, & Hartl, 2003; Whitehead & Crawford, 2006). Gene expression may even evolve more rapidly than changes in proteins since mutations affecting magnitude of expression are less likely to be deleterious than changes in protein structures (Carroll, 2005; Wray, 2007). However, it is not clear how fast plasticity in gene expression can change. Some studies reported changes in plasticity in some genes after adaptation to new conditions (Morris et al., 2014; Passow et al., 2017; von Heckel, Stephan, & Hutter, 2016), whereas others found only limited evolution of plasticity (Yampolsky, Glazko, & Fry, 2012) or less than expected (Huang & Agrawal, 2016).

In our study we used whole transcriptomes to understand the interplay between plasticity and evolution at the gene expression level using the model organism *Tribolium castaneum* (red flour beetle) and to quantify their contribution to the total phenotypic divergence between populations inhabiting different environments. More specifically, we were interested to test whether evolutionary changes in a new environment affected the same genes and were in the same direction as the ancestral plasticity. This can help us to understand how plasticity affected evolution, i.e. whether it can prevent evolution, or whether maladaptive plasticity increased strength of selection and promote evolution. Next, we tested how evolution affected gene expression reaction norms, i.e. whether adaptation to new conditions was achieved by changes in slopes or intercepts.

Material and Methods

Animal rearing, experimental evolution

We used the *Tribolium castaneum* Cro1 strain (Milutinović, Stolpe, Peuß, Armitage, & Kurtz, 2013), collected from a wild population in 2010 and adapted to lab standard conditions (32°C, 70% relative humidity (r.h.)) for more than 20 generations. Beetles were kept in 24h darkness on organic wheat flour mixed with 10% organic baker's yeast. We sterilized flour and yeast by heating them for 12h at 80°C before use. To test for adaptation to new environmental conditions we used replicate lines and exposed them to three treatment and control (CT) conditions. The conditions in the treatments were: Dry (D): 33°C and 30% r. h.; Hot (H): 37°C and 70% r. h.; Hot-Dry (HD): 37°C and 30% r. h. To generate replicate lines, we used 120 individuals (60 females and 60 males in the pupal stage) and placed them into a vial containing 80g medium. We produced six lines per selection regime (treatments plus control), resulting in a total of 24 lines. For each new generation, we randomly collected 120 pupae and placed them into a new vial. After seven to ten days, in which the pupae became adults, mated and laid eggs, adult beetles were removed by sieving the medium. We waited until the next generation (eggs/larvae in the medium) had reached the pupal stage and again collected 120 pupae per line to establish the next generation. This is similar to natural selection since individuals, depending on their fitness, do not contribute equally to the next generation. In generation 15 we produced additional mixed lines to prevent loss of genetic diversity by gene drift and inbreeding, which might impede adaptation: We mixed the six replicate lines of each selection regime in equal proportions (20 individuals from each replicate line) four times, resulting in four mixed lines with 120 individuals each. In total we had 39 lines: six normal and four mixed lines per selection regime (one line in D

became extinct). The transplant experiment to test for adaptation was conducted in generation

22.

Reciprocal transplant and fitness assay

Before testing for adaptation, all lines stayed for two generations in the same condition to

remove potential maternal or epigenetic effects (Figure 1): Beetles of generation 20 from all

selection lines were transferred to control conditions, in which they stayed for one week to mate

and lay eggs. After removal of the adults, we waited until their offspring had reached the pupal

stage and separated males and females. These individuals (generation 21) developed completely

in control conditions. When they reached the adult stage, we created 13 full-sib families per

selection line by transferring one virgin male with one virgin female of the same selection line in

15mL tubes with 1g of medium. After four days, in which the beetles could mate and lay eggs, 9g

of medium was added to provide food for the developing offspring and each mating pair was

transferred to a new vial. We repeated this three times, resulting in four vials per mating pair

containing medium and eggs. Immediately after removal of the mating pair, vials of each mating

pair were randomly assigned to the four different conditions, resulting in full-sib families split

across all conditions. As soon as offspring in these vials had reached the pupal stage, males and

females (four females and four males per family and condition) were separated and transferred

to 15 mL tubes with 5 g of medium and remained there until they were used for the fitness assay

two weeks later. These beetles were transferred to the treatments at the egg stage. They

developed completely in treatment conditions. We then assessed their performance in each

condition by estimating their fitness to test for adaptation. A virgin male and a virgin female of

the same selection line from the same condition but from different families were again placed

into a 15 mL tube with 1 g medium. After four days, the mating pair was removed. Males and

females were transferred to 1 mL Eppendorf tubes (one individual per tube), immediately frozen

in liquid nitrogen and stored at -80°C to use them for gene expression measurements. 9 g

medium was added to the mating tube. After four weeks (in CT and H) or five weeks (D and HD),

all offspring had reached the adult stage and were counted. We used the number of adult

offspring as an estimate of the fitness of a mating pair.

Statistical analysis

To test whether selection regime significantly influenced number of offspring produced and test

whether 20 generations in the treatments resulted in adaptation, we compared offspring

numbers of selection lines in their native condition to CT-lines transferred to the same condition.

We applied linear mixed models using the R-packages *lme4* (Bates, Mächler, Bolker, & Walker,

2015), and *lmertest* (Kuznetsova, Brockhoff, & Christensen, 2017) and *lsmeans* (Lenth, 2016) to

obtain *p*-values and confidence intervals. We included line and family as random factors,

selection and line type (mixed/normal) and their interaction as fixed effects. To test whether the

selection regime influenced how lines responded to the treatments, we used a linear mixed

model with offspring number in CT and treatment conditions as response variable, condition,

selection regime, line type (normal/mixed) and interactions as fixed effects and line, family and

interaction between line and condition as random effects. A significant interaction between

condition and selection regime indicates a significant effect of the selection regime on the

response to the conditions.

RNA extraction, library preparation and sequencing

208 female beetles (Table 1) p stored at -80°C were homogenized in Tri-Reagent® (Zymo

Research, California, USA) using an electric bead mill. RNA was extracted with the RNA Mini Prep

kit (Zimo Research, California, USA) following the instructions of the manufacturer. RNA-quality

was checked on a TapeStation (Agilent, Waldbronn, Germany) and concentrations were

measured with aQubit® Fluorometer (Life Technologies, California, USA). Libraries were created

with 500 ng RNA for each individual separately with the LEXOGEN mRNA-Seq Library Kit

following the manual (LEXOGEN GmbH, Vienne, Austria). Library quality was checked on a

TapeStation (Agilent, Waldbronn, Germany) and concentrations were determined by qPCR.

Libraries were diluted to the same molarity and pooled (33-36 libraries per pool). All treatments

and selection regimes were randomized during RNA-extraction, library preparation, and

sequencing. Single-end sequencing was performed in five runs on the Illumina NextSeq 500

(Illumina, Inc, California, USA) using the 75 cycles High Output Kit. After quality control using

FastQC (www.bioinformatics.bbsrc.ac.uk/projects/fastqc) reads (adaptors were trimmed and the

first 10 bases were hard trimmed, minimum average quality Q10, minimum tail quality 10,

minimum read length 20) were mapped against the reference genome

(ftp://ftp.ensemblgenomes.org/pub/release30/metazoa/gtf/tribolium_castaneum/Tribolium_ca

staneum.Tcas3.30.gtf.gz) with STAR v.2.5 (Dobin et al., 2013). We then used FeatureCounts (Liao,

Smyth, & Shi, 2014) to count the number of reads that mapped to each gene in the reference

genome. Mapping as well as read counting was performed within the data analysis framework

SUSHI (Hatakeyama et al., 2016).

196 *Gene expression analysis*

197 Gene expression analysis was done in R (R Core Team, 2017). We used the R package *edgeR*

198 (Robinson, McCarthy, & Smyth, 2010) for normalizing (method: TMM) expression data to cpm

199 (counts per million) after filtering lowly expressed genes (minimum of one cpm in at least two

samples). For subsequent differential expression analysis we used the R package *limma* (Law, Chen, Shi, & Smyth, 2014; Ritchie et al., 2015). From the differential expression analysis, we obtained the number of differentially expressed genes (DE genes) within lines between conditions (plastic changes, see Figure 1) or between lines of different origins (CT vs. selection) within conditions (evolutionary changes, see Figure 1). The total phenotypic divergence in gene expression between CT and treatments (i.e. total change TC) is the differential expression (log2-fold change) between CT-lines in CT and selection lines in the treatments (Figure 1 and 2). The ancestral plasticity (PC_{CT}) is the differential expression of CT-lines between CT and treatment conditions, while the evolved plasticity (PC_{Sel}) is the same difference measured in selection lines. The evolutionary changes are EC_T when measured as differential expression between CT and selection lines in the treatments and EC_{CT} when measured in CT (Figure 2). Finally, differences between plastic responses of CT- and selection lines (the interaction between treatment and selection regime) give the evolutionary change in plasticity. To partition TC into changes explained by ancestral plasticity (PC_{CT}) and evolutionary changes (EC_T, see Figure 2 and 3), we calculated the relative contribution of each component to the total. We obtained the log2-fold changes of each transcript to evaluate and compare the magnitude of the plastic and evolutionary changes. We accounted for non-independence among individuals from the same line by using the *duplicateCorrelation* function (Smyth et al. 2005) and added sequencing runs as batch effect. A gene is classified as differentially expressed (DE) with a FDR ≤ 5% after adjusting for multiple testing (Benjamini and Hochberg 1995).

In a second step, we used the normalized read counts (cpm, TMM-normalized) corrected for batch effects (sequencing runs) using the *removeBatchEffect* function in the *limma* R package (Ritchie et al., 2015). These counts were used to quantify original levels Lo, plastic levels Lp and adapted levels La of selection lines. This allowed us to conduct an analysis for each selection lines separately as was described by (details see below)

Evolution of reaction norms

Reaching a new phenotypic optimum can be achieved by changing the mean (intercept of the reaction norm) or by changing the plasticity of a trait (slope of reaction norm) (Figure 3). We were interested to assess which of these patterns was more prevalent in gene expression evolution. To test for an evolutionary change in plasticity (slope), we first evaluated the similarity in plasticity between CT- and selection lines by calculating the correlation between PC_{CT} and PC_{Sel} (Figure 3A) among all genes, independent of the significance. Second, to test for an evolutionary change in mean expression (intercept), we calculated the correlation between the evolved differences in CT (EC_{CT}) and in treatment conditions (EC_T). A significant correlation indicates that the overall mean has changed (Figure 3A). Further, for all genes with significant EC_T, we quantified how much of this difference could be explained by a shift in the mean or by a change in plasticity. A shift in the mean corresponds to EC_{CT}, whereas the remaining difference (EC_T-EC_{CT}) gives the evolutionary changes in plasticity (see also Stoks et al. 2016). Relative contributions of each component were obtained by dividing them by the total EC_T. To test for significant effects of the selection regime on differential expression (i.e., on number of DE genes, log2-fold change, and correlations), we used a permutation test. We randomly assigned samples and their transcriptomes to either CT selection or treatment (D, H, HD) selection (number of samples for each selection was not changed) and repeated the DE analysis. We kept the original assignment to lines and conditions and repeated the DE analysis for each permuted data set. If observed values (e.g. number of DE genes, correlations) obtained from the original dataset were higher than the most extreme 5% of the distribution calculated from permutations, they were considered significant.

Comparing plastic and evolutionary responses

To infer the relationship between ancestral plasticity and evolution we compared the direction of PC_{CT} to the direction of EC_T. Evolution may *reinforce* the plastic response when PC_{CT} is in the same direction as EC_T (Figure 2B). If EC_T is in opposite direction, it *reverses* PC_{CT} (Figure 2A). To test which of these patterns was more prevalent, we followed Ho and Zhang 2018 (Ho & Zhang, 2018). Expression levels (cpm) of CT-Lines in CT conditions represented the original stage (*Lo*), CT-Lines in treatment the plastic stage (*Lp*), and selection lines in their respective condition the adaptive stage (*La*) (Figure 2). For subsequent analysis we used genes with appreciable PC_{CT} (|*Lp* − *Lo*|> 0.2 *Lo*) and EC_T (|*La* − *Lp*|> 0.2 *Lo*) (Ho & Zhang, 2018) and calculated the proportion of plastic genes with reinforced or reversed changes.

An alternative classification is to assess whether PC_{CT} brings expression levels closer to *La*. In this case, the total change TC would be larger than EC_T (Figure 2C). Such a pattern can occur even if PC_{CT} and EC_T are in opposite directions and classified as reversion (Figure 2C). To confirm that our results were not sensitive to the applied cutoff (20% *Lo*), we repeated the analysis with a cutoff of 50% *Lo* and without any cutoff (Supporting information, Figure S1.3 A-H). Recently, it was pointed out (Mallard et al. 2018; Ho and Zhang 2019) that an excess of reversions relative to reinforcements is expected to be observed due to a statistical artefact that cannot be completely removed by permutation tests. We therefore applied an additional parametric bootstrapping test as proposed by Ho and Zhang (Ho and Zhang 2019), to compare proportion of reversions and reinforcements (Supporting information, Figure S1.3 J-L).

To better understand the relationship between the within-line proportions of reversed genes and proportions of genes with *La* closer to *Lo* with adaptation, we calculated the Spearman correlation between the proportions of reversed expression changes (or *La* closer to *Lo* respectively) and mean offspring number in seven selection lines in HD. We focused on HD

275   because it was the most extreme environment with the strongest decline in offspring number.

276   To test for significance, we used permutations: Mean offspring numbers were randomly assigned

277   to lines and correlation was calculated again. Proportion of permutations with a correlation

278   coefficient exceeding the observed value gave the respective *P*-value.

Results

Fitness assay showed evolutionary adaptation

We found that selection lines had a higher offspring number in their native condition compared to CT-lines (Dry: $F_{1,14}$= 10.04, P=0.007; Hot: $F_{1,16}$= 4.78, p=0.044; Hot-Dry: $F_{1,16}$= 23.85, p=1.660E-04), confirming that adaptation had occurred (Figure 4). In contrast to treatment conditions, there was no difference in offspring number between CT- and selection lines under CT conditions (CT-lines: 64.89 [61.49, 68.29]; D-lines: 65.76 [61.26, 70.25], H-lines: 67.41 [63.76, 71.06], HD-lines: 65.56 [62.17, 68.94]) (Figure4). Using three additional mixed models, we compared how lines from different selection regimes responded to treatments. We found significant negative effects for all stress treatments (D: $F_{1,14}$= 222.10, p = 5.21E-10; H: $F_{1,16}$= 615.10, p = 3.37E-14; HD: $F_{1,16}$= 714.39, p = 8.664e-15) (Figure4). Interaction between selection and treatments, i.e. whether the response to the treatment was different dependent on selection regime, was significant for HD-lines ($F_{1,16}$= 12.12, p=0.003051) and for D ($F_{1,14}$= 13.22, p= 0.002682), but not for H ($F_{1,16}$= 0.6879, p= 0.42). Interaction between selection regime, treatment and line type (normal/mixed) was not significant in any treatment indicating that mixing lines did not have an effect on adaptation.

Evolution of gene expression

We found many more genes with expression divergence relative to Control (TC) than genes with evolved changes in the treatments (EC_T) or in control conditions (EC_{CT}) (Table 2). Increased temperature caused expression divergence in a larger number of genes than reduced humidity (Table 2). Most DE genes with significant TC also had significant ancestral plasticity PC_{CT} (Dry: 33.9%; Hot: 72.9%, Hot-Dry: 82.2%), but only a minority showed significant evolutionary change EC_T (Dry: 4.9%; Hot: 1.1%; Hot-Dry: 1.0%, see also Figure S1.1). About half of the DE genes with

EC_T in Hot and Hot-Dry then did not diverge significantly from their ancestral expression levels in

Control (TC ~ 0). Overall, TC had a larger log2-fold change contribution from plastic changes

(PC_{CT}) than evolved changes in Hot and Hot-Dry (median proportions PC_{CT}/TC; Hot: 71.47%; Hot-

Dry: 86.30%), but not in Dry (42.90%). Many genes had PC_{CT} > TC, especially in HD (Figure 5). In

these cases, EC_T was negative and reduced TC (Figure 5).

Evolution of reaction norms

We were then interested to examine whether EC_T were due to a change in mean expression

(shift in the intercept of reaction norms) or to a change in plasticity (different slopes of reaction

norms, see Figure 3). First, we found a significant correlation between EC_{CT} and EC_T, in all three

treatments (permutation tests: $p < 0.0001$) (Table 4, Figure 5). This indicates that for a majority

of genes, evolution in the past 20 generations affected their mean expression levels by shifting

the intercept of reaction norms (see Figure 7A). Second, we also found a highly positive

correlation (Figure 7A) between PC_{CT} and PC_{Sel} (Figure 3), indicating that plasticity was mainly

preserved during adaptation to new conditions. A permutation test showed that this correlation

was not significantly different from correlations obtained from permuted datasets (Table 2),

where samples were randomly assigned to selection regimes, suggesting that the treatments did

not have specific effects on plasticity. When we quantified the relative contributions of changes

in the mean versus changes in plasticity to EC_T, we found that evolution of the intercept

explained more evolutionary divergence than evolution of the slope of reaction norms, especially

in Dry (Figure 6).

Evolution of plasticity in DE genes

We found only five genes with significantly evolved plasticity in the DE analysis, after correcting

for multiple testing. Although plasticity seemed to be preserved overall, we found evidence that

it evolved among DE genes. The DE analysis revealed a reduced number of plastic genes in

selection lines compared to plastic genes in CT-lines (Table 2), although the difference was only significant in HD (Table 3). It further showed that ancestral plasticity PC_{CT} was larger than evolved plasticity PC_{Sel}, in D and HD (Table 3), indicating a reduction of the magnitude of plastic responses after adaptation. We found the same outcome when focusing on genes with significant EC_T, in HD (Mann-Whitney U test: U = 2133, P = 2.102E-04; median: CT-lines: 0.25, HD-lines: 0.13), but not in D (U = 144, P = 0.584; median: CT-lines: 0.10, D-lines: 0.14) or H (U = 386, P = 0.158; median: CT-lines: 0.21, H-lines: 0.12) (Figure 8).

Effect of plasticity on evolutionary responses

We could not find evidence that plasticity impeded evolution. In the DE analysis genes with a significant ancestral plastic response were overrepresented among the evolved DE genes in H and HD (H: $\chi2$ = 4.51, df = 1, P = 3.38E-02; HD: $\chi2$ = 9.71, df = 1, P = 1.84E-03), but not in D (P = 0.113 Fisher's exact test). Furthermore, when we analysed each selection line separately and compared their expression levels to CT-lines, the proportion of genes exhibiting substantial PC_{CT} ($|Lp – Lo| > 0.2$ Lo) were also overrepresented among evolved genes with EC_T ($|La - Lp| > 0.2$ Lo) (Table S2.1).

Maladaptive plastic changes may be reverted by evolutionary changes when the plastic response overshoots the optimum expression level in a new environment. Among the genes with significant EC_T in the DE analysis, we found that PC_{CT} was generally in opposite direction (Figure 8). Enrichment tests revealed that genes with positive EC_T in the DE analysis were overrepresented among genes with negative PC_{CT} in H and HD (Fisher's exact test D: P = 0.076; H: P = 0.002; HD: P = 9.26e-06), and genes with negative EC_T were overrepresented among genes with positive PC_{CT}, in H and in HD as well (Fisher's exact test D: P = 0.073; H: P = 0.021; HD: P = 0.049)(Figure 6). In almost all genes that showed significant EC_T and PC_{CT}, the responses were in

opposite direction (Dry: 2 out of 2 genes; Hot: 12 out of 13; Hot-Dry: 33 out of 34). We also

found a negative correlation between PC_{CT} and EC_T in H and HD, but not in D (Supporting

information, Figure S1.2).

When we considered mean expression levels per line with (PC_{CT} > 20% *Lo* and EC_T > 20% *Lo*) we

obtained similar results as in the DE analysis. Reinforcements were present but less frequent

than reversions in all treatments (proportions of plastic responses with reversion/reinforcement:

Dry: 48.5 ± 1.0% / 34.3 ± 0.8 %, P < 2.2e-16 (binomial test); Hot: 44.6 ± 0.9% / 28.8 ± 1.2%, P <

2.2e-16; Hot-Dry: P < 2.2e-16; Figure 10A). Different cutoff values did not influence these

outcomes (Figure S1.3). Parametric bootstrapping as an additional test (Ho & Zhang, 2019) ,

confirmed our results (Supporting information, Figure S1.3). We also found that a substantial

proportion of plastic genes (PC_{CT} > 20% *Lo*) had *La* closer to *Lo* than to *Lp* (Dry: 45.4 ± 1.1%; Hot:

41.2 ± 1.2%; Hot-Dry: 40.7 ± 2.5 %)(Figure 10B). EC_T (EC_{CT} > 20% *Lo*) genes showed a higher

proportion of reversions (Dry: 34.1 ± 0.8%; Hot: 40.0 ± 0.9%; Hot-Dry: 43.0 ± 2.1 %) compared to

reinforcements (Dry: 24.1 ± 0.6%; Hot: 25.8 ± 1.1%; Hot-Dry: 26.8 ± 1.6%). 41.8 ± 0.7% (Dry), 34.2

± 0.8% (Hot), 30.2 ± 1.0 % (Hot-Dry) of EC_T genes did not show PC_{CT}.

In case of genetic assimilation, i.e. loss of plasticity in selection line and continuous expression of

the induced phenotype, evolutionary differences would not be visible in treatment condition.

We calculated the mean expression levels of selection lines in Control and defined genes as

showing assimilation if they met three criteria: no PC_{Sel} (absolute difference between selection

lines in CT and in treatment < 20% *Lo*), PC_{CT} > 20% *Lo*, and *Lp* ~*La* (absolute difference < 20% *La*).

The proportion of PC_{CT} genes showing genetic assimilation was small in all conditions (Dry: 13.5

%, Hot: 11.5 %, Hot-Dry: 14.7%; Supporting information Table S2.3).

Proportions of reversed plastic responses and association with fitness

To gain a better understanding of the adaptive value of the changes of expression levels in the

evolved lines, we tested for an association between within-line proportion of reversed or

reinforced plastic responses and the average fitness of the lines in the HD treatment. We found

that lines with a higher proportion of reversions had a higher average offspring number

(correlation: 0.82, P = 0.012, Figure 11) and we found a negative but non-significant correlation

between fitness and reinforcements (correlation: -0.43, P = 0.85). When we tested for an

association between fitness and proportion of genes with La closer to Lo, we also found a

positive correlation (correlation: 0.86, P = 0.006). Overall, better adapted lines (higher fitness in

HD) showed a higher proportion of reversed ancestral plasticity and were more similar to the

original expression levels of CT-lines in CT. Performing the analysis with gene expression data in

H and D provided similar correlations, although not significant because of lower sample sizes

(Supporting information, Table S2.2).

Discussion

We studied plastic and evolved responses in gene expression of *T. castaneum* in response to

three new environmental conditions (Dry, Hot, Hot-Dry). After 20 generation of experimental

evolution, we were able to detect adaptation and found significant evolutionary changes in

expression levels. Comparing these changes with ancestral plastic responses to the conditions

showed that a reversion of plasticity was the more frequent (ca. 60 % of all genes; genes with

significant plastic changes in DE analysis: >95%). Number of genes where ancestral plasticity was

reinforced by evolution was smaller (ca. 40%; DE analysis: <5%) and only a small proportion of

genes (8-10% of plastic genes) showed expression patterns consistent with genetic assimilation.

Of the genes with evolutionary differences in treatment conditions, a high proportion (ca. 40%)

showed expression levels closer to the ancestral than to the plastic stage suggesting that

plasticity was maladaptive. Although plasticity showed a high degree of preservation in terms of

responding genes and direction of response, we found evidence that selection lines evolved a

reduced plasticity and thus partly compensated the maladaptive ancestral response. We were

further able to show an association between the proportion of reversed plastic responses and

adaptation (mean fitness per line) in the most stressful treatment Hot-Dry.

Maladaptive plasticity

Different patterns describing the relationship between plastic and evolved changes in gene

expression have been documented. It was suggested that plasticity might help populations to

persist after environmental change, or to colonize new habitats by bringing phenotypes closer to

the new optimum. Studies found support for this hypothesis by showing that plastic responses of

non-adapted individuals diminished differences to native populations (Lohman, Stutz, & Bolnick,

2017; Mäkinen, Papakostas, Vøllestad, Leder, & Primmer, 2016). Adaptive plasticity can also be

indicated when plastic and evolutionary responses are in the same direction (Li, Li, Song, Wang,

& Zhang, 2017), or when plasticity is higher in adapted populations (Hasan et al., 2017; McCairns

& Bernatchez, 2009), suggesting that most plastic individuals were favored by selection.

However, there are also examples for the reversed pattern suggesting that plasticity was

maladaptive, at least on the long-term. In wild populations of *Fundulus heteroclitus* evolved

changes to different temperatures were opposite to plastic responses of the ancestral

population (Dayan et al., 2015). *Rhagoletis* flies shifting to a new host fruits showed evolutionary

responses opposite to plasticity of non-adapted species (Ragland et al., 2015). Experimental

evolution studies found countergradient evolution in *Drosophila* adapting to different diets

(Huang & Agrawal, 2016; Yampolsky et al., 2012). A comparative study (Ho & Zhang, 2018)

analyzing data of multiple experimental evolutions suggested that reversions of gene expression

changes might be a general pattern during adaptation.

Our study fits to these previous observations. We found a higher proportion of reversions in all

conditions indicating mostly maladaptive plasticity. An alternative explanation for the prevalence

of reversions without plastic responses being necessarily maladaptive would be that CT-lines

exhibited a response in the right direction, but overshot an optimum expression level (Figure 2C).

Fine-tuning during long-term adaptation could then lead to a partial reversion of the plastic

response. We took this possibility into account by not only focusing on a reversion of plasticity,

but also testing whether *La* was closer to *Lp* or to *Lo* (see Figure 2C). We found that adapted lines

showed a high proportion of expression levels that were closer to their ancestral level *Lo*. This

confirms that many plastic responses moved phenotypes further away from the new optimum

and evolution compensated for these detrimental changes. This pattern might even become

more pronounced after some additional generations, given that our selection lines still show

incomplete adaptations, i.e. strong reduction in offspring number compared to control levels. In

addition we were also able to show a direct association between proportion of reversions and

adaptation.

Evolution of reaction norms

The ancestral maladaptive plasticity can be compensated by shifts in the intercept or changes in

the slope of reaction norms. Both are not mutually exclusive and can occur together in the same

trait. We were aiming to quantify their relative importance for evolutionary responses in

transcriptomes.

Gene expression studies so far provided mixed results regarding the evolution of plasticity.

Drosophila populations adapted to different temperatures showed local adaptation, but there

was no evidence for evolution of thermal reaction norms of different transcripts and changes

affected mainly expression mean (Clemson, Sgrò, & Telonis-Scott, 2016). Experimental evolution

studying *Drosophila* under variable diets found no significant changes in plasticity (Yampolsky et al., 2012) or less than expected (Huang & Agrawal, 2016). In contrast, other studies found differences in temperature responses between tropical and temperate *Drosophila* populations (Levine, Eckert, & Begun, 2011; von Heckel et al., 2016). Other examples for differences in genes expression plasticity between adapted and non-adapted populations include temperature (Morris et al., 2014) and salinity (Gibbons, Metzger, Healy, & Schulte, 2017; McCairns & Bernatchez, 2009) responses of marine and freshwater sticklebacks, temperature response of killifish populations from different latitudes, as well as plastic responses to toxic hydrogen sulphide (H_2S) of fish population from H_2S rich springs versus non-toxic springs (Passow et al., 2017). There was no consistent pattern regarding the direction in which plasticity evolves: In some cases adapted population showed an increase in plasticity (Morris et al., 2014), in other cases plasticity was reduced (Huang & Agrawal, 2016; Ragland et al., 2015; von Heckel et al., 2016) or reduction and enhancement of plasticity were equally frequent (Gibbons et al., 2017; Yampolsky et al., 2014). Overall, there is evidence in multiple species that expression plasticity of some genes can evolve. However, even in some of these studies reporting evolved plasticity (Dayan et al., 2015; Gibbons et al., 2017; Morris et al., 2014) the number of transcripts with significant changes in the mean was much higher and large parts of the plastic responses showed a high degree of preservation.

In accordance with these previous findings we found that changes in the mean contributed more to the observed differences than changes in plasticity. A possible reason might be that genetic variation in mean expression was higher than genetic variation in plasticity. In addition, we did not select for changes in plasticity directly since the conditions in the treatments were constant. Selection was therefore on expression levels in the treatment and only indirectly on plasticity. Plasticity could evolve if mean expression levels were genetically correlated with plasticity.

Although we did not directly test for such correlations, we found evidence for evolution of

plasticity in DE genes that is in genes with largest changes in expression level between CT and

treatment conditions. Although plastic responses showed a high degree of preservation, we

found evidence for evolutionary changes in the magnitude of plastic responses.

Why can plasticity be maladaptive?

New stressors might disturb homeostasis resulting in inappropriate responses and long-term

adaptation therefore restores ancestral phenotypes by genetic changes, referred to as genetic

compensation (Grether, 2005) or counter-gradient variation (Conover et al., 2009). However, in

our study we applied relatively mild stressor treatments, i.e. individuals were able to survive and

reproduce. Drought and heat are also stressors, which *T. castaneum* had experienced in the past

(Sokoloff, 1972), so there had probably been selection on plastic responses to be beneficial.

However, plastic responses might be optimized for a short-term exposure: Allocation of

resources from reproduction to protection might increase survival probability and allow

individuals to continue reproduction as soon as the stress has disappeared, but this response

becomes maladaptive during continuous exposure and should therefore be under negative

selection. Expression of stress related genes is in general accompanied by a down-regulation of

genes involved in growth and reproduction due to an allocation of resources (Schwenke, Lazzaro,

& Wolfner, 2016; Sokolova, 2013). A well-studied example are heat shock proteins (hsp). Hsp are

well known for their protective function and to be crucial for survival (Feder & Hoffman, 1999),

but it was also shown that their expression came at a cost (Feder et al., 1998; Sørensen,

Kristensen, & Loeschcke, 2003). Accordingly, it was often found that hsp expression in

populations adapted to warmer climates is lower compared to non-adapted populations

(Fangue, Hofmeister, & Schulte, 2006; Narum & Campbell, 2015; J G Sørensen, Dahlgaard, &

Loeschcke, 2001). In general, other protection mechanisms independent of ancestral plasticity

may arise during long-term adaptation (e.g. enzymes, which are more stable at high

temperature) and make the costly stress response expendable.

An alternative explanation for the reduced plasticity in adapted lines is that the signal

responsible for eliciting the plastic responses is based on any kind of damage caused by heat or

stress in general (Kültz, 2005). Higher resistance in adapted lines might shift the inducing

thresholds (Sikkink, Reynolds, Ituarte, Cresko, & Phillips, 2014) above the levels we applied in the

treatments.

Interestingly, we found no differences in fitness between lines from different selection regimes

under CT conditions (Figure 4). We could detect some genes with different expression levels in

the treatments and found in general a correlated EC_{CT}, but this did not affect offspring number. It

indicates a lack of fitness trade-offs, where alleles providing a fitness advantage in one

environment are detrimental in another. Together with the observation that selection lines

evolve to bring expression closer to ancestral CT-levels, it suggests that for most genes the

optimal expression level is not different between conditions.

Potential caveats

The number of genes with significant PC_{CT} in the DE analysis was much higher compared to genes

showing EC_T. One possible explanation would be that adaptive plasticity prevented evolution. If

the plastic responses matched the optimum, no genetic changes in the selection lines are

expected to occur. However, when we analysed each line separately and considered a gene as

evolved if the mean difference between *La* and *Lp* was more than 20% of the *Lo*, we found

approximately the same number of genes with EC_T and PC_{CT} (Supporting information Table S2.2).

In the DE analysis in *limma* we did not analyse each line separately but treated them as biological

replicates. Comparisons between conditions, i.e. plastic changes, can be made within lines. They

should thus be more precise and statistical power should be higher than comparisons between

selection regimes, i.e. evolved changes, that have to be made between lines. Differences

between lines from the same selection regime lower the ability to obtain significant evolutionary

changes. These differences can arise from genetic drift. Since our population size was relatively

small (120 individuals per line) this might have been an important factor. Another explanation is

that lines from the same selection regime differed how exactly they improved their fitness in the

respective treatment. Since fitness is a highly polygenic trait, the responsible genes contributing

to an increase may not be the same. For the most extreme treatment HD, where we sequenced

seven lines, we further found considerable differences in fitness between the lines, suggesting

that not all of them were at the same stage of adaptation. It is therefore not surprising that

expression levels did not evolve in the same way.

We used two different approaches, a classical DE analysis treating lines as replicates and

analysing lines separately using mean expression level of samples. The DE analysis resulted in a

much smaller number of genes with evolved changes. Since it requires that a gene shows similar

changes in all replicate lines, it is more conservative. If the main interest of a study is to identify

promising candidate genes for future more detailed analyses it is the appropriate approach to

keep FDR as low as possible. In contrast, if the focus is more on general patterns, a less stringent

analysis using mean expression levels can give us a more complete picture. The number of

evolved genes was much higher using this approach. Many of the observed differences between

selection lines and Control-lines might be caused by drift and are therefore not significant in the

DE analysis. However, since genetic drift is random it cannot explain the observed excess of

reversions over reinforcements.

Although we demonstrated that gene expression changed during evolution, it is not clear

whether these changes are the cause for an increased fitness in these conditions or whether

they are rather the consequence of adaptation and being less stressed. One disadvantage in

studying whole transcriptomes is that not all responding genes might be of functional

importance but are correlated to other adaptive changes. Reversion of plasticity during

adaptation can implicate that responding genes were not the direct target of selection and other

mechanisms, e.g. changes in protein structure of enzymes, were responsible for adaptation of

selection lines. Future studies that manipulate expression and test for correlated changes in

offspring are needed to confirm adaptive value of expression changes.

Conclusions

Evolution compensated ancestral plasticity and reduced total phenotypic divergence between

control and treatment conditions suggesting that ancestral plasticity was maladaptive for long-

term adaptation. In the most stressful treatment, selection lines with higher fitness show a

higher proportion of reversions and a higher similarity with ancestral expression. Differences

between adapted lines and CT-lines in the treatment were mainly due to a change in mean

expression (i.e. shift in the intercept of reaction norms), while plasticity was preserved in terms

of affected genes and direction of change. However, we found that a part of the differences in

the treatments can be explained by a reduction in the magnitude of plasticity in adapted lines.

Plastic genes were clearly overrepresented among evolved genes, thus showing that plasticity

did not impede evolution. In contrast, our results suggested that maladaptive plasticity increased

selection and promoted evolution. Similar results in all three stress treatments indicated that

these findings may represent a general pattern of gene expression adaptation.

Acknowledgements

This work was supported by the Swiss National Science Foundation, grants PP00P3_1144846 and

PP00P3_176965 to FG. We thank Sonja Sbilordo, Sara Meier, Cilgia Lippuner, Tim Schoch and Tim

Emmenegger for helping during the fitness assay, Elke Karaus, Martina Berchtold, Encarnación

Lozano for RNA extraction and quality control, Lucy Poveda and Maria Domenica Moccia for

advice in library preparation and sequencing, Lennart Opitz for helping to process the RNA-seq

data.

Figure 1: Schematic representation of the experimental evolution design. Lines were derived from a single outbred
Tribolium castaneum population (collected in 2013, strain Cro1) that had lived under control conditions for ca. 30
generations. For each selection regime we used six replicate lines. In generation 15, four additional lines were created

by mixing the six lines of each selection regime in equal proportions, resulting in a total of 10 lines per selection
regime. Lines stayed in control or treatment conditions for 20 generations. After one generation in control condition
to remove potential maternal and epigenetic effects, individuals from all lines were transferred to control and
treatment conditions in the egg stage. Fitness and gene expression were measured in adult females. To get plastic
responses in gene expression, expression levels in treatment conditions were compared to levels under control
conditions of the same line. Evolved response is the difference within treatment conditions between control and
selection lines. Control conditions: 33°C, 70% relative humidity r.h.; Treatments: Dry: 33°C, 30% r.h.; Hot 37°C, 70%
r.h.; Hot-Dry: 37°C, 30% r.h.

Figure 2: Possible relations between plastic change PC and evolutionary change EC. Gene expression levels of control
lines (blue) in the ancestral control condition represent the original stage Lo, expression in the treatment the plastic
stage Lp. Expression levels of selection lines (orange) in the treatment give the adapted stage La Arrows indicate the
direction of PC (red) and EC (blue). PC can be in opposite direction to EC (reversion **(A)**) or it can be in the same
direction (reinforcement **(B))**. However, even if PC and EC are opposite to each other, PC can bring expression levels
closer to levels of the adapted lines. In this case, the total change TC, (difference between expression levels of control
lines in control conditions and selection lines in treatment) is larger than EC **(C)**.

Figure 3: Possible ways how non-adapted lines (in blue) can reach the phenotypic optimum in a new environment. We
assume that expression levels of adapted selection lines (in orange) represent the optimum level in the treatment.
During adaptation, lines could have reached the optimum by either changing mean expression, i.e. shift in the
intercept of the reaction norm **(A)** or by changing their plasticity, i.e. the slope of the reaction norm **(B)**. In case of a
change in the mean, plastic changes of Control lines PC_{CT} and selection lines PC_{Sel} as well as observed evolutionary
change EC_T in treatment and in Control EC_{CT} would be highly correlated because reaction norms (red arrows) remain
parallel **(A)**. If observed EC_T in treatment is due to a change in plasticity only, no correlation between EC_T and EC_{CT}
should exist, and the correlation between PC_{Sel} and PC_{CT} should be small **(B)**.

Figure 4: Offspring number of Control and selection lines under different conditions. Selection lines could adapt for 20 generations.

Figure 5: Relative contribution of ancestral plasticity and evolutionary change to total divergence for all genes showing significant total divergence in the DE analysis. Number of genes: Dry: 283, Hot: 1192, Hot-Dry: 2045.

Figure 6: Relative contribution of changes in the mean (shift in the intercept of reaction norm) and changes in plasticity (different slopes of reaction norms) to evolutionary differences between control-lines and selection lines in

the treatment. Only genes with significant differences in the DE analysis were included. Number of genes: Dry: 18,
Hot: 25, Hot-Dry: 55.

Figure 7: Correlation of plastic responses in gene expression of control lines and selection lines (**A**) and correlation of
evolved divergence in gene expression in control and treatment conditions (**B**). Selection lines had lived in the
respective treatment for 20 generations. After filtering lowly expressed genes (minimum of 1cpm in at least 2 samples)
11534 genes remained that were used for calculating correlations.

626
627 **Figure 8:** Plastic responses in control-lines and adapted selection-lines of genes showing significant evolutionary
628 changes in expression in the treatments Dry **(A)**, Hot **(B)** and Hot-Dry **(C)**. Plastic responses in adapted selection lines

... ve (smaller logFC). Furthermore, plastic changes are mainly opposite to evolved changes. Genes that evolved to lower expression in the treatments (represented in blue) show positive plastic changes in control-lines, i.e. they are up-regulated. Genes that evolved to higher expression levels (in red) are down-regulated in the non-adapted Control-Lines.

Figure 9: Examples of gene expression levels in control and treatment conditions of control lines (blue) and adapted selection lines (orange). **A:** example of one gene with no change in plasticity and a reversion of the plastic response, which was the commonly observed pattern. Plastic changes PC of control lines (PC_T) and selection lines (PC_{Sel}) positively correlated and very similar. Evolutionary changes EC in treatment are positively correlated with evolutionary changes in control conditions (EC_{CT}) due to a change in mean expression. In most cases, EC_{CT} smaller than EC_T . **B:** Example of a gene with significant evolutionary divergence in expression treatment conditions: If selection lines reduce their plastic response PC_{Sel}, it can increase divergence EC_T in the treatment. For most genes we observed that plastic changes (red arrows) are in the opposite direction to evolved changes (blue arrows).

Figure 10: Comparison of plastic and evolutionary changes in gene expression in response to three environmental conditions. **A:** Proportion of genes (average over selection lines) showing a reversion or reinforcement of ancestral plasticity during evolution. **B:** Proportion of genes with expression levels after evolution (adapted stage La) closer to original levels (Lo, i.e. Control-lines in control conditions), or closer to plastic levels (Lp, i.e. expression levels of Control-lines in treatment). Only genes exhibiting substantial plastic changes (|Lp-Lo| > 20% of Lo) as well as evolutionary changes (La-Lp| > 20% of Lo) were used for analysis.

Figure 11: Relationship between plastic and evolved changes in gene expression in response to hot-dry conditions. Expression levels of Control-lines in control conditions represent the ancestral stage Lo, Control-Lines in hot-dry conditions the plastic stage Lp and adapted Hot-Dry-lines in hot-dry conditions the adapted stage La. **A:** Barplots show the proportion of genes in each Hot-Dry line with a reversion (evolved response in opposite direction to plastic response of Control-Lines) or a reinforcement of the plastic response (same direction as plastic response of Control-Lines) in the left graph. The right graph shows the proportion of genes with La closer to Lp (in yellow) and La closer to Lo (in green). **B:** Relationship between the proportion of reversed plastic responses and mean fitness (=offspring number) per Hot-Dry line and between proportion of genes with La closer to Lo and mean fitness. Only genes exhibiting substantial plastic changes (|Lp-Lo| > 20% of Lo) as well as evolutionary changes (La-Lp| > 20% of Lo) were used for analysis. P-values of the spearman correlations were obtained by 10,000 permutations. 95 % Confidence intervals are based on a non-parametric bootstrap test.

Table 1**:** Number of sequenced replicate lines and individuals per selection and treatment, which were used for this study. Selection lines could adapt to conditions for 20 generations. Control (CT) conditions: 33°C, 70% relative humidity (r.h.); treatments: Dry (D): 33°C, 30% r.h.; Hot (H): 37°C, 70% r.h.; Hot-Dry (HD): 37°C, 30% r.h.

Selection	Conditions				
	Total	CT	D	H	HD
CT					
Number lines	**7**	7	5	5	7
Number Ind.	**89**	25	20	19	26
D					
Number lines	**4**	4	4	–	–
Number Ind.	**32**	16	16	–	–
H					
Number lines	**5**	5	–	5	–
Number Ind.	**40**	20	–	20	–
HD					
Number lines	**7**	7	–	–	7
Number Ind.	**56**	28	–	–	28

672
673 **Table 2:** Significantly differently expressed genes. For evolved differences, control-lines and selection-lines are
674 compared within condition (e.g. Dry-lines vs Control-lines in Dry). Plastic response gives the number of genes that
675 changed expression between control and treatment conditions (e.g. Dry-lines in Dry vs Dry-lines in control). Different
676 plasticity gives the number of genes with significant different plastic responses in lines from different selection
677 regimes. Analysis was conducted using the R package limma (Ritchie et al., 2015).

		Dry	**Hot**	**Hot-Dry**
Evolved difference in Control	down	3	0	6
	up	1	9	4
	total	**4**	**9**	**10**
Evolved difference in treatment	down	9	8	22
	up	9	17	33
	total	**18**	**25**	**55**
Plastic response Control-Lines	down	209	1765	2451
	up	156	1714	2200
	total	**365**	**3479**	**4651**
Plastic response Selection-Lines	down	28	1417	1649
	up	21	1381	1470
	total	**49**	**2798**	**3119**
Total change (Selection-lines in treatment vs Control-lines in Control)	down	154	564	1016
	up	129	628	1029
	total	**283**	**1192**	**2045**
different plasticity	total	**0**	**1**	**4**

678
679 **Table 3:** P-values obtained from permutation tests (10,000 permutations). Samples were randomly assigned to either
680 control or treatment selection and differential expression analyses were repeated. Significance was assessed by
681 calculating the proportion of permutations with more extreme values than the observed one. Control (CT) conditions:

682 33°C, 70% relative humidity r.h. Conditions in treatments: Dry (D): 33°C, 30% r.h.; Hot (H): 37°C, 70% r.h.; Hot-Dry
683 (HD): 37°C, 30% r.h.

	Selection		
	D	**H**	**HD**
CT-Lines have more genes with significant plastic responses compared to adapted selection lines	0.2981	0.3037	**0.0325**
Magnitude of plastic response is higher in CT-Lines	**0.0493**	0.1676	**0.0491**
Individuals from different selection regimes show lower similarity in plastic responses [†]	0.8608	0.3914	0.5217
Number of genes with significant differences in expression levels is higher in treatment than in CT conditions	**0.0095**	**0.0003**	**0.0031**
Differences in expression levels in CT conditions and treatment are correlated	**< 0.0001**	**<0.0001**	**< 0.0001**

684 [†]: observed correlation of plastic responses of CT- and selection lines would be lower compared to results of permutations

685
686

References

Alvarez, M., Schrey, A. W., & Richards, C. L. (2015). Ten years of transcriptomics in wild
populations: What have we learned about their ecology and evolution? *Molecular Ecology*,
24(4), 710–725. doi:10.1111/mec.13055
Barton, M., Sunnucks, P., Norgate, M., Murray, N., & Kearney, M. (2014). Co-gradient variation in
growth rate and development time of a broadly distributed butterfly. *PLoS ONE*, *9*(4), 1–8.
doi:10.1371/journal.pone.0095258
Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software*, *67*(1), 1–48. doi:10.18637/jss.v067.i01
Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and
powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B*
(Methodological), 289–300.
Carroll, S. B. (2005). Evolution at Two Levels: On Genes and Form. *PLoS Biology*, *3*(7), e245.
doi:10.1371/journal.pbio.0030245
Chevin, L. M., Lande, R., & Mace, G. M. (2010). Adaptation, plasticity, and extinction in a
changing environment: Towards a predictive theory. *PLoS Biology*, *8*(4).
doi:10.1371/journal.pbio.1000357
Clemson, A. S., Sgrò, C. M., & Telonis-Scott, M. (2016). Thermal plasticity in Drosophila
melanogaster populations from eastern Australia: quantitative traits to transcripts. *Journal*
of Evolutionary Biology, *29*(12), 2447–2463. doi:10.1111/jeb.12969
Conover, D. O., Duffy, T. A., & Hice, L. A. (2009). The covariance between genetic and
environmental influences across ecological gradients: Reassessing the evolutionary
significance of countergradient and cogradient variation. *Annals of the New York Academy*
of Sciences, *1168*(1), 100–129. doi:10.1111/j.1749-6632.2009.04575.x

Crispo, E. (2007). The Baldwin effect and genetic assimilation: Revisiting two mechanisms of
713   evolutionary change mediated by phenotypic plasticity. *Evolution*, *61*(11), 2469–2479.
714   doi:10.1111/j.1558-5646.2007.00203.x

715   Dayan, D. I., Crawford, D. L., & Oleksiak, M. F. (2015). Phenotypic plasticity in gene expression
716   contributes to divergence of locally adapted populations of *Fundulus heroclitus*.
717   *Molecular Ecology*, *24*(13), 3345–3359. doi:10.1111/mec.13188

718   de Jong, G. (2005). Evolution of phenotypic plasticity: patterns of plasticity and the emergence of
719   ecotypes. *New Phytologist*, *166*(2003), 101–118. doi:10.1111/j.1469-8137.2005.01322.x

720   de Jong, Gerdien. (2005). Evolution of phenotypic plasticity: patterns of plasticity and the
721   emergence of ecotypes. *New Phytologist*, *166*(1), 101–118. doi:10.1111/j.1469-
722   8137.2005.01322.x

723   Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., … Gingeras, T. R. (2013).
724   STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.

725   Draghi, J. A., & Whitlock, M. C. (2012). Phenotypic plasticity facilitates mutational variance,
726   genetic variance, and evolvability along the major axis of environmental variation.
727   *Evolution*, *66*(9), 2891–2902. doi:10.1111/j.1558-5646.2012.01649.x

728   Fangue, N. A., Hofmeister, M., & Schulte, P. M. (2006). Intraspecific variation in thermal
729   tolerance and heat shock protein gene expression in common killifish, *Fundulus*
730   *heteroclitus*. *The Journal of Experimental Biology*, *209*(Pt 15), 2859–72.
731   doi:10.1242/jeb.02260

732   Feder, M. E., & Hoffman, G. E. (1999). Heat-shock proteins, molecular chaperones, and the stress
733   response: evolutionary and ecological physiology. *Annual Review of Physiology*, *61*, 243–
734   282.

735   Feder, M. E., & Krebs, R. A. (1998). Natural and Genetic Engineering of the Heat-Shock Protein
736   Hsp70 in Drosophila melanogaster : Consequences for Thermotolerance. *American*
737   *Zoologist*, *38*(3), 503–517. doi:10.1093/icb/38.3.503

738   Fierst, J. L. (2011). A history of phenotypic plasticity accelerates adaptation to a new
739   environment. *Journal of Evolutionary Biology*, *24*(9), 1992–2001. doi:10.1111/j.1420-
740   9101.2011.02333.x

741   Fitzpatrick, B. M. (2012). Underappreciated consequences of phenotypic plasticity for ecological
742   speciation. *International Journal of Ecology*, *2012*, 32–37. doi:10.1155/2012/256017

743   Forsman, A. (2015). Rethinking phenotypic plasticity and its consequences for individuals,
744   populations and species. *Heredity*, *115*(4), 276–284. doi:10.1038/hdy.2014.92

745   Garland, T., & Kelly, S. A. (2008). Phenotypic plasticity and experimental evolution. *Journal of*
746   *Experimental Biology*, *211*(16), 2725–2725. doi:10.1242/jeb.022673

747   Ghalambor, C. K., Hoke, K. L., Ruell, E. W., Fischer, E. K., Reznick, D. N., & Hughes, K. A. (2015).
748   Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature.
749   *Nature*, *525*(7569), 372–375. doi:10.1038/nature15256

750   Ghalambor, C. K., McKay, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus non-
751   adaptive phenotypic plasticity and the potential for contemporary adaptation in new
752   environments. *Functional Ecology*, *21*(3), 394–407. doi:10.1111/j.1365-2435.2007.01283.x

753   Gibbons, T. C., Metzger, D. C. H., Healy, T. M., & Schulte, P. M. (2017). Gene expression plasticity
754   in response to salinity acclimation in threespine stickleback ecotypes from different salinity
755   habitats. *Molecular Ecology*, *26*(10), 2711–2725. doi:10.1111/mec.14065

756   Gibson, G. (2008). The environmental contribution to gene expression profiles. *Nature Reviews*
757   *Genetics*, *9*(8), 575.

758   Grether, G. F. (2005). Environmental Change, Phenotypic Plasticity, and Genetic Compensation.
759   *The American Naturalist*, *166*(4), E115–E123. doi:10.1086/432023

760   Guo, J., Liu, R., Huang, L., Zheng, X. M., Liu, P. L., Du, Y. S., … Ge, S. (2016). Widespread and

adaptive alterations in genome-wide gene expression associated with ecological divergence of two oryza species. *Molecular Biology and Evolution*, *33*(1), 62–78. doi:10.1093/molbev/msv196

Hasan, M. M., DeFaveri, J., Kuure, S., Dash, S. N., Lehtonen, S., Merilä, J., & McCairns, R. J. S. (2017). Sticklebacks adapted to divergent osmotic environments show differences in plasticity for kidney morphology and candidate gene expression. *The Journal of Experimental Biology*, *220*(12), 2175–2186. doi:10.1242/jeb.146027

Hatakeyama, M., Opitz, L., Russo, G., Qi, W., Schlapbach, R., & Rehrauer, H. (2016). SUSHI: An exquisite recipe for fully documented, reproducible and reusable NGS data analysis. *BMC Bioinformatics*, *17*(1), 1–9. doi:10.1186/s12859-016-1104-8

Ho, W.-C., & Zhang, J. (2018). Evolutionary adaptations to new environments generally reverse plastic phenotypic changes. *Nature Communications*, *9*(1), 1–11. doi:10.1038/s41467-017-02724-5

Ho, W.-C., & Zhang, J. (2019). Genetic gene expression changes during environmental adaptations tend to reverse plastic changes even after the correction for statistical nonindependence. *Molecular Biology and Evolution*. doi:10.1093/molbev/msz002

Hodgins-Davis, A., & Townsend, J. P. (2009). Evolving gene expression: from G to E to G × E. *Trends in Ecology and Evolution*, *24*(12), 649–658. doi:10.1016/j.tree.2009.06.011

Huang, Y., & Agrawal, A. F. (2016). Experimental Evolution of Gene Expression and Plasticity in Alternative Selective Regimes. *PLoS Genetics*, *12*(9), 1–23. doi:10.1371/journal.pgen.1006336

Krebs, R. A., & Feder, M. E. (1997). Natural Variation in the Expression of the Heat-Shock Protein HSP70 in a Population of Drosophila melanogaster and Its Correlation with Tolerance of Ecologically Relevant Thermal Stress. *Evolution*, *51*(1), 173. doi:10.2307/2410970

Kregel, K. C. (2002). Invited Review: Heat shock proteins: modifying factors in physiological stress responses and acquired thermotolerance. *Journal of Applied Physiology*, *92*(5), 2177–2186. doi:10.1152/japplphysiol.01267.2001

Kültz, D. (2005). Molecular and Evolutionary Basis of the Cellular Stress Response. *Annual Review of Physiology*, *67*(1), 225–257. doi:10.1146/annurev.physiol.67.040403.103635

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13), 1–26. doi:10.18637/jss.v082.i13

Lande, R. (2009). Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. *Journal of Evolutionary Biology*, *22*(7), 1435–1446. doi:10.1111/j.1420-9101.2009.01754.x

Laugen, A. T., Laurila, A., Räsänen, K., & Merilä, J. (2003). Latitudinal countergradient variation in the common frog (Rana temporaria) development rates – evidence for local adaptation. *J. Evol.*, *16*(5), 996–1005.

Law, C. W., Chen, Y., Shi, W., & Smyth, G. K. (2014). voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biology*, *15*(2), R29. doi:10.1186/gb-2014-15-2-r29

Lenth, R. V. (2016). Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software*, *69*(1), 1–33. doi:10.18637/jss.v069.i01

Levine, M. T., Eckert, M. L., & Begun, D. J. (2011). Whole-genome expression plasticity across tropical and temperate *Drosophila melanogaster* populations from eastern Australia. *Molecular Biology and Evolution*, *28*(1), 249–256. doi:10.1093/molbev/msq197

Levis, N. A., & Pfennig, D. W. (2016). Evaluating ' Plasticity-First ' Evolution in Nature : Key Criteria and Empirical Approaches. *Trends in Ecology & Evolution*, *31*(7), 563–574. doi:10.1016/j.tree.2016.03.012

Li, A., Li, L., Song, K., Wang, W., & Zhang, G. (2017). Temperature, energy metabolism, and
adaptive divergence in two oyster subspecies. *Ecology and Evolution*, *7*(16), 6151–6162.
doi:10.1002/ece3.3085
Liao, Y., Smyth, G. K., & Shi, W. (2014). FeatureCounts: An efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–930.
doi:10.1093/bioinformatics/btt656
Loeschcke, V., & Hoffmann, A. a. (2007). Consequences of heat hardening on a field fitness
component in Drosophila depend on environmental temperature. *American Naturalist*,
169(2), 175–183. doi:10.1086/510632
Lohman, B. K., Stutz, W. E., & Bolnick, D. I. (2017). Gene expression stasis and plasticity following
migration into a foreign environment. *Molecular Ecology*, *26*(18), 4657–4670.
doi:10.1111/mec.14234
Mahroof, R., Zhu, K. Y., & Subramanyam, B. (2005). Changes in Expression of Heat Shock Proteins
in Tribolium castaneum (Coleoptera: Tenebrionidae) in Relation to Developmental Stage,
Exposure Time, and Temperature. *Annals of the Entomological Society of America*, *98*(1),
100–107. doi:10.1603/0013-8746(2005)098[0100:CIEOHS]2.0.CO;2
Mäkinen, H., Papakostas, S., Vøllestad, L. A., Leder, E. H., & Primmer, C. R. (2016). Plastic and
evolutionary gene expression responses are correlated in European grayling (*Thymallus*
thymallus) subpopulations adapted to different thermal environments. In *Journal of*
Heredity (Vol. 107, pp. 82–89). doi:10.1093/jhered/esv069
Mallard, F., Jakšić, A. M., & Schlötterer, C. (2018). Contesting the evidence for non-adaptive
plasticity. *Nature*, *555*(7698), E21.
McCairns, R. J. S. S., & Bernatchez, L. (2009). Adaptive divergence between freshwater and
marine sticklebacks: insights into the role of phenotypic plasticity from an integrated
analysis of candidate gene expression. *Evolution*, *64*(4), 1029–1047. doi:10.1111/j.1558-
5646.2009.00886.x
Milutinović, B., Stolpe, C., Peuß, R., Armitage, S. A. O., & Kurtz, J. (2013). The Red Flour Beetle as
a Model for Bacterial Oral Infections. *PLoS ONE*, *8*(5). doi:10.1371/journal.pone.0064638
Molina-Montenegro, M. A., Peñuelas, J., Munné-Bosch, S., & Sardans, J. (2012). Higher plasticity
in ecophysiological traits enhances the performance and invasion success of Taraxacum
officinale (dandelion) in alpine environments. *Biological Invasions*, *14*(1), 21–33.
doi:10.1007/s10530-011-0055-2
Morris, M. R. J., Richard, R., Leder, E. H., Barrett, R. D. H., Aubin-Horth, N., & Rogers, S. M.
(2014). Gene expression plasticity evolves in response to colonization of freshwater lakes in
threespine stickleback. *Molecular Ecology*, *23*(13), 3226–3240. doi:10.1111/mec.12820
Narum, S. R., & Campbell, N. R. (2015). Transcriptomic response to heat stress among
ecologically divergent populations of redband trout. *BMC Genomics*, *16*(1), 103.
doi:10.1186/s12864-015-1246-5
Nussey, D. H., Postma, E., Gienapp, P., & Visser, M. E. (2005). Selection on heritable phenotypic
plasticity in a wild bird population. *Science*, *310*(5746), 304–306.
doi:10.1126/science.1117004
Passow, C. N., Henpita, C., Shaw, J. H., Quackenbush, C. R., Warren, W. C., Schartl, M., … Tobler,
M. (2017). The roles of plasticity and evolutionary change in shaping gene expression
variation in natural populations of extremophile fish. *Molecular Ecology*, *26*(22), 6384–
6399. doi:10.1111/mec.14360
Pavey, S. A., Collin, H., Nosil, P., & Rogers, S. M. (2010). The role of gene expression in ecological
speciation. *Annals of the New York Academy of Sciences*, *1206*, 110–129.
doi:10.1111/j.1749-6632.2010.05765.x
Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., & Moczek, A. P.

(2010). Phenotypic plasticity's impacts on diversification and speciation. *Trends in Ecology*
and Evolution, *25*(8), 459–467. doi:10.1016/j.tree.2010.05.006

Pichancourt, J. B., & van Klinken, R. D. (2012). Phenotypic plasticity influences the size, shape and
dynamics of the geographic distribution of an invasive plant. *PLoS ONE*, *7*(2), e32323.
doi:10.1371/journal.pone.0032323

Pigliucci, M., Murren, C. J., & Schlichting, C. D. (2006). Phenotypic plasticity and evolution by
genetic assimilation. *Journal of Experimental Biology*, *209*(12), 2362–2367.
doi:10.1242/jeb.02070

Pigliucci, Massimo. (2005). Evolution of phenotypic plasticity: Where are we going now? *Trends*
in Ecology and Evolution, *20*(9), 481–486. doi:10.1016/j.tree.2005.06.001

Price, T. D., Qvarnstrom, A., & Irwin, D. E. (2003). The role of phenotypic plasticity in driving
genetic evolution. *Proceedings of the Royal Society B: Biological Sciences*, *270*(1523), 1433–
1440. doi:10.1098/rspb.2003.2372

Price, Trevor D., Qvarnström, A., & Irwin, D. E. (2003). The role of phenotypic plasticity in driving
genetic evolution. *Proceedings of the Royal Society B: Biological Sciences*, *270*(1523), 1433–
1440. doi:10.1098/rspb.2003.2372

R Core Team. (2017). R Core Team (2017). R: A language and environment for statistical
computing. *R Foundation for Statistical Computing, Vienna, Austria. URL Http://Www.R-*
Project.Org/.

Ragland, G. J., Almskaar, K., Vertacnik, K. L., Gough, H. M., Feder, J. L., Hahn, D. A., & Schwarz, D.
(2015). Differences in performance and transcriptome-wide gene expression associated
with Rhagoletis (Diptera: Tephritidae) larvae feeding in alternate host fruit environments.
Molecular Ecology, *24*(11), 2759–2776. doi:10.1111/mec.13191

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic*
Acids Research, *43*(7), e47–e47. doi:10.1093/nar/gkv007

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics*, *26*(1), 139–
140. doi:10.1093/bioinformatics/btp616

Romero, I. G., Ruvinsky, I., & Gilad, Y. (2012). Comparative studies of gene expression and the
evolution of gene regulation. *Nature Reviews Genetics*, *13*(7), 505–516.
doi:10.1038/nrg3229

Schaum, E., Rost, B., Millar, A. J., & Collins, S. (2013). Variation in plastic responses of a globally
distributed picoplankton species to ocean acidification. *Nature Climate Change*, *3*(3), 298–
302. doi:10.1038/nclimate1774

Scheiner, S. M. (1993). Genetics and Evolution of Phenotypic Plasticity. *Annual Review of Ecology*
and Systematics, *24*(1), 35–68. doi:10.1146/annurev.es.24.110193.000343

Schwenke, R. A., Lazzaro, B. P., & Wolfner, M. F. (2016). Reproduction–Immunity Trade-Offs in
Insects. *Annual Review of Entomology*, *61*(1), 239–256. doi:10.1146/annurev-ento-010715-
023924

Sikkink, K. L., Reynolds, R. M., Ituarte, C. M., Cresko, W. A., & Phillips, P. C. (2014). Rapid
Evolution of Phenotypic Plasticity and Shifting Thresholds of Genetic Assimilation in the
Nematode Caenorhabditis remanei. *G3: Genes|Genomes|Genetics*, *4*(6), 1103–1112.
doi:10.1534/g3.114.010553

Silbermann, R., & Tatar, M. (2000). Reproductive costs of heat shock protein in transgenic
Drosophila melanogaster. *Evolution*, *54*(6), 2038–2045. doi:10.1111/j.0014-
3820.2000.tb01247.x

Smyth, G. K., Michaud, J., & Scott, H. S. (2005). Use of within-array replicate spots for assessing
differential expression in microarray experiments. *Bioinformatics*, *21*(9), 2067–2075.

doi:10.1093/bioinformatics/bti270

Sokoloff, A. (1972). *The biology of Tribolium with special emphasis on genetic aspects I.*
Clarendon Press and Oxford Univ. Press, Oxford. doi:10.1086/407731

Sokolova, I. M. (2013). Energy-limited tolerance to stress as a conceptual framework to integrate
the effects of multiple stressors. *Integrative and Comparative Biology*, *53*(4), 597–608.
doi:10.1093/icb/ict028

Sørensen, J G, Dahlgaard, J., & Loeschcke, V. (2001). Genetic variation in thermal tolerance
among natural populations of *Drosophila buzzatii*: down regulation of Hsp70 expression
and variation in heat stress resistance traits. *Functional Ecology*, *15*(3), 289–296.
doi:10.1046/j.1365-2435.2001.00525.x

Sørensen, Jesper Givskov, Kristensen, T. N., & Loeschcke, V. (2003). The evolutionary and
ecological role of heat shock proteins. *Ecology Letters*, *6*(11), 1025–1037.
doi:10.1046/j.1461-0248.2003.00528.x

Stoks, R., Govaert, L., Pauwels, K., Jansen, B., & De Meester, L. (2016). Resurrecting complexity:
The interplay of plasticity and rapid evolution in the multiple trait response to strong
changes in predation pressure in the water flea Daphnia magna. *Ecology Letters*, *19*(2),
180–190. doi:10.1111/ele.12551

Townsend, J. P., Cavalieri, D., & Hartl, D. L. (2003). Population genetic variation in genome-wide
gene expression. *Molecular Biology and Evolution*, *20*(6), 955–963.
doi:10.1093/molbev/msg106

Via, S. (1993). Adaptive Phenotypic Plasticity: Target or By-Product of Selection in a Variable
Environment? *The American Naturalist*, *142*(2), 352–365. doi:10.1086/285542

von Heckel, K., Stephan, W., & Hutter, S. (2016). Canalization of gene expression is a major
signature of regulatory cold adaptation in temperate Drosophila melanogaster. *BMC*
Genomics, *17*(1), 1–14. doi:10.1186/s12864-016-2866-0

Whitehead, A., & Crawford, D. L. (2006). Neutral and adaptive variation in gene expression.
Proceedings of the National Academy of Sciences of the United States of America\r, *103*(14),
5425–5430. doi:10.1073/pnas.0507648103

Wray, G. A. (2007). The evolutionary significance of cis-regulatory mutations. *Nature Reviews*
Genetics, *8*(3), 206–216. doi:10.1038/nrg2063

Wund, M. A. (2012). Assessing the impacts of phenotypic plasticity on evolution. *Integrative and*
Comparative Biology, *52*(1), 5–15. doi:10.1093/icb/ics050

Yampolsky, L. Y., Glazko, G. V., & Fry, J. D. (2012). Evolution of gene expression and expression
plasticity in long-term experimental populations of *Drosophila melanogaster* maintained
under constant and variable ethanol stress. *Molecular Ecology*, *21*(17), 4287–4299.
doi:10.1111/j.1365-294X.2012.05697.x

Yampolsky, L. Y., Zeng, E., Lopez, J., Williams, P. J., Dick, K. B., Colbourne, J. K., & Pfrender, M. E.
(2014). Functional genomics of acclimation and adaptation in response to thermal stress in
Daphnia. *BMC Genomics*, *15*(1), 1–12. doi:10.1186/1471-2164-15-859

Yeh, P. J., & Price, T. D. (2004). Adaptive Phenotypic Plasticity and the Successful Colonization of
a Novel Environment. *The American Naturalist*, *164*(4), 531–542. doi:10.1086/423825

Zhao, L., & Jones, W. A. (2012). Expression of heat shock protein genes in insect stress responses.
Invertebrate Survival Journal, *9*(1), 93–101.