Tensor decomposition-Based Unsupervised Feature Extraction Applied to Single-Cell Gene Expression Analysis

Y-h. Taguchi [1,*]**, Turki Turki**[2]

[1]*Department of Physics, Chuo University, Tokyo 112-8551, Japan*

[2] *King Abdulaziz University, Department of Computer Science, Jeddah 21589, Saudi Arabia*

Correspondence*:
Corresponding Author
tag@granular.com

ABSTRACT

Although single cell RNA sequencing (scRNA-seq) technology is newly invented and promising one, because of lack of enough information that labels individual cells, it is hard to interpret the obtained gene expression of each cell. Because of this insufficient information available, unsupervised clustering, e.g., t-Distributed Stochastic Neighbor Embedding and Uniform Manifold Approximation and Projection , is usually employed to obtain low dimensional embedding that can help to understand cell-cell relationship. One possible drawback of this strategy is that the outcome is highly dependent upon genes selected for the usage of clustering. In order to fulfill this requirement, there are many methods that performed unsupervised gene selection. In this study, a tensor decomposition (TD) based unsupervised feature extraction (FE) was applied to the integration of two scRNA-seq expression profiles that measure human and mouse midbrain development. TD based unsupervised FE could not only select coincident genes between human and mouse, but also biologically reliable genes. Coincidence between two species as well as biological reliability of selected genes is increased compared with principal component analysis (PCA) based FE applied to the same data set in the previous study. Since PCA based unsupervised FE outperformed other three popular unsupervised gene selection methods, highly variable genes, bimodal genes and dpFeature, TD based unsupervised FE can do so as well. In addition to this, ten transcription factors (TFs) that might regulate selected genes and might contribute to midbrain development are identified. These ten TFs, BHLHE40, EGR1, GABPA, IRF3, PPARG, REST, RFX5, STAT3, TCF7L2, and ZBTB33, were previously reported to be related to brain functions and diseases. TD based unsupervised FE is a promising method to integrate two scRNA-seq profiles effectively.

Keywords: tensor decomposition, enrichment analysis, sngle cell RNA-sequencing, midbrain development, inter-species analysis

1 INTRODUCTION

Single cell RNA sequencing (scRNA-seq) (17) is a newly invented technology that enables us to measure amount of RNA in single cell basis. In spite of its promising potential, it is not easy to interpret the

measurements. The primary reason of this difficulty is the lack of sufficient information that characterizes
individual cells. In contrast to the huge number of cells measured, which is often as many as several
thousands, the number of labeling is limited, e.g., measurement of conditions as well as the amount of
expression of key genes measured by fluorescence-activated cell sorting, whose number is typically as little
as tens. This prevents us from selecting genes that characterize the individual cell properties.

In order to deal with samples without suitable numbers of labelling, unsupervised method is frequently
used, since it does not make use of labeling information directly. K-means clustering as well as hierarchical
clustering are the popular methodology that are often applied to gene expression analysis. The popular
clustering methods specifically applied to scRNA-seq is tSNE (52) or UMAP (9), which is known to be
useful to get low dimensional embedding of a set of cells. In spite of that, the obtained clusters are highly
dependent upon genes used for clustering. Thus, the next issue is, without labeling (i.e., pre-knowledge), to
select genes that might be biologically meaningful.

The various unsupervised gene selection methods applicable to scRNA-seq were invented, e.g., highly
variable genes, bimodal genes and dpFeature and principal component analysis (PCA) based unsupervised
feature extraction (FE) (50, 34, 49, 42, 25, 31, 27, 40, 47, 4, 5, 44, 45, 13, 12, 11, 51, 39, 43, 22, 23, 24,
26, 41, 48). Chen et al. (2) recently compared genes selected by these methods and concluded that the
genes selected are very diverse and have their own (unique) biological features. In this sense, it is required
to invent more advanced unsupervised gene selection methods that can select more biologically relevant
genes.

In this paper, we propose the application of tensor decomposition (TD) based unsupervised FE (37, 46, 35,
36, 30, 32, 29, 33, 28). It is an advanced method of PCA based unsupervised FE, to scRNA-seq analysis.
For more details about PCA based unsupervised FE and TD based unsupervised FE, see the recently
published book (38). Especially focusing on the integration of two scRNA-seq profiles, the advantages of
TD based unsupervised FE when compared with PCA based unsupervised FE is as follows; The former
can integrate more than two gene expressions prior to the analysis while the latter can only integrate the
results obtained by applying the method to individual data sets.

In the following, based on the previous study (34) where PCA based unsupervised FE was employed, we
try to integrate human and mouse midbrain development gene expression profiles to obtain key genes that
contribute to this process, by applying TD based unsupervised FE. It turned out that TD based unsupervised
FE can identify biologically more relevant and more common genes between human and mouse than PCA
based unsupervised FE that outperformed other methods compared with.

2 METHODS AND MATERIALS

2.1 scRNA-seq data

2.1.1 Midbrain development of humans and mice

The first scRNA-seq data used in this study was downloaded from gene expression omnibus (GEO) under
the GEO ID GSE76381; the files named "GSE76381_EmbryoMoleculeCounts.cef.txt.gz" (for human)
and "SE76381_MouseEmbryoMoleculeCounts.cef.txt.gz" (for mouse) were downloaded. These two gene
expression profiles generated from scRNA-seq data set; One represents human embryo ventral midbrain
cells between 6 and 11 weeks of gestation (287 cells for six weeks, 131 cells for seven weeks, 331 cells for
eight weeks, 322 cells for nine weeks, 509 cells for ten weeks, and 397 cells for eleven weeks, in total
1977 cells). Another is a set of mouse ventral midbrain cells at six developmental stages between E11.5

to E18.5 (349 cells for E11.5, 350 cells for E12.5, 345 cells for E13.5, 308 cells for E14.5, 356 cells for
E15.5, 142 cells for E18.5 and 57 cells for unknown, in total 1907 cells).

2.1.2 Mouse hypothalamus with and without acute formalin stress

The second scRNA-seq data used in this study were downloaded from GEO under GEOID GSE74672;
the file named "GSE74672_expressed_mols_with_classes.xlsx.gz" was downloaded. It is generated from
snRNA-seq data set that measures mouse hypothalamus with and without acute formalin stress. Various
mete-data, which is included in the first eleven rows of data set, are available. The meta-data available
includes, sex, age, cell types (astrocytes, endothelial, ependymal, microglia, neurons, oligos and vsm) and
control vs stressed samples and so on.

2.2 TD based unsupervised FE

2.2.1 Midbrain development of humans and mice

TD based unsupervised FE is recently proposed methods successfully applied to various biological
problems. TD based unsupervised FE can be used for integration of multiple measurements applied to the
common set of genes. Suppose $x_{ij} \in \mathbb{R}^{N \times M}$ and $x_{ik} \in \mathbb{R}^{N \times K}$ are the ith expression of jth and kth cells
under the two distinct conditions (in the present study, they are human and mouse), respectively. Then
3-mode tensor, $x_{ijk} \in \mathbb{R}^{N \times M \times K}$, where $N(= 13889)$ is total number of common genes between human
and mouse, which share gene symbols, between human and mouse, $M(= 1977)$ is the number of human
cells and $K(= 1907)$ is total number of mouse cells, is defined as

$$x_{ijk} = x_{ij} \cdot x_{ik}. \tag{1}$$

86 It is Case I Type II tensor (32). Here Case I means that tensor is generated such that two matrices share the
87 samples while Type II means that summation is taken over as in eq. (2). On the other hand, tensor before
88 taking summation as in eq. (1) is Tupe I. Since it is too large to be decomposed, we further transform it
89 into type II tensor as

$$x_{jk} = \sum_{i=1}^{N} x_{ijk}, \tag{2}$$

90 where $x_{jk} \in \mathbb{R}^{M \times K}$ is now not a tensor but a matrix. In this case, TD is equivalent to singular value
91 decomposition (SVD). After applying SVD to x_{jk}, we get SVD,

$$x_{jk} = \sum_{\ell=1}^{\min(M,K)} \lambda_\ell u_{\ell j} v_{\ell k}. \tag{3}$$

92 where $u_{\ell j} \in \mathbb{R}^{M \times M}$ and $v_{\ell k} \in \mathbb{R}^{K \times K}$ are singular value vectors attributed to cells of human scRNA-seq
93 and those of mouse scRNA-seq, respectively.

Singular value vectors attributed to genes of human and mouse scRNA-seq, $u_{\ell i} \in \mathbb{R}^{N \times M}$ and $v_{\ell i} \in \mathbb{R}^{N \times K}$, are defined as

$$u_{\ell i} = \sum_{j=1}^{M} u_{\ell j} x_{ij} \tag{4}$$

$$v_{\ell i} = \sum_{k=1}^{K} v_{\ell k} x_{ik} \tag{5}$$

respectively.

In order to find genes associated with biological functions, we need to select $u_{\ell j}$ and $v_{\ell k}$ which are coincident with biological meaning. In this study, we employ time points of measurements as biological meanings. In other words, we seek genes associated with time development. Since we would like to find any kind of time dependence, we simply deal with time points as un-ordered labelling. Thus, we apply categorical regression

$$u_{\ell j} = a_{\ell} + \sum_{t=1}^{T} a_{\ell t} \delta_{jt} \tag{6}$$

(T=6; t=1 to T correspond to 6, 7, 8, 9, 10 and 11 weeks, see Methods and Materials) or

$$v_{\ell k} = b_{\ell} + \sum_{t=1}^{T} b_{\ell t} \delta_{kt} \tag{7}$$

(T=7; t=1 to T correspond to E11.5, E12.5, E13.5, E14.5, F15.5, E18.5 and unknown, see Methods and Materials) where $\delta_{jt}(\delta_{kt}) = 1$ when the jth(kth) cell is taken from the tth time point otherwise $\delta_{jt}(\delta_{kt}) = 0$. $a_{\ell}, a_{\ell t}, b_{\ell}$ and $b_{\ell t}$ are the regression coefficients.

P-values are attributed to ℓth singular value vectors using the above categorical regression (`lm` function in R (15) is used to compute P-values). P-values attributed to singular value vectors are corrected by BH criterion (1). Singular value vectors associated with corrected P-values less than 0.01 are selected for the download analysis. Hereafter, the set of selected singular value vectors of human and mouse are denoted as $\Omega_{\ell}^{\text{human}}$ and $\Omega_{\ell}^{\text{mouse}}$, respectively.

P-values are attributed to genes with assuming χ^2 distribution for the gene singular value vectors, $u_{\ell i}$ and $v_{\ell i}$, corresponding to the cell singular value vectors selected by categorical regression as

$$P_i^{\text{human}} = P_{\chi^2} \left[> \sum_{\ell \in \Omega_{\ell}^{\text{human}}} \left(\frac{u_{\ell i} - \langle u_{\ell i} \rangle}{\sigma_{\ell}^{\text{human}}} \right)^2 \right] \tag{8}$$

for human genes and

$$P_i^{\text{mouse}} = P_{\chi^2} \left[> \sum_{\ell \in \Omega_{\ell}^{\text{mouse}}} \left(\frac{v_{\ell i} - \langle v_{\ell i} \rangle}{\sigma_{\ell}^{\text{mouse}}} \right)^2 \right] \tag{9}$$

114   for mouse genes, respectively. Here

$$\langle u_{\ell i}\rangle = \frac{1}{N}\sum_{i=1}^{N} u_{\ell i} \tag{10}$$

115   and

$$\langle v_{\ell i}\rangle = \frac{1}{N}\sum_{i=1}^{N} v_{\ell i}. \tag{11}$$

116    $\sigma_\ell^{\text{human}}$ and $\sigma_\ell^{\text{mouse}}$ are the standard deviations of ℓth gene singular value vectors for human and mouse
117   respectively, $\Omega_\ell^{\text{human}}$ and $\Omega_\ell^{\text{mouse}}$ are sets of ℓs selected by categorical regression for human (eq. (6))
118   and mouse (eq.(7)), respectively, and $P_{\chi^2}[> x]$ is the cumulative probability of χ^2 distribution when
119   the argument takes values larger than x. P_i^{human} and P_i^{mouse} are corrected by BH criterion and genes
120   associated with corrected P-values less than 0.01 are selected.

121   2.2.2   Mouse hypothalamus with and without acute formalin stress

The application of TD based unsupervised FE to mouse hypothalamus is quit similar to that to mouse
and human midbrain. There are also two matrices, $x_{ij} \in \mathbb{R}^{N \times M}$ and $x_{ik} \in \mathbb{R}^{N \times K}$ that correspond to ith
expression of jth and kth cells under the two distinct conditions (in the present case, they are without and
with acute formalin stress, respectively); $N = 24341$, $M = 1785$ and $K = 1096$. Type I Case II tensor, x_{jk},
was also generated using eqs. (1) and (2) and SVD was applied to x_{jk} as eq. (3). Then singular vale vectors
attributed to genes of samples without and with acute formalin stress, $u_{\ell i}$ and $v_{\ell i}$ were computed by eqs.
(4) and (5). We alos applied categorical regressions to $u_{\ell i}$ and $v_{\ell i}$, although categories considered here are
not time points but cell types. Then categorical regressions applied to $u_{\ell i}$ and $v_{\ell i}$ in mouse hypothalamus
without and with acute formalin stress are

$$u_{\ell j} = a_\ell + \sum_{s=1}^{7} a_{\ell s}\delta_{js} \tag{12}$$

$$v_{\ell k} = b_\ell + \sum_{s=1}^{7} b_{\ell s}\delta_{ks} \tag{13}$$

131   where s stands for one of seven cell types mentioned in §2.1.2 and $\delta_{js}(\delta_{ks}) = 1$ when the jth(kth) cell is
132   taken from the sth cell types otherwise $\delta_{jt}(\delta_{kt}) = 0$. Table 1 lists the number of cells in these categories.
133   The remaining procedures to select genes associated with identified cell type dependency is exactly the
134   same as those in midbrain case.

135   **2.3   Enrichment analyses**

136   Various enrichment analysis methods are performed with separately uploading selected human and mouse
137   gene symbols, or genes selected commonly between samples without and samples with acute formalin
138   stress, to Enrichr (6).

3 RESULTS

3.1 Midbrain development of humans and mice

As a result, following the procedure described in Methods and Materials, we identified 55 and 44 singular value vectors attributed to cells, $u_{\ell j}$s and $v_{\ell k}$s, for human and mouse, respectively. One possible validation of selected $u_{\ell j}$s and $v_{\ell k}$s are coincidence. Although cells measured are not related between human and mouse at all, if SVD works well, corresponding singular value vectors (i.e., $u_{\ell j}$ and $v_{\ell k}$ sharing the same ℓs) attributed to cells should share something biological, e.g., time dependence. This suggests that it is more likely that corresponding singular value vectors attributed to cells, $u_{\ell j}$ and $v_{\ell k}$, are simultaneously associated with significant P-vales computed by categorical regression. As expected, they are highly significantly correlated. Table 2 shows confusion matrix of the coincidence of selected singular value vectors between human and mouse. For human cells, only top 1907 singular value vectors among all 1977 singular value vectors are considered, since total number of singular value vectors attributed to mouse cells is 1907. selected: corrected P-values, computed with regression analysis eqs. (6) and (7), are less than 0.01, not selected: otherwise. Odds ratio is as many as 227 and P-values computed by Fisher's exact test is 1.44×10^{-44}.

Figure 1 shows the coincidence of selected singular value vectors between human and mouse. Singular value vectors with smaller ℓs, i.e., with more contributions, are more likely selected and coincident between human and mouse. This can be the side evidence that guarantees that TD based unsupervised FE successfully integrated human and mouse scRNA-seq data.

Next, we selected genes with following the procedures described in Methods and Materials. The first validation of selected genes is the coincidence between human and mouse. In my previous study (34), more number of common genes were selected by PCA based unsupervised FE than other methods compared, i.e., highly variables genes, bimodal genes and dpFeature. Table 3 shows the confusion matrix that describes the coincidence of selected genes between human and mouse. Odds ratio is as large as 133 and P-value is 0 (i.e., less than numerical accuracy), which is significantly better than coincidence of selected genes between human and mouse (53 common genes between 116 genes selected for human and 118 genes selected mouse), previously achieved by PCA based unsupervised FE (34), which outperformed other methods, i.e., highly variable genes, bimodal genes and dpFeature.

On the other hands, most of genes selected by PCA based unsupervised FE in the previous study (34) are included in the genes selected by TD based unsupervised FE in the present study. One hundred and two genes are selected by TD based unsupervised FE among 116 human genes selected by PCA based unsupervised FE in the previous study (34) while 91 genes are selected by TD based unsupervised FE among 118 mouse genes by PCA based unsupervised FE. Thus, TD based unsupervised FE is quite consistent with PCA based unsupervised FE.

Biological significance tested by enrichment analysis is further enhanced. Most remarkable advances achieved by TD based unsupervised FE is "Allen Brain Atlas", to which only downregulated genes were enriched in the previous study (34). As can be seen in Table 4, now many enrichment associated with upregulated genes. In addition to this, most of top ranked five terms are related to paraventricular, which is adjusted to midbrain. This suggests that TD based unsupervised FE successfully identified genes related to midbrain.

In addition to this, "Jensen TISSUES" (Table 5) for Embryonic_brain is highly enhanced (i.e., more significant (smaller) P-values $\sim 10^{-100}$, which were as large as 10^{-10} to 10^{-20} in the previous study (34)).

On the other hand, "ARCHS4 tissues" also strongly supports the biological reliability of selected genes
(Table 6). The term "MIDBRAIN" is enriched highly and it is top ranked for both human and mouse.

There are some brain related enrichment found in other categories, although they are not strong enough
compared with the above three. Brain related terms in "GTEx Tissue Sample Gene Expression Profiles
up" (Table 7) are also enhanced for mouse brain (top three terms are brain), although no brain terms are
enriched within top five raked terms for human (This discrepancy cannot be understood at the moment).
Contrary, brain related terms in "MGI Mammalian Phenotype 2017" (Table 8) are enhanced for human
brain (fourth and fifth ranked), although no brain terms are enriched within top five raked terms for mouse
(This discrepancy also cannot be understood at the moment). The above observations suggest that TD
based unsupervised FE could identify genes related to mouse and human embryonic midbrain.

We also uploaded selected 456 human genes and 505 mouse genes to STRING server (21) that evaluates
protein-protein interaction (PPI) enrichment. 7488 PPI are reported among 456 human genes while expected
number of PPI is as small as 3524 (P-value is less than 1×10^{-6}). 6788 PPI are reported among 505 mouse
genes while expected number of PPI is as small as 3290 (P-value is again less than 1×10^{-6}). Thus, TD
based unsupervised FE can successfully identify significantly interacting protein coding genes.

Finally, we checked if transcription factor (TF) that targets selected genes are common between human
and mouse (Table 9). These TFs are associated with adjusted P-values less than 0.01 in "ENCODE and
ChEA Consensus TFs from ChIP-X" of Enrichr. They are highly overlapped between human and mouse
(There are ten common TFs between 16 TFs found in human and 24 TFs found in mouse). Although
selected TFs are very distinct from those in the previous study (34), they are highly interrelated with each
other (see below). These TFs are up-loaded to the regnetworkweb server (8) and TF networks shown in
Figure 2 are identified. Clearly, even in partial, these TFs interact highly with each other.

We also checked if commonly selected ten TFs (bold in Table 9) are related to brains. Lack of BHLHE40
was found to result in brain malfunction (3). The function of EGR1 was found in embryonic rat brain (54).
GABPA is essential for human cognition (16). IRF3 is related to brain disease (18). PPAR, which PPARG
belongs to, is believed to be therapeutic target of neurodegenerative diseases (53). REST is a master
regulator of neurogenesis (10). RFX5 is known to be expressive in fetal brain (20). STAT3 promotes brain
metastasis (14). TCF7L2 regulated brain gene expression (19). ZBTB33 affects the mouse behaviour
through regulating brain gene expression (7). Thus all commonly selected ten TFs are related to brains.

3.2 Mouse hypothalamus with and without acute formalin stress

Although the effectiveness of the proposed strategy toward scRNA-seq is obvious in the results shown in
the previous subsection, one might wonder if it is accidental. In order to dispel such doubts, we apply TD
based unsupervised FE to yet another scRNA-seq data set: mouse hypothalamus with and without acute
formalin stress. Contrary to the data set analyzed in the previous subsection where very distant two data
sets were anaylised, the data sets analyzed here are very close to each other. Both data sets are taken from
the same tissue of mouse, hypothalamus. The only difference is if they are stressed by formalin dope or
not. The motivation why we here specifically apply TD based unsupervised FE to two close data sets is
as follows; when two data sets are too close, it might be difficult to identify which genes are commonly
altered by additional condition, in this case, the dependence upon cell types, because all genes might
behave equally between two. Thus, it is not a bad idead to check if TD based unsupervised FE can work
well not only when very distant data sets are analyzed but also very close data set is analyzed.

With following the procedure described in the materials and methods, we identified 30 and 24 singular
value vectors attributed to cells, $u_{\ell j}$s and $v_{\ell k}$s, without and with acute formalin stress, respectively. We
again applied Fisher's exact test (Table 10). Although odds ratio is ten times larger than that in Table 2,
P-value is even smaller than that in Table 2; This suggests that TD based unsupervised FE could identify
not all of genes but only limited genes as being common between two experimental conditions: without
and with stress.

Figure 3 shows the coincidence of selected singular value vectors between samples without and with
stress. Singular value vectors with smaller ℓs, i.e., with more contributions, are more likely selected and
coincident between samples without and with stress. This can be the side evidence that guarantees that TD
based unsupervised FE successfully integrated scRNA-seq data taken from samples without and with stress
with avoiding to regard that all are coincident between two samples.

Next, we selected genes with following the procedures described in Methods and Materials. The first
validation of selected genes is the coincidence between human and mouse. Table 11 shows the confusion
matrix that describes the coincidence of selected genes between samples without and with stress. Odds
ratio is as large as 270 and P-value is 0 (i.e., less than numerical accuracy). Thus as expected, TD based
unsupervised FE could not all but limited number of genes associated with cell type dependence.

Finally, we tried to evaualte if genes selected are tissue type specific, i.e., hypothalamus. We have
uploaded commonly selected 3324 genes to Enrichr. 'GTEx Tissue Sample Gene Expression Profiles up'
suggest that all five top ranked terms are brain with high significance (adjusted P-vlaues are less than
1×10^{-130}). This suggests that TD based unsupervised FE successfully limited number of genes related to
brains even using closely related samples. In order to be more specific, we checked 'Allen Brain Atlas up'
in Enrichr. Then we found that all top ranked five terms are hypothalamic (Table 13). It is interesting that
TD based unsupervised FE could successfully identify hypothalamic specific genes only using scRNA-seq
retrieved from hypothalamic. It is usually required to use data taken from other tissues in order to identify
tissue specific genes because we need to compare targeted tissues and not targeted tissues in order to
identify genes expressive specifically in target tissues. The successful identification of genes specific to
something without using the comparison with other samples was also observed previously when tumor
specific genes are tried to be identified with TD based unsupervised FE (30). In this sense, TD based
unsupervised FE methods are effective not only when genes common between two distinct conditions are
sought but also when genes common between two closely related conditions are sought. Thus, it is unlike
that the success of TD based unsupervised applied to scRNA-seq is accideintal.

4   DISCUSSIONS AND FUTURE WORK

In this study, we applied TD based unsupervised FE to the integration of scRNA-seq data sets taken from
two species: human and mouse. In the sense of identification of biologically more relevant set of genes, TD
based unsupervised FE can outperform PCA based unsupervised FE that previously (34) could outperform
three more popular methods: highly variable genes, bimodal genes, and dpFeature. Thus, it is expected that
TD based unsupervised FE can do so, too.

For the purpose of integration of two scRNA-seq data set, TD based unsupervised FE has many advantages
than other four methods, i.e., PCA based unsupervised FE, highly variable genes, bimodal genes and
dpFeature. At first, TD based unsupervised FE can integrate two scRNA-seq data set, not after the selection
of genes, but before the selection of genes. This enables us to identify more coincident genes set between

two scRNA-seq, in this study human and mouse. As a result, we could identify more coincident results
between human and mouse.

The criterion of genes selection is quite robust; they should be dependent upon time points when they are
measured. We do not have to specify how they are actually correlated with time. It is another advantage of
TD based unsupervised FE.

With applying enrichment analysis to genes selected, we can find many valuable insights about biological
process. As a result, we can identify ten key TFs that might regulate embryonic midbrain developments.
All of ten selected TFs turn out to be related to brains.

TD based unsupervised FE turn out to be quite effective to integrate two scRNA-seq data set. This method
should be applied to various scRNA-seq data sets considering broader scope of investigations.

In future work, we plan to (1) utilize the proposed TD based unsupervised FE under the transfer learning
setting; (2) extend the proposed approach to handle the data integration from multiple related tasks; and
(3) investigate the performance of the proposed approach when coupled with machine and deep learning
algorithms.

CONFLICT OF INTEREST STATEMENT

275 No conflict of interest is declared.

AUTHOR CONTRIBUTIONS

276 YHT planed the research, performed analyses, and wrote a paper. TT discussed the results and wrote a
277 paper.

FUNDING

278 It was supported by KAKENHI, 17K00417 and 19H05270 and Okawa foundation, grant number 17-10.
279 This project was also funded by the Deanship of Scientific Research (DSR) at King Abdulaziz University,
280 Jeddah, under grant no. KEP-8-611-38. The authors, therefore, acknowledge with thanks DSR for technical
281 and financial support.

ACKNOWLEDGMENTS

282 Nothing acknowldegded.

DATA AVAILABILITY STATEMENT

283 The datasets analyzed for this study can be found in the GEO.

284   https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE76381.

285   https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE74672

REFERENCES

1 .Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful
approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)* 57,
289–300

2 .Chen, B., Herring, C. A., and Lau, K. S. (2018). pyNVR: investigating factors affecting feature selection
from scRNA-seq data for lineage reconstruction. *Bioinformatics* doi:10.1093/bioinformatics/bty950

3 .Hamilton, K. A., Wang, Y., Raefsky, S. M., Berkowitz, S., Spangler, R., Suire, C. N., et al. (2018).
Mice lacking the transcriptional regulator Bhlhe40 have enhanced neuronal excitability and impaired
synaptic plasticity in the hippocampus. *PLOS ONE* 13, 1–22. doi:10.1371/journal.pone.0196223

4 .Ishida, S., Umeyama, H., Iwadate, M., and Taguchi, Y.-h. (2014). Bioinformatic Screening of
Autoimmune Disease Genes and Protein Structure Prediction with FAMS for Drug Discovery. *Protein*
Pept. Lett. 21, 828–39

5 .Kinoshita, R., Iwadate, M., Umeyama, H., and Taguchi, Y.-h. (2014). Genes associated with genotype-
specific DNA methylation in squamous cell carcinoma as candidate drug targets. *BMC Syst Biol* 8
Suppl 1, S4

6 .Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q., Wang, Z., et al. (2016).
Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Research*
44, W90–W97. doi:10.1093/nar/gkw377

7 .Kulikov, A. V., Korostina, V. S., Kulikova, E. A., Fursenko, D. V., Akulov, A. E., Moshkin, M. P., et al.
(2016). Knockout zbtb33 gene results in an increased locomotion, exploration and pre-pulse inhibition
in mice. *Behavioural Brain Research SreeTestContent1* 297, 76 – 83. doi:https://doi.org/10.1016/j.bbr.
2015.10.003

8 .Liu, Z.-P., Wu, C., Miao, H., and Wu, H. (2015). RegNetwork: an integrated database of transcriptional
and post-transcriptional regulatory networks in human and mouse. *Database* 2015. doi:10.1093/
database/bav095

9 .McInnes, L., Healy, J., and Melville, J. (2018). Umap: Uniform manifold approximation and projection
for dimension reduction ArXiv:1802.03426

10 .Mozzi, A., Guerini, F. R., Forni, D., Costa, A. S., Nemni, R., Baglio, F., et al. (2017). REST, a
master regulator of neurogenesis, evolved under strong positive selection in humans and in non human
primates. *Scientific Reports* 7. doi:10.1038/s41598-017-10245-w

11 .Murakami, Y., Kubo, S., Tamori, A., Itami, S., Kawamura, E., Iwaisako, K., et al. (2015).
Comprehensive analysis of transcriptome and metabolome analysis in Intrahepatic Cholangiocarcinoma
and Hepatocellular Carcinoma. *Sci Rep* 5, 16294

12 .Murakami, Y., Tanahashi, T., Okada, R., Toyoda, H., Kumada, T., Enomoto, M., et al. (2014).
Comparison of Hepatocellular Carcinoma miRNA Expression Profiling as Evaluated by Next
Generation Sequencing and Microarray. *PLoS ONE* 9, e106314

13 .Murakami, Y., Toyoda, H., Tanahashi, T., Tanaka, J., Kumada, T., Yoshioka, Y., et al. (2012).
Comprehensive miRNA expression analysis in peripheral blood can diagnose liver disease. *PLoS ONE*
7, e48366

14 .Priego, N., Zhu, L., Monteiro, C., Mulders, M., Wasilewski, D., Bindeman, W., et al. (2018). STAT3
labels a subpopulation of reactive astrocytes required for brain metastasis. *Nature Medicine* 24,
1024–1035. doi:10.1038/s41591-018-0044-4

15 .R Core Team (2018). *R: A Language and Environment for Statistical Computing*. R Foundation for
Statistical Computing, Vienna, Austria

16 .Reiff, R. E., Ali, B. R., Baron, B., Yu, T. W., Ben-Salem, S., Coulter, M. E., et al. (2014). METTL23, a
transcriptional partner of GABPA, is essential for human cognition. *Human Molecular Genetics* 23,
3456–3466. doi:10.1093/hmg/ddu054

17 .Sasagawa, Y., Hayashi, T., and Nikaido, I. (2019). *Strategies for Converting RNA to Amplifiable*
cDNA for Single-Cell RNA Sequencing Methods (Singapore: Springer Singapore). 1–17. doi:10.1007/
978-981-13-6037-4_1

18 .Schultz, K. L. W., Troisi, E. M., Baxter, V. K., Glowinski, R., and Griffin, D. E. (2019). Interferon
regulatory factors 3 and 7 have distinct roles in the pathogenesis of alphavirus encephalomyelitis.
Journal of General Virology 100, 46–62

19 .Shao, W., Wang, D., Chiang, Y.-T., Ip, W., Zhu, L., Xu, F., et al. (2013). The Wnt signaling
pathway effector TCF7L2 controls gut and brain proglucagon gene expression and glucose homeostasis.
Diabetes 62, 789–800. doi:10.2337/db12-0365

20 .Sugiaman-Trapman, D., Vitezic, M., Jouhilahti, E.-M., Mathelier, A., Lauter, G., Misra, S., et al.
(2018). Characterization of the human RFX transcription factor family by regulatory and target gene
analysis. *BMC Genomics* 19, 181. doi:10.1186/s12864-018-4564-6

21 .Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J., et al. (2014).
STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids*
Research 43, D447–D452. doi:10.1093/nar/gku1003

22 .Taguchi, Y.-h. (2014). Integrative analysis of gene expression and promoter methylation during
reprogramming of a non-small-cell lung cancer cell line using principal component analysis-based
unsupervised feature extraction. In *Intelligent Computing in Bioinformatics*, eds. D.-S. Huang, K. Han,
and M. Gromiha (Heidelberg: Springer International Publishing), vol. 8590 of *LNCS*. 445–455

23 .Taguchi, Y.-h. (2015). Identification of aberrant gene expression associated with aberrant promoter
methylation in primordial germ cells between E13 and E16 rat F3 generation vinclozolin lineage. *BMC*
Bioinformatics 16 Suppl 18, S16

24 .Taguchi, Y.-h. (2016). Identification of More Feasible MicroRNA-mRNA Interactions within Multiple
Cancers Using Principal Component Analysis Based Unsupervised Feature Extraction. *Int J Mol Sci*
17, E696

25 .Taguchi, Y. H. (2016). microRNA-mRNA interaction identification in Wilms tumor using principal
component analysis based unsupervised feature extraction. In *2016 IEEE 16th International Conference*
on Bioinformatics and Bioengineering (BIBE). 71–78. doi:10.1109/BIBE.2016.14

26 .Taguchi, Y.-h. (2016). Principal component analysis based unsupervised feature extraction applied to
budding yeast temporally periodic gene expression. *BioData Min* 9, 22

27 .Taguchi, Y. H. (2016). Principal component analysis based unsupervised feature extraction applied
to publicly available gene expression profiles provides new insights into the mechanisms of action of
histone deacetylase inhibitors. *Neuroepigenetics* 8, 1 – 18. doi:http://dx.doi.org/10.1016/j.nepig.2016.
10.001

28 .Taguchi, Y.-H. (2017). Identification of candidate drugs for heart failure using tensor decomposition-
based unsupervised feature extraction applied to integrated analysis of gene expression between
heart failure and DrugMatrix datasets. In *Intelligent Computing Theories and Application* (Springer
International Publishing). 517–528

29 .Taguchi, Y.-H. (2017). Identification of candidate drugs using tensor-decomposition-based unsupervised
feature extraction in integrated analysis of gene expression between diseases and DrugMatrix datasets.
Sci Rep 7, 13733

30 .Taguchi, Y.-H. (2017). One-class differential expression analysis using tensor decomposition-based
unsupervised feature extraction applied to integrated analysis of multiple omics data from 26 lung
adenocarcinoma cell lines. In *2017 IEEE 17th International Conference on Bioinformatics and
Bioengineering (BIBE)*. 131–138. doi:10.1109/BIBE.2017.00-66

31 .Taguchi, Y.-h. (2017). Principal Components Analysis Based Unsupervised Feature Extraction Applied
to Gene Expression Analysis of Blood from Dengue Haemorrhagic Fever Patients. *Sci Rep* 7, 44016

32 .Taguchi, Y.-H. (2017). Tensor decomposition-based unsupervised feature extraction applied to matrix
products for multi-view data processing. *PLoS ONE* 12, e0183933

33 .Taguchi, Y.-H. (2017). Tensor decomposition-based unsupervised feature extraction identifies candidate
genes that induce post-traumatic stress disorder-mediated heart diseases. *BMC Med. Genomics* 10, 67

34 .Taguchi, Y.-h. (2018). Principal component analysis-based unsupervised feature extraction applied to
single-cell gene expression analysis. In *Intelligent Computing Theories and Application*, eds. D.-S.
Huang, K.-H. Jo, and X.-L. Zhang (Cham: Springer International Publishing), 816–826

35 .Taguchi, Y.-H. (2018). Tensor decomposition-based unsupervised feature extraction can identify
the universal nature of sequence-nonspecific off-target regulation of mRNA mediated by microRNA
transfection. *Cells* 7, 54. doi:10.3390/cells7060054

36 .Taguchi, Y.-H. (2018). Tensor decomposition/principal component analysis based unsupervised feature
extraction applied to brain gene expression and methylation profiles of social insects with multiple
castes. *BMC Bioinformatics* 19, 99

37 .Taguchi, Y.-h. (2019). Drug candidate identification based on gene expression of treated cells using
tensor decomposition-based unsupervised feature extraction for large-scale data. *BMC Bioinformatics*
19, 388. doi:10.1186/s12859-018-2395-8

38 .Taguchi, Y.-h. (2019). *Unsupervised Feature Extraction Applied to Bioinformatics* (Switzerland:
Springer International)

39 .Taguchi, Y.-h., Iwadate, M., and Umeyama, H. (2015). Heuristic principal component analysis-based
unsupervised feature extraction and its application to gene expression analysis of amyotrophic lateral
sclerosis data sets. In *Computational Intelligence in Bioinformatics and Computational Biology
(CIBCB), 2015 IEEE Conference on*. 1–10. doi:10.1109/CIBCB.2015.7300274

40 .Taguchi, Y. H., Iwadate, M., and Umeyama, H. (2015). Principal component analysis-based
unsupervised feature extraction applied to in silico drug discovery for posttraumatic stress disorder-
mediated heart disease. *BMC Bioinformatics* 16, 139

41 .Taguchi, Y.-h., Iwadate, M., and Umeyama, H. (2016). SFRP1 is a possible candidate for epigenetic
therapy in non-small cell lung cancer. *BMC Medical Genomics* 9, 28. doi:10.1186/s12920-016-0196-3

42 .Taguchi, Y. H., Iwadate, M., Umeyama, H., and Murakami, Y. (2017). Principal component analysis
based unsupervised feature extraction applied to bioinformatics analysis. In *Computational Methods
with Applications in Bioinformatics Analysis*, eds. J. J. P. Tsai and K.-L. Ng (Singapore: World
Scientific), chap. 8. 153–182. doi:10.1142/9789813207981_0008

43 .Taguchi, Y.-h., Iwadate, M., Umeyama, H., Murakami, Y., and Okamoto, A. (2015). Heuristic principal
component analysis-aased unsupervised feature extraction and its application to bioinformatics. In *Big
Data Analytics in Bioinformatics and Healthcare*, eds. B. Wang, R. Li, and W. Perrizo. 138–162

44 .Taguchi, Y.-h. and Murakami, Y. (2013). Principal component analysis based feature extraction
approach to identify circulating microRNA biomarkers. *PLoS ONE* 8, e66714

45 .Taguchi, Y.-h. and Murakami, Y. (2014). Universal disease biomarker: can a fixed set of blood
microRNAs diagnose multiple diseases? *BMC Res Notes* 7, 581

46 .Taguchi, Y. H. and Ng, K.-L. (2018). Tensor decomposition–based unsupervised feature extraction
for integrated analysis of TCGA data on micrRNA expression and promoter methylation of genes in
ovarian cancer. In *2018 IEEE 18th International Conference on Bioinformatics and Bioengineering*
(BIBE). 195–200. doi:10.1109/BIBE.2018.00045
47 .Taguchi, Y.-h. and Okamoto, A. (2012). Principal component analysis for bacterial proteomic analysis.
In *Pattern Recognition in Bioinformatics*, eds. T. Shibuya, H. Kashima, J. Sese, and S. Ahmad
(Heidelberg: Springer International Publishing), vol. 7632 of *LNCS*. 141–152
48 .Taguchi, Y. H. and Wang, H. (2017). Genetic association between amyotrophic lateral sclerosis and
cancer. *Genes* 8, 243. doi:10.3390/genes8100243
49 .Taguchi, Y.-h. and Wang, H. (2018). Exploring microrna biomarker for amyotrophic lateral sclerosis.
International Journal of Molecular Sciences 19. doi:10.3390/ijms19051318
50 .Taguchi, Y.-h. and Wang, H. (2018). Exploring microrna biomarkers for parkinson disease from mrna
expression profiles. *Cells* 7. doi:10.3390/cells7120245
51 .Umeyama, H., Iwadate, M., and Taguchi, Y.-h. (2014). TINAGL1 and B3GALNT1 are potential
therapy target genes to suppress metastasis in non-small cell lung cancer. *BMC Genomics* 15 Suppl 9,
S2
52 .van der Maaten, L. and Hinton, G. (2008). *Journal of Machine Learning Research* 9, 2579–2605
53 .Warden, A., Truitt, J., Merriman, M., Ponomareva, O., Jameson, K., Ferguson, L. B., et al. (2016).
Localization of PPAR isotypes in the adult mouse and human brain. *Scientific Reports* 6. doi:10.1038/
srep27618
54 .Wells, T., Rough, K., and Carter, D. (2011). Transcription mapping of embryonic rat brain reveals
EGR-1 induction in SOX2+ neural progenitor cells. *Frontiers in Molecular Neuroscience* 4, 6.
doi:10.3389/fnmol.2011.00006

FIGURE CAPTIONS

Table 1. The number of cells that belong to either without and with acute formalin stress or cell types

cell types	without acute	with stress
astrocytes	135	132
endothelial	169	71
ependymal	211	145
microglia	34	14
neurons	628	270
oligos	570	431
vsm	38	33

Table 2. Confusion matrix of coincidence between selected 55 singular value vectors selected among all 1977 singular value vectors, $u_{\ell j}$, attributed to human cells and 44 singular value vectors selected among all 1907 singular value vectors, $v_{\ell k}$, attributed to mouse cells.

		human not selected	human selected
mouse	not selected	1833	12
	selected	23	32

Figure 1. Coincidence between singular value vectors shown in Table 2. Horizontal axis: singular value vector numbering ℓ. Black open circles: ℓs selected for human, blue crosses: ℓs selected for both human and mouse, red open triangles: ℓs selected for mouse. Vertical black broken lines connect ℓs selected for both human and mouse.

Table 3. Confusion matrix of coincidence between selected 456 genes for human and selected 505 genes for mouse among all 13384 common genes. Selected: corrected P-values, computed with χ^2 distribution eqs. (8) and (9), are less than 0.01, not selected: otherwise. Odds ratio is as many as 133 and P-values computed by Fisher's exact test is 0 (i.e. less than numerical accuracy).

		human	
		not selected	selected
mouse	not selected	13233	151
	selected	200	305

(A)

(B)

Figure 2. TF network identified by regnetworkweb for TFs in Table 9. (A) human, (B) mouse.

Figure 3. Coincidence between singular value vectors shown in Table 10. Horizontal axis: singular value vector numbering ℓ. Black open circles: ℓs selected for samples without stress, blue crosses: ℓs selected for both samples without and with stress, red open triangles: ℓs selected for samples with stress. Vertical black broken lines connect ℓs selected for both samples without and with stress.

Table 4. Five top ranked terms from "Allen Brain Atlas up" by Enrichr for selected 456 human genes and 505 mouse genes.

human

Term	Overlap	P-value	Adjusted P-value
Paraventricular hypothalamic nucleus, magnocellular division, medial magnocellular part	31/301	2.68×10^{-12}	2.91×10^{-9}
Paraventricular hypothalamic nucleus, magnocellular division	31/301	2.68×10^{-12}	2.91×10^{-9}
Paraventricular hypothalamic nucleus, magnocellular division, posterior magnocellular part	28/301	3.39×10^{-10}	1.47×10^{-7}
Paraventricular hypothalamic nucleus	29/301	7.02×10^{-11}	5.08×10^{-8}
paraventricular nucleus, dorsal part	27/301	1.57×10^{-9}	4.88×10^{-7}

mouse

Term	Overlap	P-value	Adjusted P-value
Paraventricular hypothalamic nucleus, magnocellular division, medial magnocellular part	31/301	4.03×10^{-11}	2.19×10^{-8}
Paraventricular hypothalamic nucleus, magnocellular division	31/301	4.03×10^{-11}	2.19×10^{-8}
Paraventricular hypothalamic nucleus, magnocellular division, posterior magnocellular part	31/301	4.03×10^{-11}	2.19×10^{-8}
lower dorsal lateral hypothalamic area	29/301	8.40×10^{-10}	3.65×10^{-7}
Paraventricular hypothalamic nucleus, magnocellular division, posterior magnocellular part, lateral zone	31/301	4.03×10^{-11}	2.19×10^{-8}

Table 5. Enrichment of Embryonic brain by "JENSEN TISSUES" in Enrichr

Term	Overlap	P-value	adjusted P-value
human			
Embryonic_brain	330/4936	3.36×10^{-104}	4.30×10^{-102}
mouse			
Embryonic_brain	366/4936	3.59×10^{-115}	4.59×10^{-113}

Table 6. Enrichment of Embryonic brain by "ARCHS4 Tissues" in Enrichr

Term	Overlap	P-value	adjusted P-value
human			
MIDBRAIN	248/2316	1.02×10^{-129}	1.11×10^{-127}
mouse			
MIDBRAIN	248/2316	1.44×10^{-99}	1.56×10^{-97}

Table 7. Five top ranked terms from "GTEx Tissue Sample Gene Expression Profiles up" by Enrichr for selected 456 human genes and 505 mouse genes. Brain related terms are asterisked.

human

Term	Overlap	P-value	Adjusted P-value
GTEX-QCQG-1426-SM-48U22_ovary_female_50-59_years	105/1165	3.56×10^{-35}	1.04×10^{-31}
GTEX-RWS6-1026-SM-47JXD_ovary_female_60-69_years	116/1574	7.96×10^{-31}	7.74×10^{-28}
GTEX-TMMY-1726-SM-4DXTD_ovary_female_40-49_years	117/1582	2.97×10^{-31}	4.33×10^{-28}
GTEX-RU72-0008-SM-46MV8_skin_female_50-59_years	94/1103	1.99×10^{-29}	1.45×10^{-26}
GTEX-R55E-0008-SM-48FCG_skin_male_20-29_years	111/1599	3.67×10^{-27}	1.78×10^{-24}

mouse

Term	Overlap	P-value	Adjusted P-value
*GTEX-WVLH-0011-R4A-SM-3MJFS_brain_male_50-59_years	139/1957	1.93×10^{-30}	5.63×10^{-27}
*GTEX-X261-0011-R8A-SM-4E3I5_brain_male_50-59_years	135/1878	5.24×10^{-30}	7.65×10^{-27}
*GTEX-T5JC-0011-R4A-SM-32PLT_brain_male_20-29_years	129/1948	3.51×10^{-25}	3.42×10^{-22}
GTEX-R55E-0008-SM-48FCG_skin_male_20-29_years	109/1599	4.93×10^{-22}	2.40×10^{-19}
GTEX-TMMY-1726-SM-4DXTD_ovary_female_40-49_years	107/1582	2.37×10^{-21}	7.69×10^{-19}

Table 8. Five top ranked terms from "MGI Mammalian Phenotype 2017" by Enrichr for selected 456 human genes and 505 mouse genes. Brain related terms are asterisked.

human

Term	Overlap	P-value	Adjusted P-value
MP:0002169_no_abnormal_phenotype_detected	82/1674	2.52×10^{-11}	5.53×10^{-8}
MP:0001262_decreased_body_weight	63/1189	3.40×10^{-10}	3.72×10^{-7}
MP:0001265_decreased_body_size	46/774	3.20×10^{-9}	$2.33 \times s10^{-6}$
*MP:0009937_abnormal_neuron_differentiation	15/106	1.81×10^{-8}	9.90×10^{-6}
*MP:0000788_abnormal_cerebral_cortex_morphology	17/145	3.64×10^{-8}	1.60×10^{-5}

mouse

Term	Overlap	P-value	Adjusted P-value
MP:0002169_no_abnormal_phenotype_detected	89/1674	$1.36 \times s10^{-11}$	3.09×10^{-8}
MP:0011091_prenatal_lethality,_complete_penetrance	27/272	1.68×10^{-9}	1.91×10^{-6}
MP:0001262_decreased_body_weight	65/1189	3.93×10^{-9}	2.97×10^{-6}
MP:0011100_preweaning_lethality,_complete_penetrance	42/674	8.55×10^{-8}	3.88×10^{-5}
MP:0001265_decreased_body_size	46/774	8.22×10^{-8}	3.88×10^{-5}

Table 9. TFs enriched in "ENCODE and ChEA Consensus TFs from ChIP-X" by Enrichr for human and mouse. Bold TFs are common.

human	BCL3, **BHLHE40**, **EGR1**, **GABPA**, **IRF3**, **PPARG**, **REST**, **RFX5**, SP1, SP2, SRF, **STAT3**, **TCF7L2**, TRIM28, TRIM28, **ZBTB33**,
mouse	**BHLHE40**, CTCF, E2F4, E2F6, **EGR1**, ESR1, ETS1, FLI1, **GABPA**, **IRF3**, NFIC, NRF1, **PPARG**, RCOR1, **REST**, **RFX5**, SPI1, **STAT3**, **TCF7L2**, USF1, USF2, YY1, **ZBTB33**, ZNF384,

Table 10. Confusion matrix of coincidence between selected 30 singular value vectors selected among all 1096 singular value vectors, $u_{\ell j}$, attributed to samples without stress and 24 singular value vectors selected among all 1096 singular value vectors, $v_{\ell k}$, attributed to samples with stress. For samples without stress, only top 1096 singular value vectors among all 1785 singular value vectors are considered, since total number of singular value vectors attributed to samples without stress is 1096. selected: corrected P-values, computed with regression analysis eqs. (12) and (13), are less than 0.01, not selected: otherwise. Odds ratio is as many as 2483 and P-values computed by Fisher's exact test is 1.92×10^{-40}.

		with stress	
		not selected	selected
without stress	not selected	1065	1
	selected	7	23

Table 11. Confusion matrix of coincidence between selected 4150 genes for samples without stress and selected 3621 genes for samples with stress among all 24341 genes. Selected: corrected P-values, computed with χ^2 distribution that correspond to eqs. (8) and (9) in human and mouse midbrain study, are less than 0.01, not selected: otherwise. Odds ratio is as many as 270 and P-values computed by Fisher's exact test is 0 (i.e. less than numerical accuracy).

		with stress	
		not selected	selected
without stress	not selected	19894	297
	selected	826	3324

Table 12. Five top ranked terms from 'GTEx Tissue Sample Gene Expression Profiles up' by Enrichr for 3324 genes selected commonly between samples without and with stress.

Term	Overlap	P-value	Adjusted P-value
GTEX-WWYW-0011-R10A-SM-3NB35_brain_female_50-59_years	1006/2885	2.7880×10^{-151}	8.135×10^{-148}
GTEX-T6MN-0011-R1A-SM-32QOY_brain_male_50-59_years	859/2317	2.9865×10^{-144}	4.3575×10^{-141}
GTEX-QVUS-0011-R3A-SM-3GAFD_brain_female_60-69_years	963/2759	6.8195×10^{-144}	6.6325×10^{-141}
GTEX-T2IS-0011-R3A-SM-32QPB_brain_female_20-29_years	967/2792	5.5265×10^{-142}	4.0315×10^{-139}
GTEX-WZTO-0011-R3B-SM-3NMC6_brain_male_40-49_years	991/2972	2.6805×10^{-133}	1.5645×10^{-130}

Table 13. Five top ranked terms from 'Allen Brain Atlas up' by Enrichr for 3324 genes selceted commonly between samples without and with stress.

Term	Overlap	P-value	Adjusted P-value
Paraventricular hypothalamic nucleus	120/301	3.38×10^{-22}	7.41×10^{-19}
Paraventricular hypothalamic nucleus, parvicellular division	119/301	1.15×10^{-21}	1.27×10^{-18}
Paraventricular hypothalamic nucleus, parvicellular division, medial parvicellular part, dorsal zone	117/301	1.29×10^{-20}	9.42×10^{-18}
paraventricular nucleus, cap part	116/301	4.22×10^{-20}	2.31×10^{-17}
Paraventricular hypothalamic nucleus, magnocellular division	115/301	1.36×10^{-19}	5.96×10^{-17}