Nuclear factor I/B increases in prostate cancer to support androgen receptor activation

Jagpreet S. Nanda[1], Wisam N. Awadallah[1], Sarah E. Kohrt[2], Petra Popovics[1], Justin M. M.

Cates[3], Janni Mirosevich[4], Peter E. Clark[5], Giovanna A. Giannico[3], and Magdalena M.

Grabowska[1,2,6,7*]

[1] Department of Urology, Case Western Reserve University, Cleveland, OH

[2] Department of Pharmacology, Case Western Reserve University, Cleveland, OH

[3] Department of Pathology, Microbiology, and Immunology, Vanderbilt University Medical Center,

Nashville, TN

[4] Department of Urology, Vanderbilt University Medical Center, Nashville, TN

[5] Department of Urology, Levine Cancer Center/Atrium Health, Charlotte, NC

[6] Department of Biochemistry, Case Western Reserve University, Cleveland, OH

[7] Case Comprehensive Cancer Center, Case Western Reserve University, Cleveland, OH

*Address correspondence to:

Magdalena M. Grabowska

2123 Adelbert Road

Wood Research Tower; RTG00

Cleveland, OH 44106

Phone: 216-368-5736

Email: mmg126@case.edu

Abstract:

Most prostate cancers express androgen receptor (AR), and our previous studies have focused on identifying transcription factors that interact with and modify AR function. We have shown that transcription factor nuclear factor I/B (NFIB) regulates androgen receptor (AR) activity and that its expression is decreased in the luminal cells of severe benign prostatic hyperplasia. To assess whether changes in NFIB expression are associated with prostate cancer progression, we immunostained a tissue microarray including normal, hyperplastic, prostatic intraepithelial neoplasia, primary prostatic adenocarcinoma, and castration-resistant prostate cancer tissue samples for NFIB, AR, and synaptophysin, a marker of neuroendocrine differentiation. We observed increased NFIB in the nucleus and cytoplasm of prostate cancer samples independent of Gleason score. We also observed strong NFIB staining in primary small cell prostate cancer. While increased NFIB nuclear and cytoplasmic staining were not predictive of biochemical recurrence, the ratio of cytoplasmic-to-nuclear NFIB staining was predictive of earlier biochemical recurrence once the analysis was adjusted for tumor margin status. We next assessed expression of NFIB in prostate cancer cell lines and detected several isoforms (62 kDa, 57 kDa, 49 kDa, and 39 kDa). Using nuclear and cytoplasmic fractionation, we observed NFIB predominantly in the nuclear fraction of prostate cancer cells with increased cytoplasmic expression seen in castration-resistant prostate cancer cell lines. Through transient transfection of AR and NFIB into the AR and NFIB low JEG-3 cells, followed by co-immunoprecipitation, we also observed a physical interaction between AR and NFIB. In order to understand the consequences of NFIB over-expression in prostate cancer, we generated vector or 3X-FLAG-NFIB-expressing androgen-dependent LNCaP and castration-resistant C4-2B cells (a LNCaP derivative cell line). While over-expression of NFIB did not increase AR expression, it did increase prostate specific antigen (PSA) production and PSA promoter activity. In summary, we have described the expression pattern of NFIB in prostate cancer and propose that one consequence of NFIB over-expression in AR-dependent prostate cancers is increased AR activity as measured by PSA induction.

Introduction:

Since most prostate cancers express androgen receptor (AR), even advanced prostate cancer can be treated with androgen deprivation therapy[1]. Although these patients initially respond to androgen deprivation therapy, disease progression typically occurs within 30 months as these patients develop castration-resistant prostate cancer[2]. Castration-resistant prostate cancer is treated with newer androgen deprivation therapies such as abiraterone acetate (a steroidal CYP17A1 inhibitor[3]) and enzalutamide (a non-steroidal antiandrogen[4])[5,6]. However, patients eventually progress on these second generation therapies as well and develop various castration-resistant phenotypes.

Approximately 63% of castration-resistant prostate cancers maintain AR protein expression[7]. In these tumors, AR signaling continues through multiple mechanisms including AR over-expression, increased androgen production, and induction of constitutively active AR splice variants (such as AR-V7)[8]. Prostate tumors can also escape androgen deprivation therapy by bypassing AR signaling. One mechanism is neuroendocrine differentiation, whereby cells acquire neuronal markers such synaptophysin and chromogranin A and sometimes lose AR expression and activity. Focal neuroendocrine differentiation is observed in 85% of castration-resistant prostate cancer patients[9] while 8-25%[7,10-12] of heavily treated prostate cancer patients, including those treated with abiraterone or enzalutamide, will undergo neuroendocrine differentiation to develop frank therapy induced neuroendocrine prostate cancer[11,13-19], which arises from the transdifferentiation of the adenocarcinoma[10,15,18,20-23]. Foci of neuroendocrine differentiation and therapy-induced neuroendocrine prostate cancers can be AR negative, have reduced AR activity, and be less responsive to androgen deprivation therapy[11,24]. In mouse models, neuroendocrine tumors can secrete neuronal factors, driving ligand-independent AR activation[25]. Another mechanism of androgen-independence in castration-resistant prostate cancer cells is activation of the fibroblast growth factor (FGF) pathway[7]. These "double negative" tumors lack both AR and

neuronal markers and represent 21% of enzalutamide and abiraterone resistant prostate

cancers[7].

Our previous studies have focused on identifying AR co-factors, based on the premise

that these co-factors might play critical roles in prostate cancer progression. Building on the

studies of Gao *et al*, which determined Forkhead box A1 (FOXA1) interacts with AR to drive

prostate-specific gene expression[26] and that Foxa1 is required for proper murine prostate gland

development[27] and maintenance[28], we identified NFI transcription factors (NFIA, B, C, X) as

FOXA1-interacting proteins that modulate AR-target gene expression[29]. Previous studies had

demonstrated a role for NFI family members in regulation of AR-responsive promoters[30,31], AR-

target gene expression in castration-resistant prostate cancer cells[32], and that NFI-consensus

sequences were associated with AR binding sites in prostate cancer cell lines and tissues[33]. Our

subsequent transient knockdown studies in androgen-dependent LNCaP cells demonstrated that

NFIB inhibits AR-target gene expression and that NFIB is frequently associated with AR binding

sites[34]. However, little was known about the expression pattern of NFIB in prostate cancer. We

therefore performed a tissue microarray study to define the expression of NFIB in prostate cancer

and determine whether its expression was associated with time to biochemical recurrence,

neuroendocrine differentiation, or castration-resistant disease.

Methods:

Human prostate cancer samples: Following Institutional Review Board (IRB) approval

(Vanderbilt: 140838; CRWU: 20180111), we analyzed a prostate cancer tissue microarray linked

with clinical data derived from the Urologic Outcomes Database at Vanderbilt University Medical

Center. This tissue microarray has been previously described[35]. Our cohort included samples

from 56 primary prostate cancer patients who had undergone radical prostatectomy and 8

castration-resistant prostate cancer patients who underwent tumor de-bulking procedures. The

tissue microarray included replicate 1 mm cores from prostatectomy specimens and included

normal prostatic tissue, benign prostatic hyperplasia (BPH), high-grade prostatic intraepithelial

neoplasia (HGPIN), and adenocarcinoma, Gleason patterns 3-5. The tissue microarray was

stained for NFIB (HPA003956, Sigma), AR (clone N-20, sc-816, Santa Cruz), synaptophysin

(Clone 2, BD Biosciences) as a marker of neuroendocrine differentiation, and hematoxylin and

eosin (H&E). Gleason score and immunohistochemistry staining was scored by a genitourinary

pathologist (GG). H&E stained array cores were scored as normal, BPH, HGPIN, or Gleason

pattern 3-5. Immunohistochemistry was scored by intensity (0 negative, 1 weak, 2 moderate, 3

strong) and distribution (0 negative, 1 <33%, 2 33-66%, and 3 >66%) in both nuclear and

cytoplasmic compartments of luminal cells (normal, BPH) or tumor cells (HGPIN, prostate cancer,

castration-resistant prostate cancer). A composite staining score was generated by summation of

intensity and distribution scores. For correlation analysis and staining by Gleason score, we

generated average values of duplicate cores. Unique cores and duplicate cores with differing

Gleason scores were treated as unique cores. We also generated average per patient scores,

averaging all cores to generate an average value for NFIB, AR, and synaptophysin across all

conditions. These patient scores were used to evaluate expression across disease states and

evaluate prostate cancer outcomes. Statistical analysis was performed using non-parametric

tests, detailed in the figure legends. Following IRB approval (Vanderbilt: 160346; CRWU:

20180111), we also analyzed 8 de-identified patient samples of primary small cell prostate cancer

obtained from O.U.R. Labs in 2005, with 1-3 needle biopsies available for analysis, as previously

described[36]. Images were acquired using an Olympus BX43 microscope and a SC50 5 mega

Pixel color camera and CellSens software. Immunohistochemistry images were white-balanced

using Adobe Photoshop by using a tissue-less point on the slide as the reference.

Hematoxylin and Eosin staining: Tissue microarray slides were soaked in xylenes (2 x 4

minutes), 100% ethanol (2 x 3 minutes), 95% ethanol (2 x 3 minutes), 70% ethanol (1 x 3 minutes),

and 50% ethanol (1 x 3 minutes). Slides were then washed in running tap water for 3 minutes and

dipped 10 times in de-ionized water. Slides were then placed in Harris hematoxylin (Richard Allan

Scientific) for 4 minutes, followed by 3 minutes in running tap water and 10 dips in de-ionized

water. Slides were then placed in Clarifier-2 (Richard Allan Scientific) for 45 seconds and rinsed

3 minutes in running tap water and 10 dips in de-ionized water. Rinsed slides were then placed

in Bluing Reagent (Richard Allan Scientific) for 30 seconds and again rinsed 3 minutes in running

tap water and 10 dips in de-ionized water. Slides were then placed in eosin-phloxine working

solution (780ml 95% ethanol, 40ml 1% eosin Y in distilled water, 10ml 1% phloxine B in distilled

water, and 4ml glacial acetic acid) for 2 minutes, and washed through 5 100% ethanol baths, 10

dips each before finally being dehydrated in xylenes (3 x 4 minutes). Slides were cover-slipped

with Cytoseal 60.

Immunohistochemistry: All slides and antibody stains were processed the same day, with

developing times set by positive controls. Tissue microarray slides were soaked in xylenes (2 x 5

minutes), 100% ethanol (2 x 2 minutes), 95% ethanol (2 x 2 minutes), 70% ethanol (1 x 2 minutes),

50% ethanol (1 x 2 minutes). Slides were then washed in running tap water for 3 minutes and

placed in a container with citric acid-based antigen unmasking solution (H-3300, Vector). Slides

were then placed in a pressure cooker and cooked on high pressure for 25 minutes. Following

pressure release, slides were allowed to cool for 25 minutes prior to subsequent steps. Slides

were washed in 1 X PBS (3 x 10 minutes). Endogenous peroxidase activity was blocked with a

20-minute incubation of slides in peroxidase blocking solution (2.5ml of 30% hydrogen peroxide

into 250ml methanol). Slides were again washed in 1X PBS (3 x 10 minutes). The tissue

microarray was outlined with a PAP pen, and blocking solution (50µl of horse serum in 3 ml PBS)

was added to slides for 30-minutes at room temperature. Blocking solution was removed and

primary antibodies (1:1000) were added to each slide and incubated at 4°C overnight. In the

morning, slides were washed in 1X PBS (3 x 10 minutes) before secondary antibody (1: 200,

Vector) was added for an hour at room temperature. While slides were washed in 1X PBS (3 x 10

minutes), A and B components (2 drops A, 2 drops B into 5 ml PBS, VECTASTAIN Elite ABC

HRP Kit) were combined to incubate for 30 minutes. AB solution was then added to slides and

incubated at room temperature for 60 minutes, followed by washes in 1 X PBS (3 x 10 minutes).

Slides were then developed with Liquid DAB+ Substrate Chromogen System (1 drop DAB+ per

ml of buffer, DAKO), quenched with water and counterstained as follows. Positive controls

(prostate for AR, NFIB, brain for synaptophysin) were used to set the exposure time, and all slides

in an antibody series were incubated for the same amount of time in DAB. Slides were washed

for 5 minutes in running tap water and then were placed in Harris hematoxylin (Richard Allan

Scientific) for 30 seconds, followed by 1 minute in running tap water and 10 dips in de-ionized

water. Slides were then dipped in Clarifier-2 for 3 times and rinsed 1 minute in running tap water

and 10 dips in de-ionized water. Rinsed slides were then placed in Bluing Reagent (Richard Allan

Scientific) for 10 dips and again rinsed 1 minute in running tap water and 10 dips in de-ionized

water. Slides were then dipped 10 times in 70% ethanol and dehydrated through 5 100% ethanol

baths, 10 dips each before finally being dehydrated in xylenes (3 x 4 minutes). Slides were cover-

slipped with Cytoseal 60.

Cell lines: LNCaP (clone FGC, CRL-2876), JEG-3 (HTB-36), 22RV1 (CRL-2505), PC-3 (CRL-

1435), and DU145 (HTB-81) cells were purchased from American Type Culture Collection

(ATCC). C4-2B[37] cells were provided by Drs. Ruoxiang Wu and Leland Chung (Cedars-Sinai).

LNCaP, C4-2B, and 22RV1 were cultured in 10% premium fetal bovine serum (FBS, Atlanta

Biologicals S11150) in RPMI 1640 + L-glutamine (Gibco). JEG-3 and DU145 cells were cultured

in 10% FBS in MEM + Earle's salts and L-Glutamine (Gibco). PC-3 cells were cultured in 10%

FBS in F-12K Kaighn's modification + L-Glutamine (Gibco). BHPrS-1, , NHPrE-1, and BHPrE-1

cells were provided by Dr. Simon Hayward (NorthShore Research Institute)[38,39]. BHPrS-1 cells

were maintained in RPMI supplemented with 5% FBS and NHPrE-1 and BHPrE-1 cells in

F12/DMEM 1:1 medium containing 5% FBS, 1% Insulin-Transferrin-Selenium supplement, 10

ng/ml epidermal growth factor, 0.4% bovine pituitary extract and 1% antibiotic-antimycotic mix (all

from Gibco). 3D cultures of NHPrE-1 and BHPrE-1 cells were established in 8-well chamber slides

by plating 2,000 cells/well in medium containing 2% (vol/vol) Matrigel on top of an undiluted layer

of Matrigel. These cultures were grown for 9 days and cells were released from Matrigel using

Corning Cell Recovery Solution to prepare cell lysates for western blot analysis.

Plasmids: pCMV6-Entry (vector) and pCMV6-NFIB-myc-DDK (RC231240) were purchased from

Origene. The AR construct, p5HbHAR-A, used to generate AR-V7 and AR-V9[40], was provided by

Dr. Scott Dehm (University of Minnesota). The PSA-EP-luciferase[26] construct was provided by

Dr. Robert Matusik (Vanderbilt University Medical Center) and the SV40-renilla construct was

purchased from Promega (pRL-SV40, E2231).

Western blotting: Proteins from cell lines were lysed in ice-cold RIPA buffer (120 mM NaCl, 50

mM Tris (pH 8.0), 0.5% NP-40 and 1 mM EGTA) containing 1 mM phenylmethylsulphonyl fluoride,

1 mM sodium orthovanadate, 0.1 µM aprotinin and 10 µM leupeptin and kept on ice for one hour.

Supernatants were collected by centrifugation at 13,750 rpm for 20 min at 4°C. Protein

concentration was quantified and equal amounts of protein were added to 2X SDS loading buffer

with 5% beta-mercaptoethanol. Nuclear and cytoplasmic extracts of LNCaP, C4-2B, 22RV1,

JEG-3, DU-145 and PC3 cells were prepared using NE-PER Nuclear and extraction reagents

(Thermoscientific, 78833). Equal amounts of cytoplasmic and nuclear protein extracts were

resolved on NuPAGE 4-12% Bis-Tris Gels and transferred to nitrocellulose membranes.

Membranes were blocked in 5% skimmed milk in TBS/0.1% Tween-20 (TBST) for 1 hour at room

temperature followed by incubation overnight at 4°C with indicated antibodies NFIB (HPA003956,

rabbit antibody, 1:1000 dilution, Sigma); AR (ab74272, rabbit antibody, 1:1000 dilution, Abcam);

lamin B1 (ab133741, rabbit antibody, 1:2000, Abcam); GAPDH (AM4300, mouse antibody,

1:5000, Invitrogen) and α-tubulin (ab7291, mouse antibody, 1:5000, Abcam). GAPDH was used

as cytoplasmic loading control whereas lamin B1 was nuclear loading control. Following

secondary antibody incubation, proteins of interest were visualized using ECL detection reagent

(GE Healthcare, RPN3243). Images were acquired using a Bio-Rad Chemi Doc. Molecular

weights were calculated using Bio-Rad Image Lab. Average molecular weights of NFIB were

determine from three independent experiments.

Transient transfections: 400,000 (JEG-3, LNCaP, 22RV1) or 300,000 (C4-2B) cells were plated

in 6-well plates. After overnight recovery, cells were transiently transfected with Silencer Select

Pre-Designed siRNAs either as negative control (non-targeting, Silencer Select Negative Control

No. 1 and 2 siRNA) or NFIB-targeting siRNA constructs (Thermofisher, s9494, s9496) using 30

pmol of siRNA in 9 µl of Lipofectamine RNAi Max. Cells were transfected for 72 hours, collected,

and analyzed by Western blotting as above.

Co-immunoprecipitation: $2.5 \cdot 10^6$ JEG-3 cells were seeded per plate in 10 cm dishes. After 10

hours, cells were transfected with 10 µg of empty, NFIB (variant 2), full length AR plasmid alone,

or in combination using polyethylenimine (PEI) (Polysciences, Warrington, PA, USA) at a ratio of

3:1. Cells were washed with 1X PBS 48 hours after transfection and dislodged with a cell scraper.

Cells were lysed in 500 µL of ice-cold RIPA buffer (120 mM NaCl, 50 mM Tris [pH 8.0], 0.5%

NP-40 and 1 mM EGTA) containing 1 mM phenylmethylsulphonyl fluoride, 1 mM sodium

orthovanadate, 0.1 µM aprotinin and 10 µM leupeptin and kept at 4°C on a rotospin for 1 hr.

Supernatants were collected by centrifugation at 13,750 rpm for 20 min at 4°C. 80 µL of lysate

was kept aside for protein quantification and analysis of input samples. The remainder of the

lysate was used for preclearing by incubating them with 30 µL of equilibrated Pure Proteome TM

Protein A Magnetic beads (LSKMAGA10) for 1 hour on ice. Then 4 µL of NFIB antibody from

Sigma (HPA003956) was added per precleared sample for immunoprecipitation and kept on

rotospin overnight at 4°C. The next day, 50 µL of equilibrated Protein A Magnetic beads were

added to each sample and incubated for 6 hours before washing three times with RIPA buffer.

Proteins were eluted with 60 µL of 2X SDS PAGE loading buffer with 5% beta-mercaptoethanol.

Input and bead samples were heated at 70°C for 13 min and analyzed by SDS PAGE and

immunoblotting.

Stable Expression of 3X-FLAG NFIB in LNCaP and C4-2B cells: N-terminally 3X Flag tagged

human NFIB (variant 2, NM_001190738.1) was PCR amplified from GFP-NFIB plasmid (Origene,

RG231240) and cloned into BamH1 and Sal1 sites of pBABE-puro, a retroviral vector (gift from

Hartmut Land, Jay Morgenstern, and Bob Weinberg [Addgene plasmid # 1764;

http://n2t.net/addgene:1764; RRID:Addgene_1764][41]. Clones were verified by sequencing before

use. Stable LNCaP and C42B cells lines were generated by stable expression of pBABE (vector)

or 3X-Flag NFIB by retroviral infection. Briefly, $2 \cdot 10^6$ GP2-293 cells were seeded per dish in 10

cm dishes containing DMEM and 10% FBS for 18-20 hours. Cells were then transfected with 12

µg of pBABE-puro or pBABE-puro Flag NFIB and pCMV-VSV-G plasmids at a ratio of 3:1 using

polyethylenimine (PEI) (Polysciences, Warrington, PA, USA). pCMV-VSV-G was a gift from Bob

Weinberg (Addgene plasmid # 8454; http://n2t.net/addgene:8454 ; RRID:Addgene_8454)[42].

Virus-enriched medium was collected 48 and 96 hours after transfection and centrifuged at 800 x

g for 10 min and used to infect LNCaP or C4-2B cells in 60-mm dishes with 0.5 mL of RPMI1640

medium containing 10% FBS. Two rounds of retrovirus infections were performed on 2 sequential

days in presence of 4.0 µg/mL of hexadimethrine bromide (Sigma). Cells were re-plated into 10

cm dishes 48 hours after the second infection and allowed to grow in presence of puromycin

containing RPMI 1640 with 10% FBS for 4 days before they were used for further experiments. 2

µg/mL and 4 µg/mL of puromycin was used to select stable LNCaP and C4-2B cells. Whole cell

extracts from stable cells were checked for 3X-FLAG NFIB expression by western blotting using

FLAG M2 antibody (F1804, mouse antibody, 1:3000, Sigma).

Cytokine array: LNCaP and C4-2B vector and NFIB over-expressing cell lines were grown in

complete media. Cells were collected, lysed, and whole cell extracts were prepared as above.

Whole cell extracts were then applied to the Proteome Profiler Human XL Cytokine Array Kit per

manufacturer's instructions (R&D Systems). The Proteome Profiler Human XL Cytokine Array

includes 105 cytokines blotted in duplicate. Images were acquired as above, with each cell line

(vector and NFIB) imaged at the same time. Dotblot analysis was performed using ImageJ[43-45].

First, the background was subtracted (rolling ball radius of 25 pixels) and the imaged inverted. A

circle was drawn around the positive control and integrated density was measured of each visible

dot plus two negative control areas. The difference between the integrated density of each dot

and the negative control was considered the true integrated density. For analysis by cell line, we

adjusted for membrane variability by generating a ratio of vector-to-NFIB positive controls. We

multiplied all NFIB values by this number to generate an adjusted integrated density comparable

between the cell lines. For analysis between cell lines, we divided each integrated density value

by the average positive control value for each cell line. This generated a percentage of the positive

control signal, and allowed us to compare the amount of cytokine between all four cell lines.

Graphs include an average of the two adjacent dots and are provided to aid with data

interpretation. No statistical analysis was performed.

Luciferase reporter assays: LNCaP and C4-2B vector and NFIB overexpressing cells were

plated 150,000 or 70,000 cells per well of a 24-well plate, respectively. Cells were transfected

with prostate-specific enhancer-promoter construct (PSA-EP)-luciferase[26] and SV40-renilla

constructs (transfection control) in Lipofectamine 2000 for 24 hours. Cells were then collected,

lysed, and luciferase and renilla expression was quantified using the Dual-Reporter assay system

(Promega) using a SpectraMax ID3 plate reader with a dual injection system. Luciferase data was

then normalized to renilla and then vector cells.

Statistical analysis: Statistical analysis was performed using SPSS 24 (IBM) or GraphPad Prism

and utilized non-parametric tests, with P values smaller than 0.05 considered significant. Dot plots

are overlaid with mean with standard deviation.

<u>**Results:**</u>

Our previous study of a small cohort of patients demonstrated that NFIB was frequently

lost in the luminal cells of patients with severe benign prostatic hyperplasia[34]. To determine

whether this was also true in prostate cancer patients, we stained a tissue microarray and several

core biopsies with an antibody against NFIB (Figure 1, Supplemental Figure 1A, B), which we had

previously validated in *Nfib* knockout mouse tissues[34]. In order to determine whether changes in

NFIB expression were associated with changes in AR or acquisition of neuroendocrine features,

we also stained the tissue microarray for synaptophysin. Staining was then scored by a board-

certified genitourinary pathologist (GG).

Using an average patient score, we compared the expression of NFIB, AR, and

synaptophysin in normal prostate and prostate cancer (Figure 2 A-C). NFIB was over-expressed

in the nucleus (P = 0.0259) and cytoplasm (P < 0.0001) of cancer tissues versus patient matched

normal prostate tissues. While AR was not over-expressed in the nucleus, cytoplasmic AR was

increased in prostate cancer tissues versus normal prostate (P < 0.0001). Synaptophysin staining

was limited to the cytoplasm and cell membrane, and also increased in prostate cancer tissues

(P < 0.0001).

To evaluate whether NFIB, AR, and synaptophysin expression corresponded with

Gleason grade, we generated average scores per each unique core. There was no statistically

significant difference between Gleason grade 3 versus 4 or Gleason grade 3 versus 5 for AR,

NFIB, or synaptophysin (Figure 2 D-F). Using these values, we also correlated the expression of

nuclear and cytoplasmic NFIB with AR and synaptophysin. In primary prostate cancer samples,

nuclear and cytoplasmic NFIB levels were positively correlated (Spearman rho = 0.305, P =

0.001), suggesting that NFIB expression increases in both compartments and that increasing

cytoplasmic expression is not due to a shift in cytolocalization (Supplemental Figure 2A). Nuclear

NFIB was also positively correlated with nuclear AR (Spearman rho = 0.419, P < 0.0001).

Cytoplasmic NFIB was positively correlated with nuclear AR (Spearman rho = 0.248, P = 0.006),

cytoplasmic AR (Spearman rho = 0.447, P < 0.0001), and synaptophysin (Spearman rho = 0.464,

P < 0.0001). The positive correlation of cytoplasmic NFIB with synaptophysin remained intact

whether we examined all prostate cancer tissues (primary and castration-resistant) or castration-

resistant only (Supplemental Figure 2B, C).

We also examined whether NFIB expression correlated with biochemical recurrence in 51

primary prostate cancer patients for who we had biochemical recurrence data. For these patients,

we again used the average NFIB nuclear and cytoplasmic score (Figure 3A). Nuclear and

cytoplasmic NFIB staining did not correlate with biochemical recurrence (Figure 3B, C). However,

when we examined a ratio of cytoplasmic and nuclear NFIB, we found a high ratio of cytoplasmic-

to-nuclear NFIB predicted earlier biochemical recurrence (P = 0.058; hazard ratio 2.864; 95% CI

0.966 – 8.491; Figure 3D, E), an effect that achieved statistical significance when surgical

resection margin status was accounted for in the Cox regression model (P = 0.004; hazard ratio

8.930; 95% CI 1.989 – 40.086; Figure 3F). There was no difference in cytoplasmic-to-nuclear

NFIB by Gleason grade or tumor stage (Supplemental Figure 3A)

To more fully explore NFIB expression in the normal and diseased prostate, we compared

NFIB average patient scores in unmatched patient samples, using unpaired analysis (Figure 4A,

Supplemental Figure 3 B-D). Consistent with our previous report, NFIB was lost in the nucleus of

hyperplastic areas[34], with low levels of cytoplasmic NFIB observed (Supplemental Figure 3A).

While there was no statistically significant difference between nuclear NFIB staining in primary

prostate cancer versus normal tissue or castration-resistant prostate cancer, cytoplasmic NFIB

was again increased in primary prostate cancer versus normal prostate tissue (P < 0.0001) but

not castration-resistant samples. While nuclear AR expression was unchanged, cytoplasmic AR

was decreased in normal prostate samples (P = 0.0002) and increased in castration-resistant

samples (P = 0.0038), versus primary prostate cancer (Figure 4B). Synaptophysin expression

was also significantly lower in normal prostate tissue versus prostate cancer (P = 0.007), but there

was no significant difference between primary prostate cancer and castration-resistant (Figure

4C).

Because only one of our castration-resistant prostate cancer cases satisfied our criteria

for neuroendocrine features (loss of AR, strong synaptophysin staining), we also analyzed a

limited number (n=8) of primary small cell prostate cancer core biopsies. Of these, 7 (88%)

showed strong nuclear NFIB staining (Supplemental Figure 1B). Small cell prostate cancer cells

have limited cytoplasm, so NFIB expression was not scored.

In order to evaluate whether established prostate cancer cell lines recapitulate the nuclear

and cytoplasmic expression of NFIB, we performed nuclear and cytoplasmic protein isolation from

prostate cancer cell lines. We examined androgen-dependent (LNCaP), castration-resistant (C4-

2B, 22RV1), and AR-independent (DU-145, PC3) prostate cancer cells. As a control, we used

JEG-3 cells which express limited NFIB based on previous reports and gene expression

analysis[29,46]. NFIB isoforms were found in all cell lines (Figure 5A), including benign stromal and

epithelial cells (Supplemental Figure 4A). NFIB exists as four isoforms in most prostate cancer

cell lines, at 62 kDa, 57 kDa, 49 kDa, and 39 kDa (Supplemental Figure 4B).

In order to verify which of these bands are true NFIB bands, we performed transient

transfections and over-expression studies. First, we transiently transfected PC-3 cells with NFIB

over-expression or NFIB-knockdown constructs and compared which bands were impacted

(Supplemental Figure 4C). In PC-3 cells, transient NFIB over-expression gave rise to a 53kDa

band. Importantly, all other putative NFIB isoforms increased as well (62 kDa, 57 kDa, 49 kDa

and 39 kDa), suggesting that NFIB auto-regulates its own expression. Transient transfection of

an NFIB-targeting siRNA also decreased expression of the 62kDa, 57 kDa, and 49 kDa bands,

suggesting that these bands do indeed represent NFIB. We also expanded this analysis to JEG-3,

LNCaP, C4-2B, and 22RV1 cells. In the LNCaP and C4-2B cells, transient knockdown abrogated

expression of both the 57 kDa and 49 kDa bands (Supplemental Figure 4D). Neither of these

were expressed in JEG-3 cells. In 22RV1 cells, NFIB knockdown also decreased expression of

the 62 kDa band, which was not impacted in JEG-3, LNCaP, or C4-2B cells. However, as this

band increased in response to NFIB over-expression and decreased in both 22RV1 and PC-3

cells in response to transient knockdown, we consider it an NFIB isoform. Finally, although the

39 kDa band was not decreased in response to transient knockdown in the prostate cancer cells,

its expression was limited in JEG-3 cells, and its expression increased in the NFIB-overexpressing

cell lines.

Because cytoplasmic NFIB expression in prostate cancer patient tissues was higher in

castration-resistant samples, we compared NFIB localization in LNCaP, C4-2B, and 22RV1 cells

(Figure 5B). In LNCaP cells, which are androgen-dependent, NFIB is largely in the nucleus, but

as cells progress to castration resistance in C4-2B cells, NFIB expression increases in both the

nuclear and cytoplasmic fractions. These levels are even more pronounced in the unrelated

castration-resistant prostate cancer cell line, 22RV1. Similarly, cytoplasmic AR expression

increased in both castration-resistant prostate cancer tissue and cell lines.

Since AR and NFIB are frequently co-occupying the same cellular compartment, AR and

NFIB co-occupy the same genomic regions in a prostate cancer cell line, and our previous studies

determined NFIB regulates AR-target gene expression[29,34], we evaluated whether NFIB and AR

can interact directly by transfecting NFIB and AR into JEG-3 cells, which express low levels of

NFIB. Indeed, in transient transfection co-immunoprecipitation experiments, NFIB and AR co-

immunopurify (Figure 5C).

To start elucidating the roles of NFIB in primary and castration-resistant prostate cancer,

we turned to well-defined *in vitro* models. We focused on two related cell lines: LNCaP cells are

androgen-dependent, while their bone-homing metastatic derivative C4-2B cells are castration-

resistant. We generated pBABE (vector) or 3X-FLAG-NFIB over-expressing LNCaP and C4-2B

cells (Figure 5D). Cell lines were positive for the Flag-tagged NFIB (53 kDa) and also showed

increased endogenous NFIB expression (53 kDa, 49 kDa, 39 kDa). This observation was

consistent with our previous experiments, where transient over-expression of NFIB induced

increased expression of multiple NFIB isoforms in PC-3 cells (Supplemental Figure 3C).

Importantly, only one band was detected by the Flag antibody, indicating NFIB is not degraded,

but rather exists as a series of isoforms. We did not observe any changes in AR protein expression

in these cells (Figure 5D).

To examine some of the downstream targets of NFIB over-expression, we compared

cytokine expression using proteome profiling. The arrays included 102 cytokines and secreted

proteins, including prostate specific antigen (PSA) and IL-6, which has been implicated in

neuroendocrine differentiation[47-50]. In LNCaP cells, NFIB over-expression dramatically induced

PSA, along with DPPIV, GDF-15, and IGFBP-2; angiogenin, EGF, cystatin C, resistin,

pentraxin-3, and CD71 were also induced, but to a smaller extent (Figure 6A, B). In C4-2B cells,

over-expression of NFIB induced small changes in angiogenin and EGF. Importantly, over-

expression of NFIB in LNCaP cells resulted in expression of PSA, GDF-15, and IGFBP-2

comparable to their castration-resistant derivative (Figure 6C).

To confirm that PSA expression is indeed regulated by NFIB, we performed reporter

assays in our two cell lines using a prostate specific antigen enhancer and promoter (PSA-EP[26])

element linked to luciferase. In both LNCaP and C4-2B cells, over-expression of NFIB resulted in

increased PSA-EP reporter activity (Figure 6D, E), increasing AR activity by 1.2 fold and 1.4 fold

versus parental cell lines respectively (P < 0.01).

Discussion:

The goal of our study was to define the expression of NFIB in prostate cancer with respect

to AR and synaptophysin and to begin to unravel the relationship between NFIB and prostate

cancer progression. We analyzed a tissue microarray composed of cores including normal, BPH,

prostate cancer, and castration-resistant prostate cancer tissues samples. On a matched per-

patient basis, NFIB is over-expressed in both the nuclear and cytoplasmic compartments in

primary prostate cancer versus normal prostate tissue. Importantly, increasing nuclear NFIB

expression corresponds with increasing cytoplasmic expression, suggesting that NFIB continues

to occupy the nucleus in prostate cancer. Indeed, there is no statistically significant difference

between primary and castration-resistant prostate cancer in terms of nuclear or cytoplasmic NFIB

expression.

There also was not a significant difference between NFIB expression in either

compartment when compared by Gleason pattern, enabling us to evaluate the utility of NFIB as

predictor of biochemical recurrence, independent of Gleason grade. While nuclear NFIB or

cytoplasmic NFIB expression alone could not predict biochemical recurrence, a higher ratio of

cytoplasmic-to-nuclear NFIB predicted earlier time to biochemical recurrence when we accounted

for surgical margin status in the Cox regression analysis. This observation is exciting, but a

limitation to our analysis is the sample size as well as the challenge of generating both a

cytoplasmic and nuclear score for each patient. While these limitations limit the utility of NFIB as

biomarker of biochemical recurrence, these observations do suggest NFIB plays an interesting in

prostate cancer biology.

Consistent with our human tissue data, NFIB is expressed in the nuclear and cytoplasmic

fractions of prostate cancer cells. As in castration-resistant prostate cancer, castration-resistant

prostate cancer cell lines have increased levels of NFIB in both the nuclear and cytoplasmic

compartments compared to androgen-dependent prostate cancer cells. Consistent with UniProt

predicted isoforms, we report that NFIB also exists in multiple isoforms in prostate cancer cells,

with average molecular weights of 62 kDa, 57 kDa, 49 kDa, and 39 kDa.

Our observation of multiple NFIB isoforms is novel in prostate cancer cell lines, but has

been reported in other systems. The *NFIB* gene can undergo alternative splicing, giving rise to

nine variants[51], and the presence of multiple NFIB protein isoforms is consistent with reports from

UniProt[52], where NFIB has six reported isoforms (O00712 [NFIB_HUMAN], Entry version 172 [08

May 2019]). These isoforms have molecular weights of 47,442 Da (O00712-1), 22,251 Da

(O00712-2), 53,049 Da (O00712-4), 55,181 Da (O00712-5), and 33,525 Da (O00712-6). While

these do not match up exactly with our reported bands at 62 kDa, 57 kDa, 49 kDa, and 39 kDa,

NFIB a can undergo posttranslational modification like glycosylation[53] and sumoylation[54] which

could add to these molecular weights. *In vitro* translation assays have also identified NFI-B2 and

NFI-B3, at 47 kDa and 22 kDa, respectively[55], indicating that some of these splice variants are

expressed and functional. Significantly, the NFI-B3 isoform lacks the ability to bind DNA and

regulate gene expression and acts as a dominant negative factor in the presence of NFIB, NFIC,

and NFIX[55] Unfortunately, the NFIB antibody immunogen is a sequence shared by all but the

22,251 Da (O00712-2) isoform. Therefore, we do not know the status of this isoform in our

prostate cancer cell lines or prostate cancer tissues.

Although our previous studies showed that FOXA1 bridges the AR/NFIX interaction in

HeLa cells in Fluorescent Protein Förster Resonance Energy Transfer (FP-FRET) experiments[29],

we did observe some FRET signal between AR and NFIX, albeit this signal did not achieve

statistical significance. Our exogenous co-immunoprecipitation studies now demonstrate at least

one NFI, NFIB, is capable of interacting with the full length AR. What domains are responsible for

this interaction, whether this interaction occurs with AR splice variants, and what the functional

consequences of AR and NFIB interaction entail is under investigation.

In this present study, we investigated the role of NFIB in disease progression by using

LNCaP and their castration-resistant derivative cell line, C4-2B as a model of disease

progression. Over-expression of NFIB induced expression of PSA in LNCaP cells to levels

comparable with C4-2B cells. Our data supports that this is likely due to interactions with AR, as

we observed no increase in AR protein expression but instead noted increased AR activity in an

androgen-responsive promoter assay. Over-expression of NFIB also induced expression of other

secreted proteins and cytokines associated with and/or implicated in castration-resistant prostate

cancer, such as angiogenin[56], DDPIV (CD26)[57], IGFBP-2[58,59], and PSA[60].

These studies linking NFIB with more aggressive disease are more consistent with the

literature, wherein NFIB has been recently considered a proto-oncogene[61-66]. Our early studies

indicated that transient knockdown of *NFIB* increased the expression of AR-target genes[29,34], and

we believe this difference might be due to the adaptation that NFIB over-expression induces. For

example, in mouse models of lung cancer, Nfib increases chromatin accessibility[64], potentially

through interactions with FOXA1 we have demonstrated previously[29]. NFIB has also been

implicated in controlling components of the Polycomb Repressive Complex 2 (PRC2), which

includes EZH2 and EED. Over-expression of NFIB induces expression of EZH2 in melanoma[62]

and NFIB also interacts with EED[67].

Although our studies have focused on the role of NFIB in regulating AR expression and

activity, it is likely that NFIB also has roles independent of AR. For example, in mouse models of

small cell prostate cancer or lung cancer, Nfib is regularly amplified[68,69], and over-expression of

Nfib in a transgenic model of small cell lung cancer drives aggressive disease[63]. Our small cohort

of primary small cell prostate cancer samples strongly express NFIB. What the role for NFIB in

these AR-independent tumors also remains to be determined.

In summary, our study has described the expression of NFIB, AR, and synaptophysin in

a small cohort of prostate cancer patients. We report that NFIB is over-expressed in the nuclear

and cytoplasmic fractions of prostate cancer patient tissues and cell lines. We also report that

NFIB can interact with full length AR, and over-expression of NFIB can support increased

expression of canonical AR-target gene products, like PSA.

<u>Acknowledgements:</u>

We would like to thank the prostate cancer patients included in the tissue microarray for providing

their tissues and clinical data and Dr. Derek J. Taylor's laboratory for assisting with virus

generation. We would also like to thank Dr. Robert J. Matusik for his early support of these studies

and critical feedback on the manuscript, and Daniel A. Barocas whose development of the

Urologic Outcomes Database enabled linking outcomes data with this tissue microarray[35]. This

work was supported in part by the Defense Health Program, through the Prostate Cancer

Research Program Postdoctoral Training Award, under Award No. W81XWH-14-1-0312.

Opinions, interpretations, conclusions and recommendations are those of the author and are not

necessarily endorsed by the Department of Defense or U.S. Army. In the conduct of research

utilizing recombinant DNA, the investigators adhered to NIH Guidelines for research involving

recombinant DNA molecules. These studies were also supported in part by the National Cancer

Institute of the National Institutes of Health under award number K99CA197315 and

R00CA197315 (to MMG). The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health.

Contributions:

JSN, PEC, GAG, and MMG designed the research studies. JSN, WNA, SEK, PP, and MMG

conducted the experiments. JSN, WNA, SEK, and MMG acquired and analyzed data. JMMC and

GAG performed pathological analysis of tumor sections. GAG provided the tissue microarray,

which included clinical outcomes data whose collection was overseen by DAB and PEC for the

Urologic Outcomes Database maintained at Vanderbilt University Medical Center. JM, GAG, and

MMG wrote IRB protocols to acquire tissues and data. All authors contributed to writing the

manuscript.

References:

1.   Insitute, T. N. C. *PDQ Prostate Cancer Treatment*,
<http://www.cancer.gov/cancertopics/pdq/treatment/prostate/HealthProfessional>
(2014).
2.   Salonen, A. J., Taari, K., Ala-Opas, M., Viitanen, J., Lundstedt, S. & Tammela, T. L. J.
The FinnProstate Study VII: Intermittent Versus Continuous Androgen Deprivation in

Patients With Advanced Prostate Cancer. *The Journal of Urology* **187**, 2074-2081, (2012).

3. Barrie SE, P. G., Goddard PM, Haynes BP, Dowsett M, Jarman M. Pharmacology of novel steroidal inhibitors of cytochrome P450(17) alpha (17 alphahydroxylase/C17-20 lyase). *The Journal of Steroid Biochemistry and Molecular Biology* **50**, 267-273, (1994).

4. Tran, C., Ouk, S., Clegg, N. J., Chen, Y., Watson, P. A., Arora, V., Wongvipat, J., Smith-Jones, P. M., Yoo, D., Kwon, A., Wasielewska, T., Welsbie, D., Chen, C., Higano, C. S., Beer, T. M., Hung, D. T., Scher, H. I., Jung, M. & Sawyers, C. L. Development of a Second-Generation Antiandrogen for Treatment of Advanced Prostate Cancer. *Science (New York, N.Y.)* **324**, 787-790, (2009).

5. Scher, H., Beer, T., Higano, C., Anand, A., Taplin, M., Efstathiou, E., Rathkopf, D., Shelkey, J., Yu, E., Alumkal, J., Hung, D., Hirmand, M., Seely, L., Morris, M., Danila, D., Humm, J., Larson, S., Fleisher, M. & Sawyers, C. - Antitumour activity of MDV3100 in castration-resistant prostate cancer: a phase 1-2 study. *Lancet* **375**, 1437-1446, (2010).

6. Fizazi, K., Scher, H. I., Molina, A., Logothetis, C. J., Chi, K. N., Jones, R. J., Staffurth, J. N., North, S., Vogelzang, N. J., Saad, F., Mainwaring, P., Harland, S., Goodman, O. B., Sternberg, C. N., Li, J. H., Kheoh, T., Haqq, C. M. & de Bono, J. S. Abiraterone acetate for treatment of metastatic castration-resistant prostate cancer: final overall survival analysis of the COU-AA-301 randomised, double-blind, placebo-controlled phase 3 study. *The Lancet Oncology* **13**, 983-992, (2012).

7. Bluemn, E. G., Coleman, I. M., Lucas, J. M., Coleman, R. T., Hernandez-Lopez, S., Tharakan, R., Bianchi-Frias, D., Dumpit, R. F., Kaipainen, A., Corella, A. N., Yang, Y. C., Nyquist, M. D., Mostaghel, E., Hsieh, A. C., Zhang, X., Corey, E., Brown, L. G., Nguyen, H. M., Pienta, K., Ittmann, M., Schweizer, M., True, L. D., Wise, D., Rennie, P. S., Vessella, R. L., Morrissey, C. & Nelson, P. S. Androgen Receptor Pathway-Independent Prostate Cancer Is Sustained through FGF Signaling. *Cancer Cell* **32**, 474-489.e476, (2017).

8. Vis, A. N. & Schröder, F. H. Key targets of hormonal treatment of prostate cancer. Part 1: the androgen receptor and steroidogenic pathways. *BJU International* **104**, 438-448, (2009).

9. Matei, D. V., Renne, G., Pimentel, M., Sandri, M. T., Zorzino, L., Botteri, E., De Cicco, C., Musi, G., Brescia, A., Mazzoleni, F., Tringali, V., Detti, S. & de Cobelli, O. Neuroendocrine Differentiation in Castration-Resistant Prostate Cancer: A Systematic Diagnostic Attempt. *Clin.Genitourin.Cancer.*, (2012).

10. Beltran, H., Prandi, D., Mosquera, J. M., Benelli, M., Puca, L., Cyrta, J., Marotz, C., Giannopoulou, E., Chakravarthi, B. V. S. K., Varambally, S., Tomlins, S. A., Nanus, D. M., Tagawa, S. T., Van Allen, E. M., Elemento, O., Sboner, A., Garraway, L. A., Rubin, M. A. & Demichelis, F. Divergent clonal evolution of castration-resistant neuroendocrine prostate cancer. *Nature Medicine* **22**, 298-305, (2016).

11. Aggarwal, R., Huang, J., Alumkal, J. J., Zhang, L., Feng, F. Y., Thomas, G. V., Weinstein, A. S., Friedl, V., Zhang, C., Witte, O. N., Lloyd, P., Gleave, M., Evans, C. P., Youngren, J., Beer, T. M., Rettig, M., Wong, C. K., True, L., Foye, A., Playdle, D., Ryan, C. J., Lara, P., Chi, K. N., Uzunangelov, V., Sokolov, A., Newton, Y., Beltran, H., Demichelis, F., Rubin, M. A., Stuart, J. M. & Small, E. J. Clinical and Genomic Characterization of Treatment-Emergent Small-Cell Neuroendocrine Prostate Cancer: A Multi-institutional Prospective Study. *Journal of Clinical Oncology* **0**, JCO.2017.2077.6880, (2018).

12. Aparicio, A., Logothetis, C. J. & Maity, S. N. Understanding the Lethal Variant of Prostate Cancer: Power of Examining Extremes. *Cancer Discovery* **1**, 466-468, (2011).

13.   Wang, W. & Epstein, J. I. Small Cell Carcinoma of the Prostate: A Morphologic and
Immunohistochemical Study of 95 Cases. *American Journal of Surgical Pathology* **32**,
65-71, (2008).
14.   Abrahamsson, P. A. Neuroendocrine differentiation in prostatic carcinoma. *Prostate* **39**,
135-148, (1999).
15.   di Sant'Agnese, P. A. Neuroendocrine differentiation in human prostatic carcinoma.
Hum.Pathol. **23**, 287-296, (1992).
16.   Lin, D., Wyatt, A. W., Xue, H., Wang, Y., Dong, X., Haegert, A., Wu, R., Brahmbhatt, S.,
Mo, F., Jong, L., Bell, R. H., Anderson, S., Hurtado-Coll, A., Fazli, L., Sharma, M.,
Beltran, H., Rubin, M., Cox, M., Gout, P. W., Morris, J., Goldenberg, L., Volik, S. V.,
Gleave, M. E., Collins, C. C. & Wang, Y. High Fidelity Patient-Derived Xenografts for
Accelerating Prostate Cancer Discovery and Drug Development. *Cancer Research* **74**,
1272-1283, (2014).
17.   Terry, S. & Beltran, H. The many faces of neuroendocrine differentiation in prostate
cancer progression. *Frontiers in Oncology* **4**, (2014).
18.   Mosquera, J. M., Beltran, H., Park, K., MacDonald, T. Y., Robinson, B. D., Tagawa, S.
T., Perner, S., Bismar, T. A., Erbersdobler, A., Dhir, R., Nelson, J. B., Nanus, D. M. &
Rubin, M. A. Concurrent AURKA and MYCN gene amplifications are harbingers of lethal
treatment-related neuroendocrine prostate cancer. *Neoplasia* **15**, 1-10, (2013).
19.   Beltran, H., Tagawa, S. T., Park, K., MacDonald, T., Milowsky, M. I., Mosquera, J. M.,
Rubin, M. A. & Nanus, D. M. Challenges in Recognizing Treatment-Related
Neuroendocrine Prostate Cancer. *Journal of Clinical Oncology* **30**, e386-e389, (2012).
20.   Williamson, S. R., Zhang, S., Yao, J. L., Huang, J., Lopez-Beltran, A., Shen, S.,
Osunkoya, A. O., MacLennan, G. T., Montironi, R. & Cheng, L. ERG–TMPRSS2
rearrangement is shared by concurrent prostatic adenocarcinoma and prostatic small
cell carcinoma and absent in small cell carcinoma of the urinary bladder: evidence
supporting monoclonal origin. *Modern pathology : an official journal of the United States
and Canadian Academy of Pathology, Inc* **24**, 1120-1127, (2011).
21.   Beltran, H., Rickman, D. S., Park, K., Chae, S. S., Sboner, A., MacDonald, T. Y., Wang,
Y., Sheikh, K. L., Terry, S. p., Tagawa, S. T., Dhir, R., Nelson, J. B., de la Taille, A.,
Allory, Y., Gerstein, M. B., Perner, S., Pienta, K. J., Chinnaiyan, A. M., Wang, Y., Collins,
C. C., Gleave, M. E., Demichelis, F., Nanus, D. M. & Rubin, M. A. Molecular
Characterization of Neuroendocrine Prostate Cancer and Identification of New Drug
Targets. *Cancer Discovery* **1**, 487-495, (2011).
22.   Lotan, T. L., Gupta, N. S., Wang, W., Toubaji, A., Haffner, M. C., Chaux, A., Hicks, J. L.,
Meeker, A. K., Bieberich, C. J., De Marzo, A. M., Epstein, J. I. & Netto, G. J. ERG gene
rearrangements are common in prostatic small cell carcinomas. *Modern Pathology* **24**,
820-828, (2011).
23.   Mertz, K. D., Setlur, S. R., Dhanasekaran, S. M., Demichelis, F., Perner, S., Tomlins, S.,
Tchinda, J., Laxman, B., Vessella, R. L., Beroukhimt, R., Lee, C., Chinnaiyan, A. M. &
Rubin, M. A. Molecular Characterization of TMPRSS2-ERG Gene Fusion in the NCI-
H660 Prostate Cancer Cell Line: A New Perspective for an Old Model. *Neoplasia* **9**, 200-
IN203, (2007).
24.   Palmgren, J. S., Karavadia, S. S. & Wakefield, M. R. Unusual and Underappreciated:
Small Cell Carcinoma of the Prostate. *Seminars in Oncology* **34**, 22-29, (2007).
25.   Jin, R. J., Wang, Y., Masumori, N., Ishii, K., Tsukamoto, T., Shappell, S. B., Hayward, S.
W., Kasper, S. & Matusik, R. J. NE-10 Neuroendocrine Cancer Promotes the LNCaP
Xenograft Growth in Castrated Mice. *Cancer Research* **64**, 5489-5495, (2004).
26.   Gao, N., Zhang, J., Rao, M. A., Case, T. C., Mirosevich, J., Wang, Y., Jin, R., Gupta, A.,
Rennie, P. S. & Matusik, R. J. The role of hepatocyte nuclear factor-3 alpha (Forkhead

Box A1) and androgen receptor in transcriptional regulation of prostatic genes. *Molecular*
Endocrinology **17**, 1484-1507, (2003).

27. Gao, N., Ishii, K., Mirosevich, J., Kuwajima, S., Oppenheimer, S. R., Roberts, R. L.,
Jiang, M., Yu, X., Shappell, S. B., Caprioli, R. M., Stoffel, M., Hayward, S. W. & Matusik,
R. J. Forkhead box A1 regulates prostate ductal morphogenesis and promotes epithelial
cell maturation. *Development* **132**, 3431-3443, (2005).

28. DeGraff, D. J., Grabowska, M. M., Case, T. C., Yu, X., Herrick, M. K., Hayward, W. J.,
Strand, D. W., Cates, J. M., Hayward, S. W., Gao, N., Walter, M. A., Buttyan, R., Yi, Y.,
Kaestner, K. H. & Matusik, R. J. FOXA1 deletion in luminal epithelium causes prostatic
hyperplasia and alteration of differentiated phenotype. *Laboratory Investigation* **94**, 726-
739, (2014).

29. Grabowska, M. M., Elliott, A. D., DeGraff, D. J., Anderson, P. D., Anumanthan, G.,
Yamashita, H., Sun, Q., Friedman, D. B., Hachey, D. L., Yu, X., Sheehan, J. H., Ahn, J.-
M., Raj, G. V., Piston, D. W., Gronostajski, R. M. & Matusik, R. J. NFI Transcription
Factors Interact with FOXA1 to Regulate Prostate-Specific Gene Expression. *Molecular*
Endocrinology **28**, 949-964, (2014).

30. Zhang, J., Gao, N., DeGraff, D. J., Yu, X., Sun, Q., Case, T. C., Kasper, S. & Matusik, R.
J. Characterization of cis elements of the probasin promoter necessary for prostate-
specific gene expression. *The Prostate* **70**, 934-951, (2010).

31. Yeung, L. H., Read, J. T., Sorenson, P., Nelson, C. C., Jia, W. & Rennie, P. S.
Identification and characterization of a prostate-specific androgen-independent protein-
binding site in the probasin promoter. *Biochemical Journal* **371**, 843-855, (2003).

32. Jia, L., Berman, B. P., Jariwala, U., Yan, X., Cogan, J. P., Walters, A., Chen, T.,
Buchanan, G., Frenkel, B. & Coetzee, G. A. Genomic androgen receptor-occupied
regions with different functions, defined by histone acetylation, coregulators and
transcriptional capacity. *PLoS ONE* **3**, e3645, (2008). 2577007

33. Sharma, N. L., Massie, C. E., Ramos-Montoya, A., Zecchini, V., Scott, H. E., Lamb, A.
D., MacArthur, S., Stark, R., Warren, A. Y., Mills, I. G. & Neal, D. E. The androgen
receptor induces a distinct transcriptional program in castration-resistant prostate cancer
in man. *Cancer Cell* **23**, 35-47, (2013).

34. Grabowska, M. M., Kelly, S. M., Reese, A. L., Cates, J. M., Case, T. C., Zhang, J.,
DeGraff, D. J., Strand, D. W., Miller, N. L., Clark, P. E., Hayward, S. W., Gronostajski, R.
M., Anderson, P. D. & Matusik, R. J. Nfib regulates transcriptional networks that control
the development of prostatic hyperplasia. *Endocrinology* **157**, 1094-1109, (2016).

35. Goldstein, J., Borowsky, A. D., Goyal, R., Roland, J. T., Arnold, S. A., Gellert, L. L.,
Clark, P. E., Hameed, O. & Giannico, G. A. MAGI-2 in prostate cancer: an
immunohistochemical study. *Human Pathology* **52**, 83-91, (2016).

36. Mirosevich, J., Gao, N., Gupta, A., Shappell, S. B., Jove, R. & Matusik, R. J. Expression
and role of Foxa proteins in prostate cancer. *Prostate* **66**, 1013-1028, (2006).

37. Thalmann, G. N., Anezinis, P. E., Chang, S.-M., Zhau, H. E., Kim, E. E., Hopwood, V. L.,
Pathak, S., von Eschenbach, A. C. & Chung, L. W. K. Androgen-independent Cancer
Progression and Bone Metastasis in the LNCaP Model of Human Prostate Cancer.
Cancer Research **54**, 2577-2581, (1994).

38. Franco, O. E., Jiang, M., Strand, D. W., Peacock, J., Fernandez, S., Jackson, R. S., 2nd,
Revelo, M. P., Bhowmick, N. A. & Hayward, S. W. Altered TGF-β signaling in a
subpopulation of human stromal cells promotes prostatic carcinogenesis. *Cancer*
Research **71**, 1272-1281, (2011).

39. Jiang, M., Strand, D. W., Fernandez, S., He, Y., Yi, Y., Birbach, A., Qiu, Q., Schmid, J.,
Tang, D. G. & Hayward, S. W. Functional remodeling of benign human prostatic tissues
in vivo by spontaneously immortalized progenitor and intermediate cells. *Stem cells*
(Dayton, Ohio) **28**, 344-356, (2010).

40.   Dehm, S. M., Schmidt, L. J., Heemers, H. V., Vessella, R. L. & Tindall, D. J. Splicing of a
670   Novel Androgen Receptor Exon Generates a Constitutively Active Androgen
671   Receptor that Mediates Prostate Cancer Therapy Resistance. *Cancer Research* **68**,
672   5469-5477, (2008).
673   41.   Morgenstern, J. P. & Land, H. Advanced mammalian gene transfer: high titre retroviral
674   vectors with multiple drug selection markers and a complementary helper-free packaging
675   cell line. *Nucleic Acids Research* **18**, 3587-3596, (1990).
676   42.   Stewart, S. A., Dykxhoorn, D. M., Palliser, D., Mizuno, H., Yu, E. Y., An, D. S., Sabatini,
677   D. M., Chen, I. S. Y., Hahn, W. C., Sharp, P. A., Weinberg, R. A. & Novina, C. D.
678   Lentivirus-delivered stable gene silencing by RNAi in primary cells. *RNA (New York,*
679   *N.Y.)* **9**, 493-501, (2003).
680   43.   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
681   image analysis. *Nature Methods* **9**, 671, (2012).
682   44.   Abramoff, M. D., Magalhaes, P. J. & Ram, S. J. Image Processing with ImageJ.
683   *Biophotonics International* **11**, 36-42, (2004).
684   45.   Image J (U. S. National Institutes of Health, Bethesda, Maryland, USA).
685   46.   Ling, G., Hauer, C. R., Gronostajski, R. M., Pentecost, B. T. & Ding, X. Transcriptional
686   Regulation of Rat CYP2A3 by Nuclear Factor 1: IDENTIFICATION OF A NOVEL NFI-A
687   ISOFORM, AND EVIDENCE FOR TISSUE-SELECTIVE INTERACTION OF NFI WITH
688   THE CYP2A3 PROMOTER IN VIVO. *Journal of Biological Chemistry* **279**, 27888-27895,
689   (2004).
690   47.   Delk, N. A. & Farach-Carson, M. C. Interleukin-6: A bone marrow stromal cell paracrine
691   signal that induces neuroendocrine differentiation and modulates autophagy in bone
692   metastatic PCa cells. *Autophagy* **8**, 650-663, (2012).
693   48.   Nakashima, J., Tachibana, M., Horiguchi, Y., Oya, M., Ohigashi, T., Asakura, H. &
694   Murai, M. Serum Interleukin 6 as a Prognostic Factor in Patients with Prostate Cancer.
695   *Clinical Cancer Research* **6**, 2702-2706, (2000).
696   49.   Adler, H. L., McCurdy, M. A., Kattan, M. W., Timme, T. L., Scardino, P. T. & Thompson,
697   T. C. ELEVATED LEVELS OF CIRCULATING INTERLEUKIN-6 AND TRANSFORMING
698   GROWTH FACTOR-beta 1 IN PATIENTS WITH METASTATIC PROSTATIC
699   CARCINOMA. *The Journal of Urology* **161**, 182-187, (1999).
700   50.   Twillie, D. A., Eisenberger, M. A., Carducci, M. A., Hseih, W.-S., Kim, W. Y. & Simons, J.
701   W. Interleukin-6: A candidate mediator of human prostate cancer morbidity. *Urology* **45**,
702   542-549, (1995).
703   51.   Chen, K.-S., Lim, J. W. C., Richards, L. J. & Bunt, J. The convergent roles of the nuclear
704   factor I transcription factors in development and cancer. *Cancer Letters* **410**, 124-138,
705   (2017).
706   52.   Consortium, T. U. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids*
707   *Research* **47**, D506-D515, (2018).
708   53.   Mukhopadhyay, S. S. & Rosen, J. M. The C-terminal domain of the nuclear factor I-B2
709   isoform is glycosylated and transactivates the WAP gene in the JEG-3 cells. *Biochemical*
710   *and Biophysical Research Communications* **358**, 770-776, (2007).
711   54.   Yang, S. H., Galanis, A., Witty, J. & Sharrocks, A. D. An extended consensus motif
712   enhances the specificity of substrate modification by SUMO. *The EMBO Journal* **25**,
713   5083-5093, (2006).
714   55.   Liu, Y., Bernard, H.-U. & Apt, D. NFI-B3, a Novel Transcriptional Repressor of the
715   Nuclear Factor I Family, Is Generated by Alternative RNA Processing. *Journal of*
716   *Biological Chemistry* **272**, 10739-10745, (1997).
717   56.   Li, S., Hu, M. G., Sun, Y., Yoshioka, N., Ibaragi, S., Sheng, J., Sun, G., Kishimoto, K. &
718   Hu, G.-F. Angiogenin mediates androgen-stimulated prostate cancer growth and
719   enables castration resistance. *Molecular cancer research : MCR* **11**, 1203-1214, (2013).

57. Park, J. W., Lee, J. K., Witte, O. N. & Huang, J. FOXA2 is a sensitive and specific
marker for small cell neuroendocrine carcinoma of the prostate. *Modern Pathology*,
(2017).
58. Degraff, D. J., Aguiar, A. A. & Sikes, R. A. Disease evidence for IGFBP-2 as a key
player in prostate cancer progression and development of osteosclerotic lesions.
American Journal of Translational Research **1**, 115-130, (2009).
59. Kiyama, S., Morrison, K., Zellweger, T., Akbari, M., Cox, M., Yu, D., Miyake, H. &
Gleave, M. E. Castration-Induced Increases in Insulin-Like Growth Factor-Binding
Protein 2 Promotes Proliferation of Androgen-independent Human Prostate LNCaP
Tumors. *Cancer Research* **63**, 3575-3584, (2003).
60. Moreira, D. M., Howard, L. E., Sourbeer, K. N., Amarasekara, H. S., Chow, L. C.,
Cockrell, D. C., Hanyok, B. T., Aronson, W. J., Kane, C. J., Terris, M. K., Amling, C. L.,
Cooperberg, M. R., Liede, A. & Freedland, S. J. Predictors of Time to Metastasis in
Castration-resistant Prostate Cancer. *Urology* **96**, 171-176, (2016).
61. Wu, Y., Zhang, J., Hou, S., Cheng, Z. & Yuan, M. Non-small cell lung cancer: miR-30d
suppresses tumor invasion and migration by directly targeting NFIB. *Biotechnology*
Letters **39**, 1827-1834, (2017).
62. Fane, M. E., Chhabra, Y., Hollingsworth, D. E. J., Simmons, J. L., Spoerri, L., Oh, T. G.,
Chauhan, J., Chin, T., Harris, L., Harvey, T. J., Muscat, G. E. O., Goding, C. R., Sturm,
R. A., Haass, N. K., Boyle, G. M., Piper, M. & Smith, A. G. NFIB Mediates BRN2 Driven
Melanoma Cell Migration and Invasion Through Regulation of EZH2 and MITF.
EBioMedicine **16**, 63-75, (2017).
63. Wu, N., Jia, D., Ibrahim, A. H., Bachurski, C. J., Gronostajski, R. M. & MacPherson, D.
NFIB overexpression cooperates with Rb/p53 deletion to promote small cell lung cancer.
Oncotarget **7**, 57514-57524, (2016).
64. Denny, S. K., Yang, D., Chuang, C.-H., Brady, J. J., Lim, J. S., Grüner, B. M., Chiou, S.-
H., Schep, A. N., Baral, J., Hamard, C., Antoine, M., Wislez, M., Kong, C. S., Connolly,
A. J., Park, K.-S., Sage, J., Greenleaf, W. J. & Winslow, M. M. Nfib Promotes Metastasis
through a Widespread Increase in Chromatin Accessibility. *Cell* **166**, 328-342, (2016).
65. Semenova, Ekaterina A., Kwon, M.-c., Monkhorst, K., Song, J.-Y., Bhaskaran, R.,
Krijgsman, O., Kuilman, T., Peters, D., Buikhuisen, Wieneke A., Smit, Egbert F.,
Pritchard, C., Cozijnsen, M., van der Vliet, J., Zevenhoven, J., Lambooij, J.-P., Proost,
N., van Montfort, E., Velds, A., Huijbers, Ivo J. & Berns, A. Transcription Factor NFIB Is
a Driver of Small Cell Lung Cancer Progression in Mice and Marks Metastatic Disease in
Patients. *Cell Reports* **16**, 631-643, (2016).
66. Becker-Santos, D. D., Thu, K. L., English, J. C., Pikor, L. A., Martinez, V. D., Zhang, M.,
Vucic, E. A., Luk, M. T. Y., Carraro, A., Korbelik, J., Piga, D., Lhomme, N. M., Tsay, M.
J., Yee, J., MacAulay, C. E., Lam, S., Lockwood, W. W., Robinson, W. P., Jurisica, I. &
Lam, W. L. Developmental transcription factor NFIB is a putative target of oncofetal
miRNAs and is associated with tumour aggressiveness in lung adenocarcinoma. *The*
Journal of Pathology **240**, 161-172, (2016).
67. Cao, Q., Wang, X., Zhao, M., Yang, R., Malik, R., Qiao, Y., Poliakov, A., Yocum, A. K.,
Li, Y., Chen, W., Cao, X., Jiang, X., Dahiya, A., Harris, C., Feng, F. Y., Kalantry, S., Qin,
Z. S., Dhanasekaran, S. M. & Chinnaiyan, A. M. The Central Role of EED in the
Orchestration of Polycomb Group Complexes. *Nature Communications* **5**, 3127-3127,
(2014).
68. Dooley, A. L., Winslow, M. M., Chiang, D. Y., Banerji, S., Stransky, N., Dayton, T. L.,
Snyder, E. L., Senna, S., Whittaker, C. A., Bronson, R. T., Crowley, D., Barretina, J.,
Garraway, L., Meyerson, M. & Jacks, T. Nuclear factor I/B is an oncogene in small cell
lung cancer. *Genes & Development* **25**, 1470-1475, (2011).

69.   Zhou, Z., Flesken-Nikitin, A., Corney, D. C., Wang, W., Goodrich, D. W., Roy-Burman, P.
& Nikitin, A. Y. Synergy of p53 and Rb Deficiency in a Conditional Mouse Model for
Metastatic Prostate Cancer. *Cancer Research* **66**, 7889-7898, (2006).

Figure Legends

Figure 1: Expression of AR, synaptophysin, and NFIB in human prostate. Representative

images from the human tissue microarray, representing normal prostate, prostate cancer, and

castration-resistant prostate cancer. H&E: Hematoxylin and eosin; SYP: synaptophysin

Figure 2: Expression of NFIB, AR, and synaptophysin in prostate and primary prostate

cancer. Expression of NFIB (**A**), AR (**B**), and synaptophysin (**C**) in the nuclear and cytoplasmic

fractions of normal prostate and prostate cancer. Intensity and distribution scores in the nuclear

and cytoplasmic compartments of each core were added together to generate a composite

expression score for each protein of interest and patient averages were generated. Data were

analyzed using the Wilcoxon matched-pairs signed rank test. For analysis of association of

Gleason score with NFIB (**D**), AR (**E**), and synaptophysin (**F**) staining, unique core values were

used. Significant values are reported relative to Gleason score 3. Data were analyzed using the

Kruskal-Wallis test with Dunn's multiple comparisons test. SYP: synaptophysin * $P < 0.05$; **** $P<$

0.0001

Figure 3: Cox regression of NFIB expression in prostate cancer. A. Case processing

summary of NFIB as a single variable to predict time to biochemical recurrence (event) in days.

B/C. Nuclear and cytoplasmic NFIB alone do not predict biochemical recurrence. Variables in the

Cox regression model. **D**. Case processing summary of cytoplasmic-to-nuclear NFIB ratio and

surgical margin status to predict time to biochemical recurrence in days. **E.** Cox regression model

predicting time to biochemical recurrence using the cytoplasmic-to-nuclear NFIB ratio. **F.** Cox

regression model including positive margin status and nuclear-to-cytoplasmic NFIB ratio. B:

unstandardized regression coefficient; SE: standard error; Exp (B): hazard ratio; CI: confidence

interval; df: degrees of freedom

Figure 4: NFIB, AR, and synaptophysin expression in primary and castration-resistant prostate cancer. Expression of NFIB (**A**), AR (**B**), and synaptophysin (**C**) in the nuclear and cytoplasmic fractions of human prostatic tissues. Intensity and distribution scores in the nuclear and cytoplasmic compartments of each core were added together to generate a composite expression score for each protein of interest. An average value was generated for each patient. Significant values are reported between prostate cancer and normal prostate or castration-resistant prostate cancer. Data were analyzed using the Kruskal-Wallis test with Dunn's multiple comparisons test. PCa: prostate cancer; CRPCa: castration-resistant prostate cancer; SYP: synaptophysin ** P< 0.01; *** P< 0.001; **** P< 0.0001

Figure 5: NFIB is expressed in prostate cancer cell lines and interacts with AR. A. NFIB expression in prostate cancer cell lines. Expression of NFIB in NFIB-low JEG-3 cells, androgen-dependent (LNCaP), castration-resistant (C4-2B, 22RV1), and androgen independent (PC3, DU-145) prostate cancer cells. **B.** Nuclear and cytoplasmic localization of NFIB. Nuclear and cytoplasmic preparations of LNCaP, C4-2B, and 22RV1 cells, analyzed for AR and NFIB expression. Nuclear loading control: Lamin B1; cytoplasmic loading control: GAPDH; N: nuclear; C: cytoplasmic **C.** NFIB interacts with AR. Transient transfection of JEG-3 cells with Vector, NFIB, full length AR (AR-FL), or AR-FL and NFIB, followed by co-immunoprecipitation with an antibody against NFIB and Western blotting for AR and NFIB. **D.** Generation of 3X-FLAG-NFIB cells. Western blot analysis of Flag expression in vector and 3X-FLAG-NFIB LNCaP and C4-2B cells. Star: antibody heavy chain IgG. IP: immunoprecipitation

Figure 6: NFIB supports PSA expression. A. Human cytokine arrays exposed to lysates from LNCaP and C4-2B cells expressing the vector or 3X-FLAG-NFIB construct. Dashed box denotes location of DPPIV dots, solid box with tick marks outlines PSA dots. Double dots in upper left, right, and lower left are positive controls. Lower right hand corner is empty and a negative control.

B. Quantification of integrated density for proteome arrays from A, adjusted for background and

normalized respective vector positive controls. **C.** Comparison of cytokine secretion across cell

lines, presented as cytokine integrated density divided by average positive control integrated

density of each cell line. **D.** Prostate specific antigen enhancer-promoter (PSA-EP) luciferase

activity in vector and 3X-FLAG-NFIB over-expressing cells. Data presented as fold change PSA-

EP-luciferase over SV40-renilla, normalized to vector. Data were analyzed using the Mann-

Whitney U test. **E.** Prostate specific antigen enhancer promoter (PSA-EP) luciferase activity in

vector and NFIB over-expressing cells. Data presented as fold change PSA-EP-luciferase over

SV40-renilla, normalized to LNCaP vector expression. Data were analyzed using the Mann-

Whitney U test, comparing vector to NFIB luciferase changes by cell line. ** P< 0.01

Supplemental Figure 1: Expression of AR, synaptophysin, and NFIB in human prostate. A.

Representative images from the human tissue microarray, representing normal prostate, prostate

cancer, and castration-resistant prostate cancer. **B.** FOXA1 and NFIB straining in primary human

small cell prostate cancer. H&E: Hematoxylin and eosin; SYP: synaptophysin

Supplemental Figure 2: Correlations between NFIB, AR, and synaptophysin in prostate

cancer. Spearman correlations between nuclear and cytoplasmic staining for NFIB, AR, and

synaptophysin in primary prostate cancer (**A**), both primary and castration-resistant prostate

cancer (**B**), and castration-resistant only (**C**).

Supplemental Figure 3: Expression of NFIB, AR, and synaptophysin across prostate

diseases. A. Cytoplasmic-to-nuclear ratio of NFIB does not vary by Gleason pattern by core or

by patient tumor stage. **B/C/D.** Expression of NFIB (**B**), AR (**C**), and synaptophysin (**D**) in the

nuclear and cytoplasmic fractions of human prostatic tissues. Intensity and distribution scores in

the nuclear and cytoplasmic compartments of each core were added together to generate a

composite expression score for each protein of interest. Data were analyzed using the Kruskal-

Wallis test with Dunn's multiple comparisons test. Significant values are reported between normal

prostate (normal) and diseased prostate. BPH: benign prostatic hyperplasia; HGPIN: high grade

prostatic intraepithelial neoplasia; Adj. Normal: adjacent normal; PCa: prostate cancer; CRPCa:

castration-resistant prostate cancer; SYP: synaptophysin * P < 0.05; ** P< 0.01; *** P< 0.001; ****

P< 0.0001

Supplemental Figure 4: NFIB isoforms. A. Expression of NFIB in stromal and epithelial cells

derived from BPH patients (BHPrS-1, BHPrE-1) and non-hyperplastic prostate epithelial cells

(NHPrE-1). Cancer cell lines shown as control. **B.** Calculated molecular weights of NFIB. **C.** NFIB

exists in several isoforms in PC-3 cells. Transient over-expression of vector or NFIB or knockdown

of non-targeting siRNA or NFIB in PC-3 cells. **D.** NFIB exists in several isoforms. Transient

knockdown of non-targeting siRNA or NFIB in JEG-3, LNCaP, C4-2B, and 22RV1 cells.

	H&E	AR	SYP	NFIB
Normal Prostate				
Prostate Cancer (Gleason 3)				
Prostate Cancer (Gleason 5)				
Castrate Resistant Prostate Cancer (Low NFIB)				
Castrate Resistant Prostate Cancer (High NFIB)				
Therapy Acquired Neuroendocrine Prostate Cancer				

A.

Case Processing Summary

		N	Percent
Cases available in analysis	Event[a]	32	57.1%
	Censored	19	33.9%
	Total	51	91.1%
Cases dropped	Cases with missing values	5	8.9%
	Cases with negative time	0	0.0%
	Censored cases before the earliest event in a stratum	0	0.0%
	Total	5	8.9%
Total		56	100.0%

a. Dependent Variable: biochemical_recur_days

B.

Variables in the Equation

	B	SE	Wald	df	Sig.	Exp(B)	95.0% CI for Exp(B) Lower	Upper
Nuclear NFIB expression	-0.034	0.148	0.052	1	0.820	0.967	0.723	1.292

C.

Variables in the Equation

	B	SE	Wald	df	Sig.	Exp(B)	95.0% CI for Exp(B) Lower	Upper
Cytoplasmic NFIB expression	0.140	0.110	1.616	1	0.204	1.150	0.927	1.427

D.

Case Processing Summary

		N	Percent
Cases available in analysis	Event[a]	31	55.4%
	Censored	18	32.1%
	Total	49	87.5%
Cases dropped	Cases with missing values	7	12.5%
	Cases with negative time	0	0.0%
	Censored cases before the earliest event in a stratum	0	0.0%
	Total	7	12.5%
Total		56	100.0%

a. Dependent Variable: biochemical_recur_days

E.

Variables in the Equation

	B	SE	Wald	df	Sig.	Exp(B)	95.0% CI for Exp(B) Lower	Upper
NFIB Cytoplasmic to Nuclear Ratio	1.052	0.554	3.603	1	0.058	2.864	0.966	8.491

F.

Variables in the Equation

	B	SE	Wald	df	Sig.	Exp(B)	95.0% CI for Exp(B) Lower	Upper
NFIB Cytoplasmic to Nuclear Ratio	2.189	0.766	8.166	1	0.004	8.930	1.989	40.086
Positive Margin Status	2.084	0.469	19.757	1	0.00001	8.035	3.206	20.138

A.

B.

C.

A.

B.

C.

D.

Supp 1

A.

Correlations

Spearman's rho			NFIB_Tumor_Nuclear	NFIB_Tumor_Cytoplasmic	AR_Tumor_Nuclear	AR_Tumor_Cytoplasmic	SYP_Tumor_Cytoplasmic
Spearman's rho	NFIB_Tumor_Nuclear	Correlation Coefficient	1.000	.305**	.419**	0.085	0.111
		Sig. (2-tailed)		0.001	1.875E-06	0.359	0.228
		N	120	120	120	120	120
	NFIB_Tumor_Cytoplasmic	Correlation Coefficient	.305**	1.000	.248**	.447**	.464**
		Sig. (2-tailed)	0.001		0.006	3.135E-07	9.483E-08
		N	120	120	120	120	120
	AR_Tumor_Nuclear	Correlation Coefficient	.419**	.248**	1.000	0.109	.350**
		Sig. (2-tailed)	1.875E-06	0.006		0.234	0.000
		N	120	120	120	120	120
	AR_Tumor_Cytoplasmic	Correlation Coefficient	0.085	.447**	0.109	1.000	.487**
		Sig. (2-tailed)	0.359	3.135E-07	0.234		1.638E-08
		N	120	120	120	120	120
	SYP_Tumor_Cytoplasmic	Correlation Coefficient	0.111	.464**	.350**	.487**	1.000
		Sig. (2-tailed)	0.228	9.483E-08	9.050E-05	1.638E-08	
		N	120	120	120	120	120

**. Correlation is significant at the 0.01 level (2-tailed).

B.

Correlations

Spearman's rho			NFIB_Tumor_Nuclear	NFIB_Tumor_Cytoplasmic	AR_Tumor_Nuclear	AR_Tumor_Cytoplasmic	SYP_Tumor_Cytoplasmic
Spearman's rho	NFIB_Tumor_Nuclear	Correlation Coefficient	1.000	.321**	.394**	0.061	0.123
		Sig. (2-tailed)		2.176E-04	4.151E-06	0.494	0.168
		N	128	128	128	128	128
	NFIB_Tumor_Cytoplasmic	Correlation Coefficient	.321**	1.000	.228**	.381**	.473**
		Sig. (2-tailed)	2.176E-04		0.010	9.192E-06	1.766E-08
		N	128	128	128	128	128
	AR_Tumor_Nuclear	Correlation Coefficient	.394**	.228**	1.000	0.152	.279**
		Sig. (2-tailed)	4.151E-06	0.010		0.088	0.001
		N	128	128	128	128	128
	AR_Tumor_Cytoplasmic	Correlation Coefficient	0.061	.381**	0.152	1.000	.419**
		Sig. (2-tailed)	0.494	0.000	0.088		8.711E-07
		N	128	128	128	128	128
	SYP_Tumor_Cytoplasmic	Correlation Coefficient	0.123	.473**	.279**	.419**	1.000
		Sig. (2-tailed)	0.168	0.000	0.001	8.711E-07	
		N	128	128	128	128	128

**. Correlation is significant at the 0.01 level (2-tailed).

C.

Correlations

Spearman's rho			NFIB_Tumor_Nuclear	NFIB_Tumor_Cytoplasmic	AR_Tumor_Nuclear	AR_Tumor_Cytoplasmic	SYP_Tumor_Cytoplasmic
Spearman's rho	NFIB_Tumor_Nuclear	Correlation Coefficient	1.000	0.625	0.250	-0.058	0.317
		Sig. (2-tailed)		0.098	0.550	0.891	0.444
		N	8	8	8	8	8
	NFIB_Tumor_Cytoplasmic	Correlation Coefficient	0.625	1.000	-0.084	-0.320	.735*
		Sig. (2-tailed)	0.098		0.843	0.439	0.038
		N	8	8	8	8	8
	AR_Tumor_Nuclear	Correlation Coefficient	0.250	-0.084	1.000	.712*	-0.466
		Sig. (2-tailed)	0.550	0.843		0.048	0.244
		N	8	8	8	8	8
	AR_Tumor_Cytoplasmic	Correlation Coefficient	-0.058	-0.320	.712*	1.000	-0.262
		Sig. (2-tailed)	0.891	0.439	0.048		0.531
		N	8	8	8	8	8
	SYP_Tumor_Cytoplasmic	Correlation Coefficient	0.317	.735*	-0.466	-0.262	1.000
		Sig. (2-tailed)	0.444	0.038	0.244	0.531	
		N	8	8	8	8	8

*. Correlation is significant at the 0.05 level (2-tailed).

A.

B.

NFIB bands on gels	Observed molecular weight range	Mean molecular weight
Band 1	60.5 kDa to 65 kDa	62kDa
Band 2	54.7 kDa to 60 kDa	56.98 kDa
Band 3	47.5 kDa to 50.2 kDa	48.78 kDa
Band 4	37 kDa to 40 kDa	39.2 kDa

C.

D.