# Prenatal alcohol exposure programs offspring disease: Insulin resistance in adult males in a rat model of acute exposure

**Tam M T Nguyen[1], Sarah E Steane[1], Karen M Moritz[1,2], Lisa K Akison[1,2]**

[1]School of Biomedical Sciences, The University of Queensland, St Lucia, QLD, Australia.

[2]Child Health Research Centre, The University of Queensland, South Brisbane, QLD, Australia.

*Corresponding Author*:  Lisa Akison

*Address:* Child Health Research Centre, The University of Queensland, Room 408, Centre for Children's Health Research (CCHR), 62 Graham St, South Brisbane, QLD 4101, AUSTRALIA

*Phone:* +61 7 3069 7511

*Email:* l.akison@uq.edu.au

*Running Title:* Prenatal alcohol exposure and offspring metabolism

**Key points summary**

- Prenatal alcohol exposure has the potential to affect fetal development and program chronic disease in offspring.

- Previous preclinical models typically use high, chronic doses of alcohol throughout pregnancy to examine effects on offspring, particularly on the brain and behaviour.

- In this study we use a rat model of moderate, acute, prenatal alcohol exposure to determine if this can be detrimental to maintenance of glucose homeostasis in adolescent and adult offspring.

- Although female offspring were relatively unaffected, there was evidence of insulin resistance in 6-month old male offspring exposed to prenatal alcohol, suggestive of a pre-diabetic state.

- This result suggests that even a relatively low-dose, acute exposure to alcohol during pregnancy can still program metabolic dysfunction in a sex-specific manner.

**Abstract**

Alcohol consumption is highly prevalent amongst women of reproductive age. Given

that approximately 50% of pregnancies are unplanned, alcohol has the potential to affect

fetal development and program chronic disease in offspring. We examined the effect of

an acute but moderate prenatal alcohol exposure (PAE) on glucose metabolism, lipid

levels and dietary preference in adolescent and/or adult rat offspring. Pregnant Sprague-

Dawley rats received an oral gavage of ethanol (1g/kg maternal body weight, n=9 dams)

or an equivalent volume of saline (control, n=8 dams) at embryonic days 13.5 and 14.5.

PAE resulted in a blood alcohol concentration of 0.05-0.06% 1h post-gavage in dams.

Fasting blood glucose concentration was not affected by PAE in offspring at any age,

nor were blood glucose levels during a glucose tolerance test (GTT) in 6-month old

offspring ($P>0.5$). However, there was evidence of insulin resistance in PAE male

offspring at 6 months of age, with significantly elevated fasting plasma insulin ($P =$

$0.001$), a tendency for increased first phase insulin secretion during the GTT and

impaired glucose clearance following an insulin challenge ($P = 0.007$). This was

accompanied by modest alterations in protein kinase B (AKT) signalling in adipose

tissue. PAE also resulted in reduced calorie consumption by offspring compared to

controls ($P = 0.04$). These data suggest that a relatively low-level, acute PAE programs

metabolic dysfunction in offspring in a sex-specific manner. These results highlight that

alcohol consumption during pregnancy has the potential to affect the long-term health

of offspring.

*Key words:* fetal programming, glucose metabolism, insulin resistance, food preference,

gene expression

**Introduction**

It is clear from both epidemiological and animal studies that early life events can impact on development and program adult disease. This is known as the developmental origins of health and disease (DOHaD) hypothesis and incorporates perturbations throughout both fetal and early postnatal development (Barker, 2007). During pregnancy, a sub-optimal uterine environment can result from maternal insults such as malnutrition, stress, obesity, and drug use. In response to a perturbation, adaptations occur in favour of vital organs, such as the brain, to ensure short-term fetal survival; however, the development of other organs deemed relatively unnecessary, such as the kidneys, liver and pancreas, can be compromised (Heindel & Vandenberg, 2015). This can result in long-term adverse health outcomes in offspring associated with the altered function in these organs. Although both clinical studies and animal models suggest that the timing of exposure can be important, organs requiring a prolonged period of development, such as the liver (Godlewski *et al.*, 1997) and pancreas (Gittes, 2009), are particularly vulnerable to insults at multiple time-points. Therefore, metabolic dysfunction is commonly reported in offspring from animal models of maternal malnutrition, obesity and stress, as well as clinical studies with evidence of these exposures (see (Fleming *et al.*, 2018) for review). Interestingly, effects are often sex-specific, with males consistently more susceptible to adverse outcomes than females (Weinberg *et al.*, 2008; Sundrani *et al.*, 2017).

One maternal perturbation receiving increasing attention is prenatal alcohol exposure (PAE). Despite health authorities advising against alcohol consumption while pregnant or planning a pregnancy (World Health Organisation, 2004; National Health and Medical Research Council, 2009), a recent systematic review and meta-analysis estimated the global rate of alcohol consumption during pregnancy to be ~10%, although in many Western countries it is much higher (Popova *et al.*, 2017). Although PAE causes well-recognised neurological and behavioural/cognitive deficits in offspring, there is also emerging evidence for a range of deficits in other body systems, leading to long-term adverse health outcomes, including impaired glucose metabolism. While clinical studies are sparse, one study reports on a small cohort ($n = 7$) of early school-age children with fetal alcohol syndrome (FAS), providing some evidence of glucose intolerance and insulin resistance in these patients compared to 'normal'

controls (Castells *et al.*, 1981). However, compelling evidence for metabolic

dysfunction following PAE comes from preclinical studies. In our recent systematic

review, all but three of the 18 studies included reported glucose intolerance and/or

insulin resistance in PAE offspring in adulthood (see (Akison *et al.*, 2019) for details).

This was often associated with disruptions in molecular pathways involved in

gluconeogenesis, glucose transport, IGF signalling and/or insulin signalling pathways in

the liver and/or peripheral tissues. Further, recent studies, in both animal models of PAE

(Amos-Kroohs *et al.*, 2018; Dorey *et al.*, 2018) and children with FASD (Werts *et al.*,

2014; Smith *et al.*, 2015; Amos-Kroohs *et al.*, 2016), also report dysregulated eating

behaviours and food preference, potentially contributing to a propensity towards

diabetes or other metabolic dysfunction and associated comorbidities such as obesity.

However, in most of these preclinical models, chronic high doses of ethanol were

administered, resulting in daily peak blood alcohol concentrations (BAC) of 100-150

mg/dl (0.10-0.15%) (Chen & Nyomba, 2003b). This is not representative of drinking

patterns reported in the majority of pregnant women, which are typically acute, low-

moderate levels of alcohol, referred to as 'special occasion' drinking (Muggli *et al.*,

2016; McCormack *et al.*, 2017). Therefore, although the effects of substantial ethanol

consumption are well-recognised, the effects of mild exposure are less well known.

In the current study, we used a rat model of moderate, acute alcohol exposure during an

important stage for liver and pancreas development, and when fetal development is

equivalent to that of the first trimester human fetus, when the mother may still be

unaware of her pregnancy. Timing of exposure was based on the relative development

of these organs between humans and rodents (Godlewski *et al.*, 1997; Pan & Brissova,

2014). Food preference for a HFD compared to standard chow was examined at 4-5

months of age and then a glucose tolerance test (GTT) and insulin tolerance test (ITT)

were performed in offspring at 5-6 months of age. In order to elucidate potential

molecular mechanisms contributing to disease phenotypes, the effect of prenatal alcohol

exposure on hepatic expression of genes involved in glucose homeostasis and AKT-

signalling in peripheral tissues were also examined in adult offspring. We hypothesised

that our moderate, acute prenatal alcohol exposure could program glucose intolerance

and insulin resistance in adult offspring, as well as a preference for a high-fat diet.

Given previous reports of sex-specific effects, we included sex as a potential

contributing factor in all analyses.

**Materials and Methods**

*Ethical approval*

All animal experiments and procedures were approved by The University of

Queensland Anatomical Biosciences Ethics Committee (SBMS/AIBN/521/15/NHMRC)

and were conducted in accordance with the Australian Code for the Care and Use of

Animals for Scientific Purposes (2013, 8th Edition). Reporting of animal experiments

conforms to the ARRIVE guidelines (Kilkenny & Altman, 2010; Kilkenny *et al.*, 2010),

as well as the principles and regulations for animal experiment reporting and ethics

described by (Grundy, 2015).

*Animal model of acute prenatal alcohol exposure*

Outbred, nulliparous Sprague-Dawley rats were obtained from the Animal Resources

Centre, Perth WA. All animals (F0 and F1) were housed at the Australian Institute for

Bioengineering and Nanotechnology (AIBN) Animal Facility (University of

Queensland, St Lucia, QLD, Australia) in a temperature- and humidity-controlled

environment with an artificial 12 h reversed light-dark cycle and provided with standard

laboratory rat chow (Rat & Mouse Meat-Free Diet, Specialty Feeds, Glen Forrest WA,

Australia) and water ad libitum. Following transport, females were group-housed during

an initial acclimation period in open-top cages with wire lids, plastic bases and wood-

chip bedding.

At a weight threshold of 230 g (~3-4 months old), dams were assessed daily for

proestrous via vaginal electrical impedance ($\geq 4.0$ k$\Omega$) using an EC40 estrous cycle

monitor (Fine Science Tools, Foster City, CA, USA) as previously described (Jaramillo

*et al.*, 2012). Once in proestrous, dams were housed with a proven stud male for the first

5 h of the dark cycle (1200 – 1700) and successful mating confirmed by the presence of

a seminal plug, with the following morning designated as embryonic day (E) 0.5.

Pregnant dams were housed singly. Dams were weight-matched and then randomly

assigned to either receive ethanol (EtOH) or saline (Control) via oral gavage at E13.5

and E14.5. EtOH treated females ($n = 10$) received 18% v/v EtOH in saline solution

(0.9% NaCl) at a dose of 1 g/kg body weight, while Control females ($n = 8$) received an

equivalent volume of saline. Gavage was performed between 09:00-10:00 each day (i.e.

during the light cycle). Water and chow consumption were measured daily from E12.5

(one day prior to first gavage) until birth. Weight gain was monitored throughout

pregnancy. Day of birth was designated postnatal day (PN) 0, with offspring weighed

on alternate days from PN1, then weekly following weaning at PN21. Once offspring

were weaned, all dams were culled via $CO_2$ asphyxiation. Only 1 male and 1 female

were used from each litter for each experiment to remove potential litter effects, as

recommended in recent guidelines on experimental design in DOHaD studies

(Dickinson *et al.*, 2016). Where >2 pup weights were measured per litter, litter averages

were calculated and used in subsequent analyses. Figure 1 provides a flow chart

showing the dams allocated to each arm of the study and offspring that contribute to

each analysis at each age. All procedures conducted on offspring were performed during

the light cycle (0930-1200).

### *Blood and tissue collection*

Approximately 150 μl whole blood was collected from dams via a tail tip bleed at 1 h

and 5 h post gavage on E13.5 and 14.5 to measure blood alcohol concentration (BAC;

see details for measurement below) and non-fasted blood glucose level using an Accu-

Chek Performa glucometer (Roche, Bella Vista, NSW, Australia). At PN30, a subset of

offspring (see Figure 1) were weighed and then culled via $CO_2$ asphyxiation following a

16 h fasting period. Blood was collected via cardiac puncture for measurement of

fasting plasma insulin, metabolites involved in lipid metabolism and fasting blood

glucose. Blood was immediately centrifuged at 4000 rpm for 10 min at 4°C and plasma

separated, aliquoted and stored at -20°C until subsequent analysis. Fasting blood

glucose was also measured, following a 16 h fasting period, in all offspring at 10-11

weeks of age via a spot tail bleed and glucometer. Litter-mate blood glucose levels were

averaged for each sex prior to analysis for differences between PAE and control. At 6

months of age, 1 male and 1 female per litter had blood collected as part of a glucose

tolerance test (GTT) or insulin tolerance test (ITT) (see below for details). All offspring

were culled at 6-7 months of age via $CO_2$ asphyxiation, with a subset used for tissue

collection (1-2 males and 1-2 females per litter for a total of 12 per sex per treatment).

Females were assessed for proestrous (≥4.0 kΩ vaginal impedance) as described above

and were not culled at this stage of the cycle to reduce potential variability due to

hormonal changes just prior to ovulation. Body weight, abdominal circumference,

snout-rump length and tibia length were measured for calculations of size and ponderal

index (weight in g/cubed length in cm). Pancreas and liver were removed, weighed and

a section from the largest lobe of the liver snap-frozen in liquid $N_2$ and stored at -80°C

for subsequent molecular analysis. Samples of abdominal visceral fat and the left

gastrocnemius muscle were also frozen for subsequent molecular analysis.

*Measurement of blood alcohol concentration (BAC)*

Maternal blood samples were immediately centrifuged after collection at 4000 rpm for

10 min at 4°C and plasma separated, aliquoted and stored at -20°C until analysed. Blood

alcohol concentration (BAC) was measured by Pathology Queensland (Queensland

Health) using an alcohol dehydrogenase enzymatic assay (Beckman Coulter, Ref

#474947) and a Beckman Coulter AU480 Chemistry Analyzer (Beckman Coulter, Lane

Cove, NSW, Australia). The limit of detection for this assay was 5 mg/dL or 0.005%.

*Food preference study*

Food preference was assessed in offspring 4-5 months of age as previously reported

(Dorey *et al.*, 2018). Briefly, 1 male and 1 female from each litter were randomly

chosen and housed individually in cages with a divided feed hopper. After a 4-day

acclimatisation period, baseline consumption was established by measuring intake of

standard chow diet (SD, 4.8% fat) daily for a period of 4 days and calculating

consumption per gram of body weight. Offspring were then allowed free access to both

a high-fat 'Western' style chow diet (HFD; 21% fat; Specialty Feeds Diet SF00-219)

and the SD over a 4-day test period. The diets were randomly placed on either side of

the feed hopper and their positions switched after 2 days. Cage positions were also

randomised on the racks and were changed after 2 days to prevent systematic sampling

bias of Control and EtOH animals. Food and water intake were recorded daily and

consumption per gram of body weight calculated over the first 2 days (Choice Period 1,

CP1) and the second 2 days (Choice Period 2, CP2) of the two diets being offered. The

test period was split into CP1 and CP2, as a previous study has shown that the novelty

of the HFD diminishes over the test period, resulting in different eating behaviour over

the 4-day period (Dorey *et al.*, 2018).

*Glucose and insulin tolerance tests*

At 6-7 months of age, a GTT and ITT were performed as previously described (Probyn

*et al.*, 2013; Gardebjer *et al.*, 2015). Briefly, 1 male and 1 female were randomly

selected from each litter (EtOH = 9; Control = 8) for each test. For the GTT, offspring

were fasted overnight for 12-16 h and then received an intraperitoneal (ip) injection of a

50% w/v glucose solution (Baxter Healthcare, Old Toongabbie, NSW, Australia) at a

dose of 1 g/kg body weight. Blood was sampled via tail tip bleed, and blood glucose

concentrations were measured using a glucometer prior to glucose administration (-5

min), and at 5, 10, 20, 30, 45, 60 and 90 min post-bolus. Blood was also collected,

processed as described above and plasma stored for subsequent analysis of plasma

insulin and glucose levels. For the ITT, a separate group of non-fasted offspring

received an ip injection of 0.75 U/kg body weight insulin (Actrapid, Novo Nordisk

Pharmaceuticals Pty. Ltd., Baulkham Hills, NSW, Australia), and tail tip bleeds were

performed to measure blood glucose concentrations via glucometer as above at -5, 20,

40, 60, 90 and 120 min post-injection.

*Quantitative PCR and Western blotting*

Expression of genes involved in hepatic glucose metabolism and peripheral tissue

glucose transport and insulin signalling were examined using real-time quantitative

polymerase chain reaction (qPCR) (see Table 1 for details of specific genes). qPCR was

conducted and reported following the MIQE guidelines (Bustin *et al.*, 2009). Liver,

adipose tissue and gastrocnemius muscle samples collected from 6 month-old offspring

(*n* = 8 control and 9 EtOH; 1 female and/or 1 male per litter) were homogenised and

total RNA was extracted using RNeasy Mini Kits (liver; Qiagen, Chadstone, VIC,

Australia) or Trizol reagent (muscle and adipose tissue; ThermoFisher Scientific,

Richlands, QLD, Australia) according to the manufacturer's instructions. Total RNA (1

214 μg/reaction for liver and adipose tissue; 500 ng/reaction for muscle) was reverse

215 transcribed into cDNA using iScript Reverse Transcription Supermix for RT-qPCR

216 (Bio-Rad, Gladesville, NSW, Australia). qPCR reactions were performed on the

217 Applied Biosystems Quantstudio 6 Flex Real-Time PCR System (ThermoFisher

218 Scientific) using 10 ng and 20 ng of cDNA for muscle and liver/adipose tissue

219 respectively, Quantinova Probe PCR Master Mix (Qiagen) and Assay-on-Demand

220 primer/probe sets (see Table 1 for details). Reactions were multiplexed with beta-actin

221 (*Actb*) as an endogenous control. In addition, *Eif2a* and *Rpl19* were used as additional

222 endogenous control genes for liver and muscle/adipose tissue samples respectively (see

223 Table 1). All control genes were stably expressed across all samples, irrespective of

224 experimental group or sex (data not shown). The geometric mean of the two

225 endogenous control genes was used in the ΔΔCt calculation of gene expression, with

226 fold-change expressed relative to the average of the male offspring saline control group.

227 Total protein was extracted from visceral adipose tissue (100 mg), gastrocnemius

228 muscle (30 mg) and liver (100-200 mg) samples (1 male offspring only from each litter;

229 $n = 7$ control and 7 EtOH) for Western blot analysis of total and phosphorylated AKT,

230 which plays a central role in insulin signalling in hepatic and peripheral tissues

231 (Mackenzie & Elliott, 2014). Tissues were homogenised in Cell Lysis Buffer (Cell

232 Signaling Technology, Danvers, MA, USA) with protease (Sigma-Aldrich, Sydney,

233 NSW, Australia) and phosphatase (Roche) inhibitors using a FastPrep-24 5G

234 homogeniser (MP Biomedicals, Seven Hills, NSW, Australia). Homogenates were

235 centrifuged (11,000 rpm, 10 min, 4°C) and resultant supernatant assayed using a DC

236 Protein Assay Kit (Bio-Rad). A total of 20 μg of sample was loaded per well onto 12%

237 SDS-PAGE gels and subsequently transferred overnight at 4°C to Immun-Blot LF

238 PVDF membranes (Bio-Rad). Membranes were incubated overnight with one of the

239 following rabbit primary antibodies (all from Cell Signaling Technology and all cross-

240 react with rat): anti-Pan-AKT (1:1000, Cat# 5373s, 60 kDa, RRID:AB_10891424), anti-

241 phospho-AKT$_{Thr308}$ (1:1000, Cat# 9275s, 60 kDa, RRID:AB_329828), anti-phospho-

242 AKT$_{Ser473}$ (1:1000, Cat# 4060s, 60 kDa, RRID:AB_2315049). Protein expression was

243 measured as previously described using a LI-COR Odyssey CLx infrared imaging

244 system (LI-COR Biosciences, Lincoln, NE, USA, RRID:SCR_014579) following

245 exposure to LI-COR IRDye 680 goat anti-rabbit secondary antibody (LI-COR

Biosciences, Cat# 926-32221, RRID:AB_621841). Band intensity was analysed using

Image Studio Lite software (LI-COR Biosciences, v5.2, RRID:SCR_013715) and was

normalised to glyceraldehyde 3-phosphate dehydrogenase (GAPDH, 1:1000, Cat#

2118s, 37 kDa, RRID:AB_561053) or β-actin (ACTB, 1:1000, Cat#4970, 45 kDa,

RRID:AB_2223172) immunoreactivity for densitometric analysis. Bands were

quantified by an assessor blinded to treatment. Protein levels in the EtOH samples were

expressed relative to the average of the saline controls.

*Plasma insulin, glucose and lipid analysis*

Fasting plasma samples collected from offspring at PN30 and during the GTT at 6-7

months of age were analysed for insulin levels using a rat insulin radioimmunoassay kit

(SRI-13K, Millipore Australia, Kilsyth, VIC). Insulin samples were run in duplicate at

1:5, 1:10 or 1:20 dilution. Assay sensitivity was 0.03 ng/mL and inter- and intra-assay

coefficients of variation were 16.7% and 12.1% respectively. Plasma collected at

offspring culls at PN30 were analysed for triglycerides (TG), high-density lipoproteins

(HDL), and low-density lipoproteins (LDL) using a Cobas Integra 400 Plus Chemistry

Analyzer (Block Scientific, Bellport, NY, USA). In addition, plasma collected at

baseline for the GTT at 6 months of age were analysed for glucose levels using the

Cobas Analyzer, for comparison to measurement of glucose levels by the glucometer.

Blood glucose levels, as measured using the glucometer, were found to underestimate

plasma glucose levels measured using the 'gold standard' Cobas analyser (Figure 2A).

However, this underestimation was consistent across all treatment/sex combinations

(Figure 2B). Therefore, where insufficient plasma was available for Cobas analysis

(PN30), glucometer readings of blood glucose levels were adequate to compare fasting

glucose and insulin sensitivity/resistance indices between treatment/sex groups. When

sufficient plasma could be collected (6 months of age), plasma glucose levels were used

for all analyses.

*Calculations and statistical analyses*

Area under the glucose (AUGC) and insulin (AUIC) curves were calculated using the

trapezius method with baseline defined as the basal level prior to injection (Allison *et*

*al.*, 1995). Glucose concentration curves for the ITT were inverted before calculating

AUGC as previously described (Gardebjer *et al.*, 2015; Nagy & Einwallner, 2018).

Acute 1[st] phase insulin secretion was calculated as the AUIC from basal to 5 min post-

glucose bolus and 2[nd] phase insulin secretion as the AUIC from 5 to 90 min post-bolus.

Note that baseline was set as zero for the 2[nd] phase secretion analysis, given that 5 min

post-bolus was not at basal levels. Quantitative insulin sensitivity check index

(QUICKI) was calculated using $1/[\log(\text{fasting insulin (μU/mL)}) + \log(\text{fasting glucose}$

(mg/dL)] (Katz *et al.*, 2000). The homeostasis model assessment-estimated insulin

resistance (HOMA-IR) was calculated using $[\text{fasting insulin (μU/mL)} \times \text{fasting glucose}$

(mg/dL)]/2430, which has been validated for use in rats (Cacho *et al.*, 2008). Total

cholesterol (TC) was calculated by re-arranging the Friedewald equation (Friedewald *et*

*al.*, 1972) such that $TC = HDL + LDL + (TG/5)$.

Analyses were conducted using GraphPad Prism 7.0 (GraphPad Software, San Diego,

CA, USA) and all data presented as mean ± SEM. Prior to analysis, all data were tested

for normality using the D'Agnostino-Pearson omnibus or the Shapiro-Wilk normality

tests. Maternal parameters and Western blot data were analysed using a Student's t-test

(parametric data) or a Mann-Whitney U-test (non-parametric data). Where variances

were significantly different but data was normally distributed, Welch's correction was

applied. All other data were analysed using a two-way ANOVA, with offspring sex

(male/female) and treatment group (EtOH/Control) as factors. Where there was a

significant interaction, a Sidak multiple comparison test was used to determine

significantly different groups. Where data was not normally distributed, a non-

parametric Kruskal-Wallis test across all treatment/sex groups was used, with Dunn's

multiple comparison test used to determine significantly different groups but only

treatment differences within sex or sex differences with treatment are reported.

Significance level was $P<0.05$ for all statistical tests, with $0.05 \geq P \leq 1.0$ considered

indicative of trends.

**Results**

*Maternal parameters and postnatal growth*

Dams in the EtOH and Control groups were of similar weight at mating and at E13.5

(Table 2). Gestational weight gain prior to gavage and post-gavage were also not

significantly different between groups, nor were daily chow or water consumption

(Table 2). When energy intake contributed from chow consumption and the EtOH

treatment (~9-10 kJ) on the days of gavage were calculated, there was no difference in

energy intake between the EtOH and Control dams (Table 2). Blood glucose

concentrations at 1 h and 5 h post gavage on E13.5 and E14.5 were not significantly

different (Table 2). There were also no differences in pregnancy outcomes between

EtOH and Control dams, such as litter sex ratio, litter size, and number of implantation

scars (Table 2). However, one dam in the EtOH group had a litter of only 7 pups due to

a suspected blockage in one uterine horn (no implantation scars were observed) and so

this litter was excluded from subsequent analyses. BAC was measured in 4 out of 8

Control dams, with BAC below the limit of detection at each time point. In EtOH-

treated dams, mean BAC was ~50 mg/dL (~0.05%) at 1 h following gavage at E13.5

and E14.5, but by 5 h post-gavage, was below the limit of detection on both days (Table

2).

There was no difference in pup weights at PN1 between EtOH and control litters, nor

differences in weight gain until weaning at PN21 (Table 3). For the first week post-

weaning, weight gain was higher in males than in females ($P_{(sex)} = 0.03$) but there was

no effect of treatment (Table 3). Weights of offspring at the PN30 cull were also not

different (Table 3).

***Fasting blood glucose, plasma insulin and lipid levels in adolescent offspring***

Fasting blood glucose, plasma insulin and indexes for insulin resistance (HOMA-IR)

and insulin sensitivity (QUICKI) were not significantly different between groups in

either sex at PN30 (Table 4). Fasting blood glucose (mmol/L) was also not different at

10-11 weeks of age (Male EtOH: $5.5 \pm 0.2$, $n = 9$; Male Control: $5.5 \pm 0.2$, $n = 8$;

Female EtOH: $5.3 \pm 0.1$, n = 9; Female Control: $5.5 \pm 0.1$, $n = 8$. $P_{treat} = 0.69$, $P_{sex} =$

0.51, $P_{int} = 0.57$). The plasma lipid profile (HDL, LDL, TG and TC) was also not

different between EtOH and control offspring at PN30 (Table 4).

***Food preference and water consumption at 4-5 months of age***

There were no significant differences in consumption of SD or HFD between EtOH and

Control offspring at any time (Table 5). However, as the HFD was more energy dense

than the SD (19.4 MJ/kg versus 14.0 MJ/kg), there was a significant decrease in energy

intake per day by the EtOH groups compared to controls, due to a slightly reduced (non-

significant) consumption of HFD by these groups. Animals consumed ~10-fold more

HFD than SD during CP1, with a concomitant 10-fold decrease in SD consumption

compared to baseline levels. This continued through CP2, although HFD consumption

dropped to ~4-fold that of SD. However, there were sex differences in consumption

rates, with females consuming more SD at baseline and more HFD during CP1 and CP2

than males (Table 5).

Daily baseline water and SD consumption was not significantly different between

Control and EtOH offspring in either sex, but females did drink a larger volume of

water compared to males, regardless of treatment (Table 5). EtOH females also

increased their water consumption compared to controls in the 2nd half of the testing

period (CP2, Table 5), with a non-parametric Mann-Whitney U-test performed

separately for each sex indicating a significant difference between control and EtOH

females ($P = 0.003$) but no difference between groups within males ($P = 0.34$).

***Fasting plasma glucose, glucose clearance and insulin sensitivity in adult offspring***

At 6 months of age, fasting plasma glucose continued to be similar between EtOH and

Control groups and between males and females (Figure 3A). However, fasting plasma

insulin was significantly elevated in ETOH-exposed males compared to control males,

with females showing no effect of treatment (Figure 3B). HOMA-IR and QUICKI

indices were also significantly altered in EtOH compared to Control offspring,

indicative of increased insulin resistance and reduced insulin sensitivity (Figure 3C, D).

In the GTT, there was no significant difference in the response to glucose injection (as

shown in Figure 4A) and overall AUGC between EtOH and Control offspring (Figure

4B). Insulin output, as shown by the AUIC, was significantly elevated in males

compared to females during the GTT (Figure 4C, D). This was particularly evident

during the acute first-phase insulin response, with EtOH males having a tendency to

increase insulin output compared to Control male and female offspring, although this

did not reach statistical significance following the non-parametric Dunn's multiple

comparison procedure (Figure 4E). Second phase insulin production continued to be

higher in males than females, but there was no effect of PAE (Figure 4F). In addition,

there were significant differences in glucose clearance in response to exogenous insulin

in the ITT (Figure 4G, H). EtOH male offspring had significantly reduced AUGC

compared to EtOH females by Dunn's multiple comparison procedure, but this group

was also significantly different to control male offspring using a non-parametric Mann-

Whitney test ($P = 0.0079$). However, PAE had no effect on glucose clearance in females

during the ITT (Figure 4H).

### *Body condition and organ weights at 6 months of age*

At the completion of the study, all animals were culled and a subset (1-2 males and/or

females per litter for $n=12$ per sex per group) underwent detailed morphometric

analysis. There was no significant difference in body weight, measures of size (snout-

rump length, tibia length), or measures of body condition (ponderal index, absolute

abdominal circumference and abdominal circumference relative to body size

parameters) between EtOH-exposed and control offspring (Table 6). However, as would

be expected, males were significantly larger than females across all parameters (Table

6). There was also no significant difference in liver or pancreas weight (absolute or

relative to body weight) between EtOH and control offspring. Again, liver weight

(absolute and relative to BW) and absolute pancreas weight were significantly higher in

males than females. However, pancreas weight relative to body weight was significantly

higher in females than males (Table 6).

### *Expression of hepatic genes involved in insulin signalling and glucose homeostasis*

Liver tissue was collected from 6 month-old offspring for analysis of genes involved in

glucose transport (*Glut2*), homeostatic regulation of glucose levels (*G6pc*, *Gck*, *Pck1*

and *Ppargc1a*) and insulin signalling (*Insr*) (Table 7). There was no significant

difference in the expression of these genes between EtOH-exposed and control

offspring (Table 7). However, there was a sex-specific difference in expression of

*Ppargc1a* and *Insr*, and a trend for an effect of sex on *Glut2* expression, with all three

genes displaying higher relative expression in females compared with males (Table 7).

### *Hepatic and peripheral tissue insulin signalling*

Given that insulin resistance was only observed in EtOH-exposed males compared to

control males, and there was no effect of EtOH exposure on females, we focused our

molecular analysis of insulin signalling on male offspring. We examined a key molecule

involved in the insulin signalling pathway both centrally and in peripheral tissues, AKT,

also known as protein kinase B. In liver (Figure 5A-E) and skeletal muscle (Figure 5F-

J), there was no significant difference in total (pan-)AKT, $pAKT_{Ser473}$ (absolute or as a

ratio of pan-AKT) or $pAKT_{Thr308}$ (absolute or as a ratio of pan-AKT) between EtOH-

exposed and control male offspring. Muscle gene expression of the glucose transporter,

*Glut4*, and the insulin receptor, *Insr*, were also not affected by EtOH exposure (Table

8). However, there was a significant increase in pan-AKT in adipose tissue from EtOH-

exposed males compared to controls (Figure 5K) and a tendency for increased

$pAKT_{Thr308}$ (Figure 5M, $P = 0.07$), but not as a ratio of pan-AKT (Figure 5O).

Expression of genes involved in glucose transport (*Glut 4*) and insulin signalling both

upstream (*Insr*) and downstream (*Foxo1*, *Gsk3β*) of AKT were not altered in adipose

tissue from EtOH-exposed male offspring (Table 8).

**Discussion**

This study demonstrates for the first time that a low, acute dose of prenatal EtOH can

program metabolic dysfunction in a sex-specific manner. Male offspring developed

insulin resistance at 6 months of age, regardless of the lack of traditional hallmarks of

low birth weight and catch-up growth seen in fetal programming. PAE male offspring

also showed alterations in insulin signalling in adipose tissue. Given that BAC levels

following alcohol exposure were relatively moderate, at ~0.05%, these results

strengthen the message that no amount of alcohol during pregnancy is safe for long-

term offspring health.

Previous preclinical studies investigating the association between PAE and offspring

metabolic health have typically used a high dose (2 g/kg BW administered twice a day)

throughout gestation and found this can induce glucose intolerance and insulin

resistance (see (Akison *et al.*, 2019) for review). This dosage typically results in a peak

BAC of ~0.1-0.15% (Chen & Nyomba, 2003b). However, our moderate, acute dose

model, resulting in a BAC of just 0.05% one hour after gavage, is more representative

of drinking practices in pregnant women who report drinking on average 1-2 standard

drinks sporadically during pregnancy. Although the model used in this study has been

previously used by Gray et al. (Gray *et al.*, 2010), in the same species and strain, they

reported a BAC of 0.107% in treated dams 1 h after gavage. The reasons for these

differences in BAC are unknown but given these discrepancies, this highlights the

importance of measuring and reporting BAC levels, in addition to providing the dosage

administered.

In our moderate exposure model, there were no changes in fasting blood glucose levels

at any age, or fasting insulin levels in adolescent offspring exposed to prenatal alcohol.

Blood glucose concentrations during a GTT performed at 6 months of age were also not

altered. However, the major finding of our study was that male offspring exposed to

alcohol showed evidence of insulin resistance at 6 months of age, with elevated fasting

plasma insulin, resulting in an increased HOMA-IR index, indicative of insulin

resistance, and a decreased QUICKI index, indicative of reduced insulin sensitivity.

There was also a significant impairment in glucose clearance following an insulin

challenge. These are hallmarks of a pre-diabetic state. The trend for increased first

phase insulin secretion during the GTT in PAE males suggests an appropriate

compensatory response to this reduced insulin sensitivity, resulting in maintenance of

blood glucose to the same level as controls.  In a recent systematic review, 12

preclinical studies that examined glucose metabolism using a GTT and/or ITT found

evidence of insulin resistance in all but one study (Akison *et al.*, 2019). However, the

majority of these studies also reported changes in circulating glucose levels following

PAE. Interestingly, only one of these 12 studies had a low BAC (peak of ~0.03%),

comparable to that measured in our study. That study, using a chronic low dose of

alcohol throughout pregnancy, also found increased first phase insulin secretion in

response to a glucose challenge exclusively in males, without concomitant changes in

blood glucose levels (Probyn *et al.*, 2013). This suggests that the BAC may need to

reach a higher threshold to induce changes in glucose tolerance, while changes in

insulin sensitivity can potentially manifest following exposure to a much lower BAC.

Alternatively, aging may be an important '2$^{nd}$ hit' for a more overt, diabetic phenotype.

Adolescent (PN30) and early adult (10-11 week-old) offspring in our study did not have

any significant metabolic phenotype, and it was only at 6 months of age when insulin

resistance was detected. We predict that examination of offspring further aged to 12 or

18 months may also exhibit glucose intolerance, as well as insulin resistance, although

this needs to be tested.

We explored potential molecular mechanisms underlying the observed insulin resistant

phenotype. Given the importance of the liver for glucose homeostasis, we examined the

expression of a selection of hepatic genes involved in glucose transport (*Glut2*), glucose

homeostasis (*G6pc, Gck, Pck1, Ppargc1a*) and insulin signalling (*Insr*). Similar to our

low dose chronic alcohol exposure model (Probyn *et al.*, 2013), we found no

dysregulation of these genes due to PAE, and instead observed only sex-specific

differences in gene expression. We also examined AKT protein expression and

activation via phosphorylation in the liver but so no evidence of central insulin

resistance This suggested the deficits lay within peripheral tissues, skeletal muscle or

adipose tissue, with PI3K/AKT signalling known to play an essential role in insulin-

mediated glucose uptake in these peripheral tissues (Huang *et al.*, 2018). Supporting

this, we did find moderate alterations in AKT levels in adipose tissue, both total AKT

and activated AKT, measured by phosphorylation at Threonine$_{308}$. Both were

significantly elevated in adipose tissue collected from PAE male offspring at 6 months

of age. Intriguingly, our results are consistent to a previous study investigating insulin

signalling in offspring exposed to periconceptional alcohol (Gardebjer *et al.*, 2015). In

this previous study, both males and females were insulin resistant but there were

inconsistent results with regards to AKT protein levels and activation. In females, AKT

levels and phosphorylation were reduced, as you would expect given the phenotype.

However, in EtOH-exposed male offspring, total AKT levels were elevated, as we have

seen here. Also, pAKT(Thr$_{308}$) normalised to total AKT was increased in these animals.

This suggests that increased total AKT and pAKT(Thr$_{308}$) may be a potential

compensatory mechanism, operating exclusively in males, in response to increased

fasting insulin levels. This hypothesis requires further investigation. We further

interrogated the insulin signalling pathway both upstream (*Insr*) and downstream

(*Foxo1* and *Gsk3β*) in the AKT signalling pathway but found no changes in gene

expression due to ethanol exposure. We also found no differences in muscle expression

of *Glut4* or AKT signalling that have been demonstrated in other models of prenatal

alcohol exposure (Chen & Nyomba, 2003a; Chen & Nyomba, 2003b; Chen *et al.*, 2005;

Yao *et al.*, 2006), although this was only seen when tissues were collected immediately

following a glucose challenge and not when animals were in a random-fed state.

Therefore, our results and others suggest that adipose tissue is the site of insulin

resistance in male offspring in response to prenatal alcohol exposure, although further

work is needed to clarify the nature of this relationship.

Maternal malnutrition, even if relatively modest, is an established prenatal perturbation

which may result in fetal growth restriction and offspring metabolic disorders (Flynn *et*

*al.*, 2013; Gao *et al.*, 2014). However, this can be excluded in our study, as food and

water consumption throughout pregnancy did not differ between the EtOH and control

dams. Additionally, there were no differences in other maternal parameters known to

affect offspring metabolic outcome, such as obesity (Whitaker, 2004; El-Gilany &

Hammad, 2010; Janjua *et al.*, 2012), gestational weight gain (Margerison Zilko *et al.*,

2010) and blood glucose (used as an indicator for gestational diabetes) (Van Wootten &

Turner, 2002). Previous preclinical studies have reported reduced birth weight and

subsequent 'catch-up' growth in offspring exposed to prenatal alcohol that was

associated with subsequent metabolic dysfunction, specifically insulin resistance, in

adults (Chen & Nyomba, 2003b; Yao *et al.*, 2006; Dobson *et al.*, 2014; Xia *et al.*, 2014;

Gardebjer *et al.*, 2015). However, other studies highlight that growth restriction is not

required to program chronic disease, with insulin resistance arising in PAE offspring

irrespective of birth weight (Chen & Nyomba, 2004; Yao & Nyomba, 2008; Probyn *et*

*al.*, 2013; Yao *et al.*, 2013). Clinical studies have also reported that growth restriction is

not a prerequisite for development of metabolic syndrome (Euser *et al.*, 2010; Kopec *et*

*al.*, 2017). We also found no effect of PAE on offspring weight or growth, and yet

EtOH-exposed males developed insulin resistance at 6 months of age. This highlights

that even low-dose exposures can program offspring disease, irrespective of birth

weight.

Although our low-dose PAE model did not affect body condition, assessed from

morphometric analysis, at 6 months of age, we did not directly measure body fat.

Studies have previously assessed body composition in PAE offspring using dual-energy

X-ray absorptiometry (DEXA) and/or magnetic resonance imaging (MRI), with variable

results. Our result is similar to that found in a previous low-dose model in rats (Probyn

*et al.*, 2013), as well as a study in mice with EtOH exposure restricted to late gestation

(Amos-Kroohs *et al.*, 2018). However, other higher dose models have reported

increased adiposity in PAE offspring compared to controls, particularly in males

(Dobson *et al.*, 2012; Gardebjer *et al.*, 2018; Zhang *et al.*, 2018).

PAE offspring at 4 months of age did not exhibit a preference for a HFD. However,

they did eat slightly less than controls, which resulted in a significantly lower energy

consumption in PAE offspring due to the HFD being more energy rich. This was

particularly evident in the second choice period, when the novelty of the new diet had

presumably waned (Dorey *et al.*, 2018). This was contrary to a recent study in rat

offspring exposed to a higher dose of alcohol exclusively around conception, which

found that males preferentially consumed more HFD than controls (Dorey *et al.*, 2018).

However, this study was conducted in much older rats (15 months of age). Clinical

studies have found that children diagnosed with FASD tend to have hyperphagia and

abnormal eating patterns, including constant snacking and impaired satiety (Werts *et al.*,

2014; Amos-Kroohs *et al.*, 2016). Our results are more consistent with a study

conducted in mice, which found no changes in food preference as a result of PAE

(Amos-Kroohs *et al.*, 2018). Interestingly, that study also restricted PAE to later in

gestation (E12.5-17.5). Taken together, these studies and ours suggest that a change in

eating behaviour is not consistently associated with prenatal alcohol exposure but may

relate to the timing and severity of the exposure as well as offspring age. Our study also

suggests that altered food consumption or a propensity for consuming a HFD does not

underlie the metabolic dysfunction observed in male offspring. Future studies

examining the impact of PAE on offspring metabolism should include analysis of

feeding behaviours to further define any associations.

A possible limitation of this study was that the saline gavage did not control for the

additional energy provided by the EtOH. A recent study highlighted that the typical

maltodextrin control, used as a substitute for the calories provided by EtOH, is not

metabolised in the same way, and when a medium chain triglycerides solution was

included as a control, no changes in metabolic outcomes could be attributed to PAE

(Amos-Kroohs *et al.*, 2018). However, our dose of EtOH per gavage was so small (~9-

10 kJ), that when we calculated the combined energy intake attributed to the EtOH and

chow on the days of gavage, there was no significant difference between the EtOH and

Control dams in our study. Therefore, our saline treatment adequately controlled for the

potential effects of the gavage treatment and did not result in a lower overall energy

intake when compared with our low-dose EtOH treatment group.

*Conclusions*

This study supports the programming of offspring metabolic health as a result of a

relatively moderate PAE during a critical period of development. Although ethanol was

administered during mid-pregnancy in the rat, this equates to ~week 10-12 of a human

pregnancy when a woman may not yet be aware she is pregnant, particularly if the

pregnancy was not planned. The pre-diabetic, insulin resistant phenotype reported in

male offspring of PAE dams occurred despite a relatively low BAC and standard

postnatal care, and without the '2$^{nd}$ Hit' of a high fat diet as used by others to 'unmask'

disease. Programmed alterations were also independent of maternal parameters known

to affect offspring metabolic health, such as gestational diabetes and obesity. Therefore,

this study highlights the importance of abstaining from alcohol consumption during

pregnancy or when planning a pregnancy.

**Additional information**

*Competing interests*

The authors have no conflicts of interest to disclose.

*Author contributions*

All experiments were performed in the Developmental Programming in Disease Lab,

School of Biomedical Sciences, The University of Queensland. K.M.M. and L.K.A.

were responsible for the conception and design of the experiments. T.M.T.N., L.K.A.

and S.E.S. were responsible for the collection of the data. All authors were involved in

data analysis and/or interpretation. All authors contributed to drafting the work and/or

revising it critically for important intellectual content. All authors approve of the final

version of the manuscript and agree to be accountable for all aspects of the work in

ensuring that questions related to the accuracy or integrity of any part of the work are

appropriately investigated and resolved. All persons designated as authors qualify for

authorship, and all those who qualify for authorship are listed.

*Funding*

Funding for this project was provided by the University of Queensland Early Career

Researcher Grants Scheme (to L.K.A.) and the National Health and Medical Research

Council (to K.M.M., APP1078164).

**Acknowledgements**

We would like to acknowledge Natasha Steiger (Animal Endocrinology Lab, School of

Biomedical Sciences, University of Queensland) for analysis of plasma insulin; Dave

Herne and Barb Arnts (University of Queensland Biological Resources) for assistance

with animal treatments and husbandry; Elizabeth McReight (School of Biomedical

Sciences, University of Queensland) for assistance with animal work; and Jacobus

Ungerer (Pathology Queensland, Queensland Health) for analysis of BAC.

**Supporting information**

All underlying data used in figures and tables reported in this study and can be accessed

from the following URL: https://doi.org/10.5281/zenodo.3406863.

**References**

Akison LK, Reid N, Wyllie M & Moritz KM. (2019). Adverse health outcomes in
    offspring associated with fetal alcohol exposure: A systematic review of clinical
    and preclinical studies with a focus on metabolic and body composition
    outcomes. *Alcohol Clin Exp Res*, DOI: 10.1111/acer.14078.

Allison DB, Paultre F, Maggio C, Mezzitis N & Pi-Sunyer FX. (1995). The use of areas
    under curves in diabetes research. *Diabetes Care* **18,** 245-250.

Amos-Kroohs RM, Fink BA, Smith CJ, Chin L, Van Calcar SC, Wozniak JR & Smith
    SM. (2016). Abnormal eating behaviors are common in children with fetal
    alcohol spectrum disorder. *J Pediatr* **169,** 194-200 e191.

Amos-Kroohs RM, Nelson DW, Hacker TA, Yen C-LE & Smith SM. (2018). Does

prenatal alcohol exposure cause a metabolic syndrome?(Non-) evidence from a

mouse model of fetal alcohol spectrum disorder. *PLoS One* **13,** e0199213.

Barker DJ. (2007). The origins of the developmental origins theory. *J Intern Med* **261,**

412-417.

Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan

T, Pfaffl MW, Shipley GL, Vandesompele J & Wittwer CT. (2009). The MIQE

guidelines: minimum information for publication of quantitative real-time PCR

experiments. *Clin Chem* **55,** 611-622.

Cacho J, Sevillano J, de Castro J, Herrera E & Ramos MP. (2008). Validation of simple

indexes to assess insulin sensitivity during pregnancy in Wistar and Sprague-

Dawley rats. *Am J Physiol Endocrinol Metab* **295,** E1269-1276.

Castells S, Mark E, Abaci F & Schwartz E. (1981). Growth retardation in fetal alcohol

syndrome. Unresponsiveness to growth-promoting hormones. *Dev Pharmacol*

*Ther* **3,** 232-241.

Chen L & Nyomba B. (2003a). Glucose intolerance and resistin expression in rat

offspring exposed to ethanol in utero: modulation by postnatal high-fat diet.

*Endocrinology* **144,** 500-508.

Chen L & Nyomba BLG. (2003b). Effects of prenatal alcohol exposure on glucose

tolerance in the rat offspring. *Metab Clin Exp* **52,** 454-462.

Chen L & Nyomba BLG. (2004). Whole body insulin resistance in rat offspring of
mothers consuming alcohol during pregnancy or lactation: comparing prenatal
and postnatal exposure. *J Appl Physiol* **96,** 167-172.

Chen L, Yao XH & Nyomba BL. (2005). In vivo insulin signaling through PI3-kinase is
impaired in skeletal muscle of adult rat offspring exposed to ethanol in utero. *J*
*Appl Physiol* **99,** 528-534.

Dickinson H, Moss TJ, Gatford KL, Moritz KM, Akison L, Fullston T, Hryciw DH,
Maloney CA, Morris MJ, Wooldridge AL, Schjenken JE, Robertson SA,
Waddell BJ, Mark PJ, Wyrwoll CS, Ellery SJ, Thornburg KL, Muhlhausler BS
& Morrison JL. (2016). A review of fundamental principles for animal models
of DOHaD research: an Australian perspective. *J Dev Orig Health Dis* **7,** 449-
472.

Dobson CC, Mongillo DL, Brien DC, Stepita R, Poklewska-Koziell M, Winterborn A,
Holloway AC, Brien JF & Reynolds JN. (2012). Chronic prenatal ethanol
exposure increases adiposity and disrupts pancreatic morphology in adult guinea
pig offspring. *Nutr Diabetes* **2,** e57.

Dobson CC, Thevasundaram K, Mongillo DL, Winterborn A, Holloway AC, Brien JF
& Reynolds JN. (2014). Chronic prenatal ethanol exposure alters expression of
central and peripheral insulin signaling molecules in adult guinea pig offspring.
*Alcohol* **48,** 687-693.

Dorey ES, Cullen CL, Lucia D, Mah KM, Manchadi MR, Muhlhausler BS & Moritz
KM. (2018). The impact of periconceptional alcohol exposure on fat preference
and gene expression in the mesolimbic reward pathway in adult rat offspring. *J*
*Dev Orig Health Dis* **9,** 223-231.

El-Gilany A-H & Hammad S. (2010). Body mass index and obstetric outcomes in Saudi
Arabia: a prospective cohort study. *Ann Saudi Med* **30,** 376-380.

Euser AM, Dekker FW & Hallan SI. (2010). Intrauterine growth restriction: no unifying
risk factor for the metabolic syndrome in young adults. *Eur J Cardiovasc Prev*
*Rehabil* **17,** 314-320.

Fleming TP, Watkins AJ, Velazquez MA, Mathers JC, Prentice AM, Stephenson J,
Barker M, Saffery R, Yajnik CS, Eckert JJ, Hanson MA, Forrester T, Gluckman
PD & Godfrey KM. (2018). Origins of lifetime health around the time of
conception: causes and consequences. *Lancet* **391,** 1842-1852.

Flynn ER, Alexander BT, Lee J, Hutchens ZM, Jr. & Maric-Bilkan C. (2013). High-
fat/fructose feeding during prenatal and postnatal development in female rats
increases susceptibility to renal and metabolic injury later in life. *Am J Physiol*
*Regul Integr Comp Physiol* **304,** R278-285.

Friedewald WT, Levy RI & Fredrickson DS. (1972). Estimation of the concentration of
low-density lipoprotein cholesterol in plasma, without use of the preparative
ultracentrifuge. *Clin Chem* **18,** 499-502.

Gao F, Liu Y, Li L, Li M, Zhang C, Ao C & Hou X. (2014). Effects of maternal
undernutrition during late pregnancy on the development and function of ovine
fetal liver. *Anim Reprod Sci* **147,** 99-105.

Gardebjer EM, Anderson ST, Pantaleon M, Wlodek ME & Moritz KM. (2015).
Maternal alcohol intake around the time of conception causes glucose
intolerance and insulin insensitivity in rat offspring, which is exacerbated by a
postnatal high-fat diet. *FASEBJ* **29,** 2690-2701.

Gardebjer EM, Cuffe JSM, Ward LC, Steane S, Anderson ST, Dorey ES, Kalisch-Smith
JI, Pantaleon M, Chong S, Yamada L, Wlodek ME, Bielefeldt-Ohmann H &
Moritz KM. (2018). The effects of periconceptional maternal alcohol intake and
a postnatal high-fat diet on obesity and liver disease in male and female rat
offspring. *Am J Physiol Endocrinol Metab* **315,** E694-E704.

Gittes GK. (2009). Developmental biology of the pancreas: a comprehensive review.
*Dev Biol* **326,** 4-35.

Godlewski G, Gaubert-Cristol R, Rouy S & Prudhomme M. (1997). Liver development
in the rat and in man during the embryonic period (Carnegie stages 11-23).
*Microsc Res Tech* **39,** 314-327.

Gray SP, Denton KM, Cullen-McEwen L, Bertram JF & Moritz KM. (2010). Prenatal
exposure to alcohol reduces nephron number and raises blood pressure in
progeny. *J Am Soc Nephrol* **21,** 1891-1902.

Grundy D. (2015). Principles and standards for reporting animal experiments in The
Journal of Physiology and Experimental Physiology. *J Physiol* **593,** 2547-2549.

Heindel JJ & Vandenberg NL. (2015). Developmental origins of health and disease: a
paradigm for understanding disease cause and prevention. *Curr Opin Pediatr* **27,**
248-253.

Huang X, Liu G, Guo J & Su Z. (2018). The PI3K/AKT pathway in obesity and type 2
diabetes. *Int J Biol Sci* **14,** 1483-1496.

Janjua NZ, Mahmood B, Islam MA & Goldenberg RL. (2012). Maternal and early
childhood risk factors for overweight and obesity among low-income

predominantly black children at age five years: A prospective cohort study. *J*

*Obes* **2012,** 457173.

Jaramillo LM, Balcazar IB & Duran C. (2012). Using vaginal wall impedance to

determine estrous cycle phase in Lewis rats. *Lab Animal* **41,** 122.

Katz A, Nambi SS, Mather K, Baron AD, Follmann DA, Sullivan G & Quon MJ.

(2000). Quantitative insulin sensitivity check index: a simple, accurate method

for assessing insulin sensitivity in humans. *J Clin Endocrinol Metab* **85,** 2402-

2410.

Kilkenny C & Altman DG. (2010). Improving bioscience research reporting: ARRIVE-

ing at a solution. *Lab Anim* **44,** 377-378.

Kilkenny C, Browne W, Cuthill IC, Emerson M, Altman DG & Group NCRRGW.

(2010). Animal research: reporting in vivo experiments: the ARRIVE

guidelines. *Br J Pharmacol* **160,** 1577-1579.

Kopec G, Shekhawat PS & Mhanna MJ. (2017). Prevalence of diabetes and obesity in

association with prematurity and growth restriction. *Diabetes Metab Syndr Obes*

**10,** 285-295.

Mackenzie RW & Elliott BT. (2014). Akt/PKB activation and insulin signaling: a novel

insulin signaling pathway in the treatment of type 2 diabetes. *Diabetes Metab*

*Syndr Obes* **7,** 55-64.

Margerison Zilko CE, Rehkopf D & Abrams B. (2010). Association of maternal

gestational weight gain with short- and long-term maternal and child health

outcomes. *Am J Obstet Gynecol* **202,** 574.e571-578.

McCormack C, Hutchinson D, Burns L, Wilson J, Elliott E, Allsop S, Najman J, Jacobs
S, Rossen L, Olsson C & Mattick R. (2017). Prenatal alcohol consumption
between conception and recognition of pregnancy. *Alcohol Clin Exp Res* **41,**
369-378.

Muggli E, O'Leary C, Donath S, Orsini F, Forster D, Anderson PJ, Lewis S, Nagle C,
Craig JM, Elliott E & Halliday J. (2016). "Did you ever drink more?" A detailed
description of pregnant women's drinking patterns. *BMC Public Health* **16,** 683.

Nagy C & Einwallner E. (2018). Study of in vivo glucose metabolism in high-fat diet-
fed mice using oral glucose tolerance test (OGTT) and insulin tolerance test
(ITT). *J Vis Exp* **131,** e56672.

National Health and Medical Research Council. (2009). Australian Guidelines to
Reduce Health Risks from Drinking Alcohol, pp. 181. Commonwealth of
Australia, Canberra, ACT.

Pan FC & Brissova M. (2014). Pancreas development in humans. *Curr Opin Endocrinol*
*Diabetes Obes* **21,** 77-82.

Popova S, Lange S, Probst C, Gmel G & Rehm J. (2017). Estimation of national,
regional, and global prevalence of alcohol use during pregnancy and fetal
alcohol syndrome: a systematic review and meta-analysis. *Lancet Glob Health*
**5,** e290-e299.

Probyn ME, Parsonson KR, Gardebjer EM, Ward LC, Wlodek ME, Anderson ST &
Moritz KM. (2013). Impact of low dose prenatal ethanol exposure on glucose
homeostasis in Sprague-Dawley rats aged up to eight months. *PLoS One* **8,**
e59718.

Smith S, Amos-Kroohs R, Chin L, Fink B & Wozniak J. (2015). Disordered eating

behaviors and nutritional issues in children with Fetal Alcohol Spectrum

Disorders (FASD). *FASEBJ* **29**.

Sundrani DP, Roy SS, Jadhav AT & Joshi SR. (2017). Sex-specific differences and

developmental programming for diseases in later life. *Reprod Fertil Dev* **29,**

2085-2099.

Van Wootten W & Turner RE. (2002). Macrosomia in neonates of mothers with

gestational diabetes is associated with body mass index and previous gestational

diabetes. *J Am Diet Assoc* **102,** 241-243.

Weinberg J, Sliwowska JH, Lan N & Hellemans KG. (2008). Prenatal alcohol exposure:

foetal programming, the hypothalamic-pituitary-adrenal axis and sex differences

in outcome. *J Neuroendocrinol* **20,** 470-488.

Werts RL, Van Calcar SC, Wargowski DS & Smith SM. (2014). Inappropriate feeding

behaviors and dietary intakes in children with fetal alcohol spectrum disorder or

probable prenatal alcohol exposure. *Alcohol Clin Exp Res* **38,** 871-878.

Whitaker RC. (2004). Predicting preschooler obesity at birth: the role of maternal

obesity in early pregnancy. *Pediatrics* **114,** e29-36.

World Health Organisation. (2004). Department of Mental Health and Substance

Abuse: Alcohol Policy. Geneva.

Xia L, Shen L, Kou H, Zhang B, Zhang L, Wu Y, Li X, Xiong J, Yu Y & Wang H.

(2014). Prenatal ethanol exposure enhances the susceptibility to metabolic

syndrome in offspring rats by HPA axis-associated neuroendocrine metabolic

programming. *Toxicol Lett* **226,** 98-105.

Yao X-H, Nguyen HK & Nyomba BG. (2013). Prenatal ethanol exposure causes

glucose intolerance with increased hepatic gluconeogenesis and histone

deacetylases in adult rat offspring: reversal by tauroursodeoxycholic acid. *PLoS*

*One* **8,** e59680.

Yao XH, Chen L & Nyomba BL. (2006). Adult rats prenatally exposed to ethanol have

increased gluconeogenesis and impaired insulin response of hepatic

gluconeogenic genes. *J Appl Physiol* **100,** 642-648.

Yao XH & Nyomba BLG. (2008). Hepatic insulin resistance induced by prenatal

alcohol exposure is associated with reduced PTEN and TRB3 acetylation in

adult rat offspring. *Am J Physiol Regul Integr Comp Physiol* **294,** R1797-R1806.

Zhang CR, Kurniawan ND, Yamada L, Fleming W, Kaminen-Ahola N, Ahola A,

Galloway G & Chong S. (2018). Early gestational ethanol exposure in mice:

Effects on brain structure, energy metabolism and adiposity in adult offspring.

*Alcohol* **75,** 1-10.

**Figure legends**

**Figure 1: Flow chart of animals treated and offspring used in each experiment.**

Pregnant dams were treated with either saline or 1g/kg body weight ethanol (EtOH) via

gavage. Note that one EtOH litter was excluded from further experiments given the

litter size was 7 (mean, range for other litters: 15, 11-19). $^\wedge$ <8 females available due to

litter size/sex-ratio constraints and use in another study; available females were

prioritised for adult experiments; 2 litters could not be used in plasma analyses as not

fasted. $^{\wedge\wedge}$ <9 females available due to litter size/sex-ratio constraints and use in another

study; available females were prioritised for adult experiments; 1 litter had no animals

used at this time-point. $^+$ Only males analysed; sample size reduced to allow all samples

to be run on one gel. F = female; M = male; E = embryonic day; BW = body weight;

PM = post-mortem; GTT = glucose tolerance test; ITT = insulin tolerance test; qPCR =

real-time quantitative polymerase chain reaction.

**Figure 2: Comparison of blood glucose concentrations measured by glucometer**

**and plasma glucose concentrations measured by Cobas analyser.** Dual-analysis was

conducted on $n = 8$ control litters and $n = 9$ ethanol litters, with 1 male and 1 female

analysed per litter. (A) Bland-Altman plot to visualise the difference in the glucometer

and Cobas readings versus the average of the readings. (B) Comparison of glucose

levels across each treatment/sex group by method. White bars indicate control and black

bars indicate ethanol-exposed (EtOH) animals within each sex. Data analysed using a

two-way ANOVA, with treatment/sex (Control Male, EtOH Male, Control Female,

EtOH Female) and method (glucometer/Cobas) as factors.

**Figure 3: Effect of prenatal alcohol exposure on fasting plasma glucose and plasma**

**insulin levels in adult offspring.** Offspring were prenatally exposed to ethanol (grey)

or saline control (white). (A) Fasting plasma glucose and (B) plasma insulin were

measured at 6 months of age. Corresponding homeostatic model assessment of insulin

resistance (HOMA-IR) (C) and quantitative insulin-sensitivity check index (QUICKI)

(D) were calculated. Data are presented as mean ± SEM. $n = 8$ control litters and $n = 9$

ethanol litters, with 1 male and 1 female analysed per litter. Data for A) and B) were

analysed with two-way ANOVA; while data for C) and D) were analysed using a non-

parametric Kruskal-Wallis test due to non-normal data distribution (indicated by ^). **

= significantly different to control male by Sidak's multiple comparison post-hoc

procedure.

**Figure 4: Effect of prenatal alcohol exposure on glucose clearing and insulin**

**sensitivity.** Offspring were prenatally exposed to ethanol (grey) or saline (white). (A)

Blood glucose concentration curve during a glucose tolerance test (GTT); (B) area

under the glucose curve (AUGC) generated from the GTT; (C) plasma insulin secretion

during the GTT; (D) area under the insulin curve (AUIC) generated from the GTT; (E)

AUIC for 1st phase insulin secretion; and (F) AUIC for 2nd phase insulin secretion. (G)

Blood glucose concentrations following an insulin tolerance test (ITT) and (H) AUGC

from inverted ITT curves. Control males (white circles), ethanol males (black circles),

control females (white squares) and ethanol females (black squares) (A, C, G). Data

represented as mean ± SEM; $n = 8$ control litters and $n = 9$ ethanol litters, with 1 male

and 1 female analysed per litter. Data were analysed with two-way ANOVA (B, D, F)

or a non-parametric Kruskal-Wallis test (indicated by ^) due to non-normal data

distribution (E, H). # indicates significant difference between sexes within treatment

($P<0.05$), where significance was determined by Dunn's multiple comparisons test.

**Figure 5: Effect of prenatal alcohol on AKT protein levels in liver and peripheral**

**tissues in male offspring at 6 months of age.** Protein levels and phosphorylation state,

expressed relative to ACTB, in liver (A-E) and expressed relative to GAPDH in

gastrocnemius muscle (F-J) and adipose tissue (K-O). Offspring were prenatally

exposed to ethanol (grey) or saline (white). Data represented as mean ± SEM. $n = 7$

litters per group with 1 male per litter. Data were analysed with an unpaired t-test

(normally distributed data) or a Mann-Whitney U-test (non-normal distribution,

indicated by ^). + indicates Welch's correction was applied due to unequal variances.

ACTB = β-actin; GAPDH = glyceraldehyde 3-phosphate dehydrogenase.

**Table 1: Primers used for real-time quantitative PCR analysis of metabolic gene expression.** All primers were Assay-on-Demand primer/probe sets from ThermoFisher Scientific (Richlands, QLD, Australia).

| Gene name | Gene symbol | Assay ID | Amplicon size (bp) | Accession number(s) |
|---|---|---|---|---|
| Beta actin[*] | *Actb* | Cat #4351319 | 61 | NM_031144.3 |
| Eukaryotic translation initiation factor 2A[*] | *Eif2a* | Rn01494813_m1 | 82 | NM_001109339.1 |
| Forkhead box O1 | *Foxo1* | Rn01494868_m1 | 99 | NM_001191846.2 |
| Glucose-6-phosphatase, catalytic subunit | *G6pc* | Rn00689876_m1 | 64 | NM_013098.2 |
| Glucokinase | *Gck* | Rn00561265_m1 | 58 | NM_001270849.1 NM_001270850.1 NM_012565.2 |
| Glucose transporter 2 [solute carrier family 2 (facilitated glucose transporter), member 2] | *Glut2 (Slc2a2)* | Rn00563565_m1 | 76 | NM_012879.2 |
| Glucose transporter 4 | *Glut4* | Rn00562597_m1 | 75 | NM_012751.1 |
| Glycogen synthase kinase 3 beta | *Gsk3β* | Rn_00583429_m1 | 71 | NM_032080.1 |
| Insulin receptor | *Insr* | Rn01403321_m1 | 76 | NM_017071.2 |
| Phosphoenolpyruvate carboxykinase 1 | *Pck1* | Rn01529014_m1 | 87 | NM_198780.3 |
| Peroxisome proliferator-activated receptor gamma, coactivator 1 alpha | *Ppargc1a* | Rn00580241_m1 | 94 | NM_031347.1 |
| Ribosomal protein L19[*] | *Rpl19* | Rn00821265_g1 | 57 | NM_031103.1 |

[*] Endogenous control.

**Table 2: Maternal parameters for dams treated with saline (Control) or ethanol (EtOH).** All animals treated by oral gavage on embryonic day 13.5 and 14.5.

| Maternal parameters | Treatment group | | *P* value |
| | Control ($n = 8$) | EtOH ($n = 10$) | |
| --- | --- | --- | --- |
| Body weight when mated (g) | $251 \pm 4$ | $255 \pm 4$ | 0.61 |
| Body weight at E13.5 (g) | $310 \pm 5$ | $312 \pm 5$ | 0.77 |
| Total weight gain (g)* | $152 \pm 9$ | $148 \pm 5$ | 0.98 |
| Pre-gavage (mating to E12.5) | $59 \pm 3$ | $58 \pm 4$ | 0.85 |
| Post-gavage (E15.5 to birth) | $91 \pm 6$ | $89 \pm 5$ | 0.77 |
| $H_2O$ consumption (mL/day) | | | |
| Pre-gavage (E12.5) | $31.9 \pm 1.7$ | $29.6 \pm 1.8$ | 0.37 |
| During gavage (E13.5-E14.5) | $25.8 \pm 1.3$ | $25.3 \pm 1.4$ | 0.79 |
| Post-gavage (E15.5 to birth) | $36.9 \pm 1.1$ | $34.8 \pm 1.6$ | 0.33 |
| Chow consumption (g/day) | | | |
| Pre-gavage (E12.5) | $24.7 \pm 1.2$ | $24.4 \pm 0.8$ | 0.79 |
| During gavage (E13.5-E14.5) | $20.9 \pm 1.0$ | $20.4 \pm 1.0$ | 0.70 |
| Post-gavage (E15.5 to birth) | $26.6 \pm 0.7$ | $26.0 \pm 0.9$ | 0.61 |
| Energy consumption (kJ/day) | | | |
| E13.5* | $299.3 \pm 14.6$ | $283.4 \pm 6.2$ | 0.70 |
| E14.5 | $286.0 \pm 18.8$ | $305.0 \pm 24.0$ | 0.56 |
| Blood glucose (mmol/L) | | | |
| 1h post E13.5 gavage* | $5.8 \pm 0.2$ | $6.1 \pm 0.2$ | 0.13 |
| 5h post E13.5 gavage | $5.9 \pm 0.1$ | $6.1 \pm 0.2$ | 0.38 |
| 1h post E14.5 gavage | $5.7 \pm 0.2$ | $6.0 \pm 0.1$ | 0.09 |
| 5h post E14.5 gavage* | $6.2 \pm 0.2$ | $6.0 \pm 0.1$ | 0.62 |
| Litter sex ratio (M:F) | $1.1 \pm 0.2$ | $1.2 \pm 0.2$ | 0.78 |
| Litter size | 15[†] | 14[†] | 0.47 |
| Number of implantation scars* | 16[†] | $15 \pm 1$[†] | 0.99 |
| Blood alcohol concentration (BAC) (mg/dL) | | | |

| | | | |
|---|---|---|---|
| 1h post E13.5 gavage | <LD ‡ | 48.2 ± 7.5 | - |
| 5h post E13.5 gavage | <LD ‡ | <LD | - |
| 1h post E14.5 gavage | <LD ‡ | 52.9 ± 4.5 | - |
| 5h post E14.5 gavage | <LD ‡ | <LD | - |

Data are presented as mean ± SEM. *P* values were obtained using an unpaired Student's t-test.

<LD = below the limit of detection; E = embryonic day; EtOH = ethanol; M = male; F = female.

* Indicates data not normally distributed and analysed with a non-parametric Mann-Whitney U-test.

† Denotes parameters expressed as whole pups/scars.

‡ Measured in 4 animals only.

**Table 3: Postnatal weight and growth of offspring from saline (Control) and ethanol (EtOH) treated litters during the first month.**

All offspring from dams treated by oral gavage at embryonic day 13.5 and 14.5. Weights at PN1 and PN30 are average body weight (g). At other ages, average total weight gain over the specified age ranges are shown (g).

| | Male | | Female | | Statistics | | |
|---|---|---|---|---|---|---|---|
| | **Control** | **EtOH** | **Control** | **EtOH** | | | |
| **Age** | ($n = 8$) | ($n = 9$) | ($n = 8$) | ($n = 9$) | $P_{(trt)}$ | $P_{(sex)}$ | $P_{(int)}$ |
| PN1$^*$ | $6.6 \pm 0.1$ | $7.0 \pm 0.3$ | $6.4 \pm 0.1$ | $6.7 \pm 0.3$ | 0.15 | 0.32 | 0.84 |
| PN1-10 | $10.2 \pm 0.4$ | $10.5 \pm 0.5$ | $10.0 \pm 0.3$ | $10.3 \pm 0.5$ | 0.50 | 0.60 | 0.95 |
| PN11-21 | $16.4 \pm 0.5$ | $17.9 \pm 1.2$ | $15.6 \pm 0.5$ | $17.4 \pm 1.2$ | 0.11 | 0.51 | 0.89 |
| PN22-28 | $31.9 \pm 0.8$ | $33.4 \pm 2.2$ | $29.4 \pm 0.6$ | $29.6 \pm 1.0$ | 0.54 | 0.03 | 0.66 |
| PN30$^\dagger$ | $73.1 \pm 1.3$ | $74.1 \pm 3.6$ | $67.2 \pm 2.0$ | $62.6 \pm 4.4$ | 0.89 | 0.70 | 0.69 |

Data are presented as mean $\pm$ SEM, with littermates averaged. $n$ = litter number. $P$ values were obtained using a two-way ANOVA, with treatment (Control/EtOH) and sex (male/female) as factors.

EtOH = ethanol; int = interaction; PN = postnatal day; trt = treatment.

$^*$ Birth weight was not recorded at PN0 to minimise disturbance to litters.

$^\dagger$ Only a subset of the litters were weighed prior to cull (see Figure 1 for details); total body weight shown at this age.

**Table 4: Effect of prenatal ethanol exposure on fasting blood glucose, plasma insulin and plasma lipid levels in adolescent offspring.** Offspring were prenatally exposed to ethanol (EtOH) or saline (control) at embryonic day 13.5 and 14.5. All parameters were measured at postnatal day 30.

| Parameters | Male | | Females | | Statistics | | |
| --- | --- | --- | --- | --- | --- | --- | --- |
| | **Control** | **EtOH** | **Control** | **EtOH** | | | |
| | ($n = 6$) | ($n = 8$) | ($n = 3$) | ($n = 3$) | $P_{(trt)}$ | $P_{(sex)}$ | $P_{(int)}$ |
| Fasting blood glucose (mmol/L) | $5.6 \pm 0.2$ | $5.9 \pm 0.2$ | $6.1 \pm 0.2$ | $5.9 \pm 0.1$ | 0.64 | 0.69 | 0.83 |
| Fasting plasma insulin (ng/mL) | $0.55 \pm 0.07$ | $0.65 \pm 0.06$ | $0.64 \pm 0.11$ | $0.63 \pm 0.07$ | 0.58 | 0.67 | 0.53 |
| HOMA-IR | $0.71 \pm 0.11$ | $0.81 \pm 0.07$ | $0.83 \pm 0.15$ | $0.79 \pm 0.07$ | 0.78 | 0.65 | 0.54 |
| QUICKI | $0.31 \pm 0.007$ | $0.30 \pm 0.003$ | $0.30 \pm 0.007$ | $0.30 \pm 0.003$ | 0.55 | 0.50 | 0.50 |
| Plasma HDL (mmol/L) | $1.07 \pm 0.15$ | $0.98 \pm 0.21$ | $0.87 \pm 0.16$ | $0.80 \pm 0.18$ | 0.73 | 0.41 | 0.95 |
| Plasma LDL (mmol/L) | $0.43 \pm 0.06$ | $0.46 \pm 0.04$ | $0.52 \pm 0.07$ | $0.33 \pm 0.01$ | 0.26 | 0.80 | 0.11 |
| Plasma triglycerides (mmol/L) | $0.92 \pm 0.10$ | $1.09 \pm 0.21$ | $0.83 \pm 0.20$ | $0.78 \pm 0.14$ | 0.53 | 0.22 | 0.99 |
| Plasma total cholesterol (mmol/L)[*] | $1.68 \pm 0.18$ | $1.65 \pm 0.23$ | $1.55 \pm 0.20$ | $1.29 \pm 0.16$ | 0.59 | 0.35 | 0.65 |

Data presented as mean ± SEM, with 1 male and 1 female used per litter ($n$ = individual offspring). Sample sizes were lower than the number of dams treated due to availability of animals of each sex for each parameter measured (see Figure 1 for details). $P$ values were obtained using a two-way ANOVA, with treatment (Control/EtOH) and sex (male/female) as factors. $P<0.05$ was considered significant.
EtOH = ethanol; HDL = high density lipoprotein; HOMA-IR = homeostatic model assessment of insulin resistance; int = interaction; LDL = low density lipoprotein; QUICKI = quantitative insulin-sensitivity check index; trt = treatment.
[*] Calculated using HDL + LDL + (triglycerides/5).

**Table 5: Offspring food and water consumption throughout the food preference study.** All animals were housed individually and acclimatised to the study cage for 4 days. Following acclimatisation, baseline measurements of standard chow diet (SD) and water were conducted over days 1-4. A choice of high fat diet (HFD) or SD was then offered for the next 4 days, divided into choice period (CP)1 (days 5-6) and CP2 (days 7-8).

| Parameter | Male | | Female | | Statistics | | |
| --- | --- | --- | --- | --- | --- | --- | --- |
| | Control | EtOH | Control | EtOH | | | |
| | ($n = 8$) | ($n = 9$) | ($n = 8$) | ($n = 9$) | $P_{(trt)}$ | $P_{(sex)}$ | $P_{(int)}$ |
| *$H_2O$ consumption (mL/g BW/day)* | | | | | | | |
| Baseline | $0.064 \pm 0.006$ | $0.060 \pm 0.002$ | $0.070 \pm 0.005$ | $0.082 \pm 0.005$ | 0.34 | 0.007 | 0.12 |
| CP1 | $0.049 \pm 0.004$ | $0.047 \pm 0.002$ | $0.058 \pm 0.011$ | $0.068 \pm 0.004$ | 0.37 | <0.001 | 0.19 |
| CP2[*] | $0.046 \pm 0.003$ | $0.049 \pm 0.003$ | $0.055 \pm 0.003$ | $0.070 \pm 0.004$[†] | | $P = 0.0004$ | |
| *Standard chow diet (SD) consumption (g/g BW/day)* | | | | | | | |
| Baseline | $0.047 \pm 0.001$ | $0.046 \pm 0.001$ | $0.053 \pm 0.002$ | $0.054 \pm 0.001$ | 0.99 | <0.001 | 0.51 |
| CP1 | $0.007 \pm 0.003$ | $0.005 \pm 0.001$ | $0.008 \pm 0.004$ | $0.003 \pm 0.003$ | 0.25 | 0.92 | 0.63 |
| CP2 | $0.019 \pm 0.005$ | $0.017 \pm 0.003$ | $0.010 \pm 0.004$ | $0.013 \pm 0.003$ | 0.87 | 0.11 | 0.44 |
| *High fat diet (HFD) consumption (g/g BW/day)* | | | | | | | |
| CP1 | $0.055 \pm 0.004$ | $0.055 \pm 0.003$ | $0.077 \pm 0.006$ | $0.072 \pm 0.004$ | 0.54 | <0.0001 | 0.64 |
| CP2 | $0.046 \pm 0.004$ | $0.043 \pm 0.004$ | $0.067 \pm 0.004$ | $0.060 \pm 0.004$ | 0.27 | <0.0001 | 0.61 |
| *Energy intake (kJ/day)* | | | | | | | |
| Baseline | $357.8 \pm 8.4$ | $348.3 \pm 10.3$ | $230.4 \pm 10.4$ | $222.1 \pm 6.5$ | 0.33 | <0.0001 | 0.95 |
| CP1 | $628.3 \pm 20.6$ | $605.8 \pm 26.9$ | $502.2 \pm 27.1$ | $430.4 \pm 20.1$ | 0.06 | <0.0001 | 0.31 |
| CP2 | $579.8 \pm 25.6$ | $538.5 \pm 26.6$ | $457.1 \pm 22.5$ | $396.0 \pm 17.6$ | 0.04 | <0.0001 | 0.67 |

Data are presented as mean $\pm$ SEM, with 1 male and 1 female used per litter ($n$ = individual offspring). $P$ values were obtained using a two-way ANOVA, with treatment (Control/EtOH) and sex (male/female) as factors.

CP = choice period; HFD = high-fat diet; BW = body weight.

[*] Indicates data not normally distributed and analysed with a non-parametric Kruskal-Wallis Test across all 4 groups.

[†] Indicates significantly different to corresponding treatment group in males ($P$=0.006) from Dunn's multiple comparison procedure.

**Table 6: Body condition and organ weights of offspring from saline (Control) and ethanol (EtOH) treated litters at 6 months of age.** Offspring were prenatally exposed to ethanol (EtOH) or saline (control) at embryonic day 13.5 and 14.5.

| Parameters | Male | | Females | | Statistics | | |
| --- | --- | --- | --- | --- | --- | --- | --- |
| | Control | EtOH | Control | EtOH | | | |
| | ($n = 12$) | ($n = 12$) | ($n = 12$) | ($n = 12$) | $P_{(trt)}$ | $P_{(sex)}$ | $P_{(int)}$ |
| Body weight (g) | $618 \pm 25$ | $600 \pm 16$ | $345 \pm 13$ | $328 \pm 7$ | 0.29 | <0.0001 | 0.99 |
| Abdominal circumference (cm) | $24.7 \pm 0.7$ | $24.0 \pm 0.4$ | $19.8 \pm 0.5$ | $19.1 \pm 0.3$ | 0.16 | <0.0001 | 0.97 |
| Snout-rump length (cm) | $28.7 \pm 0.3$ | $28.7 \pm 0.2$ | $24.5 \pm 0.2$ | $24.6 \pm 0.2$ | 0.82 | <0.0001 | 0.66 |
| Tibia length (cm) | $6.07 \pm 0.05$ | $6.09 \pm 0.04$ | $5.32 \pm 0.04$ | $5.28 \pm 0.05$ | 0.89 | <0.0001 | 0.52 |
| Ponderal index (g/cm$^3$)$^*$ | $26.0 \pm 0.5$ | $25.6 \pm 0.9$ | $23.5 \pm 0.7$ | $22.0 \pm 0.4$ | 0.15 | <0.0001 | 0.38 |
| Abdominal circumference: snout-rump length (cm/cm) | $0.86 \pm 0.02$ | $0.84 \pm 0.02$ | $0.81 \pm 0.02$ | $0.78 \pm 0.01$ | 0.09 | <0.001 | 0.61 |
| Abdominal circumference: tibia length (cm/cm) | $0.41 \pm 0.01$ | $0.40 \pm 0.01$ | $0.37 \pm 0.01$ | $0.36 \pm 0.01$ | 0.26 | <0.001 | 0.89 |
| Liver weight (g) | $22.24 \pm 1.20$ | $20.68 \pm 0.74$ | $11.47 \pm 0.40$ | $10.41 \pm 0.21$ | 0.08 | <0.0001 | 0.74 |
| Liver:BW ratio (mg/g) | $35.91 \pm 0.80$ | $34.44 \pm 0.60$ | $33.41 \pm 0.94$ | $31.91 \pm 0.81$ | 0.07 | 0.003 | 0.99 |
| Pancreas weight (g) | $1.00 \pm 0.06$ | $0.95 \pm 0.06$ | $0.77 \pm 0.03$ | $0.71 \pm 0.03$ | 0.22 | <0.001 | 0.92 |
| Pancreas:BW ratio (mg/g)$^†$ | $1.63 \pm 0.07$ | $1.60 \pm 0.12$ | $2.24 \pm 0.10^‡$ | $2.17 \pm 0.11^‡$ | $P$ <0.0001 | | |

Data are presented as mean ± SEM. $n$=12 per group, as 1-2 rats per sex per litter (across 8 control litters and 9 EtOH litters; $n$ = individual offspring). $P$ values were obtained using a two-way ANOVA, with treatment (Control/EtOH) and sex (male/female) as factors.
BW = body weight.
$^*$ Ponderal index calculated as body weight (g) divided by snout-rump length cubed (cm$^3$).
$^†$ Indicates data not normally distributed and analysed with a non-parametric Kruskal-Wallis Test across all 4 groups.
$^‡$ Indicates significantly different to corresponding treatment group in males ($P$<0.001) from Dunn's multiple comparison procedure.

**Table 7: Expression of hepatic genes involved in glucose and insulin signalling.** Offspring were prenatally exposed to ethanol (EtOH) or saline (control) at embryonic day 13.5 and 14.5. Liver tissue was collected at 6 months of age.

| Gene of Interest | Male | | Females | | Statistics | | |
|---|---|---|---|---|---|---|---|
| | Control ($n = 8$) | EtOH ($n = 9$) | Control ($n = 8$) | EtOH ($n = 9$) | $P_{(trt)}$ | $P_{(sex)}$ | $P_{(int)}$ |
| *Glucose transport* | | | | | | | |
| *Glut2* | $1.00 \pm 0.05$ | $0.93 \pm 0.07$ | $1.15 \pm 0.08$ | $1.10 \pm 0.12$ | 0.51 | 0.08 | 0.93 |
| *Gluconeogenesis and glycolysis* | | | | | | | |
| *G6pc* | $1.00 \pm 0.17$ | $0.69 \pm 0.16$ | $0.94 \pm 0.14$ | $0.75 \pm 0.14$ | 0.12 | 0.98 | 0.70 |
| *Gck*[*] | $1.00 \pm 0.17$ | $0.75 \pm 0.18$ | $0.70 \pm 0.16$ | $0.53 \pm 0.06$ | | $P = 0.26$ | |
| *Pck1* | $1.00 \pm 0.15$ | $1.20 \pm 0.13$ | $0.80 \pm 0.10$ | $1.08 \pm 0.18$ | 0.10 | 0.26 | 0.80 |
| *Ppargc1α*[*] | $1.00 \pm 0.16$ | $0.96 \pm 0.20$ | $2.01 \pm 0.15$ | $2.40 \pm 0.33$[†] | | $P = 0.001$ | |
| *Insulin signalling* | | | | | | | |
| *Insr* | $1.00 \pm 0.08$ | $0.97 \pm 0.08$ | $1.47 \pm 0.15$ | $1.42 \pm 0.17$ | 0.74 | <0.001 | 0.95 |

Data are presented as mean $\pm$ SEM, with 1 male and 1 female used per litter ($n =$ individual offspring). *P* values were obtained using a two-way ANOVA, with treatment (Control/EtOH) and sex (male/female) as factors.
[*] Indicates data not normally distributed and analysed with a non-parametric Kruskal-Wallis Test across all 4 groups.
[†] Indicates significantly different to corresponding treatment group in males (*P*=0.008) from Dunn's multiple comparison procedure.

**Table 8: Expression of genes involved in glucose transport and insulin signalling in peripheral tissues.** Offspring were prenatally exposed to ethanol (EtOH) or saline (control) at embryonic day 13.5 and 14.5. Gastrocnemius muscle and adipose tissue were collected at 6 months of age.

| | Treatment group | | |
|---|---|---|---|
| | **Control** | **EtOH** | |
| **Gene of Interest** | $(n = 8)$ | $(n = 9)$ | **$P$ value** |
| *Gastrocnemius muscle* | | | |
| Glut4 | $1.00 \pm 0.30$ | $1.11 \pm 0.33$ | 0.81 |
| Insr$^*$ | $1.00 \pm 0.16$ | $0.99 \pm 0.72$ | 0.38 |
| *Adipose tissue* | | | |
| Glut4 | $1.00 \pm 0.11$ | $0.85 \pm 0.12^{\dagger}$ | 0.38 |
| Insr | $1.00 \pm 0.08$ | $0.83 \pm 0.08^{\dagger}$ | 0.16 |
| Foxo1 | $1.00 \pm 0.05$ | $0.96 \pm 0.05^{\dagger}$ | 0.57 |
| Gsk3b | $1.00 \pm 0.04$ | $0.96 \pm 0.05^{\dagger}$ | 0.58 |

Data are presented as mean $\pm$ SEM. $P$ values were obtained using an unpaired Student's t-test.
$^*$ Indicates data not normally distributed and analysed with a non-parametric Mann-Whitney U-test.
$^{\dagger}$ $n = 8$.

**Figure 1**

**Figure 2**

**Figure 3**

**Figure 4**

**Figure 5**