Title: Functional dissociation of theta oscillations in the frontal and visual cortices and their long-range network during sustained attention

Hio-Been Han,[1,2,§] Ka Eun Lee,[2,3,§] Jee Hyun Choi[2,4, *]

[1] Program of Brain and Cognitive Engineering, Korea Advanced Institute of Science and Technology, Daejeon, Republic of Korea

[2] Center for Neuroscience, Korea Institute of Science and Technology, Seoul, Republic of Korea

[3] College of Liberal Studies, Seoul National University, Seoul, Republic of Korea

[4] Department of Neuroscience, University of Science and Technology, Daejeon, Republic of Korea

[§] First authors

[*] Corresponding author

Address correspondence to:

Jee Hyun Choi, Ph.D.

Center for Neuroscience, Korea Institute of Science and Technology

Seoul 02792, Republic of Korea

(Tel) +82-2-958-6952

(E-mail) jeechoi@kist.re.kr

AUTHOR CONTRIBUTIONS

Conceptualization, HBH and JHC; Methodology, HBH; Investigation, HBH and KEL; Formal Analysis, HBH and KEL; Writing – Original Draft, HBH and KEL; Writing – Review & Editing, JHC; Supervision, JHC

ACKNOWLEDGMENTS

This research was supported by the National Research Council of Science and Technology of Korea for the project Development of Solution for Diagnosis, Treatment, and Care System of Dementia (No. CRC-15-04-KIST) and the National Research Foundation of Korea (NRF) grant funded by the Korea government (2017R1A2B3012659).

ABSTRACT

Theta-band (4–12 Hz) activities in the frontal cortex have been thought to be a key mechanism of sustained attention and goal-related behaviors, forming a phase-coherent network with task-related sensory cortices for integrated neuronal ensembles. However, recent visual task studies found that selective attention attenuates stimulus-related theta power in the visual cortex, suggesting a functional dissociation of cortical theta oscillations. To investigate this contradictory behavior of cortical theta, a visual Go/No-Go task was performed with electroencephalogram recording in mice. During the No-Go period, transient theta oscillations were observed in both the frontal and visual cortices, but theta oscillations of the two areas were prominent in different trial epochs. By separating trial epochs based on subjects' short-term performance, we found that frontal theta was prominent in good-performance epochs, while visual theta was prominent in bad-performance epochs, exhibiting a functional dissociation of cortical theta rhythms. Furthermore, the two theta rhythms also showed a heterogeneous pattern of phase-amplitude coupling with fast oscillations, reflecting their distinct architecture in underlying neuronal circuitry. Interestingly, in good-performance epochs, where visual theta was relatively weak, stronger fronto-visual long-range synchrony and shorter posterior-to-anterior temporal delay were found. These findings highlight a previously overlooked aspect of long-range synchrony between distinct oscillatory entities in the cerebral cortex and provide empirical evidence of a functional dissociation of cortical theta rhythms.

KEYWORDS

Theta rhythms, frontal theta, visual theta, electroencephalogram (EEG), Go/No-Go task

IN BRIEF

Previous literature emphasized the pro-cognitive role of coherent oscillatory networks between distal brain regions, such as the fronto-visual theta synchrony. However, such a conceptual framework has been challenged as recent findings revealed distinct behavioral correlates of theta oscillations found in different cortical regions, especially in the frontal and visual cortices. Here, we show that frontal and visual theta represent distinct cortical processes and that the functional connectivity between them increases during sustained attention, especially when one of the two theta rhythms is relatively suppressed. The data presented here highlight a novel aspect of neural long-range synchrony between distinct cortical oscillators with distinct functional significance in task performance.

INTRODUCTION

In the mammalian brain, oscillatory networks in the lower frequency bands (< 12 Hz) such as the theta band (4–12 Hz) are often thought to be a key mechanism of brain-wide interaction between distal regions (1). In particular, increased theta-rhythmic activity of the frontal cortex is a hallmark of improved task performance and sustained attention, known to reflect the level of cognitive demand of the given task (2, 3). In addition, theta oscillations of the frontal cortex have been known to form a phase-coherent network with task-related brain regions, including the sensory cortices (4). The emergence of a theta oscillation network reflects active neuronal processing for cognitive functions such as sustained attention (2), cognitive control (3), and working memory (5). For instance, Liebe *et al.* (6) reported local-field potentials and unit activities that were synchronous across the prefrontal and visual areas in the theta band during visual working memory tasks.

On the other hand, recent findings suggested that theta activities of the visual area are distinct from those of the frontal cortex and that visual theta is also associated with attention but in an opposite manner: stimulus-evoked theta power decreased with increased attention. For instance, Spyropoulos *et al.* (7) recently reported that theta oscillations in the visual cortex elicited by visual stimuli decreased in amplitude when visual attention was directed. In addition, another recent study found arousal-reduced theta power in the high-order visual area (i.e., posterior parietal cortex) of the ferret brain (8) despite its enhanced long-range theta synchrony with the prefrontal cortex during sustained attention reported by an earlier study (9). In fact, a pioneering neuro-feedback study in the 1970s reported similar results of an association between occipital theta suppression and improved performance in a visual detection task (10). There have also been several reports of enhanced theta in the posterior brain region being predictive of inattentiveness such as after sleep deprivation (11) or an attentional lapse (12). Although the term "attention" of each study referred to different aspects of attention (e.g., internal or external attention (13)) on different time scales (e.g., minute or second levels), evidence generally suggests that the employment of attention yields contrary patterns of theta oscillations in the frontal and visual cortices.

This attentional modulation of frontal and visual theta in opposite directions poses a question about the functionality of theta oscillations in different brain regions, especially in terms of their long-range functional connectivity. If cortical theta oscillations of different regions have different roles with opposing behavioral correlates, how could they form a coherent oscillatory network? A possible explanation of this counterintuitive interaction between cortical oscillators is that the oscillations of one region are suppressed to achieve synchronization with those of the other

region. Here, we hypothesize that enhanced fronto-visual theta synchrony can occur during the suppression of frontal or visual theta.

To address this hypothesis, we designed a forced-paced (i.e., the opposite of self-paced) visual Go/No-Go task for head-fixed mice and analyzed the electroencephalogram (EEG) oscillations of each area. With this design, we aimed to capture the dynamic interaction between frontal and visual theta within a single protocol. In particular, in No-Go trials, we expected the recruitment of both frontal and visual theta oscillations, depending on the attentive state of the animal. This is because elaborate sustained attention (i.e., remaining alert while inhibiting a Go response) has been known to elicit control-related theta in the frontal cortex (2, 9), while visual perception without attention has been reported to elicit theta oscillations in the visual area (7). By tracking short-term behavioral performance within a single experimental session, we investigated state-dependent changes of frontal and visual theta oscillations and their coupling with higher frequency oscillations (e.g., gamma oscillations) to confirm their functional dissociation. Then, we characterized the property of functional connectivity between frontal and visual theta to show how distinct cortical oscillators are coupled to form a coherent long-range network.

RESULTS

Changes of behavioral performance during Go/No-Go task

We trained mice (n = 5) to associate rightward motion (S+) with reward (water) and leftward

motion (S-) with punishment (air puff) (Figs. 1A–B). The visual stimulus was ceased at licking,

as exemplified in Fig. 1C. After training (see Materials and Methods for details), we monitored

the performance of the mice during the Go/No-Go test sessions for 10 successive days. During the

test days, overall discrimination performance measured by d-prime (M ± SE, $d' = 0.475 \pm 0.08$)

remained above the chance level (i.e., $d' = 0$, Wilcoxon signed-rank test, $Z = 5.845$, $p < .001$, see

Extended Data Fig. 1-1 for details). There was no statistically significant difference in the

discrimination performance across the test days (Kruskal-Wallis test, $\chi^2 = 1.179$, $p = .999$).

Moreover, a paired comparison between performance of the first half (1–5 d) versus the last half

(6–10 d) did not yield any statistically significant difference (Wilcoxon signed-rank test, $Z =$

0.405, $p = .686$), indicating the absence of a learning effect (Fig. 1-1A).

On the other hand, the animals showed clear changes in performance within a single session,

probably due to the change of motivational factors (e.g., fatigue and quenched thirst) (Fig. 1-1B–

C). The time-dependent patterns of performance were analyzed by dividing the session into early,

middle, and late phases. When we classified the trials by Hit (licking to S+), CR (correct

rejection; no licking to S-), Miss (no licking to S+), and FA (false alarm; licking to S-) (see Fig.

1D for the ratios and numbers of classified trials), the rate of correct answers did not show any

time dependency: sum of Hit and CR ratios was 57.1%, 59.9%, and 57.4% in the early, middle,

and late phases, respectively. Instead, the rates of licking decreased over time, shown through the

sum of Hit and FA ratios which was 64.7%, 51.6%, and 37.1% in the early, middle, and late

phases, respectively, possibly due to the decreased motivation for the task (see Fig. 1-1 for

details).

Prominent theta rhythms in frontal and visual cortices during correct rejection

All analyses of EEG data were focused on the time period from 0.5 s after stimulus onset until

the cessation of stimulus (4 s after stimulus onset) during CR trials (n = 2,411 from five mice). As

expected, our results showed that during the stimulus period of CR trials, theta-band activities

increased in both the frontal and visual cortices (Figs. 1E–F). Elicitation of gamma activities in

the low gamma band (around 40 Hz) was also observed in the visual cortex (Fig. 1E–F). The peak

frequency of theta amplitudes was 7.62 Hz (FWHM = 6.26 Hz) and 6.64 Hz (FWHM = 4.96 Hz)

in the frontal and visual cortices, respectively. The stimulus-locked and response-locked

spectrograms in all trials are summarized in Fig. 1-2. Interestingly, although the behavior was the

118 same (i.e., absence of licking response), Miss trials did not display such prominent theta band

119 activity (Fig. 1-2).

Assessment of attentional state using short-term task performance

Because of the nature of the Go/No-Go task, some trial epochs without active engagement in

the task (e.g., attentional lapse) could be counted as correct No-Go responses. To distinguish

those *by-accident* CR trials from the intended, *elaborate* No-Go responses, we tracked the

changes in attentional state within a single session. During a task, attentional states change and

correspondingly, performance fluctuates over time (2). In human and non-human primate studies,

the short-term history of correctness (6) or reaction time (RT) (14) is often used to track the

change of behavioral state. Similarly, we tracked the short-term proportion correct, p_c, by defining

it as the ratio of correct trials (i.e., Hit and CR) across the 10 previous trials (see Materials and

Methods for details). With a rough range of 2–5 minutes, p_c monitors the momentarily

fluctuating, varying trial-to-trial, attentional state. The distribution of p_c for all CR trials exhibited

a positively shifted Gaussian function (Fig. 2A, $\mu = .66$, $\sigma = .23$, $R^2 = .98$) whose mean was above

the chance level (i.e., $\mu = .5$) (one-sample Z test, $Z = 23.532$, $p < .001$). Using the chance level as

a threshold, we regarded the trial epochs as either attentive (i.e., p_c above the chance level; $p_c >$

0.5) or inattentive states (i.e., $p_c \leq 0.5$). For CR trials, the lowest value of p_c was 0.2 and the

highest value was 1, and attentive states were observed twice as frequently as inattentive states

(1,550 attentive states, 861 inattentive states). When we drew RT as a function of p_c for licking

cases (i.e., Hit and FA), RT decreased in the case of Hit trials (from 1.93 s for $p_c = 0.2$ to 1.14 s

for $p_c = 1$, $Z = 3.047$, $p = .002$, Fig. 2B). On the other hand, RT in the case of FA trials did not

show any dependence on p_c. In this way each trial was labeled with p_c, which successfully

reflected the change of attentional state, for further analysis.

Opposite effects of attentional state on frontal and visual theta amplitudes

The primary aim of the experiment was to articulate the ongoing theta activity signatures with

respect to attention, which were reported to act in an opposite way in the frontal (2) and visual (7)

cortices. Thus, we were interested in the differences in ongoing theta activities between two

attentional states. The scatterplot in Fig. 2C shows a positive correlation between frontal and

visual theta amplitudes. On the other hand, the mean of each p_c presented as a colored error bar

shows a negative correlation, with a movement in their distribution from the lower-right to the

upper-left corner as p_c is increased. This opposite behavior of frontal and visual theta is

manifested in the plot of theta amplitudes as a function of p_c (Fig. 2D). In the frontal cortex, the

correlation between theta amplitude and p_c was positive, reaching its maximum at $p_c = 1$ (Pearson's $r = +.08$, $p < .001$). In contrast, the relationship was negative for visual theta, reaching its minimum at $p_c = 1$ (Pearson's $r = -.10$, $p < .001$). The average theta amplitudes for attentive versus inattentive, analyzed in the time-frequency domain, demonstrates the opposite effect of attention on the frontal and visual theta amplitudes (Fig. 2E). In the frontal cortex, theta amplitude in attentive states was significantly larger compared to that in inattentive states ($96.5 \pm 0.8 \, \mu$V in inattentive, $99.1 \pm 0.7 \, \mu$V in attentive; Wilcoxon rank-sum test, $Z = 3.189$, $p < .001$). On the other hand, in the visual cortex, the amplitude of theta in attentive states was significantly smaller compared to that in inattentive states ($86.7 \pm 1.0 \, \mu$V in inattentive, $82.2 \pm 0.9 \, \mu$V in attentive; Wilcoxon rank-sum test, $Z = -2.582$, $p < .001$). These contrasting correlations suggest that frontal and visual theta activities might have different functional roles during the engagement of cognitive processes required for the No-Go response, raising a question about the neurological signature of sustained attention.

Distinct pattern of phase-amplitude coupling with fast oscillations

To further investigate the role of theta in each region, we performed the phase-amplitude cross-frequency coupling (CFC) analysis, whose characteristics are suggested to reflect the architecture and function of underlying neuronal circuits (15). The amplitude maps of fast oscillations plotted with respect to theta phase revealed strong but different CFC patterns across different p_c values and regions (Fig. 3A). For example, in the high-performance epochs with higher p_c, the amplitude of fast oscillations was strongly modulated by the phase of theta oscillation in the frontal cortex compared to in the low-performance epochs. To analyze the effect of behavioral state on CFC, we calculated the modulation index (MI) (16) as an indicator of CFC strength. As depicted in Fig. 3B, the MI in the frontal cortex showed frequency-specific modulation of behavioral state, while the visual cortex did not show such modulation. Between theta and low gamma (20–40 Hz), the attentional enhancement in the frontal CFC was statistically significant (inattentive: $6.1 \times 10^{-4} \pm 0.1 \times 10^{-4}$, attentive: $6.6 \times 10^{-4} \pm 0.1 \times 10^{-4}$; Wilcoxon rank-sum test, $Z = 2.187$, $p = .029$) (Fig. 3C). For the faster frequency band (high gamma, 80–160 Hz) in the frontal cortex, attentional reduction was observed (inattentive: $7.4 \times 10^{-4} \pm 0.2 \times 10^{-4}$, attentive: $7.0 \times 10^{-4} \pm 0.1 \times 10^{-4}$; Wilcoxon rank-sum test, $Z = -2.546$, $p = .011$). However, in the visual cortex, the attentional effect on CFC was not statistically significant for either low gamma ($Z = -1.443$, $p = .149$) or high gamma ($Z = 1.121$, $p = .262$).

These region-specific behaviors of CFC were further analyzed by investigating the theta phase at which gamma amplitudes peaked (i.e., peak preference). When these phase angles were pooled

separately for each region, their distributions differed (see polar plots in Fig. 3D). In the frontal cortex, the preferred theta phase (i.e., a phase bin of theta where maximum low gamma amplitude was detected) was observed in the middle of the rising phase ($109.7 \pm 2.0°$), whereas in the visual cortex, the preferred theta phase was observed near the start of the falling phase ($191.1 \pm 1.7°$). The distributions of the preferred theta phase of low gamma were statistically different for the frontal cortex and the visual cortex (circular Kruskal-Wallis test, $\chi^2 = 251.140$, $p < .001$). The angle distance of the preferred theta between them was $81.4°$, corresponding to 32.3 ms at 7 Hz. In addition to the results from distinct behavioral modulation (Figs. 2C–E), the result of our CFC analysis presented here provides more reliable evidence that the theta rhythms of each area have distinct functional roles in the cortical microcircuit.

Enhanced fronto-visual functional connectivity during attentive state

Phase synchrony analysis allows the examination of signals that are cooperative between brain areas. To investigate the dynamic interplay of the fronto-visual theta network as attention fluctuates, we first examined the phase relationship between two theta rhythms using a joint phase histogram (see Materials and Methods for details). Fig. 4A highlights the systematic change of the instantaneous phase relationship as a function of p_c. In general, the joint phase of frontal and visual theta showed the stronger colocalization along the diagonal axis regardless of behavioral state, which indicates their intimate phase relationship. Furthermore, the colocalization along the diagonal axis became more robust as p_c increased, suggesting a higher level of phase synchrony between the two theta rhythms. The phase locking value (PLV) (17), an amplitude-independent parameter, elucidates the brain state-dependent synchrony between frontal and visual theta oscillations (Fig. 4B). Compared to the inattentive state, the PLV was significantly higher in the attentive state (PLV = $0.614 \pm .004$ in inattentive, $0.624 \pm .003$ in attentive; Wilcoxon rank-sum test, $Z = -2.313$, $p = .021$). Our result of a fronto-visual theta synchrony associated with task performance is consistent with a previous finding in a non-human primate study (6). While an opposite relationship between frontal and visual theta amplitudes was found in association with the behavioral state, a stronger fronto-visual long-range functional network was established, especially when frontal theta was relatively increased and when visual theta was relatively suppressed.

The asymmetrical joint phase distribution (i.e., bottom-right corner of each joint phase histogram in Fig. 4A) raised the possibility of a driver–responder relationship between frontal and visual theta. Therefore, we investigated the time lag, τ, of frontal theta with respect to visual theta by finding the peaks at the cross-correlation functions (CCFs) of the two theta rhythms. Fig. 3C

illustrates an example CCF of frontal and visual theta rhythms. τ was obtained at each trial and

the distributions of τ were plotted for different p_c values (Fig. 4D). Among all trials, a majority of

the trials (84.61%) had positive τ, indicating that visual theta predominantly precedes frontal

theta. In addition, as p_c increased, the distribution of τ shifted to shorter phase lag. The peak τ in

Fig. 4D systematically decreased as the state changed from inattentive to attentive, with values of

14.83 and 4.33 ms at p_c of 0.2 and 1, respectively. τ was 8.25 ± 0.24 and 6.97 ± 0.17 ms for

inattentive and attentive states, respectively (Wilcoxon rank-sum test, $Z = 4.832$, $p < .001$). In

addition, we found that the peak frequency of the cross-spectral density function was significantly

higher for the attentive state (5.84 ± 0.03 Hz) compared to the inattentive state (5.76 ± 0.04 Hz)

(Wilcoxon rank-sum test, $Z = 2.052$, $p = .040$) (Fig. 2-1). The increase of synchrony and shift to

shorter phase lag during the attentive state suggests the contribution of theta oscillations to task

engagement and sustained attention via their long-range functional network.

DISCUSSION

We found that the frontal and visual theta induced by the task had functionally distinct features. Whereas the frontal theta amplitude was positively correlated with behavioral performance, the visual theta amplitude was negatively correlated. In the frequency domain, the peaks of the two theta oscillations differed by almost 1 Hz, and their phases modulated fast cortical oscillations in different ways. Nonetheless, the frontal-visual theta synchrony increased as a function of performance despite the weakened amplitude of visual theta, implying a critical role of phase-locked oscillations in successful task engagement. It is unlikely that frontal and visual theta have common driving mechanisms; instead, it is more likely that the two theta have independent origins but are weakly coupled to each other. We should note that there is a distinction between phase locking (i.e., coupled oscillation with a constant phase difference) and phase trapping (i.e., un-locked oscillation with the same frequency) in nonlinear oscillator systems (18). Indeed, the frontal and visual theta in our observation showed a phase transition from a stable phase-trapping system to a non-symmetric phase-locking system as the attention level increased. Although the differences in the values were small, we observed that there were abrupt changes in the interaction parameters (e.g., max in joint phase histogram, synchrony, and lag in Fig. 4) near $p_c = 0.8$. Furthermore, the behavioral parameters (i.e., RT in Fig. 2B) and the neural dynamics parameters (i.e., theta amplitude in Fig. 2D, theta–gamma coupling in Figs. 3A and 3C) presented their characteristic features near $p_c = 0.8$. It is highly possible that the theta oscillations changed their dynamic trajectory at the highly task-engaged brain state in their phase portrait; however, determining whether this transition contributed to sustained attention or vice versa was beyond the scope of this study.

Our phase dynamics analysis showed that visual theta led frontal theta throughout the recording sessions. At the same time, frontal, not visual, theta increased its amplitude at higher p_c accompanied by a reduction in its phase delay to visual theta. This confounding behavior at higher p_c could be an influence of top-down directed frontal theta for cognitive control such as motor inhibition (3) or phase resetting of theta oscillations by a stimulus in a highly task-engaged brain state (19). In terms of the *Communication through Coherence* theory (20), the tight interplay of two oscillations renders neuronal communication more effective, precise, and selective. It is possible that a new common source arose around $p_c = 0.8$ to enhance their synchrony and that the newly adjusted phase lag was simply differential conduction delay. However, the different patterns of theta–gamma coupling suggest this was not the case. Recently, Zhang *et al.* reported that human cortical oscillations were regionally distinctive and had the properties of weakly coupled oscillations rather than those of oscillations driven by a common

source or feedback loop (21). Although attentional modulation was not included in their study, it was consistent with our observations, in that the visual rhythm was faster than the frontal rhythm by roughly 1 Hz and propagated to the anterior cortex. However, there was a mismatch in the propagation delay in the human visual to frontal cortex, which was tenfold slower compared to our observation in mice or in monkeys (22). The leading of visual theta might reflect the bottom-up relay of sensory information, and the systematic reduction of the delay is functionally relevant to its purpose of faster relaying of sensory information. The delay is interpreted as a designated temporal shift for the effective relay of information, transmitting a presynaptic spike at the peak depolarization phases of postsynaptic neurons (22, 23). Metaphorically speaking, to selectively improve the traffic in one direction, it may be optimal for the green lights of successive crossways to be slightly delayed considering the amount of time it takes to drive through, rather than operating with zero-lag synchrony.

Then what is the generation mechanism of frontal theta observed here? Considering that our task requires an effort to withhold impulsive actions, frontal theta might originate from the interaction of the cortico-basal ganglia circuit (i.e., anterior-cingulate/medial-frontal and subthalamic nucleus) related to the conflict-related decision process (3, 24, 25). Or the frontal theta may reflect communication between hippocampal structures and the frontal cortex (26, 27), as the task required the subjects to maintain visuospatial information (i.e., direction of motion). Regarding the role of frontal theta oscillations in sustained attention, a large and consistent body of literature has discussed the top-down role of frontal theta in cognitive control (3) and sustained attention (2). All these works included task-engaged mental processes (e.g., keeping one's goal in mind, overriding the prepotent response, monitoring the goal-related sensory cues). Our results provide further evidence for this top-down role of frontal theta oscillations, as we found that frontal theta increased during sustained attention in mice. Importantly, phase synchrony between frontal and visual theta increased in high-performance trials in a critical manner. Thus, frontal theta appears to be behaviorally relevant as its top-down control mode. It should be noted that the substantial work conducted in the frontal cortex has found gamma oscillations to play an important role in pro-cognitive functions (28, 29). On the neuronal spike levels, the spiking activities of the frontal cortex in the high gamma band were locked to theta oscillations in the frontal and posterior parietal cortices during sustained attention (9). In our study, we found effective modulation of low gamma, but not high gamma, by frontal theta using cross-frequency coupling analysis. Continued work with neuronal recording or optogenetic interrogation will be needed to clarify the specific role of frontal theta on neuronal activities during cognitive processing.

The counteracting behavior of visual theta compared to frontal theta was counterintuitive, but it was consistently and significantly observed across animals and days. This phenomenon of "attentional reduction" in visual theta is in agreement with recent findings in monkeys (7, 30) and ferrets (8). In humans, it has been reported that alpha activities (8–12 Hz) in the occipital cortex reflect functional inhibition of task-irrelevant cortical areas (31). To the best of our knowledge, wakeful relaxation related alpha activities have not been reported in mice unlike other cross-species frequency-preserved rhythms (32). In addition, compared to baseline activity, visual theta in our study increased during stimulus presentation (Fig. 2E), whereas alpha usually decreases (33), which is similar to the finding of Spyropoulos *et al.* (7). Therefore, it seems more plausible to regard the oscillatory activities observed in this study as visual theta rather than occipital alpha. However, the functional significance of such attentional reduction in the cortical microcircuit remains unknown.

What is the role of theta rhythmicity in the visual system? Recently, it has been suggested that the theta rhythmicity in the perceptual domain reflects the periodic sampling process of the sensory system (i.e., rhythmic sampling (34, 35)). In this framework, attention periodically fluctuates around the theta-band to sample sensory stimuli, resulting in theta-rhythmic fluctuations of relevant neural activities and behavioral responses (36–38). Still, the suppression of visual theta by attention is not explained, particularly when we consider that theta rhythms reflect the attentional sampling process. According to Spyropoulos *et al.*, the attentional reduction of visual theta was suggested to reflect "more continuous processing" of attended stimuli, and a shift to the faster rhythm was predicted (7). Moreover, Harris *et al.* reported that the phase modulation of visual theta on visual detection performance was more effective for a non-attended than attended target in humans (39). These results suggest that visual theta might be in charge of processing outside the attentional spotlight rather than involved in neural processing in the focal attentional field.

While a growing number of studies have revealed the modulation effects of visual theta on neural excitability, the neural basis for the generation of visual theta remains unknown (see 34). Recently, a predominant feedforward directionality of visual theta was found in monkeys, from lower to higher levels of the visual cortex, and to the temporal cortex (7). In addition, a colocalization of stimulus-induced gamma oscillations and visual theta oscillations was revealed (30). Considering that the feedforward directionality of gamma oscillations in the primary sensory cortex is relatively well known, it is believed that visual theta reflects bottom-up sensory processing rather than top-down influences from higher-order visual areas. Moreover, simultaneous recording of the visual thalamic area and the posterior-parietal cortex in the ferret

brain also demonstrated a strong feedforward directionality in the theta frequency band (8), but it

is not clear whether it is generated from the thalamo-cortical network or instead is locally driven.

If visual theta propagates in a feedforward direction, how can it be modulated by top-down

processes such as attention? One recent study using ferret thalamus recordings suggests an answer

to this question. It found that spiking activities in the pulvinar are modulated by attention, and

these neurons signal in the theta frequency band during sustained attention (40). This area exhibits

bidirectional anatomical connections with higher-order visual cortices (41), and even with the

prefrontal cortex (42). As there has also been a report that unit activities in the primate visual area

V4 are modulated by the external theta driver in the prefrontal cortex (6), it seems highly

plausible that the top-down modulation of theta is generated via a cortico-thalamic input, which in

turn leads to a change of thalamo-cortical connectivity. Indeed, attentional modulation of

neuronal activities in the visual cortices has attracted a wide range of interest among cognitive

scientists. Considering the genetic techniques uniquely applicable to the mouse model, future

studies may take advantage of mouse models to understand the functional architecture and

underlying circuitry of visual theta rhythmicity in association with its attentional modulation.

METHODS

All procedures were approved by the Institutional Animal Care and Use Committees at the <u>XXX (intentionally blinded for double-blind review process)</u> and complied with NIH guidelines. Detailed experimental procedures were as follows.

Animal Preparation. Male C57BL/6 mice were used in the experiments. The animal care and surgery were performed as described in *Extended Data - Materials and Methods*.

Go/No-Go Task. After mice learned to discriminate the direction of the random-dot motion on a screen, their task performance was tested by randomly presenting rightward ("Go") or downward ("No-Go") motion of random dots that stopped at licking or after 4 s when licking was not detected. Each mouse performed 10 experimental sessions on successive days. The design of the stimulus, training, and task are described in *Extended Data - Materials and Methods*.

Rating Task Performance. The ongoing task performance of mice was assessed by the ratio of correct answers (Hit or correct rejection) in 10 successive trials, as described in *Extended Data - Materials and Methods*.

Electroencephalography. EEG responses were recorded at a 2-kHz sampling rate with microscrew electrodes implanted on the skull above the frontal and visual cortices. Ground and reference electrodes were implanted on the occipital bone, as described in *Extended Data - Materials and Methods*.

EEG Analysis. Before analysis, the data were high-pass filtered at 1 Hz and then segmented from 2 s before stimulus onset to 2 s after the cessation of the stimulus. Theta oscillations were investigated by a bandpass filter at 4–12 Hz. For the time-frequency analysis, a fast Fourier transform at a 1-Hz interval with a 512-ms sliding Hanning window with a step size of 100 ms was performed. To eliminate the effect of early evoked potentials, the first 500 ms of each epoch was excluded from the main analyses. Possible cross-frequency coupling between theta and gamma oscillations was studied using the modulation index described by Tort *et al.* (16). The modulation index for phase frequency in the theta band and that for amplitude frequency in the fast frequency band (20–40 Hz for low gamma, 80–160 Hz for high gamma) were calculated. Further information on data analysis can be found in *Extended Data - Materials and Methods*.

Statistical Testing. Non-parametric versions of the t-test (two-sided), one-way ANOVA, and circular centrality test with $\alpha = 0.05$ were performed for statistical tests. For the correlation analysis, Pearson's method was used with $\alpha = 0.05$. Sample size was not previously estimated for the experiments, and blinding or randomization was not needed in this work, as the design was decided by subjects' performance.

REFERENCES

1. von Stein A, Sarnthein J (2000) Different frequencies for different scales of cortical integration: from local gamma to long range alpha/theta synchronization. *Int J Psychophysiol* 38(3):301–313.

2. Clayton MS, Yeung N, Kadosh RC (2015) The roles of cortical oscillations in sustained attention. *Trends Cogn Sci* 19(4):188–195.

3. Cavanagh JF, Frank MJ (2014) Frontal theta as a mechanism for cognitive control. *Trends Cogn Sci* 18(8):414–421.

4. Benchenane K, Tiesinga PH, Battaglia FP (2011) Oscillations in the prefrontal cortex: a gateway to memory and attention. *Curr Opin Neurobiol* 21(3):475–485.

5. Sauseng P, Griesmayr B, Freunberger R, Klimesch W (2010) Control mechanisms in working memory: a possible function of EEG theta oscillations. *Neurosci Biobehav R* 34(7):1015–1022.

6. Liebe S, Hoerzer GM, Logothetis NK, Rainer G (2012) Theta coupling between V4 and prefrontal cortex predicts visual short-term memory performance. *Nat Neurosci* 15(3):456–U150.

7. Spyropoulos G, Bosman CA, Fries P (2018) A theta rhythm in macaque visual cortex and its attentional modulation. *Proc Natl Acad Sci USA* 115(24):E5614–E5623.

8. Stitt I, Zhou ZC, Radtke-Schuller S, Frohlich F (2018) Arousal dependent modulation of thalamo-cortical functional interaction. *Nat Commun* 9(1):2455.

9. Sellers KK*, et al.* (2016) Oscillatory dynamics in the frontoparietal attention network during sustained attention in the ferret. *Cell Rep* 16(11):2864–2874.

10. Beatty J, Greenberg A, Deibler WP, Ohanlon JF (1974) Operant control of occipital theta rhythm affects performance in a radar monitoring task. *Science* 183(4127):871–873.

11. Vyazovskiy VV, Tobler I (2005) Theta activity in the waking EEG is a marker of sleep propensity in the rat. *Brain Res* 1050(1–2):64–71.

12. Mazaheri A, Nieuwenhuis ILC, van Dijk H, Jensen O (2009) Prestimulus alpha and mu activity predicts failure to inhibit motor responses. *Hum Brain Mapp* 30(6):1791–1800.

13. Chun MM, Golomb JD, Turk-Browne NB (2011) A taxonomy of external and internal attention. *Annu Rev Psychol* 62:73–101.

14. Esterman M, Noonan SK, Rosenberg M, Degutis J (2013) In the zone or zoning out? Tracking behavioral and neural fluctuations during sustained attention. *Cereb Cortex* 23(11):2712–2723.

15. Hyafil A, Giraud AL, Fontolan L, Gutkin B (2015) Neural Cross-frequency coupling: connecting architectures, mechanisms, and functions. *Trends Neurosci* 38(11):725–740.

16. Tort AB, Komorowski R, Eichenbaum H, Kopell N (2010) Measuring phase-amplitude coupling between neuronal oscillations of different frequencies. *J Neurophysiol* 104(2):1195–1210.

17. Lachaux JP, Rodriguez E, Martinerie J, Varela FJ (1999) Measuring phase synchrony in brain signals. *Hum Brain Mapp* 8(4):194–208.

18. Aronson DG, Ermentrout GB, Kopell N (1990) Amplitude response of coupled oscillators. *Physica D* 41(3):403–449.

19. Rizzuto DS*, et al.* (2003) Reset of human neocortical oscillations during a working memory

task. *Proc Natl Acad Sci USA* 100(13):7931–7936.

20.   Fries P (2015) Rhythms for cognition: communication through coherence. *Neuron* 88(1):220–
235.

21.   Zhang HH, Watrous AJ, Patel A, Jacobs J (2018) Theta and alpha oscillations are traveling
waves in the human neocortex. *Neuron* 98(6):1269–+.

22.   Gregoriou GG, Gotts SJ, Zhou H, Desimone R (2009) High-frequency, long-range coupling
between prefrontal and visual cortex during attention. *Science* 324(5931):1207–1210.

23.   Nowak LG, Bullier J (1997) The timing of information transfer in the visual system. *Extrastriate*
cortex in primates, (Springer), pp 205–241.

24.   Cavanagh JF*, et al.* (2011) Subthalamic nucleus stimulation reverses mediofrontal influence over
decision threshold. *Nat Neurosci* 14(11):1462–1467.

25.   Cohen MX (2014) A neural microcircuit for cognitive conflict detection and signaling. *Trends*
Neurosci 37(9):480–490.

26.   Siapas AG, Lubenov EV, Wilson MA (2005) Prefrontal phase locking to hippocampal theta
oscillations. *Neuron* 46(1):141–151.

27.   Jones MW, Wilson MA (2005) Phase precession of medial prefrontal cortical activity relative to
the hippocampal theta rhythm. *Hippocampus* 15(7):867–873.

28.   Cho KK*, et al.* (2015) Gamma rhythms link prefrontal interneuron dysfunction with cognitive
inflexibility in Dlx5/6(+/-) mice. *Neuron* 85(6):1332–1343.

29.   Kim H, Ahrlund-Richter S, Wang X, Deisseroth K, Carlen M (2016) Prefrontal parvalbumin
neurons in control of attention. *Cell* 164(1–2):208–218.

30.   van Kerkoerle T*, et al.* (2014) Alpha and gamma oscillations characterize feedback and
feedforward processing in monkey visual cortex. *Proc Natl Acad Sci USA* 111(40):14332–14341.

31.   Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity:
gating by inhibition. *Front Hum Neurosci* 4.

32.   Buzsaki G, Logothetis N, Singer W (2013) Scaling brain size, keeping timing: evolutionary
preservation of brain rhythms. *Neuron* 80(3):751–764.

33.   Fries P, Womelsdorf T, Oostenveld R, Desimone R (2008) The effects of visual stimulation and
selective visual attention on rhythmic neuronal synchronization in macaque area v4. *J Neurosci*
28(18):4823–4835.

34.   VanRullen R (2018) Attention cycles. *Neuron* 99(4):632–634.

35.   VanRullen R (2016) Perceptual cycles. *Trends Cogn Sci* 20(10):723–735.

36.   Busch NA, VanRullen R (2010) Spontaneous EEG oscillations reveal periodic sampling of visual
attention. *Proc Natl Acad Sci USA* 107(37):16048–16053.

37.   Fiebelkorn IC, Pinsk MA, Kastner S (2018) A dynamic interplay within the frontoparietal network
underlies rhythmic spatial attention. *Neuron* 99(4):842–+.

38.   Kienitz R*, et al.* (2018) Theta rhythmic neuronal activity and reaction times arising from cortical
receptive field interactions during distributed attention. *Curr Biol* 28(15):2377–+.

39.   Harris AM, Dux PE, Mattingley JB (2018) Detecting unattended stimuli depends on the phase of
pre-stimulus neural oscillations. *J Neurosci* 3006–3017.

40. Yu C, *et al.* (2018) Theta oscillations organize spiking activity in higher-order visual thalamus

during sustained attention. *eNeuro* 5(1).

41. Jones EG (2001) The thalamic matrix and thalamocortical synchrony. *Trends Neurosci*

24(10):595–601.

42. Romanski LM, Giguere M, Bates JF, Goldman-Rakic PS (1997) Topographic organization of

medial pulvinar connections with the prefrontal cortex in the rhesus monkey. *J Comp Neurol*

379(3):313–332.

 FIGURE LEGENDS

Figure 1. Theta oscillations during Go/No-Go task in frontal and visual cortices

(**A**) Schematic illustration of experimental setup. Mice performed Go/No-Go task in custom-built head
restrainer. (**B**) Experimental procedures. Task required subject to discriminate direction of random-dot
motion (rightward: S+, downward: S-). "Go" response (i.e., licking) to S+ and S- motion resulted in
reward and punishment, respectively. There was no particular reinforcing stimulus following "No-Go"
response (i.e., no licking). (**C**) Example EEG data of two trials. (Upper) For S+ motion, licking response
was counted as "Hit," and water reward was followed instantaneously. Motion stimulus was ceased at first
lick (RT). (Lower) For S- motion, No-Go response was counted as "Correct rejection (CR)." Scale bars on
bottom right indicate 500 ms and 300 µV. (**D**) Summary of behavioral results. From all subjects (five
mice), and all sessions (10 days), data of 8,390 trials were collected. In this study, only the trials from CR
(n = 2,411) were analyzed. (**E** and **F**) (**E**) Time-frequency representation of grand-averaged event-related
spectral perturbation (ERSP) of each cortical area. In both the frontal and visual cortices, prominent theta
(4–12 Hz) oscillations were observed during 4 s of stimulus presentation (No-Go period). (**F**) Temporal
average (0.5 to 4 s) of theta oscillations in each cortical area. Numbers denote peak frequency of each
channel. FA: false alarm, Fr: frontal cortex, Vis: visual cortex, Amptd: amplitude, Freq: frequency.

Figure 2. Task performance and theta amplitude

(**A**) Distribution of p_c (proportion correct) in all CR trials (n = 2,411). p_c was calculated across 10 previous trials. With a threshold of chance level ($p_c = 0.5$), each trial was classified as either attentive ($p_c > 0.5$) or inattentive ($p_c \leq 0.5$) state. (**B**) Change of mean RT as function of p_c. The RT of Hit trials decreased as p_c increased (Pearson's $r = -.051$, $p < .05$), suggesting the change of attentional level within the Hit trial epochs. (**C**) Scatterplot of frontal/visual theta amplitude with color-coding of p_c. Each gray dot represents the observed theta amplitude value of a single CR trial. Gray solid lines indicate the probability density function of each channel. Along with the changes of p_c, the mean amplitudes of frontal and visual theta showed a negative relationship. (**D**) Mean amplitude of frontal/visual theta as function of p_c. The linear correlation between theta amplitude and p_c was positive in the frontal ($r = .076$, $p < .001$) and negative in the visual ($r = -.099$, $p < .001$), suggesting their distinct behavioral correlates. (**E**) Time-frequency representation of oscillatory amplitudes highlighting state- and region-dependent change of theta amplitudes. Black dashed lines indicate the start/end of the visual stimuli. Error bars represent ±1 *SD* (**C**) and *SE* (**B**, **D**) of the means. p_c: probability correct, RT: reaction time.

Figure 3. Task performance and theta-gamma phase-amplitude coupling

(**A**) Color-coded fast-frequency amplitudes as function of theta phase in frontal (top) and visual (bottom) and as function of p_c (left to right). (**B, C**) MI for local theta-gamma coupling as function of p_c (left: frontal, right: visual). In the frontal, the effect of attention on CFC was frequency-specific (increased CFC with low gamma, decreased CFC with high gamma). The attentional modulation of the visual coupling was not significant. (**D**) Grand-averaged amplitudes of fast oscillations as function of theta phase, showing noticeable regional difference of preferred theta phase (left: frontal, right: visual). (**E**) Trial histogram of preferred theta phase of 20–40 Hz oscillations (left). Hexagons and dotted lines denote the peak and mean of each polar histogram, respectively. Mean preferred angle of each area (right). The distances between preferred angles over sites were both significant. Error bars represent ±1 *SEM*. * $p < .05$, *** $p < .001$. CFC: cross-frequency coupling. INA: inattentive, ATT: attentive. MI: modulation index.

Figure 4. Synchrony and lags of theta oscillations

(**A**) Two-dimensional representation of joint phase histogram of frontal and visual theta oscillations as function of p_c. The white color of each histogram indicates the higher probability of colocalization of the phase of two oscillations, suggesting a higher level of synchrony between two oscillations. The in-set text indicates the maximum value of colocalization probability in each p_c bin. (**B**) Mean fronto-visual phase synchrony calculated by PLV as function of p_c. The phase synchrony was higher for the attentive state than the inattentive state. (**C**) Cross-correlation analysis of example EEG epoch. (Top) To calculate temporal lag (τ), raw time traces of EEG data from two channels were band-pass filtered in the theta band. (Bottom) After filtering, the cross-correlation function was obtained and τ was defined as a temporal lag where the correlation coefficient reaches its maximum. Positive and negative values of τ indicate a lead of the visual and frontal theta, respectively. (**D**) Trial histograms of τ between frontal and visual theta. The dots denote the peak of each histogram. Color-coded texts indicate the peak position from the lowest p_c (green) and to the highest p_c (red) bin. (**E**) Mean τ as function of p_c. The τ was smaller in the attentive state than in the inattentive state. Error bars represent $\pm$ 1 *SEM*. * $p < .05$, *** $p < .001$. INA: inattentive, ATT: attentive, PLV: phase locking value.

Extended Data

This PDF file includes:

- Extended Data - Materials and Methods

- Extended Data - Figures 1-1, Figure 1-2, and Figure 2-1

Extended Data - Materials and Methods

Animals and surgery

Five male C57BL/6 mice (4–8 weeks old at time of surgery, body weight $M = 26.1$ g, $SD = 4.7$ g) were used, maintained under a 12:12 h light:dark cycle (lights on at 8:00 a.m.) in a temperature- and humidity-controlled environment. Food was available *ad libitum*, and water was mildly restricted (1 mL per day, additional water available as reward from task) from two weeks after surgery. Health status of the animals was assessed daily according to a systematic assessment protocol (1). All procedures were conducted in compliance with the Institutional Animal Care and Use Committee of XXX (intentionally blinded for double-blind review process), conforming to the NIH Guide for Care and Use of Laboratory Animals (NIH Publication No. 86-23, revised 1985).

For stereotaxic surgery, mice were anesthetized with a ketamine and xylazine cocktail (120 and 6 mg/kg, respectively) by intraperitoneal injection and positioned on a stereotaxic apparatus (Model 957, Kopf Instruments, CA, USA). After shaving the head, an incision was made to expose the skull and implant sterilized screw (Asia Bolt, South Korea) electrodes above the frontal (anterior-posterior, 1.9-2.1 mm; medial-lateral, ±2.1 mm from bregma) and visual cortices (anterior-posterior, -3.7 mm; medial-lateral, ±3.7 mm from bregma). For selection of recording sites, we referred to a previous study which targeted the mouse prelimbic area to record theta oscillations using surface electrodes *in vivo* (2), and our pilot study using high-density surface EEG array (3) which showed that the selected mouse visual area exhibited most prominent motion-specific oscillatory responses to random-dot motion stimuli. Reference and ground electrodes were implanted on the occipital bone (4). Dental cement (Vertex Self-Curing, Vertex Dental, Netherlands) was applied over the electrodes for fixation and two polycarbonate nuts (inner diameter 3 mm, Nippon Chemi-Con, Japan) were attached to the caudal edges of the cement, for use as anchors of the custom-built head-fixation apparatus. After surgery, mice were treated with antibiotics and analgesics.

Stimuli and apparatus

All experiments were conducted in a lightproof Faraday cage. Mice were restrained in a custom-built acrylic tube (3.5 cm in diameter) and head fixed by the polycarbonate nuts. For a reward, a water drop (3 μL) was released through a pressure equalizer tube (2 μm in diameter) with a programmable syringe pump (Fusion 200, Chemyx, TX, USA). Licking behavior was counted with a custom-built lickometer made of stainless steel wire (3 μm in diameter) and a data

acquisition system (cDAQ-9147, National Instruments, TX, USA). For punishment, a single puff
of air was delivered to the animal's cheek (air pressure = 15–20 p.s.i., duration = 0.2 s), supplied
with a compressed air tank and controlled by a solenoid valve. A schematic diagram of the
experimental setup is shown in Figure 1a.

For visual stimuli, a random-dot motion was generated and presented through Matlab
(Mathworks, Natick, MA, USA). The dot motion was composed of 100 small dots (full
coherence, size = 0.5 ° in radius, field size = 60 ° in diameter, speed = 42 °/s, dot life = 1.67 ±
0.33 s from a Gaussian distribution, orientation = 90 ° or 180 °) and presented on a LCD screen
(LG E1910PM-SN, 19 inches, 1280x1024 resolution), located 15 cm away from the animal's
eyes. As for the validity of using random-dot motions in the mouse model, previous studies have
shown rodents are able to discriminate the direction of the random-dot motion (5) and exhibit
brain oscillatory responses similar to those found in humans (6).

Training procedure

The behavioral training procedure was consisted of three stages: (1) habituation (7 d), (2)
response shaping (2–3 d), and (3) conditioning (2–4 d). The habituation stage includes animal
handling (10–15 min/d) and provides animals the opportunity to adapt to the head-fixation setup
inside the experimental apparatus (30 min/d) without behavioral requirements. The response
shaping stage was aimed to teach animals that water can be accessed by licking the tube. During
this period, a water reward was given for every lick (lick port sensor time-out = 20 s). Once the
animal started to robustly lick the tube, they advanced to the conditioning stage. The conditioning
stage trained animals to associate a visual stimulus with a water reward. Each trial began with a
monotone sound cue (10 kHz, 70 dB, 100 ms) and a white central fixation cross (size = 3 °). After
2–3 s of the cue, a rightward moving-dots stimulus (S+, 'Go'-motion) was presented on the
screen, which subsequently turned black for 30 s for the water-tube licking. Each stimulus did not
last longer than 2 min, and 100 trials were given in one day. Our pilot study of reaction time (RT),
validated through video recording, showed that intentional licking results in a typical RT
distribution with a peak near 1 s, whereas spontaneous licking results in a scattered RT
distribution over a 2 min time period. Also, we found that licking rates decreased over time,
potentially due to thirst satiation. Thus, we considered the conditioning was successfully
established when the RT of the first 80 trials reached a typical RT distribution. Overall, the
percentage of mice licking in response to the visual stimulus on the last day of the conditioning
stage was roughly 70%.

Go/No-Go task

In the Go/No-Go sessions (10 d, consecutive), the animals were required to discriminate the direction of the moving dots (Fig. 1B). Each trial began with a monotone sound cue (10 kHz, 70 dB, 100 ms) and a white central fixation cross (size = 3 °). After 2–3 s of the cue, the stimulus was presented for 4 s with either a rightward- (S+, 'Go'-motion) or downward- (S-, 'No-Go'- motion) motion which was terminated when licking was detected. The direction of the motion of each trial was randomly determined (50%) with a restriction that the same stimulus was not presented more than 4 times in a row. A licking response during the S+, which was counted as a Hit, resulted in instant reward. If the mouse did not take the reward within 20 s, the water in the tube was suctioned out. On the other hand, a licking response to the S- was counted as a FA, resulting in instant punishment and a 20 s timeout. Trials without licking responses were counted as a Miss or CR for the S+ or S-, respectively. To guide the animals to react more consciously, rather than reflexively, the response window started 0.5 s after the onset of motion (Fig. 1B). Inter-trial intervals (ITI, black screen) were 5–8 s long. Any spontaneous licking during the ITI or cue period before the response window reset the trial starting from the ITI.

The number of trials per session was determined by the animal's performance that day (287 trials/d on average). Each session was terminated once the animal stopped licking for 30 successive trials, which we regarded as a lack of motivation for the task, and were excluded from the EEG analysis. We unexpectedly observed an obsessive licking behavior, during which the animal would indiscriminately lick the tube regardless of stimulus type. If animals licked for more than 30 trials in a row, those trials were excluded from EEG analysis. Overall, around 41% of trials were excluded.

Behavioral state rating

As previous human and animal studies have shown, performance in cognitive tasks varies over time depending on the behavioral state, for instance due to an attentional lapse (7, 8). To investigate the relationship between cortical theta activities and sustained attention, it was important to infer correctly the behavioral state, as either attentive or inattentive. This is particularly important in Go/No-Go tasks because mind wandering for a short period of time (i.e., attentional lapse) can be counted as a correct No-Go response. To rate each trial according to attentional state, the consistency of performance over time was analyzed. Because the span of attentional state in humans was reported to range between 2–7 min (8) or exceed 10 min (9), we assumed that short-term attentional state in mouse would be similar or shorter than that of human. Then, we assumed the level of attentional state is reflected through performance in the previous

659 trials (in the order of a few minutes). Finally, we identified fluctuations in performance over 10

660 successive trials by calculating the probability of correct answers (p_c; Hit and CR), ranging from 0

661 (0 correct trials) to 1 (10 correct trials) with the chance level 0.5. p_c of k-th trial was defined as

662 follows,

$$p_c(k) = \frac{1}{10} \sum_{n=k-9}^{k} C(n),$$ Eq. (1)

where the binary function, $C(n)$, indicates the outcome of the trial as correct (1) or not (0). In our

paradigm, 10 trials lasted approximately 2–5 min. We subsequently classified the trials into

attentive state (p_c larger than the chance level) or inattentive state (p_c equal to or smaller than the

chance level).

Statistical analysis

Non-parametric versions of t-test (Wilcoxon signed-rank and rank-sum test), ANOVA

(Analysis of variance; Kruskal-Wallis test), and circular-median test using CircStat Toolbox (10)

were performed ($\alpha = .05$, two-tailed). For correlation analysis, Pearson's method was used with

the same alpha level.

EEG data acquisition and processing

EEG data were collected through Grass 8-17C amplifiers (Grass Technologies, RI, USA) with

60 Hz notch filters (gain = 50,000) and were digitized using a data acquisition system (cDAQ-

9147). After band-pass filtering (1–200 Hz), the EEG data were processed on a trial-by-trial basis.

To eliminate hemisphere-specific activities, EEG signals from bilateral electrodes were averaged

in the time domain. To minimize the effect of early evoked activities (e.g., event-related

potentials), only the EEG activities from 0.5 to 4 s after motion onset were used for statistical

testing of oscillation amplitudes. We computed the mean and two-tailed 99% confidence intervals

of the distribution of the overall amplitude spectrum, and the values lying outside of these

confidence intervals were excluded. A time-frequency amplitude spectrogram was obtained by

applying fast Fourier transform with a sliding Hanning window (bin size = 512 ms, step size =

100 ms) and by taking absolute real parts from the epochs, which resulted in a 1 Hz frequency

resolution. For the time-frequency spectrogram visualization, the amplitudes of the pre-stimulus

periods (2 to 0.5 s before stimulus onset) were averaged and subtracted in a frequency-wise

manner.

Cross-frequency coupling between theta and gamma oscillations

To investigate cross-frequency coupling between theta and gamma oscillations, we calculated

the modulation index (MI) as described in Tort et al. (11). In brief: (i) raw EEG epochs were band

passed (phase frequency: 4–12 Hz, amplitude frequencies: 6 to 200 Hz with 2.5 Hz-step sliding

window and 5 Hz width) with 5^{th} order Butterworth filters. (ii) The time series of the phase of the

theta oscillations denoted as $\phi_{f_\theta}(t)$, and the time series of the amplitude of the gamma

oscillations denoted as $A_{f_\gamma}(t)$, were extracted via Hilbert transform. (iii) The phase of the theta

oscillations was divided into N bins from -π to π. In each bin, the average of $A_{f_\gamma}(t)$ was

calculated and then normalized by dividing the sum over the N bins and was denoted as P_j for the

j-th bin. Here, P_j equals the probability density function of the gamma amplitude distribution over

the theta phase ranging from 0 to 1. (iv) An entropy index, H, was calculated by the definition of

Shannon entropy, given by

$$H = -\sum_j^N P_j \log P_j. \qquad\qquad \text{Eq. (2)}$$

702   If the gamma oscillations are independent from theta phase, P_j is $1/N$; therefore $H = \log(N)$. (v)

703   The statistical deviance of the measured distribution (P) from the uniform distribution was

704   calculated by adopting Kullback-Leibler distance by the following formula,

$$D_{KL} = \log(N) - H \qquad\qquad \text{Eq. (3)}$$

705   where D_{KL} stands for the Kullback-Leibler distance. (vi) Finally, MI was defined with a Z-score

706   of observed entropy as the following formula indicates,

$$MI = \frac{D_{KL}}{\log(N)} \qquad\qquad \text{Eq. (4)}$$

In the current study, we set N equal to 30 (i.e., bin size = 12°).

Phase synchrony analysis of theta activities

To quantify the interaction between theta oscillations in the frontal and visual areas, phase

synchrony was estimated by joint phase histograms and by calculating phase locking values

(PLV) (12). To obtain joint phase histograms, a Matlab built-in function for two-dimensional

histogram (`hist2.m`) was used. To calculate PLV, first, instantaneous phase of the theta

oscillations was calculated via Hilbert transform and was denoted as $\phi_{Fr}(t)$ and $\phi_{Vis}(t)$ for the

frontal and visual areas, respectively. Second, PLV was calculated in each stimulation period

excluding the first 0.5 s (i.e., from $t = 0.5$ to $t = 4$) as follows,

$$PLV = \frac{1}{T}\left|\sum_{t=0}^{T-1} \exp(i\Delta\phi(t))\right| \qquad\qquad \text{Eq. (5)}$$

717   where $\Delta\phi(t) = \phi_{\mathrm{Fr}}(t) - \phi_{\mathrm{Vis}}(t)$. PLV ranges from 0 (i.e., absence of phase locking) and 1

718   (i.e., perfect phase locking).

719   A cross-correlation function $\rho_{X,Y}(\Delta t)$ of visual and frontal theta oscillations, $X(t)$ and $Y(t)$,

720   respectively, was calculated in each trial excluding the first 0.5 s (i.e., from $t = 0.5$ to $t = 4$),

721   according to

$$\rho_{X,Y}(\Delta t) = \frac{1}{\sigma_X \sigma_Y} \cdot \frac{1}{T} \sum_{t=0}^{T-1} (X(t) - \mu_X)(Y(t + \Delta t) - \mu_Y) \qquad \text{Eq. (6)}$$

722   in which μ and σ indicate the mean and standard deviation. The time delay, τ, was obtained from;

$$\tau = \underset{\Delta t \in [-1/f_\theta, 1/f_\theta]}{\arg\max} \rho_{X,Y}(\Delta t) \qquad \text{Eq. (7)}$$

723   where arg max indicates the arguments of the maxima (i.e., the points that the cross-correlation

724   functions are maximized). Positive values of τ indicate the temporal lead of visual theta over

725   frontal theta and vice versa. The resolution of τ was 0.5 ms (2,000 Hz sampling).

 Extended Data - Figures

Fig. 1-1. Performances changing across- and within-session.

(**A**) Performance varies across sessions. Across the 10 days of sessions, overall behavioral performance

(d', $M = 0.475$, $SE = 0.08$) was significantly larger than chance level, $Z = 5.845$, $p < .001$. Learning effect,

calculated by the difference of mean d' between first five days ($M = 0.399$, $SE = 0.07$) and last five days

($M = 0.551$, $SE = 0.21$), was not significant, $Z = 0.405$, $p = .686$. Error bars represent ± 1 SEM. (**B**)

Performance varies within a session. Splitting trials of single session into three different groups, initial

(first 1/3), middle (second 1/3), and final (last 1/3), change of behavioral performance within session was

analyzed. In general, proportion of Go response trials (*i.e.*, Hit and FA) was higher at initial phase, and

gradually decreased, suggesting the change of animals' motivational level to water reward. Interestingly,

despite the decrease of probability Go response, p(Go), overall probability of correct (*i.e.*, Hit and CR),

p(Correct), showed relatively small variation within session. (**C**) Behavioral state changes across- and

within-session. Changes of proportion of 'attentive'-labeled trials ($p_c > 0.5$) as a function of experimental

session (left) and as a function of within-session progress (right) were analyzed. To calculate the effect

size, one-way ANOVA (Kruskal-Wallis test) was performed. We found experimental session explained

17% of the variance, while the within-session progress explained 35% of the variance, suggesting the

larger effect size of motivation-related factors relative to the learning-related factors. Error bars represent ±

1 *SEM*. (**D**) Duration and transitions of behavioral state. Behavioral state showed transitions (i.e., transition

from inattentive to attentive, and vice versa) and stationary phase with certain duration (inattentive: 6.60

trials in average, attentive: 7.81 trials in average) (left). Using Markov-chain model with two-state process,

this state transition was summarized in a state diagram (right).

Fig. 1-2. Grand-averaged EEG responses during Go/No-Go task.

(**A**) Stimulus-locked grand-averaged amplitude spectrogram of four behavioral outcomes. Black dashed

lines indicate the onset of visual stimulus. Hit and FA trials exhibited relatively large changes in

oscillatory amplitudes, compared to those of Miss and CR trials. These large activities were considered as

reinforcer-related activities (i.e., water reward, air puff punishment, motor movement, etc.) as they were

locked to the moment of reaction (See panel c).

(**B**) Averaged (t = 0.5 – 4 s after motion onset) amplitude spectrum of spectrograms in (a).

(**C**) Same as (a), but response-locked. For the trials which do not contain behavioral response (CR, Miss),

mean RTs of the session from Hit and FA trials were used to align the epoch.

(**D**) Same as (b), but response-locked.

Fig. 2-1. Baseline-uncorrected amplitude- and cross-spectral densities.

(**A**) Mean amplitude- or cross-spectrum of CR trials (n = 2,411, t = 0.5–4 s) calculated from the frontal.

Error bars represent ± 1 *SEM*.

(**B**) Same as (**A**), from the visual.

(**C**) Same as (**A**), from cross-spectrum of the frontal and visual.

(**D**) Mean frontal theta (4–12 Hz) amplitude. Error bars represent ± 1 *SEM*.

(**E**) Same as (**D**), from the visual.

(**F**) Same as (**D**), from cross-spectrum of the frontal and visual.

(**G**) Mean peak frequency of the frontal theta amplitude spectrum. Peak values were calculated in trial-by-

trial basis. Colored numbers illustrate the duration of one cycle of each frequency. Error bars represent ± 1

SEM.

(**H**) Same as (**G**), from the visual.

(**I**) Same as (**G**), from cross-spectrum of the frontal and visual.
777
778

References

1. Guo ZCV*, et al.* (2014) Procedures for Behavioral Experiments in Head-Fixed Mice. *Plos One* 9(2).
2. Sauer JF, Struber M, & Bartos M (2015) Impaired fast-spiking interneuron function in a genetic mouse model of depression. *eLife* 4.
3. Choi JH, Koch KP, Poppendieck W, Lee M, & Shin HS (2010) High resolution electroencephalography in freely moving mice. *J Neurophysiol* 104(3):1825-1834.
4. Paxinos G & Franklin KB (2004) *The mouse brain in stereotaxic coordinates* (Gulf professional publishing).
5. Douglas RM, Neve A, Quittenbaum JP, Alam NM, & Prusky GT (2006) Perception of visual motion coherence by rats and mice. *Vision Res* 46(18):2842-2847.
6. Han HB, Hwang E, Lee S, Kim MS, & Choi JH (2017) Gamma-Band Activities in Mouse Frontal and Visual Cortex Induced by Coherent Dot Motion. *Sci Rep-Uk* 7.
7. Liebe S, Hoerzer GM, Logothetis NK, & Rainer G (2012) Theta coupling between V4 and prefrontal cortex predicts visual short-term memory performance. *Nat Neurosci* 15(3):456-U150.
8. Esterman M, Noonan SK, Rosenberg M, & Degutis J (2013) In the zone or zoning out? Tracking behavioral and neural fluctuations during sustained attention. *Cerebral cortex* 23(11):2712-2723.
9. Clayton MS, Yeung N, & Kadosh RC (2015) The roles of cortical oscillations in sustained attention. *Trends Cogn Sci* 19(4):188-195.
10. Berens P (2009) CircStat: a MATLAB toolbox for circular statistics. *J Stat Softw* 31(10):1-21.
11. Tort AB, Komorowski R, Eichenbaum H, & Kopell N (2010) Measuring phase-amplitude coupling between neuronal oscillations of different frequencies. *J Neurophysiol* 104(2):1195-1210.
12. Lachaux JP, Rodriguez E, Martinerie J, & Varela FJ (1999) Measuring phase synchrony in brain signals. *Hum Brain Mapp* 8(4):194-208.