Homo- and heterodimerization of bHLH transcription factors balance stemness and bipotential differentiation

Aleix Puig-Barbé[1] and Joaquín de Navascués[1,2]

[1] European Cancer Stem Cell Research Institute, School of Biosciences, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff CF24 4HQ, UK.

[2] School of Life Sciences, University of Essex, Wivenhoe Park, Colchester CO4 3SQ, UK.

Correspondence: j.denavascues@essex.ac.uk

Running title: Da, Emc and Sc balance stemness and differentiation in the fly intestine

Key words: bHLH / stemness / differentiation / bipotent stem cell / intestinal stem cell

ABSTRACT

Multipotent adult stem cells must balance self-renewal with differentiation into various mature cell types. How this activity is molecularly regulated is poorly understood. By using genetic and molecular analyses in vivo, we show that a small network of basic Helix-Loop-Helix (bHLH) transcription factors controls both stemness and bi-potential differentiation in the *Drosophila* adult intestine. We find that homodimers of Daughterless (Da, homolog to mammalian E proteins) maintain the self-renewal of intestinal stem cells and antagonise the activity of heterodimers of Da and Scute (Sc, homolog to ASCL and known to promote intestinal secretory differentiation). We find a novel role for the HLH factor Extramacrochaetae (Emc, homolog to Id proteins), titrating Da and Sc to promote absorptive differentiation. We further show that Emc prevents committed absorptive progenitors from de-differentiating, revealing the plasticity of these cells. This mechanism of interaction partner-switching enables the active maintenance of stemness, but primes stem cells for differentiation along two alternative fates. Such regulatory logic could be recapitulated in other bipotent stem cell systems.

 INTRODUCTION

The regulation of stem cell fate decisions hinges on the control of transcription. Central to this control are gene regulatory networks, whose activity and dynamics steer cells along particular differentiation pathways (Graf & Enver, 2009; Levine & Davidson, 2005; Moris *et al*, 2016). These pathways can often be construed as a succession of binary steps that are regulated by the cross antagonism of transcription factor pairs (Graf & Enver, 2009; Simon *et al*, 2018). However, for multipotent stem cells, several fate options may be simultaneously available through higher-dimensional switches (Cinquin & Demongeot, 2002; 2005). To understand the choice between several alternative fates, knowledge of the interactions between transcriptional regulators is essential.

Intestinal stem cells (ISCs) are a clear example of multipotency in adult tissues. In *Drosophila* and mammals, ISCs face a triple choice between self-renewal and differentiation into one of two mature lineages: secretory or absorptive (reviewed in Jiang & Edgar, 2012; Crosnier *et al*, 2006; Philpott & Winton, 2014). The secretory lineage comprises several cell types in vertebrates, but only enteroendocrine (EE) cells in *Drosophila*. Absorptive cells are enterocytes (ECs). In *Drosophila*, ISCs produce lineage-specific precursors through distinct molecular triggers. High Notch signalling induces formation of enteroblasts (EBs), which will give rise to ECs. Expression of the bHLH (basic Helix-Loop-Helix) transcription factors Scute (Sc) or Asense (Ase), members of the *achaete-scute Complex* (*AS-C*, homologs of *ASCL* mammalian genes) induces the formation of EE precursor cells (pre-EEs) and their transition to mature EE cells (Micchelli & Perrimon, 2006; Ohlstein & Spradling, 2006; Bardin *et al*, 2010; Zeng & Hou, 2015; Chen *et al*, 2018). This argues for ISCs facing a multiple choice, rather than a sequence of binary decisions. However, very little is known about the molecular mechanisms that can simultaneously regulate both self-renewal and bi-potential differentiation.

Transcription factors of the bHLH family control fate in multiple developmental contexts, from sex determination and mesoderm specification, to neurogenesis and immune cell type specification (reviewed in Baylies *et al*, 1997; Murre *et al*, 1994; Murre, 2005; Bertrand *et al*, 2002; Ohtsuka & Kageyama, 2010). Their HLH motif mediates dimerization, while the preceding region, rich in basic amino acids, allows DNA binding (Massari & Murre, 2000). bHLH factors are subdivided into several

classes, with class I (E proteins) comprising factors such as E2A, E2.2 and HEB (encoded by *TCF3/4/12*; a single gene, *daughterless* (*da*), in *Drosophila*), usually with broad expression patterns; and class II containing more restrictedly expressed factors such as MYOD, TWIST and ASCL (MyoD, Twi, and AS-C proteins in *Drosophila*). Generally, class II factors heterodimerise with a class I member to bind DNA, while class I factors can either homo- or heterodimerise (Murre *et al*, 1994). The class V comprises HLH factors lacking the basic domain, known as *Inhibitors of DNA binding* (Id proteins; a single gene, *extra macrochaetae* (*emc*), in *Drosophila*), as heterodimerisation with them renders dimers unable to bind DNA (Murre *et al*, 1994). Combinatorial, antagonistic or cooperative interactions confer unique properties to regulatory networks based on these factors (Amoutzias *et al*, 2008), potentially including multi-stable dynamical behaviour (Cinquin & Page, 2007; Cinquin & Demongeot, 2005).

Class II bHLH factors are known to regulate differentiation in the metazoan intestine (reviewed in Philpott & Winton, 2014; Hartenstein *et al*, 2017). In *Drosophila*, Sc and Ase can initiate EE differentiation (Bardin *et al*, 2010; Amcheslavsky *et al*, 2014; Chen *et al*, 2018; Guo & Ohlstein, 2015), while other bHLH genes maintain EE function (*dimmed*, homolog of *NeuroD*; Beebe *et al*, 2015), or promote their functional diversity (*tap*, homolog of *Neurogenins*; Hartenstein *et al*, 2017). On the other hand, Da is required for ISC maintenance, as ISCs mutant for *da* differentiate (Bardin *et al*, 2010; Lan *et al*, 2018). However, which interaction partner binds to Da to maintain stemness is not known, and how different bHLH factors dimerise to allow differentiation has not been explored. Here we identify the Da homodimer as the critical bHLH complex maintaining ISC self-renewal, elucidate the role of the HLH factor Emc in titrating Da and Sc to promote absorptive differentiation, and find a functional antagonism between Da:Da and Da:Sc dimers. Our results suggest that a network of bHLH factors acts as a three-way switch to regulate self-renewal and bipotential differentiation in the adult fly gut.

RESULTS

emc is expressed preferentially in the absorptive lineage

bHLH factors are essential for correct differentiation during adult intestinal homeostasis across metazoans (Philpott & Winton, 2014; Hartenstein *et al*, 2017). This suggests that *emc*, the only *Drosophila* Id factor (Garrell & Modolell, 1990; Campuzano, 2001), may have a regulatory function in the adult fly intestine, where it is strongly transcribed (Chintapalli *et al*, 2007; Brown *et al*, 2014; Dutta *et al*, 2015). We examined its expression pattern using the homozygous viable protein trap line *emc$^{CPTI-002740}$* (Lowe *et al*, 2014) and detected Emc in all EBs and most ECs in the posterior midgut (Fig 1A-B). Some EEs and ISCs showed very low levels of expression (Fig 1C-D). This expression pattern, with the described role of several bHLH proteins in intestinal homeostasis, suggested a function of *emc* in promoting EC differentiation.

Emc is required for ISC differentiation

We examined the effect of *emc* loss in the adult posterior midgut by generating *emc* mutant MARCM clones (Lee & Luo, 1999). Clones of the null allele *emc^{AP6}* (Ellis, 1994; Fig S1A) were depleted in differentiated cells and enriched in ISCs, as revealed by expression of the Notch ligand, Delta (Dl) (Ohlstein & Spradling, 2007) (Fig 1E-F, J and Table S1). Clones for the strong hypomorphic allele *emc^1* and the predicted transcriptional null *emc^{LL02590}* (Fig S1A) showed similar enrichment in Dl$^+$ ISCs respect to wild-type clones (Fig 1E, G-J). We then knocked-down *emc* in ISCs and EBs, and traced their offspring using the *escargot* flip-out (FO) approach ("*esg^{TS}-FO*"; Jiang *et al*, 2009; Fig 2A). Loss of *emc* blocked the differentiation of ISCs and EBs, and sometimes the typically small nests of 1-3 *esg$^+$* cells (Ohlstein & Spradling, 2006; de Navascués *et al*, 2012) expanded into larger clusters (Fig 2C). This was suggestive of impairment of Notch signalling. However, these larger clusters did not display the unchecked growth and invasive behaviour observed when Notch signalling is deregulated in this tissue (Micchelli & Perrimon, 2006; Patel *et al*, 2015; see also Fig 4A). Moreover, knockdown of *emc* in the ISC/EB compartment led to the loss of many *esg$^+$* cell nests (identified by their expression of Hdc, an RNA binding protein expressed like *esg*; Resende *et al*, 2017; Fig S1B) (Fig S1C, compare with Fig 2B). This was due to apoptosis, since co-expression of the caspase inhibitor p35 and *emc*-specific RNAi prevents the loss of *esg$^+$* cells (Fig S1D). We conclude that *emc* is necessary to allow differentiation of ISCs and EBs, as well as for their survival.

emc maintains the commitment of EBs

The increased number of Dl+ cells in *emc* depletion conditions conflicted with *emc* expression being predominantly in EBs and ECs. To test whether the loss of *emc* in EBs could induce their expression of Dl, we expressed RNAi against *emc* specifically in EBs using *NRE-Gal4*, *UAS-GFP*, *tub-Gal80^TS* (Zeng *et al*, 2010). Indeed, *emc* knockdown led to EBs reacquiring Dl expression, often while displaying reduced GFP signal (Fig 3A-B). This suggested reduced expression of the EB-specific *NRE-Gal4* and that *emc*-depleted EBs were reverting their identity. To verify this, we used a flip-out lineage tracing approach to irreversibly label EBs ("*NRE^TS-FO*"; Fig 3C) and stained for the mitotic marker phospho-Histone3. In contrast with unmodified EBs, which are post-mitotic and only produce ECs (Zeng & Hou, 2015; Micchelli & Perrimon, 2006; Ohlstein & Spradling, 2006), *emc*-depleted EBs undergo cell division and generate Dl+, proliferative cells (Fig 3D-E). This suggests that *emc*-deficient EBs can not only activate Dl expression but also de-differentiate into functional ISCs, capable of cell division and EE differentiation.

emc acts in parallel to Notch to direct EC differentiation

We next assessed the capacity of *emc* to impose differentiation by overexpressing it with *esg^TS-FO*. Indeed, all ISCs and EBs differentiated into ECs (Fig 2E), as previously reported (Lan *et al*, 2018). This establishes Emc as both necessary and sufficient to force differentiation into the absorptive fate.

This phenotype coincides with that of over-activating the Notch pathway, a known regulator of ISC fate (Micchelli & Perrimon, 2006; Ohlstein & Spradling, 2006; Bardin *et al*, 2010). This suggests that *emc* might be mediating the effects of Notch signalling. To explore this possibility, we tested whether *emc* could force EC differentiation in the absence of Notch. Knockdown of *Notch* with *esg^TS-FO* leads to an invasive expansion of Dl+ ISC-like and Pros+ EE-like cells (Micchelli & Perrimon, 2006; Patel *et al*, 2015; see Fig 4A). When we co-expressed *emc* and *N^RNAi* together, all ISCs and EBs turned into ECs (Fig 4B), strongly suggesting that *emc* operates downstream of Notch. This phenotype, however, could also be due to the excess of Emc quickly driving cells irreversibly into differentiation, before the knockdown of *Notch* resulted in effective clearance of Notch protein. To reduce Notch signalling with the kinetics of protein synthesis, rather than the additive kinetics of RNAi biogenesis, RNA interference, and Notch decay, we overexpressed Hairless (H), the co-repressor associated to the

nuclear effector of N[ICD], Su(H) (reviewed in Bray & Furriols, 2001), with esg^{TS}-FO. As expected, this leads to expansion of Dl+ and Pros+ cells, with a strong reduction of EC differentiation (Fig 4C, G; see also Bardin *et al*, 2010). By contrast, co-expression of *emc* and *H* led to a sharp increase in the levels of EC differentiation (Fig 4D, G). This further indicates that Emc can induce differentiation in the absence of Notch signalling.

Next, we tested whether *emc* mediates the effects of Notch signalling. Activating the pathway by knocking-down *H* leads to an increase in EC differentiation (Bardin *et al*, 2010; Fig 4E) and to the expression of the NRE reporter in all the progenitor, diploid cells, including those expressing Dl (Fig 4H); this is likely due to an increased baseline expression of the NRE reporter, as expected in the absence of H (Furriols & Bray, 2001). Simultaneously knocking down *H* and *emc* did not reduce the increase in EC differentiation induced by overexpression of *H*-specific RNAi, but actually enhanced it. (Fig 4F-H). Taken together, our data suggest that both Emc and Notch signalling can induce EC differentiation independently of each other.

Emc antagonises self-renewal by disrupting Daughterless homodimers

The molecular function of Emc consists in titrating transcriptional activators. Archetypal targets of this inactivation are the class I and II bHLH factors such as Da or Sc, respectively, to which Emc binds directly in vitro and in vivo (Van Doren *et al*, 1991; Cabrera *et al*, 1994; Alifragis *et al*, 1997; Giot *et al*, 2003; Shokri *et al*, 2019). Therefore, the intestinal phenotypes we observed with the loss of *emc* may be due to increased activity of Da and its dimerization partners. Indeed, overexpression of *da* with esg^{TS}-FO recapitulated precisely the effect of *emc* knockdown in *esg+* cell nests. Some of these nests (labelled by *hdc* expression) were expanded (Fig 5A-B, I), and many were lost, their survival rescued by co-expression of p35 (Fig 5B and S2C). Conversely, loss of *da* leads to EC differentiation (Fig S2A-B and 5I; as previously described (Bardin *et al*, 2010; Lan *et al*, 2018) and mimics the excess of Emc (Fig 2E). This suggests that the role of Emc is largely to repress one or more Da-containing bHLH dimers, which in turn maintain stemness and survival. Epistasis experiments further confirmed this, as the simultaneous loss of Emc and Da led to the differentiation of ISCs and EBs into ECs (Fig 5E-H).

We sought to determine the identity of the Da partners involved in maintaining stemness. Most *Drosophila* class II bHLH factors (Moore *et al*, 2000) are expressed in the intestine at very low levels (Fig S2D); several of them are already known not to

have an essential role in ISC maintenance (Bardin *et al*, 2010; Beebe *et al*, 2015; Hartenstein *et al*, 2017). However, Da can form homodimers (Cabrera *et al*, 1994; Cabrera & Alonso, 1991) with demonstrated or suspected functional relevance in several contexts, and often antagonised by Emc (Tapanes-Castillo & Baylies, 2004; Tanaka-Matakatsu *et al*, 2014; D'Rozario *et al*, 2016; Bhattacharya & Baker, 2011; Andrade-Zapata & Baonza, 2014; Troost *et al*, 2015). To test whether Da:Da dimers were involved in preventing differentiation, we overexpressed forced Da homodimers using the *UAS-da:da* tethered construct (Tanaka-Matakatsu *et al*, 2014; Neuhold & Wold, 1993) with *esg^{TS}-FO*. This blocked the differentiation of ISCs and EBs, which accumulated in larger clusters (Fig 5C-D; Table S2). Expression of Da:Da did not lead to *esg^+* cell death (Fig 5D, compare nest density with 5B). This shows that Da homodimers prevent differentiation; it does not rule out that other Da-containing complexes are also necessary to maintain self-renewal. To test this, we took advantage of the *UAS-da^{RNAi}* transgene *P{TRiP.JF02488}*; this targets the 5'UTR of the endogenous *da* transcript, absent in the *UAS-da:da* construct (Fig S2E). Thus, we co-expressed *da:da* and *da^{JF02488}* in ISCs and EBs to evaluate the capacity of Da:Da to prevent differentiation in the absence of endogenous, monomeric Da. The effects were identical to the overexpression of *UAS-da:da* alone (Fig S2F, compare with Fig 5D). This shows that Da can promote ISC self-renewal solely as homodimers, which are antagonised by Emc to allow EC differentiation. Furthermore, our results suggest that Da binds to an unknown partner, to form complexes that are titrated by Emc and whose excess promotes cell death in esg^+ cells.

Basal levels of Sc in esg^+ cells confer ISC properties

While the maintenance of self-renewal corresponds to Da homodimers, the loss of *emc* mimics best the overexpression of monomeric Da, which increases all Da-containing complexes. As *emc* loss induces the elevation of Dl levels, another Da partner, such as Scute, could be mediating this effect. Detectable levels of Sc have recently been found in esg^+ cells (Chen *et al*, 2018; Doupé *et al*, 2018), and members of the *AS-C* are proposed to promote Dl expression during neurogenesis (Kunisch *et al*, 1994; Hinz *et al*, 1994). Therefore, we tested whether the increased levels of Dl observed upon *emc* loss (Figs 2C-D) could be caused by elevated activity of Sc or any of the other bHLH factors belonging to the *AS-C*. We induced *emc* knockdown in MARCM clones, which were either wild-type or homozygous for *sc^{B57}*, a deficiency that removes the entire *AS-C*, and compared their respective levels of Dl expression. Indeed, we observed an

average four-fold decrease in Dl levels in *emc*-depleted clones when they were homozygous for *sc^B57*, indicating that basal Sc levels (and possibly other members of the *AS-C*) can boost Dl expression in ISCs (Fig 6A-C).

These observations prompted us to consider whether Sc could be contributing to additional stem cell properties in *esg^+* cells. Expression of Sc with *esg^TS-FO* clearly leads to increased differentiation into Pros^+ EEs (Bardin *et al*, 2010; Chen *et al*, 2018) and Fig 6D, H). To uncover a role in inducing stem properties, we overexpressed Sc with the EB-specific driver *NRE^TS-FO*. Sc-expressing EBs were capable of division, and their offspring included both Pros^+ and Dl^+ expressing cells, indicating the acquisition of ISC-like properties (Fig 6I, compare with 3D). These functions of Sc depend on the formation of Sc:Da heterodimers, as the overexpression of Sc in *da*-depleted cells prevents the induction of EE differentiation and of Dl expression (Fig S3E).

Sc:Da and Da:Da antagonise each other in the ISC-to-EE transition

We have shown so far that Da homodimers maintain ISC self-renewal, with EC differentiation simply triggered by the Emc-mediated titration of Da. EE differentiation, however, requires the transition from the transcriptional program of Da:Da to that of Da:Sc. Two main scenarios are possible. Sc:Da could initiate a program that silenced, or was epistatic to, that maintained by Da:Da ("succession"). Alternatively, the relative strengths of the two programs could determine the fate outcome for the cell ("antagonism"). To distinguish between these alternatives, we evaluated the ability of an excess of Da monomer to suppress EE differentiation induced by overexpression of Sc with *esg^TS-FO*. Overexpression of *sc* in ISCs and EBs leads to a quick expansion of Pros^+ cells (Fig 6D) (Bardin *et al*, 2010; Chen *et al*, 2018). These were often mitotic (Fig S3A, D), and the number of GFP^+ cells generated in only 3 days of lineage tracing was much higher than those generated during normal tissue replacement (Fig 6D, compare with 2A and 5A). This suggests that some of these cells have been trapped into a pre-EE state, where cells express Pros and Dl and are also capable of mitosis (Zeng & Hou, 2015; Chen *et al*, 2018). However, the co-expression of Da with Sc greatly reduced the number of extra Pros^+ Dl^− cells and led to an increase in Dl^+ cells (Fig 6E, H), while maintaining mitotic figures (Fig S3B, D). Under these conditions, the amount of Sc:Da is expected to be either similar or higher than with the overexpression of Sc alone (as endogenous *da* is weakly expressed (Bardin *et al*, 2010), likely in limiting amounts), with a simultaneous increase in Da homodimers. Thus, the reduction

of EE differentiation and increase of ISC population is indicative of the antagonism, rather than the succession, of their respective transcriptional programs. If that was true, we would expect that the co-expression of tethered Da:Da and Sc would result in even less EE differentiation, as this construct cannot contribute Da monomers to bind to Sc. This was the case, resulting in an even higher amount of Dl+ cells at the expense of Pros+ cell formation (Fig 6F, H), with a concomitant increase in mitotic figures (Fig S3C, D).

To further test the capacity of Da homodimers to antagonise EE differentiation, we co-expressed *da:da* and *Notch^RNAi* with *esg^TS-FO*, as the reduction of Notch leads to the formation of abundant Pros+ EE-like cells (Ohlstein & Spradling, 2006; Bardin *et al*, 2010). Indeed, Da:Da opposed the formation of extra Pros+ cells under conditions of *Notch* loss (Fig 6G, compare with Fig 4A). This further shows that Da:Da can oppose EE differentiation, and suggests that Sc:Da dimers need to overcome the activity of Da:Da dimers to induce the EE fate.

Da:Da promotes ISC identity while Da:Sc confers mixed ISC/EE properties

Our data indicate that Da:Da and Sc:Da activate distinct and antagonistic transcriptional programs to maintain self-renewal and initiate EE differentiation, respectively. However, during the development of the peripheral nervous system, both Da homodimers and Sc:Da heterodimers seem to be able to direct the same cell fate (sensory organ precursor) (Troost *et al*, 2015). To determine the transcriptional signature induced by these factors, we performed RNAseq analysis of the overexpression of either Da, Da:Da or Sc, in ISCs, EBs and ECs simultaneously.

Each overexpression experiment gave a distinct signature (Fig 7A, Fig S4A and Table S3), with Sc inducing modest changes in gene expression, and Da and Da:Da affecting far more genes (Fig 7B). Interestingly, the three conditions had a modest overlap, and monomeric Da induced gene expression changes in many genes unaffected by either Da:Da or Sc, suggesting that it may participate in additional complexes (Fig 7B). We first considered whether the genes upregulated in the three conditions were located in the vicinity of regulatory elements containing the conserved binding site of the bHLH factor, the E-box. Indeed, using the target and enhancer prediction tool i-*cis*Target (Herrmann *et al*, 2012; Imrichova *et al*, 2015) we observed that E-boxes were overrepresented in the predicted regulatory elements of the genes with elevated

expression after Da, Da:Da or Sc overexpression (Tables S4, S5 and S6), suggesting that a good fraction of these genes might be primary targets.

Overexpressing Da or Da:Da led to increased expression of many genes known to be expressed specifically in ISC/EBs and/or required for their maintenance and proliferation (Table S7) like *spdo, polo, Cdk1, mira, HmgD, klu, zfh1, sna, ttk* and multiple components of the Notch pathway (*N, Su(H), H, neur*, and several members of the *E(spl)* complex of HLH/BFM transcription factors (*E(spl)-C*), typical targets and effectors of the pathway); it also led to decreased expression of known mediators of differentiation into EEs (*pros, numb*) or ECs (*Myo31DF*) (Fig 7C-D). By contrast, overexpression of Sc led to increased expression of genes known to promote EE differentiation (*ase, pros, phyl*), as well as of genes expressed specifically in ISCs and/or being required for ISC function, such as *Dl, neur, spdo, polo, Cdk1, pon, sna, mira* and *HmgD* (Fig 7E). This is all in agreement with our phenotypic observations. However, Da:Da or Da overexpression also induced increased expression of genes required for differentiation and/or specifically expressed in differentiated cells (*nub/pdm1, ck*) while reducing expression of other ISC and/or EB marker genes (*Smvt, Oatp58Dc, Myc, bun, insc, Zip71B*) (Fig 7C-D), which shows that the response of individual genes is insufficient to characterise the molecular phenotypes of Da:Da, Da and Sc.

To characterise the molecular signature induced by Da:Da, Da and Sc more systematically, we used Gene Set Enrichment Analysis (GSEA; Subramanian *et al*, 2005) against previously described transcriptional profiles of midgut cell types (Dutta *et al*, 2015) and genes reported to be required for normal homeostasis in the *Drosophila* gut (Zeng *et al*, 2015). Genes upregulated by Da:Da, Da and Sc were enriched in ISC-specific signatures, in line with the genetic requirements of Da as well as our observation that Sc contributes to conferring ISC properties (Fig 7F and Fig S4B-C). Genes regulated by Da and Da:Da also showed depletion of the transcriptional signature of ECs and genes common to differentiated cells, while Da:Da showed specifically a reduction in EE-specific genes (Fig 7F and Fig S4D-E). In turn, Sc-regulated genes showed enrichment in genes specifically expressed in EEs (Fig 7F and Fig S4F) and required for their differentiation (Fig 7G and Fig S4B-C).

We next analysed the transcriptional signatures induced by Da, Da:Da and Sc more generally, to determine whether they indicated changes in specific cellular processes.

Using the Gene List Annotation for Drosophila (GLAD) resource (Hu *et al*, 2015), the transcriptional signatures of Da:Da, Da and Sc overexpression showed enrichment in categories involved in regulatory mechanisms (receptors, transcription factors and DNA binding, GCPRs and signalling pathways, with Notch and EGFR/receptor tyrosine kinases (RTKs) singled out) (Fig 7H and Fig S4I). Interestingly, they also showed depletion of genes involved in metabolism, ribosome, mitochondrial function and oxidative phosphorylation (Fig 7H and Fig S4G-H). This is unlikely to reflect a switch towards glycolytic metabolism, as most glycolytic genes (including *Pfk*, which codes for the rate-limiting enzyme) are generally either not affected or mildly downregulated under these conditions (Table S8). These two features (higher decision-making pathways and lower metabolism) were confirmed with a GSEA of Gene Ontology terms (Biological Process), which show an enrichment of terms related to fate specification and morphogenesis (specifically Notch and RTK pathways), and depletion of metabolic and mitochondrial-related genes. Interestingly, this analysis also shows an enrichment of genes regulating cell cycle and cytokinesis (Table S9 and Fig S4J); in the case of Sc, this is in agreement with its recently described role as a mitotic inducer in the intestine (Chen *et al*, 2018).

These results suggest that Da and Sc, while enhancing largely overlapping developmental and cell cycle regulatory processes, induce distinct signatures which actively promote the ISC and EE identities, respectively. However, Sc can also induce ISC-specific genes. In parallel, Da and Sc promote a state of lower metabolic signature.

Da:Da is antagonised by Notch and maintains self-renewal in parallel to Esg

Esg, a Zinc finger factor of the Snail family, is required to prevent ISC/EB differentiation, mostly through transcriptionally repressing differentiation genes (Korzelius *et al*, 2014; Loza-Coll *et al*, 2014). Da:Da seems to have a similar effect, through the activation of ISC-identity genes. Therefore, we sought to test whether they may be acting in combination to maintain stemness. Knockdown of *esg* in ISCs and EBs leads to differentiation, with an overrepresentation of EEs (Fig S5C and 8F), as expected from the described antagonism between Esg and Sc at the *pros* locus (Li *et al*, 2017). Simultaneous overexpression of *UAS-esg^RNAi* and *UAS-da:da* prevented most differentiation, and specifically reduced EE cell production (Fig 8D and 8F), indicating that Da:Da can maintain ISCs and oppose EE differentiation independently

of *esg*. In turn, overexpression of *esg* blocked differentiation irrespective of the presence of *da* (Fig 8C, F and Fig S5B), indicating that Da:Da and Esg largely act independently in ISC maintenance. Furthermore, expression of *da* and *esg* are independent of each other (Fig S5D, E).

We next sought to determine whether Da:Da operates as part of a pathway with the known regulators of ISC fate. Our previous epistatic analysis with *emc* and Notch signalling suggested that Da:Da is required independently of the Notch pathway. Indeed, when we knocked-down both *N* and *da* using *esg^{TS}-FO*, all ISCs/EBs became ECs (Fig S5A). Similar results were obtained by simultaneously knocking-down *da* and inhibiting Notch signalling by overexpressing *H* (Fig 8A, compare with Fig 4C, and Fig 8E). However, expressing *UAS-da:da* prevented the differentiation induced by increasing Notch signalling by knockdown of *H* (Fig 8B, compare with Fig 4E, and Fig 8E), suggesting that Da:Da is antagonised by Notch signalling. Together, our data indicate that Da homodimers operate downstream of Notch and in parallel to Esg to maintain ISC self-renewal.

DISCUSSION

Multipotent adult stem cells must be able to change their transcriptional state from self-renewal to several alternative differentiation programs. bHLH transcription factors seem well placed to coordinate these changes molecularly; they have been described to operate synergistically, antagonistically or combinatorially (Amoutzias *et al*, 2008; Bhattacharya & Baker, 2011; Troost *et al*, 2015; Sallee *et al*, 2017); their capacity for dimerization allows reduction in molecular noise (Bundschuh *et al*, 2003). Here we show that a bHLH factor network controls three alternate fates in the adult *Drosophila* intestine. Class I homodimers (Da:Da) promote the progenitor state of ISCs/EBs. Progenitor cells will acquire the absorptive fate by titration of Da by the HLH factor Emc, or the secretory fate by expression of the class II bHLH Sc, which heterodimerises with Da to initiate EE differentiation. Moreover, Emc is required in EBs to maintain their committed state, while low levels of Sc seem to contribute to boost the ISC transcriptional program. Our results argue for a central and conserved role of bHLH factors in the acquisition and maintenance of multiple fates in the metazoan intestine.

Three fates regulated by a dimerization network

Our data show that Da homodimers maintain ISCs and EBs undifferentiated, but this situation can be easily swayed towards either EC or EE differentiation by Da:Da swapping partners with Emc or Sc (Fig 9). Similar networks involving class I, II and V bHLH factors have been found to regulate the development of the *Drosophila* retina (Bhattacharya & Baker, 2011) and the peripheral nervous system (PNS) (Cubas *et al*, 1991; Van Doren *et al*, 1991; Troost *et al*, 2015). However, in these cases only one new fate is accessed (neural), and Da:Da dimers seem to have a relatively 'accessory' role. In the retina, Da:Da dampens the titration by Emc of the bHLH proneural factor Ato (Bhattacharya & Baker, 2011; Li & Baker, 2018). In the PNS, Da:Da cooperate with Da:Ac/Sc heterodimers to overcome Emc inhibition and induce neural fate, which both Da:Ac/Sc and Da:Da can induce (Troost *et al*, 2015). By contrast, in the adult midgut Da:Da and Da:Sc support distinct fates (progenitor and secretory, respectively) while Emc titrates both dimers to allow EC differentiation, enabling the network to regulate a three-way fate decision. This is possible because Da:Da and Da:Sc induce specific but partially overlapping transcriptional profiles (Fig 7) which yet seem to be able to act antagonistically (Fig 6). By contrast, Emc simply prevents the DNA binding of either Da

or Sc (Van Doren *et al*, 1991; Cabrera *et al*, 1994), which suggests that EC is the default fate when Da:Da dimers do not operate (Fig 5).

Interestingly, bHLH transcription factors of classes I-II-V (or E/A/Id) are also active in the mammalian intestine. Class II bHLH factors promote EE differentiation (reviewed in Philpott & Winton, 2014) and ISC maintenance (van der Flier *et al*, 2009). However, class I factors E2a and Heb are specifically expressed in crypt cells in the mouse intestine (van der Flier *et al*, 2009). Our work suggests that they may function as class I dimers as well as dimerization partners for class II factors. Meanwhile, Emc homologs Id2 and Id3 are expressed specifically in the intestinal differentiated cells (Wice, 1998), suggesting similar functions. By contrast, Id1 is required in ISCs for regeneration (Zhang *et al*, 2014) and can act as an oncogene (Wice, 1998), which is the opposite behaviour to that of Emc.

A balance of bHLH factors regulates intestinal stem cell fate

Our observations indicate that additional work on the mechanisms that regulate the transcription of both *emc* and *sc* would be very informative. The *cis*-regulatory region of *emc* seems to be complex and spreads across a large genomic region (Li & Baker, 2019), so it is likely that is subject to multiple regulatory inputs. An obvious candidate for regulation of *emc* transcription is Notch signalling, as shown in other contexts (Baonza *et al*, 2000; Baonza & Freeman, 2001; Adam & Montell, 2004; Bhattacharya & Baker, 2009; Spratford & Kumar, 2015), and considering the importance of Notch for EB establishment and EC differentiation (Micchelli & Perrimon, 2006; Ohlstein & Spradling, 2006; Bardin *et al*, 2010). However, our epistasis analyses suggest that Emc and Notch signalling can induce EC differentiation independently of each other, observation that makes Notch an unlikely regulator of *emc* in the midgut. A major regulator of Sc is the transcriptional repressor Ttk69 (Wang *et al*, 2015), whose stability is in turn regulated by the E3 ubiquitin ligase Sina and its adaptor Phyl (Yin & Xi, 2018). Therefore, Phyl expression could be the trigger to allow the accumulation of Sc and initiate EE differentiation, but how *phyl* is regulated is not known yet. On the other hand, Chen et al. (2018) observed that Sc activates itself as well as *E(spl)m8*, a member of the *Enhancer of split Complex* (which comprises several transcription factors, many of which are often both targets and effectors of the Notch pathway; reviewed in Delidakis *et al*, 2014). As E(spl)m8 is able to repress Sc, the authors proposed that a cell-autonomous oscillator involving Sc and *E(spl)-C* members would

create pulses of Sc expression which would set the pace of EE production (Chen *et al*,
2018). However, while we too observe induction of *E(spl)-C* genes upon Sc forced
expression, so we do as a result of Da:Da and Da overexpression (Fig 7). This
suggests that the regulation of *E(spl)-C* (and therefore of *sc*) may be more
complicated. Additional work will be necessary to determine the sequence and timing
of regulatory inputs that initiate fate-inducing levels of Sc or Emc.

Moreover, the Emc/Da/Sc network clearly does not work as a simple boolean switch.
The three genes are expressed, and seem to be functionally relevant, in both ISCs and
EBs (Bardin *et al*, 2010; Chen *et al*, 2018; Doupé *et al*, 2018; this work). This argues
against a mechanism that simply initiates Emc or Sc expression to induce the EC or
EE fates. Moreover, the functions of Emc and Sc seem to be more complex than acting
as mere fate switches. Adequate levels of Emc are required in the EB, as its depletion
induces de-differentiation into ISCs (Fig 3), but its overexpression accelerates EB
terminal differentiation into ECs (Fig 2). Intriguingly, it has recently been proposed that
the Emc homolog Id1 protects pluripotency specifically during the peri-implantation
transition (Malaguti *et al*, 2019), which suggests that Emc could be part of a timing
mechanism in the transition between ISC and EC. As for Sc, our data suggests that
low levels of expression in ISCs (Chen *et al*, 2018; Doupé *et al*, 2018) induce the
expression of ISC-specific genes (Fig 7) without eliciting a conversion into pre-EE cell.
Higher levels would be required to initiate EE differentiation. Therefore, the amounts of
Da, Sc and Emc must be maintained in balance. In the case of Emc, this is likely to be
achieved through post-transcriptional mechanisms. During neurogenesis, Emc is
stabilised when bound to Da, and degrades when Da frees itself to bind a proneural
factor (Bhattacharya & Baker, 2011; Li & Baker, 2018). This may be the reason why
Emc shows higher levels of accumulation in EBs and ECs (Fig 1), where Da is
expressed but other bHLH factors are at minimal levels.

Da and Sc cooperate to endow cells with ISC properties

Our data shows a critical requirement for Da:Da in ISC maintenance, but also that Sc
can contribute to enhance ISC features (proliferation, expression of Dl). This is likely an
active promotion of ISC fate, rather than just prevention of differentiation, as our
transcriptomic analysis shows an enrichment of the ISC transcriptional signature when
overexpressing either Da:Da, Da or Sc, together with a depletion of differentiation
genes (except for EE-specific genes and Sc). This active role is further supported by

the capacity of Sc to induce EBs to de-differentiate into ISCs and re- or transdifferentiate into EEs (Fig 6). The transcriptional changes induced by Sc, Da and Da:Da expression also show an enrichment in receptors and ligands of signalling pathways (Fig 7). This is in agreement with the recent proposal that ISCs are sources of autocrine signals required for their own maintenance (Doupé *et al*, 2018). We also find evidence that mitochondrial function is reduced by expression of Sc, Da and Da:Da (Fig 7), which fits well the observation that ISCs display an 'empty mitochondria' phenotype under the electron microscope (Hung *et al*, 2018). Together, this suggests that low levels of Sc:Da dimers contribute non-critically to the maintenance of the ISC identity and function, and we surmise that Emc expression in EBs might be required to dampen this Sc activity.

Regulation of the committed state of the enteroblast

Under normal conditions, EBs will only give rise to mature ECs without further division (Zeng & Hou, 2015; Wang *et al*, 2015; Yin & Xi, 2018). Depletion of Emc specifically in EBs coupled to lineage tracing results in cells proliferating and expressing Dl (Fig 3), which likely represents a de-differentiation process. This may partially reflect the need to control Sc activity in the EB, as the expression of Sc can induce the same effects (Fig 6). Intriguingly, EB-specific depletion of Ttk69 or Klu or overexpression of Phyl seem to induce trans-differentiation, as these conditions produce Pros[+] cells (Wang *et al*, 2015; Yin & Xi, 2018). This difference could be because Emc is also influencing the activity of multiple Da-containing complexes in the EB, or because of the activity of additional targets of Klu and the Phyl/Ttk69 pathway than Sc. The reacquisition of proliferation and Delta expression is also controlled by other factors, such as Sox21a (Zhai *et al*, 2015), or the global co-repressor Gro, acting downstream of Notch (Guo *et al*, 2019) (though surprisingly Notch itself does not seem to be required to maintain the EB; Siudeja *et al*, 2015). Together with our observations, this indicates a previously unappreciated plasticity of the EB, which is consistent with its similarity to the ISC in transcriptome (Hung *et al*, 2018) and chromatin accessibility landscape (Aughey *et al*, 2018). Importantly, this resembles the behaviour of EC precursors in the mammalian intestine, which can dedifferentiate and repopulate the intestinal crypt during regeneration (Tetteh *et al*, 2016).

MATERIALS AND METHODS

Drosophila culture and genetics

Flies were raised and maintained on standard cornmeal/yeast medium at 25°C unless specifically indicated. Experiments were conducted in mated females, collected daily and maintained in fresh vials with added yeast until 4-7 days old. For experiments using Gal80^{TS}, flies were reared and aged to gut maturity (~7 days) at 18°C, then transferred to 29 °C. For induction of MARCM and flip-out clones, 4-7 days old flies were treated at 37°C for 60 or 15 min, respectively. Flies were aged for 7 days after induction treatment before dissection, unless otherwise indicated. See Appendix for fly strains.

Immunohistofluorescence, confocal microscopy and image analysis

Stainings were performed as in Bardin *et al*, (2010). See Appendix for details on primary and secondary antibodies. DNA dye was Hoechst (Sigma-Aldrich B2261 used 1:5000 from a stock solution at 10 mg/ml). Mounting medium was 4% N-propyl-galate, 80% glycerol. Confocal stacks were obtained in a Zeiss LSM 710 with an EC Plan-Neofluar 40X oil immersion objective (numerical aperture 1.3). All stack positions were acquired in the posterior midgut. Typically, three positions along the anterior-posterior axis of the posterior midgut were acquired for each organ. In MARCM clone experiments, stacks were acquired from all clones in the posterior midgut. Quantification of anti-Dl staining is detailed in Images and figures were assembled using Adobe Photoshop and Illustrator CS6.

Cell counts and statistics

For evaluating the proportion of cell types in GFP$^+$ clones or labelled tissue, confocal stacks were maximum-intensity projected using FIJI/ImageJ (Schindelin *et al*, 2012), cells of the relevant types were counted with the Cell Counter plugin. In the experiments overexpressing *UAS-sc* and *UAS-da:da*, the associated increase in proliferation generated large, highly densely populated cell clusters which could not be counted with single-cell precision. Therefore, for this genotype we estimated the proportion of each cell population in each field of view separately. See the Appendix for details on cell type identification.

Statistical tests were performed in Prism 8 or in R 3.5.1 (R Core Team, 2018) using RStudio 1.1.383. Statistical significance of changes in proportions of cell types was assessed by binomial logistic regression in basic R. We used Firth's bias reduced logistic regression (package *logistf*; Heinze and Ploner, 2013) in experiments with zero observations in one or more of the genotypes considered, to avoid the nonsensical results arising from the 'complete separation' of data (Albert & Anderson, 1984).

RNA-seq

Flies bearing either *UAS-da:da*, *UAS-da* or *UAS-sc* as well as *esg-Gal4*, *Myo1A-Gal4* and *tub-Gal80^{TS}* were reared at 18ºC until 4-7 days old, transferred to 29ºC for 24h and their posterior midguts dissected, collected in lysis buffer and frozen at -80ºC, then thawed and extracted for total RNA. Libraries from three biological replicates per condition were prepared and ~26 million reads per library were generated using Illumina technology. See the Appendix for additional details.

ACKNOWLEDGEMENTS

We thank Nicholas Baker, Allison Bardin, Antonio Baonza, Sonsoles Campuzano, Sangbin Park, Bruce Edgar, Mike Taylor, Shinya Yamamoto, the Bloomington *Drosophila* Stock Center, the Vienna *Drosophila* Resource Center, the *Drosophila* Genetics Resource Center (Kyoto) and the National Institute of Genetics (Japan) for providing fly stocks, and the Developmental Studies Hybridoma Bank (University of Iowa) for supplying antibodies.

We would like to thank Juan Modolell, Sonsoles Campuzano, Catherine Hogan, Fernando dos Anjos-Afonso, Florian Siebzehnrubl and Terrence Trinca for critical comments on the manuscript, Sonia López de Quinto, Helen White-Cooper, Mike Taylor and Wynand van der Goers van Naters for useful discussions during the development of the project, and the data analysis team of the College of Biomedical and Life Sciences and the Genomics facility of the School of Biosciences of Cardiff University for technical assistance. This work was supported by Cardiff University.

AUTHOR CONTRIBUTIONS

Conception of the study and supervision: JdN. Data acquisition, analysis and interpretation of data: AP and JdN. Experimental design; writing, revising and approval of the manuscript: JdN and AP.

CONFLICT OF INTEREST

The authors declare no competing interest.

FIGURE LEGENDS

Figure 1. *emc* is expressed in the absorptive lineage and is required for

differentiation.

A-D. Projected confocal stacks showing endogenous Emc:GFP expression. Emc is expressed in EBs (marked with *NRE-lacZ*, arrowheads in A, A') and most ECs (positive for *myoIA-lacZ*, solid arrowheads in B, B') but not all (empty arrowhead in B, B'). *emc* expression can also be detected in some ISCs (Dl⁺) and EEs (Pros⁺) (solid arrowheads and asterisks, respectively, in C-D') but not all (empty arrowheads and asterisks in C-D').

E-I. Differentiation is impaired in *emc* mutant clones. Control clones display more polyploid Dl⁻/Pros⁻ ECs (asterisks in E, H) than clones homozygous of for the mutant alleles *emc^{AP6}* (F), *emc^1* (G) and *emc^{LL02590}* (I), which are enriched in Dl⁺ ISCs (solid arrowheads) and diploid Dl⁻/Pros⁻ EBs (empty arrowheads) with respect to controls.

J. Stacked bars plots summarising the cell type composition of clones by genotype. Numbers (cells/clones) are: 194/106, 213/118, 310/117, 112/95 and 208/200 for control *FRT2A*, *emc^{LL02590}*, control *FRT80B*, *emc^1*, and *emc^{AP6}* respectively. p-values are <0.05, <0.01 and <0.001 for one, two or three asterisks (see Tables S1 and S2 for further details).

Scale bar: 20µm in all panels.

Figure 2. *emc* is necessary for differentiation and sufficient for selection of the EC fate.

A. The *esg^{TS}-FO* method allows the temporal control of expression in progenitor cells (esg⁺ ISCs and EBs) and their lineage tracing to access their fate potential. *UAS-GFP* and *UAS-FLP* are expressed specifically in ISCs and EBs after induction by temperature switch. This leads to the progenitor-restricted expression of the (potentially) ubiquitous *Actin5C-Gal4* by excision of a transcriptional terminator. *Actin5C-Gal4* maintains *UAS-GFP* expression in the ISC/EB lineage when differentiation occurs and *esg* expression stops.

B-E. Emc is required for differentiation and sufficient for induce the EC fate. ISCs and EBs in control guts produce both EC (Dl⁻/Pros⁻) and EE (Dl⁻/Pros⁺) GFP⁺ cells (B, note the difference in intensity levels between Dl and Pros, respectively solid and empty arrowheads), while differentiation is reduced when ISCs/EBs express *emc*-specific RNAi , shown by increased number of Dl⁺/GFP⁺ cells (C, D). Reduction of *emc* also induces increased Dl expression: compare Dl-expressing with Pros-expressing cells

(respectively solid and empty arrowheads in C, D, compare with those similarly marked

in B). Forced expression of *emc* in ISCs/EBs forces differentiation into ECs (E).

F. Stacked bars plots summarising the cell type composition of *esg^{TS}-FO* tissue.

Numbers (total GFP$^+$ cells / fields of view) are 1046/15, 104/21, 832/15 and 812/10 for

control, *UAS-emc^{RNAi}_{NIG}, UAS-emc^{RNAi}_{KK+HMS}, and UAS-emc*, respectively. p-values are

<0.05, <0.01 and <0.001 for one, two or three asterisks (see Tables S1 and S2).

Scale bar: 20μm in all panels.

Figure 3. *emc* is required for maintenance of the EB committed state.

A-B. The *NRE-Gal4* driver is expressed in EBs at different levels in normal intestines

(solid and empty arrowheads in A for higher and lower-expressing EBs), but never

co-express Dl. When *emc* is depleted with *NRE-Gal4*, many EBs co-express Dl (empty

arrowheads, B) but not all (solid arrowheads, B). Also, EEs develop close to

emc-depleted EBs (arrow, B).

C. Use of *NRE^{TS}-FO* allows the simultaneous manipulation and tracing of the EB

lineage. *NRE-Gal4, UAS-GFP* is expressed specifically in EBs after induction and

activates specifically in EBs the (potentially) ubiquitous *Actin5C-Gal4*, which in turn will

maintain *UAS-GFP* expression if differentiation or decommitment occur.

D-E. Loss of *emc* in EBs leads to their decommitment. Activation of *NRE^{TS}-FO* in

control guts leads exclusively to the labelling of EBs and ECs, with no co-expression of

GFP and Dl or Pros (solid and empty arrowheads, respectively, in D; some ISCs and

EEs in the micrograph seem inside the GFP territory due to the Z-projection of the

confocal stack). By contrast, in *NRE^{TS}-FO* guts where *emc* is depleted, EB-derived

cells express Dl and are capable of mitosis, as judged by phospho-Histone 3 staining

(arrowhead in E).

Scale bar: 20μm in all panels.

Figure 4. *emc* acts in parallel to Notch signalling.

A-D. *emc* can direct EC differentiation in the absence of Notch signalling. EC

differentiation is abolished or strongly impaired with knockdown of *N* (A) or

overexpression of H (C), respectively, with accumulation of Dl$^+$ or Pros$^+$ cells (solid and

empty arrowheads, respectively, in A, C). Simultaneous overexpression of *emc*

induces EC differentiation in both cases (arrowheads in B, D), but to a lesser an extent

in a background of *H* overexpression (asterisks in D, compare with B).

E-F. Notch signalling can direct EC differentiation in the absence of *emc*. Loss of EC

differentiation caused by depletion of *emc* (see panel H and Fig. 2C-D, F) is rescued by

simultaneous knockdown of *H* (F). Depletion of *H* can induce EC differentiation on its own (E).

G-H. Stacked bars plots summarising the cell type composition of *esg^{TS}-FO* tissue. Numbers (total GFP$^+$ cells / fields of view) are 1810/21 and 1292/27 for *UAS-H* and *UAS-H+UAS-emc*, respectively (G) and 984/15 and 126/16 for *UAS-H$^{RNAi}_{HMS}$* and *UAS-H$^{RNAi}_{HMS}$+UAS-emc$^{RNAi}_{NIG}$*, respectively (H). p-values are <0.05, <0.01 and <0.001 for one, two or three asterisks (see Tables S1 and S2).

Scale bar: 20μm in all panels.

Figure 5. Da homodimers prevent differentiation and are antagonised by Emc.

A-D. Da homodimers maintain ISCs/EBs undifferentiated. Solid arrowheads: ISCs/EBs; empty arrowheads: ECs/EEs. Overexpression of Da with *esg^{TS}-FO* prevents formation of ECs, though it allows EE differentiation (B, compare with A). Expression of a forced Da:Da dimer in esg$^+$ cells prevents all differentiation, as evaluated with Hdc/NRE-lacZ (C) or Dl (D).

E-H. *da* is epistatic over *emc*. Solid arrowheads: ECs; empty arrowheads: Dl$^+$ ISCs. MARCM clones mutant for *da^{10}* differentiate as ECs (E). This phenotype is recapitulated when these clones simultaneously express *emc*-specific RNAi (F). Likewise, impaired differentiation in *emc^{LL02590}* clones (G) is rescued by simultaneous knockdown of *da*, which leads to EC formation (H).

I. Stacked bars plots summarising the cell type composition of *esg^{TS}-FO* tissue. Numbers (total GFP$^+$ cells / fields of view) are 1135/15, 549/14, 693/13, 803/15 and 1086/19 for control, *UAS-da*, *UAS-da:da*, *UAS-da$^{RNAi}_{JF}$* and *UAS-da$^{RNAi}_{HMS}$*, respectively. P-values are <0.05, <0.01 and <0.001 for one, two or three asterisks (see Tables S1 and S2).

Scale bar: 20μm in all panels.

Figure 6. Sc:Da and Da:Da antagonise each other in EE formation but collaborate in ISC maintenance.

A-C. Emc antagonises Sc in inducing Dl expression. Loss of the *AS-C* using the *sc^{B57}* deficiency reduces the increase in Dl expression observed with knockdown of *emc* (B, compare with A). Dl levels are quantified by comparison with those of Pros (see Appendix). Numbers (Dl$^+$ cells/Pros$^+$ cells) are 210/146 and 718/210 for *UAS-emc$^{RNAi}_{NIG}$* and *UAS-emc$^{RNAi}_{NIG}$, Df(1)sc^{B57}*, respectively. P-values are <0.05, <0.01 and <0.001 for one, two or three asterisks (Mann-Whitney test).

D-F. Antagonism/collaboration between Sc and Da. Solid arrowheads: Dl[+]/Pros[+] pre-EEs; empty arrowheads: Dl[+] ISCs; asterisks: Pros[+] EEs. Overexpression of Sc leads to a dramatic increase of Dl[+]/Pros[+] pre-EEs and Pros[+] EEs, but maintains a population of Dl[+] ISCs (D). The population of ISCs progressively increases by co-expression of monomeric Da (E) and dimeric Da:Da (F), at the cost of EE differentiation (E, F) and pre-EE formation (F). Quantified in H.

G. Da:Da antagonises EE formation. While reduction of Notch signalling leads to an excess of EEs, simultaneous overexpression of Da:Da and *N*-specific RNAi results in very few Pros[+] cells (solid arrowheads, compare with Fig 4A).

H. Stacked bars plots summarising the cell type composition of *esg[TS]-FO* tissue. Numbers (total GFP[+] cells / fields of view) are 1241/6, 2131/10 and not determined/11 for *UAS-sc, UAS-sc, UAS-da*, and *UAS-sc, UAS-da:da*, respectively. p-values are <0.05, <0.01 and <0.001 for one, two or three asterisks (see Tables S1 and S2).

I. Sc can induce decommitment of EBs. Expression of Sc with *NRE[TS]-FO* leads to reacquisition of ISC properties (Dl expression, mitosis assessed with phospho-Histone 3, empty arrowhead) and pre-EE formation (Pros expression and mitosis, solid arrowhead).

Scale bar: 20µm in all panels.

Figure 7. Da and Sc cooperate to induce expression of the ISC signature.

A-B. Da:Da, Da and Sc induce specific but overlapping transcriptional signatures. Hierarchical clustering of differentially expressed genes (fold change > 4) (A) shows groups of genes specifically increased or reduced in Da, Da:Da and Sc respect to the wild-type. UpSetR plots (B) indicate the degree of overlap between these signatures, which is highest between Da and Da:Da.

C-E. MA plots ($\log_2$ fold change vs mean expression in transcripts per million, tpm) for the overexpression of Da:Da, Da and Sc, respectively. Some genes of interest are identified because of their role or expression pattern.

F-H. Heatmaps of gene set enrichment scores for sets of genes that are: expressed in specific cell types (Dutta *et al*, 2015) (F), known to be functionally relevant in different aspects of intestinal homeostasis (Zeng *et al*, 2015) (G) or annotated because of their broad molecular function in *Drosophila* (Hu *et al*, 2015). Black asterisks indicate the enrichment scores with p-values < 0.05 in that heatmap. White crosses indicate the enrichment scores with p-values > 0.05 in that heatmap; the rest have p-values < 0.05.

Figure 8. *da* is epistatic over Notch signalling but acts in parallel with *esg*.

Solid arrowheads: ISCs (Hdc$^+$/NRE$^-$ or Dl$^+$/NRE$^-$); empty arrowheads: EEs (Pros$^+$).

A-B. *da* is epistatic over Notch signalling. Loss of *da* leads to EC and EE differentiation when Notch signalling is impaired by overexpression of *H* (A), while overexpression of *da:da* maintains cell undifferentiated even when Notch signalling is exacerbated by knockdown of *H* (B).

C-D. *da* and *esg* function in parallel to prevent differentiation. Overexpression of *esg* can prevent differentiation and compensate for the loss of *da* (C), while overexpression of *da:da* can similarly compensate for the loss of *esg*, including prevention of EE formation (D).

E-F. Stacked bars plots summarising the cell type composition of *esg*^{TS}-*FO* tissue. Numbers (total GFP$^+$ cells / fields of view) are 1510/21, 1810/21, 613/8 and 984/15 for *UAS-H, UAS-H+UAS-da$^{RNAi}_{JF}$, UAS-H$^{RNAi}_{HMS}$* and *UAS-H$^{RNAi}_{HMS}$+UAS-da:da*, respectively (E) and 850/9, 1236/12, 628/18 and 939/19 for for *UAS-esg UAS-esg+UAS-da$^{RNAi}_{JF}$, UAS-esg$^{RNAi}_{HMS}$* and *UAS-esg$^{RNAi}_{HMS}$+UAS-da:da*, respectively (F). p-values are <0.05, <0.01 and <0.001 for one, two or three asterisks (see Tables S1 and S2 for further details). Scale bar: 20μm in all panels.

Fig 9. Sc, Da and Emc regulate self-renewal and bipotential differentiation in the intestine.

Orange arrows: Da homodimers maintain ISCs and EBs undifferentiated, antagonising EC differentiation and pre-EE formation by inducing expression of ISC-specific genes. Pink arrows: An increase of Sc expression induces formation of Da:Sc heterodimers, which steer the transcriptional program towards pre-EE formation; Sc can also induce expression of the ISC-specific signature, potentially leading to EB decommitment if its activity is not prevented by titration by Emc. Green arrows: Emc increased expression in EBs maintains commitment for EC differentiation and can direct terminal EC differentiation by interruption of Da:Da and Da:Sc transcriptional activities. This occurs in parallel to Esg prevention of differentiation and Notch signalling being upstream and opposing Da:Da to induce EC differentiation.

REFERENCES

Adam JC & Montell DJ (2004) A role for extra macrochaetae downstream of Notch in follicle cell differentiation. *Development* **131:** 5971–5980

Albert A & Anderson JA (1984) On the existence of maximum likelihood estimates in logistic regression models. *Biometrika* **71:** 1–10

Alifragis P, Poortinga G, Parkhurst SM & Delidakis C (1997) A network of interacting transcriptional regulators involved in *Drosophila* neural fate specification revealed by the yeast two-hybrid system. *Proc Natl Acad Sci USA* **94:** 13099–13104

Amcheslavsky A, Song W, Li Q, Nie Y, Bragatto I, Ferrandon D, Perrimon N & Ip YT (2014) Enteroendocrine cells support intestinal stem cell-mediated homeostasis in *Drosophila*. *Cell Reports* **9:** 1–8

Amoutzias GD, Robertson DL, Van de Peer Y & Oliver SG (2008) Choose your partners: dimerization in eukaryotic transcription factors. *Trends Biochem. Sci.* **33:** 220–229

Andrade-Zapata I & Baonza A (2014) The bHLH factors Extramacrochaetae and Daughterless control cell cycle in *Drosophila* imaginal discs through the transcriptional regulation of the cdc25 phosphatase *string*. *PLoS Genet* **10:** e1004233–16

Aughey GN, Estacio Gomez A, Thomson J, Yin H & Southall TD (2018) CATaDa reveals global remodelling of chromatin accessibility during stem cell differentiation in vivo. *eLife* **7:** 6061

Baonza A & Freeman M (2001) Notch signalling and the initiation of neural development in the *Drosophila* eye. *Development* **128:** 3889–3898

Baonza A, de Celis JF & Garcia-Bellido A (2000) Relationships between extramacrochaetae and Notch signalling in *Drosophila* wing development. *Development* **127:** 2383–2393

Bardin AJ, Perdigoto CN, Southall TD, Brand AH & Schweisguth F (2010) Transcriptional control of stem cell maintenance in the *Drosophila* intestine. *Development* **137:** 705–714

Baylies MK, Bate M & Ruiz-Gómez M (1997) The specification of muscle in Drosophila. *Cold Spring Harb Symp Quant Biol* **62:** 385–393

Beebe K, Park D, Taghert PH & Micchelli CA (2015) The *Drosophila* prosecretory transcription factor *dimmed* is dynamically regulated in adult enteroendocrine cells and protects against gram-negative infection. *G3 (Bethesda)* **5:** 1517–1524

Bertrand N, Castro DS & Guillemot F (2002) Proneural genes and the specification of neural cell types. *Nat Rev Neurosci* **3:** 517–530

Bhattacharya A & Baker NE (2009) The HLH protein Extramacrochaetae is required for R7 cell and cone cell fates in the *Drosophila* eye. *Developmental Biology* **327:** 288–300

Bhattacharya A & Baker NE (2011) A network of broadly expressed HLH genes regulates tissue-specific cell fates. *Cell* **147:** 881–892

Bray S & Furriols M (2001) Notch pathway: Making sense of Suppressor of Hairless. *Curr Biol* **11:** R217–R221

Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, Ben W Booth, Wen J, Park S, Suzuki AM, Wan KH, Yu C, Zhang D, Carlson JW, Cherbas L, Eads BD, Miller D, Mockaitis K, Roberts J, Davis CA, et al (2014) Diversity and dynamics of the *Drosophila* transcriptome. *Nature* **512:** 393–399

Bundschuh R, Hayot F & Jayaprakash C (2003) The role of dimerization in noise reduction of simple genetic networks. *J Theor Biol* **220:** 261–269

Cabrera CV & Alonso MC (1991) Transcriptional activation by heterodimers of the achaete-scute and daughterless gene products of *Drosophila*. *EMBO J* **10:** 2965–2973

Cabrera CV, Alonso MC & Huikeshoven H (1994) Regulation of scute function by extramacrochaete in vitro and in vivo. *Development* **120:** 3595–3603

Campuzano S (2001) Emc, a negative HLH regulator with multiple functions in *Drosophila* development. *Oncogene* **20:** 8299–8307

Chen J, Xu N, Wang C, Huang P, Huang H, Jin Z, Yu Z, Cai T, Jiao R & Xi R (2018) Transient Scute activation via a self-stimulatory loop directs enteroendocrine cell pair specification from self-renewing intestinal stem cells. *Nature Cell Biology* **20:** 152–161

Chintapalli VR, Wang J & Dow JAT (2007) Using FlyAtlas to identify better Drosophila melanogaster models of human disease. *Nat Genet* **39:** 715–720

Cinquin O & Demongeot J (2002) Positive and negative feedback: Striking a balance between necessary antagonists. *J Theor Biol* **216:** 229–241

Cinquin O & Demongeot J (2005) High-dimensional switches and the modelling of cellular differentiation. *J Theor Biol* **233:** 391–411

Cinquin O & Page KM (2007) Generalized, switch-like competitive heterodimerization networks. *Bull. Math. Biol.* **69:** 483–494

Crosnier C, Stamataki D & Lewis J (2006) Organizing cell renewal in the intestine: stem cells, signals and combinatorial control. *Nat. Rev. Genet.* **7:** 349–359

Cubas P, de Celis JF, Campuzano S & Modolell J (1991) Proneural clusters of achaete-scute expression and the generation of sensory organs in the *Drosophila* imaginal wing disc. *Genes & Development* **5:** 996–1008

de Navascués J, Perdigoto CN, Bian Y, Schneider MH, Bardin AJ, Martinez-Arias A & Simons BD (2012) *Drosophila* midgut homeostasis involves neutral competition between symmetrically dividing intestinal stem cells. *EMBO J* **31:** 2473–2485

Delidakis C, Monastirioti M & Magadi SS (2014) E(spl): Genetic, developmental, and evolutionary aspects of a group of invertebrate Hes proteins with close ties to Notch signaling. *Curr Top Dev Biol* **110:** 217–262

Doupé DP, Marshall OJ, Dayton H, Brand AH & Perrimon N (2018) *Drosophila* intestinal stem and progenitor cells are major sources and regulators of homeostatic niche signals. *Proceedings of the National Academy of Sciences* **115:** 12218–12223

Dutta D, Dobson AJ, Houtz PL, Gläßer C, Revah J, Korzelius J, Patel PH, Edgar BA & Buchon N (2015) Regional cell-specific transcriptome mapping reveals regulatory complexity in the adult *Drosophila* midgut. *Cell Reports* **12:** 1–13

D'Rozario M, Zhang T, Waddell EA, Zhang Y, Sahin C, Sharoni M, Hu T, Nayal M, Kutty K, Liebl F, Hu W & Marenda DR (2016) Type I bHLH Proteins Daughterless and Tcf4 Restrict Neurite Branching and Synapse Formation by Repressing Neurexin in Postmitotic Neurons. *Cell Reports* **15:** 386–397

Ellis HM (1994) Embryonic expression and function of the Drosophila helix-loop-helix gene, extramacrochaetae. *Mech Dev* **47:** 65–72

Furriols M & Bray S (2001) A model Notch response element detects Suppressor of Hairless-dependent molecular switch. *Curr Biol* **11:** 60–64

Garrell J & Modolell J (1990) The Drosophila extramacrochaetae locus, an antagonist of proneural genes that, like these genes, encodes a helix-loop-helix protein. *Cell* **61:** 39–48

Giot L, Bader JS, Brouwer C, Chaudhuri A, Kuang B, Li Y, Hao YL, Ooi CE, Godwin B, Vitols E, Vijayadamodar G, Pochart P, Machineni H, Welsh M, Kong Y, Zerhusen B, Malcolm R, Varrone Z, Collis A, Minto M, et al (2003) A protein interaction map of Drosophila melanogaster. *science (New York, NY)* **302:** 1727–1736

Graf T & Enver T (2009) Forcing cells to change lineages. *Nature* **462:** 587–594

Guo X, Huang H, Yang Z, Cai T & Xi R (2019) Division of labor: roles of Groucho and CtBP in Notch-mediated lateral inhibition that controls intestinal stem cell differentiation in *Drosophila*. *Stem Cell Reports* **12:** 1–17

Guo Z & Ohlstein B (2015) Bidirectional Notch signaling regulates *Drosophila* intestinal stem cell multipotency. *Science (New York, NY)* **350:** aab0988-1–8

Hartenstein V, Takashima S, Hartenstein P, Asanad S & Asanad K (2017) bHLH proneural genes as cell fate determinants of entero-endocrine cells, an evolutionarily conserved lineage sharing a common root with sensory neurons. *Developmental Biology* **431:** 36–47

Heinze G & Ploner M (2013) Firth's Bias-Reduced Logistic Regression [R package logistf version 1.23]. Available at: https://CRAN.R-project.org/package=logistf

Herrmann C, Van de Sande B, Potier D & Aerts S (2012) i-cisTarget: an integrative genomics method for the prediction of regulatory features and cis-regulatory modules. *Nucleic Acids Res.* **40:** e114–e114

Hinz U, Giebel B & Campos-Ortega JA (1994) The basic-Helix-Loop-Helix domain of *Drosophila* Lethal of Scute protein is sufficient for proneural function and activates neurogenic genes. *Cell* **76:** 77–87

Hu Y, Comjean A, Perkins LA, Perrimon N & Mohr SE (2015) GLAD: an online database of gene list annotation for *Drosophila*. *J. Genomics* **3:** 75–81

Hung R-J, Hu Y, Kirchner R, Li F, Xu C, Comjean A, Tattikota SG, Song WR, Sui SH & Perrimon N (2018) A cell atlas of the adult *Drosophila* midgut. *bioRxiv*: 1–30

Imrichova H, Hulselmans G, Kalender Atak Z, Potier D & Aerts S (2015) i-cisTarget 2015 update: generalized cis-regulatory enrichment analysis in human, mouse and fly. *Nucleic Acids Res.* **43:** W57–W64

Jiang H & Edgar BA (2012) Intestinal stem cell function in *Drosophila* and mice. *Curr. Opin. Genet. Dev.* **24:** 1–7

Jiang H, Patel PH, Kohlmaier A, Grenley MO, McEwen DG & Edgar BA (2009) Cytokine/Jak/Stat signaling mediates regeneration and homeostasis in the *Drosophila* midgut. *Cell* **137:** 1343–1355

Korzelius J, Naumann SK, Loza-Coll MA, Chan JS, Dutta D, Oberheim J, Gläßer C, Southall TD, Brand AH, Jones DL & Edgar BA (2014) Escargot maintains stemness and suppresses differentiation in *Drosophila* intestinal stem cells. *EMBO J* **33:** 2967–2982

Kunisch M, Haenlin M & Campos-Ortega JA (1994) Lateral inhibition mediated by the *Drosophila* neurogenic gene *Delta* is enhanced by proneural proteins. *Proc Natl Acad Sci USA* **91:** 10139–10143

Lan Q, Cao M, Kollipara RK, Rosa JB, Kittler R & Jiang H (2018) FoxA transcription factor Fork head maintains the intestinal stem/progenitor cell identities in *Drosophila*. *Developmental Biology* **433:** 324–343

Lee T & Luo L (1999) Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. *Neuron* **22:** 451–461

Levine M & Davidson EH (2005) Gene regulatory networks for development. *Proc Natl Acad Sci USA* **102:** 4936–4942

Li K & Baker NE (2018) Regulation of the *Drosophila* ID protein Extra macrochaetae by proneural dimerization partners. *eLife* **7:** e33967

Li K & Baker NE (2019) Transcriptional and post-transcriptional regulation of extra macrochaetae during *Drosophila* adult peripheral neurogenesis. *Developmental Biology* **449:** 41–51

Li Y, Pang Z, Huang H, Wang C, Cai T & Xi R (2017) Transcription factor antagonism controls enteroendocrine cell specification from intestinal stem cells. *Sci Rep* **7:** 988

Lowe N, Rees JS, Roote J, Ryder E, Armean IM, Johnson G, Drummond E, Spriggs H, Drummond J, Magbanua JP, Naylor H, Sanson B, Bastock R, Huelsmann S, Trovisco V, Landgraf M, Knowles-Barley S, Armstrong JD, White-Cooper H,

Hansen C, et al (2014) Analysis of the expression patterns, subcellular localisations and interaction partners of *Drosophila* proteins using a pigP protein trap library. *Development* **141:** 3994–4005

Loza-Coll MA, Southall TD, Sandall SL, Brand AH & Jones DL (2014) Regulation of *Drosophila* intestinal stem cell maintenance and differentiation by the transcription factor Escargot. *EMBO J* **33:** 2983–2996

Malaguti M, Migueles RP, Blin G, Lin C-Y & Lowell S (2019) Id1 stabilizes epiblast identity by sensing delays in nodal activation and adjusting the timing of differentiation. *Dev Cell* **50:** 1–22

Massari ME & Murre C (2000) Helix-loop-helix proteins: regulators of transcription in eucaryotic organisms. *Molecular and Cellular Biology* **20:** 429–440

Micchelli CA & Perrimon N (2006) Evidence that stem cells reside in the adult *Drosophila* midgut epithelium. *Nature* **439:** 475–479

Moore AW, Barbel S, Jan LY & Jan YN (2000) A genomewide survey of basic helix-loop-helix factors in *Drosophila*. *Proc Natl Acad Sci USA* **97:** 10436–10441

Moris N, Pina C & Arias AM (2016) Transition states and cell fate decisions in epigenetic landscapes. *Nat. Rev. Genet.* **17:** 693–703

Murre C (2005) Helix-loop-helix proteins and lymphocyte development. *Nat Immunol* **6:** 1079–1086

Murre C, Bain G, Vandijk MA, Engel I, Furnari BA, Massari ME, Matthews JR, Quong MW, Rivera RR & Stuiver MH (1994) Structure and Function of Helix-Loop-Helix Proteins. *Biochim Biophys Acta* **1218:** 129–135

Neuhold LA & Wold B (1993) HLH forced dimers: Tethering MyoD to E47 generates a dominant positive myogenic factor insulated from negative regulation by Id. *Cell* **74:** 1033–1042

Ohlstein B & Spradling A (2006) The adult *Drosophila* posterior midgut is maintained by pluripotent stem cells. *Nature* **439:** 470–474

Ohlstein B & Spradling A (2007) Multipotent *Drosophila* intestinal stem cells specify daughter cell fates by differential notch signaling. *science (New York, NY)* **315:** 988–992

Ohtsuka T & Kageyama R (2010) The Basic Helix-Loop-Helix Transcription Factors in Neural Differentiation. In *Cell Cycle Regulation and Differentiation in Cardiovascular and Neural Systems*, Giordano A & Galderisi U (eds) pp 15–34. New York, NY: Springer New York

Patel PH, Dutta D & Edgar BA (2015) Niche appropriation by *Drosophila* intestinal stem cell tumours. *Nature Cell Biology* **17:** 1182–1192

Philpott A & Winton DJ (2014) Lineage selection and plasticity in the intestinal crypt. *Curr Opin Cell Biol* **31:** 39–45

R Core Team (2018). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available online at https://www.R-project.org/

Resende LPF, Truong ME, Gomez A & Jones DL (2017) Intestinal stem cell ablation reveals differential requirements for survival in response to chemical challenge. *Developmental Biology* **424:** 10–17

Sallee MD, Littleford HE & Greenwald I (2017) A bHLH code for sexually dimorphic form and function of the *C elegans* somatic gonad. *Curr Biol* **27:** 1853–1860.e5

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P & Cardona A (2012) Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9:** 676–682

Shokri L, Inukai S, Hafner A, Weinand K, Hens K, Vedenko A, Gisselbrecht SS, Dainese R, Bischof J, Furger E, Feuz J-D, Basler K, Deplancke B & Bulyk ML (2019) AcComprehensive *Drosophila* melanogaster transcription factor interactome. *Cell Reports* **27:** 955–970.e7

Simon CS, Hadjantonakis A-K & Schröter C (2018) Making lineage decisions with biological noise: Lessons from the early mouse embryo. *WIREs Dev Biol* **7:** e319–16

Siudeja K, Nassari S, Gervais L, Skorski P, Lameiras S, Stolfa D, Zande M, Bernard V, Frio TR & Bardin AJ (2015) Frequent somatic mutation in adult intestinal stem cells drives neoplasia and genetic mosaicism during aging. *Cell Stem Cell* **17:** 1–12

Spratford CM & Kumar JP (2015) Inhibition of Daughterless by Extramacrochaetae mediates Notch-induced cell proliferation. *Development* **142:** 2058–2068

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES & Mesirov JP (2005) Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA* **102:** 15545–15550

Tanaka-Matakatsu M, Miller J, Borger D, Tang WJ, Du W (2014) Daughterless homodimer synergizes with Eyeless to induce Atonal expression and retinal neuron differentiation. *Developmental Biology* **392:** 256–265

Tapanes-Castillo A & Baylies MK (2004) Notch signaling patterns *Drosophila* mesodermal segments by regulating the bHLH transcription factor twist. *Development* **131:** 2359–2372

Tetteh PW, Basak O, Farin HF, Wiebrands K, Kretzschmar K, Begthel H, van den Born M, Korving J, de Sauvage F, van Es JH, van Oudenaarden A & Clevers H (2016) Replacement of lost Lgr5-positive stem cells through plasticity of their enterocyte-lineage daughters. *Cell Stem Cell* **18:** 203–213

Troost T, Schneider M, Klein T (2015) A Re-examination of the Selection of the Sensory Organ Precursor of the Bristle Sensilla of *Drosophila melanogaster. PLoS Genet* **11:** e1004911–21

van der Flier LG, van Gijn ME, Hatzis P, Kujala P, Haegebarth A, Stange DE, Begthel H, van den Born M, Guryev V, Oving I, van Es JH, Barker N, Peters PJ, van de Wetering M & Clevers H (2009) Transcription factor achaete scute-like 2 controls intestinal stem cell fate. *Cell* **136:** 903–912

Van Doren M, Ellis HM & Posakony JW (1991) The Drosophila extramacrochaetae protein antagonizes sequence-specific DNA binding by daughterless/achaete-scute protein complexes. *Development* **113:** 245–255

Wang C, Guo X, Dou K, Chen H & Xi R (2015) Ttk69 acts as a master repressor of enteroendocrine cell specification in *Drosophila* intestinal stem cell lineages. *Development* **142:** 3321–3331

Wice BM (1998) Forced expression of Id-1 in the adult mouse small intestinal epithelium is associated with development of adenomas. *Journal of Biological Chemistry* **273:** 25310–25319

Yin C & Xi R (2018) A Phyllopod-mediated feedback loop promotes intestinal stem cell enteroendocrine commitment in *Drosophila*. *Stem Cell Reports* **10:** 43–57

Zeng X & Hou SX (2015) Enteroendocrine cells are generated from stem cells through a distinct progenitor in the adult *Drosophila* posterior midgut. *Development* **142:** 644–653

Zeng X, Chauhan C & Hou SX (2010) Characterization of midgut stem cell- and enteroblast-specific Gal4 lines in *Drosophila*. *Genesis* **48:** 607–611

Zeng X, Han L, Singh SR, Liu H, Neumüller RA, Yan D, Hu Y, Liu Y, Liu W, Lin X & Hou SX (2015) Genome-wide RNAi screen identifies networks involved in intestinal stem cell regulation in *Drosophila*. *Cell Reports* **10:** 1226–1238

Zhai Z, Kondo S, Ha N, Boquete J-P, Brunner M, Ueda R & Lemaitre B (2015) Accumulation of differentiating intestinal stem cell progenies drives tumorigenesis. *Nat Commun* **6:** 10219–27

Zhang N, Yantiss RK, Nam H-S, Chin Y, Zhou XK, Scherl EJ, Bosworth BP, Subbaramaiah K, Dannenberg AJ & Benezra R (2014) ID1 is a functional marker for intestinal stem and progenitor cells required for normal response to injury. *Stem Cell Reports* **3:** 716–724

F

Figure 7