The origin of the Gravettians: genomic evidence from a 36,000-year-old Eastern

European

E. Andrew Bennett[1], Sandrine Prat[2], Stéphane Péan[3], Laurent Crépin[3], Alexandr Yanevich[4],

Simon Puaud[2], Thierry Grange[1], Eva-Maria Geigl[1]

[1]Institut Jacques Monod, CNRS, Université de Paris, 75013 Paris, France; [2]UMR 7194 (HNHP),

MNHN/CNRS/UPVD, Alliance Sorbonne Université, Musée de l'Homme, Palais de Chaillot, 17

Place du Trocadéro, 75116, Paris, France; [3]UMR 7194 (HNHP), MNHN/CNRS/UPVD, Muséum

national d'Histoire naturelle, Alliance Sorbonne Université, Institut de Paléontologie Humaine, 1

rue René Panhard, 75013, Paris, France; [4]Institute of Archaeology, National Academy of Sciences

of Ukraine, Heroiv Stalingrada 12, 04210 Kyiv, Ukraine.

The Gravettian technocomplex was present in Europe from more than 30,000 years ago until

the Last Glacial Maximum, but the source of this industry and the people who manufactured

it remain unsettled. We use genome-wide analysis of a ~36,000-year-old Eastern European

individual (BuranKaya3A) from Buran-Kaya III in Crimea, the earliest documented

occurrence of the Gravettian, to investigate relationships between population structures of

Upper Palaeolithic Europe and the origin and spread of the culture. We show BuranKaya3A

to be genetically close to both contemporary occupants of the Eastern European plain and

the producers of the classical Gravettian of Central Europe 6,000 years later. These results

support an Eastern European origin of an Early Gravettian industry practiced by members

of a distinct population, who contributed ancestry to individuals from much later Gravettian

sites to the west.

Since the 19th century, archaeologists have defined similarities identified across material remains

as an archaeological culture, complex, or in some cases, a "people", and used these definitions to

trace the mobility, interactions, and technical development of past populations. The burgeoning

field of ancient DNA analysis allows past populations to be studied directly, and the genetic

relationships of the manufacturers of materially defined cultures can now be characterized. While

palaeogenomics and archaeological culture describe two separate past phenomena, they are not

always unrelated, and Upper Palaeolithic (UP) Europe has shown surprising correlations between

closely-related genetic clusters and archeologically defined material industries[1]. The Gravettian

technocomplex defines European Mid-Upper Palaeolithic (MUP) industries characterized by a

suite of shared innovations, along with specific modalities of their production, such as stone

Gravette points, backed blades and bladelets, personal ivory and shell ornaments, ochre, and antler

or bone tools. The Gravettian became widespread throughout Europe beginning after ca. 36,000[2]

until ca. 23,000[3] cal BP years ago (ca. 32-21 ka [14]C BP). The technocomplex defined by this term,

however, is by no means homogeneous, describing rather shared practices across many regional

facies and evolving *stages*[3] (See Supplementary Text). While the Gravettian is considered to be a

local European industry, its origins continue to be the subject of ongoing research and debate[4]. The

density of Gravettian sites found in the Danubian valley radiocarbon dated as early as ca. 36,000

years cal BP have been used to argue for a local evolution of the Gravettian from the technical

legacy of the Aurignacian in the upper valley[5], or through acculturation with post-Mousterian leaf-

point transition industries, such as the Szeletian, in the Middle Danubian basin[6]. However, long-

recognized similarities between features of Gravettian lithic traditions and Near Eastern industries,

such as the Ahmarian, which is found 10,000 years earlier in the Levant, as well as similar Early-

Upper Palaeolithic (EUP) micro-laminar industries found in both northern and southern slopes of

the Caucasus suggest an earlier influence from the southeast[7–9].

Although the integration of Eastern European EUP and MUP traditions into an archaeological

framework defined in Central and Western Europe has historically been problematic, recent

reassessments of EUP assemblages from the Caucasus[10–12] emphasize similar lithic traditions

among roughly contemporaneous (40-30,000 cal BP) layers at sites such as Dzudzuana Cave (layer

D), Ortvale Klde (layer 4c), Mezmaiskaya Cave (layer 1C), and the six layers of Buran Kaya III

(layer 6-5 to 5-2) in Crimea[13,14] (see Supplementary Text for the stratigraphy of Buran-Kaya III).

Some authors suppose that the appearance of these Caucasian and Eastern European EUP

assemblages, which include backed blades and bladelets, are primarily based on the distribution

and transformation of Ahmarian traditions from Near East to Europe, probably through the

Caucasus, independent of that which brought the proto-Aurignacian to Central and Southern

Europe and the Mediterranean area[10,14–16]. The parallels between these Eastern European industries

and the Gravettian appearing later in Central Europe and the Danubian Valley have led some to

propose the term "Early Gravettian" to describe these industries to distinguish them from the

classical Gravettian[8,16,17]. However, the relationship of this Early Gravettian identified in Eastern

Europe to the development of the Gravettian in the west, as well as the extent to which it may have

involved transfers and adaptations of technology and movements of people, are unclear. The direct

study of the populations associated with these industries through genomic analysis of skeletal

remains from key archaeological sites will likely clarify this process and answer important

questions concerning how technologies disperse and change in UP societies.

Several human remains associated with the classical Gravettian from Central Europe, Belgium and

Italy were previously analysed genetically and found to share more genetic drift with each other

than individuals associated with other material cultures[1]. This group was termed the *Vestonice*

cluster, after Vestonice16, the best-covered genome from the ca. 30,000-year-old (cal BP)

Gravettian site Dolní Věstonice II in the Czech Republic. Interestingly, Vestonice16 and other

high-coverage genomes from the Gravettian sites Krems-Wachtberg (Austria) and Ostuni (Italy)

have been found to be admixed, deriving between 40 to 90% of their ancestry from a lineage

related to Kostenki14[1,18], an individual living ca. 7,000 years earlier at the Kostenki-14 site found

in the Borshchyovo archaeological complex of Western Russia, linking Gravettians of Central

Europe and the Apennine peninsula with an earlier population farther east.

While the Kostenki14 burial has no associated cultural material, contemporaneous Gravettian

artefacts have been recovered in Eastern Europe from three cultural layers at Buran-Kaya III in

Crimea and dated between ca. 38-34,000 cal BP[2,19]. Discovered in 1990, the Buran-Kaya III rock

shelter contains stratigraphic layers spanning from the Middle Palaeolithic to the Middle Ages.

The Gravettian layers have yielded backed microliths, microgravette points, ochre, body

ornaments of ivory, shell, and teeth, as well as multiple human skull fragments showing signs of

post-mortem processing[2]. Buran-Kaya III is the earliest site bearing Gravettian material[20] (see

Supplementary Text), yet it remains a distant outlier, both geographically and temporally, to the

more densely clustered appearances of the Gravettian occurring several thousand years later and

2,000 km to the west in the Danubian valley and the Swabian Jura, such as Krems-Hundssteig and

Geiβenklösterle, respectively.

To genetically characterize one of the earliest manufacturers of the Gravettian complex and how

they relate to other known MUP populations, as well as to further explore the relationships between

UP populations and the material culture they made, we present genome-wide data from a human

parietal bone fragment, BuranKaya3A, aseptically excavated in 2009 from layer 6-1 of Buran-

Kaya III (see Supplementary Text). Radiocarbon (AMS) dates from a different human cranial

fragment from this layer range from 36,260 to 35,280 cal BP (31,900+210-220 ^{14}C BP)[2] (Figure 1

and Table S1), which is located between two other Gravettian layers, while all dated material (n=4)

from the same layer ranges from 37,560-33,850 cal BP[19].

Results

Initial genetic characterization of BuranKaya3A revealed extremely poor preservation of

endogenous DNA, both in fragment length and in low quantity relative to environmental DNA,

but little contamination with modern human DNA as assessed by quantitative PCR. In order to

identify the most efficient material for mitochondrial enrichment and shotgun genome sequencing,

our strategy was to 1) screen single-stranded DNA libraries by both shallow shotgun sequencing

and mitochondrial enrichment from eight extractions taken from four different areas of the sample

and subjected to two different extraction treatments, 2) identify the extract that performed best in

terms of highest sequence complexity, endogenous content, and lowest modern human

contamination, and 3) construct a second series of single-stranded libraries with and without UNG

for deeper mitochondrial enrichment and shotgun sequencing. The best candidate identified from

the screening results contained 0.34% endogenous DNA (see *Methods* section and Extended Data

Figure 1). The second series of libraries from this extract was used to generate 82-fold coverage

of the mitochondrial genome and shotgun nuclear data (Table S2).

Mitochondrial and Y-chromosome haplogroups

The mitochondrial haplogroup of BuranKaya3A was determined to belong to an early branch of the N lineage, N1. Surprisingly, this assignment falls outside of the lineages previously reported for UP Europe, nearly all of which derive from later N branches (U and R haplogroups, Figure 2). The N1 of BuranKaya3A is notably distinct from the mitochondrial haplogroup N identified from the roughly 40,000-year-old mandible from Peștera cu Oase in Romania, which belongs to a more basal branch that has no modern descendants[21]. In addition, the N1 of BuranKaya3A carries three of the eight mutations occurring prior to N1b, a rare haplogroup most highly concentrated in the Near East, yet appearing broadly from western Eurasia to Africa. The descendants of the N1b node include N1b2, currently found only in Somalia[22], and N1b1b, found in nearly 10% of Ashkenazi Jewish haplogroups[23]. These three mutations allow us to place BuranKaya3A on a lineage apart from that which has been proposed to later enter Europe from Anatolia during the Neolithic (N1a1a)[24]. Among ancient samples, the mitochondrial sequence of an 11,000-year-old Epipalaeolithic Natufian from the Levant ("Natufian9")[25] is also a later derivative of this N1b branch. Thus, mitochondrial sequences branching both upstream and downstream of the BuranKaya3A sequence can be traced to the Near East, and the modern presence in Europe of haplogroups descended from the N1 (N1b1b and N1a1a) branch to which BuranKaya3A belongs appear to be due to later migrations from the Near East (Extended Data Figure 2). We determined the genetic sex of BuranKaya3A to be male using both the ratio of chromosome X and Y mapped reads, giving an R_y value of 0.0893-0.097 (95% CI, SE 0.002)[26], as well as a ratio of chromosome X mapped reads to the average of autosomal reads of 0.55 (a ratio near 1.0 would indicate diploid for X). From the reads mapping to the Y chromosome, six out of six Single Nucleotide Polymorphisms (SNPs) that overlap with diagnostic sites for Y-haplogroup BT all carry the

derived allele, allowing a minimum assignment to BT, which has origins in Africa, with additional

derived alleles suggesting an eventual placement of CT or C, found in Asia and the Epipalaeolithic

Near East[25]. Additional ancestral alleles make an assignment of C1a2 or C1b, which appear in UP

Europe[1], unlikely (see Table S3 for a summary and comparative placement of Palaeolithic Y-

haplogroups, and Supplementary Data 1 for a complete list of Y diagnostic SNPs).

Neanderthal ancestry

Neanderthal settlements, attributed to the Micoquian, Kiik-Koba type, at Buran-Kaya III (layer B)

has been dated by faunal bone fragments to 43.5 to 39.6 ka cal BP[19], demonstrating an earlier

Neanderthal presence at the site just prior to the Campanian Ignimbrite eruption (39,280±110 years

cal BP (^{40}Ar/^{39}Ar))[27]. Anatomically Modern Human (AMH) remains dating to this period in

Romania have documented local admixture with late Neanderthals in Eastern Europe[21], leading us

to investigate whether admixture with local Neanderthals in Crimea could be detected in early

AMHs living ca. 4,000 years after the presumed disappearance of Neanderthals from the

region[19,28]. Neanderthal ancestry calculated using ancestry informative SNPs[1] on libraries from

four independent preparations determined that BuranKaya3A possesses 3.4% (SD 0.008)

Neanderthal ancestry (Table S4). This level of Neanderthal ancestry is typical for

contemporaneous West Eurasians and shows no evidence of late, local Neanderthal admixture in

Crimea with AMHs ancestral to the population to which BuranKaya3A belonged.

Genomic relationships of UP Europe and their archaeological cultures

The genetic relationship between BuranKaya3A and other UP individuals for which sufficient

genomic data exist can be measured using outgroup $f3$-statistics and various D-statistical tests[29].

These were performed using SNPs from 52 previously published individuals dating from the Initial

Upper Palaeolithic (IUP) to the Mesolithic that had been reprocessed in the same pipeline used for

BuranKaya3A to reduce possible artefacts that may arise from data coming from disparate sources.

27,774 SNPs having base quality 30 or above were identified in BuranKaya3A that overlapped

with the nearly 3M SNPs from the combined panels in Fu *et al.* 2016[1]. To minimize standard

errors, we restricted the results of this analysis to include only individuals having more than

300,000 SNPs overlapping with the panel.

Outgroup $f3$-statistics quantify the amount of shared genetic drift between two populations relative

to an outgroup population and have been used to reveal genetic affinities among UP individuals[1].

Of the 21 ancient individuals (x) including at least 1000 SNPs in the calculation: $f3$(BuranKaya3A,

x; Mbuti) or $f3$(BuranKaya3A, x; Han), the highest $f3$ values for both outgroups were found when

x was either Sunghir3 (SIII), a 34,000-year-old (cal BP) group burial characterized as either

Streletskian[30] or Eastern Gravettian[31,32] (see Supplementary Text) found ca. 1,300 km to the

northeast of Buran-Kaya III, Vestonice16, from the context of a Gravettian *facies* known as the

Pavlovian, who lived ca. 6,000 years later and ca. 1,400 km to the northwest, and Kostenki14, a

burial predating BuranKaya3A by ca. 1,000 years and lying ca. 1,100 km to the northeast (Figure

3). Notably, BuranKaya3A shows less affinity for the 35,000-year-old (cal BP) GoyetQ116-1 from

Goyet cave in Belgium, with no direct cultural association, but dated to the Aurignacian period in

this area[1,33], as well as the El Miron cluster, which corresponds to European late glacial

(Magdalenian) hunter-gatherers[1]. BuranKaya3A was also found to be particularly distant from the

Villabruna cluster, representing European post-glacial hunter-gatherers (Figure 3 and Extended

Data Figure 3A-B). The individual from Buran-Kaya III can thus be defined as a member of a

population making an Early Gravettian in Eastern Europe whose closest known genetic

relationships are with populations living within a 3,000-year window (ca. 37,000-34,000 cal BP)

farther northeast on the Eastern European plains (Sunghir3 and Kostenki14), as well as with

Gravettian populations who appeared ca. 6,000 years later in Central Europe (Vestonice16).

We then applied D-statistics to measure shared drift between BuranKaya3A and the individuals

tested above (x) as compared to the 45,000-year-old (cal BP) Central Asian Ust-Ishim ($D(x$, Ust-

Ishim; BuranKaya3A, Mbuti)), shown to be basal to all western MUP Eurasians studied to date[34]

(Figure 4a). Substituting the modern Han for Ust-Ishim gives additional support for the above

relationships with greater significance (Z-scores > 2.5) (Figure 4b). Full results for all tested

samples using both all SNPs and transversions only are given in Extended Data Figures 4 and 5.

The D-statistics support the results of the outgroup $f3$-statistical tests, except that here Sunghir3 is

replaced by Vestonice16 as sharing the most alleles with BuranKaya3A, although we note results

from both Vestonice16 and Sunghir3 overlap within one standard error. The additional D-statistic

$D(w$, x; BuranKaya3A, Mbuti), where w and x are various well-covered ancient individuals

representing previously defined Eurasian populations, was unable to significantly resolve which

of these two individuals is most closely related genetically to BuranKaya3A: the ca. 34,000-year-

old (cal BP) East European Sunghir3 or the ca. 30,000-year-old (cal BP) Central European

Vestonice16 (Extended Data Figure 6).

Genomic data from less well covered MUP Europeans allows us to better define the relative

position of BuranKaya3A with more individuals from Gravettian contexts, including

representatives of two of the three previously identified *Vestonice* genetic sub-clusters

corresponding to Gravettian sites[1]. Comparing two low-coverage individuals requires more

caution in interpreting the results due to the low number of SNPs used for the calculations.

However, we were able to establish additional support for the relationships by applying a strict

filtering process whereby 1) we performed the analysis using two datasets, one including all SNPs

and the other considering only the transversions. The transversion-only dataset reduces the number

of SNPs used, but eliminates statistical noise from deaminated cytosines, a common form of

damage in ancient DNA. 2) We excluded samples using less than 50 SNPs for f3 analysis in the

smaller transversion-only dataset. 3) We required the resulting values from both datasets to be

within 30% of each other. This approach allowed us to define genetic affinities between

BuranKaya3A and the following individuals: Kostenki12, a 32,500-year-old (cal BP) neonate

whose remains were found at the Kostenki-Borshchyovo archaeological complex in Russia and

associated with a cultural layer attributed to a local industry defined as the Gorodtsovian[35];

Vestonice43 and Pavlov1, two Central European Gravettian (Pavlovian) burials similar in age and

location to Vestonice16 and belonging to the "Vestonice Central European" genetic subcluster;

Paglicci133, a ca. 33,000-year-old (cal BP) tooth from the Apulia region of Southern Italy found

in association with Early Gravettian cultural material; and Ostuni1, a ca. 27,500 year-old (cal BP)

Gravettian burial, also from the Apulia region and both belonging to the "Vestonice Italian"

genetic subcluster. No members of the "Vestonice Goyet" subcluster from Goyet cave in Belgium

passed our filters to be included in the analysis (full descriptions of these samples and clusters are

given in the Supplementary Information of Fu et al. 2015[1]). Visual representation of the outgroup

f3-statistics, f3(BuranKaya3A, *MUP*; Mbuti), for all samples passing filters is shown in Extended

Data Figure 7 and full results for all tested samples are given in Extended Data Figure 8.

Of the less well-covered samples, Kostenki12 is shown to have a high affinity with BuranKaya3A.

This is in agreement with the previously reported affinity between Kostenki12 and Sunghir3[36] and

closely links BuranKaya3A to all EUP and MUP individuals currently known from the Eastern

European plains. To the west, BuranKaya3A shows close genetic affinity with Pavlov1, followed

more distantly by Vestonice43 and Ostuni1, and at a greater distance, the older Paglicci133. In

summary, these results allow us to firmly establish a close genetic relationship between

BuranKaya3A with contemporary EUP and MUP populations of the Eastern European plains, as

well as the broader population manufacturing the Pavlovian *facies* of the Gravettian in Central

Europe ca. 6,000 years later, but less so with those associated with the Gravettian of southern Italy.

This implies a network of gene flow across the Eastern European plains and the Danubian Valley

between 37,000-30,000 years ago (cal BP) which, apart from the local Gorodtsovian culture found

at Kostenki 12, can be associated with various *facies* of the Gravettian technocomplex. The results

from Ostuni1 and Paglicci133 indicate a more indirect relationship between these populations and

BuranKaya3A. Higher coverage genetic information from early Gravettian sites in Italy will be

required to better understand to what extent the movements of people and ideas lead to the

appearance of the Gravettian in the southern Apennine Peninsula.

Low Common West Eurasian ancestry in UP Eastern Europe

Ancestry from a population which split from all non-Africans prior to their separation from each

other, termed Basal Eurasian, had not been known in Europe until after the Last Glacial Maximum

(LGM)[1]. However, two 24,000-27,000-year-old (cal BP) individuals from layer C at Dzudzuana

Cave in the southern Caucasus have recently been reported to share ~70% common ancestry with

Villabruna and ~30% ancestry deriving from this Basal-Eurasian source[18]. Additionally, varying

layers of Villabruna ancestry, which did not enter Europe in an unmixed form prior to ca. 14,000

years ago (cal BP)[1], have been found in members of the Gravettian Vestonice cluster as well as

the Magdalenian El Miron cluster, indicating a degree of shared ancestry in these groups with a

"Common West Eurasian" population[18]. We calculated the relative level of Basal Eurasian

ancestry using D-statistic $D(EUP\ East\ Asian, UP, \text{Ust-Ishim}, \text{Mbuti})$, where a positive value

indicates Basal Eurasian ancestry as allele sharing between the UP individual and Africans. This

analysis showed BuranKaya3A, like other pre-LGM UP Europeans, to be lacking Basal Eurasian

ancestry (Extended Data Figure 9). Similar tests examine the relative affinities of Palaeolithic and

Mesolithic populations to Villabruna as opposed to either modern East Asians, $D(x, \text{Han};$

Villabruna, Mbuti), or BuranKaya3A, $D(x, \text{BuranKaya3A}; \text{Villabruna}, \text{Mbuti})$ (Extended Data

Figures 10 and 11, respectively). These results show levels of Villabruna ancestry in Eastern

European EUP individuals (Sunghir3, Kostenki12, Kostenki14, and BuranKaya3A) below that of

later UP Central and Western Europeans from Gravettian contexts (the Vestonice cluster).

Intriguingly, this includes a proportion of shared ancestry with Villabruna in BuranKaya3A similar

to that found in Ust-Ishim and both ancient and modern East Asians (Extended Data Figures 10

and 11), which should be insignificant based on previous admixturegraph analysis[18].

Discussion

The results of the genome-wide analysis of BuranKaya3A offer important evidence linking the

previously established genetic signature of the manufacturers of the Gravettian in Central Europe

to a much earlier appearance of the Gravettian in Eastern Europe. The absence of the "Common

West Eurasian" ancestry, as represented by Villabruna, in BuranKaya3A marks a key genetic

distinction between the Gravettian inhabitants of Buran-Kaya III, possibly including the broader

populations of EUP Eastern Europe as well, and the UP populations of Western and Central

Europe, which is characterized by a West-to-East reduction in "Common West Eurasian" ancestry

(seen in Extended Data Figure 10). The association we show here of this eastern genetic character

with the cultural material of the Gravettian of Buran-Kaya III, which has been compared to nearby

contemporaneous Early Upper Palaeolithic assemblages from the Caucasus[11,15] indicated in Figure

3, collectively support an eastern advance of AMHs during the EUP into Europe through the

Caucasus as has been previously proposed based on archaeological evidence alone[7,8,16]. Such a

population would have had to have split from the settlers of Central Europe and the Mediterranean

prior to their acquisition of the Common West Eurasian component as represented by Villabruna.

In this scenario, the technical adaptations required for the challenging environment of the open

Eastern European plains, a dryer landscape with little natural shelter, as well as possible cultural

exchanges with local populations, may have played a role in the development of the Early

Gravettian industry[8]. The individuals recently characterized genetically from layer C of

Dzudzuana cave in the Caucasus (data not yet available), who were found to contain ancestry (both

Basal Eurasian and Common West Eurasian) that was absent ca. 9,000 years earlier in Crimea,

may represent more recent immigration into Eastern Europe. A higher resolution of these

movements awaits genetic analysis of more EUP and MUP sites from this region.

Numerous parallels in lithic industries, such as microblade-knapping methods, backed blades, and

analogous stone blade, point and tool morphology (such as the partly backed Ahmarian el Wad-

points and Gravette points of Europe) have suggested earlier Near Eastern cultures as possible

precursors to Gravettian techniques[7,8]. Despite both uni-parental markers being shared between

BuranKaya3A and Epipalaeolithic Natufians in the Near East, we were unable to detect extensive

genome-wide allele-sharing between BuranKaya3A and the Natufians in our analysis. We note,

however, that the high level of Basal Eurasian ancestry in Natufian genomic sequences (38-54%[25])

limits our sensitivity when comparing populations lacking this component. For example, negative

D-statistics seen in Figure 4 involving the Mesolithic Caucasus Hunter-Gatherer (CHG)

individuals Kotias and Satsurblia[37], as well as the Natufians and the Iranian Hotu[25], are due to the

Basal Eurasian content reported for these individuals, which appears in the statistic as an affinity

toward the outgroup Mbuti. Also, given the more than 20,000-year age difference between

BuranKaya3A and the Natufians, it is unlikely that the Epipalaeolithic Natufians are the best

surrogates for EUP Near Eastern populations. While influences from the Ahmarian can be seen in

the Gravettian[17], it is the Early Ahmarian that has separately been proposed as a source for the

wave of AMHs bringing the Proto-Aurignacian west, possibly by way of the Balkans, into Central

and Southern Europe beginning prior to the Campanian Ignimbrite eruption[13]. Populations

associated with the Early Ahmarian and the Ahmarian may thus suggest better candidate source

populations to investigate further the hypothesis of a population split behind separate routes in the

settling of Europe, the Balkans/Mediterranean to the west and the Caucasus to the east, each

associated with accompanying industries (Proto-Aurignacian and Early Gravettian, respectively).

The shared ancestry, as well as cultural similarities, demonstrated between the settlements at

Buran-Kaya III, later Gravettians in Central Europe, and to some extent, Sunghir, suggest a broad

and long-lasting network of social exchange in the EUP across Eastern and Central Europe, from

the Eastern European plains to the Danubian corridor. Given this background, the appearance of

the Gravettian in Central Europe in the MUP, where it later would blossom, is likely to have

resulted from this input from the east. While we show that the Gravettian, *sensu lato*, was not

practiced by a single genetically uniform cluster across all *facies*, the close genetic relationship

between BuranKaya3A and the Kostenki individuals raise further questions as to the origin of the

local culture termed Gorodtsovian, which is found at Kostenki 12 and unknown outside the

Kostenki-Borshchevo region. Given the genetic affinities we report, the previous assignment by

some authors of the Late Streletskian industry of Sunghir being a local *facies* of an "Eastern"

Gravettian[31,32] should lead to a closer examination of the possible influences underlying the

appearance of the Gorodtsovian, with appreciation for the impact that climate, the specific

landscape of the site, and site-specific activities may have had on the individual tool requirements

of the assemblages[8]. Alternatively, the Gorodtsovian and other local UP industries of Eastern

Europe may represent distinct cultures practiced on the Eastern European plains, and the genomic

affinities between the individuals from Kostenki, Sunghir and BuranKaya3A may show only a

relationship to a common source population among the occupants of Eastern Europe branching

more recently than those present in Western Europe.

This study, the genomic analysis of the oldest AMH from an archeologically defined context,

demonstrates an underlying genetic continuity between manufacturers of various *facies* of the

Gravettian spanning ca. 9,000 years. A geographical divergence among groups entering Europe

more than 37,000 years ago is supported by the finding that the earliest appearance at Buran-Kaya

III is associated with a population unadmixed with the Common West Eurasian component already

present in Europe to some degree, and is thus distinct. Regional features such as micro-laminar

industries and Gravette points found in EUP assemblages in the Caucasus, both archeologically

comparable and contemporary with Buran-Kaya III, suggest the Caucasus as a possible route for

this diffusion, and a role of these industries in the development of the Early Gravettian. A more

comprehensive understanding of both genomic information and archaeological assemblages of UP

sites in the Caucasus and Near East will allow more precise identification of the origins of both

this population and, potentially, the Gravettian.

Methods

Dating

All radiocarbon dates were recalibrated using the software OxCal v4.3.2 based on the IntCal13 calibration data set[38]. The calibrated dates are rounded to 5.

Sample handling, DNA extraction, library construction, and sequencing

A human parietal fragment was excavated aseptically from layer 6-1 during the 2009 excavation season at Buran Kaya III (see Supplementary Text). All pre-amplification sample preparation was performed in the dedicated ancient DNA facility using decontamination and clean-room protocols as described in Bennett, et al[39]. All buffers and solutions were prepared using water decontaminated by gamma-irradiation (8 kGy). After first removing the surface of the areas to be sampled with a sterile scalpel, between 47 and 114 mg of bone powder was recovered from four different places of the bone using a variable-speed drill at low speed to reduce overheating (Dremel, Mount Prospect, IL, USA). Two of these samplings were each divided into two equal portions, one of which was subject to phosphate buffer pre-treatment as described in Korlevic, *et al*[40]. Phosphate buffer washes for each sample were collected and combined for DNA purification. Both phosphate and non-phosphate buffer treated samples, including reagent-only mocks, were then incubated in 1.5 mL LoBind microcentrifuge tubes (Eppendorf, Hamburg, Germany) with 1 mL 0.5 M EDTA , pH 8.0 (Sigma-Aldrich, St. Louis, MO), with 0.25mg/mL proteinase K (Sigma-Aldrich) and 0.05% UV-irradiated Tween-20 (Sigma-Aldrich), at 37°C for 24 H. Following incubation, all tubes were centrifuged at maximum speed for 10 min, and supernatant was mixed with 10 times its volume of "2M70" binding buffer (2 M guanidine hydrochloride and 70%

isopropanol) in a 15 mL tube and passed through QIAquick silica columns (Qiagen, Hilden,

Germany) using 25 mL tube extenders (Qiagen) and a vacuum manifold (Qiagen) as described[39,41].

2M70 binding buffer has been shown to retain the smaller DNA fragments lost during purification

with traditional binding buffers[42]. Columns were washed twice with 1 mL PE Buffer (Qiagen) then

transferred to a micro-centrifuge and dried by spinning 1 minute at 16,100 × g, turning tubes 180°

and repeating. DNA was eluted in a total of 60 μl of 10 mM EBT (Tris-HCl pH 8.0 containing

0.05% Tween-20) performed in two elutions of 30 μl each, by spinning 16,100 × g for 1 minute

after a 5-minute incubation.

For the screening step, single-stranded libraries were constructed using either 2 μl (for screening)

or 6 μl (for mitochondrial capture) of the eluted DNA, including mocks of all treatments, water

only samples, and a positive control oligo following the protocol of Gansauge, *et al.*[43] using the

splinter oligonucleotide TL110, and eluting in 50 μl EBT. Either 40 μl (for mitochondrial capture)

or 4 μl (for screening) of each library was used for bar-coding amplification using dual-barcoded

single-stranded library adapters[44] as primers in the following 100 μl volume reaction: 10 μl 10x

PCR Buffer + MgCl2 (Roche, Basel, Switzerland) 0.4 μM of each primer, 80 μM dNTPs (Roche),

15 units of FastStart Taq (Roche). Reactions were heated 95°C for 5 min, followed by 35 cycles

of 95°C 20 s, 53°C for 45 s, 68°C for 45 s, and then 68°C for 5 min. Heteroduplexes that could

confound size selection were resolved by diluting the PCR product 1:5 in a 100 μl reaction

containing 20 μl of the initial reaction, 8 μl PCR Buffer + MgCl2, 0.4 μM of standard Illumina

primers P5 and P7, and 80 μM dNTPs, and amplified a single cycle of 95°C for 1 minute, 60°C

for 2 min and 68°C for 5 min. Products were then purified and size-selected using NucleoMag

beads (Macherey-Nagel, Düren, Germany) for two rounds of purification/size selection according

to the supplied protocol at a ratio of bead solution 1.3 times the reaction volume and eluted in 30

µl EBT. Purified libraries were quantified using a Nanodrop ND-1000 spectrophotometer (Thermo

Fisher Scientific, Waltham, Massachusetts, USA), Bioanalyzer2100 (Agilent, Santa Clara,

California, USA), Qubit 2.0 Fluorometer (Thermo Fisher Scientific), and qPCR reaction. 46 to 148

ng of DNA were enriched for human mitochondrial sequence in two rounds of capture using 1200

ng of biotinylated RNA baits reverse transcribed from human mitochondrial PCR products

(courtesy of L. Cardin and S. Brunel.) following the protocol described in Massilani *et al.*[45], but

with four changes: (1) for hybridization and wash steps, 60°C was used instead of 62°C. (2)

DNA/RNA-bait solution was incubated 96 H instead of 48 H. (3) Elution of the bead-bound

enriched DNA was performed with a 5 minute incubation in 30 µl EBT at 95°C followed by a

magnetic bead separation and the transfer of the eluate to a new tube rather than a 0.1 N NaOH

elution followed by silica column purification. (4) All post-capture amplifications were performed

for 35 cycles followed by a heteroduplex resolution step as described above. Enriched DNA was

then quantified as above, and products from all libraries were pooled in equimolar amounts and

sequenced on an Illumina MiSeq using a v3 reagent kit for 2x76 cycles, substituting primer CL72

for the Read1 sequencing primer as described[44].

Eight additional libraries along with positive and negative controls were made from 3-8 µl each of

the remaining extract BK_A4B, which had the highest relative endogenous DNA content. It should

be noted that this extract was derived from the portion of the cranial fragment which included the

suture. Libraries were prepared as described above except one library was first treated with USER

enzyme (New England Biolabs, Ipswich, Massachusetts, USA) for 30 min to remove deaminated

cytosine damage. Prior to the barcoding amplification, a 6-cycle amplification of pre-barcoded

libraries was performed using 45 µl of each library with 45 µl OneTaq 2X Master Mix with

Standard Buffer (New England Biolabs), and 0.1 µM internal primers CL72[44] and CL130[40] with

the above PCR conditions. Three pairs of different dual-barcoded adapters were then used to

amplify 20 μl of each amplified library, followed by purification and size selection as described

above, which allowed the later pooling of two enrichment protocols and non-enriched DNA from

the same library. An average of 1.2 μg of DNA to 1 μg RNA-baits was used for each mitochondrial

enrichment, as above, however, an additional alternative "touch-down" hybridization protocol

consisting of 60°C for 12 H, 59°C for 12 H, 58°C for 12 H, 57°C for 12 H, and 56°C for 48 H was

tested for each library. For select libraries, an alternative wash protocol described in Fu *et al.*[46] was

also tested. Neither of these alternative protocols had a substantial impact on the results obtained.

Enriched and shotgun libraries were then pooled separately and size selected on an E-Gel

SizeSelect 2% agarose gel (Thermo Fisher). Enriched libraries were then sequenced on an Illumina

MiSeq, as above, and shotgun libraries were sequenced on an Illumina NextSeq using a NextSeq

500/550 High Output Kit v2 (2x75 cycles).

Data analysis

Paired-end sequencing results were merged and adapters trimmed using leeHom[47], and reads were

then aligned to the human genome (hg37d5) with BWA (v0.7.12) aln, parameters -n 0.01 -l 0,

followed by samse[48]. Reads shorter than 28 bp long were then removed directly from sam files

with an awk command[39]. PCR duplicates were removed using MarkDuplicates (v2.9.0)[49] and reads

mapping with quality less than 25 were removed with SAMtools (v1.7)[50]. Reads mapping to the

nuclear genome with a mapping quality score of 25 or greater were locally realigned around known

indels using GATK (v3.7-0) IndelRealigner[51]. Following this step, mapped reads less than 35 bp

containing indels were removed[52]. This step reduced aberrant SNP calls due to spurious alignments

of short fragments in our dataset, where the base did not match either of two expected alleles, from

1.1% to 0.6%. In comparison, raising the minimum length of all reads to 30 bp reduced aberrant

SNP calls to 0.4%, but reduced informative SNPs by 18.4%, demonstrating the utility of retaining

a short minimum read length while excluding short indel-containing reads for this sample. All

libraries showed extremely poor preservation of genetic material. The average fragment length of

mapped reads after the above treatment was 38 bp, and the first position C>T transition rate from

damage was 55% at the 5' end of the molecule as calculated by mapDamage (v2.0.6)[53] (Extended

Data Figure 12).

Reads mapping to the mitochondrion were used to determine the posterior probability for

contaminating modern human sequences with Schmutzi[54] and found to have a distribution

maximum at 1%. A mitochondrial consensus sequence was called from the majority of bases at

each position using Geneious (v8.1.9)[55], which exactly matched that generated from Schmutzi.

The 5'-most 100 bp of this consensus sequence was duplicated at the 3' end and used as a reference

for a new alignment using the enriched reads. The haplogroup was called using Phy-Mer with

Build 16 rCRS-based haplogroup motifs[56] and verified by manual analysis of sequence changes.

A Bayesian tree of 25 ancient and 3 modern mitochondrial sequences, excluding the hypervariable

regions, was constructed using MrBayes[57] using a GTR+i+G nucleotide substitution model, which

gave the lowest log-likelihood for the tree out of all models tested (GTR and HKY with all

combinations of 4 invariant sites and gamma distributions), agreeing with the results of

JModelTest2[58]. The chain was run for 1,100,000 iterations, subsampled every 200 after discarding

a　9%　burn-in　period　and　visualized　using　FigTree　v.　1.4.3

(http://tree.bio.ed.ac.uk/software/figtree). A list of sources for the mitochondrial sequences is

given in Table S5.

Genetic sex was determined using the ratio of sequences aligning to the X and Y chromosomes,

given as the R_y value, as described in Skoglund, *et al.*[26]. An additional calculation of the ploidy of

the X chromosome using read counts mapping to the X chromosome to the autosomes,

($Reads_X/\overline{Reads}_{autosomes}$) was also performed. SNPs informative for the Y-chromosome haplogroup

were identified using Yleaf (v1.0)[59] for reads with mapping quality scores of both 10 and 20

(Supplementary Data 1).

Neanderthal content

Neanderthal ancestry was calculated using the ancestry informative SNP method[1] calculated

separately for Neanderthal- or AMH-derived SNPs on four libraries prepared from two

independent extractions, treated either with or without UDG. The combined libraries overlapped

a total of 6,252 SNPs reported to have derived in either the Neanderthal (2,538 SNPs) or AMH

(3,714 SNPs) lineages[60]. The combined libraries averaged 3.5% (SD 0.0074) Neanderthal

ancestry. To test the sensitivity of these SNPs to determine Neanderthal ancestry, this SNP subset

was then used to recalculate Neanderthal ancestry from several previously reported UP genomes[1].

The results agreed with previously reported values (Table S4).

SNP calling and f-statistics

BuranKaya3A bam files for all barcoded libraries were merged with SAMtools merge[50] and PCR

duplicates removed with MarkDuplicates[49]. SNPs were called for positions overlapping with the

combined SNP panels from Fu *et al.*[1] using SAMtools mpileup[50], requiring a SNP base quality

score of 30 or greater, and choosing one allele at random when necessary with pileupcaller[61]. This

resulted in 27,740 SNPs out of 2,990,848 being called for BuranKaya3A. To monitor the impact

cytosine deamination may have on the SNP calls and resulting analyses of ancient samples,

alternative datasets removing all C to T or G to A transitions were generated and all statistical

analyses were performed on both data sets. Results with >30% disagreement between the two

datasets were excluded from the analyses, but both datasets are included in the supplements.

To calculate f-statistics, data from 52 previously reported genomes (in either fastq or bam format)

were downloaded and realigned to the human genome (hg37d5), and SNPs were called following

the identical pipeline used for BuranKaya3A, with the exception that diploid SNP calls were

retained for high-coverage individuals used as an outgroup in $f3$ analyses (list of samples and

references given in Table S6). Modern humans used in statistical analyses are from Mallick, et

al.[62]. "Mbuti" is a population of three modern Mbuti individuals. $f3$-statistics and D-statistics were

computed using ADMIXTOOLS[29] qp3Pop (v412) qpDstat (v712), respectively. Standard error

was estimated using a block jackknife with 0.050 centiMorgan blocks. Full analyses performed

and results for $f3(x, y;$ Mbuti) and $D(w, y;$ BuranKaya3, Mbuti) are given in Table S7 and

Supplementary Data 2, respectively.

Data Availability

Sequence data generated in this study will be made available upon publication.

References

1. Fu, Q. *et al.* The genetic history of Ice Age Europe. *Nature* **534**, 200–205 (2016).

2. Prat, S. *et al.* The Oldest Anatomically Modern Humans from Far Southeast Europe: Direct

Dating, Culture and Behavior. *PLoS ONE* **6**, e20834 (2011).

3. Noiret, P. De quoi Gravettien est-il le nom ? in *Les Gravettians* 29–64 (Paris: Editions Errance,

2013).

4. Kozlowski, J. K. The origin of the Gravettian. *Quat. Int.* **359–360**, 3–18 (2015).

5. Conard, N. J. & Moreau, L. Current Research on the Gravettian of the Swabian Jura.

Mitteilungen Ges. Für Urgesch. **13**, 29–59 (2004).

6. Karel Valoch. L'origine du Gravettien de l'Europe centrale. in *The Upper Palaeolithic (ed*

Palma Di Cesnola, A., Montet-White, A. & Valoch, K.) **6**, 203–211 (1996).

7. Svoboda, J. A. On Modern Human Penetration to Northern Eurasia: the Multiple Advances

Hypothesis. in *Rethinking the human revolution : new behavioural and biological perspectives*

on the origin and dispersal of modern humans **19**, 329–339 (McDonald Institute for

Archaeological Research, Cambridge, UK : Oxford, UK ; Oakville, 2007).

8. Hoffecker, J. F. & Holliday, V. T. Landscape Archaeology and the Dispersal of Modern

Humans in Eastern Europe. in *The Upper Paleolithic of Northern Eurasia and America: Sites,*

Cultures, Traditions 140–158 (Petersburg 2014).

9. Golovanova, L. V. & Doronichev, V. B. VII - The Early Upper Paleolithic of the Caucasus in

the West Eurasian Context. in *The Aurignacian of Yafteh Cave and its context (2005-2008*

excavations) 137–160 (Liège University, 2012).

10.   Bar-Yosef, O. *et al.* Dzudzuana: an Upper Palaeolithic cave site in the Caucasus foothills

(Georgia). *Antiquity* **85**, 331–349 (2011).

11.   Golovanova, L. V. *et al.* The Early Upper Paleolithic in the northern Caucasus (new data

from Mezmaiskaya Cave, 1997 excavation). *Eurasian Prehistory* **4**, 43–78 (2012).

12.   Hoffecker, J. F. The early upper Paleolithic of eastern Europe reconsidered. *Evol.*

Anthropol. Issues News Rev. **20**, 24–39 (2011).

13.   Hublin, J.-J. The modern human colonization of western Eurasia: when and where? *Quat.*

Sci. Rev. **118**, 194–210 (2015).

14.   Demidenko, Y. E. Crimean Upper Paleolithic. in *Encyclopedia of Global Archaeology* (ed.

Smith, C.) 1782–1791 (Springer New York, 2014). doi:10.1007/978-1-4419-0465-2_1864

15.   Bar-Yosef, O., Belfer-Cohen, A. & Adler, D. S. The Implications of the Middle-Upper

Paleolithic Chronological Boundary in the Caucasus to Eurasian Prehistory. *Anthropologie*

XLIV, 49–60 (2006).

16.   Hoffecker, J. F. A New Framework for the Upper Paleolithic of Eastern Europe. (2012).

17.   Svoboda, J. A. The Gravettian on the Middle Danube. *PALEO Rev. Archéologie*

Préhistorique **19**, 203–220 (2007).

18.   Lazaridis, I. *et al.* Paleolithic DNA from the Caucasus reveals core of West Eurasian

ancestry. Preprint at bioRxiv doi: https://doi.org/10.1101/423079.

19.   Péan, S., Puaud, S., Crépin, L., Prat, S. & Quiles, A. The Middle to Upper Paleolithic

Sequence of Buran-Kaya III (Crimea, Ukraine): New Stratigraphic, Paleoenvironmental, and

Chronological Results. *Radiocarbon* **55**, 1454–1469 (2013).

20.   Yanevich, A. Les occupations gravettiennes de Buran-Kaya III (Crimée) : contexte

archéologique. *L'Anthropologie* **118**, 554–566 (2014).

21.   Fu, Q. *et al.* An early modern human from Romania with a recent Neanderthal ancestor.

Nature **524**, 216–219 (2015).

22.   Fernandes, V. *et al.* The Arabian Cradle: Mitochondrial Relics of the First Steps along the

Southern Route out of Africa. *Am. J. Hum. Genet.* **90**, 347–355 (2012).

23.   Costa, M. D. *et al.* A substantial prehistoric European ancestry amongst Ashkenazi

maternal lineages. *Nat. Commun.* **4**, (2013).

24.   Haak, W. *et al.* Ancient DNA from the First European Farmers in 7500-Year-Old Neolithic

Sites. *Science* **310**, 1016–1018 (2005).

25.   Lazaridis, I. *et al.* Genomic insights into the origin of farming in the ancient Near East.

Nature **536**, 419–424 (2016).

26.   Skoglund, P., Storå, J., Götherström, A. & Jakobsson, M. Accurate sex identification of

ancient human remains using DNA shotgun sequencing. *J. Archaeol. Sci.* **40**, 4477–4482

(2013).

27.   Fitzsimmons, K. E., Hambach, U., Veres, D. & Iovita, R. The Campanian Ignimbrite

Eruption: New Data on Volcanic Ash Dispersal and Its Potential Impact on Human Evolution.

PLoS ONE **8**, e65839 (2013).

28.   Higham, T. *et al.* The timing and spatiotemporal patterning of Neanderthal disappearance.

Nature **512**, 306–309 (2014).

29.   Patterson, N. *et al.* Ancient Admixture in Human History. *Genetics* **192**, 1065–1093

(2012).

30.   Bradley, B., Anikovich, M. & Engenii, G. Early Upper Paeolithic in the Russian Plain:

Streletskayan flaked stone artefacts and technology. *Antiquity* **69**, 989–998 (1995).

31.   Dobrovolskaya, M., Richards, M.-P. & Trinkaus, E. Direct radiocarbon dates for the Mid

Upper Paleolithic (eastern Gravettian) burials from Sunghir, Russia. *Bull. Mém. Société*

Anthropol. Paris **24**, 96–102 (2012).

32.   Pettitt, P. B. & Bader, N. O. Direct AMS radiocarbon dates for the Sungir mid Upper

Palaeolithic burials. *Antiquity* **74**, 269–270 (2000).

33.   Posth, C. *et al.* Pleistocene Mitochondrial Genomes Suggest a Single Major Dispersal of

Non-Africans and a Late Glacial Population Turnover in Europe. *Curr. Biol.* **26**, 827–833

(2016).

34.   Lipson, M. & Reich, D. A working model of the deep relationships of diverse modern

human genetic lineages outside of Africa. *Mol. Biol. Evol.* **34**, 889–902 (2017).

35.   Sinitsyn, A. A. Les Sepultures de Kostenki: Chronologie, Attribution Culturelle, Rite

Funéraire. in *La Spiritualité, Actes du colloque de la commission 8 de l'UISPP (Paléolithique*

supérieur), Liège, 10-12 décembre 2003. 237–244 (Etudes et Recherches Archeologiques de

l'Universite de Liege, 2004).

36.   Sikora, M. *et al.* Ancient genomes show social and reproductive behavior of early Upper

588   Paleolithic foragers. *Science* **358**, 659–662 (2017).

589   37.   Jones, E. R. *et al.* Upper Palaeolithic genomes reveal deep roots of modern Eurasians. *Nat.*

590   *Commun.* **6**, (2015).

591   38.   Reimer, P. J. *et al.* INTCAL13 and MARINE13 radiocarbon age calibration curves 0–

592   50,000 years CAL BP. *Radiocarbon* **55**, 1869–1887 (2013).

593   39.   Bennett, E. A. *et al.* Library construction for ancient genomics: single strand or double

594   strand? *BioTechniques* **56**, 289–298 (2014).

595   40.   Korlević, P. *et al.* Reducing microbial and human contamination in DNA extractions from

596   ancient bones and teeth. *BioTechniques* **59**, (2015).

597   41.   Gorgé, O. *et al.* Analysis of Ancient DNA in Microbial Ecology. *Methods Mol. Biol.*

598   *Clifton NJ* **1399**, 289–315 (2016).

599   42.   Glocke, I. & Meyer, M. Extending the spectrum of DNA sequences retrieved from ancient

600   bones and teeth. *Genome Res.* **27**, 1230–1237 (2017).

601   43.   Gansauge, M.-T. *et al.* Single-stranded DNA library preparation from highly degraded

602   DNA using T4 DNA ligase. *Nucleic Acids Res.* **45**, e79 (2017).

603   44.   Gansauge, M. T. & Meyer, M. Single-stranded DNA library preparation for the sequencing

604   of ancient or damaged DNA. *Nat Protoc* **8**, 737–48 (2013).

605   45.   Massilani, D. *et al.* Past climate changes, population dynamics and the origin of Bison in

606   Europe. *BMC Biol.* **14**, 93 (2016).

607   46.   Fu, Q. *et al.* DNA analysis of an early modern human from Tianyuan Cave, China. *Proc.*

608   *Natl. Acad. Sci.* **110**, 2223–2227 (2013).

47.   Renaud, G., Stenzel, U. & Kelso, J. leeHom: adaptor trimming and merging for Illumina
sequencing reads. *Nucleic Acids Res.* **42**, e141–e141 (2014).

48.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754–60 (2009).

49.   http://broadinstitute.github.io/picard/.

50.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–9 (2009).

51.   Van der Auwera, G. A. *et al.* From FastQ Data to High-Confidence Variant Calls: The
Genome Analysis Toolkit Best Practices Pipeline: The Genome Analysis Toolkit Best Practices
Pipeline. in *Current Protocols in Bioinformatics* 11.10.1-11.10.33 (John Wiley & Sons, Inc.,
2013).

52.   de Filippo, C., Meyer, M. & Prüfer, K. Quantifying and reducing spurious alignments for
the analysis of ultra-short ancient DNA sequences. *BMC Biol.* **16**, (2018).

53.   Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P. L. F. & Orlando, L. mapDamage2.0:
fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* **29**,
1682–1684 (2013).

54.   Renaud, G., Slon, V., Duggan, A. T. & Kelso, J. Schmutzi: estimation of contamination
and endogenous mitochondrial consensus calling for ancient DNA. *Genome Biol.* **16**, (2015).

55.   Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software platform
for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–9 (2012).

56.   Navarro-Gomez, D. *et al.* Phy-Mer: a novel alignment-free and reference-independent
mitochondrial haplogroup classifier. *Bioinforma. Oxf. Engl.* **31**, 1310–1312 (2015).

57.   Huelsenbeck, J. P. & Ronquist, F. MRBAYES: Bayesian inference of phylogenetic trees.

Bioinformatics **17**, 754–755 (2001).

58.   Darriba, D., Taboada, G. L., Doallo, R. & Posada, D. jModelTest 2: more models, new

heuristics and parallel computing. *Nat. Methods* **9**, 772 (2012).

59.   Ralf, A., Montiel González, D., Zhong, K. & Kayser, M. Yleaf: Software for Human Y-

Chromosomal Haplogroup Inference from Next-Generation Sequencing Data. *Mol. Biol. Evol.*

35, 1291–1294 (2018).

60.   Prufer, K. *et al.* The complete genome sequence of a Neanderthal from the Altai

Mountains. *Nature* **505**, 43–49 (2014).

61.   https://github.com/stschiff/sequenceTools.

62.   Mallick, S. *et al.* The Simons Genome Diversity Project: 300 genomes from 142 diverse

populations. *Nature* **538**, 201–206 (2016).

Acknowledgments

We thank Olivier Gorgé for assistance with DNA sequencing. We thank also the National

Academy of Sciences of Ukraine for permission to excavate at Buran-Kaya III, and all the team

members of the 2001 and 2009–2011 excavation seasons. EAB was supported by the CNRS and

the Labex "Who am I". SPr was supported by the Foundation Fyssen. SPe was supported by the

French National Research Agency (ANR-05-JCJC-0240-01). The field work was supported by the

Muséum National d'Histoire Naturelle (MNHN, Paris), the CNRS and French Ministry of Foreign

Affairs.The paleogenomic facility obtained support from the University Paris Diderot within the

program "Actions de recherches structurantes". The sequencing facility of the Institut Jacques

Monod, Paris, was supported by grants from the University Paris Diderot, the Fondation pour la

Recherche Médicale (DGE20111123014), and the Région Ile-de-France (11015901).

Author contributions

EMG, SPr, and SP initiated the project, EMG and EAB designed the study. EAB performed the

laboratory work, EAB, TG, and EMG analysed the data, LC performed aseptic excavation of the

sample. SPr performed the anthropological analysis. AY, SP and SPu provided archaeological

data. EAB wrote the manuscript with input from EMG, TG, SPr, SP, LC, and AY.

Competing Interests

The authors declare that they have no competing interests that might have influenced the work

described in this manuscript.

Correspondence should be addressed to EAB (eabennett@gmail.com)

Figure 1. Recalibration of comparative AMS ¹⁴C dates of Buran-Kaya III, layer 6-1 with Early
and Mid-Upper human Palaeolithic European samples. All dates were recalibrated using the
software OxCal v4.3.2 based on the IntCal13 calibration data set[38]. For complementary
information on cultural contexts, see Table S1.

Figure 2. Bayesian phylogenetic tree of mitochondrial sequences, excluding Hyper-Variable Regions, from Early and Mid-Upper Palaeolithic individuals including BuranKaya3A. Posterior probability indicated at the nodes. Non-European individuals are in parentheses. General ages shown as either closed or open circles, and pre-glacial material cultures, when known, indicated by colour. A question mark by the Aurignacian indicates a cultural assignment by dating rather than direct association. Scale bar denotes substitutions per site. Sources for mitochondrial sequences are listed in Table S5.

Figure 3. Location and heatmap of *f*3(BuranKaya3A, *x*; Mbuti) with archaeological sites mentioned in the text. Values of high genomic coverage ancient samples (higher values show more shared alleles with BuranKaya3A). Boxed text indicates samples associated with a Gravettian archaeological context (including Sunghir3, alternatively described as Streletskian). Non-European samples are given along the right margin at their approximate latitudes, their relative distances indicated by black arrows. Archaeological sites for which no human genetic data are available that contain micro-laminar industries comparable with Buran-Kaya III at contemporaneous layers (in parentheses) indicated by black dots. A broad green arrow shows the proposed EUP route introducing the Early Gravettian into Europe, suggested by the similar features of these assemblages and the lack of a Common Western Eurasian genomic component in BuranKaya3A.

Figure 4. D-statistics results for a) D(x, Ust-Ishim; BuranKaya3A, Mbuti) and b) D(x, Han; BuranKaya3A, Mbuti) where x = selected UP and Mesolithic Eurasians. Starred individuals denote a significant Z-score >2. Approximate ages, being the mean of the latest calibrated published date rounded to the nearest ky, are appended to the names. More positive values represent increased allele sharing between x and *BuranKaya3A* relative to *Ust-Ishim* or *Han*, with *Mbuti* as an outgroup. Results from all SNPs shown. All Z-scores and results from transversions only are given in Extended Data Figure 4. The values for Kotias, Satsurblia, and the Natufians are distorted due to Basal Eurasian content in these samples (see Extended Data Figure 9). Error bars = one standard error.

Extended Data Figure 1. Flowchart of screening methods and results of mitochondrial capture and

phosphate enrichment for each library. Results show the percent of unique reads out of all reads

28 nucleotides or greater mapping to the nuclear and mitochondrial references as well as the

percent of PCR duplicates for each library. The suffix of sample numbers is as follows, A = no

phosphate treatment, B = phosphate treated, Bp = phosphate treatment buffer. Extract BKA4B was

selected for subsequent analysis. Original bone sample and regions used for each library is shown

above. It has been noted that the region containing the darker suture yielded the best results.

Extended Data Figure 2. Detailed mutation map of mitochondrial N clade with the positions of BuranKaya3A, the Epipalaeolithic Natufian (Natufian9), and lineages discussed in the text. Grey italics represent the population or geographical region where the clade is prominent. Inset indicates the maximum likelihood estimation of TMRCA values of the nodes with 95% CI from Fernandes et al.[22]. Mutation tree calculated by mtphyl version 5.003 (https://sites.google.com/site/mtphyl/home) compared to the rCRS. Underlined positions represent back mutations.

Extended Data Figure 3. Outgroup *f*3-statistics results showing the degree of shared alleles between BuranKaya3A and high-coverage UP Eurasians for outgroups Mbuti (a), Han (b). Left axis: *f*3-statistic corresponding to circles with error bars. Right axis: Z-score corresponding to "x". Results for all SNPs are in blue, transversions only in red. The degree of overlap between blue and red shows the degree of agreement between the two datasets for a given sample combination. Approximate ages are appended to the names. Boxed names indicate individuals associated with a Gravettian context. Error bars = one standard error.

Extended Data Figure 4. D-statistics results for a) $D(x$, Ust-Ishim; Burankaya3A, Mbuti) and b)

$D(x$, Han; BuranKaya3A, Mbuti) where $x =$ high coverage UP individuals. Left axis: D-statistic

corresponding to circles with error bars. Right axis: Z-score corresponding to "x". Results for all

SNPs are in blue (shown also in Figure 4), transversions only in red. The degree of overlap between

blue and red shows the degree of agreement between the two datasets for a given sample

combination. Approximate ages are appended to the names. Boxed names indicate individuals

associated with a Gravettian context. The values for Kotias, Satsurblia, and the Natufians are

distorted in (a) due to Basal Eurasian content in these samples. Error bars = one standard error.

Extended Data Figure 5. D-statistics results for $D(x,$ Ust-Ishim; Burankaya3A, Mbuti) where $x =$ both high and low coverage UP individuals and a modern East Asian (Han). Only samples using at least 100 transversion-only SNPs for the calculation are shown. Left axis: D-statistic corresponding to circles with error bars. Right axis: Z-score corresponding to "x". Results for all SNPs are in blue, transversions only in red. The degree of overlap between blue and red shows the degree of agreement between the two datasets for a given sample combination. Approximate ages are appended to the names. Boxed names indicate individuals associated with a Gravettian context. The values for Kotias, Satsurblia, and the Natufians are distorted due to Basal Eurasian content in these samples. Error bars = one standard error.

a

b

Extended Data Figure 6. D-statistics results for $D(w, x;$ BuranKaya3A, Mbuti) where w and x are

high-coverage UP Eurasians. More positive values show an excess of allele sharing of w over x

with BuranKaya3A. Left axis: D-statistic corresponding to circles with error bars. Right axis: Z-

score corresponding to black dots. Significance (Z-score > 2) is indicated by horizontal dashed

lines. The values for Kotias are distorted due to Basal Eurasian content in this sample. Error bars

= one standard error. a) Results for all SNPs, b) results for transversions only.

Extended Data Figure 7. Location and heatmap of *f*3(BuranKaya3A, *x*; Mbuti) with archaeological

sites mentioned in the text. Values, as in Figure 3, but adding results from *x* including the lower-

coverage samples whose calculations made use of at least 50 transversion-only SNPs and were

within 30% agreement of values calculated using all SNPs. Boxed text indicates samples

associated with a Gravettian archaeological context (including Sunghir3, alternatively described

as Streletskian). Non-European samples are given along the right margin at their approximate

latitudes, their relative distances indicated by arrows. Archaeological sites for which no human

genetic data are available that contain micro-laminar industries comparable with Buran-Kaya III

at contemporaneous layers (in parentheses) indicated by black dots. A broad green arrow shows

the proposed EUP route introducing the Early Gravettian into Europe, suggested by the similar

features of these assemblages and the lack of a Common Western Eurasian genomic component

in BuranKaya3A.

Extended Data Figure 8. Outgroup *f3*-statistics results showing the degree of shared alleles

between BuranKaya3A and UP Eurasians for outgroups Mbuti, $f3$(BuranKaya3A, *x*; Mbuti), as in

Extended Data Figure 3A but including all samples for which values could be calculated. Left axis:

f3-statistic corresponding to circles with error bars. Right axis: Z-score corresponding to "x".

Results for all SNPs are in blue, transversions only in red. The degree of overlap between blue and

red shows the degree of agreement between the two datasets for a given sample combination.

Approximate ages are appended to the names. Boxed names indicate individuals associated with

a Gravettian context. Error bars = one standard error.

Extended Data Figure 9. Relative Basal Eurasian content as demonstrated by the statistic

D(*Modern or Ancient East Asian*, *x*; Ust-Ishim, Mbuti)[1] using the transversions only SNP dataset.

Left axis: D-statistic corresponding to circles with error bars. Right axis: Z-score corresponding to

"x". *x* =Han, blue, *x*=Tianyuan, red. Significance (Z-score = 2) is indicated by a horizontal dashed

line. Estimated Basal Eurasian content percentages given as suffixes to the names (a) estimated

from ADMIXTUREGRAPH[1], (b) one standard error range estimated from the f4 ratio[25]. Error

bars = one standard error.

Extended Data Figure 10. Relative Common West Eurasian content (as represented by Villabruna) given as $D(x$, Han; Villabruna, Mbuti) for ancient samples (x) using more than 1,000 SNPs for the calculation. More positive values signify an excess of shared alleles between x and Villabruna as compared to Han. Reported genetic clusters[1] given as suffixes to the names: VN, Vestonice; EM, ElMiron; VB, Villabruna; SS, Satsurblia; MA, Mal'ta. Boxed names indicate individuals associated with Gravettian contexts. Left axis: D-statistic corresponding to circles with error bars. Right axis: Z-score corresponding to "x". Results for all SNPs are in blue, transversions only in red. The degree of overlap between blue and red shows the degree of agreement between the two datasets for a given sample combination. This value will be distorted for samples having Basal Eurasian content (see Extended Data Figure 9). Significance (Z-score = 2) is indicated by a horizontal dashed line. Error bars = one standard error.

Extended Data Figure 11. Measuring the excess of shared alleles between Common West Eurasian

(as represented by Villabruna) and *x* over BuranKaya3A using the D-statistic *D*(*x*, BuranKaya3A;

Villabruna, Mbuti). More positive values signify an excess of shared alleles between *x* and

Villabruna compared to BuranKaya3A. Reported genetic clusters[1] given as suffixes to the names:

VN, Vestonice; EM, ElMiron; VB, Villabruna; SS, Satsurblia; MA, Mal'ta. Boxed names indicate

individuals associated with Gravettian contexts. Left axis: D-statistic corresponding to circles with

error bars. Right axis: Z-score corresponding to "x". Results for all SNPs are in blue, transversions

only in red. The degree of overlap between blue and red shows the degree of agreement between

the two datasets for a given sample combination. This value will be distorted for samples having

Basal Eurasian content (Extended Data Figure 9). Significance (Z-score = 2) is indicated by a

horizontal dashed line. Error bars = one standard error.

Extended Data Figure 12. a) Size distribution of recovered BuranKaya3A DNA fragments mapping to the nuclear genome. b) The frequency of C to T mismatches due to cytosine deamination of the first 25 nucleotides from the 5' and 3' ends of the DNA molecules. Graph generated by mapDamage (v2.0.6)[53].