Manipulating immune pathways differentially impact emotionality: dissociable effects of complement C3 and C3aR on learned fear and innate anxiety

Laura J. Westacott[1,4†], Trevor Humby[1,2,4†], Niels Haan[1,4], Sophie A. Brain[2], Emma-Louise Bush[2], Margarita Toneva[2], Anna L. Moon[1], Jack Reddaway[1,4], Michael J. Owen[1,] Jeremy Hall[1,4], Timothy R. Hughes[3,4], B. Paul Morgan[3,4,6], William P. Gray[1,4,5*] & Lawrence S. Wilkinson[1,2,4*]

[1] Neuroscience and Mental Health Research Institute, MRC Centre for Neuropsychiatric Genetic and Genomics, School of Medicine, Hadyn Ellis Building, Cardiff University, Cardiff, CF24 4HQ, UK.[2] Behavioural Genetics Group, Schools of Psychology and Medicine, Cardiff University, Cardiff, CF10 3AT, UK. [3] Complement Biology Group, Systems Immunity Research Institute, School of Medicine, Cardiff University, CF14 4XW, Cardiff UK. [4] Hodge Centre for Neuropsychiatric Immunology, School of Medicine, Cardiff University, Cardiff CF24 4HQ, UK. [5] Brain Repair and Intracranial Therapeutics (BRAIN) Unit, School of Medicine, Cardiff University, CF24 4HQ, UK. [6] UK Dementia Research Institute, Cardiff University, Cardiff, CF24 4HQ, UK.

*Corresponding authors: Lawrence S. Wilkinson & William P. Gray

Email address: WilkinsonL@cardiff.ac.uk; GrayWP@cardiff.ac.uk

Address: Neuroscience and Mental Health Research Institute, School of Medicine, Cardiff University, Cardiff, CF24 4HQ, UK

[†] These authors contributed equally to the work

Key words: *Complement system, Fear, Anxiety, Stress response, Behaviour, Neuroimmunology*

Abstract length: 244/250

Article length: 3965 / 4000

Number of Figures: 8

Number of Tables: 0

Number of Supplementary Files: 3

Abstract

Background Recent findings implicate complement, a key component of the

immune system, in healthy brain functioning and risk for psychiatric disorders.

Altered emotional function, in particular maladaptive fear and anxiety, is a pervasive

and clinically important symptom across several neuropsychiatric disorders. In this

work we examined the extent to which different complement pathways impact on

dissociable components of anxiety and learned fear.

Methods: $C3^{-/-}$ and $C3aR^{-/-}$ mice were assessed in a battery of tests assaying innate

anxiety and learned fear. The effects of systemic diazepam (2mg/kg) and the C3aR

antagonist SB290157 (10mg/kg) on behaviour was assessed. Expression of genes

linked to anxiety, stress responses and risk for psychiatric disorder were assessed in

brain regions implicated in fear and anxiety.

Results: The data indicated a robust and highly reproducible double-dissociation of

C3 and C3aR on learned fear and innate anxiety, respectively. We probed

downstream complement pathways responsible for the C3-specific effects on

learned fear and excluded contributions from the terminal pathway, C6 to C9. The

selective anxiety phenotype in $C3aR^{-/-}$ mice was resistant to the anxiolytic properties

of diazepam and acute SB290157 administration in wildtypes did not phenocopy

constitutive $C3aR$ knockout, suggesting $C3aR^{-/-}$ anxiety phenotypes resulted from

long-term $C3aR$ deficiency. Expression of the calcium voltage-gated channel subunit

Alpha1c (*Cacna1c*) mRNA, an established psychiatric risk gene, was selectively

increased in $C3aR^{-/-}$ mice.

Conclusions: These findings reveal novel functions and unexpected dissociations

between the C3 and C3aR pathways impacting on specific emotional behaviours

relevant to psychiatric disorder.

Introduction

The complement system is a key component of the immune system that plays a

pivotal role in inflammation and host-defence. Complement activation occurs via

several pathways, all of which lead to cleavage of the central protein C3 (see

Supplementary Figure S1)(1). Activation of C3 generates the fragments C3a and

C3b. C3a is an anaphylatoxin which upon binding to its canonical G-protein coupled

receptor C3aR, induces both pro- and anti-inflammatory effects(2). The C3b

fragment plays a role in opsonization by macrophages and microglia via complement

receptor 3 (CR3) and propagates further complement activation by contributing to

the cleavage of complement component 5 (C5). Akin to C3, C5 cleavage yields C5a,

a ligand for the C5a receptor (C5aR), and C5b, which triggers the terminal

complement pathway by binding proteins C6, C7, C8 and C9 which form the

membrane attack complex (MAC), leading to lytic cell death or activation(3).

Neurons, astrocytes and microglia express complement receptors and are capable

of synthesising complement proteins(4; 5), the expression patterns of which vary

over the course of brain development 6).

It is increasingly recognised that complement has versatility beyond its canonical

immune functions(7). In the CNS, evidence has implicated complement in

neurodevelopmental processes including neurogenesis(8), migration(9) and synaptic

pruning(10), and in the pathogenesis of neurodevelopmental, neurodegenerative and

psychiatric conditions. In Alzheimer's disease, genetic variants in complement

related loci have been associated with increased disease risk(11; 12), and

complement knockout mice exhibit reduced age-related synapse loss(13) and

neuropathology(14). Alterations in complement proteins and activation have also

been reported in serum from individuals with autism-spectrum disorder(15)

schizophrenia(16), major depressive disorder(17), bipolar disorder(18) and post-

traumatic stress disorder(19). In the case of schizophrenia, an important finding

comes from elegant genetic work demonstrating that structural variation in the *C4*

locus is associated with risk of developing the disease(20). C4 cleavage generates

fragments that contribute to the activation of C3, yielding C3a and C3b. Given the

known roles for C3b/CR3 in synaptic pruning(10; 21), it has been suggested that *C4*

variants may impact on psychiatric risk via this mechanism, leading to abnormal

connectivity and disruption of neural networks(20). Variants in *C3* and putative

complement-control genes *CSMD1* and *CSMD2* have also been implicated in

genetic susceptibility for schizophrenia(22; 23).

Altered emotional function, in particular excessive fear and anxiety, is a pervasive

and clinically important symptom that is comorbid across several of the DSM-5 and

ICD-11 defined disorders in which complement has been implicated. Anxiety is

characterised by sustained arousal, negative affect, apprehension and

hypervigilance surrounding anticipated or potential threats, while fear is an acute

response to an experienced, imminent danger(24). It is important to note that while

these processes share some underlying neurobiology, they also have contributions

from distinct neuronal circuitries(25). Although links between anxiety and

inflammation have been documented(26), there has been little investigation of the

specific role of complement in anxiety and fear. In animal models, prenatal

manipulations increasing C3 deposition in the fetal brain are associated with adult

anxiety phenotypes(27). Additionally, complement inhibitors reduced anxiety in a

model of Alzheimer's disease(28) and aged *C3* deficient mice exhibit reduced

anxiety alongside enhanced contextual fear responses(13). These studies suggest

that complement influences fear and anxiety; however, the precise relationship

between signalling within discrete pathways of the complement system (e.g.,

C3a/C3aR or C3b/CR3 axes) and their relationship to dissociable aspects of anxiety

and fear and underlying brain function is unknown.

In this work we used knockout mouse models to manipulate C3 and C3aR signalling

in order to functionally parse anxiety and fear related phenotypes. In homozygous

C3 knockout mice[29] complement cannot be activated beyond C3, and therefore

these animals lack C3 activation fragments (C3a, C3b) and downstream activation

products (C5a, C5b) thus cannot activate the terminal complement pathway.

Phenotypic effects in this model could therefore be the result of disturbance to any

downstream effector molecule(s). We compared the $C3^{-/-}$ model with homozygous

C3aR knockout mice[30].   In these mice, complement is intact apart from the

capacity for C3a to bind to its canonical receptor C3aR and here we tested the

extent to which phenotypic effects were the result specifically of C3a/C3aR

signalling. *A priori*, because C3a is an obligate cleavage fragment of C3 we

hypothesised that any phenotypes dependent on the binding of C3a to C3aR would

be apparent in both $C3^{-/-}$ and $C3aR^{-/-}$ models.

In contrast to this initial expectation we found completely different effects of the two

models on emotional behaviours. $C3aR^{-/-}$ mice displayed a profound innate anxiety

phenotype across a range of measures that was lacking in the $C3^{-/-}$ model. When we

examined learned fear (where a previously neutral cue generates a fear response as

a result of predicting an aversive outcome) we found that the dissociation was

reversed, with the $C3^{-/-}$ mice exhibiting an enhanced fear response to a conditioned

cue, but no differences between wildtype and $C3aR^{-/-}$mice. We probed downstream

complement pathways responsible for the C3-specific effects on learned fear and

excluded contributions from the terminal pathway by using $C6^{-/-}$ mice. These findings

point to a hitherto unrecognized complexity of complement effects on brain function

and behaviour of potential relevance to emotional dysfunction in psychopathology.

Materials and Methods

For additional materials and methods, see Supplementary Information.

Animals. Wildtype and $C3^{-/-}$ strains were sourced from The Jackson Laboratory

(strain B6.PL-Thy1^a/CyJ stock#000406 and B6;129S4-C3tm1Crr/J stock#003641

respectively), whereas $C3aR^{-/-}$ mice were provided by Craig Gerard of Boston

Children's Hospital, USA and $C6^{-/-}$ mice were an in-house strain. $C3^{-/-}$, $C3aR^{-/-}$ and

$C6^{-/-}$ strains were maintained via homozygous breeding on a C57Bl/6J background.

Male mice (3-8 months old) in littermate groups (2-5 mice/cage) were kept in a

temperature and humidity-controlled vivarium (21±2°C and 50±10%, respectively)

with a 12-hour light-dark cycle (lights on at 07:00hrs/lights off at 19:00hrs). Home

cages were environmentally enriched with cardboard tubes, soft wood blocks and

nesting materials. Standard laboratory chow and water were available *ad libitum*. All

procedures were performed in accordance with the requirements of the UK Animals

(Scientific Procedures) Act (1986). Sample sizes for each experiment are provided in

the corresponding Figure.

General behavioural methods. Testing took place between the hours of 09:00 and

17:00, with random distribution of testing for subjects of different genotypes

throughout the day. Mice were habituated to the test rooms for 30 min prior to

testing. All assays involved individual testing of mice and apparatus was cleaned

thoroughly with a 70% ethanol solution between subjects. Data for the EPM, EZM

and Open Field were collected using EthoVision XT software (Noldus Information

Technology, Netherlands).

The elevated plus maze (EPM). Subjects were placed at the enclosed end of a

closed arm and allowed to freely explore for 5 min. Data from each pair of arms

were combined to generate single open and closed arm values. In addition, the

following parameters were manually scored: number of stretch-attend postures

(SAPs; defined as the animal slowly and carefully reaching towards the open arms in

a low, elongated body posture(31; 32)), number of head dips (looking over the edge

of an open arm) and grooming frequency.

The elevated zero maze (EZM). Subjects were placed at the border of one of the

open and closed regions and allowed to freely explore for 5 min. Data from each

pair of regions were combined to generate single open and closed region values. In

addition, the number of head dips (as above) were measured.

Assessment of fear potentiated startle (FPS). FPS was assessed using a pair of

ventilated and soundproofed SR-LAB startle chambers (San Diego Instruments, CA,

USA). FPS consisted of three separate sessions presented over a two-day period

(see Figure 7A). FPS was determined as the fold change between pulse-alone trials

and pulse+CS trials within the post-conditioning session.

Pharmacological manipulations

Diazepam A three-day dosing regimen of diazepam (2 mg/kg, i.p., Hameln

Pharmaceuticals, UK) or an equivalent volume of vehicle (0.1 M phosphate buffered

saline, pH 7.4) was used. Following 2 days of pre-treatment, diazepam or vehicle

was administered 30 min prior to testing on the EPM on the 3[rd] day.

C3aR antagonist Two dosing regimens were used to assess effects of

SB290157(33) (N2-[2-(2,2-Diphenylethoxy)acetyl]-L-arginine trifluoroacetate salt; 10

mg/kg, i.p, Sigma Aldrich, UK) or equivalent volume of vehicle (0.1 M phosphate

buffered saline, pH 7.4, with 1% DMSO) on behaviour.

Acute SB290157 or vehicle was administered 1 hour prior to testing on the EPM.

Semi-chronic SB290157 or vehicle was injected twice daily for five days and

subjects were tested on the EPM on the 6th day.

Corticosterone measurements in mice subjected to the EPM. Testing took place

between the hours of 10:00 and 14:00 to account for the diurnal pattern of

corticosterone release(34). Subjects were culled 20 minutes post-EPM by cervical

dislocation. Control mice were removed from their home cage and immediately

culled. Plasma was collected from trunk blood and frozen at -80°c until required.

Corticosterone ELISA was performed according to manufacturer's instructions (ADI-

900-097, Enzo Life Sciences, UK).

Quantitative Polymerase Chain Reaction (qPCR) Tissue from behaviourally naïve

mice of each genotype was used to assess gene expression using standardised

quantitative qPCR methods (see Supplementary Information), with quantification

using the $2^{-\Delta\Delta Ct}$ method (35). All samples were run in triplicate and samples differing

by >0.3 Cts were excluded. The change in expression of genes of interest, after

normalisation to two house-keeping genes (ΔCt) was transformed to yield $2^{-\Delta\Delta Ct}$

values. Relative changes from wildtype expression were calculated for each gene of

interest.

Statistical analysis. All statistical analyses were carried out using GraphPad Prism

8.0 (GraphPad Software, CA, USA). Data was assessed for equality of variances

using the Brown-Forsythe test and then appropriate parametric (one-way or two-way

ANOVA) or non-parametric (Kruskal-Wallis) tests used.   *Post hoc* pairwise

comparisons were performed using the Tukey or Dunn's tests for parametric or non-

parametric analyses, respectively. Exact p values were reported unless p<0.0001. All

p values were multiplicity adjusted(36). Data are expressed as mean + standard

error of the mean.

Results

Increased innate anxiety in *C3aR*[-/-] but not *C3*[-/-] mice

We assessed emotional reactivity in the elevated plus maze (EPM), a well-

established test of innate anxiety in rodents(37; 38). As demonstrated in Figure 1A,

heatmaps indexing overall maze exploration indicated significant differences in open

arm exploration between genotypes (see Supplementary Video 1 for representative

examples). In comparison to wildtype and *C3*[-/-] mice, *C3aR*[-/-] mice took longer to first

enter the open arms (Figure 1B) and spent less time there per entry (Figure S2A),

leading to a reduced overall duration spent in the open arms (Figure 1C). The

ethological parameters of head dips and SAPs also differed (Figure 1D,E), with

C3aR[-/-] mice exhibiting a significant reduction in the former and significant increase

in the latter, consistent with increased anxiety(39).   We noted a significantly

increased frequency of head dipping in *C3*[-/-] mice (Figure 1D) which may indicate

reduced anxiety(40). These initial data were consistent with an anxiogenic

phenotype present in *C3aR*[-/-] mice but absent in *C3*[-/-] mice. As might be predicted

from a heighted state of anxiety, we observed reduced locomotion around the maze

in $C3aR^{-/-}$ mice (Figure S2B). Given that several measures were dependent on

movement, it was important to eliminate potential locomotor confounds. To address

this issue, we measured activity independently in a non-anxiety provoking

environment over three days and found no group differences (Figure S2C) indicating

that EPM measures were not influenced by movement confounds.

In a further cohort of animals, we assessed behaviour in the elevated zero maze

(EZM), another established assay of anxiety related-behaviour(41). These data

recapitulated the pattern of findings seen in the EPM with the $C3aR^{-/-}$ mice avoiding

the open arms of the maze to a much greater extent than the wildtype and $C3^{-/-}$

animals (Figure 1F-J). Additional data from the open field test were consistent with

the maze data in demonstrating a $C3aR^{-/-}$ specific anxiety phenotype (Figure S3).

Excessive grooming in $C3aR^{-/-}$ mice

During behavioural testing we noticed an increase in grooming in the $C3aR^{-/-}$ mice

that was present in all $C3aR^{-/-}$ cohorts tested. We quantified this behaviour in one of

the EPM sessions, as shown in Figure 2A&B. In the context of the EPM, grooming

may be a displacement behaviour(42) and is positively correlated with anxiogenic

phenotypes(43). Furthermore, excessive grooming in the $C3aR^{-/-}$ mice was chronic

and present in the home-cage environment, resulting in substantial hair loss and

occasionally skin lesions (see Figure 2C for representative examples). Again, this

phenotype was not observed in $C3^{-/-}$ or wildtype mice.

Altered sensitivity of *C3aR^{-/-}* and *C3^{-/-}* mice to diazepam in the elevated plus

maze

In a separate cohort of animals, we repeated the EPM experiment and in addition

tested sensitivity of anxiety-related behaviour to 2mg/kg diazepam, an established

clinically effective anxiolytic drug(37; 44). Vehicle-treated animals replicated the

initial findings across all behavioural indices of anxiety, again showing a dissociation

between the *C3aR^{-/-}* and *C3^{-/-}* mice (Figure 3A). In wildtype mice diazepam led to a

trend for increased time on the open arms (Figure 3B), an effect often held to index

reduced anxiety(37; 44; 45). While these effects did not reach statistical significance

there was a highly significant reduction in SAPs consistent with anxiolysis(40; 45)

Figure 3C). In contrast to wildtypes, diazepam was without effects in *C3aR^{-/-}* mice

(Figure 2A,B,C) and produced an anxiogenic pattern of effects in *C3^{-/-}* mice (Figure

2A,B and Figure S4B,C). Locomotor activity monitored across all areas of the maze

(Figure S4D) indicated that wildtype and *C3aR^{-/-}* mice were unlikely to have been

influenced by sedation. In *C3^{-/-}* mice however, activity was significantly suppressed

under drug conditions indicating a possible sedative effect.

Enhanced corticosterone response to the EPM in *C3aR^{-/-}* mice

In a sub-cohort of wildtype, *C3^{-/-}* and *C3aR^{-/-}* mice treated with vehicle in the

diazepam study we assayed plasma corticosterone 30 minutes after exposure to the

EPM and compared levels to those of a group of animals who remained in their

home-cages (baseline condition). There were no genotype differences in basal

levels of corticosterone (Figure 4); however, being placed on the EPM increased

plasma corticosterone 6-15-fold in all genotypes, suggesting that the EPM was a

potent stressor. *Post hoc* analyses indicated a significantly greater EPM-evoked

corticosterone response in the *C3aR$^{-/-}$* animals, consistent with their anxiety-like

behaviour observed on the maze.

Changes in expression of stress and anxiety related genes in *C3$^{-/-}$* and *C3aR$^{-/-}$*

mice

Using qPCR we measured expression of genes linked to anxiety and the stress

response in brain regions implicated in fear and anxiety(25). Both knockouts

demonstrated significantly greater cerebellar expression of *Nr3c1* encoding the

glucocorticoid receptor (Figure 5A) and there was a trend in *C3aR$^{-/-}$* mice for

increased expression in the ventral hippocampus (vHPC) and medial prefrontal

cortex (mPFC). Expression of *Crhr1* (encoding the corticotrophin-releasing hormone

receptor 1) and *Gabra2*, which encodes the GABA$_A$ receptor $\alpha 2$ subunit responsible

for mediating benzodiazepine anxiolysis (46) was unaltered (Figure 5C,D). Since

activation of C3aR is known to stimulate calcium influx(47; 48), we also investigated

the expression of *Cacna1c,* a gene which encodes the Ca$_v$1.2 subunit of the L-type

calcium channels and has established links to psychiatric disorder and anxiety(49;

50). We found a significant increase in expression (Figure 5B) that was genotype

and brain region specific; with selective increases in expression in *C3aR$^{-/-}$* mice in

the vHPC and cerebellum, but not the mPFC.

Administration of the C3aR antagonist SB290157 did not phenocopy *C3aR^{-/-}*
mice

We next examined the extent to which systemic administration of the C3aR
antagonist SB290157(33) to wildtype mice, at doses which have been shown to
influence brain phenotypes(8; 48), would phenocopy *C3aR^{-/-}* mice. We tested the
effects of both acute and semi-chronic SB290157 treatment in the EPM. While the
previous finding of enhanced anxiety was replicated in vehicle-treated *C3aR^{-/-}* mice
in both acute (Figure S5) and semi-chronic dosing regimens (Figure 6) neither
SB290157 treatment had any effects on behaviour in wildtype mice.

Enhanced learned fear in *C3^{-/-}* but not *C3aR^{-/-}* mice

In a separate group of animals, we next assessed fear potentiated startle (FPS; see
Figure 7A)(51; 52), an established means of assaying learned (or conditioned)
fear(53; 54). In the pre-conditioning session, pulse-alone trials indicated increased
basal startle reactivity in both *C3aR^{-/-}* and *C3^{-/-}* animals relative to wildtype (Figure
7B). Increased reactivity to the unconditioned foot shock stimulus (in the absence of
any startle stimulus) during the conditioning session was also observed in both
knockouts (Figure 7C). The potentiation of the startle response by the CS in the
post-conditioning session is illustrated in Figure 7D. Here, in contrast to the previous
assays of innate anxiety an opposite pattern of effects was observed, with *C3^{-/-}*
animals showing a selective enhancement of conditioned fear relative to the *C3aR^{-/-}*
and wildtype animals.

No effects on learned fear in *C6^{-/-}* mice

The demonstration of enhanced fear learning in *C3^{-/-}*, but not *C3aR^{-/-}* mice suggested

that effects in *C3^{-/-}* mice were mediated by molecules downstream of C3,

independent from the C3a/C3aR axis. As there is no activity beyond the C3

convertase in *C3^{-/-}* mice, C3 cannot be cleaved and therefore *C3^{-/-}* mice lack

C3a/C3b and C5a/C5b in addition to the terminal pathway (C6-C9). Therefore, it is

possible that indirect loss of function of these molecules may contribute to the

observed phenotype in *C3^{-/-}* mice, although it is unclear which one. We therefore

compared *C3^{-/-}* with *C6^{-/-}* mice (which have intact C3b/CR3, C5b and C5a/C5aR

function but lack the terminal pathway) and hypothesised that should *C6^{-/-}* mice show

comparable behaviour to *C3^{-/-}* mice then the phenotype would likely be due to a loss

of terminal pathway activity. On the other hand, if *C6^{-/-}* mice demonstrated no fear

learning phenotype, this would confine the likely mediating molecules to either C3b

or the downstream protein C5 and its breakdown products. Using a separate cohort,

we replicated the previous *C3^{-/-}* finding of enhanced learned fear (Figure 8).

However, across all measures *C6^{-/-}* mice were comparable to wildtype mice,

suggesting that loss of C3b/CR3 or C5a/C5aR signaling, but not the terminal

pathway, was involved in the *C3^{-/-}* fear learning phenotype. Additionally, we did not

observe innate anxiety-like phenotypes in *C6^{-/-}* mice (Figure S6).

Discussion

We have used knockout models manipulating specific complement proteins to reveal

dissociable effects on emotional behaviours. A main finding was a profound innate

anxiety phenotype in the *C3aR^{-/-}* mice that was absent in *C3^{-/-}* mice. The specificity of

the anxiety phenotype exhibited by *C3aR*[-/-] mice at the behavioural level was

paralleled by EPM-evoked corticosterone levels, confirming the validity of the EPM

as an index of anxiety-like behaviour. *C3aR*[-/-] mice also displayed a pathological

grooming phenotype, which has previously been linked to the immune system and

microglial dysfunction (55). A second main finding was a double dissociation in terms

of learned fear. Here we observed the opposite pattern of effects whereby *C3*[-/-] and

not *C3aR*[-/-] mice displayed enhanced fear reactivity. Although the data presented

here was in males, we can report that the same highly specific double dissociation

was also seen in females (Figures S7,S8 & S9) extending the broader applicability of

our findings. Together, these data are suggestive of distinct effects mediated by

closely related elements of the complement system that impact differentially on the

neural mechanisms underlying innate anxiety and learned fear.

Since C3a is solely produced via C3 cleavage, and C3aR is the canonical receptor

for C3a, we expected that any phenotypes dependent on the binding of C3a to C3aR

would be apparent in both knockout models. Given the divergence in phenotypes

observed, one explanation is that the *C3aR*[-/-] phenotypes are C3a independent and

instead mediated by an alternative ligand. TLQP-21, a cleavage fragment of the

neuropeptide precursor protein VGF (non-acronymic; 56; 57) was recently reported

to bind the C3aR. This peptide has pleiotropic roles including the stress

response(58) and its precursor VGF is widely expressed throughout the CNS with

high levels found in the hypothalamus, a region associated with stress reactivity and

abundant C3aR expression(59; 60). Determining whether TLQP-21/C3aR

interactions mediate the phenotypes observed will be a priority for future work.

We probed mechanisms underpinning the *C3aR*[-/-] innate anxiety phenotype by

assessing the effects of the anxiolytic drug diazepam. We found that a dose of

diazepam that was anxiolytic (i.e. reduced SAPs) in wildtype mice had no effect in

C3aR[-/-] mice. SAPs are highly sensitive to pharmacological manipulation(40; 61) and

in agreement with our own findings, diazepam has been shown to specifically

decrease SAPs in the absence of effects on open arm exploration (62). Importantly,

C3aR[-/-] mice consistently performed more SAPs than other genotypes, and therefore

floor effects cannot account for the pattern of results observed. Benzodiazepines act

on GABA$_A$ receptors(46) however we found no significant changes in expression of

Gabra2, a GABA$_A$ receptor subunit responsible for anxiolytic actions of

benzodiazepines in tests of innate anxiety(46). We did however see changes in

glucocorticoid receptor gene expression and a highly specific increase in expression

of *Cacna1c* in *C3aR*[-/-] brain tissue. The latter finding is of particular interest given the

strong evidence implicating *CACNA1C* variants in susceptibility for psychiatric

disorders including schizophrenia and bipolar disorder(49). Alterations in anxiety

have also been reported in human risk variant carriers(63) and in *Cacna1c* animal

models(50; 64). Hence, it may be speculated that since C3aR activation is known to

trigger Ca^{2+} influx(47), alterations in calcium channels may be a compensatory

mechanism resulting from *C3aR* deletion.

Another question relating to potential mechanisms was whether anxiety phenotypes

seen in the *C3aR*[-/-] mice were a result of acute, ongoing effects of *C3aR* deletion in

the adult brain or instead the consequence of neurodevelopmental impacts of *C3aR*

deficiency. At the outset, both were possibilities on the basis of previous findings

implicating C3aR in developmental neurogenesis(65) and acute changes in C3aR

expression in the adult brain after behavioural manipulations(66). The C3aR

antagonist SB290157 had no effects on behaviour in wildtype animals and hence

adult anxiety phenotypes in C3aR[-/-] mice were likely a consequence of long-term

constitutive knockout of C3aR, suggestive of developmental mechanisms. It should

be noted that SB290157 has attracted criticism due to evidence of agonist

activity(67). However, the negative results obtained in this work were not consistent

with agonist effects which would be hypothesized to reduce anxiety in the EPM.

In contrast to innate anxiety, we observed a specific effect of *C3* knockout on

conditioned fear.   Lack of comparable phenotypes in *C3aR*$^{-/-}$ and *C6*$^{-/-}$ mice

suggested that the effects were unlikely to be dependent on either C3a/C3aR or the

terminal pathway. These observations lead to several possibilities regarding the

mediating pathways. Firstly, effects could be mediated via loss of C3b/CR3

dependent synaptic elimination(10; 21), suggesting a developmental mechanism. In

support, complement mediated synaptic pruning has now been demonstrated in

brain regions relevant to fear learning such as the amygdala(68). The C3b/CR3

pathway could also be involved acutely in learning, as *C3* mRNA is upregulated

during discrete stages of fear learning(66) and microglial CR3 is implicated in long

term depression(69). Another possibility is that lack of the C5a/C5aR axis, which has

a range of developmental roles(65), mediates enhanced fear-learning phenotypes in

C3$^{-/-}$ mice, although there is less evidence for a role of C5a/C5aR in this context at

present.

While our experiments have focused on anxiety and learned fear these

endophenotypes are commonly comorbid with depression, and recent preclinical

work has suggested a protective role for C3a/C3aR in chronic-stress induced

depressive behaviour(70). Given the common co-occurrence of anxiety and

depression, our findings of enhanced anxiety in *C3aR*$^{-/-}$ mice might seem at odds

with the reported resilience of *C3aR*$^{-/-}$ to depressive phenotypes(70). However, the

chronic unpredictable-stress paradigm used by Crider et al (2018) is likely to evoke

significant inflammation, and therefore the extent to which our data in acutely

stressed animals can be compared is questionable. In summary, our findings add

significantly to the evidence that perturbations of the complement system, whether

reduced complement activation as in the present work or excessive activation as is

predicted by *C4* variants(20; 71), have major and dissociable effects on brain and

behavioral phenotypes of relevance to core clinical symptoms of psychiatric disease.

Author's contributions

LJW, TH, WPG and LSW designed the study. LJW and TH performed behavioural experiments with assistance from NH, SAB, EB and MT. ALM, SAB, EB, MT and LJW performed molecular analyses. BPM and TRH provided animals. JH, MJO, JR, WPG, NH, TRH, BPM, LSW and TH assisted in data interpretation and conceptualisation of experiments. LJW, TH, WPG and LSW drafted the manuscript. All authors approved the final manuscript.

Acknowledgements

The authors thank Professor Craig Gerard for providing the *C3aR$^{-/-}$* mice, and Rhys Perry, Helen Read and staff at JBIOS for their invaluable animal care and husbandry. This work was supported by a Wellcome Trust Integrative Neuroscience PhD Studentship awarded to LJW (099816/Z/12/Z), a Waterloo Foundation Early Career Fellowship awarded to LJW, a Hodge Centre for Neuropsychiatric Immunology Seed Corn and Project grant awarded to LJW and a Wellcome Trust Strategic Award 100202/Z/12/Z (DEFINE) held by the Neuroscience and Mental Health Research Institute at Cardiff University.

Competing financial interests

The authors declare no competing financial interests.

Materials and correspondence

All data from this study are available from the corresponding authors upon reasonable request.

References

1. Beek J, Elward K, Gasque P (2003): Activation of Complement in the Central Nervous System. *Ann N Y Acad Sci*. 992: 56–71.

2. Coulthard LG, Woodruff TM (2015): Is the Complement Activation Product C3a a Proinflammatory Molecule? Re-evaluating the Evidence and the Myth. *J Immunol*. 194: 3542–3548.

3. Morley BJ, Walport MJ (1999): *The Complement FactsBook*. Academic Press.

4. Nataf S, Stahel PF, Davoust N, Barnum SR (1999): Complement anaphylatoxin receptors on neurons: new tricks for old receptors? *Trends Neurosci*. 22: 397–402.

5. Veerhuis R, Nielsen HM, Tenner AJ (2011): Complement in the brain. *Molecular Immunology*. 48: 1592–1603.

6. Bénard M, Gonzalez BJ, Biological MSJO, 2004 (n.d.): Characterization of C3a and C5a receptors in rat cerebellar granule neurons during maturation neuroprotective effect of C5a against apoptotic cell death. *ASBMB*. doi: 10.1074/jbc.M404124200.

7. Kemper C, Köhl J (2018): Back to the future – non-canonical functions of complement. *Seminars in Immunology*. 37: 1–3.

8. Rahpeymai Y, Hietala MA, Wilhelmsson U, Fotheringham A, Davies I, Nilsson A-K, *et al.* (2006): Complement: a novel factor in basal and ischemia-induced neurogenesis. *EMBO J*. 25: 1364–1374.

9. Gorelik A, Sapir T, Haffner-Krausz R, Olender T, Woodruff TM, Reiner O (2017): Developmental activities of the complement pathway in migrating neurons. *Nature Communications*. 8: 15096.

10. Stevens B, Allen NJ, Vazquez LE, Howell GR, Christopherson KS, Nouri N, *et al.* (2007): The classical complement cascade mediates CNS synapse elimination. *Cell*. 131: 1164–1178.

11. Harold D, Abraham R, Hollingworth P, Sims R, Gerrish A, Hamshere ML, *et al.* (2009): Genome-wide association study identifies variants at CLU and PICALM associated with Alzheimer's disease. *Nature Genetics*. 41: 1088–1093.

12. Jansen IE, Savage JE, Watanabe K, Bryois J, Williams DM, Steinberg S, *et al.* (2019): Genome-wide meta-analysis identifies new loci and functional pathways influencing Alzheimer's disease risk. *Nature Genetics*. 1.

13. Shi Q, Colodner KJ, Matousek SB, Merry K, Hong S, Kenison JE, *et al.* (2015): Complement C3-Deficient Mice Fail to Display Age-Related Hippocampal Decline. *Journal of Neuroscience*. 35: 13029–13042.

14. Zhou J, Fonseca MI, Pisalyaput K, Tenner AJ (2008): Complement C3 and C4 expression in C1q sufficient and deficient mouse models of Alzheimer's disease. *Journal of Neurochemistry*. 106: 2080–2092.

15. Corbett BA, Kantor AB, Schulman H, Walker WL, Lit L, Ashwood P, *et al.* (2007): A proteomic study of serum from children with autism showing differential expression of apolipoproteins and complement proteins. *Mol Psychiatry*. 12: 292.

16. Maes M, Delange J, Ranjan R, Meltzer HY, Desnyer R (1997): Acute phase proteins in schizophrenia, mania and major depression: modulation by psychotropic drugs. *Pharmacology Biochemistry and Behaviour*. 1–11.

17. Ruland T, Chan MK, Stocki P, Grosse L, Rothermundt M, Cooper JD, *et al.* (2016): Molecular serum signature of treatment resistant depression. *Psychopharmacology*. 1–9.

18. Song YR, Wu B, Yang YT, Zhang LJ, Zhang ZW, Shi HY, *et al.* (n.d.): Specific alterations in plasma proteins during depressed, manic, and euthymic states of bipolar disorder. *Brazilian Journal of Medical and Biological Research*. 48: 973–982.

19. Oganesyan LP, Mkrtchyan GM, Sukiasyan SH, Boyajyan AS (2009): Classic and alternative complement cascades in post-traumatic stress disorder. *Bull Exp Biol Med*. 148: 859–861.

20. Sekar A, Bialas AR, de Rivera H, Davis A, Hammond TR, Kamitaki N, *et al.* (2016): Schizophrenia risk from complex variation of complement component 4. *Nature*. 1–17.

21. Schafer DP, Lehrman EK, Kautzman AG, Koyama R, Mardinly AR, Yamasaki R, *et al.* (2012): Microglia Sculpt Postnatal Neural Circuits in an Activity and Complement-Dependent Manner. *Neuron*. 74: 691–705.

22. Håvik B, Le Hellard S, Rietschel M, Lybæk H, Djurovic S, Mattheisen M, *et al.* (2011): The Complement Control-Related Genes CSMD1 and CSMD2 Associate to Schizophrenia. *BPS*. 70: 35–42.

23. Zhang S, Zhou N, Liu R, Rao W, Yang M, Cao B, *et al.* (2018): Association Between Polymorphisms of the Complement 3 Gene and Schizophrenia in a Han Chinese Population. *Cell Physiol Biochem*. 46: 2480–2486.

24. Perusini JN, Fanselow MS (2015): Neurobehavioral perspectives on the distinction between fear and anxiety. *Learn Mem*. 22: 417–425.

25. Tovote P, Fadok JP, Lüthi A (2015): Neuronal circuits for fear and anxiety. *Nat Rev*. 16: 317–331.

26. Vogelzangs N, Beekman ATF, de Jonge P, Penninx BWJH (2013): Anxiety disorders and inflammation in a large adult cohort. *Translational Psychiatry*. 3: e249–e249.

27. Girardi G, Fraser J, Lennen R, Vontell R, Jansen M, Hutchison G (2015): Imaging of activated complement using ultrasmall superparamagnetic iron oxide particles (USPIO)-conjugated vectors: An in vivo in utero non-invasive method to predict placental insufficiency and abnormal fetal brain development. *Mol Psychiatry*. 20: 1017–1026.

28. Kulkarni AP, Govender DA, Kotwal GJ, Kellaway LA (2011): Modulation of anxiety behavior by intranasally administered Vaccinia virus complement control protein and curcumin in a mouse model of Alzheimer's disease. *Current Alzheimer Research*. 8: 95–113.

29. Fischer MB, Ma M, Goerg S, Zhou X, Xia J (n.d.): Regulation of the B cell response to T-dependent antigens by classical pathway complement. *JImmunol*.

30. A role for the C3a anaphylatoxin receptor in the effector phase of asthma. (2000): A role for the C3a anaphylatoxin receptor in the effector phase of asthma. 406: 998–1001.

31. Molewijk HE, van der Poel AM, Olivier B (n.d.): The ambivalent behaviour "stretched approach posture" in the rat as a paradigm to characterize anxiolytic drugs. *Psychopharmacology*. 121: 81–90.

32. Mackintosh JH, Grant EC (1963): A Comparison of the Social Postures of Some Common Laboratory Rodents. *Behaviour*. 21: 246–259.

33. Ames RS, Lee D, Foley JJ, Jurewicz AJ, Tornetta MA, Bautsch W, *et al.* (2001): Identification of a selective nonpeptide antagonist of the anaphylatoxin C3a receptor that demonstrates antiinflammatory activity in animal models. *J Immunol*. 166: 6341–6348.

34. Dickmeis T (2008): Glucocorticoids and the circadian clock. *Journal of*
Endocrinology. 200: 3–22.
35. Livak KJ, Schmittgen TD (2001): Analysis of Relative Gene Expression Data
Using Real-Time Quantitative PCR and the 2–ΔΔCT Method. *Methods*. 25: 402–
408.
36. Wright SP (1992): Adjusted p-values for simultaneous inference. *Biometrics*. 48:
1005–1013.
37. Pellow S, Chopin P, File SE, Briley M (1985): Validation of open : closed arm
entries in an elevated plus-maze as a measure of anxiety in the rat. *J Neurosci*
Methods. 14: 149–167.
38. Walf AA, Frye CA (2007): The use of the elevated plus maze as an assay of
anxiety-related behavior in rodents. *Nat Protoc*. 2: 322–328.
39. Ennaceur A (2014): Tests of unconditioned anxiety - pitfalls and
disappointments. *Physiology & Behavior*. 135: 55–71.
40. Rodgers RJ (1997): Anxiety, defence and the elevated plus-maze. *Pharmacology*
Biochemistry and Behaviour. doi: 10.1016/S0149-7634(96)00058-9.
41. Braun AA, Skelton MR, Vorhees CV, Williams MT (2011): Comparison of the
elevated plus and elevated zero mazes in treated and untreated male Sprague-
Dawley rats: Effects of anxiolytic and anxiogenic agents. *Pharmacology*
Biochemistry and Behavior. 97: 406–415.
42. Rodgers R (1995): Factor analysis of spatiotemporal and ethological measures in
the murine elevated plus-maze test of anxiety. *Pharmacology Biochemistry and*
Behaviour. 52: 297–303.
43. Kalueff AV, Tuohimaa P (2005): The grooming analysis algorithm discriminates
between different levels of anxiety in rats: potential utility for neurobehavioural
stress research. *J Neurosci Methods*. 143: 169–177.
44. Lepicard EM, Joubert C, Hagneau I, Perez-Diaz F, Chapouthier G (2000):
Differences in anxiety-related behavior and response to diazepam in BALB/cByJ
and C57BL/6J strains of mice. *Pharmacology Biochemistry and Behavior*. 67:
739–748.
45. Albrechet-Souza L, Cristina de Carvalho M, Rodrigues Franci C, Brandão ML
(2007): Increases in plasma corticosterone and stretched-attend postures in rats
naive and previously exposed to the elevated plus-maze are sensitive to the
anxiolytic-like effects of midazolam. *Hormones and Behavior*. 52: 267–273.
46. Smith KS, Engin E, Meloni EG, Rudolph U (2012): Benzodiazepine-induced
anxiolysis and reduction of conditioned fear are mediated by distinct GABAA
receptor subtypes in mice. *Neuropharmacology*. 63: 250–258.
47. Sayah S, Jauneau AC, Patte C, Tonon MC, Vaudry H, Fontaine M (2003): Two
different transduction pathways are activated by C3a and C5a anaphylatoxins on
astrocytes. *Molecular Brain Research*. 112: 53–60.
48. Lian H, Li Y, Lu H-C, Zheng H (2015): NFκB-Activated Astroglial Release of
Complement C3 Compromises Neuronal Morphology and Function Associated
with Alzheimer's Disease. 85: 101–115.
49. Hamshere ML, Walters JTR, Smith R, Richards AL, Green E, Grozeva D, *et al.*
(2019): Genome-wide significant associations in schizophrenia to ITIH3/4,
CACNA1C and SDCCAG8, and extensive replication of associations reported by
the Schizophrenia PGC. *Mol Psychiatry*. 1–5.
50. Kabir ZD, Che A, Fischer DK, Rice RC, Rizzo BK, Byrne M, *et al.* (2017): Rescue
of impaired sociability and anxiety-like behavior in adult *cacna1c*-deficient mice
by pharmacologically targeting eIF2α. *Mol Psychiatry*. 22: 1096–1109.

51. Risbrough VB, Geyer MA (2005): Anxiogenic treatments do not increase fear-potentiated startle in mice. *BPS*. 57: 33–43.

52. Falls WA, Carlson S, Turner JG, Willott JF (1997): Fear-potentiated startle in two strains of inbred mice. *Behavioral Neuroscience*. 111: 855–861.

53. Davis M (2006): Neural systems involved in fear and anxiety measured with fear-potentiated startle. *Am Psychol*. 61: 741–756.

54. Cryan JF, Holmes A (2005): The ascent of mouse: advances in modelling human depression and anxiety. *Nature Reviews Drug Discovery*. 4: 775–790.

55. Chen S-K, Tvrdik P, Peden E, Cho S, Wu S, Spangrude G, Capecchi MR (2010): Hematopoietic origin of pathological grooming in Hoxb8 mutant mice. *Cell*. 141: 775–785.

56. Hannedouche S, Beck V, Leighton-Davies J, Beibel M, Roma G, Oakeley EJ, *et al.* (2013): Identification of the C3a Receptor (C3AR1) as the Target of the VGF-derived Peptide TLQP-21 in Rodent Cells. *Journal of Biological Chemistry*. 288: 27434–27443.

57. Cero C, Vostrikov VV, Verardi R, Severini C, Gopinath T, Braun PD, *et al.* (2014): The TLQP-21 peptide activates the G-protein-coupled receptor C3aR1 via a folding-upon-binding mechanism. *Structure*. 22: 1744–1753.

58. Razzoli M, Bo E, Pascucci T, Pavone F, Amato FRD, Cero C, *et al.* (2012): Implication of the VGF-derived peptide TLQP-21 in mouse acute and chronic stress responses. *Behav Brain Res*. 229: 333–339.

59. Lewis B, Francis K, Gasque P, Scanlon M, Ham J (2002): Functional complement C3a receptors in the rat pituitary gland. *Endocrine Abstracts*. 3: 198.

60. Francis K (2003): Complement C3a receptors in the pituitary gland: a novel pathway by which an innate immune molecule releases hormones involved in the control of inflammation. *The FASEB Journal*. 1–4.

61. Holly KS, Orndorff CO, Murray TA (2016): MATSAP: An automated analysis of stretch-attend posture in rodent behavioral experiments. *Scientific Reports*. 6. doi: 10.1038/srep31286.

62. Genewsky A, Albrecht N, Bura SA, Kaplick PM, Heinz DE, Nußbaumer M, *et al.* (2019): How much fear is in anxiety? 1–42.

63. Roussos P, Giakoumaki SG, Georgakopoulos A, Robakis NK, Bitsios P (2011): The CACNA1C and ANK3 risk alleles impact on affective personality traits and startle reactivity but not on cognition or gating in healthy males. *Bipolar Disorders*. 13: 250–259.

64. Lee AS, Ra S, Rajadhyaksha AM, Britt JK, De Jesus-Cortes H, Gonzales KL, *et al.* (2012): Forebrain elimination of *cacna1c* mediates anxiety-like behavior in mice. *Mol Psychiatry*. 17: 1054–1055.

65. Coulthard LG, Hawksworth OA, Woodruff TM (2018): Complement: The Emerging Architect of the Developing Brain. *Trends Neurosci*. 41: 373–384.

66. Scholz B, Doidge AN, Barnes P, Hall J, Wilkinson LS, Thomas KL (2016): The Regulation of Cytokine Networks in Hippocampal CA1 Differentiates Extinction from Those Required for the Maintenance of Contextual Fear Memory after Recall. (K. Felmingham, editor) *PLoS ONE*. 11: e0153102–29.

67. Therien AG, Baelder R, Köhl J (2005): Agonist Activity of the Small Molecule C3aR Ligand SB 290157. *J Immunol*. 174: 7479–7480.

68. Kopec AM, Smith CJ, Ayre NR, Sweat SC, Bilbo SD (2018): Microglial dopamine receptor elimination defines sex-specific nucleus accumbens development and social behavior in adolescent rats. *Nature Communications*. 9: 3769.

69. Zhang J, Malik A, Choi HB, Ko RWY, Dissing-Olesen L, MacVicar BA (2014):
Microglial CR3 Activation Triggers Long-Term Synaptic Depression in the
Hippocampus via NADPH Oxidase. *Neuron*. 82: 195–207.
70. Crider A, Feng T, Pandya CD, Davis T, Nair A, Ahmed AO, *et al.* (2018):
Complement component 3a receptor deficiency attenuates chronic stress-
induced monocyte infiltration and depressive-like behavior. *Brain, Behavior, and*
Immunity. 70: 246–256.
71. Sellgren CM, Gracias J, Watmuff B, Biag JD, Thanos JM, Whittredge PB, *et al.*
(2019): Increased synapse elimination by microglia in schizophrenia patient-
derived models of synaptic pruning. *NatNeurosci*. 1.

699 **Figures and legends**

Elevated plus maze (EPM)

Figure 1. *C3aR⁻/⁻*, but not *C3⁻/⁻* mice show increased anxiety-like behaviour in

the elevated plus maze (EPM;A-E) and elevated zero maze (EZM;F-J). (A)

Heatmaps displaying relative time per zone of the EPM across genotypes **(B)** There

was a significant difference between genotypes in the latency to first open arm visit

(wildtype 8.21±1.53s, *C3*$^{/-}$ 14.1±2.52s, *C3aR*$^{/-}$ 27.6±8.31s, H_2=10.5, p=0.005). *Post*

hoc tests demonstrated that *C3aR*$^{/-}$ mice took significantly longer to first enter the

open arms than wildtype mice (p=0.0045). **(C)** *C3aR*$^{/-}$ mice distributed their time

across the EPM differently to wildtype and *C3*$^{/-}$ mice (GENOTYPE×ZONE,

$F_{4,62}$=17.7, p=0.0001) spending less time in the open arms (*C3aR*$^{/-}$ 16.70±3.73s vs.

wildtype 75.78±8.86s, p<0.0001, *C3aR*$^{/-}$ vs. *C3*$^{/-}$ 93.86±9.59s p<0.0001) and

significantly more time in the closed arms (*C3aR*$^{/-}$ 221.88±12.06s vs. wildtype

146.01±7.01s, p<0.0001, and *C3*$^{/-}$ 140.04±8.61s p<0.0001). **(D)** *C3aR*$^{/-}$ (14.90±2.22)

mice performed significantly fewer head dips than wildtype (37.92±2.53, p<0.0001)

and *C3*$^{/-}$ mice (49.17±2.37, p<0.0001), whereas *C3*$^{/-}$ mice performed significantly

more head dips than wildtype mice (p=0.0061; overall ANOVA $F_{2,31}$=48.0, p<0.0001).

(E) *C3aR*$^{/-}$ mice performed significantly more stretch attend postures (SAPs;

15.30±1.80) than wildtype (7.58±1.66, p=0.0042) and *C3*$^{/-}$ mice (5.67±0.94,

p=0.0004; overall ANOVA $F_{2,31}$=10.3, p=0.0004). **(F)** Heatmaps displaying relative

exploration of the open segments of the elevated zero maze, across genotypes.

Note that due to the height of the walls in the closed regions it was not possible to

track mice or observe ethological behaviours such as grooming or SAPs. **(G)** There

was a significant difference in the latency to first enter the open arms (wildtype

101.00±31.00s, *C3*$^{/-}$ 42.00±2.52s, *C3aR*$^{/-}$ 204.00±40.40s, H_2=8.13, p=0.0171). *Post*

hoc tests revealed that *C3aR*$^{/-}$ mice took significantly longer than *C3*$^{/-}$ mice to

initially enter the open region (p=0.0140). **(H)** The number of entries made to open

regions differed between genotypes (wildtype 7.69±1.69, *C3*$^{/-}$ 11.5±1.43s, *C3aR*$^{/-}$

2.10±1.15, $F_{2,30}$=8.96, p=0.0009). *C3aR*$^{/-}$ mice made significantly fewer entries to the

open areas than wildtype (p=0.0324) and *C3*$^{/-}$ mice (p=0.0006) and **(I)** spent

significantly less time on the open arms (1.77±1.29) compared to wildtype

(35.7±8.43s, p=0.0132) and $C3^{-/-}$ (57.7±9.32s, p<0.0001; overall Kruskal-Wallis test

H_2=19.2, p<0.0001).   **(J)** $C3^{-/-}$ mice performed significantly more head dips

(20.7±1.62) than wildtype (13.9±1.89, p=0.048) and $C3aR^{-/-}$ mice (10.3±2.42,

p=0.0034; overall ANOVA $F_{2,27}$=6.86,p=0.0039). Data are mean ± S.E.M. *, **, ***

and **** represent p≤0.05, p≤0.01, p≤0.001 and p≤0.0001 for *post-hoc* genotype

comparisons, respectively.

Figure 2. Excessive grooming in *C3aR^/-* mice (A) There were significant differences in the frequency of grooming bouts during the 5-minute elevated plus maze (EPM) test (wildtype 2.75±0.55, *C3aR^/-* 5.20±0.72, *C3^/-* 3.17±0.64, $F_{2,31}$=3.94, p=0.0298). *C3aR^/-* mice engaged in grooming significantly more frequently than wildtype mice (p=0.0320). **(B)** There was also a trend towards a greater duration of time spent grooming in *C3aR^/-* mice on the EPM, however this did not reach significance ($F_{2,31}$=2.59, p=0.0909, wildtype 8.11±1.61s, *C3aR^/-* 13.8±2.10, *C3^/-* 9.70±1.63s). **(C)** Representative images of the effects of chronic homecage excessive grooming in *C3aR^/-* mice resulting in bald patches. By 5 months of age, 80% of *C3aR^/-* mice displayed such signs of excessive grooming. In contrast, no wildtype or *C3^/-* mice demonstrated signs of excessive grooming. This phenotype was observed in all cohorts of *C3aR^/-* mice assessed. Data are mean + S.E.M. * represents p≤0.05 for *post-hoc* genotype comparisons.

Figure 3. Altered sensitivity to diazepam in *C3aR*^{-/-} and *C3*^{-/-} mice. Behaviourally

naïve mice were treated with either diazepam (2mg/kg, i.p) or vehicle injections once

daily for 2 days and then 30 minutes prior to testing. (**A)** Heatmaps demonstrating

duration spent in zones of the maze by vehicle treated and diazepam treated

animals **(B)** Plots showing duration spent on open arms in 1-minute time bins (start-

01:00 was excluded due to effect of diazepam in delaying initial entry to open arms,

see Supplementary Figure 4a). There was a trend for wildtype diazepam treated

animals to spend more time on the open arms throughout the task although this did

not reach significance (main effect of DRUG, $F_{1,38}$=1.41, p=0.2462). In *C3*^{-/-} mice

there was a strong tendency for drug treated animals to explore the open arms less

than vehicle treated *C3$^{-/-}$* mice (main effect of DRUG, F$_{1,22}$=1.25, p=0.2764). A

similar, though less pronounced pattern was seen in *C3aR$^{-/-}$* mice (main effect of

DRUG, F$_{1,19}$=1.55, p=0.2284) **(C)** There were genotype differences in SAPs (main

effect of GENOTYPE, F$_{2,79}$=10.7, p<0.0001), a main effect of DRUG (F$_{1,79}$=4.13,

p=0.0454) and a significant GENOTYPE $\times$ DRUG interaction (F$_{2,79}$=4.64, p=0.0138).

Post hoc tests showed that diazepam significantly reduced the number of SAPs in

wildtype mice only (wildtype vehicle 9.00$\pm$1.44 vs. wildtype diazepam 3.09$\pm$0.71,

p=0.0006, *C3$^{-/-}$* vehicle 4.67$\pm$1.24 vs. *C3$^{-/-}$* diazepam 5.00$\pm$1.51, p=0.9975, *C3aR$^{-/-}$*

vehicle 11.45$\pm$1.49 vs. *C3aR$^{-/-}$* diazepam 10.50$\pm$1.34, p=0.9558). Data are mean +

S.E.M. *, **, and *** represent p$\leq$0.05, p$\leq$0.01 and p$\leq$0.001 for *post-hoc* genotype

comparisons, respectively.

Figure 4. Elevated corticosterone response to the elevated plus maze in *C3aR$^{-/-}$*

mice. Exposure to the EPM significantly elevated corticosterone in all genotypes

(main effect of CONDITION, $F_{1,54}$=143, p<0.0001; baseline 344.66±63.70 vs. EPM

3553.84±274.13). There was a significant GENOTYPE × CONDITION interaction

($F_{2,54}$=4.64, p=0.0138). *Post hoc* analysis showed that after the EPM, *C3aR$^{-/-}$* mice

demonstrated significantly higher corticosterone levels (4640.27±376.13) than

wildtype (3033.78±535.06, p=0.0127) and *C3$^{-/-}$* mice (2948.00±374.87, p=0.0032).

Post hoc tests also indicated that there were no baseline differences between

genotypes (wildtype 216.54±63.2 vs. *C3aR$^{-/-}$* 316.17±111.60 p>0.9999, wildtype vs.

C3$^{-/-}$ p=0.9927, and *C3$^{-/-}$* 485.60±130.35 vs.*C3aR$^{-/-}$* mice p=0.9992). Data are mean

+ S.E.M. *, **, and **** represent p≤0.05, p≤0.01 and p≤0.0001 for *post-hoc*

genotype comparisons, respectively.

Figure 5. mRNA expression of stress and anxiety related genes. The medial prefrontal cortex (mPFC), ventral hippocampus (vHPC) and cerebellum were dissected from behaviourally naïve mice and were subjected to quantitative PCR analysis for the transcript expression of candidate genes associated with anxiety and the stress response. **(A)** There were trends towards increased expression of the glucocorticoid receptor gene *Nr3c1* in the mPFC and vHPC of *C3aR*$^{-/-}$ mice but these were not significant (mPFC; $F_{2,56}$=1.33, p=0.2723, N wildtype=20, *C3*$^{-/-}$=20, *C3aR*$^{-/-}$= 19 vHPC; $F_{2,62}$=1.11, p=0.3345, N wildtype=23, *C3*$^{-/-}$=20, *C3aR*$^{-/-}$= 22). Expression of the glucocorticoid receptor (*Nr3c1*) was significantly increased in the cerebellum of both *C3*$^{-/-}$ and *C3aR*$^{-/-}$ groups however ($F_{2,61}$=10.3, p=0.0002, *C3*$^{-/-}$ vs. wildtype p=0.0023, *C3aR*$^{-/-}$ vs. wildtype p=0.0002, N wildtype=19, *C3*$^{-/-}$=20, *C3aR*$^{-/-}$= 15). **(B)** There were no significant differences in the expression of Corticotrophin releasing

hormone receptor 1 (*Crhr1*) in any region, across genotypes (mPFC $F_{2,53}$=0.587,

p=0.5597, N wildtype=20, *C3*$^{-/-}$=17, *C3aR*$^{-/-}$= 19; vHPC $F_{2,49}$=0.169, p=0.8450, N

wildtype=20, *C3*$^{-/-}$=15, *C3aR*$^{-/-}$= 17; cerebellum $F_{2,47}$=0.0482, p=0.8346, N

wildtype=19, *C3*$^{-/-}$=17, *C3aR*$^{-/-}$= 14). **(C)** There were also no significant changes in

the expression of the GABA$_A$ receptor α2 subunit (*Gabra2*) in any region, across

genotypes (mPFC H_2=1.04, p=0.5939, N wildtype=20, *C3*$^{-/-}$=19, *C3aR*$^{-/-}$= 16; vHPC

$F_{2,49}$=0.721, p=0.4914, N wildtype=20, *C3*$^{-/-}$=13, *C3aR*$^{-/-}$= 19; cerebellum $F_{2,47}$=0.221,

p=0.8026, N wildtype=18, *C3*$^{-/-}$=17, *C3aR*$^{-/-}$= 15). **(D)** Expression levels of the

Calcium voltage-gated channel subunit Alpha 1c (*Cacna1c*) were significantly

increased in *C3aR*$^{-/-}$ mice in a regionally specific manner in the vHPC ($F_{2,47}$=3.20,

p=0.0496, *C3*$^{-/-}$ vs. wildtype p=0.6895, *C3aR*$^{-/-}$ vs. wildtype p=0.0295, N wildtype=21,

C3$^{-/-}$=13, *C3aR*$^{-/-}$= 16) and the cerebellum ($F_{2,54}$=5.84, p=0.0051, *C3*$^{-/-}$ vs. wildtype

p=0.1613, *C3aR*$^{-/-}$ vs. wildtype p=0.0024, N wildtype=20, *C3*$^{-/-}$=20, *C3aR*$^{-/-}$= 17).

There were no significant changes in the mPFC ($F_{2,52}$=1.04, p=0.5939, N

wildtype=20, *C3*$^{-/-}$=19, *C3aR*$^{-/-}$= 16). Data represent fold change from wildtype +

SEM.

Figure 6. Semi-chronic pharmacological C3aR blockade in wildtype mice does not recapitulate *C3aR^-/-* phenotypes in the elevated plus maze. SB290157 (C3aRA) was administered twice daily for five days (i.p. 10mg/kg). **(A)** Heatmaps demonstrating duration spent in zones of the maze **(B)** There was a trend for greater latencies to first open arm visit in *C3aR^-/-* mice (26.06±15.73s vs. wildtype vehicle 8.75±2.50s, wildtype C3aRA 13.38±6.70s) approaching significance ($F_{2,18}$=2.73, p=0.0923). **(C)** *C3aR^-/-* mice again distributed their time across the EPM differently to wildtype-vehicle and wildtype-C3aRA mice (GENOTYPE×ZONE, $F_{4,54}$=5.67, p=0.0007) spending less time in the open arms (*post hoc:* wildtype-vehicle 55.39±12.11 vs. *C3aR^-/-* 12.85±5.69s p=0.0157, wildtype-C3aRA 67.14±10.86 vs. *C3aR^-/-* p=0.0007), and significantly more time in the closed arms than wildtype-C3aRA mice (*C3aR^-/-* 242.21±37.72 vs. wildtype-C3aRA 207.18±14.55 p=0.0363, wildtype-vehicle 214.99±13.97s vs. *C3aR^-/-* p=0.1667). **(D)** *C3aR^-/-* mice made fewer head dips (8.57±5.50) than wildtype-vehicle (21.67±2.62, p=0.0057) and wildtype-

C3aRA mice (25.63 ± 3.04, p=0.0002, overall ANOVA $F_{2,18}$=13.4, P = 0.0003) and **(F)**

displayed more SAPs (17.29 ± 4.72) than wildtype-vehicle (13.22 ± 5.19, p=0.0853)

and wildtype-C3aRA treated mice (7.45 ± 2.28, p=0.0018, overall ANOVA $F_{2,18}$=8.50,

P = 0.0025). For all measures, there were no significant differences between

wildtype-vehicle and wildtype-C3aRA mice. Data are mean + S.E.M. *, **, *** and

**** represent p$\leq$0.05, p$\leq$0.01, p$\leq$0.001 and p$\leq$0.0001 for *post-hoc* genotype

comparisons, respectively.

Figure 7. Enhanced fear-potentiated startle in *C3*[^-/-] but not *C3aR*[^-/-] mice. (A) Flow

chart depicting the FPS protocol used, which took place in three separate sessions

over two consecutive days. Baseline startle reactivity to a range of pulse intensities

was assessed in the pre-conditioning session, immediately preceding the

conditioning session in which a visual stimulus (light) was paired with 3 weak foot

shocks. 24 hours later, subjects were re-introduced to the same chamber and startle

reactivity was compared between pulse-alone trials and pulse+CS trials to determine

the degree of FPS. On all trials, the peak startle response was recorded and

normalised for body weight differences using Kleiber's 0.75 mass exponent[74], and

fold-changes calculated. **(B)** There was a significant main effect of GENOTYPE

($F_{2,66}$=9.04, p=0.0003) and a significant GENOTYPE $\times$ STIMULUS INTENSITY

interaction ($F_{4,132}$=7.55, p<0.0001).

C3$^{-/-}$ and *C3aR*$^{-/-}$ mice demonstrated increased levels of startle responding relative to

wildtype mice at 100dB (*C3*$^{-/-}$ 11.34±1.51 vs. wildtype 2.63±0.26, p<0.0001, *C3aR*$^{-/-}$

9.12±2.63 vs. wildtype p=0.0174) and 110dB (*C3*$^{-/-}$ 12.69±1.55 vs. wildtype

3.74±0.50, p<0.0001, *C3aR*$^{-/-}$ 10.58±3.58 vs. wildtype p=0.0111) **(C)** *C3*$^{-/-}$ and *C3aR*

$^{-/-}$ mice also showed increased startle responding to the footshock+CS (*C3*$^{-/-}$

75.18±5.73, *C3aR*$^{-/-}$ 73.14±7.78) pairings relative to wildtype mice (44.34±5.29, *C3*$^{-/-}$

vs. wildtype p=0.0006, *C3aR*$^{-/-}$ vs. wildtype p=0.0137, overall ANOVA F$_{2,66}$=8.7,

p=0.0004), although it should be noted that responses were much greater to these

stimuli in all mice than to the startle stimuli in the pre-conditioning session. **(D)** In the

post-conditioning session, all mice demonstrated increases to the pulse+CS stimuli

in comparison to pulse-alone stimuli, as demonstrated by the fold-change increase in

startle responding, however, this effect was significantly increased in *C3*$^{-/-}$ mice

(3.72±0.27) relative to wildtype (1.99±0.11, p<0.0001) and *C3aR*$^{-/-}$ mice (2.27±0.24,

p=0.0056, overall Kruskal-Wallis test H$_2$=27.7, p<0.0001). Data are mean + S.E.M. *,

, * and **** represent p≤0.05, p≤0.01, p≤0.001 and p≤0.0001 for *post-hoc*

genotype comparisons, respectively.

Figure 8. Enhanced fear-potentiated startle in *C3$^{-/-}$* but not *C6$^{-/-}$* mice. A separate cohort of wildtype and *C3$^{-/-}$* mice were assessed alongside *C6$^{-/-}$* mice. **(A)** There was a significant main effect of GENOTYPE ($F_{2,20}$=5.87, p=0.0098). *C3$^{-/-}$* mice (90 db: 4.98±0.79; 100 db: 7.24±0.94; 110 db: 6.97±0.84) demonstrated elevated startle responding at all intensities relative to wildtype (90 db: 2.55±0.30 p=0.1846; 100 db: 2.69±0.24 p=0.0042; 110 db: 3.74±0.51 p=0.0545) and *C6$^{-/-}$* mice (90 db: 2.68±0.31 p=0.2023; 100 db: 3.19±0.48 p=0.0098; 110 db: 3.03±0.544 p=0.0123). **(B)** *C3$^{-/-}$* mice demonstrated a trend towards greater startle reactivity (78.35±10.25) to the footshock+CS pairings during the conditioning phase of training (wildtype 38.50±5.41, *C6$^{-/-}$* 43.49±7.57, H_2=4.66, p=0.0971). **(C)** Post-conditioning, *C3$^{-/-}$* mice showed the greatest fold change (2.48±0.12) in startle responding between pulse-alone stimuli and pulse+CS stimuli, which was significantly greater than that shown by wildtype (1.70±0.08, p=0.0094) and *C6$^{-/-}$* mice (1.60±0.15, p=0.0025, overall ANOVA $F_{2,29}$=8.08, p=0.0016). There were no differences in responding between *C6$^{-/-}$* and wildtype mice on any measures. Data are mean + S.E.M. * and ** represent p≤0.05 and p≤0.01 for *post-hoc* genotype comparisons, respectively.