Different complement immune pathways mediate innate and learned aversion

Laura J. Westacott[1,4†], Trevor Humby[1,2†], Niels Haan[1,4], Sophie A. Brain[2], Emma-Louise Bush[2], Margarita Toneva[2], Andreea-Ingrid Baloc[2], Nicola Lander[7], Anna L. Moon[1], Jack Reddaway[1,4], Michael J. Owen[1,] Jeremy Hall[1,4], Timothy R. Hughes[3,4], B. Paul Morgan[3,4,6], William P. Gray[1,4,5$] & Lawrence S. Wilkinson[1,2,4*$]

[1] Neuroscience and Mental Health Research Institute, MRC Centre for Neuropsychiatric Genetic and Genomics, School of Medicine, Hadyn Ellis Building, Cardiff University, Cardiff, CF24 4HQ, UK. [2] Behavioural Genetics Group, Schools of Psychology and Medicine, Cardiff University, Cardiff, CF10 3AT, UK. [3] Complement Biology Group, Systems Immunity Research Institute, School of Medicine, Cardiff University, CF14 4XW, Cardiff UK. [4] Hodge Centre for Neuropsychiatric Immunology, School of Medicine, Cardiff University, Cardiff CF24 4HQ, UK. [5] Brain Repair and Intracranial Therapeutics (BRAIN) Unit, School of Medicine, Cardiff University, CF24 4HQ, UK. [6] UK Dementia Research Institute, Cardiff University, Cardiff, CF24 4HQ, UK. [7] School of Biosciences, Cardiff University, Cardiff, CF10 3AT, UK.

[†] These authors contributed equally to the work
[$] Joint senior authors: Lawrence S. Wilkinson & William P. Gray

[*] Corresponding author: Lawrence S. Wilkinson

Email address: WilkinsonL@cardiff.ac.uk

Address: Neuroscience and Mental Health Research Institute, School of Medicine, Cardiff University, Cardiff, CF24 4HQ, UK

Key words: *Neuroimmunology, Behaviour, Complement system, Anxiety, Fear, Stress response.*

Abstract

Complement is a key component of the immune system. Here we show that

complement pathways with canonical roles in inflammation and host defence also

impact on emotional responses to aversive events in a highly selective manner.

Using mouse models deficient in a panel of complement proteins and focusing on

the central role of C3 in complement signalling, we first demonstrated that *C3aR*

deletion markedly increased physiological and behavioural responses to innate

anxiety-provoking stimuli. These effects, which were resistant to benzodiazepines,

drugs used in the treatment of anxiety, were independent of C3a the ligand for

C3aR, indicating the existence of an alternative C3aR ligand mediating anxiety

responses. In contrast, manipulation of C3aR had no influence on learned fear,

where a previously neutral cue generates a fear response as a result of predicting an

aversive outcome. Instead increases in learned fear were found only as a

consequence of *C3* deletion and subsequent downstream disruption to iC3b/CR3

signalling, a complement pathway previously associated with synaptic plasticity via

effects on synaptic elimination. Our findings indicate an unanticipated complexity and

subtlety to the way in which particular complement proteins, acting through distinct

signalling pathways, can influence emotional dysfunction of relevance to many

common psychopathologies, as such they add evidence to the emerging links

between complement and risk for psychiatric disorder.

The complement system is a key component of the immune system that plays a

pivotal role in inflammation and host-defence. Complement activation occurs via

several pathways, all of which lead to cleavage of the central protein, C3 (see Figure

S1). Activation of C3 generates the fragments C3a and C3b. C3a is an

anaphylatoxin that signals via its canonical G-protein coupled receptor C3aR[1].

Activation of this receptor has been demonstrated to trigger calcium mobilization[2-4],

stimulating an array of intracellular signalling pathways to induce both pro- and anti-

inflammatory effects[1,5]. C3b on the other hand propagates further complement

activation by contributing to the cleavage of complement component 5 (C5)

downstream of C3 and, after further cleavage to iC3b, plays a role in opsonisation by

macrophages and microglia via complement receptor 3 (CR3). Akin to C3, C5

cleavage generates C5a (another anaphylatoxin and a ligand for the C5a receptor,

C5aR) and C5b, which triggers the terminal complement pathway by sequentially

binding proteins C6, C7, C8 and C9. These proteins subsequently congregate to

assemble the membrane attack complex (MAC) which ultimately results in

destruction of the target cell or pathogen via cell lysis[6].

In the central nervous system evidence is emerging that complement has functions

beyond its canonical immune roles[7]. Neurons, astrocytes and microglia express

complement receptors and regulators, and are also capable of synthesising

complement proteins[8,9]. The expression patterns of these vary over the course of

brain development[10]. Complement impacts a number of neurodevelopmental

processes including neurogenesis[11], migration[12] and synaptic pruning[13] as well as

ongoing synaptic plasticity processes underlying learning and memory in the adult

brain[14]. Furthermore, there is increasing evidence that complement is causally

involved in the pathogenesis of neurodegenerative and psychiatric conditions. In

Alzheimer's disease, genetic variants in complement related loci have been

associated with increased disease risk[15,16], and complement knockout mice exhibit

reduced age-related synapse loss[17] and neuropathology[18]. Alterations in

complement proteins and activation have also been reported in sera from individuals

with autism-spectrum disorder[19] schizophrenia[20], major depressive disorder[21],

bipolar disorder[22] and post-traumatic stress disorder[23].

In the case of schizophrenia, an important finding comes from elegant genetic work

demonstrating that structural variation in the *C4* locus is associated with risk of

developing the disease[24]. C4 cleavage generates fragments that contribute to the

activation of C3, yielding C3a and C3b. Given the known roles for the iC3b/CR3

pathway in developmental synaptic pruning[13,25], it has been suggested that *C4*

variants may impact on psychiatric risk via this mechanism, with excessive synaptic

elimination leading to abnormal connectivity and disruption of neural networks[24].

Variants in *C3* and putative complement-control genes *CSMD1* and *CSMD2* have

also been implicated in genetic susceptibility for schizophrenia[26,27].

Altered emotional function, in particular maladaptive anxiety and fear, is a pervasive

and clinically important symptom that is comorbid across several of the DSM-5 and

ICD-11 defined disorders in which complement has been implicated. Anxiety and fear

are distinct aversive states elicited by differing environmental factors, that result in

distinguishable behavioural outputs[28]. Anxiety is characterised by sustained arousal,

hypervigilance and risk assessment surrounding anticipated or potential threats,

while fear is an acute response to an experienced, imminent danger resulting in

immediate avoidance, fight or freezing behaviour[29,30]. Whilst there is significant

overlap in the neurocircuitry underlying these states, there are also contributions

from distinct neuronal circuitries[28,31].   Although links between anxiety and

inflammation have been documented[32], there has been little investigation of the

specific role of complement in either anxiety or fear. In animal models, anxiety

phenotypes arise from heightened pre-natal complement activity[33] and complement

inhibitors reduce anxiety associated with neurodegeneration[34]. Furthermore, aged

C3 deficient mice exhibited lower levels of anxiety alongside enhanced contextual

fear responses[17]. These latter studies suggest that complement can influence both

innate and learned components of aversive behaviours, however, the precise

relationship between signalling within discrete pathways of the complement system

and their relationship to dissociable aspects of emotionality and underlying brain

function is unknown.

Utilising the central role of C3 in complement signalling, we used a combination of

complement knockout mice to functionally parse anxiety and fear related

phenotypes. In homozygous *C3* knockout mice ($C3^{-/-}$)[35] complement cannot be

activated beyond C3, and therefore these animals lack C3 activation fragments

(C3a, C3b) and downstream activation products (C5a, C5b) and thus cannot activate

the terminal complement pathway. Phenotypes in this model could therefore be the

result of loss of any of these downstream effector molecules. We compared the $C3^{-/-}$

model with homozygous *C3aR* knockout mice ($C3aR^{-/-}$)[36].   In these mice,

complement is intact apart from the capacity for C3a to bind its canonical receptor

C3aR and hence we tested the extent to which phenotypic effects were the result

specifically of disrupted C3a/C3aR signalling. A priori, because C3a is an obligate

cleavage fragment of C3, we hypothesised that any phenotypes dependent on the

binding of C3a to C3aR would be apparent in both $C3^{-/-}$ and $C3aR^{-/-}$ models.

Contrary to our hypotheses we found clear phenotypic dissociations between the $C3^{-}$

$^{/-}$ and $C3aR^{-/-}$ models which were dependent on the nature of the emotional

behaviour. $C3aR^{-/-}$ mice displayed a profound innate anxiety phenotype that was

lacking in the $C3^{-/-}$ model. In contrast, when we examined learned fear, where a

previously neutral cue generates a fear response as a result of predicting an

aversive outcome, we found that the dissociation was reversed with the $C3^{-/-}$ mice

exhibiting an enhanced fear response to a conditioned cue, but no differences

between wildtype and $C3aR^{-/-}$ mice. This observed double dissociation in emotionality

was linked to distinct underlying complement signalling mechanisms, with anxiety

phenotypes in C3aR mice likely to be independent of C3a/C3aR signalling raising

the possibility of an alternative ligand for C3aR, and $C3^{-/-}$ effects on learned fear the

result of alterations in iC3b-CR3 pathway activity. These findings point to a hitherto

unrecognized complexity of complement effects on brain function and behaviour of

relevance to emotional dysfunction in psychopathology.

Results

Increased innate anxiety in $C3aR^{-/-}$ but not $C3^{-/-}$ mice

Full details of the complement knockout models used can be found in the Methods.

Using a cohort of male wildtype, $C3^{-/-}$ and $C3aR^{-/-}$ mice we first assessed emotional

reactivity in the elevated plus maze (EPM), a well-established test of innate anxiety

in rodents which exploits the conflict between the drive to explore novel

environments and the innate aversion towards open, brightly lit spaces[37,38].

Heatmaps indexing overall maze exploration over a 5-minute session demonstrated

major differences in open arm exploration between genotypes (Figure 1A; see

Supplementary Video 1 for representative examples). Notably, in comparison to

wildtype and $C3^{-/-}$ mice, $C3aR^{-/-}$ mice took significantly longer to first enter the open

arms (Figure 1B) and spent less time on the open arm per entry (Figure S2A),

leading to a reduced overall duration spent in the open arms (Figure 1C), findings

consistent with increased anxiety. The ethological parameters head dips and stretch

attend postures (SAPs) also differed between genotypes (Figure 1D,E), with $C3aR^{-/-}$

mice exhibiting decreases in the former and increases in the latter, a pattern of

results again consistent with heightened anxiety[39]. We also noted a significantly

increased frequency of head dipping in $C3^{-/-}$ mice (Figure 1D), consistent with

reduced levels of anxiety relative to both wildtype and $C3aR^{-/-}$ mice[40].

These initial data were consistent with an anxiogenic phenotype present in $C3aR^{-/-}$

mice but absent in $C3^{-/-}$ mice. We confirmed the findings in two further independent

tests of anxiety using additional cohorts of animals. First we used the elevated zero

maze (EZM, see Methods), another test of anxiety-like behaviour which similarly

probes behavioural responses to exposed, illuminated spaces[41]. The data

recapitulated the pattern of findings seen in the EPM (Figure 1F-J). Additional data

from the open field test, where $C3aR^{-/-}$ mice were more likely to avoid the centre of

the arena, were also consistent in demonstrating a specific anxiety-like phenotype in

$C3aR^{-/-}$ but not $C3^{-/-}$ mice (Figure S3). Given that several of the measures indexing

anxiety were dependent on movement around the apparatus it was important to

eliminate potential locomotor confounds. To address this issue, we measured

activity independently in a non-anxiety provoking environment and found no

differences in locomotor activity between genotypes (Figure S2C), demonstrating

that anxiety measures were not influenced by movement confounds.

Neuroendocrine response and neuronal activation in *C3aR*$^{-/-}$ and *C3*$^{-/-}$ mice

following exposure to the elevated plus maze

We next tested whether the behavioural measures of anxiety were paralleled by

changes in plasma levels of the stress hormone corticosterone. In a separate cohort

of wildtype, *C3*$^{-/-}$ and *C3aR*$^{-/-}$ mice, we assayed plasma corticosterone 30 minutes

after exposure to the EPM and compared levels to those of a group of animals who

remained in their home-cages.   There were no genotype differences in basal

corticosterone levels; however, being placed on the EPM increased plasma

corticosterone 6-15-fold in all genotypes, demonstrating that the EPM was a potent

stressor (Figure 2A). *Post hoc* analyses showed a significantly greater EPM-evoked

corticosterone response in the *C3aR*$^{-/-}$ animals, consistent with their increased

anxiety-like behaviour observed on the maze.

Neuronal activity within the ventral hippocampus (vHPC) has been linked to anxiety

in rodents[42,43] therefore we also investigated mRNA expression of two immediate

early genes (IEGs), *c-Fos* and *zif268*, in a further cohort of behaviourally naïve

wildtype, C3$^{-/-}$ and C3aR$^{-/-}$ animals. There were no effects on *zif268* expression but

5-minute exposure to the EPM was found to significantly increase *c-Fos* expression

30 minutes later, compared to animals who remained in their home-cages (Figure

2B). In wildtype animals there was a 4.95-fold increase in *c-Fos* expression over

baseline and increases in EPM-evoked *c-Fos* expression in *C3*$^{-/-}$ and *C3aR*$^{-/-}$ models

of 7.77-fold and 6.54-fold, respectively (Figure 2B). Thus, whilst being placed on the

EPM led to significant general increases in *c-Fos* in vHPC, the enhanced activity

seen in the knockout models did not differentiate between *C3*[-/-] and *C3aR*[-/-] and did

not predict the observed dissociations in anxiety-related behaviours.

Altered sensitivity of *C3aR*[-/-] and *C3*[-/-] mice to diazepam in the elevated plus

maze

In a further independent cohort of animals, we tested the sensitivity of EPM induced

anxiety-like behaviour to the benzodiazepine diazepam, an established clinically

effective anxiolytic drug[37,44]. Our initial behavioural findings were replicated in

vehicle-treated animals across all behavioural indices of anxiety, again showing an

anxiogenic phenotype in *C3aR*[-/-] but not *C3*[-/-] mice (Figure 3A). As anticipated, in

wildtype mice 2mg/kg diazepam led to a trend for increased time on the open arms

(Figure 3B) and a significant reduction in SAPs which are considered to reflect risk

assessment behaviour[45-47](Figure 3C). These effects were therefore consistent with

reduced anxiety[40,47]. In contrast, the same dose of drug that was effective in eliciting

anxiolysis in wildtypes was without effects in *C3aR*[-/-] mice (Figure 3A,B,C) and

produced a seemingly anxiogenic (increased anxiety) pattern of effects in *C3*[-/-] mice

(Figure 3A,B and Figure S4B,C). Locomotor activity monitored across all the maze

(Figure S4D) indicated that wildtype and *C3aR*[-/-] mice were unlikely to have been

influenced by diazepam-induced sedation. In *C3*[-/-] mice however, activity was

significantly suppressed under drug conditions indicating a possible sedative effect.

Together, these data indicated a fundamentally altered reactivity to diazepam in both

C3[-/-] and *C3aR*[-/-] models.

Enhanced fear learning in *C3$^{-/-}$* but not *C3aR$^{-/-}$* mice

Psychiatric disorders are associated with maladaptive responses to both innate, unconditioned and learned aversive stimuli[48,49]. We therefore extended our analysis to investigate whether the behavioural dissociations in innate anxiety observed between *C3$^{-/-}$* and *C3aR$^{-/-}$* mice would also apply to learned or conditioned fear, where a previously neutral cue generates a fear response as a result of predicting an aversive outcome. In a further group of animals, we used the fear-potentiated startle (FPS) paradigm[30,50] a well-established method of generating learned fear responses to an acute and imminent danger signal that is characteristic of fear. In this paradigm (see Figure 4A and Methods) fear learning is indexed by an enhanced response to a startling noise in the presence of a cue (the conditioned stimulus or CS) previously paired with mild foot shock (the unconditioned stimulus). In the pre-conditioning session, pulse-alone trials revealed increased basal startle reactivity in both *C3aR$^{-/-}$* and *C3$^{-/-}$* animals relative to wildtype (Figure 4B). Increased reactivity to the unconditioned foot shock stimulus (in the absence of any startle stimulus) during the conditioning session was also observed in both knockouts (Figure 4C). However, these common effects of genotype were not seen in the fear-potentiated startle measures which index fear learning. Whilst all groups showed the expected enhancement of the startle response in the presence of the CS, the effect of the CS was significantly greater in *C3$^{-/-}$* animals relative to the *C3aR$^{-/-}$* and wildtype animals (Figure 4D), indicating enhanced learning of the fear related-cue by the *C3$^{-/-}$* mice. This was the opposite pattern of effects to those observed in the tests of innate anxiety and showed a double dissociation in the impact of manipulating C3 and C3aR function that depended fundamentally on the nature of the aversion.

Complement signalling pathways underlying abnormal learned fear

phenotypes in $C3^{-/-}$ mice

Given the central role of C3 within the complement system, its deletion affects a

number of distal pathways (Figure 5A), with the activity of the C3a/C3aR, C3b/CR3,

C5a/C5aR and terminal pathways affected. Therefore, the loss of function of any of

these pathways may have contributed to the observed fear learning phenotype in $C3^{-/-}$

mice. However, it is possible to exclude effects due to loss of the C3a/C3aR

pathway since the fear learning phenotype was specific to $C3^{-/-}$ and not $C3aR^{-/-}$ mice

(Figure 4D). This left iC3b/CR3 signalling and/or pathways downstream of C5 (i.e.

C5a/C5aR, terminal pathway) as the remaining possibilities. In order to distinguish

between these pathways, we repeated the FPS experiment with the addition of $C5^{-/-}$

mice. This model has intact C3a/C3aR and iC3b/CR3 signalling, but lacks C5a/C5b

and terminal pathway activity, as do $C3^{-/-}$ mice. We hypothesised that if $C5^{-/-}$ mice

also displayed enhanced fear-potentiated startle, then the phenotype in $C3^{-/-}$ mice

would likely be due to a loss of C5a/C5aR signalling or the terminal pathway. On the

other hand, if $C5^{-/-}$ mice demonstrated normal fear-potentiated startle, this would

confine the likely mediating pathway in $C3^{-/-}$ mice to iC3b/CR3 signalling.

Results from the pre-conditioning session demonstrated increases in the startle

response of $C5^{-/-}$ mice, (Figure 5B) although in this instance the previously observed

enhanced startle reactivity in $C3^{-/-}$ mice (Figure 4B) was not replicated. In the

conditioning session, there was again evidence of increased startle responses to

shock (Figure 5C) in $C3^{-/-}$ mice and responses were of a similar magnitude in $C5^{-/-}$

mice, although these were not significantly different to wildtype. We replicated the

previous finding of enhanced fear-potentiated startle in $C3^{-/-}$ mice (Figure 5D), but

critically $C5^{-/-}$ mice showed no evidence of enhanced fear learning and were

comparable to wildtypes, indicating that loss of iC3b/CR3 signalling, but not loss of

C5a/C5aR and the terminal pathway, was involved in the $C3^{-/-}$ fear learning

phenotype. Additionally, we did not observe innate anxiety-like phenotypes in $C5^{-/-}$

mice (Figure S5).

This pattern of effects allowed us to distinguish between the likely mechanisms

underlying the enhanced learned fear in $C3^{-/-}$ mice, as we could exclude concomitant

loss of C5a/C5aR signalling or molecules downstream of C5, and hence also

exclude an explanation based on effects of C5a/C5aR signalling on developmental

neurogenesis[51,52]. Instead, these data raised the possibility of an explanation based

on the established effects of the iC3b/CR3 pathway on synaptic pruning, a

mechanism involving microglia mediated elimination of synapses impacting on

neurodevelopment and learning-related synaptic plasticity[13,14,53].

Differential expression of stress and anxiety related genes in $C3aR^{-/-}$ and $C3^{-/-}$
mice

We next sought to assess whether the dissociations in innate anxiety and learned

fear between $C3aR^{-/-}$ and $C3^{-/-}$ models were associated with differential gene

expression in brain regions associated with emotional behaviours. We assayed gene

expression in three regions with known roles in stress and anxiety; the medial

prefrontal cortex (mPFC), ventral hippocampus (vHPC) and cerebellum[28,54](Figure

6A). Given our previous data showing differential corticosterone responses and

altered sensitivity to diazepam in both knockouts, we first measured expression of

the glucocorticoid receptor *Nr3c1* and the corticotrophin-releasing hormone receptor

1 *Crhr1*, together with *Gabra2* which encodes the GABA$_A$ receptor α2 subunit

responsible for mediating benzodiazepine anxiolysis[55]. There were no effects of

genotype on *Crhr1* and *Gabra2* mRNA expression in any of the brain regions

assayed (Figure 6B,C). *Nr3c1* expression did however show effects of genotype with

strong trends indicating specific increases in *C3aR*[-/-] mice in mPFC and vHPC and

significantly increased expression in cerebellum that was common to both knockouts

(Figure 6D). As activation of C3aR has been shown to stimulate calcium influx from

the extracellular space[2-4,56,57] and calcium channel subunit variants have strong links

to risk for psychiatric and neurological disorders, as well as anxiety phenotypes[58,59],

we also investigated a panel of voltage-gated calcium channels. Expression of

Cacna1d, which encodes the Cav1.3 channel of L-type calcium gated voltage

channels, was increased in the mPFC of *C3aR*[-/-] mice (Figure 6E) and in both *C3*[-/-]

and *C3aR*[-/-] mice there was upregulation of cerebellar *Cacna1e,* which encodes the

Cav2.3 channel of R-type voltage gated calcium channels (Figure 6F). We also

investigated the expression of *Cacna1c* which encodes the Ca$_v$1.2 subunit of L-type

voltage gated calcium channels that forms the channel pore allowing calcium entry[60].

We found genotype and brain region specific changes in *Cacna1c* expression, with

selective increases in expression in *C3aR*[-/-] mice in the vHPC and cerebellum, but

not the mPFC (Figure 6G).

Discussion

We have used knockout models manipulating specific complement proteins to reveal

dissociable effects of complement pathways on distinct elements of aversive

behaviours. A main finding was a profound innate anxiety phenotype in the *C3aR*[-/-]

mice that was absent in *C3*[-/-] mice. The specificity of the anxiety phenotype exhibited

by $C3aR^{-/-}$ mice at the behavioural level was paralleled by EPM-evoked

corticosterone levels, confirming the validity of the EPM as an index of anxiety-like

behaviour. A second main finding was a double dissociation in terms of fear learning,

whereby $C3^{-/-}$ and not $C3aR^{-/-}$ mice displayed enhanced conditioned fear reactivity.

Using $C5^{-/-}$ mice, we concluded that the mechanisms accounting for the fear learning

phenotype in $C3^{-/-}$ mice are likely to arise from lack of iC3b/CR3 pathway activity.

These data indicate distinct effects mediated by closely related elements of the

complement system that differentially impact upon the neural mechanisms

underlying innate anxiety and learned fear.

Our use of specific complement knockout models allowed us to pinpoint the likely

complement pathways and potential mechanisms underlying the observed

phenotypes. Since C3a is solely produced via C3 cleavage, and C3aR is the

canonical receptor for C3a, we expected that any phenotypes dependent on the

binding of C3a to the C3aR would be apparent in both $C3^{-/-}$ and $C3aR^{-/-}$ models.

However, this was not borne out in our data. Given the divergence in innate anxiety

phenotypes seen, one explanation is that the $C3aR^{-/-}$ phenotypes are independent of

C3a and instead mediated by an alternative ligand. It has long been speculated that

there may be promiscuity of the C3aR due to its unusually large second extracellular

loop[61]. Indeed, a cleavage fragment of the neuropeptide precursor protein VGF (non-

acronymic), TLQP-21, was recently reported to bind the C3aR[62,63]. This peptide has

pleiotropic roles including in the stress response[64] and its precursor VGF is widely

expressed throughout the CNS[65] and in regions associated with stress reactivity and

C3aR expression such as the hypothalamus[66,67]. Determining whether the

mechanisms underlying innate anxiety phenotypes in *C3aR*[/-] mice are dependent on

TLQP-21/C3aR interactions will be a priority for future work.

Whether the *C3aR*[/-] phenotypes described here are the result of ongoing effects of

C3aR deletion in the adult brain or instead the enduring consequence of

neurodevelopmental impacts of *C3aR* deficiency also remains to be determined. On

the basis of previous findings implicating C3aR in both developmental

neurogenesis[68] and in acute brain changes following behavioural manipulations[69],

both are possibilities. One strategy would be to test whether acute administration of

the C3aR antagonist SB290157[70] phenocopies the constitutive knockout of *C3aR*,

though at present no data is available on the CNS penetration of SB290157 and this

molecule has received criticism due to evidence of agonist activity[71].

We also probed mechanisms underpinning the *C3aR*[/-] innate anxiety phenotype by

assessing the effects of the anxiolytic drug diazepam. We found that a dose of

diazepam that was anxiolytic in wildtype mice had no effect in *C3aR*[/-] mice. Stretch

attend postures, postulated to reflect risk assessment behaviour[46], are highly

sensitive to pharmacological manipulation[40,72] and in agreement with our own

findings, diazepam has been shown to specifically decrease SAPs in the absence of

effects on open arm exploration[73]. Importantly, *C3aR*[/-] mice consistently performed

more SAPs than other genotypes, and therefore floor effects cannot account for the

pattern of results observed. Benzodiazepines act on $GABA_A$ receptors[55] however we

found no significant changes in expression of *Gabra2*, a $GABA_A$ receptor subunit

responsible for anxiolytic actions of benzodiazepines in tests of innate anxiety[55], in

the brain regions sampled. This raises the possibility of alternative molecular

mechanisms mediating the anxiety phenotypes seen in the $C3aR^{-/-}$ model. Whatever

the molecular underpinnings of the dissociable anxiety phenotypes, our data show a

profoundly altered effect of diazepam in both knockouts; a lack of response in $C3aR^{-/-}$

and an apparent paradoxical anxiogenic effect of the drug in $C3^{-/-}$ mice, though this

interpretation needs to take into account an apparent selective sedative effect in $C3^{-/-}$

mice.

In contrast to innate anxiety, we observed a specific effect of $C3$ knockout on

conditioned fear. The absence of a comparable phenotype in $C3aR^{-/-}$ and $C5^{-/-}$ mice

suggested that these effects were unlikely to be due to loss of either C3a/C3aR,

C5a/C5aR, or the terminal pathway, and instead that enhanced fear learning

phenotypes in $C3^{-/-}$ mice were likely dependent on loss of the iC3b/CR3 pathway.

This pathway has been strongly implicated in activity dependent synaptic elimination

during neurodevelopment[13,25] and in age-dependent synapse loss[74]. While

demonstrations of complement mediated synaptic pruning during development have

centered on the visual system, complement-mediated microglial phagocytosis of

dopamine D1 receptors has been demonstrated in the nucleus accumbens with

functional impacts on social behaviour[75]. It remains to be seen whether complement

mediated processes of this nature, within brain regions linked to fear processing

such as the ventral hippocampus, amygdala and prefrontal cortex are responsible for

enhanced fear learning in $C3^{-/-}$ mice, or whether this phenotype is a general

consequence of altered synaptic elimination throughout the $C3^{-/-}$ brain. In addition to

developmental processes, the iC3b/CR3 pathway could also be involved acutely in

fear learning. $C3$ mRNA is upregulated during discrete stages of fear learning[69] and

microglial CR3 is implicated in long term depression[76]. Furthermore, complement-

mediated engulfment of synapses by microglia may be important in the forgetting of

fear memories[14].

At the gene expression level, we found some changes which were common to both

knockouts, and one result that was specific to *C3aR*[-/-] mice. Regarding the latter,

there was a highly specific increase in expression of *Cacna1c* in the ventral

hippocampus and cerebellum of *C3aR*[-/-] mice. This finding is of potential interest

given the strong evidence implicating *CACNA1C* variants in genetic risk for a broad

spectrum of psychiatric disorders including schizophrenia and bipolar disorder[58], with

anxiety phenotypes reported in both human risk variant carriers[77] and animal

models[78-81]. Whether alterations in *Cacna1c* are of functional relevance to the *C3aR*

[-/-] anxiety phenotypes observed here remains to be determined. We also observed

increased cerebellar expression of the glucocorticoid receptor in both *C3*[-/-] and

C3aR[-/-] mice, suggesting that these alterations may result from the absence of

C3a/C3aR signalling. Expression of these genes did not differentiate between

models and therefore were unlikely to contribute to the dissociable effects of the

knockouts on behaviour and stress hormone physiology. Neuronal activity in the

ventral hippocampus as indexed by *c-Fos* gene expression after exposure to the

EPM also failed to differentiate between genotypes, and hence activity in other brain

regions in particular the amygdala and hypothalamus may be more informative in

terms of functional neuroanatomy underlying the anxiety-related behavioural and

physiological differences seen in the knockout models.

Anxiety and fear are commonly comorbid with depression and recent preclinical work

has suggested a protective role for C3a/C3aR in chronic-stress induced depressive-

behaviour[82]. Given the common co-occurrence of anxiety and depression, our

findings of enhanced anxiety in *C3aR*[-/-] mice might seem at odds with the reported

resilience of this strain[-] to depression-related phenotypes[82]. However, the chronic

unpredictable-stress paradigm used in these studies is likely to evoke significant

inflammation, and therefore the extent to which our data in acutely stressed animals

can be compared is questionable. In summary, our findings add significantly to the

evidence that perturbations of the complement system, whether reduced

complement activation as in the present work or excessive activation as is predicted

by *C4* genetic variants[24,83], have major and dissociable effects on brain and

behavioural phenotypes of relevance to core clinical symptoms of psychiatric

disease.

Materials and Methods

Mouse models and husbandry. Wildtype and *C3*[-/-] strains were sourced in-house

from Professor B. Paul Morgan and Dr Timothy Hughes (strains originally from The

Jackson Laboratory; B6.PL-Thy1[a]/CyJ stock#000406 and B6;129S4-C3tm1Crr/J

stock#003641 respectively); *C3aR*[-/-] mice were provided by Professor Craig Gerard

of Boston Children's Hospital, USA (strain originally from The Jackson Laboratory,

strain C.129S4-C3ar1[tm1Cge]/J, stock#005712, subsequently backcrossed to

C57Bl/6J. *C5*[-/-] mice (as described in [84]) were kindly donated by Professor Marina

Botto, Imperial College London. This strain originated from naturally C5-deficient

DBA/2J mice, that had been backcrossed to C57Bl/6J. *C3*[-/-], *C3aR*[-/-] and *C5*[-/-] strains

were maintained via homozygous breeding on a C57Bl/6J background and in all

experiments were compared to wildtype mice also on a C57Bl/6J background. Male

mice (3-8 months old during experimental testing) were kept in a temperature and

humidity-controlled vivarium (21±2°C and 50±10%, respectively) with a 12-hour light-

dark cycle (lights on at 07:00hrs/lights off at 19:00hrs). Home cages were

environmentally enriched with cardboard tubes, soft wood blocks and nesting

materials and animals were housed in single sex littermate groups (2-5 mice/cage).

Standard laboratory chow and water were available *ad libitum*. All procedures were

performed in accordance with the requirements of the UK Animals (Scientific

Procedures) Act (1986).

General behavioural methods. Testing took place between the hours of 09:00 and

17:00, with random distribution of testing for subjects of different genotypes

throughout the day. Mice were habituated to the test rooms for 30 min prior to

testing. All assays involved individual testing of mice and apparatus was cleaned

thoroughly with a 70% ethanol solution between subjects.

Data collection. Data for the EPM, EZM and Open Field were collected using

EthoVision XT software (Noldus Information Technology, Netherlands) via a video

camera mounted above the centre of each piece of apparatus. Tracking of each

subject was determined as the location of the greater body-proportion (12 frames/s)

in the specific virtual zones of each piece of apparatus.

The elevated plus maze (EPM). The maze, positioned 300 mm above the floor and

illuminated evenly at 15 lux, was constructed of opaque white Perspex and consisted

of two exposed open arms (175 x 78 mm^2, length x width, no ledges) and two

equally sized enclosed arms, which had 150 mm high walls [85]. Equivalent arms were

arranged opposite one another. Subjects were placed at the enclosed end of a

closed arm and allowed to freely explore for 5 minutes. Data from each pair of arms

were combined to generate single open and closed arm values (number and

duration of arm entries and latency of first entry to each arm).   In addition, the

following parameters were manually scored (by an experimenter positioned at a

computer in the same room as the maze, watching the live-video stream of the test);

number of stretch-attend postures (SAPs; defined as the animal slowly and carefully

reaching towards the open arms in a low, elongated body posture[86,87]) and number

of head dips from the open arms (looking down over the edge of an open arm).

The elevated zero maze (EZM). The maze, positioned 520 mm above the floor and

illuminated evenly at 15 lux, was constructed of wood and consisted of two exposed

open regions (without ledges; 52 mm wide) and two equally sized enclosed regions

(also 52 mm wide), which had 200 mm high grey opaque walls. The diameter of the

maze was 600mm. Equivalent regions were arranged opposite one another.

Subjects were placed at the border of one of the open and closed regions and

allowed to freely explore for 5 min. Data from each pair of regions were combined to

generate single open and closed region values (number and duration of region

entries and latency of first entry to each region).   In addition, the number of head

dips (as above) were measured. Due to the high walls of the enclosed sections of

the maze, subjects were not visible to the experimenter when in the closed regions

and therefore these parameters were scored only when a subject was on the open

regions.

Locomotor activity (LMA).   LMA was measured in an apparatus consisting of

twelve transparent Perspex chambers (each 210 x 360 x 200 mm, width x length x

height). Two infrared beams were embedded within the walls of each chamber,

which crossed the chamber 30 mm from each end and 10 mm from the chamber

floor. Individual subjects were placed in a designated chamber for a 120 min duration

on three consecutive days. Beam breaks were recorded as an index of activity,

using a computer running custom written BBC Basic V6 programme with additional

interfacing by ARACHNID (Cambridge Cognition Ltd, Cambridge, UK). Data were

analysed as the total number of beam breaks per session per day.

Fear-potentiated startle (FPS). FPS was assessed using startle chamber

apparatus which consisted of a pair of ventilated and soundproofed SR-LAB startle

chambers (San Diego Instruments, CA, USA) each containing a non-restrictive

Plexiglas cylinder (35 mm in diameter), mounted on a Perspex plinth, into which a

subject was placed. The motor responses of subjects to white noise stimuli

(generated from a speaker 120 mm above the cylinder) were recorded via a

piezoelectric accelerometer, attached centrally below the Plexiglas cylinder, which

converted flexion plinth vibration into electrical signals. The peak startle response,

within 200ms from the onset of each startle presentation, in each trial, was

normalized for body weight differences using Kleiber's 0.75 mass exponent[88] as

per[85]. A computer running SR-Lab software (Version 94.1.7.48) was used to

programme trials and record data. A foot shock grid connected to a shock generator

(San Diego Instruments, CA, USA) was inserted into the Plexiglas cylinder before

conditioning sessions.

FPS consisted of three separate sessions presented over a two-day period (see

Figure 4A). On the first day, mice were given a pre-conditioning session immediately

followed by the conditioning session. The pre-conditioning session started with a 5

min acclimatisation phase followed by presentation of 3 no-stimulus trials, and then a

block of pulse-alone trials presented at 90, 100 and 110dB (5 of each at 40 ms

duration). Trials were randomly distributed throughout the session and presented

with a 60 s random interval (range 36 s to 88 s). After the pre-conditioning session

was complete, mice were removed from the startle chambers, restraint tubes

cleaned, and shock grids were placed into the Plexiglas cylinders prior to

commencing the conditioning session. The mice were then returned to the startle

chambers and subjected to a session consisting of a 5 min acclimatisation phase

followed by 3 CS+shock trials, with 3 no stimulus trials before and after, presented

with a 2min random interval (range 1.5 to 3min). The scrambled 0.14 mA, 0.5 s foot

shock was delivered in the final 0.5 s of the 30 s visual CS. Following a 24hr delay,

subjects were assessed for FPS in the post-conditioning session. This session

followed the same format as the pre-conditioning session (5 min acclimatisation

phase followed by presentation of 3 no-stimulus trials, and then a block of pulse-

alone trials presented at 90, 100 and 110dB, with 5 of each at 40 ms duration)

however the final block of trials also included pulse+CS trials at 90, 100 and 110 dB

(5 of each), with the startle pulse presented in the final 40 ms of the CS. FPS was

determined as the fold change between pulse-alone trials and pulse+CS trials within

the post-conditioning session.

Corticosterone measurements. Testing took place between the hours of 10:00 and

14:00 to account for the diurnal pattern of corticosterone release[89]. Mice were

allowed to freely explore the EPM for 5 min, after which they were placed in a

holding cage for a further 25 min before being culled by cervical dislocation. Control

mice were removed from their home cage and immediately culled. There was an

equal distribution of subjects of different genotypes, counterbalanced between the

two test conditions and throughout the testing period. Trunk blood was collected into

heparin tubes (Becton Dickinson, USA) and immediately centrifuged at 4000 rpm for

10 min, and the supernatant removed and frozen at -80°c until further use. A

corticosterone ELISA was performed according to manufacturer's instructions (ADI-

900-097, Enzo Life Sciences, UK) and analysed using a four-parameter logistic

curve plug in (https://www.myassays.com/four-parameter-logistic-curve.assay).

Immediate early gene (IEG) experiment. Mice were allowed to freely explore the

EPM for 5 min, after which they were placed in a holding cage for a further 25 min

before being culled by cervical dislocation. Control mice were removed from their

home cage and immediately culled. Brains were removed and the ventral

hippocampus (vHPC; the whole hippocampus was dissected and the posterior third

was collected) was dissected and frozen at -80° until further use. Tissue was

processed for qPCR analyses as detailed below.

Diazepam study. Wildtype, $C3^{-/-}$ and $C3aR^{-/-}$ were used and were randomly

assigned to either vehicle or drug conditions within each genotype. A three-day

dosing regimen of diazepam (2 mg/kg, i.p., Hameln Pharmaceuticals, UK) or an

equivalent volume of vehicle (0.1 M phosphate buffered saline, pH 7.4) was used.

Following 2 days of pre-treatment, diazepam or vehicle was administered 30 min

prior to testing on the EPM on the 3[rd] day.

Tissue for gene expression analysis. Mice were removed from their home cage and immediately culled via cervical dislocation. Brains were removed from mice in both conditions, and the following regions dissected: medial prefrontal cortex (mPFC), ventral hippocampus (vHPC) and cerebellum (see Figure 6A) and frozen at -80° until further use.

Quantitative Polymerase Chain Reaction (qPCR). Gene expression was analysed using standardised qPCR methods with quantification using the $2^{-\Delta\Delta Ct}$ method[90]. Mouse brain tissue from the mPFC, vHPC and the cerebellum was analysed. RNA was extracted using the RNeasy kit (QIAGEN) and was subsequently treated with DNAse to remove genomic DNA (TURBO DNA-free kit, Thermo Fisher Scientific). RNA was then converted to cDNA (RNA to cDNA EcoDry Premix, Random Hexamers, Clontech, Takara). cDNA samples were run in triplicate in 96 well reaction plates using SYBR-Green-based qPCR (SensiFast, HI-ROX, Bioline) according to manufacturer's instructions using a StepOnePlus System (Applied Biosystems, Thermo Fisher Scientific). Genotypes were counterbalanced across plates and genes of interest were run alongside housekeeping genes *Gapdh* and *Hrpt1* for each sample, within the same reaction plate. All samples were run in triplicate and samples differing by >0.3 Cts were excluded. The change in expression of genes of interest, after normalisation to the two house-keeping genes (ΔCt) was transformed to yield $2^{-\Delta\Delta Ct}$ values. Relative changes from wildtype animals were calculated for each gene of interest.

Primers

Primers were designed to span at least one exon-exon junction and to match the

target sequence only in mouse (Primer-Blast, NCBI) and were synthesised

commercially (Sigma Aldrich). Primer efficiency was determined separately through

a dilution series of cDNA samples from wildtype hippocampus, cerebellum and

cortex. Primers with an efficiency between 90-110% were selected.

Table 1. List of primer sequences used.

Gene	Species	Forward	Reverse
Gapdh	Mouse	GAACATCATCCCTGCATCCA	CCAGTGAGCTTCCCGTTCA
Hprt1	Mouse	TTGCTCGAGATGTCATGAAGGA	AATGTAATCCAGCAGGTCAGCAA
Gabra2	Mouse	AAGCCACTGGAGGAAAACATCT	TTAGCCAGCACCAACCTGAC
Crhr1	Mouse	CTTCAACTCTTTCCTGGAGTCCT	TGGCAGAGCGGACCTCA
Nr3c1	Mouse	AAACTCTGCCTGGTGTGCTC	GGTAATTGTGCTGTCCTTCCAC
Cacna1c	Mouse	ATGGTTCTTGTCAGCATGTTGCGG	TGCAAATGTGGAACCGGTAAGTG
Cacna1d	Mouse	AGAGGACCATGCGAACGAG	CCTTCACCAGAAATAGGGAGTCT
Cacna1e	Mouse	CTCATGTCACCACCGCTAGG	TCTGTCTGCACCACCTTTGG
cFos	Mouse	CTCCAAGCGGAGACAGATCAA	TCGGTGGGCTGCCAAAATAA
Zif268	Mouse	GACCACAGAGTCCTTTTCTGACA	CTGAAAAGGGGTTCAGGCCA

Genotyping Genotyping was performed on post-mortem tail tip samples. Qiagen

DNeasy Blood and Tissue Kits (Qiagen, Manchester, UK) were used to extract

genomic DNA (gDNA) as per the manufacturers standard protocol. For $C3^{-/-}$ mice,

JAX protocol 27746 was used (common; ATCTTGAGTGCACCAAGCC, wildtype;

GGTTGCAGCAGTCTATGAAGG, mutant; GCCAGAGGCCACTTGTATAG) and for

C3aR[−/−] JAX protocol 27638 was used (common; AGCCATTCTAGGGGCGTATT,

wild   type   reverse;   CATGGTTTGGGGTTATTTCG,   mutant   reverse;

TTGATGTGGAATGTGTGCGAG). For both genotypes, a touchdown cycling protocol

was used (see JAX protocols for details). Genotyping for *C5*[−/−] mice was performed

as described in [84].

Statistical analysis. All statistical analyses were carried out using GraphPad Prism

8.4.1 (GraphPad Software, CA, USA). Data was assessed for equality of variances

using the Brown-Forsythe test and then appropriate parametric (*t* test, one-way or

two-way ANOVA) or non-parametric (Kruskal-Wallis) tests used. *Post hoc* pairwise

comparisons were performed using the Tukey or Dunn's tests for parametric or non-

parametric analyses, respectively. For all analyses, alpha was set to 0.05 and exact

p values were reported unless $p < 0.0001$. All p values were multiplicity adjusted[91].

Data are expressed as mean ± standard error of the mean.

The main between-subjects' factor for all ANOVA analyses was GENOTYPE (WT,

C3[−/−], *C3aR*[−/−], or *C5*[−/−]). For the EPM, LMA and FPS experiments, there were within-

subject factors of ZONE (open, closed, middle), DAY (1,2,3) and STIMULUS

INTENSITY (90, 100, 110 dB) respectively. Analysis of plasma corticosterone by

two-way ANOVA included an additional between subject factor of CONDITION

(baseline, EPM), and for the diazepam experiment, there was an additional between

subject factor of DRUG (diazepam, vehicle). For qPCR analyses, ΔCt values were

analysed by one-way ANOVA.

Author's contributions

The study was designed by LJW, TH, BPM, WPG and LSW. LJW and TH performed behavioural experiments with assistance from NH. Molecular analyses were performed by SAB, EB, MT, ALM, SAB, EB, MT, NL, AIB and LJW. Data interpretation were carried out by JH, MJO, JR, WPG, NH, TRH, BPM, LSW and TH. The manuscript was drafted by LJW, TH, WPG and LSW. All authors approved the final manuscript.

Acknowledgements

The authors thank Professor Craig Gerard and Professor Marina Botto for provision of the $C3aR^{-/-}$ and $C5^{-/-}$ strains respectively, and to Rhys Perry, Pat Mason, Helen Read and other staff at JBIOS for their invaluable animal care and husbandry. Thanks also to Dr Emma Watts for her feedback on the manuscript. This work was supported by a Wellcome Trust Integrative Neuroscience PhD Studentship awarded to LJW (099816/Z/12/Z), a Waterloo Foundation Early Career Fellowship awarded to LJW, a Hodge Centre for Neuropsychiatric Immunology Seed Corn and Project grant

awarded to LJW and a Wellcome Trust Strategic Award 100202/Z/12/Z (DEFINE)

held by the Neuroscience and Mental Health Research Institute at Cardiff University.

Competing financial interests

The authors declare no competing financial interests.

Materials and correspondence

All data from this study are available from the corresponding authors upon

reasonable request.

References

1.   Coulthard, L. G. & Woodruff, T. M. Is the Complement Activation Product
C3a a Proinflammatory Molecule? Re-evaluating the Evidence and the
Myth. *J. Immunol.* **194,** 3542–3548 (2015).
2.   Norgauer, J. *et al.* Complement fragment C3a stimulates Ca2+ influx in
neutrophils via a pertussis-toxin-sensitive G protein. *Eur. J. Biochem.*
217, 289–294 (1993).
3.   Möller, T., Nolte, C., Burger, R., Verkhratsky, A. & Kettenmann, H.
Mechanisms of C5a and C3a Complement Fragment-Induced [Ca2+]i
Signalling in Mouse Microglia. *Journal of Neuroscience* **17,** 615–624
(1997).
4.   Lian, H., Li, Y., Lu, H.-C. & Zheng, H. NFκB-Activated Astroglial Release
of Complement C3 Compromises Neuronal Morphology and Function
Associated with Alzheimer's Disease. *Neuron,* **85,** 101–115 (2015).
5.   Sayah, S. *et al.* Two different transduction pathways are activated by C3a
and C5a anaphylatoxins on astrocytes. *Molecular Brain Research* **112,**
53–60 (2003).
6.   Morley, B. J. & Walport, M. J. *The Complement FactsBook.* (Academic
Press, 1999).
7.   Kemper, C. & Köhl, J. Back to the future – non-canonical functions of
complement. *Seminars in Immunology* **37,** 1–3 (2018).
8.   Nataf, S., Stahel, P. F., Davoust, N. & Barnum, S. R. Complement
anaphylatoxin receptors on neurons: new tricks for old receptors? *Trends*
Neurosci. **22,** 397–402 (1999).
9.   Veerhuis, R., Nielsen, H. M. & Tenner, A. J. Complement in the brain.
Molecular Immunology **48,** 1592–1603 (2011).
10.   Bénard, M., Gonzalez, B. J., Biological, M. S. J. O.2004. Characterization
of C3a and C5a receptors in rat cerebellar granule neurons during

maturation neuroprotective effect of C5a against apoptotic cell death. *J*
Biol. Chem, **42***,* 43487-43496 (2004).

11. Rahpeymai, Y. *et al.* Complement: a novel factor in basal and ischemia-
induced neurogenesis. *EMBO J* **25,** 1364–1374 (2006).

12. Gorelik, A. *et al.* Developmental activities of the complement pathway in
migrating neurons. *Nature Communications* **8,** 15096 (2017).

13. Stevens, B. *et al.* The classical complement cascade mediates CNS
synapse elimination. *Cell* **131,** 1164–1178 (2007).

14. Wang, C. *et al.* Microglia mediate forgetting via complement-dependent
synaptic elimination. *Science,* **367,** 688-694 (2020).

15. Harold, D. *et al.* Genome-wide association study identifies variants at
CLU and PICALM associated with Alzheimer's disease. *Nature Genetics*
41, 1088–1093 (2009).

16. Jansen, I. E. *et al.* Genome-wide meta-analysis identifies new loci and
functional pathways influencing Alzheimer's disease risk. *Nature Genetics*
51, 404–413 (2019).

17. Shi, Q. *et al.* Complement C3-Deficient Mice Fail to Display Age-Related
Hippocampal Decline. *Journal of Neuroscience* **35,** 13029–13042 (2015).

18. Zhou, J., Fonseca, M. I., Pisalyaput, K. & Tenner, A. J. Complement C3
and C4 expression in C1q sufficient and deficient mouse models of
Alzheimer's disease. *Journal of Neurochemistry* **106,** 2080–2092 (2008).

19. Corbett, B. A. *et al.* A proteomic study of serum from children with autism
showing differential expression of apolipoproteins and complement
proteins. *Mol. Psychiatry* **12,** 292 (2007).

20. Maes, M., Delange, J., Ranjan, R., Meltzer, H. Y. & Desnyer, R. Acute
phase proteins in schizophrenia, mania and major depression:
modulation by psychotropic drugs. *Psychiatry Research,* **66,** 1–11 (1997).

21. Ruland, T. *et al.* Molecular serum signature of treatment resistant
depression. *Psychopharmacology* 1–9 (2016).

22. Song, Y. R. *et al.* Specific alterations in plasma proteins during
depressed, manic, and euthymic states of bipolar disorder. *Brazilian*
Journal of Medical and Biological Research **48,** 973–982, (2015).

23. Oganesyan, L. P., Mkrtchyan, G. M., Sukiasyan, S. H. & Boyajyan, A. S.
Classic and alternative complement cascades in post-traumatic stress
disorder. *Bull. Exp. Biol. Med.* **148,** 859–861 (2009).

24. Sekar, A. *et al.* Schizophrenia risk from complex variation of complement
component 4. *Nature,* **530,** 117–183 (2016).

25. Schafer, D. P. *et al.* Microglia Sculpt Postnatal Neural Circuits in an
Activity and Complement-Dependent Manner. *Neuron* **74,** 691–705
(2012).

26. Håvik, B. *et al.* The Complement Control-Related Genes CSMD1 and
CSMD2 Associate to Schizophrenia. *BPS* **70,** 35–42 (2011).

27. Zhang, S. *et al.* Association Between Polymorphisms of the Complement
3 Gene and Schizophrenia in a Han Chinese Population. *Cell. Physiol.*
Biochem. **46,** 2480–2486 (2018).

28. Tovote, P., Fadok, J. P. & Lüthi, A. Neuronal circuits for fear and anxiety.
Nat. Rev. **16,** 317–331 (2015).

29. Perusini, J. N. & Fanselow, M. S. Neurobehavioral perspectives on the
distinction between fear and anxiety. *Learn. Mem.* **22,** 417–425 (2015).

30. Davis, M. Neural systems involved in fear and anxiety measured with
fear-potentiated startle. *Am Psychol* **61,** 741–756 (2006).
31. Engin, E. *et al.* Modulation of anxiety and fear via distinct
intrahippocampal circuits. *eLife,* **5:e14120**, (2016).
32. Vogelzangs, N., Beekman, A. T. F., de Jonge, P. & Penninx, B. W. J. H.
Anxiety disorders and inflammation in a large adult cohort. *Translational*
Psychiatry **3,** e249–e249 (2013).
33. Girardi, G. *et al.* Imaging of activated complement using ultrasmall
superparamagnetic iron oxide particles (USPIO)-conjugated vectors: An
in vivo in utero non-invasive method to predict placental insufficiency and
abnormal fetal brain development. *Mol. Psychiatry* **20,** 1017–1026 (2015).
34. Kulkarni, A. P., Govender, D. A., Kotwal, G. J. & Kellaway, L. A.
Modulation of anxiety behavior by intranasally administered Vaccinia
virus complement control protein and curcumin in a mouse model of
Alzheimer's disease. *Current Alzheimer Research* **8,** 95–113 (2011).
35. Fischer, M. B., Ma, M., Goerg, S., Zhou, X. & Xia, J. Regulation of the B
cell response to T-dependent antigens by classical pathway complement.
J.Immunol, **157,** 549-556, (1996).
36. Humbles, A. A. *et al.* A role for the C3a anaphylatoxin receptor in the
effector phase of asthma. *Nature,* **406,** 998–1001 (2000).
37. Pellow, S., Chopin, P., File, S. E. & Briley, M. Validation of open : closed
arm entries in an elevated plus-maze as a measure of anxiety in the rat.
J. Neurosci. Methods **14,** 149–167 (1985).
38. Walf, A. A. & Frye, C. A. The use of the elevated plus maze as an assay
of anxiety-related behavior in rodents. *Nat Protoc* **2,** 322–328 (2007).
39. Ennaceur, A. Tests of unconditioned anxiety - pitfalls and
disappointments. *Physiology & Behavior* **135,** 55–71 (2014).
40. RODGERS, R. J. Anxiety, defence and the elevated plus-maze.
Neuroscience and Biobehavioural Reviews, **21,** 801-810, (1997).
41. Braun, A. A., Skelton, M. R., Vorhees, C. V. & Williams, M. T.
Comparison of the elevated plus and elevated zero mazes in treated and
untreated male Sprague-Dawley rats: Effects of anxiolytic and anxiogenic
agents. *Pharmacology Biochemistry and Behavior* **97,** 406–415 (2011).
42. Kheirbek, M. A. *et al.* Differential Control of Learning and Anxiety along
the Dorsoventral Axis of the Dentate Gyrus. *Neuron* **77,** 955–968 (2013).
43. Kubik, S., Miyashita, T. & Guzowski, J. F. Using immediate-early genes
to map hippocampal subregional functions. *Learning & Memory* **14,** 758–
770 (2007).
44. Lepicard, E. M., Joubert, C., Hagneau, I., Perez-Diaz, F. & Chapouthier,
G. Differences in anxiety-related behavior and response to diazepam in
BALB/cByJ and C57BL/6J strains of mice. *Pharmacology Biochemistry*
and Behavior **67,** 739–748 (2000).
45. Grewal, S. S., SHEPHERD, J. K., Bill, D. J., Fletcher, A. & Dourish, C. T.
Behavioural and pharmacological characterisation of the canopy
stretched attend posture test as a model of anxiety in mice and rats.
Psychopharmacology **133,** 29–38, (1997).
46. Kaesermann, H. P. Stretched attend posture, a non-social form of
ambivalence, is sensitive to a conflict-reducing drug action.
Psychopharmacology **89,** 31–37, (1986).

47. Albrechet-Souza, L., Cristina de Carvalho, M., Rodrigues Franci, C. &
Brandão, M. L. Increases in plasma corticosterone and stretched-attend
postures in rats naive and previously exposed to the elevated plus-maze
are sensitive to the anxiolytic-like effects of midazolam. *Hormones and*
Behavior **52,** 267–273 (2007).
48. Holt, D. J., Coombs, G., Zeidan, M. A., Goff, D. C. & Milad, M. R. Failure
of neural responses to safety cues in schizophrenia. *Arch Gen Psychiatry*
69, 893–903 (2012).
49. Grillon, C., Morgan, C. A., Southwick, S. M., Davis, M. & Charney, D. S.
Baseline startle amplitude and prepulse inhibition in Vietnam veterans
with posttraumatic stress disorder. *Psychiatry Research* **64,** 169–178
(1996).
50. Falls, W. A., Carlson, S., Turner, J. G. & Willott, J. F. Fear-potentiated
startle in two strains of inbred mice. *Behavioral Neuroscience* **111,** 855–
861 (1997).
51. Coulthard, L. G. *et al.* Complement C5aR1 Signalling Promotes
Polarization and Proliferation of Embryonic Neural Progenitor Cells
through PKCζ. *J. Neurosci.* **37,** 5395–5407 (2017).
52. Hawksworth, O. A., Coulthard, L. G., Taylor, S. M., Wolvetang, E. J. &
Woodruff, T. M. Brief Report: Complement C5a Promotes Human
Embryonic Stem Cell Pluripotency in the Absence of FGF2. *Stem Cells,*
32, 3278-3284, (2014).
53. Schafer, D. P. *et al.* Microglia sculpt postnatal neural circuits in an activity
and complement-dependent manner. *Neuron* **74,** 691–705 (2012).
54. Apps, R. & Strata, P. Neuronal circuits for fear and anxiety - the missing
link. *Nat. Rev.* **16,** 642 (2015).
55. Smith, K. S., Engin, E., Meloni, E. G. & Rudolph, U. Benzodiazepine-
induced anxiolysis and reduction of conditioned fear are mediated by
distinct GABAA receptor subtypes in mice. *Neuropharmacology* **63,** 250–
258 (2012).
56. Ahamed, J., Venkatesha, R. T., Thangam, E. B. & Ali, H. C3a Enhances
Nerve Growth Factor-Induced NFAT Activation and Chemokine
Production in a Human Mast Cell Line, HMC-1. *J. Immunol.* **172,** 6961–
6968 (2004).
57. Wu, F. *et al.* Complement component C3a plays a critical role in
endothelial activation and leukocyte recruitment into the brain. *J*
Neuroinflammation **13,** 23 (2016).
58. Hamshere, M. L. *et al.* Genome-wide significant associations in
schizophrenia to ITIH3/4, CACNA1C and SDCCAG8, and extensive
replication of associations reported by the Schizophrenia PGC. *Mol.*
Psychiatry **18,** 708-712, (2019).
59. Kabir, Z. D. *et al.* Rescue of impaired sociability and anxiety-like behavior
in adult *cacna1c*-deficient mice by pharmacologically targeting eIF2α.
Mol. Psychiatry **22,** 1096–1109 (2017).
60. Simms, B. A. & Zamponi, G. W. Neuronal Voltage-Gated Calcium
Channels: Structure, Function, and Dysfunction. *Neuron* **82,** 24–45
(2014).
61. Chao, T. H. *et al.* Role of the second extracellular loop of human C3a
receptor in agonist binding and receptor function. *Journal of Biological*
Chemistry **274,** 9721–9728 (1999).

62.   Hannedouche, S. *et al.* Identification of the C3a Receptor (C3AR1) as the
868   Target of the VGF-derived Peptide TLQP-21 in Rodent Cells. *Journal of*
869   *Biological Chemistry* **288,** 27434–27443 (2013).
870   63.   Cero, C. *et al.* The TLQP-21 peptide activates the G-protein-coupled
871   receptor C3aR1 via a folding-upon-binding mechanism. *Structure* **22,**
872   1744–1753 (2014).
873   64.   Razzoli, M. *et al.* Implication of the VGF-derived peptide TLQP-21 in
874   mouse acute and chronic stress responses. *Behav. Brain Res.* **229,** 333–
875   339 (2012).
876   65.   van den Pol, A. N., Bina, K., Decavel, C. & Ghosh, P. VGF expression in
877   the brain. *J. Comp. Neurol.* **347,** 455–469 (1994).
878   66.   Lewis, B., Francis, K., Gasque, P., Scanlon, M. & Ham, J. Functional
879   complement C3a receptors in the rat pituitary gland. *Endocrine Abstracts*
880   **3,** 198 (2002).
881   67.   Francis, K. Complement C3a receptors in the pituitary gland: a novel
882   pathway by which an innate immune molecule releases hormones
883   involved in the control of inflammation. *The FASEB Journal,* **17,** 2266-
884   2268 (2003).
885   68.   Coulthard, L. G., Hawksworth, O. A. & Woodruff, T. M. Complement: The
886   Emerging Architect of the Developing Brain. *Trends Neurosci.* **41,** 373–
887   384 (2018).
888   69.   Scholz, B. *et al.* The Regulation of Cytokine Networks in Hippocampal
889   CA1 Differentiates Extinction from Those Required for the Maintenance
890   of Contextual Fear Memory after Recall. *PLoS ONE* **11,** e0153102–29
891   (2016).
892   70.   Ames, R. S. *et al.* Identification of a selective nonpeptide antagonist of
893   the anaphylatoxin C3a receptor that demonstrates antiinflammatory
894   activity in animal models. *J. Immunol.* **166,** 6341–6348 (2001).
895   71.   Therien, A. G., Baelder, R. & Köhl, J. Agonist Activity of the Small
896   Molecule C3aR Ligand SB 290157. *J. Immunol.* **174,** 7479–7480 (2005).
897   72.   Holly, K. S., Orndorff, C. O. & Murray, T. A. MATSAP: An automated
898   analysis of stretch-attend posture in rodent behavioral experiments.
899   *Scientific Reports* **6,** (2016).
900   73.   Genewsky, A. *et al.* How much fear is in anxiety? Preprint at
901   https://doi:10.1101/385823 (2018).
902   74.   Hong, S. *et al.* Complement and microglia mediate early synapse loss in
903   Alzheimer mouse models. *Science* **352,** 712–716 (2016).
904   75.   Kopec, A. M., Smith, C. J., Ayre, N. R., Sweat, S. C. & Bilbo, S. D.
905   Microglial dopamine receptor elimination defines sex-specific nucleus
906   accumbens development and social behavior in adolescent rats. *Nature*
907   *Communications* **9,** 3769 (2018).
908   76.   Zhang, J. *et al.* Microglial CR3 Activation Triggers Long-Term Synaptic
909   Depression in the Hippocampus via NADPH Oxidase. *Neuron* **82,** 195–
910   207 (2014).
911   77.   Roussos, P., Giakoumaki, S. G., Georgakopoulos, A., Robakis, N. K. &
912   Bitsios, P. The CACNA1C and ANK3 risk alleles impact on affective
913   personality traits and startle reactivity but not on cognition or gating in
914   healthy males. *Bipolar Disorders* **13,** 250–259 (2011).
915   78.   Bader, P. L. *et al.* Mouse model of Timothy syndrome recapitulates triad
916   of autistic traits. *Proc. Natl. Acad. Sci. U.S.A.* **108,** 15432–15437 (2011).

79.   Lee, A. S. *et al.* Forebrain elimination of cacna1c mediates anxiety-like
behavior in mice. *Nat.Neurosci* **17,** 1054–1055 (2012).
80.   Dao, D. T. *et al.* Mood Disorder Susceptibility Gene CACNA1C Modifies
Mood-Related Behaviors in Mice and Interacts with Sex to Influence
Behavior in Mice and Diagnosis in Humans. *Biological Psychiatry* **68,**
801–810 (2010).
81.   Dedic, N. *et al.* Cross-disorder risk gene CACNA1C differentially
modulates susceptibility to psychiatric disorders during development and
adulthood. *Nat.Neurosci* **23,** 533–543 (2018).
82.   Crider, A. *et al.* Complement component 3a receptor deficiency
attenuates chronic stress-induced monocyte infiltration and depressive-
like behavior. *Brain, Behavior, and Immunity* **70,** 246–256 (2018).
83.   Sellgren, C. M. *et al.* Increased synapse elimination by microglia in
schizophrenia patient-derived models of synaptic pruning. *Nat.Neurosci*,
22, 374-385, (2019).
84.   Wang, Y. *et al.* A role for complement in antibody-mediated inflammation:
C5-deficient DBA/1 mice are resistant to collagen-induced arthritis. *J.*
Immunol. **164,** 4340–4347 (2000).
85.   Mikaelsson, M. A., Constancia, M. C. A., Dent, C. L., Wilkinson, L. S. &
Humby, T. Placental programming of anxiety in adulthood revealed by
Igf2-null models. *Nature Communications* **4,** 1–9 (2013).
86.   Molewijk, H. E., van der Poel, A. M. & Olivier, B. The ambivalent
behaviour 'stretched approach posture' in the rat as a paradigm to
characterize anxiolytic drugs. *Psychopharmacology* **121,** 81–90, (1995).
87.   Mackintosh, J. H. & Grant, E. C. A Comparison of the Social Postures of
Some Common Laboratory Rodents. *Behaviour* **21,** 246–259 (1963).
88.   Kleiber, M. Body size and metabolism. *Hilgardia Journal of Agricultural*
Science, **6,** 315-353 (1932).
89.   Dickmeis, T. Glucocorticoids and the circadian clock. *Journal of*
Endocrinology **200,** 3–22 (2008).
90.   Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression
Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method.
Methods **25,** 402–408 (2001).
91.   Wright, S. P. Adjusted p-values for simultaneous inference. *Biometrics*
48, 1005–1013 (1992).

Figure 1. *C3aR^{-/-}*, but not *C3^{-/-}* mice show increased anxiety-like behaviour in the elevated plus maze (EPM;A-E) and elevated zero maze (EZM;F-J). (A) Heatmaps displaying relative time per zone of the EPM across genotypes **(B)** Latency to first open arm visit; wildtype 8.21±1.53s, *C3^{-/-}* 14.1±2.52s, *C3aR^{-/-}* 27.6±8.31s, (H$_2$=10.5, p=0.005). *Post hoc* tests demonstrated that *C3aR^{-/-}* mice took

significantly longer to first enter the open arms than wildtype mice (p=0.0045). **(C)**

C3aR$^{-/-}$ mice distributed their time across the EPM differently to wildtype and *C3*$^{-/-}$

mice (GENOTYPE×ZONE, $F_{4,62}$=17.7, p=0.0001) spending less time in the open

arms (*C3aR*$^{-/-}$ 16.70±3.73s vs. wildtype 75.78±8.86s, p<0.0001, *C3aR*$^{-/-}$ vs. *C3*$^{-/-}$

93.86±9.59s p<0.0001) and significantly more time in the closed arms (*C3aR*$^{-/-}$

221.88±12.06s vs. wildtype 146.01±7.01s, p<0.0001, and *C3*$^{-/-}$ 140.04±8.61s

p<0.0001). **(D)** *C3aR*$^{-/-}$ (14.90±2.22) mice performed significantly fewer head dips

than wildtype (37.92±2.53, p<0.0001) and *C3*$^{-/-}$ mice (49.17±2.37, p<0.0001),

whereas *C3*$^{-/-}$ mice performed significantly more head dips than wildtype mice

(p=0.0061; overall ANOVA $F_{2,31}$=48.0, p<0.0001). **(E)** *C3aR*$^{-/-}$ mice performed

significantly more stretch attend postures (SAPs; 15.30±1.80) than wildtype

(7.58±1.66, p=0.0042) and *C3*$^{-/-}$ mice (5.67±0.94, p=0.0004; overall ANOVA

$F_{2,31}$=10.3, p=0.0004). **(F)** Heatmaps displaying relative exploration of the open

segments of the elevated zero maze, across genotypes. Note that due to the height

of the walls in the closed regions it was not possible to track mice or observe

ethological behaviours such as grooming or SAPs. **(G)** There was a significant

difference in the latency to first enter the open arms (wildtype 101.00±31.00s, *C3*$^{-/-}$

42.00±2.52s, *C3aR*$^{-/-}$ 204.00±40.40s, H_2=8.13, p=0.0171). *Post hoc* tests revealed

that *C3aR*$^{-/-}$ mice took significantly longer than *C3*$^{-/-}$ mice to initially enter the open

region (p=0.0140). **(H)** The number of entries made to open regions differed between

genotypes (wildtype 7.69±1.69, *C3*$^{-/-}$ 11.5±1.43s, *C3aR*$^{-/-}$ 2.10±1.15, $F_{2,30}$=8.96,

p=0.0009). *C3aR*$^{-/-}$ mice made significantly fewer entries to the open areas than

wildtype (p=0.0324) and *C3*$^{-/-}$ mice (p=0.0006) and **(I)** spent significantly less time on

the open arms (1.77±1.29) compared to wildtype (35.7±8.43s, p=0.0132) and *C3*$^{-/-}$

(57.7±9.32s, p<0.0001; overall Kruskal-Wallis test H_2=19.2, p<0.0001). **(J)** *C3*[^-/-] mice

performed significantly more head dips (20.7±1.62) than wildtype (13.9±1.89,

p=0.048) and *C3aR*[^-/-] mice (10.3±2.42, p=0.0034; overall ANOVA

$F_{2,27}$=6.86,p=0.0039). Data are mean ± S.E.M. *, **, *** and **** represent p≤0.05,

p≤0.01, p≤0.001 and p≤0.0001 for *post-hoc* genotype comparisons, respectively.

1036
1037
1038

Figure 2. Neuroendocrine response and neuronal activation following exposure to the elevated plus maze (A) 5-minute exposure to the EPM significantly elevated corticosterone in all genotypes (main effect of CONDITION, $F_{1,54}$=143, p<0.0001; baseline 344.66±63.70 vs. EPM 3553.84±274.13). There was a significant GENOTYPE × CONDITION interaction ($F_{2,54}$=4.64, p=0.0138). *Post hoc* analysis showed that after the EPM, *C3aR*[/-] mice demonstrated significantly higher corticosterone levels (4640.27±376.13) than wildtype (3033.78±535.06, p=0.0127) and *C3*[/-] mice (2948.00±374.87, p=0.0032). *Post hoc* tests also indicated that there were no baseline differences between genotypes (wildtype 216.54±63.2 vs. *C3aR*[/-] 316.17±111.60 p>0.9999, wildtype vs. *C3*[/-] p=0.9927, and *C3*[/-] 485.60±130.35 vs.*C3aR*[/-] mice p=0.9992). **(B)** A separate group of animals were taken from their home-cages and culled immediately or placed on the open arm of the EPM for 5 minutes and culled 25 minutes later. *c-Fos* mRNA was significantly upregulated after exposure to the EPM across genotypes (main effect of CONDITION, $F_{1,46}$=45.4, p<0.0001). There was no significant main effect of GENOTYPE ($F_{2,46}$=1.27, p=0.2894) and no CONDITION × GENOTYPE interaction ($F_{2,46}$=0.0004, p=0.9996). *Zif268* mRNA was not altered in EPM exposed animals, nor between genotypes. Data represent mean + S.E.M or fold change from wildtype home-cage + SEM. *, **,

1057   and **** represent p≤0.05, p≤0.01 and p≤0.0001 for *post-hoc* genotype comparisons,

1058   respectively.

Figure 3. Altered sensitivity to diazepam in *C3aR*^-/- and *C3*^-/- mice. Behaviourally

naïve mice were treated with either diazepam (2mg/kg, i.p) or vehicle injections once

daily for 2 days and then 30 minutes prior to testing. (**A**) Heatmaps demonstrating

duration spent in zones of the maze by vehicle treated and diazepam treated

animals (**B**) Plots showing duration spent on open arms in 1-minute time bins (start-

01:00 was excluded due to effect of diazepam in delaying initial entry to open arms

across genotypes, see Supplementary Figure 4a). There was a trend for wildtype

diazepam treated animals to spend more time on the open arms throughout the task

although this did not reach significance (main effect of DRUG, $F_{1,38}$=1.41, p=0.2462).

In $C3^{/-}$ mice there was a strong tendency for drug treated animals to explore the

open arms less than vehicle treated $C3^{/-}$ mice (main effect of DRUG, $F_{1,22}$=1.25,

p=0.2764). A similar, though less pronounced pattern was seen in $C3aR^{/-}$ mice

(main effect of DRUG, $F_{1,19}$=1.55, p=0.2284) **(C)** There were genotype differences in

SAPs (main effect of GENOTYPE, $F_{2,79}$=10.7, p<0.0001), a main effect of DRUG

($F_{1,79}$=4.13, p=0.0454) and a significant GENOTYPE $\times$ DRUG interaction ($F_{2,79}$=4.64,

p=0.0138). *Post hoc* tests showed that diazepam significantly reduced the number of

SAPs in wildtype mice only (wildtype vehicle 9.00±1.44 vs. wildtype diazepam

3.09±0.71, p=0.0006, $C3^{/-}$ vehicle 4.67±1.24 vs. $C3^{/-}$ diazepam 5.00±1.51,

p=0.9975, $C3aR^{/-}$ vehicle 11.45±1.49 vs. $C3aR^{/-}$ diazepam 10.50±1.34, p=0.9558).

Data are mean + S.E.M. *, **, and *** represent p≤0.05, p≤0.01 and p≤0.001 for

post-hoc genotype comparisons, respectively.

Figure 4. Enhanced fear-potentiated startle in *C3⁻/⁻* but not *C3aR⁻/⁻* mice. (A) Flow

chart depicting the FPS protocol used, which took place in three separate sessions

over two consecutive days. Baseline startle reactivity to a range of pulse intensities

was assessed in the pre-conditioning session, immediately preceding the

conditioning session in which a visual stimulus (light) was paired with 3 weak foot

shocks. 24 hours later, subjects were re-introduced to the same chamber and startle

reactivity was compared between pulse-alone trials and pulse+CS trials to determine

the degree of FPS. On all trials, the peak startle response was recorded and

normalised for body weight differences using Kleiber's 0.75 mass exponent, and

fold-changes calculated. **(B)** There was a significant main effect of GENOTYPE

($F_{2,66}$=9.04, p=0.0003) and a significant GENOTYPE $\times$ STIMULUS INTENSITY

interaction ($F_{4,132}$=7.55, p<0.0001). *C3⁻/⁻* and *C3aR⁻/⁻* mice demonstrated increased

levels of startle responding relative to wildtype mice at 100dB (*C3⁻/⁻* 11.34±1.51 vs.

wildtype 2.63±0.26, p<0.0001, *C3aR*$^{-/-}$ 9.12±2.63 vs. wildtype p=0.0174) and 110dB

(*C3*$^{-/-}$ 12.69±1.55 vs. wildtype 3.74±0.50, p<0.0001, *C3aR*$^{-/-}$ 10.58±3.58 vs. wildtype

p=0.0111) **(C)** *C3*$^{-/-}$ and *C3aR*$^{-/-}$ mice also showed increased startle responding to

the footshock+CS (*C3*$^{-/-}$ 75.18±5.73, *C3aR*$^{-/-}$ 73.14±7.78) pairings relative to wildtype

mice (44.34±5.29, *C3*$^{-/-}$ vs. wildtype p=0.0006, *C3aR*$^{-/-}$ vs. wildtype p=0.0137, overall

ANOVA $F_{2,66}$=8.7, p=0.0004), although it should be noted that responses were much

greater to these stimuli in all mice than to the startle stimuli in the pre-conditioning

session. **(D)** In the post-conditioning session, all mice demonstrated increases to the

pulse+CS stimuli in comparison to pulse-alone stimuli, as demonstrated by the fold-

change increase in startle responding, however, this effect was significantly

increased in *C3*$^{-/-}$ mice (3.72±0.27) relative to wildtype (1.99±0.11, p<0.0001) and

C3aR$^{-/-}$ mice (2.27±0.24, p=0.0056, overall Kruskal-Wallis test H_2=27.7, p<0.0001).

Data are mean + S.E.M. *, **, *** and **** represent p≤0.05, p≤0.01, p≤0.001 and

p≤0.0001 for *post-hoc* genotype comparisons, respectively.

1149

1150

1151

A

1152

Figure 5. Pathways underlying fear learning phenotypes in *C3^{-/-}* mice (A) C3

activation leads to generation of cleavage fragments C3a and C3b. The former

signals via C3aR whereas the latter signals via complement receptor 3 (CR3). C3b is

also necessary for forming the convertase enzyme that cleaves C5. Upon cleavage,

C5 generates the fragments C5a and C5b (not shown). C5a signals via the C5aR,

whereas C5b propagates activity of the terminal complement pathway via C6. Since

C3 cannot be activated in *C3^{-/-}* mice, the action of all these pathways (C3a/C3aR,

C3b/CR3, C5a/C5aR, terminal pathway) is absent. By using *C5^{-/-}* mice, which lack

C5a/C5aR and terminal pathway activity, we examined whether lack of C3b/CR3,

C5a/C5aR or the terminal pathway was responsible for fear learning phenotypes in

$C3^{-/-}$ mice. **(B)** In the pre-conditioning session there were significant main effects of

GENOTYPE ($F_{2,27}$=18.4, p<0.0001) and STIMULUS INTENSITY ($F_{2,54}$=19.0,

p<0.0001) and a significant GENOTYPE × STIMULUS INTENSITY interaction

($F_{4,54}$=7.00, p<0.0001). $C5^{-/-}$ mice demonstrated increased levels of startle

responding relative to wildtype and $C3^{-/-}$ mice at all stimulus intensities (90dB; WT

2.55±0.26 vs. $C5^{-/-}$ 13.92±3.14, p=0.0069, $C3^{-/-}$ 4.92±1.40 vs. $C5^{-/-}$ p=0.0405, WT vs.

$C3^{-/-}$ p=0.7919; 100dB; WT 3.23±0.45 vs. $C5^{-/-}$ 20.83±4.07, p<0.0001, $C3^{-/-}$

4.924.17±0.88 vs. $C5^{-/-}$ p<0.0001, WT vs. $C3^{-/-}$ p=0.9639; 110dB; WT 4.92±2.03 vs.

$C5^{-/-}$ 29.78±4.29, p<0.0001, $C3^{-/-}$ 8.07±2.76 vs. $C5^{-/-}$ p<0.0001, WT vs. $C3^{-/-}$

p=0.6643). **(C)** There were no significant differences in startle responses to the

footshock+CS pairings during the conditioning session (WT 20.23±4.76, $C3^{-/-}$

33.10±5.74, $C5^{-/-}$ 31.08±3.59, $F_{2,27}$=2.10, p=0.1421). **(D)** In the post-conditioning

session, all mice demonstrated increases to the pulse+CS stimuli in comparison to

pulse-alone stimuli, as demonstrated by the fold-change increase in startle

responding, however, this effect was again significantly increased only in $C3^{-/-}$ mice

(2.92±0.43) relative to wildtype (1.47±0.17, p=0.0020) and $C5^{-/-}$ mice (1.23±0.08,

p=0.0004, overall ANOVA $F_{2,27}$=11.5, p=0.0002). Data represent mean + S.E.M. *,

, * and **** represent p≤0.05, p≤0.01, p≤0.001 and p≤0.0001 for *post-hoc*

genotype comparisons, respectively.

1182

1183

1184

Figure 6. mRNA expression of stress and anxiety related genes. (A) Animals

were culled and the medial prefrontal cortex (mPFC), ventral hippocampus (vHPC)

and cerebellum were dissected. **(B)** There were no significant differences in the

expression of Corticotrophin releasing hormone receptor 1 (*Crhr1*) in any region,

across genotypes (mPFC $F_{2,53}=0.587$, p=0.5597, N wildtype=20, *C3$^{-/-}$*=17, *C3aR$^{-/-}$*=

19; vHPC $F_{2,49}=0.169$, p=0.8450, N WT=20, *C3$^{-/-}$*=15, *C3aR$^{-/-}$*= 17; cerebellum

$F_{2,47}=0.0482$, p=0.8346, N WT=19, *C3$^{-/-}$*=17, *C3aR$^{-/-}$*= 14). **(C)** There were also no

significant changes in the expression of the GABA$_A$ receptor α2 subunit (*Gabra2*) in

any region, across genotypes (mPFC $H_2=1.04$, p=0.5939, N wildtype=20, *C3$^{-/-}$*=19,

$C3aR^{/-}$= 16; vHPC $F_{2,49}$=0.721, p=0.4914, N WT=20, $C3^{/-}$=13, $C3aR^{/-}$= 19;

cerebellum $F_{2,47}$=0.221, p=0.8026, N WT=18, $C3^{/-}$=17, $C3aR^{/-}$= 15). **(D)** Expression

of the glucocorticoid receptor (*Nr3c1*) was significantly increased in the cerebellum of

both $C3^{/-}$ and $C3aR^{/-}$ groups ($F_{2,61}$=10.3, p=0.0002, $C3^{/-}$ vs. wildtype p=0.0023,

$C3aR^{/-}$ vs. wildtype p=0.0002, N wildtype=19, $C3^{/-}$=20, $C3aR^{/-}$= 15). There were

trends towards increased expression of the glucocorticoid receptor gene *Nr3c1* in the

mPFC and vHPC of $C3aR^{/-}$ mice but these were not significant (mPFC; $F_{2,56}$=1.33,

p=0.2723, N wildtype=20, $C3^{/-}$=20, $C3aR^{/-}$= 19, vHPC; $F_{2,62}$=1.11, p=0.3345, N

wildtype=23, $C3^{/-}$=20, $C3aR^{/-}$= 22). **(E)** Calcium voltage-gated channel subunit

Alpha 1d (*Cacna1d*) expression was changed in the mPFC ($F_{2,36}$=7.52, p=0.0407)

owing to altered expression between $C3^{/-}$ and $C3aR^{/-}$ mice (p=0.0314; N

wildtype=11, $C3^{/-}$=13, $C3aR^{/-}$= 15). There were no differences in the vHPC

($F_{2,31}$=2.27, p=0.1199, N wildtype=14, $C3^{/-}$=10, $C3aR^{/-}$= 10) or cerebellum

($F_{1,39}$=0.648, p=0.5286, N wildtype=14, $C3^{/-}$=16, $C3aR^{/-}$= 12). **(F)** Expression of the

Calcium voltage-gated channel subunit Alpha 1e (*Cacna1e*) was significantly

upregulated in the cerebellum of both knockouts ($F_{2,39}$=7.52, p=0.0017, wildtype vs.

$C3^{/-}$ p=0.0082, wildtype vs. $C3aR^{/-}$ p=0.0032; N wildtype=14, $C3^{/-}$=16, $C3aR^{/-}$= 12).

There were borderline significant changes in expression in the vHPC ($F_{2,32}$=3.15,

p=0.0565, N wildtype=14, $C3^{/-}$=11, $C3aR^{/-}$= 10) and no significant changes in the

mPFC (H_2=3.43, p=0.1802, N wildtype=11, $C3^{/-}$=12, $C3aR^{/-}$= 15).**(G)** Expression

levels of the Calcium voltage-gated channel subunit Alpha 1c (*Cacna1c*) were

significantly increased in $C3aR^{/-}$ mice in a regionally specific manner in the vHPC

($F_{2,47}$=3.20, p=0.0496, $C3^{/-}$ vs. wildtype p=0.6895, $C3aR^{/-}$ vs. wildtype p=0.0295, N

wildtype=21, $C3^{/-}$=13, $C3aR^{/-}$= 16) and the cerebellum ($F_{2,54}$=5.84, p=0.0051, $C3^{/-}$

vs. wildtype p=0.1613, $C3aR^{/-}$ vs. wildtype p=0.0024, N wildtype=20, $C3^{/-}$=20,

C3aR$^{/-}$= 17). There were no significant changes in the mPFC ($F_{2,52}$=1.04, p=0.5939,

N wildtype=20, *C3*$^{/-}$=19, *C3aR*$^{/-}$= 16). Data represent fold change from wildtype +

SEM. *, **, *** represent p≤0.05, p≤0.01 and p≤0.001 for *post-hoc* genotype

comparisons, respectively.

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234