Dynamics and variability in regenerative potential of neuronal subtypes in the

***Nematostella* nerve net**

Jamie A. Havrilak[1], Layla Al-Shaer[1], Nesli Akinci[1], Aldine Amiel[2], Eric Röttinger[2],

Michael J. Layden[1*]

[1] Lehigh University, Department of Biological Sciences, Bethlehem, PA

[2] Université Nice Sophia Antipolis UMR 7284, CNRS UMR 7284, INSERM U1081,
Institute for Research on Cancer and Aging, Nice, France

[*] Corresponding Author: layden@lehigh.edu

Keywords: *Nematostella*, Regeneration, Neurogenesis, Nerve net, LWamide,

Regenerative neurogenesis, Nervous system scalability

Abstract:

The cell dynamics and responses of individual neuronal cell types during whole-

body axis regeneration are not well understood in any system. We exploited transgenic

technology to track individual neuronal subtypes within regenerating *Nematostella*

vectensis nerve nets *in vivo*. Individual neuronal subtypes had specific responses that

included always, never, and conditional regeneration. Regenerates were smaller than the

pre-amputated animal, and the nerve net was reduced in neuronal number. The reduced

nerve net in regenerates led us to investigate whether the nerve net scaled with changes in

body size. Neuronal number decreased as animals shrunk in response to starvation.

Conversely, neurons increased in fed animals as they increased in length. The increase

and decrease of neurons was reversible by switching fed animals to starvation and vice

versa. Regenerates and starved/fed animals with scaled-down or scaled-up nervous

systems responded to external mechanical cues. These data suggest that the *Nematostella*

nerve net is dynamic, capable of scaling with changes to overall body size, and that

individual neuronal subtypes display differential regenerative potential which is likely

linked to the scale-state of regenerating nervous system.

Introduction:

How entire nervous systems reform in the context of whole-body axis

regeneration is not well understood. Work in hydrozoan cnidarians and planarians have

provided insights on some of the general molecular mechanisms that promote

regenerative neurogenesis (Cebrià et al. 2002; Currie & Pearson 2013; Chen et al. 2013;

Deochand et al. 2016; Gahan et al. 2017; Galliot et al. 2007; Miljkovic-Licina et al. 2007;

Siebert et al. 2018; Roberts-Galbraith et al. 2016), but details about when and how

individual neuronal subtypes respond are currently lacking. For example, it is currently

unknown if all neuronal subtypes regenerate new neurons lost by amputation. In

planarians, the total number of neurons for some neuronal fates positively correlates with

the length of the animal (Takeda et al. 2009), but tracking individual neuronal fates *in*

vivo in planarians is currently not feasible. As such it is not clear if neuronal number is

dynamic within any one individual. The lack of understanding about the cellular

dynamics during whole-body axis regeneration limits our ability to interpret functional

studies aimed at understanding the molecular mechanisms that re-pattern whole nervous
systems.

The cnidarian sea anemone *Nematostella vectensis* offers the potential to track
individual neuronal fates throughout the process of regeneration. *Nematostella* is an
optically clear diploblastic animal. The primary axis is the oral-aboral axis. The oral pole
possesses a single mouth surrounded by up to 16 tentacles (Fritz et al. 2013). Eight
segments radially situated around the mouth run the length of the oral-aboral axis. Each
radial segment contains a mesentery, which contains gonads, a number of differentiated
cells, longitudinal tracts of neurites, and longitudinal myoepithelial cells (Tucker et al.
2011; Renfer et al. 2010; Jahnel et al. 2014; Steinmetz et al. 2017; Williams 1975). The
Nematostella nervous system is a typical cnidarian nerve net, which initially forms during
embryonic and larval stages in both the ectoderm and endoderm (Rentzsch et al. 2016;
Nakanishi et al. 2012). A number of transgenic lines exist, which allows for *in vivo*
tracking of individual cell types (Layden et al. 2016; Nakanishi et al. 2012; Richards &
Rentzsch 2014; Renfer et al. 2010; Havrilak et al. 2017; Busengdal & Rentzsch 2017).
Using *in vivo* characterization of the *NvLWamide-like::mCherry* transgene, we identified
at least five neuronal subtypes, determined that some of those neuronal subtypes form in
a highly stereotyped manner, and that the number of some neuronal subtypes correlates
with length (Layden et al. 2016; Havrilak et al. 2017).

Here, we exploited the ability to track individual neuronal subtypes using
NvLWamide-like::mCherry transgenic animals during whole-body axis regeneration and
during feeding and starvation treatments. *Nematostella* regenerates were smaller than
their parent uncut animal, and their nervous systems had reduced numbers of at least
some neuronal subtypes. Neurons were classified into at least three classes based on their
specific regenerative responses: Class I neurons, which regenerated under all
circumstances; Class II neurons, which did not appear to regenerate; and Class III
neurons, which displayed differential regenerative potential that appeared to depend on
the size of the remnant fragment. The reduced size of the regenerated nervous system led
us to investigate scalability of the nerve net. Fed or starved *Nematostella* grew or shrunk
respectively, and the nervous system responded in part by modifying the number of some
neuronal subtypes to scale with changing polyp length. Despite their reduced nervous

systems, regenerates and animals that had been starved still responded to being touched

and were capable of feeding, suggesting that the reduction does not inhibit functionality.

These data suggest that the *Nematostella* nerve net is highly dynamic and scales with

body size. To the best of our knowledge, these findings are the first evidence that

regenerative responses of individual neurons are variable during whole-body axis

regeneration. These data indicate that the molecular mechanisms enacted to promote

regenerative neurogenesis likely vary based on the size of the remnant fragment and

which neuronal subtypes must regenerate.

Results:

Neuronal subtypes display differential regenerative potentials.

To determine how neuronal subtypes regenerate, our goal was to bisect animals

into two equal halves (*Oral* and *Aboral* remnants) and measure the neuronal response in

Aboral remnants (Figure 1A). We focused on *Aboral* remnants because there are

morphological indicators for when regeneration has completed and a well-established

regenerative time course (Passamaneck & Martindale 2012; Amiel et al. 2015; Bossert et

al. 2013). One challenge was that animals do not always relax completely or uniformly

along the oral-aboral axis when treated with $MgCl_2$, and the wound site on both remnants

curls in, which makes using measurements to ensure the cut site is always in the same

position unreliable. However, the number of longitudinal neurons in the *NvLWamide-*

like::mCherry transgenic animals scale with animal length and appear to be evenly

distributed along the oral-aboral axis (Havrilak et al. 2017). We reasoned that if evenly

distributed, quantifying longitudinal neurons pre- and post-amputation could be used as a

proxy for length, and allow us to confirm amputation occurred in the intended location.

The oral-aboral axis was divided into 4 equal quarters, and the longitudinal neurons were

quantified in each quarter in animals of various lengths. Each quarter contained similar

numbers of longitudinal neurons with no statistical difference between the numbers of

neurons in any of the quarters (Figure 1B). Thus, we were justified in using longitudinal

number to track amputation sites and as a proxy for length.

To quantify regeneration of individual neuronal subtypes we used animals that

were 2.5 to 8.5 mm long (Supplemental Figure 1A), because this range represents the

sizes of naturally occurring wild-caught animals found along the East Coast of the United

States (Adam Reitzel, personal communication, 11/2018). Animals in this size range also

have a manageable number of longitudinal and tripolar neurons for quantification. We

focused on longitudinal, tripolar, pharyngeal, and tentacular neuron subtypes described

by the *NvLWamide-like::mCherry* transgene (Havrilak et al. 2017). Tentacular and

pharyngeal neurons are specifically located at the oral end of the animal, whereas

longitudinal and tripolar are evenly distributed along the oral-aboral axis in the endoderm

and ectoderm, respectively (Figure 1C) (Havrilak et al. 2017; Layden et al. 2016). We

excluded the *NvLWamide-like::mCherry+* mesentery neurons from this analysis. This is

because the ruffled mesentery structure and large number of mesentery neurons made

quantification unreliable. With the *NvLWamide-like::mCherry* line it is possible to easily

score for the regeneration of oral specific neurons (tentacular and pharyngeal), as well as

track regeneration of an individual neuronal subtype in both the endoderm and ectoderm.

To characterize neuronal subtype regeneration animals were allowed to relax,

$MgCl_2$ was added to relaxed animals, and the number of longitudinal and tripolar neurons

were quantified (*uncut*) (Figure 1C, F, and H). Animals were then bisected in half and

longitudinal and tripolar neurons were quantified in the *Aboral* remnant (*time 0 cut*)

(Figure 1D, F-I; Supplemental Figure 2). This process takes less than 20 minutes/animal.

$MgCl_2$ was then washed out and animals were allowed to regenerate in 1/3x artificial sea

water, (hereafter called *Nematostella* medium). Because extensive cell death occurs over

the first 24 hours of regeneration (Warner et al. 2019), regenerating *Aboral* remnants

were paralyzed using $MgCl_2$ and neuronal number was reassessed again at 24 hours post

amputation (*24hpa*) (Figure 1F-I; Supplemental Figure 2). $MgCl_2$ was again washed out

and animals were allowed to regenerate until 7 days post amputation (dpa) when neuronal

numbers were quantified a final time (*7dpa*) (Figure 1E, F-I; Supplemental Figure 2). All

oral structures including mouth, pharynx and tentacles regenerated in the aboral remnant

by *7dpa* (Figure 1E; Supplemental Figure 3), which is consistent with previous findings

and suggests that $MgCl_2$ treatments did not impact regenerative timing (Amiel et al. 2015;

Passamaneck & Martindale 2012). Regenerates were reduced in size compared to pre-

amputated animals, as indicated by a reduction in overall length (Figure 1 C and E;

Supplemental Figure 1A-C) and decreased longitudinal and tripolar neuronal numbers

(Figure 1F-I; Supplemental Figure 2). On average regenerates from animals bisected at

the midpoint were ~65% of their pre-amputated length at *7dpa*, with the caveat that

precise length measurements were difficult (Supplemental Figure 1A and B).

Nevertheless, it is clear that regenerates did not return to their precut length, and that

longitudinal and tripolar neurons were decreased compared to the uncut starting number

for all regenerates. We conclude that the nerve net of regenerates is reduced in terms of

neuronal number compared to the uncut parental animal.

We observed pharyngeal and tentacular neuron regeneration in 100% of 7dpa

regenerates (N = 73; Figure 1E; Supplemental Figure 3). Further characterization

suggested that tentacular neurons were visible in the forming tentacle buds by 72hpa, and

that pharyngeal neurons were first detected between 48 and 72hpa (Supplementary Figure

3). We concluded that both tentacular and pharyngeal neurons regenerated and their

regeneration coincides with the re-appearance of their corresponding tissues.

To better understand regeneration of longitudinal and tripolar neurons, which are

present in both the *Aboral* remnant and regenerates, we quantified changes in neuronal

number throughout regeneration (Figures 1F-I). Since the number of longitudinal neurons

never returned to the initial starting numbers in any of the animals, we focused our

investigation on the time between when they were cut (*time 0 cut*) and after they fully

regenerated (*7dpa*) (Figures 1 G and I, boxed regions in F and H, respectively).

We initially treated all regenerates equal regardless of size, but observed

differential regeneration of longitudinal neuronal subtypes that appeared to be size

dependent. For the remainder of analysis we will describe the longitudinal and tripolar

regeneration in four different size groups, based on the starting number of longitudinal

neurons present in the uncut animal. We categorized animals as small (20-40 longitudinal

neurons or ~2.5-5 mm), medium (41-60 longitudinal neurons or ~3.5-5.5 mm), medium-

large (61-80 longitudinal neurons or ~5-7.5mm), and large (81-100 longitudinal neurons

or ~6.5-8.5 mm) (Supplemental Figure 1A). Both size and longitudinal number were used

to classify animals for the reasons discussed above.

Longitudinal neurons regenerated in small and medium animals, but not medium-

large or large animals. Between *time 0 cut* and *24hpa* the number of longitudinal neurons

initially decreased in all size groups (Figure 1G, all $p < 0.001$), likely due to increased

apoptosis that occurs over the first 24hpa (Warner et al. 2019). The number of

longitudinal neurons significantly increased between *24hpa* to *7dpa* in small and medium

animals ($p < 0.001$ and $p = 0.001$, respectively). The numbers of longitudinal neurons in

7dpa regenerates were statistically similar to the numbers observed in *time 0 cut* animals

for both small and medium groups (small: $p = 0.77$, medium: $p = 1.00$), suggesting

longitudinal neurons regenerated back to their numbers present in the remnant fragment

at the time of bisection (Figure 1G, red and green lines). No regeneration of longitudinal

neurons was observed in medium-large and large animals between *24hpa* and *7dpa*

(Figure 1G, yellow and blue lines; $p = 0.249$ and $p = 1.00$, respectively). Despite the

differences in regenerative potentials of the longitudinal neurons, there was no significant

difference in the percent length of regenerates at *7dpa* relative to their starting size

(Supplemental Figure 1B).

In contrast to longitudinal neurons, there was no regeneration of tripolar neurons

in any of the animal size groups. The number of tripolar neurons decreased between *time*

0 cut and *24hpa* (Figure 1I, $p < 0.001$) but failed to return to the *time 0 cut* numbers by

7dpa (Figure 1I, $p < 0.001$).

We conclude that longitudinal neurons display differential regenerative potential

that corresponds to the size of the pre-amputated animal, and that tripolar neurons do not

demonstrate any regenerative potential.

Differential regenerative potential of longitudinal neurons depends on the size of the

remnant fragment.

We wondered what was influencing the variable regenerative potential of

longitudinal neurons. All animals, including our large group animals (Supplemental

Figure 4), show increased proliferation in both the ectoderm and endoderm of

regenerating tissue. This observation is consistent with the finding that *Nematostella*

regeneration has been described to be dependent on cell proliferation (Amiel et al. 2015;

Passamaneck & Martindale 2012). Two obvious factors that might influence the

regenerative potential of longitudinal neurons are the animal's pre-amputation size or the

size of the *Aboral* remnant. If the remnant size plays a role in regenerative potential, then

shifting the amputation site in large and small animals to create larger or smaller remnant

fragments should alter the regenerative potential of longitudinal neurons. However, if the

animal's pre-amputation size dictates regenerative potential, then altering the amputation

site should not alter regenerative potential of longitudinal neurons.

When smaller *Aboral* fragments were generated from large animals by aborally

shifting the amputation site to remove 75% of the oral-aboral axis (Figure 2A),

longitudinal neurons regenerated between *24hpa* and *7dpa* (Figure 2C, *p = 0.046*).

Interestingly, the remnant fragment generated by the aboral shift resulted in a similar

number of longitudinal neurons at *time 0 cut* and *24hpa* as the fragment from medium

animals bisected into equal halves (Figure 2D, compare black and faded green lines).

Alternatively, orally shifting the amputation site of medium sized animals to remove only

25% of the oral-aboral axis (Figure 2B) increased the remaining number of remaining

longitudinal neurons at *24hpa* similar to those remaining in medium-large and large

animals bisected into two equal halves (Figure 2 E and F, compare black and faded blue

and yellow lines), and similarly resulted in no longitudinal neuron regeneration between

24hpa and *7dpa* (*p = 1.00*). When the amputation site of small animals is orally shifted

(Figure 2B), the increased *Aboral* remnant fragment has a similar number of remaining

longitudinal neurons as the fragment generated by bisecting a medium animal into two

equal halves (Figure 2G and H, black lines). Longitudinal neurons showed a trend

towards regeneration in small animals with an orally shifted amputation site, similar to

the medium sized animals (Figure 2G, H, compare black and faded green lines) but the

increase in number between *24hpa* and *7dpa* was not statistically significant (*p = 0.469*)

However, the number of neurons at *7dpa* is also not statistically significant from the

number at *time 0 cut*, (*p = 0.119*), but the *time 0 cut* and *24hpa* values are different (*p <*

0.001), suggesting that the trend towards regeneration is real and that variability in

responses of small animals by *7dpa* are the source for lack of statistical significance.

The size of the remnant fragment appears to play a larger role in dictating the

regenerative potential of longitudinal neurons than the starting size of the pre-amputated

animal. Overall, these data suggest there are at least three regenerative responses for

neuronal subtypes in *Nematostella.* Class I neurons that regenerate 100% of the time

following amputation (tentacular and pharyngeal *NvLWamide-like* neurons); Class II

neurons, which do not regenerate by *7dpa* (tripolar *NvLWamide-like* neurons); and Class

III neurons that have differential regenerative abilities likely controlled in part by the size

of the remnant fragment (longitudinal *NvLWamide-like* neurons).

Longitudinal and tripolar neuron numbers are dynamic and scale with size in adult

animals.

The reduced size of the regenerated nerve net following amputation led us to

investigate scalability. The number of longitudinal and tripolar neurons positively

correlate with animal length during growth from juvenile polyps into adult stages

(Havrilak et al. 2017). Because regenerates are smaller than pre-amputated animals, and

the number of longitudinal and tripolar neurons are reduced in regenerates (Figures 1 and

2; Supplemental Figure 2), we wondered if longitudinal and tripolar numbers might

fluctuate in adult animals as they grow and shrink in size. *Nematostella* increase in size

when fed (Figure 3 A, B, and G)(Hand & Uhlinger 1992; Layden et al. 2016), and our

observations demonstrate that they decrease in size when starved (Figure 3A, B, and G).

Thus, we set out to determine if tripolar and longitudinal neuronal numbers are dynamic

in adults that are increasing and decreasing in size.

To compare responses, we performed two replicate experiments that varied the

number of starting longitudinal and tripolar neurons and therefore the starting size of the

animals (Figure 3, smaller animals; Supplemental Figure 5, larger animals). The initial

experimental design was to track neuronal number in animals over fourteen weeks.

Animals would be either starved or fed for seven weeks, then the feeding regimen would

be changed for the remaining seven weeks. However, we found our largest animals that

were initially fed (the feed-starve treatment group) became too large to accurately count

within the time constraints of our paralytic after 3 weeks into our feeding regimen. We

therefore switched the animals to starvation for the remaining 11 weeks in both

independent experiments (Figure 3; Supplemental Figure; black lines).

After the initial 3 weeks of feeding, the fed then starved animals (Figure 3 and

Supplemental Figure 5, black lines) had significantly increased in length (Figure 3A and

G, $p = 0.008$), and the number of both longitudinal neurons (Figure 3C and D;

Supplemental Figure 5A and B; $p < 0.001$ for both) and tripolar neurons (Figure 3E and

F, $p = 0.32$; Supplemental Figure 5C and D, $p < 0.001$) significantly increased. Following

the switch to a starvation regime, the fed then starved animals significantly shrunk in size

after 11 weeks (Figure 3A and G, $p = 0.008$), and lost a significant number of

longitudinal (Figure 3C and D, $p = 0.003$; Supplemental Figure 5A and B, $p = 0.001$) and

tripolar neurons (Figure 3E and F, $p = 0.031$; Supplemental Figure 5C and D, $p < 0.001$).

Animals placed on the opposite feeding regime had opposite effects on their size

and neuronal number (Figure 3B-G; Supplemental Figure 5A-D, blue lines). The animals

that were initially starved for 7 weeks (starved then fed group) got smaller in size (Figure

3B) and their measured length showed a trend toward a shorter length (Figure 3G, $p =$

0.098), suggesting the animal is slower to shrink in size than grow. Starved then fed

animals showed significant decreases in longitudinal neurons during the starvation

regime (Figure 3C and D, $p = 0.004$; Supplemental Figure 5A and B; $p = 0.031$). Tripolar

neurons significantly decreased in the animals that were initially smaller to start (Figure

3E and F, $p = 0.029$), and their decrease approached significance in animals that were

larger to start (Supplemental Figure 5 C and D, $p = 0.094$). When switched to feeding, the

starved then fed animals significantly increased in size (Figure 3B, 3G, $p < 0.001$) and

both neuronal subtypes significantly increased in number (Figure 3C-F, Supplemental

Figure 5 A-D; $p \leq 0.001$ for all).

We conclude that *Nematostella* scales at least part of its nervous system by

altering neuronal subtype number coincidently with increases or decreases in its size.

Increased Caspase 3 activity in neuron-like cells is observed in starved animals

Increase in neuronal number is due to birth of new neurons, and increased

apoptosis is the most likely cause for decrease in neuronal number. To assess if the loss

of neurons in starved animals is correlated with increased apoptosis, we stained fed and

starved animals with an α−Caspase 3 antibody that recognizes the p17 fragment of active

Caspase-3, a potential marker of apoptosis. Caspase-3 is an effector of the apoptosis

pathway (Elmore 2016) and has been identified as a key mediator in neuronal

programmed cell death (D'Amelio et al. 2012; Kuida et al. 1996). The *Nematostella*

genome possesses at least three *caspase 3* orthologs (Moya et al. 2016). Using the active

Caspase-3 antibody on fed *versus* 3-week starved animals, we observed activated

Caspase 3 staining in 15 out of 15 of starved animals (Figure 4), which is in contrast to

the Caspase-3 staining only being observed in 2 of the 16 fed animals. Activated

Caspase-3 was detected in cells located along the longitudinal tracts of starved animals

(Figure 4E-H, arrow), as well as in cells throughout the body column of starved animals

(Figure 4E-H, arrowhead). Interestingly, some of the activated Caspase3 positive cells

had a neuronal-like morphology, suggesting that neuronal death does occur during

periods of starvation.

Our data suggest that apoptosis is not only playing a role in the loss of cells,

including some neuron populations in the starved animals, but also that apoptosis is

restricted primarily to the physa, which is the most aboral region of the body. We

conclude that starvation induces an increase in Caspase 3 activity in neuronal and non-

neuronal cells, which suggests that decreases in animal size are due to induced cell death.

Scaled down nervous systems maintain the ability to respond to external cues

Lastly, we wondered if the *Nematostella* nervous system retained function in each

scale state. However, *Nematostella* are known to retract when challenged with a

mechanical stimulus. Thus, we decided to determine if *Nematostella* retained the ability

to respond to mechanical cues in different scale states. Because there are no quantified

behavioral assays available yet in *Nematostella*, we developed an action-reaction Poking

assay using a toothpick to induce a reaction response. Poking an animal near the tentacles

induced a contraction of the animal away from the stimulus (see Supplemental Movies 1-

6). Each animal was poked three times in the most oral region of the body, and a positive

response was scored when the animal retracted their tentacles and/or their body at least

once in response to being poked. We observed variable responses to the Poking assay

(anywhere from entire body retraction to only retracting 1-2 tentacles). Therefore any

attempt to move their body away from the toothpick was classified as a positive response.

We first examined the behavioral response in pre- and post- regenerated animals

(Figure 5; Table 1). For all the cases (N = 8), the same animal responded to the Poking

assay both pre-bisection (Figure 5A-D) and *7dpa* (Figure 5E-H; Table 1). After

regeneration, the nerve net and the number of neurons it contained were decreased

compared to the initial non-bisected body (Figures 1 and 2; Supplemental Figure 2). The

maintenance of the behavioral response in the regenerated animal suggests that the

reduction in the nervous system size did not limit the animal's ability to coordinate a
response to external mechanical cues.
We next performed the Poking assay in fed *versus* starved animals (Figure 6;
Table1). We observed that animals starved (6-7 weeks) or fed (3 weeks) both responded
to the Poking assay (Figure 6A-E, H-M; Table 1). As expected, we observed a significant
decrease of the numbers of longitudinal and tripolar neurons in the starved animals
(Figures 6F and G) and a significant increase in these neuronal numbers in the fed
animals (Figure 6N and O). We conclude that retraction in response to a mechanical cue
behavior is not dependent on the size of the nervous system. These preliminary findings
suggest that the neurocircuitry for at least some behaviors are retained regardless of the
overall size of the nerve net.
353
354 **Discussion:**
355
Variability in regenerative potential of individual cell types during whole-body axis
regeneration.
This study represents the first study we are aware of to track neuronal subtype
regeneration *in vivo* in individuals undergoing whole-body axis regeneration. We found
that individual neuronal subtypes have differential potential to regenerate new neurons,
and this is based in part on the size of the remnant fragment. We therefore grouped
neurons into three classes based on regenerative response. Class I neurons that
regenerated 100% of the time, Class II neurons that did not regenerate following
amputation, and Class III neurons, which are perhaps the most interesting, because they
are capable of regenerating under some but not all circumstances. We were initially
concerned that the lack of regeneration observed in Class II and sometimes Class III
neurons was due to delayed regeneration, particularly in larger animals. Regeneration
occurred within seven days regardless of the size of the animal. Larger animals would
need to regrow more tissue to go from ~50% (following amputation) to ~65% pre-cut
length (after regeneration) (Supplemental Figure 1). To accomplish this, they would need
to proliferate more rapidly than smaller animals and an increase in proliferation rates
could delay longitudinal differentiation, whereas tripolar regeneration could be delayed in

all cases. We did check regenerates at 12 and 14dpa and we saw no change from the

neuronal numbers observed at 7dpa (data not shown), suggesting that there is no delay in

longitudinal specification in larger animals nor any delay in tripolar regeneration. We

argue that there is fundamental difference in why longitudinal neurons regenerate in

smaller but not larger remnant fragments, and that Class II and Class III neurons are both

part of the normal regenerative response.

It is likely that all three neuronal classes described here are present in other

animals. Class I neurons were neurons that were present in structures that needed to

completely reform in regenerates. Within planarian whole-body axis regeneration, Class I

neurons would likely be any of the neuronal subtypes present specifically in the brain or

eyes that needed to regenerate following amputation of anterior regions. It is not clear

how widespread the presence of Class II neurons is in animals, because tracking of

individual neuronal subtypes during regeneration in the same animal has not been

reported in planarians or *Hydra*. Given that both planarians and *Nematostella* have been

shown to have nervous systems that scale with size (Figures 3; Supplemental Figure 5)

(Takeda et al. 2009), it is likely that Class II neurons are not a *Nematostella* specific

phenomenon. It should be noted that Class II neurons are capable of regenerating in some

regenerative paradigms. Tripolar neurons that were lost as animals shrunk during

starvation repopulated in animals that regrew once they were switched back to feeding

(Figure 3 and Supplemental Figure 5). This suggests that the regenerative potential of any

neuronal subtype is not uniform across all regeneration paradigms. There is evidence for

Class III-like regenerative potential in other animals. For example, populations of

dopaminergic neurons in the brains of adult zebrafish have been shown to be diverse in

their regenerative capacity (Caldwell et al. 2019). Based on the fact that regenerates are

smaller than the pre-cut animal, that degrees of nervous system scaling is likely common

in animals, and that some amputations result in complete loss of some neuronal subtypes,

we predict that all three neuronal classes are common to all animals that undergo whole-

body axis regeneration.

***Nematostella* nervous system scales with animal length.**

Nematostella modulated the size of its nerve net by increasing or decreasing

neuronal number in response to feeding or starvation periods, and these changes were

reversible (Figure 3). The correlation between neuronal number and body size suggests

that *Nematostella* scales its nervous system with its body size. Importantly, *Nematostella*

maintains normal behaviors in each scale state (Figures 5 and 6). The potential scalability

of nervous systems has been suggested by studies using molecular markers to track

neuronal subtypes in planarians. Specific cell types, including visual cells and neural cells

in the brain, were shown to correlate with body length, and there is also a constant ratio

maintained between several types of neurons in the brain regardless of overall number

(Takeda et al. 2009). Although this planarian study was performed in fixed animals, their

findings are consistent with the dynamic and scalable nervous system found in

Nematostella. It should be noted that we did not address, and do not we rule out, that the

Nematostella nervous system may also scale by modulating connectivity of existing

neurons.

During both regeneration and body scaling, *Nematostella* retains normal body

form (Figures 1 and 3), the ability to respond to a mechanical cue (Figures 5, 6), and the

ability to eat (Passamaneck & Martindale 2012; Layden et al. 2016). Additionally, even

with a smaller nerve net, we can infer that these animals are telling the difference

between a potential predator (toothpick) and prey (brine shrimp), because while the

animals responded to the Poking assay with a defensive retraction, they readily capture

and ingest the food. Taken together these observations suggest that the nervous system is

functional in all scale states.

Nervous system scalability may influence regenerative potential of neuronal

subtypes.

The fact that nervous systems scale with size in *Nematostella,* and that

regenerates are smaller than pre-cut animals, suggests that the differential regenerative

responses of individual neurons are linked to the scale state of the regenerate. The two

neuronal subtypes that we confirmed have reduced numbers in regenerates are the

longitudinal (Class III) and tripolar neurons (Class II). Coincidently, the total number of

neurons for both of these neuronal subtypes correlate with overall length and scale in

number as animal length increases or decreases (Figure 3; Supplemental Figure

5)(Havrilak et al. 2017). While it is logical that regenerates that are smaller than the pre-

cut animal have less tripolar and longitudinal neurons, it is not clear why tripolar neurons

do not show regenerative potential, but longitudinal neurons do under some

circumstances. One possible explanation for why tripolar neurons do not regenerate is

that the number of neurons present in the *Aboral* remnant are within an acceptable range

for the scale state of the regenerate. As a result, tripolar neurons do not need to increase

in numbers as part of a regenerative response. An alternative explanation is that

regeneration of tripolar neurons may not be necessary because they may function

similarly to tentacular neurons. Tripolar and tentacular neurons are both ectodermal

neurons that share a similar morphology (Havrilak et al. 2017). The reduced size of

regenerates suggests that the majority of regenerated tissue is contributing to pharyngeal

and tentacular structures. If tentacular neurons have a similar function to the tripolar

neurons they may compensate until the regenerate starts to feed and tripolar neurons

begin populating the body column with growth. However, it should be noted that tripolar

and tentacular neurons are specified at different times during development (Havrilak et

al. 2017), and the tissues they arise from have distinct gene expression profiles of

transcription factors known to pattern neuronal fates in *Nematostella* and traditional

model systems (Layden et al. 2010; Sinigaglia et al. 2013; Leclère et al. 2016; Marlow et

al. 2013; Röttinger et al. 2012). Thus, it isn't clear if they are similar neurons that are

patterned using variable molecular programs during development, or if they are distinct

neuronal fates.

The differential response of longitudinal neuron regeneration observed in

different sized remnant fragments is interesting. We have thought of two hypotheses to

explain the response. The first hypothesis is simply that the scale state of longitudinal

neurons is different than that of tripolar neurons, and that in smaller remnants the number

of longitudinal neurons is below a minimum threshold, which results in induction of

regenerative response after amputation. In larger fragments, the number of longitudinal

neurons is above the minimum threshold for the new scale state of the nervous system,

and thus no regenerative response is induced. The alternative hypothesis is that there are

differences in regenerative mechanisms between large and smaller animals. Although cell

proliferation is required for *Nematostella* oral regeneration (Passamaneck & Martindale

2012), some remodeling of remnant tissue occurs (Amiel et al. 2015). These findings

suggest *Nematostella* uses a hybrid of cell proliferation and morphallaxis to regenerate.

EdU labeling suggests that larger animals are still regenerating using both cell

proliferation and morphallaxis, but perhaps they shift mechanisms to favor morphallaxis

and smaller animals favor cell proliferation. Future experiments to help distinguish

between potential different mechanisms of regeneration based on animal size should be

done to better understand *Nematostella* regeneration. Our limited studies favor the

threshold scale model. First, we did not observe additional longitudinal neurons in

regenerates from large animals bisected in half at 14dpa. Even if large animals do rely

more heavily on morphallaxis, they are still increasing in size at equal proportions to

small animals. Thus, the prediction is that longitudinal neurons should increase in number

unless there is a sufficient number of neurons to be at the "low" end of the acceptable

scale state in the regenerate. Additionally, when cut sites were shifted the regenerative

response of the remnant tissue was dictated by its size. This suggests that there are

particular scale states that need to be maintained in different sized animals. Regardless of

the exact mechanism, the data support the hypothesis that once we identify the key ratio

of neuronal number to length for a particular neuronal subtype, it should be possible to

predict its regenerative response based on the size and number of neurons present in the

remnant fragment.

Scalability of adult *Nematostella* nerve net

Nematostella must have a mechanism to sense and regulate the ratio of neuronal

number to body size. Possible mechanisms include the nervous system self-regulating

subtype numbers autonomously, or that neural numbers are influences by signaling from

neural and other tissues/cell types within the body. It is unlikely that changes in size also

influence the amount of other cell types besides neurons. Non-neuronal cells had

detectable activated Caspase-3 (Figure 4), which suggests other cell types decrease as the

animal length decreases during starvation. This is similar to observations in planarians

where an overall increase or decrease in cell number across multiple cell types

corresponds to larger or smaller length, respectively (Baguna et al. 1981; Romero &

BAGUÑÀ 1991; Takeda et al. 2009; Oviedo et al. 2003). The global change of other

neuronal cell types suggests that signaling between cell types may coordinate scaling of

the entire animal simultaneously. One example of regulation of cell number being

mediated by signaling between different cell types is the regulation of hematopoietic stem

cell number by Notch and BMP signaling from osteoblasts (Zhang et al. 2003; Bowers et

al. 2015; Calvi et al. 2003). However, microglia have been shown to self-regulate their

numbers during periods of resolution following neurodegeneration in both zebrafish (van

Ham et al. 2014) and mice (Tay et al. 2017). This occurs via cell egress and apoptosis

(Tay, 2017), and phagocytosis of disease state microglia undergoing apoptosis by other

viable/healthy microglia has also been demonstrated (van Ham, 2014). At this point it is

unclear if scaling of the nervous system occurs autonomously, is directed by signaling

from other tissues, or via both mechanisms. Future work characterizing *Nematostella*

growth and shrinking and the scaling responses of differentiated cell types is an exciting

potential area of research.

Conclusion:

The ability to observe and quantify individual neuronal subtypes makes

Nematostella a powerful model to investigate neurogenesis in the context of both

regeneration and whole-body scaling. Our data suggest that the regenerative responses of

neuronal subtypes in *Nematostella* are variable and depend both on individual subtypes

and the size of the remnant fragment. These data imply that there is differential

deployment of the molecular pathways that specify specific subtypes that may be

dependent on the scale-state of the nervous system in the regenerate. How *Nematostella*

controls the generation and incorporation of new neurons into the existing nerve net, and

how they scale the numbers of these neurons to body size in real time remains an

unknown, but both are exciting areas for future investigation. Lastly, our data suggest that

experiments aimed at uncovering the molecular mechanisms regulating neural

regeneration in *Nematostella* require careful consideration of the differential regenerative

potential of individual neuronal subtypes when designing experiments and interpreting

results.

Methods:

Animal care

All animals were kept at room temperature and fed rotifers or crushed brine

shrimp until they reached the young juvenile adult stage, at which time they were

transferred to 17°C and switched to solely brine shrimp. All juvenile adult and adult

animals were housed at 17°C in the dark and fed brine shrimp 4x/week, unless otherwise

noted for experimental purposes. *Nematostella* were maintained in 1/3x artificial sea

water, (*Nematostella* medium) at a pH of 8.1-8.2 that was changed once per week.

Nematostella were relaxed in 7.14% (wt/vol) $MgCl_2$ in *Nematostella* medium for

~10 minutes prior to quantification, imaging, or bisection. The lengths of relaxed animals

were measured in mm with a ruler placed under a Petri dish containing the animal.

Quantification of NvLWamide-like::mCherry-expressing neurons

NvLWamide-like::mCherry tripolar and longitudinal neurons were counted on live

animals under a dissection microscope (Nikon SMZ1270), on live animals placed under a

slide with a raised coverslip on a compound microscope (Nikon NTi), or on full-length

images of *Nematostella*. All quantifications of longitudinal and tripolar neurons were

obtained by counting the neurons from 2-4 segments on each individual animal that were

then averaged together.

Regeneration experiments

Longitudinal and tripolar neurons were quantified first on the uncut animal, then

immediately following bisection (time 0 cut), 24 hours post amputation (hpa), and 7 days

post amputation (dpa). Animals were either bisected in half with the cut site at oral-aboral

midpoint (~50% body removed), bisected below the pharynx ("oral shift", ~25%

removed), or bisected at the aboral end ("aboral shift", ~75% removed). Following

quantification and/or bisection, the remnant aboral fragment was washed with

Nematostella medium, then placed in the dark at room temperature. Prior to

quantification, animals were starved 4 days (Amiel et al. 2015; Passamaneck &

Martindale 2012). Animals were not fed for the duration of the experiment.

Feed/Starve experiments

For each animal, the number of longitudinal and tripolar neurons were counted at Time 0, at the time of their feeding regime switch (starved to fed, 6-7 weeks later; fed then starved 3 weeks later), and at the end of the 14 week experiment. All animals were maintained on the same normal feeding schedule until Time 0 of the experiment. At this time animals were randomly assigned to either the starve-feed group or feed-starve group, then they were imaged, and their initial starting neural counts were taken. Animals on the feeding regime were fed brine shrimp 2x/week plus oyster 1x/week, and were given weekly water changes.

Poking assay

Animals were transferred to a depression slide containing *Nematostella* medium. Once the animal had relaxed its tentacles a toothpick was slowly inserted into the medium so as to not disturb the animal prematurely. The animal was then "poked" with the tip of the toothpick at the oral region. The entire assay was video recorded and analyzed to determine the animal's response. Following the "Poking assay", the animal was left alone in the dish until it relaxed again, upon which the Poking assay was performed again. The assay was performed 3 consecutive times. A positive response was determined if the animal responded to being touched with a toothpick by retracting some or all of its tentacles, scrunching or retracting in its mouth (oral region) or retracting its entire body. To be considered responsive the animal had to react in at least 1 out of the 3 poking assays performed.

EdU Stain

EdU incorporation was carried out as previously described with modifications (Fritz et al. 2013). Animals were incubated with 330uM EdU in *Nematostella* medium for 30 minutes, then fixed as previously described (Fritz et al. 2013) and dehydrated to 100% MeOH and stored at -20 °C. Following rehydration, samples were permeabilized in PBT (PBS with 0.5% Triton X-100, Sigma) for ~2 hours. Animals were then treated with the reaction cocktail (prepared following the Click-It protocol provided by the manufacturer,

with the exception of doubling the amount of Alexa Fluor azide) at room temperature in

the dark for 20 minutes. Following 5 long washes in PBT the animals were labeled with

propidium iodide for 1 hour while rocking, washed 5X in PBT and finally transferred to

90% glycerol for imaging.

Immunohistochemistry for activated Caspase-3

Preparation for staining was carried out as previously described (Amiel et al.

2015). To analyze for activated Caspase-3 (apoptosis stain), we used a polyclonal rabbit

active Caspase-3 antibody (ABCAM, ab2302). Secondary Alexa Fluor 488 donkey anti-

rabbit (Life Technologies, A21206) was used.

Imaging

Live images of *NvLWamide-like::mCherry*-expressing animals were taken on a

Nikon NTi with a Nikon DS-Ri2 color camera with Nikon Elements software. Large full-

length images are comprised of several images stitched together. Confocal images were

captured using a Zeiss LSM 880 with LSM software and processed using Imaris 8.4.1

software (Bitplane LLC) to create 3D images from serial optical sections (z-stacks).

Images were then cropped and assembled using Photoshop and Illustrator (Adobe).

Live video was captured on a Nikon Eclipse E1000 microscope with Nikon

Imaging software. Movies were processed with ImageJ/Fiji software and screen shots

were obtained from individual frames, then cropped and assembled using Adobe

Photoshop and Illustrator.

Statistical analysis

For the regeneration and feed/starve experiments, mixed analysis of variance

(ANOVA) tests were used to analyze each dependent variable (i.e. the number of

longitudinal and tripolar neurons quantified). For both experiments, each observation of

an individual over time served as the within-subject factor (Regeneration experiment:

time 0 cut, 24hpa, 7dpa; Feed/starve: week 1, week of feeding regime switch, week 14).

The between subject factors used were animal size category (small: 20-40mm, medium:

41-60mm, medium-large: 61-80mm, large: 81-100mm) and assigned feeding group

(Feed-Starve or Starve-Feed) for the regeneration and feed/starve experiments

respectively. A mixed ANOVA was also used to analyze changes in oral-aboral length

during the feed/starve experiment, with each observation of an individual serving as the

within-subject factor (week 1, week of feeding regime switch, and week 14) and assigned

feeding group as the between-subject factor (Feed-Starve or Starve-Feed). Data used for

ANOVA tests met assumptions of normality and homogeneity. Greenhouse-Geisser

corrected F-statistics were reported if sphericity was violated (i.e. Figure 1I). When

omnibus ANOVA results were significant, post-hoc testing was performed using

Bonferroni corrections in order to maintain tight control over the type 1 error rate. Effect

sizes are reported as partial eta^2 (η_p^2). Full AVONA model results are reported in

Supplemental Document 1. Paired t-tests were used to compare individual changes in the

number of neurons or length at two time points. All data were analyzed using SPSS

version 21.0.

Acknowledgements

We would like to thank David Balli and Omar Ahmed for providing help with R

software and construction of the line graphs. We also thank Dylan Faltine-Gonzalez,

Noor Baban, and Brianna Maslonka for their help with maintenance of the animal stocks.

This work was funded by the National Institutes of Health Award Number

R01GM127615.

Competing Interests Statement

The authors declare that they have no financial or non-financial competing interests

Figure 1: Differential regeneration of *NvLWamide-like* neurons depends on the specific neuronal subtype. (A) Schematic for oral bisection at ~50% body length. (B) Quantification of the number of longitudinal neurons along the oral-aboral axis in each of the 4 equal quarters. Each line represents an individual animal. (C) Uncut *NvLWamide-like::mCherry* expressing animal. (D) Same animal as C at time 0 cut. (D) Same animal as C and D at 7 days post amputation (dpa), with regenerated oral structures. (F-I**)** Quantification of longitudinal and tripolar neurons before amputation (uncut), immediately following bisection at the O-A midline (time 0 cut), 24hpa, and at completion of regeneration (7dpa) in small (red), medium (green), medium-large (orange), and large (blue) sized animals. The numbers of longitudinal neurons (F) or tripolar neurons (H) never regenerate back to their original pre-cut numbers. (G) Longitudinal neurons regenerate back to numbers similar to time 0 cut in the small and medium sized animals, but do not regenerate in the larger animals by 7dpa. (I) Tripolar neurons do not regenerate by 7dpa. N = 40 animals. Scale bar = 1mm. ((G) Mixed ANOVA: interaction effect between animal size and observation time: $F_{6,70} = 10.27$, $p < 0.001$, $\eta_p^2 = 0.47$; (I) Mixed ANOVA: main effect of observation time Tripolar: $F_{1.4,46.9} = 20.70$, $p = < 0.001$, $\eta_p^2 = 0.38$). Data points in F-I represent mean ± SEM.

Figure 2: Regeneration of longitudinal neurons depends on the size of the

regenerating fragment. (A-B) Schematics demonstrating the shifts in cut sites used to

obtain remnant fragments of varying sizes. (C-H) Quantification of longitudinal neurons

in regenerates with shifted cut sites at time 0 cut, 24hpa, and 7dpa. Regenerates from an

aborally-shifted cut site in large animals regenerated their longitudinal neurons (C),

resembling regenerates of small and medium sized animals cut at the midline of the O-A

axis (D). Regenerates from an orally-shifted cut site in medium animals did not

regenerate their longitudinal neurons (E), resembling regenerates from large animals cut

at the midline of the O-A axis (F). Regenerates from an orally-shifted cut site in small

animals still regenerated their longitudinal neurons (G), resembling regenerates from

medium animals cut at the midline of the O-A axis (H). N = 8 large; 13 medium; 13

small. Arrows indicate longitudinal neurons and arrowheads indicate tripolar neurons

(Repeated measure ANOVA comparing observation times: (C) Large, $F_{2,14} = 5.58$, $p =$

0.017, $\eta_p^2 = 0.44$; (E) Medium, $F_{2,24} = 21.67$, $p < 0.001$, $\eta_p^2 = 0.64$; (G) Small, $F_{2,24} =$

8.56, $p = 0.002$, $\eta_p^2 = 0.42$). Data points in C-H represent mean ± SEM.

Figure 3: The *Nematostella* nerve net scales with changes in length. (A-B)

Visualization of *NvLWamide-like* neurons in relation to body size during periods of

feeding and starvation. (C-F) Quantification of *NvLWamide-like* neurons during periods

of growth and starvation with both mean (C, E) and population (D, F) data presented.

Animals fed for 3 weeks rapidly grew longer in length (A, 3 weeks fed), and numbers of

both longitudinal (C and D, black lines) and tripolar (E and F, black lines) neurons

significantly increased. Following 11 weeks starvation, the same animal then shrunk in

size (A, 11 weeks starved) and the numbers of both longitudinal and tripolar neurons

significantly decreased (C-F, black lines). Animals initially starved for 7 weeks shrunk in

size (B, 7 weeks starved) and number of both longitudinal (C and D, blue lines) and

tripolar (E and F, blue lines) significantly decreased. Upon being switched to feeding

regime for 7 weeks the same animals grew longer (B, 7 weeks starved) and the numbers

of both longitudinal and tripolar neurons significantly increased (C-F, blue lines). Oral

end is to the left. Scale bar = 1mm. N = 5 animals per treatment group. Arrows indicate

longitudinal neurons and arrowheads indicate tripolar neurons. (Mixed ANOVA of

interaction between assigned feeding regime and observation time: (C) Longitudinal,

$F_{2,16} = 102.01$, $p < 0.001$, $\eta_p^2 = 0.93$; (E) Tripolar, $F_{2,16} = 48.60$, $p < 0.001$, $\eta_p^2 = 0.86$).

(G) The measured length of the animals significantly decreased during the starvation

period and increased during the feeding period (Mixed ANOVA: interaction effect

between assigned feeding regime and observation time: $F_{2,34} = 25.75$, $p < 0.001$, $\eta_p^2 =$

0.60). Data points in C, E and G represent mean ± SEM.

752
753
754

Figure 4: Increased apoptosis in 3-week starved animals. (A-D) Normally fed animals
did not have activated Caspase-3 staining in the most aboral region in 14 out of 16
animals examined (2/16 had positive staining, data not shown). (E-H) Animals that were
starved for 3 weeks had positive Caspase-3 visible in the most aboral region in 15 out of
15 animals examined. (F-H) Caspase-3 was located along the endodermal longitudinal
tracks in the most aboral region (arrow) and in the aboral body column (arrowhead).

774
775
776
777 **Figure 5: Response to poking assay is maintained in regenerating animals.** (A-D)
778 *Nematostella* respond to the touch stimulus by retracting its tentacles when touched with
779 a toothpick (N = 8/8). (E-H) At 7 days post amputation the same animal again responded
780 to being poked with a toothpick by retracting its tentacles (N = 8/8).

Figure 6: Starved and fed animals respond to the Poking assay by retracting their tentacles and body. (A-E) A normally fed animal responded to the touch stimulus from a toothpick by retracting its tentacles and oral region (N = 17/19). (D-E) Following 6-7 weeks starvation, the same animal again responded to being touched with a toothpick by retracting its tentacles and oral region (N = 17/19). (F-G) Quantification of the number of longitudinal and tripolar neurons in individual animals following each poking experiment. The numbers of both longitudinal (I: t_{16} = 7.97, $p < 0.001$, η_p^2 = 0.80) and tripolar (J: t_{16} = 6.99, $p < 0.001$, η_p^2 = 0.75) neurons significantly decreased after 6-7 weeks starvation. (H-J) A normally fed animal responded to the touch stimulus from a toothpick by retracting its tentacles and body (N = 13/13). (K-M) Following 3 weeks feeding, the same animal again responded to being touched with a toothpick by retracting its tentacles and oral region (N = 12/13). (N-O) Quantification of the number of longitudinal and tripolar neurons in individual animals following each poking experiment. The numbers of both longitudinal (G: t_{12} = -9.15, $p < 0.001$, η_p^2 = 0.88) and tripolar (H: p = t_{12} = -12.76, $p < 0.001$, η_p^2 = 0.93) neurons significantly increased following 3 weeks feeding.

826   **Table 1:** Summary of the analysis for the action-reaction Poking assay in regenerated,

827   starved, and fed animals.

Treatment	Timepoint	Number of animals that responded to the touch stimulus	Percent of animals that responded to the touch stimulus
Regeneration	Uncut	8/8	100%
	7dpa	8/8	100%
Starved	Time 0	17/19	89%
	6-7 weeks starved	17/19	89%
Fed	Time 0	13/13	100%
	3 weeks fed	12/13	92%

Supplemental Figure 1: Regenerated *Nematostella* did not return to their original

length by 7dpa. Length was measured with a ruler under a dissection microscope at the

start of the experiment before oral bisection ("Uncut"). The animals never re-grew back

to their original length, and there was no difference in the percent length of the

regenerates relative to their body size by 7dpa. (A) Percent length for animals bisected in

half. N = 11 small; 8 medium; 6 medium-large; 6 large (One way ANOVA: $F_{3,27} = 0.23$,

$p = 0.875$, $\eta_p^2 = 0.025$). (B) Population data showing lengths for individual animals

before they were cut (uncut) and 7dpa. Small animals are represented in red, medium in

green, medium-large in orange, and large in blue ($t_{30} = 8.65$, $p < 0.001$, $\eta_p^2 = 0.71$). N =

31 animals. (C) Percent length for animals with the shifted cut sites. N = 13 small oral

shift; 11 medium oral shift; 7 large aboral shift (One way ANOVA: $F_{2,17} = 8.12$, $p =$

0.003, $\eta_p^2 = 0.488$). Bars in A and C represent mean ± SEM.

Supplemental Figure 2: Quantification of neuronal numbers during oral regeneration per individual regenerating animal. Population data showing the time course of the average number of (A) longitudinal neurons and (B) tripolar neurons per 2-3 radial segments in individual regenerating animals bisected in half. Neuronal numbers do not return to levels present in the uncut animal. Quantifications were taken before cutting (uncut), time 0 cut, 24hpa, and 7dpa. N = 40 animals. Data represented here is the same dataset grouped by starting size and averaged in Figure 1.

Supplemental Figure 3. Regeneration of oral structures and neurons in

NvLWamide-like::mCherry **animals following bisection along the O-A axis.** (A)

Regenerated oral structures, including a mouth, pharynx (arrow), and tentacles at 7dpa.

(A') New neurons are observed in the regenerated oral structures, including pharyngeal

(arrow) and tentacular (arrowhead) neurons. (B) Remnant fragment at 24hpa. No new

structures are observed at this timepoint. (C) Tentacle buds are visible in the remnant

fragment at 48hpa. Longitudinal (green arrowhead) and tripolar (purple arrowhead)

neurons are observed. Regenerated pharyngeal neurons are also present in the

regenerating pharynx (orange arrowhead) at 48hpa. (D) Pharyngeal (orange arrowhead)

and tentacular (red arrowhead) neurons are clearly visible in these regenerating structures

by 72hpa. (E) Clearly regenerated pharyngeal (orange arrowhead) and tentacular (red

arrowhead) neurons are seen at 96hpa. Longitudinal neurons (green arrowhead) also

populate the regenerated tissue, but if these are regenerated neurons or from the remnant

is undetermined. Asterisks indicate oral opening.

891
892
Supplemental Figure 4: Detection of proliferating cells in "large" *Nematostella*

during oral regeneration. (A) EdU (green) is detected at 48hpa in the oral-most region

of "large" regenerating animals bisected in half. (B) EdU overlaid with propidium iodide

(PI, red) stain in the same animal as A, with yellow color showing co-expression. (C)

Higher magnification image of the same animal, showing EdU detection in both the

ectoderm and endoderm, suggesting both tissue layers proliferate during oral

regeneration. N = 3 large animals. Scale bars for A and B = 200um; C = 50um.

Supplemental Figure 5: Independent experiment demonstrating *Nematostella* nerve net scales with changes in length. (A-D) Quantification of *NvLWamide-like* neurons during periods of feeding and starvation with both mean (A,C) and population (B,D) data presented. The numbers of both longitudinal (A and B, black lines) and tripolar (C and D, black lines) neurons significantly increased with 3 weeks of feeding, and then both longitudinal and tripolar neurons significantly decreased with 11 weeks starvation (A-D, black lines). The numbers of both longitudinal (A and B, blue lines) and tripolar (C and D, blue lines) significantly decreased following 6 weeks starvation. Upon being switched to feeding regime for 8 weeks the numbers of both longitudinal and tripolar neurons significantly increased (A-D, blue lines) (Mixed ANOVA: interaction effect between assigned feeding regime and observation time: (A) Longitudinal, $F_{2,16} = 38.88$, $p < 0.001$, $\eta_P^2 = 0.83$; (C) Tripolar, $F_{2,16} = 206.13$, $p < 0.001$, $\eta_P^2 = 0.96$). Data points in A and C represent mean ± SEM.

938 **References**
939

Amiel, A.R. et al., 2015. Characterization of Morphological and Cellular Events
Underlying Oral Regeneration in the Sea Anemone, Nematostella vectensis.
International Journal of Molecular Sciences, 16(12), pp.28449–28471.

Baguna, J., Romero, R.Romero, 1981. Quantitative anlaysis of cell types during growth,
degrowth, and regeneration in the planarians Dugesia mediterranea and Dugesia
tigrina. *Hydrobiologia*, 84, pp.181–194.

Bossert, P.E., Dunn, M.P. & Thomsen, G.H., 2013. A staging system for the regeneration
of a polyp from the aboral physa of the anthozoan Cnidarian Nematostella vectensis.
Developmental Dynamics, 242(11), pp.1320–1331.

Bowers, M. et al., 2015. BLOOD582924 2678..2688. *Blood*, 125(17), pp.2678–2688.

Busengdal, H. & Rentzsch, F., 2017. Unipotent progenitors contribute to the generation
of sensory cell types in the nervous system of the cnidarian Nematostella vectensis.
Developmental biology, 431(1), pp.59–68.

Caldwell, L.J. et al., 2019. Regeneration of Dopaminergic Neurons in Adult Zebrafish
Depends on Immune System Activation and Differs for Distinct Populations. *The
Journal of Neuroscience*, 39(24), pp.4694–4713.

Calvi, L.M. et al., 2003. Identification of the haematopoietic stem cell niche and control
of the niche size. *Nature*, 425(6960), pp.841–846.

Cebrià, F. et al., 2002. Dissecting planarian central nervous system regeneration by the
expression of neural-specific genes. *Development, Growth & Differentiation*, 44(2),
pp.135–146.

Chen, C.C.G., Wang, I.E. & Reddien, P.W., 2013. pbx is required for pole and eye
regeneration in planarians. 140(4), pp.719–729.

Currie, K.W. & Pearson, B.J., 2013. Transcription factors lhx1/5-1 and pitx are required
for the maintenance and regeneration of serotonergic neurons in planarians.
Development, 140(17), pp.3577–3588.

Deochand, M.E., Birkholz, T.R. & Beane, W.S., 2016. Temporal regulation of planarian
eye regeneration. *Regeneration*, 3(4), pp.209–221.

D'Amelio, M., Sheng, M. & Cecconi, F., 2012. Caspase-3 in the central nervous system:
beyond apoptosis. *Trends in Neurosciences*, 35(11), pp.700–709.

Elmore, S., 2016. Apoptosis: A Review of Programmed Cell Death. *Toxicologic
Pathology*, 35(4), pp.495–516.

Fritz, A.E. et al., 2013. Mechanisms of tentacle morphogenesis in the sea anemone

Nematostella vectensis. *Development*, 140(10), pp.2212–2223.

Gahan, J.M. et al., 2017. Functional studies on the role of Notch signaling in Hydractinia
975   development. *Developmental biology*, 428(1), pp.224–231.

976   Galliot, B. et al., 2007. RNAi gene silencing affects cell and developmental plasticity in
977   hydra. *Comptes Rendus Biologies*, 330(6-7), pp.491–497.

978   Hand, C. & Uhlinger, K.R., 1992. The Culture, Sexual and Asexual Reproduction, and
979   Growth of the Sea Anemone *Nematostella vectensis*. *Biological Bulletin*, 182,
980   pp.169–176.

981   Havrilak, J.A. et al., 2017. Characterization of NvLWamide-like neurons reveals
982   stereotypy in Nematostella nerve net development. *Developmental biology*, pp.1–0.

983   Jahnel, S.M., Walzl, M. & Technau, U., 2014. Development and epithelial organisation
984   of muscle cells in the sea anemone Nematostella vectensis. 11(1), pp.1–15.

985   Kuida, K. et al., 1996. Decreased apoptosis in the brain and premature lethality in
986   CPP32-deficient mice. *Nature*, 384, pp.368–372.

987   Layden, M.J. et al., 2010. Expression and phylogenetic analysis of the zic gene family in
988   the evolution and development of metazoans. *EvoDevo*, 1(1), p.12. Available at:
989   http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=21054859
990   &retmode=ref&cmd=prlinks.

991   Layden, M.J. et al., 2016. MAPK signaling is necessary for neurogenesis in Nematostella
992   vectensis. *BMC Biology*, 14(1), p.61.

993   Leclère, L. et al., 2016. Development of the aboral domain in Nematostella requires β-
994   catenin and the opposing activities of Six3/6 and Frizzled5/8. *Development*, 143(10),
995   pp.1766–1777.

996   Marlow, H., Matus, D.Q. & Martindale, M.Q., 2013. Ectopic activation of the canonical
997   wnt signaling pathway affects ectodermal patterning along the primary axis during
998   larval development in the anthozoan Nematostella vectensis. *Developmental biology*,
999   380(2), pp.324–334.

1000   Miljkovic-Licina, M. et al., 2007. Head regeneration in wild-type hydra requires de novo
1001   neurogenesis. 134(6), pp.1191–1201.

1002   Moya, A. et al., 2016. Functional conservation of the apoptotic machinery from coral to
1003   man: the diverse and complex Bcl-2 and caspase repertoires of Acropora millepora.
1004   *BMC Genomics*, pp.1–21.

1005   Nakanishi, N. et al., 2012. Nervous systems of the sea anemone Nematostella vectensis
1006   are generated by ectoderm and endoderm and shaped by distinct mechanisms.
1007   *Development*, 139(2), pp.347–357.

Oviedo, N.J., Newmark, P.A. & Sánchez Alvarado, A., 2003. Allometric scaling and
proportion regulation in the freshwater planarian Schmidtea mediterranea.
Developmental Dynamics, 226(2), pp.326–333.

Passamaneck, Y.J. & Martindale, M.Q., 2012. Cell proliferation is necessary for the
regeneration of oral structures in the anthozoan cnidarian Nematostella vectensis.
BMC Developmental Biology, 12(1), p.34.

Renfer, E. et al., 2010. A muscle-specific transgenic reporter line of the sea anemone,
Nematostella vectensis. *Proceedings of the National Academy of Sciences of the*
United States of America, 107(1), pp.104–108.

Rentzsch, F., Layden, M. & Manuel, M., 2016. The cellular and molecular basis of
cnidarian neurogenesis. *Wiley Interdisciplinary Reviews: Developmental Biology*,
pp.1–19.

Richards, G.S. & Rentzsch, F., 2014. Transgenic analysis of a SoxB gene reveals neural
progenitor cells in the cnidarian Nematostella vectensis. *Development*, 141(24),
pp.4681–4689.

Roberts-Galbraith, R.H., Brubacher, J.L. & Newmark, P.A., 2016. A functional genomics
screen in planarians reveals regulators of whole-brain regeneration. *eLIFE*, pp.1–31.

Romero, R. & BAGUÑÀ, J., 1991. Quantitative cellular analysis of growth and
reproduction in freshwater planarians (Turbellaria; Tricladida). I. A cellular
description of the intact organism. *Invertebrate Reproduction & Development*, 19(2),
pp.157–165.

Röttinger, E., Dahlin, P. & Martindale, M.Q., 2012. A Framework for the Establishment
of a Cnidarian Gene Regulatory Network for "Endomesoderm" Specification: The
Inputs of ß-Catenin/TCF Signaling M. C. Mullins, ed. *PLoS Genetics*, 8(12),
p.e1003164.

Siebert, S. et al., 2018. Stem cell differentiation trajectories in Hydra resolved at single-
cell resolution. pp.1–29.

Sinigaglia, C. et al., 2013. The bilaterian head patterning gene six3/6 controls aboral
domain development in a cnidarian. *PLoS Biology*, 11(2), p.e1001488.

Steinmetz, P.R.H. et al., 2017. Gut-like ectodermal tissue in a sea anemone challenges
germ layer homology. *Nature Ecology & Evolution*, 1(10), pp.1–0.

Takeda, H., Nishimura, K. & Agata, K., 2009. Planarians Maintain a Constant Ratio of
Different Cell Types During Changes in Body Size by Using the Stem Cell System.
Zoological Science, 26(12), pp.805–813.

Tay, T.L. et al., 2017. A new fate mapping system reveals context-dependent random or
clonal expansion of microglia. *Nature Neuroscience*, 20(6), pp.793–803.

Tucker, R.P., Shibata, B. & Blankenship, T.N., 2011. Ultrastructure of the mesoglea of
the sea anemone Nematostella vectensis (Edwardsiidae). *Invertebrate Biology*,
130(1), pp.11–24.

van Ham, T.J. et al., 2014. Intravital correlated microscopy reveals differential
macrophage and microglial dynamics during resolution of neuroinflammation.
Disease Models & Mechanisms, 7(7), pp.857–869.

Warner, J.F. et al., 2019. Regeneration is a partial redeployment of the embryonic gene
network. *bioRxiv*, 121, pp.1731–28.

Williams, R.B., 1975. A redescription of the brackish-water sea anemone *Nematostella*
vectensis Stephenson, with an appraisal of congeneric species. *Journal of Natural*
History, 9(1), pp.51–64.

Zhang, J. et al., 2003. Identification of the haematopoietic stem cell niche and control of
the niche size. *Nature*, 425(6960), pp.836–841.

1057