Experience shapes activity dynamics and stimulus coding of VIP inhibitory and excitatory cells

Garrett M[*], Manavi S, Roll K, Ollerenshaw D, Groblewski P, Kiggins J, Jia X, Casal L, Mace K, Williford A, Leon A, Mihalas S, Olsen SR[*]

Allen Institute for Brain Science, Seattle, WA 98109, USA

*Corresponding authors: marinag@alleninstitute.org, shawno@alleninstitute.org

ABSTRACT

Cortical circuits are flexible and can change with experience and learning. However, the effects of experience on specific cell types including distinct inhibitory types are not well understood. Here we studied how excitatory and VIP inhibitory cells in layer 2/3 of mouse visual cortex were impacted by visual experience in the context of a behavioral task. Mice learned to perform an image change detection task with a set of eight natural scene images. Subsequently, during 2-photon imaging experiments, mice performed the task with these familiar images and three additional sets of novel images. Familiar images evoked less overall activity in both excitatory and VIP populations, and excitatory cells showed higher selectivity for familiar images. The temporal dynamics of VIP cells differed markedly between novel and familiar images: VIP cells were stimulus-driven for novel images but displayed ramping activity during the inter-stimulus interval for familiar images. Moreover, when a familiar stimulus was omitted, VIP cells showed extended ramping activity until the subsequent image flash. This prominent shift in response dynamics suggests that VIP cells may adopt different modes of processing during familiar versus novel conditions.

HIGHLIGHTS

- Experience with natural images in a change detection task reduces overall activity of cortical excitatory and VIP inhibitory cells
- Encoding of natural images is sharpened with experience in excitatory neurons
- VIP cells are stimulus-driven by novel images but show pre-stimulus ramping for familiar images
- VIP cells show strong ramping activity during the omission of an expected stimulus

INTRODUCTION

It is increasingly clear that cortical activity, even in primary sensory areas, is influenced by behavioral context, internal states, and other factors beyond external stimulus features including movement, arousal, motivation, and task engagement (Batista-Brito et al., 2018; Busse et al., 2017; Gilbert and Li, 2013; Khan and Hofer, 2018; Kuchibhotla and Bathellier, 2018; McGinley et al., 2015; Pakan et al., 2018). Moreover, cortical circuits are not fixed but change with experience and learning, leading to the modification of sensory representations (De Lange et al., 2018; LeMessurier and Feldman, 2018; Ranganath and Rainer, 2003), including changes in response gain, selectivity, correlations, and population dynamics (Jurjut et al., 2017; Khan et al., 2018; Makino and Komiyama, 2015; Poort et al., 2015; Weskelblatt and Niell, 2019; Woloszyn and Sheinberg, 2012). Sensory experience has been also associated with the emergence of predictive activity in visual cortex such as reward anticipation (Poort et al., 2015; Shuler and Bear, 2006), spatial expectation (Fiser et al., 2016), pattern completion (Gavornik and Bear, 2014), and prediction error signals (Fiser et al., 2016; Hamm and Yuste, 2016; Homann et al., 2017). Experience-dependent changes in sensory cortex can involve top-down feedback (Fiser et al., 2016; Makino and Komiyama, 2015; Petro et al., 2014; Zhang and Dan, 2014) and neuromodulatory inputs (Chubykin et al., 2013; Pinto et al., 2013), and may be associated with a shift in the balance of bottom-up sensory and top-down contextual signals conveying internal state and learned expectations.

Vasoactive intestinal peptide (VIP) expressing cells comprise a major class of inhibitory neurons and are well-positioned to mediate top-down influences on local circuits in sensory cortex. VIP cells receive long-range projections from frontal areas (Lee et al., 2013; Wall et al., 2016; Zhang and Dan, 2014) as well as cholinergic and noradrenergic inputs (Alitto and Dan, 2013; Fu et al., 2014). VIP cells are highly active during states of arousal (Reimer et al., 2014), are modulated by running (Fu et al., 2014; Reimer et al., 2014) and task engagement (Kuchibhotla et al., 2017), and are responsive to behavioral reinforcement (Pi et al., 2013). In the local cortical circuitry, VIP cells primarily inhibit another major class of inhibitory interneuron, somatostatin (SST) cells (Lee et al., 2013; Munoz et al., 2017; Pfeffer et al., 2013; Pi et al., 2013), which can result in disinhibition of excitatory neurons (Fu et al., 2017; Lee et al., 2013; Letzkus et al., 2011). SST cells target the apical dendrites of pyramidal neurons (Kepecs and Fishell, 2014) and removal of this inhibition may facilitate the association of top-down and bottom-up input by pyramidal cells (Chen et al., 2015; Larkum, 2012; Makino and Komiyama, 2015). However, little is known about how VIP cell activity is modified by experience. One recent study examined VIP cells before and after orientation discrimination learning, but did not document major activity changes in this population (Khan et al., 2018). VIP cells are known to exhibit strong surround suppression (Dipoppa et al., 2018), and examining the effects of experience in visual tasks with naturalistic stimuli may be critical to understanding the role of VIP cells in regulating the flow of information.

Here we investigated how long-term behavioral experience with natural scene images alters activity in cortical excitatory and VIP inhibitory cells in layers 2/3 of mouse visual cortex. We trained mice to perform a change detection task in which images are flashed in a periodic manner and mice are rewarded for detecting changes in image identity. Mice learned the task with one set of eight natural images, which were viewed thousands of times during training and were thus highly familiar. During 2-photon imaging, these familiar images as well as three novel image sets were tested. Familiar images were associated with lower overall population activity in both excitatory and VIP cells, and excitatory cells were more stimulus selective. Strikingly, VIP inhibitory cells had distinct activity dynamics during sessions with familiar versus novel images. VIP cells were stimulus-driven by novel images but displayed ramping activity between stimulus flashes when presented with familiar images. Cells with pre-stimulus ramping showed even greater activity when an expected stimulus was omitted from the periodic image sequence. These results demonstrate experience-dependent changes in responsiveness, selectivity, and activity dynamics across two cortical cell classes, and suggest that VIP cells may adopt different modes of processing during familiar versus novel conditions.

RESULTS

Visual change detection task with familiar and novel images

We trained GCaMP6f-expressing transgenic mice to perform a go/no-go visual change detection task with natural scene stimuli (Figure 1A). In this task, mice are presented with a continuous stream of flashing stimuli (250 ms image flash separated by 500 ms gray screen; Figure 1B). On 'go' trials, a change in image identity occurs at a time unknown to the animal. To earn water rewards, mice must report the image change by licking a reward spout within a 750 ms response window. False alarms are quantified during 'catch' trials in which the repeated image does not change, but licking behavior is measured in a similar time window (Figure 1B; see Methods). Behavioral training proceeded through a series of stages: mice first learned task rules with simple full-field gratings of two different orientations with no intervening gray period, followed by introduction of a 500ms gray screen period between grating flashes, and lastly, training of natural scene stimuli (Figure 1C). During the natural image stage, mice were trained with one set of eight images (image set A) for an extended number of sessions (range = 6-46 sessions with image set A, median = 17 sessions; Figure 1F, Supplemental Figure 1A). Mice viewed each of the eight scenes from the familiar image set an average of 10,350 times prior to the 2-photon imaging stage (range: 944-26,784 individual image presentations per image).

The behavioral task flow for one example session with natural images is shown in Figure 1G. The stimulus frequency is periodic, with an image flash occurring every 750ms (vertical colored lines denote different image identities, transitions between colors denote changes in image identity). Rewards are delivered following a correct licking response within 750ms of the image change. To test whether expectation signals are present in the visual cortex during this task, we randomly omitted ~5% of all non-change flashes during the 2-photon imaging sessions (flashes were never omitted during training). This appears as an extended gray period to the mouse and is denoted by a gap in the regular stimulus frequency, as shown in Figure 1G.

Mice rapidly generalized change detection behavior to novel image sets (Figure 1E,H,I), and detected image changes with similar hit and low false alarm rates across image sets (Figure 1H; statistics are described in figure legends, see Methods for description of statistical tests used). Reaction times were also similar for familiar and novel image sets (Figure 1I). During the task, mice are free to run on a circular disk and typically stop running when emitting a licking response. There was no difference in average running speed or pattern of running behavior during novel and familiar image sessions (Supplemental Figure 1D-E). Together, these results indicate that overall behavior was similar for novel and familiar image sets.

Imaging excitatory and inhibitory cortical populations during task performance

To explore differences between cell classes, we measured activity in transgenic mice expressing the calcium indicator GCaMP6f in either excitatory pyramidal neurons (Slc17a7-IRES2-Cre; CaMKII-tTA; Ai93-GCaMP6f) or VIP inhibitory neurons (VIP-IRES-Cre; Ai148-GCaMP6f) (Figure 2A,B). We imaged on average 181+/-77 (mean +/-SD) Slc17a7+ cells or 15+/-10 VIP+ cells per session (Supplemental Figure 2A). Overall, we collected data from a total of 12-13 sessions per transgenic line for each image set (Supplemental Figure 2A). We measured activity in the primary visual cortex, VISp, and one higher visual area, VISal. However, no major differences were found between populations in these two areas for the metrics we evaluated (data not shown), so we combined cells across these areas for all analyses. In each imaging session, one of the 4 image sets of natural scenes was shown (Figure 1C). In some cases, calcium imaging experiments failed to meet our quality control criteria (see Methods), and the session was retaken on the following day. As a result, some sessions with novel image sets B, C or D were the second or third exposure (67% were first exposure, 27% were the second exposure, 6% were the third or fourth exposure). This amount of exposure was still much less than for the familiar image set (median of 17 sessions with image set A).

Example imaging sessions for both excitatory and VIP cells with either familiar or novel image sets are shown in Figure 2C-F. Excitatory cells were typically responsive to only one or a few images in each set. When presented with their preferred stimulus, most excitatory cells responded with an increase in fluorescence shortly after stimulus onset, while other cells responded after stimulus offset (Figure 2C,D right panels). VIP cells were generally less selective for image identity than excitatory cells and were more similar with each other in their pattern of activity. Notably, VIP responses across different flashes of the same stimulus could be highly variable in magnitude (Figure 2E,F left panels). Moreover, VIP cells showed clear differences in their activity to novel versus familiar images. Stimulus-driven activity was apparent during sessions with novel image sets (Figure 2F) but was reduced or absent with familiar images (Figure 2E). In sessions with familiar image sets, many VIP neurons showed ramp-like responses that preceded stimulus presentation, and these ramping responses were even more pronounced when image flashes were omitted (Figure 2E, right column of right panel). In contrast, in novel image sessions VIP cells showed little activity when stimuli were omitted (Figure 2C,D, right column of right panels). Additional examples of average image responses from different sessions are shown in Supplemental Figure 2E-H.

Qualitatively, we observed that both Slc17a7 excitatory and VIP inhibitory neuronal populations showed differences in stimulus responsiveness, image tuning, and neural dynamics during behavioral sessions with novel versus familiar images. These results are further explored and quantified in the sections below.

A larger fraction of cells are stimulus responsive for novel images

To quantify image responsiveness for each cell, we compared the mean dF/F value in a 500ms window after each stimulus flash with a shuffled distribution of dF/F values taken from the omitted flash periods when no stimulus was shown. If a cell had a statistically significant response for >25% of stimulus flashes for its preferred image, it was considered image responsive (see Methods and Supplemental Figure 2). Then, the fraction of cells that met this criterion within each session was computed (Figure 2G). Most excitatory cells were image responsive for novel stimulus sets (range = 67-77% for image sets B, C & D), whereas a lower fraction of cells responded to familiar images (35%). VIP cells showed some stimulus responsiveness for novel images (range = 31-40%), but essentially no reliable stimulus specific activity for image set A (1%). To evaluate whether responsiveness differed for the change flash specifically, rather than across all flashes of an image, we computed change responsiveness in a similar manner, now restricting the analysis to the first flash after a change to each cell's preferred image (Figure 2H). A large fraction of excitatory cells were change responsive for novel images (range = 88-92% for image sets B, C & D), with a smaller fraction for the familiar image set A (61%). For VIP cells, a subset of cells were change responsive for novel images (range = 24-28%), with only 7% for familiar images. To quantify responsiveness during omitted flashes we compared activity during stimulus omission (dF/F values in 750ms window after the expected stimulus presentation time) to the preceding time window during which stimuli were presented (750ms prior to the expected stimulus time) (Figure 2I). If >25% of omission trials had a significantly larger response during the omission window compared to the previous image flash, the cell was deemed omission responsive. VIP populations showed a high fraction of cells with increased activity following stimulus omission for the familiar image set (72%), and a smaller fraction of omission responsive cells for the novel image sets (range = 17-30% for image sets B, C & D). Less than 1.5% of excitatory cells were omission responsive across all image sets.

Mean activity is lower but stimulus selectivity is higher for familiar image sets

The fraction of stimulus responsive cells was reduced for familiar versus novel stimulus sets. This could represent a sparsening of neural representations for the familiar images. More generally, several broad categories of sparseness have been identified (Willmore et al., 2019), including overall activity,

population sparseness, and lifetime sparseness. Previous studies have shown these metrics of sparseness reflect distinct coding properties and do not necessarily co-vary (Willmore and Tolhurst, 2001; Willmore et al., 2011). Thus, in the analysis below, we assessed whether each of these sparseness metrics was altered by image familiarity.

We computed the mean response to the eight images tested in each session for each cell, across all stimulus presentations for each image. Visualizing the tuning curves across all cells in the dataset as a heatmap (Figure 3A,B) confirmed that fewer cells (both excitatory and VIP) had image evoked activity for the familiar image set A. Quantification of mean response magnitude for the preferred image across cells also demonstrated that overall activity levels were reduced for image set A compared to B, C, and D within each cell class (Figure 3C). The strength of evoked activity across the VIP population was ~4-5 times higher compared to excitatory cells on average (Figure 3C).

To examine the population representation for familiar and novel images, we computed a population sparseness value for each image on a session by session basis (example in Figure 3E), and then took the average across all images within each session (Figure 3D). Population sparseness provides a measure of how selectively a population of simultaneously recorded neurons responds to any one stimulus, independently of the overall level of activity (Willmore et al., 2019). In the limit of an infinitely large population, sparseness of 0 indicates that all neurons respond equally to a given stimulus, while a sparseness of 1 indicates that only a single cell responds to the stimulus. VIP inhibitory populations showed lower population sparseness values than excitatory populations, as previously reported (de Vries et al., 2018). This indicates that image responses are more correlated across the population. However, no significant differences were observed across image sets within each cell class (Figure 3D).

Many excitatory and VIP inhibitory cells had a preferential response for some images over others. To quantify this, we computed an image selectivity metric for cells that met our image responsiveness criteria (Figure 2G, Supplemental Figure 2B). Lifetime sparseness (Vinje and Gallant, 2000) provides a measure of selectivity on a single neuron basis; at the extremes, this metric takes a value of 0 for cells responding equally to all images and a value of 1 for cells responding to only one image (Figure 3F). We found that excitatory populations had higher lifetime sparseness values for the familiar image set compared to the novel image sets (Figure 3G). Plotting the population tuning curve for each image set revealed sharper tuning in excitatory cells for familiar images due to a selective increase in the preferred image response (Figure 3H), consistent with previous literature (Woloszyn and Sheinberg, 2012). A comparison of lifetime sparseness across familiar and novel image sets for VIP inhibitory cells was not possible, as very few VIP cells met our criteria for image responsiveness (Figure 2I).

Thus, while overall population activity levels were reduced for familiar images, single cell selectivity was sharpened due to stronger responses to the preferred image. The higher overall level of activity for novel images, combined with lower selectivity indicates that more cells were recruited to respond to novel images, but the peak response was typically not as high as for images that were experienced during training. Interestingly, population sparseness, which is a measure of the shape as opposed to magnitude of the population activity distribution, was less impacted by experience, indicating that familiar and novel images can be represented by a similar pattern of activity but with a different overall gain.

To evaluate sparseness of the change response, we performed a similar analysis, limited to the first image presentation after a change (Supplemental Figure 3). Overall, activity for image changes was stronger compared to activity averaged across all image presentations (compare Figure 3C and Supplemental Figure 3C). The mean response across excitatory cells was 0.12 dF/F for the change flash, compared to 0.06 for the average of all flashes, and the mean change response for VIP inhibitory cells was 0.37 dF/F, compared to 0.26 for all flashes. Comparing across image sets, excitatory cells again showed higher sparseness for familiar images, but this difference was driven by a broadening of

selectivity for the novel image sets; there was stronger activity for non-preferred image changes relative to the optimal image (Supplemental Figure 3E,F). The stimulus tuning of VIP inhibitory cells for the change flash was markedly different than for all flashes, no longer showing preference for specific images and instead responding broadly to multiple images (Supplemental Figure 3B). Direct comparison of population tuning curves computed from the average of all flashes and population tuning curves for the change flash confirm that selectivity for image identity is broader following changes, for both excitatory and VIP cells (Supplemental Figure 2G,H). These results indicate that there is a difference in the pattern of activity for the change flash compared to subsequent stimulus repetitions (i.e. when averaging across all flashes).

Enhanced activity following stimulus changes

To explore response dynamics following an image change and subsequent image repetitions, we examined the trial averaged trace aligned to the change time for each cell's preferred image (Figure 4A,B). For both excitatory and VIP inhibitory cells, a large fraction of the population showed some level of activity following a change (seen as a vertical band of activity after time = 0 sec, Figure 4A,B). In excitatory cells, the strength of the response typically decreased on subsequent image flashes, consistent with stimulus specific adaptation (Figure 4C). In VIP cells, activity during the preferred image sometimes increased in magnitude with repeated flashes (Figure 4C, Supplemental Figure 4A); this increase in activity with stimulus repetition could reflect true facilitation of spiking, or a buildup of calcium activity associated with repetitive stimulation with the preferred stimulus. This stronger response to repeated image flashes across the VIP population was not observed when averaging the response to all images (Supplemental Figure 4B). This suggests that VIP cells have generally enhanced change responses across many images, consistent with the broader tuning of these cells for changes (Supplemental Figure 3G,H), but that any enhancement for the preferred image may be masked by the strong responses to subsequent presentations of the preferred image. To quantify these observations across cells, we computed a change modulation index to compare the response to image changes relative to the 10th repetition of the image for each cell (see Methods), then averaged across cells within each session. For the preferred image, excitatory cells typically had positive values of this index, whereas VIP had negative change modulation indexvalues (Figure 4D). When considering all images, both cell classes had positive values of the index (Supplemental Figure 4C), indicating stronger change responses.

These results show that the activity of excitatory and VIP cells is modulated by stimulus repetition following a change. The observed differences in activity following stimulus repetition could reflect stimulus specific adaptation, enhancement associated with a global change signal, or a combination of these factors.

Activity is more strongly stimulus modulated during sessions with novel images

The trial-to-trial reliability of neural activity can be modulated by bottom-up salience, behavioral context, and learning. Here we found that stimulus evoked responses were most reliable following a stimulus change, especially for novel images. We quantified reliability for image responsive cells by computing the average pairwise Pearson correlation coefficient across flashes, using the dF/F trace in a 500ms window after stimulus onset (Figure 4E), separately for the 1st, 5th, 10th, and 15th flashes after a change to each cell's preferred image, then took the average within each session (Figure 4F). This demonstrates that both stimulus novelty and stimulus repetition can modulate the reliability of neural responses in visual cortex. We also found that reliability is higher for novel images even when only considering the most selective cells in the population (Supplemental Figure 4D).

We noticed that the VIP population response for the familiar image set was only weakly stimulus driven, in comparison to the novel images. To quantify this, we computed a cell by cell metric of stimulus modulation (Matteucci et al., 2019), based on the power at the stimulus frequency (1/750ms) over an 8

second window after a change to the preferred image for each cell (Figure 4G). The stimulus modulation index is larger when activity is strongly coupled to the stimulus frequency across repetitive flashes and is lower when it is not (example cells with high and low modulation shown in Figure 4G). While many VIP cells were strongly stimulus modulated for novel images, modulation at the stimulus frequency was very weak for familiar images (Figure 4H, bottom panel), consistent an overall reduction in stimulus driven activity with experience. Excitatory cells also showed reduced stimulus modulation for familiar images, although the effect was not as pronounced as for VIP cells (Figure 4H, top panel).

Together, these results show that neural activity during task performance with novel, unfamiliar images is stronger across the population, more reliable across trials, and more strongly stimulus driven, compared to activity for images that were extensively experienced during training.

Inter-stimulus activity dynamics of VIP cells are altered by training history

During sessions with familiar images we observed that VIP cells displayed activity that began ramping up prior to image onset (see Figure 4C right panel). To examine these dynamics and their relationship to experience, we examined the average population response for excitatory and VIP cells averaged over all image flashes (Figure 5A). Excitatory neurons showed a sharp stimulus-locked increase in activity following image onset for all image sets. Although the response magnitude was lower with familiar images, as expected, the dynamics of the excitatory population were consistent across images sets. In contrast, the dynamics of the VIP population were almost anti-correlated for familiar and novel images sets. With novel images, VIP cells showed activity that increased in response to stimulus onset, but for familiar images, the population response increased during the inter-stimulus interval and peaked at stimulus onset. Consistent with this effect, the distribution of peak response times across VIP cells was shifted earlier in time for familiar versus novel images sets (Supplemental Figure 5A).

To characterize the diversity of single cell dynamics, we made use of a ramping index (Makino and Komiyama, 2015) to quantify how activity increases or decreases within the pre-stimulus and stimulus epochs (250ms windows, indicated by light and dark gray shading, respectively, in Figures 5A,B,E). This index compares activity between the early and late portions of a window, giving a positive value for activity increases and a negative value for activity decreases (Figure 5A, right panel inset). While positive pre-stimulus ramping was much less prevalent in excitatory cells than in VIP cells, both cell classes showed increased pre-stimulus activity and decreased stimulus evoked activity for the familiar image set (Supplemental Figure 5B,C).

Individual cell dynamics could be characterized by the relative value of the pre-stimulus and stimulus ramping indices (Figure 5B). The values of the ramp index for these two windows were typically anti-correlated with each other (Figure 5C; VIP: r = -0.72, -0.78, -0.61, -0.83 for image sets A-D; excitatory: r = -0.29, -0.35, -0.32, -0.26 for image sets A-D; p<0.05 for all conditions). Cells were naturally divided into four groups based on their ramping behavior before and after stimulus onset (Figure 5C; the four groups correspond to the four quadrants of this scatter plot). For example, cells in quadrant 1 (Q1) had increasing dF/F values in the stimulus window (corresponding to a positive ramp index) and decreasing values for the pre-stimulus window (corresponding to a negative ramp index), consistent with a stimulus evoked response profile. Cells in quadrant 4 (Q4) were the opposite, having increasing activity prior to stimulus onset (positive pre-stimulus ramp index) and decreasing activity thereafter (negative stimulus ramp index). The average population response profile across cells belonging to each of these quadrants is shown in Figure 5E.

We quantified the fraction of cells in each quadrant and found that the majority of excitatory cells for both novel and familiar images were located in Q1, consistent with strong stimulus evoked activity (72-84% of cells across image sets A-D were in Q1; Figure 5D). The majority of VIP inhibitory cells were also in Q1 for novel image sets (69-78% for image sets B-D). However there was a large increase in the proportion of VIP cells in Q4 showing pre-stimulus activity for the familiar image set (55% for image

set A compared to 7-9% for image sets B-D; Figure 5D), representing an experience-dependent increase in inter-stimulus activity across the VIP population. Cells belonging to these distinct response types also differed in their stimulus-evoked response properties. Q1 cells, in line with being sensory-driven, were generally more strongly stimulus modulated across repeated image flashes, more reliable across trials, and more image selective than Q4 inter-stimulus ramping type cells (Figures 5G-I). In contrast, Q4 cells had earlier response times (Figure 5F) and were unselective for image identity (Figure 5I).

VIP cells show strong ramping activity during omission of an expected stimulus

Would cells with pre-stimulus ramping continue to ramp if an image was omitted? To assess this, we analyzed activity during periods in which the stimulus flash was randomly omitted (the change flash and flash immediately prior to the change were never omitted; see Methods). Such trials made up 5% of all stimulus flashes.

Strikingly, during stimulus omission in sessions with familiar images, VIP population activity continued to ramp up until the subsequent stimulus flash, more than doubling in response magnitude within the omission window (Figure 6A; prior image peak response: 0.11 dF/F, omission peak response: 0.23 dF/F for VIP with image set A). This ramping was much stronger during familiar compared to novel image sessions, although single VIP cells in sessions with either familiar or novel image sets could show ramping, and some excitatory cells also showed ramping behavior (Figure 6B). To evaluate the prevalence of ramping across all cells in the dataset, we examined heatmaps showing activity aligned to the time of stimulus omission (Figure 6C,D). For familiar image sessions, nearly all cells in the VIP population showed an increase in activity during the omission period that lasted for ~750ms, up to the time of the subsequent stimulus flash (Figure 6D). For novel image sessions, activity in VIP cells was concentrated outside the omission period, with visible stimulus-locked activity in the surrounding timepoints. Still, a subset of VIP cells for image sets B-D showed some degree of activity during stimulus omission, but the onset was later than for novel image sessions. Activity during the omission period was very rare in excitatory cells (Figure 6C).

To quantify ramping activity associated with stimulus omission, we computed the ramping index over the stimulus omission window. Here we focused on cells that were active during the omission, with an absolute dF/F of > 0.05 dF/F (3-6% of excitatory cells and 68-76% of VIP inhibitory cells met this criterion; Figure 6E). Most VIP cells that met the minimum activity level had positive values of the omission ramp index for sessions with familiar images (Figure 6F, right panel). Overall, 80% of VIP cells showed omission ramping for familiar sessions, compared to 40.5% of cells for novel sessions on average. While ramping was rare in excitatory cells, the fraction of cells with ramping during omission was slightly increased for sessions with familiar images (1.1% for familiar, 0.57% for novel on average).

When looking at the response profiles of individual cells (Figure 6B), we noticed a relationship between omission ramping and stimulus related dynamics. Typically, cells that had strong stimulus-evoked responses showed little to no activity in the omitted window, while cells with pre-stimulus activity had the strongest omitted stimulus ramping (Figure 6B). Indeed, there was a correlation between pre-stimulus ramping index and the omitted stimulus ramping index, particularly for VIP cells (Figure 6G). However, the population of VIP cells that show omission related ramping during sessions with novel image sets had a more sensory driven response profile upon image presentation (having both positive omitted ramp index and positive stimulus ramp index; Figure 6H; example cell in upper right panel of Figure 6B). This indicates that during visual stimulation with novel images, VIP cells can show a combination of sensory driven activity and ramping during omission. These results demonstrate that the dynamics of VIP cells in visual cortex are strongly impacted by experience.

DISCUSSION

By imaging cortical activity in L2/3 excitatory and VIP inhibitory neurons during a visual change detection task with familiar and novel images, we identified several changes associated with training history. Extended behavioral experience with natural scene images resulted in reduced overall activity levels and increased selectivity of single neuron representations. In contrast, responses to novel images were strong and reliable, particularly for stimulus changes. Remarkably, VIP cells displayed distinct activity dynamics when tested with familiar versus novel images. Novel images drove stimulus-locked activity in VIP cells, but this cell type was only weakly stimulus responsive to familiar images. Together these results indicate that both stimulus coding and temporal dynamics of cortical circuits can be impacted by experience in a cell type-specific manner.

Distinct activity and tuning in excitatory versus VIP cells

Across all conditions, VIP cells had higher overall average activity levels compared to excitatory neurons. In contrast, excitatory neurons were more strongly stimulus modulated and had more reliable stimulus-evoked responses. Excitatory cells were sharply tuned for natural images, whereas VIP cells responded more broadly, and were more correlated with each other. These results are consistent with previous studies showing overall activity and selectivity differences of VIP compared to excitatory cells (Kerlin et al., 2010; de Vries et al., 2018). We also found that stimulus repetition influenced activity. When averaging over all images tested in a session, both cell classes showed stronger responses to changes relative to subsequent stimulus repetitions. In addition, the change response was more broadly tuned than the 10th stimulus repetition for both excitatory and VIP cells. These differences could be a result of stimulus specific adaptation (Grill-Spector et al., 2006; Nelken and Ulanovsky, 2007), or novelty enhancement (Hamm and Yuste, 2016; Homann et al., 2017; Vinken et al., 2017), potentially reflecting a combination of bottom-up, top-down, or neuromodulatory influences (Ranganath and Rainer, 2003).

Experience reduces overall activity and sharpens tuning

While stimulus repetition was associated with reduced activity on short timescales (~seconds), we also observed reduced activity in both VIP and excitatory cells due to long-term experience with highly familiar images (~days-weeks). The fraction of image responsive cells (Figure 2) and response magnitude across the population (Figure 3, Figure 4) were lower for familiar compared to novel images. Previous studies have shown reductions in activity with experience (Anderson et al., 2008; Mruczek and Sheinberg, 2007; Woloszyn and Sheinberg, 2012), and enhancement for novelty (Hamm and Yuste, 2016; Homann et al., 2017; Ranganath and Rainer, 2003). A recent study showed that experience-induced reductions in the fraction of active cells can be specific to distinct functional subpopulations: 'transient' cells were reduced with experience, but 'sustained' and 'suppressed by contrast' cells showed no change in the fraction of active cells (Weskelblatt and Niell, 2019). Reduced activity for highly familiar stimuli may serve to more efficiently code for stimuli which are predictable, utilizing a smaller population of cells to represent learned information (LeMessurier and Feldman, 2018). On the other hand, enhanced activity for novel stimuli may be involved in the detection of salient and behaviorally meaningful events by augmenting output to downstream targets and facilitating associative plasticity (Ranganath and Rainer, 2003). Consistent with this idea, we observe a nearly two fold increase in the number of cells that respond to novel images compared to familiar ones, for both excitatory and VIP inhibitory populations (Figure 2G,H).

We found that stimulus selectivity of single cells, as measured by lifetime sparseness, was higher in the L2/3 excitatory cells for familiar images. Stimulus experience and task learning have been shown to increase orientation selectivity in the visual cortex of mice (Fiser et al., 2016; Frenkel et al., 2006; Jurjut et al., 2017; Khan et al., 2018; Poort et al., 2015) and selectivity for natural images including complex objects in primates (Freedman et al., 2006; Ghose et al., 2002; Meyer et al., 2014; Schoups et al.,

2001; Yang and Maunsell, 2004). While few studies have explored cell type-specific differences in stimulus tuning following experience, one study in macaque IT showed that both regular spiking (RS) and fast spiking (FS; putative inhibitory) cells showed increased stimulus selectivity for familiar images (Woloszyn and Sheinberg, 2012)**.** A study in mice reported increased selectivity with experience in excitatory and multiple inhibitory cell types, including PV and SST, but similar to our results, did not find a change in selectivity for VIP cells (Khan et al., 2018).

Many studies have shown that experience-dependent changes in responsiveness and selectivity can underlie improvements in perception and behavior. Here, we did not find significant changes in behavioral performance between familiar and novel image sets. Mice rapidly generalized performance on task to novel images and had comparable hit and false alarm rates, reaction times, and patterns of running behavior across image sets (Figure 1, Supplemental Figure 1). Accordingly, it is unlikely that the differences in response properties for familiar versus novel images are the consequence of differences in animal behavior or task performance. That we observed consistent task performance despite differences in stimulus coding suggests the increased activity for novel images could help maintain task performance in a new stimulus context, or alternatively, that a more efficient representation is sufficient for performance with familiar images.

Experience alters temporal dynamics of VIP cells

One of the main findings of our study is that extended visual experience with temporally predictable natural scene stimuli is associated with altered activity dynamics in VIP inhibitory cells in layer 2/3 of visual cortex (Figure 5). The VIP population displayed largely stimulus-driven activity for novel images but showed prominent inter-stimulus activity for familiar images. This represents a major experience-induced change in response dynamics, specific to VIP cells, that has not been demonstrated previously. Further, cells with inter-stimulus ramping typically continue to increase their activity following omission of a stimulus, up until the time of the next stimulus onset (Figure 6). The magnitude of these omission related signals could be several times larger than the strength of stimulus-driven activity, suggesting that they are meaningful signals that could strongly influence network activity. By comparing a ramping index for both pre-stimulus and stimulus periods, we found an inverse linear relationship between inter-stimulus ramping and stimulus-triggered increases in VIP cells, indicating there is trade-off between sensory-driven and pre-stimulus activity in these cells (Figure 5).

What does pre-stimulus and stimulus-omission ramping activity in VIP cells represent? One possibility is that this ramping activity reflects the temporal structure of the behavioral task such that these ramping signals encode predictions about stimulus timing or reward expectation, or serve as a general attentional signal. Previous studies have shown stimulus, reward, and spatial expectation signals in the visual cortex of rodents. An early example in the rat visual cortex showed that after pairing visual stimulation with temporally predictable reward delivery, a subset of cells began to signal the timing of reward delivery (Shuler and Bear, 2006). Cholinergic signaling has been implicated in mediating these changes (Chubykin et al., 2013), as well as other attention and learning dependent effects (Hasselmo, 1995; Lee and Dan, 2012). Given the strong neuromodulatory drive to VIP cells, including cholinergic input, neuromodulation is one candidate mechanism to underlie the shift in response dynamics we observe across the VIP population.

Visual cortex has also been shown to learn experience-dependent stimulus predictions for repeated sequences of visual stimuli (Gavornik and Bear, 2014; Xu et al., 2012). However, in these studies, the predictive signal peaked at the expected time of the predicted event on omission trials, whereas our results show a continued ramping past the expected time of stimulus onset on omission trials. This suggests that the pre-stimulus ramping and associated omission ramping signals we observe may represent something other than a pure prediction of stimulus or reward timing.

In virtual navigation paradigms in which mice locomote along a linear track, V1 neurons have been found to predict upcoming stimuli at specific locations, with separate populations of cells that respond when expected stimuli are omitted (Fiser et al., 2016). In contrast, we observe pre-stimulus and omission related activity in the same cells, and these responses are generally unselective for image identity. Another study using virtual navigation with a visual discrimination task found pre-stimulus ramping activity specifically in the subpopulation of excitatory cells that encoded the rewarded stimulus, suggestive of reward anticipation (Poort et al., 2015). The emergence of an experience dependent ramping profile in anticipation of an imminent foot shock has also been documented (Makino and Komiyama, 2015). However, without an omission condition, it is unclear how these cells would behave if the expectation was violated. Further modifications of our change detection task to include omitted rewards, or to vary the inter-stimulus interval, could help to distinguish between coding of stimulus timing versus reward anticipation. Another open question is whether pre-stimulus ramping behavior emerges following passive exposure to the same set of familiar images in the absence of reward, or whether active task performance and reinforcement are necessary for the emergence of this phenomenon.

It is important to note that most prior studies documenting predictive or ramping activity were measurements of excitatory neurons, and thus may not be directly comparable to our results in VIP cells. In fact, it is surprising that we do not observe many excitatory cells with pre-stimulus or omission related activity given these prior results. One study of inhibitory cell activity in visual cortex did observe pre-stimulus ramping behavior in VIP cells in an orientation discrimination task, but did not note a major change in VIP activity dynamics before and after learning (Khan et al., 2018). It is possible that our experimental design, detecting changes in familiar and novel images, is sufficiently different as to not be directly comparable. In the prefrontal cortex of the mouse, VIP cell activity during the delay period of an auditory go/no-go discrimination task was found to be important for behavior, and activation of VIP cells during the delay improved task performance by improving coding in the excitatory population (Kamigaki and Dan, 2017). The inter-stimulus activity of VIP cells during sessions with familiar images in the change detection task may be similarly important for enhancing the responses of subsets of excitatory cells to an expected change stimulus, via disinhibition through SST cells. In contrast, during sessions with novel images, VIP activity is driven by stimulus presentation and could serve to increase the gain of stimulus evoked excitatory responses to facilitate learning of new images via associative plasticity (LeMessurier and Feldman, 2018; Ranganath and Rainer, 2003). Future studies examining the evolution of VIP activity across multiple behavior sessions as novel images become familiar, as well as concurrent recordings of VIP and excitatory cells and other inhibitory classes including SST cells, will be critical to determine the time course and mechanistic nature of these interactions.

Predictive coding and experience-dependent changes in activity

Predictive processing has emerged as a powerful paradigm for understanding brain function and may help reconcile the traditional view of sensory processing with increasing evidence for experience and context dependent modulation in early sensory areas. This family of theories posits that the brain constructs an internal model of the environment based on experience, and that incoming sensory information is compared with learned expectations to continually update the model (De Lange et al., 2018; Keller and Mrsic-Flogel, 2018; Lochmann and Deneve, 2011; Rao and Ballard, 1999). This dynamic updating with experience may be associated with a shift in the balance of bottom-up sensory and top-down predictive pathways as internal representations become more effective at predicting the external causes of sensations. As stimuli become familiar with learning, predictive signals are thought to 'explain away' incoming information by suppressing bottom-up input, resulting in a sparse code. In contrast, novel or surprising stimuli are expected to robustly drive neural activity, signaling deviations from learned predictions. Our results are consistent with this model, demonstrating reduced activity with long-term experience, in addition to within-trial stimulus specific adaptation. Further, the observation of a switch between stimulus evoked activity and inter-stimulus ramping in VIP cells provides a novel demonstration of an experience dependent change in activity dynamics in a specific subtype of

inhibitory interneuron that may regulate the balance between top-down predictive and bottom-up sensory information. Future studies examining the activity and impact of the diverse inputs to VIP cells, including neuromodulatory inputs from subcortical structures and feedback projections from other cortical regions, are necessary to establish the function of and mechanism behind the shift in VIP dynamics with experience.

ACKNOWLEDGEMENTS

We thank Jerome Lecoq and Kevin Takasaki for technical help with the 2-photon microscope, Derric Williams for help with behavior and stimulus control software, Saskia de Vries, Brain Hu, and Christof Koch for comments on the manuscript.

REFERENCES

Alitto HJ, Dan Y. 2013. Cell-type-specific modulation of neocortical activity by basal forebrain input. *Front Syst Neurosci* **6**:1–12. doi:10.3389/fnsys.2012.00079

Anderson B, Mruczek REB, Kawasaki K, Sheinberg D. 2008. Effects of familiarity on neural activity in monkey inferior temporal lobe. *Cereb Cortex* **18**:2540–2552. doi:10.1093/cercor/bhn015

Batista-Brito R, Zagha E, Ratliff JM, Vinck M. 2018. Modulation of cortical circuits by top-down processing and arousal state in health and disease. *Curr Opin Neurobiol* **52**:172–181. doi:10.1016/j.conb.2018.06.008

Busse L, Cardin JA, Chiappe ME, Halassa MM, McGinley MJ, Yamashita T, Saleem AB. 2017. Sensation during Active Behaviors. *J Neurosci* **37**:10826–10834. doi:10.1523/JNEUROSCI.1828-17.2017

Chen SX, Kim AN, Peters AJ, Komiyama T. 2015. Subtype-specific plasticity of inhibitory circuits in motor cortex during motor learning. *Nat Neurosci* **18**:1109–1115. doi:10.1038/nn.4049

Chubykin AA, Roach EB, Bear MF, Shuler MGH. 2013. A Cholinergic Mechanism for Reward Timing within Primary Visual Cortex. *Neuron* **77**:723–735. doi:10.1016/j.neuron.2012.12.039

De Lange FP, Heilbron M, Kok P. 2018. How Do Expectations Shape Perception? Perceptual Consequences of Expectation **xx**:1–16. doi:10.1016/j.tics.2018.06.002

de Vries SEJ de, Lecoq J, Buice MA, Groblewski PA, Ocker GK, Oliver M, Feng D, Cain N, Ledochowitsch P, Millman D, Roll K, Garrett M, Keenan T, Kuan L, Mihalas S, Olsen S, Thompson C, Wakeman W, Waters J, Williams D, Barber C, Berbesque N, Blanchard B, Bowles N, Caldejon S, Casal L, Cho A, Cross S, Dang C, Dolbeare T, Edwards M, Galbraith J, Gaudreault N, Griffin F, Hargrave P, Howard R, Huang L, Jewell S, Keller N, Knoblich U, Larkin J, Larsen R, Lau C, Lee E, Lee F, Leon A, Li L, Long F, Luviano J, Mace K, Nguyen T, Perkins J, Robertson M, Seid S, Shea-Brown E, Shi J, Sjoquist N, Slaughterbeck C, Sullivan D, Valenza R, White C, Williford A, Witten D, Zhuang J, Zeng H, Farrell C, Ng L, Bernard A, Phillips JW, Reid RC, Koch C. 2018. A large-scale, standardized physiological survey reveals higher order coding throughout the mouse visual cortex. *bioRxiv* 359513. doi:10.1101/359513

Dipoppa M, Ranson A, Krumin M, Pachitariu M, Carandini M, Harris KD. 2018. Vision and Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. *Neuron* 1–14. doi:10.1016/j.neuron.2018.03.037

Fiser A, Mahringer D, Oyibo HK, Petersen A V., Leinweber M, Keller GB. 2016. Experience-dependent spatial expectations in mouse visual cortex. *Nat Neurosci* **19**:1658–1664. doi:10.1038/nn.4385

Freedman DJ, Riesenhuber M, Poggio T, Miller EK. 2006. Experience-dependent sharpening of visual shape selectivity in inferior temporal cortex. *Cereb Cortex* **16**:1631–1644. doi:10.1093/cercor/bhj100

Frenkel MY, Sawtell NB, Diogo ACM, Yoon B, Neve RL, Bear MF. 2006. Instructive Effect of Visual Experience in Mouse Visual Cortex. *Neuron* **51**:339–349. doi:10.1016/j.neuron.2006.06.026

Fu Y, Kaneko M, Tang Y, Alvarez-Buylla A, Stryker MP. 2017. A cortical disinhibitory circuit for enhancing adult plasticity. *Elife* 1–12. doi:10.7554/eLife.05558

Fu Y, Tucciarone JM, Espinosa JS, Sheng N, Darcy DP, Nicoll RA, Huang ZJ, Stryker MP. 2014. A cortical circuit for gain control by behavioral state. *Cell* **156**:1139–1152. doi:10.1016/j.cell.2014.01.050

Garrett ME, Nauhaus I, Marshel JH, Callaway EM. 2014. Topography and Areal Organization of Mouse Visual Cortex. *J Neurosci* **34**:12587–12600. doi:10.1523/JNEUROSCI.1124-14.2014

Gavornik JP, Bear MF. 2014. Learned spatiotemporal sequence recognition and prediction in primary

visual cortex. *Nat Neurosci* **17**:732–737. doi:10.1038/nn.3683

Ghose GM, Yang T, Maunsell JHR. 2002. Physiological correlates of perceptual learning in monkey V1 and V2. *J Neurophysiol* **87**:1867–1888. doi:10.1152/jn.00690.2001

Gilbert CD, Li W. 2013. Top-down influences on visual processing. *Nat Rev Neurosci* **14**:350–363. doi:10.1038/nrn3476

Goldey GJ, Roumis DK, Glickfeld LL, Kerlin AM, Reid RC, Bonin V, Andermann ML. 2014. Versatile cranial window strategies for long-term two-photon imaging in awake mice. *Nat Protoc* **9**:2515–2538. doi:10.1038/nprot.2014.165

Grill-Spector K, Henson R, Martin A. 2006. Repetition and the brain: Neural models of stimulus-specific effects. *Trends Cogn Sci* **10**:14–23. doi:10.1016/j.tics.2005.11.006

Guo Z V., Hires SA, Li N, O'Connor DH, Komiyama T, Ophir E, Huber D, Bonardi C, Morandell K, Gutnisky D, Peron S, Xu N, Cox J, Svoboda K. 2014. Procedures for Behavioral Experiments in Head-Fixed Mice. *PLoS One* **9**:e88678. doi:10.1371/journal.pone.0088678

Hamm JP, Yuste R. 2016. Somatostatin Interneurons Control a Key Component of Mismatch Negativity in Mouse Visual Cortex. *Cell Rep* **16**:597–604. doi:10.1016/j.celrep.2016.06.037

Hasselmo ME. 1995. Neuromodulation and cortical function: modeling the physiological basis of behavior. *Behav Brain Res* **67**:1–27. doi:10.1016/0166-4328(94)00113-T

Homann J, Koay SA, Glidden AM, Tank DW, Berry Ii MJ. 2017. Predictive Coding of Novel versus Familiar Stimuli in the Primary Visual Cortex. *bioRxiv*. doi:10.1101/197608

Jurjut O, Georgieva P, Busse L, Katzner S. 2017. Learning Enhances Sensory Processing in Mouse V1 before Improving Behavior. *J Neurosci* **37**:6460–6474. doi:10.1523/JNEUROSCI.3485-16.2017

Kamigaki T, Dan Y. 2017. Delay activity of specific prefrontal interneuron subtypes modulates memory-guided behavior. *Nat Neurosci* **20**:854–863. doi:10.1038/nn.4554

Keller GB, Mrsic-Flogel TD. 2018. Predictive Processing: A Canonical Cortical Computation. *Neuron* **100**:424–435. doi:10.1016/j.neuron.2018.10.003

Kepecs A, Fishell G. 2014. Interneuron cell types are fit to function. *Nature* **505**:318–326. doi:10.1038/nature12983

Kerlin AM, Andermann ML, Berezovskii VK, Reid RC. 2010. Broadly tuned response properties of diverse inhibitory neuron subtypes in mouse visual cortex. *Neuron* **67**:858–71.

Khan AG, Hofer SB. 2018. Contextual signals in visual cortex. *Curr Opin Neurobiol* **52**:131–138. doi:10.1016/j.conb.2018.05.003

Khan AG, Poort J, Chadwick A, Blot A, Sahani M, Mrsic-flogel TD, Hofer SB. 2018. Distinct learning-induced changes in stimulus selectivity and interactions of GABAergic interneuron classes in visual cortex - Supplemental. *Nat Neurosci*. doi:10.1038/s41593-018-0143-z

Kuchibhotla K, Bathellier B. 2018. Neural encoding of sensory and behavioral complexity in the auditory cortex. *Curr Opin Neurobiol* **52**:65–71. doi:10.1016/j.conb.2018.04.002

Kuchibhotla K V., Gill J V., Lindsay GW, Papadoyannis ES, Field RE, Sten TAH, Miller KD, Froemke RC. 2017. Parallel processing by cortical inhibition enables context-dependent behavior. *Nat Neurosci* **20**:62–71. doi:10.1038/nn.4436

Larkum M. 2012. A cellular mechanism for cortical associations: An organizing principle for the cerebral cortex. *Trends Neurosci* **36**:141–151. doi:10.1016/j.tins.2012.11.006

Lee S-H, Dan Y. 2012. Neuromodulation of brain states. *Neuron* **76**:209–22. doi:10.1016/j.neuron.2012.09.012

Lee S, Kruglikov I, Huang ZJ, Fishell G, Rudy B. 2013. A disinhibitory circuit mediates motor integration in the somatosensory cortex. *Nat Neurosci* **16**:1662–1670. doi:10.1038/nn.3544

LeMessurier AM, Feldman DE. 2018. Plasticity of population coding in primary sensory cortex. *Curr Opin Neurobiol* **53**:50–56. doi:10.1016/j.conb.2018.04.029

Letzkus JJ, Wolff SBE, Meyer EMM, Tovote P, Courtin J, Herry C, Lüthi A. 2011. A disinhibitory microcircuit for associative fear learning in the auditory cortex. *Nature* **480**:331–5. doi:10.1038/nature10674

Lochmann T, Deneve S. 2011. Neural processing as causal inference. *Curr Opin Neurobiol* **21**:774–781. doi:10.1016/j.conb.2011.05.018

Makino H, Komiyama T. 2015. Learning enhances the relative impact of top-down processing in the

visual cortex. *Nat Neurosci* **18**:1–10. doi:10.1038/nn.4061

Marshel JH, Garrett ME, Nauhaus I, Callaway EM. 2011. Functional Specialization of Seven Mouse Visual Cortical Areas. *Neuron* **72**:1040–1054. doi:10.1016/j.neuron.2011.12.004

Matteucci G, Bellacosa Marotti R, Riggi M, Rosselli FB, Zoccolan D. 2019. Nonlinear processing of shape information in rat lateral extrastriate cortex. *J Neurosci* **39**:1938–18. doi:10.1523/JNEUROSCI.1938-18.2018

McGinley MJ, Vinck M, Reimer J, Batista-Brito R, Zagha E, Cadwell CR, Tolias AS, Cardin JA, McCormick DA. 2015. Waking state: rapid variations modulate neural and behavioral responses. *Neuron* **87**:1143–1161. doi:10.1016/j.neuron.2015.09.012

Meyer T, Walker C, Cho RY, Olson CR. 2014. Image familiarization sharpens response dynamics of neurons in inferotemporal cortex. *Nat Neurosci* **17**:1388–1394. doi:10.1038/nn.3794

Mruczek REB, Sheinberg DL. 2007. Context Familiarity Enhances Target Processing by Inferior Temporal Cortex Neurons. *J Neurosci* **27**:8533–8545. doi:10.1523/jneurosci.2106-07.2007

Munoz W, Tremblay R, Levenstein D, Rudy B. 2017. Layer-specific modulation of neocortical dendritic inhibition during active wakefulness **959**:954–959.

Nelken I, Ulanovsky N. 2007. Mismatch negativity and stimulus-specific adaptation in animal models. *J Psychophysiol* **21**:214–223. doi:10.1027/0269-8803.21.34.214

Olmos A, Kingdom FAA. 2004. A biologically inspired algorithm for the recovery of shading and reflectance images. *Perception* **33**:1463–1473. doi:10.1068/p5321

Oommen BS, Stahl JS. 2008. Eye orientation during static tilts and its relationship to spontaneous head pitch in the laboratory mouse. *Brain Res* **1193**:57–66. doi:10.1016/j.brainres.2007.11.053

Pakan JMJ, Francioni V, Rochefort NNL. 2018. Action and learning shape the activity of neuronal circuits in the visual cortex. *Curr Opin Neurobiol* **52**:88–97. doi:10.1016/j.conb.2018.04.020

Petro LS, Vizioli L, Muckli L. 2014. Contributions of cortical feedback to sensory processing in primary visual cortex. *Front Psychol* **5**:1–8. doi:10.3389/fpsyg.2014.01223

Pfeffer CK, Xue M, He M, Huang ZJ, Scanziani M. 2013. Inhibition of inhibition in visual cortex: the logic of connections between molecularly distinct interneurons. *Nat Neurosci* 1–12. doi:10.1038/nn.3446

Pi H-J, Hangya B, Kvitsiani D, Sanders JI, Huang ZJ, Kepecs A. 2013. Cortical interneurons that specialize in disinhibitory control. *Nature* **503**:521–4. doi:10.1038/nature12676

Pinto L, Goard MJ, Estandian D, Xu M, Kwan AC, Lee S-H, Harrison TC, Feng G, Dan Y. 2013. Fast modulation of visual perception by basal forebrain cholinergic neurons. *Nat Neurosci* 1–10. doi:10.1038/nn.3552

Poort J, Khan AG, Pachitariu M, Nemri A, Orsolic I, Krupic J, Bauza M, Sahani M, Keller GB, Mrsic-Flogel TD, Hofer SB. 2015. Learning Enhances Sensory and Multiple Non-sensory Representations in Primary Visual Cortex. *Neuron* **86**:1478–1490. doi:10.1016/j.neuron.2015.05.037

Ranganath C, Rainer G. 2003. Cognitive neuroscience: Neural mechanisms for detecting and remembering novel events. *Nat Rev Neurosci* **4**:193–202. doi:10.1038/nrn1052

Rao RPN, Ballard DH. 1999. Predictive coding in the visual cortex: A functional interpretation of some extra-classical receptive-field effects. *Nat Neurosci* **2**:79–87. doi:10.1038/4580

Reimer J, Froudarakis E, Cadwell CR, Yatsenko D, Denfield GH, Tolias AS. 2014. Pupil Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness. *Neuron* **84**:355–362. doi:10.1016/j.neuron.2014.09.033

Schoups A, Vogels R, Qian N, Orban G. 2001. Practising orientation identification improves orientation coding in V1 neurons. *Nature* **412**:549–553.

Shuler MG, Bear MF. 2006. Reward timing in the primary visual cortex. *Science (80-)* **311**:225–240. doi:10.1016/b978-0-408-01434-2.50020-6

Strasburger H, Rentschler I, Jüttner M. 2011. Peripheral vision and pattern recognition: A review. *J Vis* **11**:1–82. doi:10.1167/11.5.13.Contents

van Hateren JH, van der Schaaf a. 1998. Independent component filters of natural images compared with simple cells in primary visual cortex. *Proc Biol Sci* **265**:359–366. doi:10.1098/rspb.1998.0303

Vinje WE, Gallant JL. 2000. Natural Vision Sparse Coding and Decorrelation in Primary Visual Cortex During Sparse Coding and Decorrelation in Primary Visual Cortex During Natural Vision. *Science*

(80-) **287**:1273–1276. doi:10.1126/science.287.5456.1273

Vinken K, Vogels R, Op de Beeck H. 2017. Recent Visual Experience Shapes Visual Processing in Rats through Stimulus-Specific Adaptation and Response Enhancement. *Curr Biol* **27**:914–919. doi:10.1016/j.cub.2017.02.024

Wall NR, De La Parra M, Sorokin JM, Taniguchi H, Huang ZJ, Callaway EM, Parra MD La, Sorokin JM, Taniguchi H, Huang ZJ, Callaway EM. 2016. Brain-Wide Maps of Synaptic Input to Cortical Interneurons. *J Neurosci* **36**:4000–4009. doi:10.1523/jneurosci.3967-15.2016

Weskelblatt J, Niell CM. 2019. Distinct functional classes of excitatory neurons in mouse V1 are differentially modulated by learning and task engagement. *bioRxiv*. doi:.1037//0033-2909.I26.1.78

Willmore B, Tolhurst DJ. 2001. Network : Computation in Neural Systems Characterizing the sparseness of neural codes **6536**:255–270. doi:dx.doi.org/10.1080/net.12.3.255.270

Willmore BDB, Mazer JA, Gallant JL. 2019. Sparse coding in striate and extrastriate visual cortex 2907–2919. doi:10.1152/jn.00594.2010.

Willmore BDB, Mazer JA, Gallant JL. 2011. Sparse coding in striate and extrastriate visual cortex 2907–2919. doi:10.1152/jn.00594.2010.

Woloszyn L, Sheinberg DL. 2012. Effects of Long-Term Visual Experience on Responses of Distinct Classes of Single Units in Inferior Temporal Cortex. *Neuron* **74**:193–205. doi:10.1016/j.neuron.2012.01.032

Xu S, Jiang W, Poo MM, Dan Y. 2012. Activity recall in a visual cortical ensemble. *Nat Neurosci* **15**:449–455. doi:10.1038/nn.3036

Yang T, Maunsell JHR. 2004. The effect of perceptual learning on neuronal responses in monkey visual area V4. *J Neurosci* **24**:1617–1626. doi:10.1523/JNEUROSCI.4442-03.2004

Zhang S, Dan Y. 2014. Long-range and local circuits for top-down modulation of visual cortex processing. *Science (80-)* **2**:1970.

MATERIALS AND METHODS

Mice

All experiments and procedures were performed in accordance with protocols approved by the Allen Institute Animal Care and Use Committee. We used male and female transgenic mice expressing GCaMP6f in VIP inhibitory interneurons (double transgenic: VIP-IRES-Cre x Ai148 mice; https://www.jax.org/strain/010908; https://www.jax.org/strain/030328) or in excitatory glutamatergic neurons (triple transgenic: Slc17a7-IRES2-Cre x CaMKII-tTA x Ai93; https://www.jax.org/strain/023527, https://www.jax.org/strain/010712, https://www.jax.org/strain/024108) were used in these experiments. Mice were single housed and maintained on a reverse 12-hour light cycle (off at 9am, on at 9pm) and all experiments were performed during the dark cycle.

Surgery

Surgical procedures were performed as described in (de Vries et al., 2018). Headpost and cranial window surgery was performed on healthy mice that ranged in age from 5-12 weeks. Pre-operative injections of dexamethasone (3.2 mg/kg, S.C.) were administered at 12h and 3h before surgery. Mice were initially anesthetized with 5% isoflurane (1-3 min) and placed in a stereotaxic frame (Model# 1900, Kopf, Tujunga, CA), and isoflurane levels were maintained at 1.5-2.5% for surgery. An incision was made to remove skin, and the exposed skull was levelled with respect to pitch (bregma-lambda level), roll and yaw. The stereotax was zeroed at lambda using a custom headframe holder equipped with stylus affixed to a clamp-plate. The stylus was then replaced with the headframe to center the headframe well at 2.8 mm lateral and 1.3 mm anterior to lambda. The headframe was affixed to the skull with white Metabond and once dried, the mouse was placed in a custom clamp to position the skull at a rotated angle of 23° such that visual cortex was horizontal to facilitate the craniotomy. A circular piece of skull 5 mm in diameter was removed, and a durotomy was performed. A coverslip stack (two 5 mm and one 7 mm glass coverslip adhered together) was cemented in place with Vetbond (Goldey et al., 2014). Metabond cement was applied around the cranial window inside the well to secure the glass window. Post-surgical brain health was documented using a custom photo-documentation system and at one, two, and seven days following surgery, animals were assessed for overall health (bright, alert and responsive), cranial window clarity, and brain health. After 1-2 week recovery from surgery animals underwent intrinsic signal imaging for retinotopic mapping, then entered into behavioral training.

Intrinsic signal imaging

Intrinsic signal imaging (ISI) was performed as described in (de Vries et al., 2018) to produce a retinotopic map to define visual area boundaries and target *in vivo* two-photon calcium imaging experiments to the center of visual space in each imaged area. Mice were lightly anesthetized with 1-1.4% isoflurane administered with a somnosuite (model #715; Kent Scientific, CON). Vital signs were monitored with a Physiosuite (model # PS-MSTAT-RT; Kent Scientific). Eye drops (Lacri-Lube Lubricant Eye Ointment; Refresh) were applied to maintain hydration and clarity of eye during anesthesia. Mice were headfixed for imaging.

The brain surface was illuminated with two independent LED lights: green (peak λ=527nm; FWHM=50nm; Cree Inc., C503B-GCN-CY0C0791) and red (peak λ=635nm and FWHM of 20nm; Avago Technologies, HLMP-EG08-Y2000) mounted on the optical lens. A pair of Nikon lenses lens (Nikon Nikkor 105mm f/2.8, Nikon Nikkor 35mm f/1.4), provided 3.0x magnification (M=105/35) onto an Andor Zyla 5.5 10tap sCMOS camera. A bandpass filter (Semrock; FF01-630/92nm) was used to only record reflected red light onto the brain.

A 24" monitor was positioned 10 cm from the right eye. The monitor was rotated 30° relative to the animal's dorsoventral axis and tilted 70° off the horizon to ensure that the stimulus was perpendicular to the optic axis of the eye (Oommen and Stahl, 2008). The visual stimulus for mapping retinopy was a 20° x 155° drifting bar containing a checkerboard pattern, with individual square sizes measuring 25º, that alternated black and white as it moved across a mean-luminance gray background.

The bar moved in each of the four cardinal directions 10 times. The stimulus was warped spatially so that a spherical representation could be displayed on a flat monitor (Marshel et al., 2011).

After defocusing from the surface vasculature (between 500 μm and 1500 μm along the optical axis), up to 10 independent ISI timeseries were acquired and used to measure the hemodynamic response to the visual stimulus. Averaged sign maps were produced from a minimum of 3 timeseries images for a combined minimum of 30 stimulus sweeps in each direction (Garrett et al., 2014).

The resulting ISI maps were automatically segmented by comparing the sign, location, size, and spatial relationships of the segmented areas against those compiled in an ISI-derived atlas of visual areas. A cost function, defined by the discrepancy between the properties of the matched areas, was minimized to identify the best match between visual areas in the experimental sign map and those in the atlas, resulting in an auto-segmented and annotated map for each experiment. Manual correction and editing of the results included merging and splitting of segmented and annotated areas to correct errors. Finally, target maps were created to guide *in vivo* two-photon imaging location using the retinotopic map. The center of retinotopic space was computed from azimuth and altitude maps and adjusted for variability in eye position relative to the monitor by zeroing to the anatomical center V1. The corresponding retinotopic location was identified for each visual area, and two-photon imaging was targeted to a region within 20° of the center of gaze.

Behavior Training

Water restriction and habituation

Throughout behavior training mice were water-restricted in order to maintain consistent motivation to learn and perform the behavioral task (Guo et al., 2014). Prior to water restriction mice were weighed once daily for three days to obtain a stable, initial baseline weight. During the first week of water restriction mice were handled daily and habituated to increasing duration of head fixation in the behavior enclosure over a five-day period. Thus, the first day of behavior training occurred after 10 days of water restriction. Mice were trained 5 days per week and could earn as much water as possible during the daily one hour sessions; supplemental water was provided if earned volume fell below 1.0 mL and/or body weight fell under 80-85% of their initial baseline weight. On non-training days mice were weighed and received enough water provision to reach their target weight of 80-85% (never less than 1.0 mL per day).

Apparatus

Headposted mice were trained in custom-designed, sound-attenuating behavior enclosures. Visual stimuli were displayed on a 24" LCD monitor (ASUS, Model # PA248Q) placed at a ~15cm distance from the mouse's right eye. The monitor was rotated 30° relative to the animal's dorsoventral axis and tilted 70° off the horizon to ensure that the stimulus was perpendicular to the optic axis of the eye (Oommen and Stahl, 2008). A behavior stage was placed in a consistent location using a kinematic mount and consisted of a standardized headframe clamp to enable repeatable positioning of the mouse relative to the monitor, and a 6.5" running wheel tilted upwards by 10-15 degrees. Running behavior was measured by a rotational encoder. Water rewards were delivered using a solenoid (NResearch, Model #161K011) that allowed for a calibrated volume of fluid to pass through a blunted, 17g hyperdermic needle (Hamilton) positioned approximately2-3mm from the animal's mouth. Licks were detected by a capacitive sensor coupled to the reward delivery spout. Running speed, lick times, and reward delivery times were recorded on a NI PCI-6612 digital IO board and sampled at the frequency of the visual display (60 Hz).

Behavioral training procedure

Mice were trained for 1 hour/day, 5 days/week using an automated training algorithm. Briefly, mice were trained to lick when the identity of a flashed visual stimulus changed. If mice responded correctly within a short, post-change response window (750ms) a water reward (5-10uL) was delivered. On Day 1 of the automated training procedure mice received a short, 15-min "open loop" conditioning

session during which non-contingent water rewards were delivered coincident with 90 degree changes in orientation of a full-field, static square-wave grating. This session was intended to 1) introduce the mouse to the fluid delivery system and, 2) provide the technician an opportunity to identify the optimal lick spout position for each mouse and 3) condition the association between stimulus changes and reward delivery. Each session thereafter was run in "closed loop" mode, and progressed through 3 stages of the operant task (schematized in Figure 1D): 1) static, full-field square wave gratings (changes between 0 and 90 degrees), 2) flashed, full-field square-wave gratings (changes between 0 and 90 degrees) presented for 250ms with an 500ms inter stimulus gray period, and 3) flashed full-field natural scenes (8 natural images from the Allen Institute Brain Observatory) presented for 250ms with an 500ms inter stimulus gray period. Progression through each stage required mice to achieve a session maximum d' of 2 on two of the last 3 sessions. Thus, the shortest amount of time to reach the final stage of training was 5 sessions. Once in stage 3, mice were considered 'ready for imaging' when 2 out of 3 sequential sessions had a d' greater than 1 and mice performed at least 100 contingent trials. Mice were transitioned to behavior on the two-photon rig as scheduled time on the microscope became available. This resulted in a variable training duration in stage 3 across mice (Supplemental Figure 1A).

Session and trial structure

Each behavior session consisted of a continuous series of GO and CATCH trials, schematized in Figure 1B. Briefly, prior to the start of each trial a change-type and change-time were selected. Change-type was chosen based on predetermined frequencies such that GO and CATCH trials occurred with equal probabilities for sessions with 2 oriented gratings. For the natural image phase in which there were 64 change-pair possibilities, CATCH frequency was set to 12.5% (1/8 of the number of image transitions). To ensure even sampling of all stimulus transitions, a transition path is selected at random from a matrix of 1000 pre-generated paths. Each path takes a pre-determined route through each of the 64 possible transitions, including same-to-same, or catch, transitions. Once a transition path is completed, another path is chosen at random.

Change times were selected from an exponential distribution ranging from 2.25 to 8.25 seconds (mean of 4.25 seconds) following the start of a trial. Catch trial times were drawn from the same distribution such that false alarm rates were measured with the same temporal statistics as change trials, to account for any learning of the temporal distribution of change times. On trials when a mouse licked prior to the change or catch time, the trial was restarted with the same scheduled change or catch time. To prevent mice from getting stuck on a single trial, the number of times a trial could be repeated was limited to five. In all, this trial structure permits equal sampling of GO and CATCH trials, that when combined with mouse's licking response, yields HIT, MISS, FALSE ALARM, and CORRECT REJECTION trials. In addition to the four trial types described above, behavior sessions contained a subset of "free reward" trials (GO trials followed immediately by delivery of a non-contingent reward). Behavior sessions across all phases began with 5 free-reward trials. Additionally, to promote continued task engagement, one of these free rewards was delivered after 10 consecutive MISS trials.

Two photon imaging during behavior

Visual Stimulation

Visual stimuli were generated using custom Python scripts written in PsychoPy (Peirce, 2007, 2008) and were displayed using an ASUS PA248Q LCD monitor, with 1920 x 1200 pixels. Stimuli were presented monocularly, and the monitor was positioned 15 cm from the mouse's eye, and spanned 120° X 95° of visual space. The monitor was rotated 30° relative to the animal's midline and tilted 70° off the horizon to ensure that the stimulus was perpendicular to the optic axis of the eye (Oommen and Stahl, 2008).

The monitor was gamma corrected and had a mean luminance of 50 cd/m^2. To account for the close viewing angle of the mouse, a spherical warping was applied to all stimuli to ensure that the apparent size, speed, and spatial frequency were constant across the monitor as seen from the mouse's perspective (Marshel et al., 2011). Visual stimuli were presented at 60Hz frame rate.

Visual stimuli consisted of a subset of the natural scene images used in the publicly available Brain Observatory dataset (https://observatory.brain-map.org/visualcoding/). The 32 natural images that we used originated from 3 different databases of natural scene images: the Berkeley Segmentation Dataset (images 000, 005, 012, 013, 024, 031, 034, 035, 036, 044, 047, 045, 054, 057) (Strasburger et al., 2011), the van Hateren Natural Image Dataset (images 061, 062, 063, 065, 066, 069, 072, 073, 075, 077, 078, 085, 087, 091) (van Hateren and van der Schaaf, 1998), and the McGill Calibrated Colour Image Database (images 104, 106, 114, 115) (Olmos and Kingdom, 2004). The images were presented in grayscale, contrast normalized, matched to have equal mean luminance, and resized to 1174 X 918 pixels.

Behavior apparatus

Running speed measurement, lick detection and reward delivery were performed as described above for behavioral training. The monitor was placed in a fixed location relative to the behavior stage to ensure a consistent relationship between the mouse's eye and the screen. Running speed, lick times, and reward delivery times were recorded on a NI PCI-6612 digital IO board and sampled at the frequency of the visual display (60 Hz).

Two-photon calcium imaging during behavior

Calcium imaging was performed using a Scientifica Vivoscope (https://www.scientifica.uk.com/products/scientifica-vivoscope). Laser excitation was provided by a Ti:Sapphire laser (Chameleon Vision – Coherent) at 910 nm. Pre-compensation was set at -10,000 fs2. Movies were recorded at 30Hz using resonant scanners over a 400 µm field of view (512x512 pixels). Temporal synchronization of calcium imaging, visual stimulation, reward delivery and behavioral output (lick times and running speed) was achieved by recording all experimental clocks on a single NI PCI-6612 digital IO board at 100 kHz.

Behavior sessions under the two-photon microscope were 1 hour in duration, with task parameters identical to stage 3 of the behavior training procedure as described above. In addition, during most two-photon imaging sessions, 5% of stimulus flashes were randomly omitted, excluding the change flash and the flash immediately prior to the change. These omitted flashes were added to the experimental protocol partway into the experiment, resulting in 86/101 (85%) imaging sessions including omitted flashes. The 15 sessions without omitted flashes included data from one Slc17a7-IRES2;CaMKII-tTA;Ai93 mouse (4 sessions in VISp), and two Vip-IRES-Cre;Ai148 mice (3 sessions from VISal, and 8 sessions from VISp). Sessions without omitted flashes were excluded from any analysis depending on stimulus omission.

Movies of fluorescence were acquired near the center of retinotopic space in VISp and VISal, using ISI target maps and vasculature images as a guide. Once a cortical region was selected, the objective was shielded from stray light coming from the stimulus monitor using opaque black tape. All recordings were made at a depth of ~175um from the brain surface. Once a field of view was selected, a combination of PMT gain and laser power was selected to maximize laser power (based on a look-up table against depth) and dynamic range while avoiding pixel saturation (max number of saturated pixels <1000). Immersion water was occasionally supplemented while imaging using a micropipette taped to the objective (Microfil MF28G67-5 WPI) and connected to a 5 ml syringe via an extension tubing. At the end of each experimental session, a z-stack of images (+/- 30 µm around imaging site, 0.1 µm step) was collected to evaluate cortical anatomy and evaluate z-drift during experiment. Experiments with z-drift above 10µm over the course of the entire session were excluded.

For each field of view, imaging and behavior sessions were conducted using each of the 4 image sets shown in Figure 1C, including the familiar image set A used during behavior training, and 3 novel image sets first experienced by the mouse during the imaging phase of the experiment. On subsequent imaging days for a given field of view, we returned to the same location by matching (1) the pattern of vessels in epi-fluorescence with (2) the pattern of vessels in two photon imaging and (3) the pattern of cellular labelling in two photon imaging at the previously recorded location.Typically, only one field of view was imaged per mouse, however in 3 out of the 21 mice, fields of view were recorded in both VISp and VISal. In cases where an imaging session failed our QC criteria (for example for z-drift >10um, or

due to hardware issues such as dropped stimulus of imaging frames; see below), the session was retaken. As a result, some sessions with 'novel' image sets B, C or D were the second or third exposure (67% were first exposure, 27% were the second exposure, 6% were the third or fourth exposure). In contrast, mice were exposed to familiar image set A for average of 17 +/- 14 sessions during training.

Quality control

All data streams were required to pass an initial integrity check. Frame sync times for 2-photon can have no more than 6 dropped frames, and a mean physiology period (frame rate) between 0.032 - 0.034. The visual stimulus presentation sync times can have no more than 60 dropped frames and an average frame interval between 0.0165 and 0.0167. The display lag of the monitor is measured using a photodiode to compare with recorded frame times, and the display lag must not exceed 150ms. The running wheel encoder data stream is examined for any visible artifacts (such as a spike in the trace or a flat trace despite running activity). The average intensity of the 2-photon field of view may not drift more than 10% over the course of a session.The acquired movie is checked for saturation to ensure that no more than 500 saturated pixels are present across the duration of the recording session. Z-drift is quantified by performing phase correlation between the frames of a 100um z-stack taken after the imaging session and a 500 frame average from the beginning of the 2-photon movie and a 500 frame average at the end of the movie. If the distance between the z-stack frames found to be most correlated with the beginning and end of the movie is greater than 10um, the session is retaken.

Data processing

All data processing was performed as described in de Vries et al., 2018.

For each two-photon imaging session, the image processing pipeline included the following steps: 1) motion correction, 2) image normalization to minimize confounding random variations between sessions, 3) segmentation of ROIs, and 4) ROI filtering. Motion correction was performed using phase correlation and rigid translation. Segmentation was performed by morphological filtering on normalized periodic average images constructed from 400 frame blocks, followed by unification of masks across all blocks. ROI filtering was performed to remove segmented regions that were unlikely to correspond to cell somas, based on attributes including size and shape (for example, small ROIs likely corresponding to apical dendrites were removed).

Following identification of cell ROIs, the following steps were performed to obtain $\Delta F/F$ (dF/F) traces: 1) neuropil subtraction, 2) trace demixing, 3) $\Delta F/F$ computation. For each ROI, a neuropil mask was created, consisting of a 13 pixel ring around the cell soma, excluding any other ROIs. The raw fluorescence trace was generated by averaging all pixels within each cell ROI and the neuropil mask. A neuropil contamination ratio was computed for each ROI and the calcium trace was modeled as $F_M = F_C + rF_N$, where F_M is the measured fluorescence trace, F_C is the unknown true ROI fluorescence trace, F_N is the fluorescence of the surrounding neuropil, and r is the contamination ratio. After determination of r, we computed the true trace as $F_C = F_M - rF_N$, which is used in all subsequent analysis. To avoid artificially correlating neurons' activity by averaging fluorescence over two spatially overlapping ROIs, we demixed the activity of all recorded ROIs, as described de Vries et al., 2018. Finally, the global dF/F trace for each cell was computed, with the baseline F_0 determined by a rolling median filter of 180 seconds across the raw fluorescence trace.

Temporal alignment was performed to link two-photon acquisition frames (30Hz frame rate) with visual stimulation frames (60Hz frame rate) and associated behavioral signals (licking, running speed, reward delivery, sampled at 60Hz frame rate of visual stimulus). The visual stimulus time nearest to each two-photon (2P) frame time was computed, with the condition that the visual stimulus time must be before the 2P acquisition time, to ensure that dF/F responses were not attributed to stimulus or behavior events occurring after the change in the calcium signal.

Data Analysis

Behavior

Response rates for GO and CATCH trials were calculated by evaluating the fraction of trials of each type where a lick was registered within the 750ms response window following the change or sham change time (Figure 1H). The fraction of GO trials with a response is the hit rate and the fraction of CATCH trials with a lick response is the false alarm rate. The d-prime value for each session was computed as:

$$d' = Z[hit\ rate] - Z[false\ alarm\ rate]$$

Where Z is the inverse of the cumulative distribution function (using scipy.stats.norm.ppf).

Reaction time was calculated as the time to first lick after the start of the change time on GO trials. Mean run speed was calculated by taking the average of the running speed trace in a +/-2 second window around the image change time for each GO trial, then averaging across all GO trials in each session (Supplemental Figure 1D).The average running speed trace across sessions (Supplemental Figure 1E) was computed by averaging the running speed trace across all GO trials in a [-2,6] second window around the change time for GO trials or the sham change time for CATCH trials.

Calculation of all behavior metrics was limited to the portion of the session where the mouse was actively engaged in the behavioral task, where engagement was defined those periods during which the mouse earned at least 2 rewards per minute. Mice performed 248 engaged GO trials per session on average (range = 83-335).

Physiology

All analysis was performed on the global dF/F traces (where baseline F was computed as the mode of a rolling 3-minute window, described in data processing section above).

Neural responses were analyzed for three main conditions of interest – the time of image presentations (across all stimulus flashes), the time of image changes (on GO trials, first flash after an image transition), and the time of stimulus omissions (5% of all non-change image flashes were randomly omitted). For each individual image presentation (or image omission), the mean response in a 500ms window after stimulus onset (or the time of omission) was computed (including the 250ms stimulus duration and 250ms after, to account for the slow decay of GCaMP6 responses and to include cells with delayed responses or off responses after stimulus offset).

The preferred stimulus for a cell was identified as the image evoking the largest trial-averaged response for a given condition. When considering all image presentations, the preferred stimulus is the image that evoked the largest mean response, averaged across all stimulus presentations of each image. For image changes, the preferred stimulus is the image that evoked the largest mean change response, considering only the first image presentation after a change in stimulus identity. Thus, an individual cell may have a different preferred image when considering all image flashes versus only the change flash. For some analyses of stimulus omissions, we considered omission trials where the preceding image was the cell's preferred image defined across all stimulus presentations. Other analyses were agnostic to the identity of the image before and after the omission. Any analyses considering the image prior to the omission are indicated in text and figure legends.

Responsiveness was evaluated on a trial by trial basis (here an individual image presentation is considered as a 'trial'). The mean response for each image flash was compared to a shuffled distribution of dF/F values taken from omission periods (the longest period of extended gray screen during the session) for each cell. A p-value was computed by resampling the shuffled distribution 10,000 times and determining the fraction of comparisons where the mean response was larger than the shuffled values. If the trial had a significantly larger response compared to the shuffled distribution, that trial was deemed responsive. For a cell to be considered *image responsive*, at least 25% of all stimulus presentations for the preferred image for that cell must be significant per the above definition (Figure 2G). The distribution of fraction significant image trials (for each cell's preferred image) is shown in Supplemental Figure 2B. For a cell to be considered *change responsive*, at least 25% of all change flashes of the preferred image must be significant (Figure 2H). The distribution of fraction of significant change trials (for each cell's preferred change image) is shown in Supplemental Figure 2C.

For omission responsiveness, a different definition was used. In this case, a p-value was computed comparing the dF/F trace during the omission window (750ms after the omitted stimulus onset time, up to the time of the next stimulus onset) with a 750ms second baseline period prior to the omission (which includes the prior stimulus presentation and one inter-stimulus gray period). If 25% of all stimulus omission trials had a p-value > 0.05, with the mean omission response being greater than the baseline response, the cell was considered *omission responsive* (Figure 2I). The distribution of fraction of responsive omission trials is shown in Supplemental Figure 2C.

To evaluate image selectivity, we created tuning curves of mean response across the 8 images shown in each session, for each cell, for both the change flash (Supplemental Figure 3) and across all flashes (Figure 3). We generated a normalized image tuning curve for each image set by sorting each cell's tuning curve by the strength of the mean response to each image, normalizing to the max, and averaging across cells (Figure 3G, Supplemental Figure 3E). Only cells meeting the criteria for image or change responsiveness described above were included in the calculation of the image tuning curve. An insufficient number (<10) of VIP cells were image responsive for image set A, thus a tuning curve was not included for this condition in Figure 3G. To quantify selectivity for individual cells, we used a lifetime sparseness metric, computed using the definition in (Vinje and Gallant, 2000):

$$Sparseness = \frac{1 - \frac{1}{N}\frac{(\sum_i r_i)^2}{\sum_i r_i^2}}{1 - \frac{1}{N}}$$

where N is the number of images and r_i is the response of the neuron to image i averaged across trials. Lifetime sparseness was only computed for image or change responsive cells. Population sparseness was computed with the same metric, but where N is the number of neurons and r_i is average response vector of neuron i to all images (Figure 3D, Supplemental Figure 3C). Lifetime and population sparseness were computed for the change flash only (Supplemental Figure 3), and for the mean response across all image flashes (Figure 3).

We computed a reliability metric by taking the average of the trial-to-trial cross correlation of the dF/F trace in a [-0.5, 0.75] second window around the stimulus onset time for the preferred image for each cell (see example cells in Figure 4D). This metric was evaluated under several conditions: for change responsive cells (Supplemental Figure 4E), for cells that were image selective (lifetime sparseness value > 0.3) vs. unselective (lifetime sparseness < 0.3) (Supplemental Figure 4F), and as a function of stimulus repetition number after a change for the preferred image (repetition 1 = first flash after a change, repetition = 10 is the 10^{th} flash in the sequence for that image) (Figure 4E).

To better understand the difference in response strength between the first flash after a change in image identity compared to after multiple repetitions of a given image, we computed a change modulation index (CMI):

$$change\ modulation\ index = \frac{(R_1 - R_{10})}{(R_1 + R_{10})}$$

Where R_1 is the trial averaged response to the 1^{st} flash after a change and R_{10} is the trial averaged response to the 10^{th} flash after a change. This metric was computed for each cell, either taking the trial average across all images (Supplemental Figure 4D) or the trial average only for each cell's preferred image (Supplemental 4C) then averaged across all cells within a session to produce the plots in Supplemental Figures C&D. For Supplemental Figure 4B, the CMI listed in the title for each plot was computed using the mean of the population average trace across all images for the 1^{st} and 10^{th} repetitions.

A stimulus modulation index, measuring the modulation of the response at the stimulus frequency, was computed as in (Matteucci et al., 2019):

$$stimulus\ modulation\ index = \left| \frac{PS(f_1) - \langle PS \rangle_f}{\sqrt{\langle PS^2 \rangle_f - \langle PS \rangle_f^2}} \right|,$$

Where PS is the power spectral density (computed using scipy.signal.welch() with nperseg=128 and fs = 1/0.75) of the trial-averaged response over an 8 second window after the time of the change to the cell's preferred stimulus (time = 0-8sec in example figures in Figure 4G), f_1 is the stimulus frequency (1/0.75, images are presented every 750ms), and $\langle \ \rangle_f$ indicates the average over frequencies. This metric quantifies the difference between the power at the stimulus frequency and the average of the power spectrum. The distribution of the signal modulation index value for the preferred image across cells is shown in Figure 4H.

The dynamics of cell responses were evaluated by computing a ramp index over different time windows of interest, similar to (Makino and Komiyama, 2015):

$$Ramp\ index = \log_2(R_{late)} - R_{early)}$$

Where $R_{late)}$ is the mean response in the first half of a defined window of time, and $R_{early)}$ is the second half of the window. This index provides a measure of the magnitude and direction of a change in a signal within the window. For Figure 5, the ramp index was computed for two windows: the pre-stimulus window (250ms prior to stimulus onset; Figures 5A&B light shading) and the stimulus window (250ms after stimulus offset; Figures 5A&B dark shading) for the mean dF/F trace for each cell across all flashes of its preferred image. If the dF/F values are increasing during the window, the ramp index is positive. If the dF/F values are decreasing during the window, the ramp index is negative. The ramp index was only computed for cells with a mean dF/F value of >0.05 in the stimulus window, as the ramp index for a flat, unchanging signal can result in extreme values with little meaning.

The pre-stimulus and stimulus ramp indices were plotted against each other on a cell by cell basis (for cells with a minimum level of activity as described above) and found to be correlated by least squares linear regression between the two measures, performed separately for each image set (scipy.stats.linregress). Cells fell into four quadrants of this graph based on the relative sign of pre-stimulus and stimulus ramping. Cells with positive values of the stimulus ramp index and negative values of the pre-stimulus ramp index, indicating stimulus evoked activity with no pre-stimulus ramping, belonged to quadrant 1 (Q1). Most cells fell into this category. At the other extreme, cells with negative values of the stimulus ramp index and positive values of the pre-stimulus ramp index fall in Q4, indicating increasing activity prior to stimulus onset, and decreasing activity after stimulus offset. These cells showed the largest change in proportion between the familiar and novel image sets, particularly for VIP cells (Figure 5D). The fraction of cells belonging to each quadrant was computed for each imaging session, then averaged across sessions, in figure 5D. To illustrate the response dynamics associated with these cell groups, the mean response across all cells belonging to each quadrant was computed, then normalized to its max, for the plots in Figure 5E.

After grouping cells by their response dynamics, several metrics were computed, as described previously, for each group, separated by Cre line and image set, including: time to peak response for the average of all image flashes (Figure 5F), the stimulus modulation index (Figure 5G), reliability across all image flashes (Figure 5H), and lifetime sparseness across all flashes (Figure 5I). Only cells with a minimum level of activity (<0.05 dF/F) were included in this analysis. Plots in Figures 5F-I show the mean+/-95% confidence intervals across included cells.

The ramp index described above was again used to quantify the increase in activity during stimulus omission (Figure 6). For cells with a minimum level of activity during the omission period (>0.05 mean dF/F; Figure 6E), the ramp index was computed over the 750ms window, taking 250ms starting at the time where the omitted stimulus would have been presented as the early portion of the window, and the 250ms prior to the beginning of the next image presentation as the late portion of the window (note that 250ms in the middle of this 750ms window was unused). A positive value of this index indicates increasing activity over the omission period. The distribution of the ramp index values over this window across cells with a minimum level of activity during omission quantified in Figure 6F. The relationship between pre-stimulus ramping and omitted ramping is shown in Figure 5G on a cell by cell basis, with the strength and significance of the correlation determined using least squares linear

regression (scipy.stats.linregress) as before. The fraction of cells having both increasing activity during the pre-stimulus period and increasing activity following stimulus omission are quantified in Figure 5H.

Statistics and data visualization

All statistical comparisons were made across image sets, within each Cre line. ANOVA (scipy.stats.f_oneway) was used to test for an effect of image set, followed by a paired t-test (scipy.stats.ttest_ind) with Bonferroni correction for multiple comparisons for each individual image set pair. p-values are reported throughout the text and figure legends, and significance of comparisons ($p < 0.05$) is indicated by an asterisks in figure insets. Point plots (seaborn.pointplot()) show individual sessions as gray points and the mean +/- 95% confidence intervals in color. Point plots lacking gray points are the mean +/- 95% confidence intervals across cells. Cumulative distributions across cells were generated with seaborn.distplot() with hist=False, hist_kws={'cumulative':True}. Barplot show either the mean across cells or the mean +/-SEM across cells. Heatmaps of cell responses were created using seaborn.heatmap().

Figure 1: Natural image change detection task with familiar and novel images

Figure 1: Natural image change detection task with familiar and novel images. A) Schematic of stimulus presentation during task. Images are presented for 250ms followed by 500ms of gray screen. **B)** Schematic of trial structure. On go trials, a change in image identity occurs and mice must lick within the 750ms response window to receive a water reward. On catch trials, no stimulus change occurs and the behavioral response is measured to quantify guessing behavior. **C)** Schematic of training sequence. Mice are initially trained with static gratings of 2 orientations, first with no gray screen in between grating presentations (stage 1), then introducing the 500ms inter stimulus interval (stage 2). After achieving consistent performance with flashed gratings, mice are transitioned to change detection with 8 natural scene images (image set A). During the imaging portion of the experiment, mice are tested with image set A as well as 3 novel image sets (B, C, D) on subsequent days. **D)** Four sets of 8 natural images were shown, in separate sessions. **E)** Example training trajectory for one mouse. Colors correspond to stages in panel D. Triangles represent behavior-only and circles represent behavior + imaging sessions. **F)** Number of sessions spent in each stage across mice. Mean +/- 95% confidence intervals in color, individual mice in gray. **G)** Example behavior performance across 5 minutes of one session, separated into 5 one minute epochs. Colored vertical bars indicate stimulus presentations, different colors are different images. Licks are shown as green tick marks. Licks within 750ms of a change (transition between two colors) result in reward delivery, shown as purple triangles. 5% of all non-change image flashes are omitted, visible as a gap in the otherwise periodic stimulus sequence. **H)** Response rates for go and catch trials are similar across image sets, demonstrating that mice generalize task performance to novel images. Individual behavior sessions are shown in gray and average +/- 95% confidence intervals across sessions for each image set are shown in colors corresponding to image sets in C (p>0.05 for all image set comparisons). **I)** No significant difference in average response latency across image sets (p>0.05 for all image set comparisons). Mean +/- 95% confidence intervals in color, individual sessions in gray.

Supplemental Figure 1: Behavior performance across image sets

Supplemental Figure 1: Behavior performance across image sets. A) Time in each training stage for all mice in the study. Colors correspond to training stages in Figure 1D. **B)** Hit rate for each of the 8 images in the 4 image sets. Mean and 95% confidence intervals are shown in black with gray points for individual behavior sessions. **C)** Mean d-prime is similar across image sets (no significant difference across image set pairs except A-B, where p=0.002). **D)** Mean running speed is similar across image sets (p>0.05 for all comparisons). Error bars in B-D are 95% confidence intervals. **E)** Running behavior is similar across image sets, average +/- SEM for go and catch trials (n=6260+/-294 go trials, n=941+/-40 catch trials).

Figure 2: Activity in L2/3 excitatory and VIP inhibitory cells during image change detection behavior

Figure 2: Activity in L2/3 excitatory and VIP inhibitory cells during image change detection behavior. A) Maximum intensity projection from a 2-photon imaging session in L2/3 of an Slc17a7-IRES2-Cre;CaMk2-tTa;Ai93 mouse expressing GCaMP6f in excitatory neurons. **B)** Same as in A for L2/3 inhibitory neurons in a Vip-IRES-Cre; Ai148(GCaMP6f) mouse. **C)** Examples cells. Left panel shows dF/F traces from 4 excitatory cells from a 90 second epoch of a behavior session with the familiar image set A in VISal. Scale bars on the left of each row indicate 150% dF/F. Colored vertical bars indicate presentation of the 8 natural images. Bottom row shows licking behavior in green with reward delivery in purple. Right panel shows the response of the same 4 cells to each image, averaged across all presentations of each image, as well as the average response when flashes are omitted (right column, gray indicates time of stimulus omission). Scale bars indicate 25% dF/F. Bottom row of right panel shows the 8 natural images for this session. Note the selectivity of excitatory responses and temporal dynamics relative to stimulus onset. **D)** Same as C) for 4 excitatory cells from a session with novel image set C in VISal. Left panel scale bars indicate 150% dF/F, right panel scale bars indicate 25% dF/F. **E)** Same as C) for 4 VIP cells from a session with familiar image set A in VISal. Note the dynamics of the response relative to stimulus onset and stimulus omission. Left panel scale bars indicate 250% dF/F, right panel scale bars indicate 25% dF/F. (legend continued on next page)

Figure 2: Activity in L2/3 excitatory and VIP inhibitory cells during image change detection behavior (continued). F) Same as C), F for 4 VIP cells for a session with novel image set C in VISal. Note that stimulus evoked responses are unselective and highly correlated across the population. Left panel scale bars indicate 250% dF/F, right panel scale bars indicate 25% dF/F. **G)** A larger fraction of cells are image responsive for novel image sets compared to the training set A. Responsiveness is defined for each cell as having >25% of preferred image flashes with a significant response compared to a shuffled distribution of stimulus omissions. The fraction of responsive cells is shown for each session in gray, with mean +/- 95% confidence intervals in color. $p < 0.0005$ for all comparisons with image set A for excitatory cells and $p < 0.01$ for all comparisons with image set A for VIP inhibitory cells. **H)** A larger fraction of cells are responsive to the first flash after an image change for novel image sets. Responsiveness computed as described in G, limited to the change flash only. $p < 0.0006$ for all comparisons with image set A for excitatory cells, $p > 0.05$ for all image set comparisons for VIP cells. **I)** Cells with elevated activity during stimulus omission are most common in VIP populations for the trained image set A. Omission responsiveness is defined as having >25% of omissions with a significantly larger response than the preceding image flash. $p < 0.02$ for A-B and A-C for VIP cells.

Supplemental Figure 2: Image, change, and omission responsiveness

Supplemental Figure 2: Image, change, and omission responsiveness. A) Top panel, number of segmented ROIs per imaging session by cell class and image set. Bottom panel, total number of sessions, mice and cells per image set and cell class. **B)** Cumulative distribution of the fraction of significant image flashes for the preferred image across cells, showing higher image responsiveness for novel images. Dotted lines indicate the mean of each distribution. Inset shows image set comparisons where p<0.0002. **C)** Cumulative distribution of the fraction of significant change trials for the preferred image across cells, showing higher change responsiveness for novel images. Inset shows image set comparisons where p<0.0002. **D)** Cumulative distribution of the fraction of responsive omission trials across cells,. Very few excitatory cells respond during stimulus omission, while a large number of VIP cells show omission related activity, especially for the familiar image set. **E)** Average response across all presentations of each image (columns, colored bars indicate stimulus presentation) and during stimulus omission (right column, gray bar indicates time of stimulus omission) in dark gray, and during the change flash for each image in light gray, for 10 excitatory cells from one session with familiar image set A in VISp. Cells with the top 10 SNR values were selected for display, where SNR is defined as the ratio of the mean and standard deviation of the dF/F trace. **F)** Same as panel E for 10 excitatory cells from a session with novel image set D in VISp. **G)** Same as panel E for 10 VIP inhibitory cells from a session with familiar image set A in VISp. **H)** Same as panel E for 10 VIP inhibitory cells from a session with novel image set B in VISp. Scalebars indicate 25% dF/F for E-H.

Figure 3: Response sparseness for familiar and novel images

Figure 3: Response sparseness for familiar and novel images. A) Mean response across images (columns) for all cells (rows) for familiar image set A and novel image sets B-D in L2/3 excitatory cells, for a 500ms window after stimulus onset, averaged across all presentations of each image. Note elevated activity for novel image sets. **B)** Mean response across images for all VIP inhibitory cells. **C)** Average response magnitude across the population is larger for novel image sets. The mean response across all flashes for each cells preferred image was computed before averaging across all cells. Barplot shows mean +/-SEM. **F)** Example tuning curves from one excitatory cell (top panel) and one VIP inhibitory cell (bottom panel) demonstrating high and low values of lifetime sparseness. **D)** Mean population sparseness across sessions for each cell class and image set. Population sparseness is first computed for each image, then averaged across images within a session. No significant differences are observed across image sets. Gray points are individual sessions, colors indicate mean +/- 95% confidence intervals. **E)** Example of mean population activity from one session in excitatory cells (top panel) and one session for VIP inhibitory cells (bottom panel), demonstrating high and low population sparseness values. Each plot shows population activity for a single image. **G)** Cumulative distribution of lifetime sparseness for image responsive cells in excitatory (top) and VIP inhibitory (bottom) populations, showing increased sparseness for familiar images in excitatory cells. For excitatory cells, inset shows comparisons where p<0.0005. For inhibitory cells, image set A was not included because there were insufficient image responsive cells, and there were no significant differences between novel image sets. **H)** Population tuning curve shows increased selectivity for familiar images for excitatory cells. Cell tuning curves were ranked according to their mean response across images before averaging over image responsive cells. Tuning curve for image set A in VIP cells is not shown because there were very few cells (<10) that met the image responsiveness criteria.

Supplemental Figure 3: Reduced sparseness following stimulus changes

Supplemental Figure 3: Reduced sparseness following stimulus changes. A) Mean response in a 500ms window after stimulus changes for all images (columns) for all cells (rows) for familiar image set A and novel image sets B-D in L2/3 excitatory cells. Note elevated activity for novel image sets. **B)** Mean response for image changes for all VIP inhibitory cells. **C)** Average change response magnitude across the population is larger for novel image sets. The mean response across all changes for each cells preferred image was computed before averaging across all cells. Barplot shows mean +/-SEM. **D)** Mean population sparseness for the change flash across sessions for each cell class and image set. No significant differences are observed across image sets. Gray points are individual sessions, colors indicate mean +/- 95% confidence intervals. **E)** Cumulative distribution of lifetime sparseness for change responsive cells in excitatory (top) and VIP inhibitory (bottom) populations, showing increased sparseness for familiar images following a change. Dotted lines indicate the mean of each distribution. For excitatory cells, inset shows comparisons where p<0.01. For inhibitory cells, inset shows comparisons where p<0.05. **F)** Population tuning curve for image changes shows increased selectivity for familiar images for excitatory cells. Cell tuning curves were ranked by image response magnitude before averaging. Plot only includes cells with a significant change response. **G)** Image tuning curves for familiar images comparing activity averaged across all presentations of each image (light gray) and activity for the change flash for each image (dark gray). Only image or change responsive cells were included. Tuning curve for all flashes for VIP inhibitory cells is not shown because very few cells (<10) met the criteria for image responsiveness. **H)** Selectivity following a stimulus change is broader compared to activity averaged across all flashes for novel images. Population tuning curve for the change response is shown in dark gray and for all flashes in light gray. Only image and change responsive cells were included.

Figure 4: Increased stimulus modulation and reliability for novel images

Figure 4: Increased stimulus modulation and reliability for novel images. A) Trial averaged response to the preferred image across all excitatory cells, aligned to the time of the change to the preferred image (averaging across multiple images prior to time = 0s). Note the larger fraction of cells with a visible change response for novel image sets B-D. **B)** Trial averaged responses of VIP inhibitory cells for their preferred stimulus, aligned to the change time. Note the increase in stimulus evoked activity for novel image sets B-D. **C)** Population average response for all cells to their preferred image for excitatory and VIP inhibitory populations, aligned to the time of the change to the preferred image. Population response strength is consistently higher for novel image sets compared to the familiar image set A. For excitatory cells, the change response has the largest magnitude followed by a reduction in activity with stimulus repetition. Traces show mean+/-SEM across all cells. **D)** The change modulation index compares the response to the first stimulus following a change with the 10th repetition of that image. The index is computed for the preferred image of each cell, then averaged within a session (gray points). Colored points show mean +/- 95% confidence intervals. No significant differences across image sets for excitatory or VIP. **E)** Example cell responses for the first image flash after a change, showing individual trials for the cell's preferred stimulus in gray and the trial average in black, demonstrating high reliability (top panel, average trial to trial correlation of the dF/F trace = 0.94), and lower reliability (lower panel, average trial to trial correlation = 0.39). **F)** Responses to novel images and stimulus changes have the highest response reliability. Boxplots show the distribution of mean reliability across sessions, computed first for each cell as a function of stimulus repetition number, then averaged within a session. Repetition 1 is the first flash after an image change and repetition 10 is the 10th image presentation after a change. **G)** Example responses demonstrating strong modulation at the stimulus frequency (top panel, modulation index: 5.65 for the 0-8s period after the change time) and weak stimulus modulation (bottom panel, modulation index: 1.94 for t=0-8s). See methods for description of stimulus modulation index. **H)** Cumulative distribution of the stimulus modulation index across cells for each image set by cell class. Responses are more strongly stimulus modulated for novel image sets, especially for VIP populations. Dotted lines indicate the mean of each distribution.

Supplemental Figure 4: Change modulation and image preference

Supplemental Figure 4: Change modulation and image preference. A) Population response across each cell's preferred preferred image for each image set, plotted separately for the change flash (in color), and the 10th flash after a change (gray, adapted). Responses are normalized to the max of the change condition. CMI indicates the change modulation index comparing the strength of the response to the change to the adapted response. Higher values indicate stronger change response relative to the 10th repetition. Traces show mean +/-SEM across all cells. **B)** Population response across all images for each image set, plotted separately for the change flash (in color), and the 10th flash after a change (gray, adapted). Responses are normalized to the max of the change condition. CMI indicates the change modulation index comparing the strength of the response to the change to the adapted response (see Methods). Higher values indicate stronger change response relative to the 10th repetition. Traces show mean +/-SEM across all cells. **C)** Both excitatory and VIP populations show stronger activity for the change stimulus compared to following stimulus repetition when considering all images. Plots show the change modulation index computed for the mean response across all images for each cell, then averaged within sessions (gray points). Colored points show mean +/- 95% confidence intervals. For excitatory cells, p<0.005 for A-B and A-D. No significant differences across image sets for VIP cells. **D)** Responses to novel images (blue box plots) show higher trial to trial reliability compared to familiar images (red blox plots), for both image selective (lifetime sparseness > 0.3) and unselective (lifetime sparseness < 0.3) populations. Boxplots show distribution across responsive cells for each condition. Asterix on plot indicates comparisons where p<0.05.

Figure 5: Experience-dependent shift in the dynamics of VIP inhibitory cells

Figure 5: Experience-dependent shift in the dynamics of VIP inhibitory cells. A) Population average response across all stimulus flashes for all cells. The VIP population response shows image evoked activity for novel image sets. For familiar images, VIP population activity ramps up during the pre-stimulus period and peaks at the time of stimulus onset. Traces show mean+/-SEM across all cells. Inset, a ramping metric is used to quantify the relative change in activity within the pre-stimulus (-250-0 seconds prior to stimulus onset) and stimulus (0-250ms after stimulus onset) periods. The index compares the dF/F values in the first (early) and second (late) portions of the window. **B)** Example cells demonstrating response dynamics and associated ramping index values for the pre-stimulus and stimulus periods. Positive values of the ramp index indicate increasing dF/F values in the window, while negative values indicate decreasing values. Note the relationship between the ramp index in the pre-stimulus and stimulus windows. **C)** Relationship of pre-stimulus ramping index (x-axis) with stimulus ramping index (y-axis) shows a tradeoff between stimulus predictive and stimulus evoked signals. Cells can be separated into 4 quadrants depending on the relative value of pre- and stimulus ramping indices (Q1-4). For excitatory cells, r=-0.47 for image set A, r=-0.47 for B, r=-0.42 for C, r=-0.35 for D. p<0.005 for all image sets for excitatory cells. For VIP cells, r=-0.61 for image set A, -0.75 for B, -0.66 for C, -0.83 for D. p<0.005 for all image sets for VIP. **D)** Fraction of cells belonging to each quadrant across cell class and image set. Cells with predictive responses (Q4, increasing activity during pre-stimulus period, negative during stimulus period) are more prevalent for familiar images, particularly for VIP cells, whereas stimulus evoked responses (Q1, negative ramping during pre-stimulus window, positive during stimulus window) are more prevalent for novel image sets. **E)** Normalized population response for cells belonging to each quadrant from C). **F)** Time to peak is lower for Q4 pre-stimulus ramping cells, using the average response across all flashes for each cell to compute peak time. **G)** Stimulus modulation index is stronger for Q1 stimulus ramping type cells on average, except for familiar images with VIP (see Methods for definition of stimulus modulation index). H) Trial to trial response reliability across all flashes is higher for stimulus driven Q1 cells compared to stimulus predictive Q4 cells. **I)** Image selectivity is higher for Q1 cells compared to Q4 cells for excitatory populations, consistent with being a sensory driven population. Neither group is strongly selective for VIP populations. Only cells with a minimum level of activity (> 0.05 dF/F) in the stimulus window are included in panels C-I, as the ramp index is not meaningful for windows with a flat response profile.

Supplemental Figure 5: Stimulus and pre-stimulus ramping activity depend on experience

Supplemental Figure 5: Stimulus and pre-stimulus ramping activity depend on experience. A) Distribution of time to peak response across cells is shifted towards the right for familiar images (red) across the VIP population. Note the bi-modality of response timing for excitatory cells, likely corresponding to on- and off-responsive populations. No significant differences are observed across image sets for excitatory cells. For VIP cells, p<0.0005 for all comparisons with image set A. **B)** Distribution of pre-stimulus ramp index is shifted towards more positive values in sessions with image set A for both excitatory and VIP inhibitory populations, indicating an increased prevalence of predictive signals. For excitatory cells, p<0.0005 for all comparisons across image sets except B-D. For VIP cells, p<0.0005 for all comparisons with image set A, p<0.01 for B-D. **C)** Distribution of stimulus ramp index is shifted towards more positive values for novel image sets for both excitatory and VIP inhibitory populations, consistent with increased stimulus responsiveness for novel images. For excitatory cells, p<0.001 for all comparisons across image sets except B-D. For VIP cells, p<0.0005 for all comparisons with image set A.

Figure 6: VIP cells show strong ramping activity during stimulus omission

Figure 6: VIP cells show strong ramping activity during stimulus omission. A) Population average response following stimulus omission. On average, no change in activity is observed during stimulus omission in excitatory cells for either familiar or novel images (left panel). In sessions with novel images, the VIP population also shows little change in activity following stimulus omission (right panel, image sets B-D). In contrast, activity of the VIP population in sessions with familiar images (image set A) continues to ramp up until the time of the next stimulus presentation. Inset, window over which an omission ramp index is computed. Note the presence of image evoked activity before and after the omission for excitatory populations and VIP inhibitory populations for novel images, but not the familiar image set A. **B)** Example cells exhibiting a variety of response dynamics in response to stimulus (gray bars) and omission (after time = 0 sec). Cells typically show stimulus evoked activity and no omission response or show pre-stimulus ramping and strong omission responses. Some cells (example in top right) show a combination of stimulus evoked activity and omission related activity. **C)** Heatmap of activity around the time of stimulus omission across all excitatory cells, sorted by magnitude of the response in the omission window, averaged for omissions where the preceding image presentation was the cell's preferred image. Image responses are visible in the time before and after stimulus omission (t = 0 sec. white vertical line), but very few excitatory cells are active during stimulus omission. **D)** Heatmap of mean omission related activity in VIP cells, averaging across all omission trials for each cell. Time of omission indicated with a white line at t = 0 sec. Most VIP cells show ramping during omission for image set A, with fewer cells exhibiting omission ramping in sessions with novel image sets. **E)** Cumulative distribution of mean dF/F responses during the omission window across cells. Very few excitatory cells show activity during the omission window. Dotted lines indicate the mean of each distribution. For excitatory cells, p<0.0005 for all comparisons with image set A. For VIP cells, p<0.002 for all comparisons with image set A. **F)** Cumulative distribution of ramp index during stimulus omission showing increased ramping activity for VIP cells with image set A. Note that the omitted ramp index is only computed for cells with a mean dF/F value > 0.05 dF/F during the omitted window. No significant differences between image sets for excitatory cells. For VIP cells, p<0.0005 for all comparisons with image set A. **G)** The strength of the omitted ramp index (y-axis) and pre-stimulus ramp index (x-axis) are positively correlated, indicating that cells with stimulus predictive activity (positive pre-stimulus ramp index) typically also show enhancement following stimulus omission (positive omission ramp index). Fits were not significant for excitatory cells. For VIP, p<0.005 for all image sets, r=0.38 for image set A, r=0.62 for B, r=0.53 for C, r=0.58 for D. **H)** Fraction of omitted ramping cells with positive stimulus evoked responses is higher for novel image sets, especially for VIP cells, indicating that a subset of cells show a combination of stimulus evoked activity and enhancement during omission, similar to the example cell in the top row, right column of panel B.