1 # Expansion of apical extracellular matrix underlies the
2 # morphogenesis of a recently evolved structure
3

4 Sarah Jacquelyn Smith[1], Lance A. Davidson[2], Mark Rebeiz[1]*
5
6 [1] University of Pittsburgh Department of Biological Sciences, Pittsburgh, PA 15260, United States.
7 [2] University of Pittsburgh, Department of Bioengineering, Pittsburgh, PA 15260, United States.
8
9 *Corresponding author: rebeiz@pitt.edu

10
11

Abstract

One of the fundamental gaps in our knowledge of the evolution of novel structures is
understanding how the morphogenetic processes that form these structures arise. Here, we
traced the cellular development of a morphological novelty, the posterior lobe of *D.*
melanogaster. We found that this genital outgrowth forms through an extreme increase in cell
height. By examining the apical extracellular matrix (aECM), we uncovered a vast network
associated with the developing genitalia of lobed and non-lobed species. We observed that cells
which will form the posterior lobe show expanded expression of the aECM protein Dumpy which
connects them to the ancestral aECM network. Further analysis demonstrated a required role
for Dumpy in cell height increase during development. We propose that the aECM presents a
rich reservoir for generating morphological novelty, in addition to highlighting a yet unseen role
for aECM in regulating extreme cell height.

24
25

Introduction

Biologists have long been mesmerized by the appearance of morphological novelties, new
structures that appear to lack homologs in other species groups (Moczek, 2008; Günter et al.,
2010). To understand the origins of these novel structures, significant effort has focused on
determining how spatial and temporal patterning of genes are altered during evolution (Peter &
Davidson, 2015; Rebeiz, et al., 2015; Wagner, 2014). This has indicated how developmental
programs are often associated with morphological novelties, and they are frequently co-opted
from other tissues. However, limited attention has been directed to how novel structures form at
the cellular level. Understanding how a structure physically forms is important, as it can help
explain which morphogenetic processes might be targeted during evolution. In addition,
because most morphological novelties arose in the distant past, it is likely that the causative

genetic changes will be obscured by additional changes scattered throughout relevant gene
regulatory networks (Liu, Y., 2019). Hence, understanding the morphogenetic basis of a novelty
is critical to identifying the most important aspects of the gene regulatory networks that
contributed to its origin.
41
Most studies of morphogenetic evolution have focused on structures subject to diversification,
illuminating processes that contributed to their modification, as opposed to origination. For
example, studies of tooth morphogenesis have elucidated how both internal mechanisms, such
as cell shape changes (Li et al., 2016), and external forces, such as the pressure from the
surrounding jaw (Renvoisé et al., 2017) could be contributing factors in their diversification. An
examination of the enlarged ovipositor of *Drosophila suzukii* revealed how a 60% increase in
length was associated with increases in apical area and anisotropic cellular rearrangement
(Green et al., 2019). In addition, differences in early morphogenetic mechanisms between
distantly related species are observed in both the development of breathing tubes on the
Drosophilid eggshell (Osterfield et al., 2015) and migration of sex comb precursors on
Drosophila male forelegs (Atallah et al., 2009; Tanaka et al., 2009), together highlighting how
rapid changes in morphogenetic mechanisms can evolve to form the same structure. Overall,
these studies have illustrated how evolutionary comparative approaches can reveal
morphogenetic processes critical to the sculpting of anatomical structures.
56
Morphogenesis is the product of both cell intrinsic processes, such as those conferred by the
cytoskeleton or cell-cell junctions, and external forces from the environment in which the cell
resides. Extracellular mechanics are relatively understudied compared to intracellular
mechanics (Paluch & Heisenberg, 2009). An important component of the microenvironment of a
cell is the extracellular matrix (ECM) which can be subdivided into two populations of ECM, the
basal ECM and the apical ECM (aECM) (Brown, 2011; Daley & Yamada, 2013; Linde-Medina &
Marcucio, 2018; Loganathan et al., 2016). While comparatively understudied, recent work has
defined vital roles for aECM in the morphogenesis of structures, such as the *Drosophila* wing
(Diaz-de-la-Loza et al., 2018; Etournay et al., 2015; Ray et al., 2015), denticles (Fernandes et
al., 2010), and trachea (Dong et al., 2014), as well as in *C. elegans* neurons (Heiman et al.,
2009; Low et al., 2019). Despite recent interest in the aECM, its role in the evolution of
morphogenetic processes is currently unknown.
69

Genital traits represent a particularly advantageous system in which to study the morphogenetic

basis of novel structures. The study of morphological novelty is often difficult because most

structures of interest evolved in the distant past, rendering it difficult to understand the ancestral

ground state from which the novelty emerged. Genitalia are noted for their rapid evolution

(Eberhard, 1985), and thus bear traits among closely-related species that have recently evolved

in the context of a tissue that is otherwise minimally altered. For example, the posterior lobe, a

recently evolved anatomical structure present on the genitalia of male flies of the *melanogaster*

clade (Kopp & True, 2002) (Figure 1A), is a three-dimensional outgrowth that is required for

genital coupling (Frazee & Masly, 2015; Jagadeeshan & Singh, 2006; LeVasseur-Viens et al.,

2015). Besides the posterior lobe, the genitalia of lobed and non-lobed species are quite similar

in composition, providing an excellent context in which to examine the morphogenesis of the

ancestral structures from which the posterior lobe emerged.

Here, we find cell shape changes which increase cell height along the apico-basal axis drive

morphogenesis of the posterior lobe. We investigated internal and external factors that might

contribute to this height increase and find a correlation between the aECM protein Dumpy and

the height of posterior lobe cells. Comparisons to non-lobed species uncovered the presence of

a conserved aECM network on the genitalia that has expanded to cells that form the posterior

lobe. Our work shows how the formation of a morphological novelty depends upon novel aECM

attachments, integrating cells into a larger pre-existing aECM network.

Results

The posterior lobe grows from the lateral plate epithelium

The male genitalia of *Drosophila* is a bilaterally symmetrical anatomical structure which forms

from the genital disc during pupal development. In adults, the posterior lobe protrudes from a

structure called the lateral plate (also known as the epandrial ventral lobe (Rice et al., 2019))

(Figure 1A,D; Figure 1 - video 1). In *D. melanogaster*, prior to posterior lobe formation, the

lateral plate is fully fused to a neighboring structure called the clasper (also known as the

surstylus (Rice et al., 2019)) (Figure 1B) (Glassford et al., 2015). The lateral plate begins to

separate from the clasper around 32 hours after pupal formation (APF) in *D. melanogaster*

(Figure 1 – supplement 1)*.* Approximately 4 hours later, the posterior lobe begins to project from

the plane of the lateral plate and achieves its final shape by 52 hours APF (Figure 1D; Figure 1-

supplement 1). During posterior lobe development, cleavage of the lateral plate from the clasper

continues, dropping the tip of the lateral plate behind the clasper and separating both tissues

(Figure 1D; Figure 1 - supplement 1). Full separation of the lateral plate and clasper stops

slightly above (ventral to) the posterior lobe (Figure 1 - supplement 1). By contrast, the lateral

plate in the non-lobed species *D. biarmipes* remains flat throughout development, but all other

morphogenetic events are very similar, forming on a schedule that is approximately 4 hours

behind *D. melanogaster* (Figure 1C,E; Figure 1 - supplement 1).

Posterior lobe cells increase in height to protrude from the lateral plate

To investigate which cellular behaviors are unique to lobed species, we examined how the

posterior lobe grows from the lateral plate in both lobed and non-lobed species. First, we looked

at cell proliferation, which commonly contributes to morphogenesis through patterned and/or

oriented cell division (Heisenberg & Bellaïche, 2013), such as observed during branching

morphogenesis in the lung where oriented cell division expands the bud before it bifurcates into

two branches (Schnatwinkel & Niswander, 2013). During stages prior to the development of the

posterior lobe morphogenesis, we observed widespread cell proliferation throughout the entire

genital epithelium (Figure 2 - supplement 1). However, proliferation declines tissue-wide and all

cell proliferation is essentially absent during posterior lobe development (Figure 2 - supplement

1). Similar dynamics in proliferation are also observed in non-lobed species (Figure 2 -

supplement 1), suggesting that proliferation is not a major contributor to the morphogenesis of

the posterior lobe.

Next we tested the possibility that cell intercalation could contribute to posterior lobe

morphogenesis. Such processes may play a role in tissue elongation (Guirao & Bellaïche, 2017;

Tada & Heisenberg, 2012; Walck-Shannon & Hardin, 2014), such as in germ-band extension in

Drosophila where directed cell intercalation results in a reduction in the number of cells on the

anterior-posterior axis and an increase in the number of cells along the dorsal-ventral axis,

elongating the tissue along the dorsal-ventral axis (Irvine & Wieschaus, 1994). To test this, we

utilized live cell tracking during posterior lobe development. Initial observations of the outer face

of the posterior lobe revealed few cell rearrangement events. When cell rearrangements did

occur it was in response to a cell being removed from the apical surface (Figure 2 - video 1).

Due to the limited number of cell rearrangement events observed during posterior lobe

morphogenesis, cell intercalation does not appear to be a major driver of posterior lobe

morphogenesis, causing us to instead examine changes in cell shape.

Changes to cell shape are quite common during tissue morphogenesis, as classically illustrated

by the process of apical constriction that deforms tissues during many developmental processes

(Lecuit & Lenne, 2007; Martin & Goldstein, 2014). To examine cell shape, we utilized the

Raeppli system to label individual cells with a fluorescent marker (mTFP1) (Kanca et al., 2014).

We observed that cells within the posterior lobe are tall and thin, spanning from the basal to the

apical surface of the epithelium (Figure 2A). Because cells span the full thickness of this tissue,

we can approximate the height of the tallest cells in the posterior lobe by measuring tissue

thickness. For these measurements, we used the lateral plate as an in-sample comparison,

since it represents the tissue from which the posterior lobe protrudes and should differ from the

lobe in morphogenetic processes. We observed a pronounced increase in thickness of the

posterior lobe compared to the lateral plate (Figure 2B-C,F; Figure 2 - supplement 2). The

posterior lobe more than doubles in thickness with an average increase of 145.3% (+ 47.5µm),

while the lateral plate only increases by 22.6% (+ 7.9µm) overall. In contrast, when non-lobed

species are examined, no thickness changes are observed in the location where a posterior

lobe would form, indicating that this increase in tissue thickness is unique to the posterior lobe

(Figure 2B-E,G; Figure 2 - supplement 2). Interestingly, this increase in thickness is a dynamic

process during development. During the first 12 hours of posterior lobe development the lateral

plate thickness decreases by 5.1µm, but the posterior lobe increases in thickness by 16.5µm on

average (Figure 2F). By contrast, during the last 4 hours of development, rapid increases in

thickness occur in both the posterior lobe and lateral plate, which increase on average by

31.0µm and 14.6µm respectively (Figure 2F). These observations reveal a slow phase of cell

height increase during the first 12 hours of posterior lobe development, and fast phase during

the last four hours of posterior lobe development. Together this data suggests that the cells of

the posterior lobe undergo an extreme cell shape change to increase in length along their apico-

basal axis, driving the posterior lobe cells to project out of the plane of the lateral plate.

Cytoskeletal components increase in concentration in posterior lobe cells

Elongation of cells along their apico-basal axes appears to be a major contributor to posterior

lobe formation. To understand potential internal forces contributing to this cell shape change, we

examined the organization of cytoskeletal components. As expected for a polarized epithelium,

we found F-actin strongly localized to the apical cortex overlapping with E-cadherin throughout

the entire genitalia (Figure 3A). In contrast with the adjacent tissues, F-actin is also

concentrated along the apico-basal axis of posterior lobe cells (Figure 3A). This F-actin

localization was unique to the posterior lobe, as it is less intense in neighboring structures, such

as the lateral plate, clasper, and sheath, as well as in non-lobed species (Figure 3A; Figure 3 -

supplement 1). Next we evaluated microtubules by examining two post-translational

modifications that appear on tubulin, acetylation of α-tubulin on lysine40, a stabilizing

modification (Roll-Mecak, 2019; Xu et al., 2017), and tyrosinated tubulin, which has been

associated with rapid microtubule turnover (Roll-Mecak, 2019; Webster, Gundersen et al.,

1987). In the posterior lobe, acetylated tubulin levels are highest at the apex of the posterior

lobe and weaken towards the basal side of the lobe (Figure 3B-C). Compared to other

structures in the genitalia, acetylated tubulin is greatly increased specifically in the posterior lobe

(Figure 3B-C). In contrast, the levels of acetylated tubulin in non-lobed species are similar

throughout the genitalia (Figure 3 - supplement 1). We found tyrosinated tubulin has a more

consistent signal along the entire apico-basal axis in the posterior lobe (Figure 3B&D). The

amount of tyrosinated tubulin in posterior lobe cells is increased compared to neighboring

structures, but is weaker relative to the observed differences in acetylated tubulin. In non-lobed

species the levels of tyrosinated tubulin are consistent across the entire genitalia (Figure 3-

supplement 1). Collectively, these results suggest that changes in assembly and/or dynamics of

both F-actin and microtubule cytoskeletal networks could be contributing factors in changing the

shape of posterior lobe cells to increase its height along the apico-basal axis.

An apical extracellular matrix associates with posterior lobe cells

In addition to investigating cell autonomous mechanisms leading to increases in tissue

thickness, we also sought to identify sources of external forces which could play a role in

posterior lobe morphogenesis. Extrinsic roles for the basal and apical extracellular matrix have

been established in the pupal wing of D. melanogaster (Diaz-de-la-Loza et al., 2018; Etournay

et al., 2015; Ray et al., 2015). We first attempted to characterize the basal ECM by analyzing a

GFP-tagged version of Collagen IV (Viking:GFP). We observed that Viking:GFP, while present

at very early stages of genital morphogenesis, is weakly present during posterior lobe formation

across the entire genitalia (Figure 4 - supplement 1), suggesting that minimal basal ECM is

present at this time point. To further test for the presence of basal ECM, we examined another

basal ECM component, Perlecan (Perlecan:GFP), and also observed weak signal (Figure 4 –

supplement 1). Together, this data suggests that the basal ECM is globally decreased in the

genitalia during early pupal development, such that it is very weak during posterior lobe

morphogenesis.

We next sought to determine if an aECM is present, and if so, whether it could potentially

influence posterior lobe morphogenesis. A major component of the aECM is Dumpy, which is a

gigantic (2.5 MDa) zona pellucida domain-containing glycoprotein (Wilkin et al., 2000). We

examined a line in which Dumpy is endogenously tagged with a Yellow Fluorescent Protein

(Dumpy:YFP). Dumpy:YFP forms a complex three-dimensional network over the pupal genitalia

and is closely associated with cells of the posterior lobe (Figure 4; Figure 4 - video 1). At certain

points in the genitalia, this aECM network of Dumpy can extend up to 39.4 μm on average

above the cells, which is taller than the thickness of posterior lobe cells at the beginning of

development (Figure 4 - supplement 2). The intricate complex morphology of this aECM

network is hard to fully appreciate in flattened images due to its three-dimensional shape and

spatially varying levels of Dumpy:YFP, making it difficult to see weaker populations of Dumpy

without over-saturating more concentrated deposits.

In late pupal wing development, Dumpy anchors the wing to the surrounding cuticle, preventing

the tissue from retracting away from the cuticle, which is important to properly shape the wing

(Etournay et al., 2015; Ray et al., 2015). This same mechanism has been hypothesized to also

occur in the leg and antennae (Ray et al., 2015), however, in the posterior lobe we do not find

discrete anchorage points to the cuticle. Instead, we observed a large tether of Dumpy

emanating from the anal plate and connecting with the pupal cuticle membrane that encases the

entire pupa (Figure 4 - supplement 3, video 2) (Bainbridge & Bownes, 1981). This tether does

not come in direct contact with posterior lobe associated Dumpy or other nearby structures such

as the lateral plate, clasper, sheath, or phallus, suggesting that if Dumpy is contributing to

posterior lobe evolution and morphogenesis, it is likely through a mechanism which does not

depend on a direct mechanical linkage with the overlying pupal cuticle.

To investigate the role that Dumpy may play in posterior lobe morphogenesis, we examined its

localization throughout development. Prior to posterior lobe development, future cells of the lobe

lack apical Dumpy, and yet an intricate network associated with the clasper is observed (Figure

4A). However, from the early stages of posterior lobe development, as it first protrudes from the

lateral plate, we observe large deposits of Dumpy associated with future lobe cells (Figure 4B).

These deposits persist throughout most of its development (Figure 4C), becoming more

restricted to the apex of the posterior lobe towards the end of posterior lobe development

(Figure 4D). Throughout development, the posterior lobe associated Dumpy population is

connected to the complex network of Dumpy attached to more medial structures such as the

phallus (Figure 4 A2-D2), indicating that the posterior lobe is interconnected via the aECM with

nearby structures (Figure 4). In contrast to the posterior lobe, the lateral plate has minimal

Dumpy associated with it (Fig. 4 A1-D1). Only when we oversaturate the Dumpy:YFP signal can

we observe a weak population of Dumpy associated with the lateral plate (Figure 4 -

supplement 4). Together, this indicates that the cells of the posterior lobe and the lateral plate

substantially differ in the levels of associated Dumpy, suggesting a potential role in the

morphogenesis of the posterior lobe.

Expansion of Dumpy expression is correlated with the evolution of the posterior

lobe

The association of the posterior lobe with Dumpy suggests that changes in the expression of

dumpy may have been significant during the evolution of the posterior lobe. To test if posterior

lobe-associated Dumpy is a unique feature of species which produce a posterior lobe, we

compared the spatial distribution of its mRNA in *D. melanogaster* with *D. biarmipes,* a species

which lacks this structure. Early in pupal genital development at 32 hours APF we observe very

similar expression patterns of *dumpy* between *D. melanogaster* and *D. biarmipes*, with

expression at the base of the presumptive lateral plate-clasper (Figure 5A, Figure 5 -

supplement 1). From 36 to 40 hours APF, when the posterior lobe begins to develop, this

pattern becomes restricted to a small region at the base of the lateral plate and clasper, near

the anal plate in *D. biarmipes*, but is expanded in lobed species (Figure 5B, Figure 5 -

supplement 1). By 44 hours APF, expression of *dumpy* is reduced in the posterior lobe, as well

as in non-lobed species, with strongest expression associated with the clasper in *D. biarmipes*

(Figure 5A-B, Figure 5 - supplement 1). Overall, these results indicate that expression of *dumpy*

is expanded in a lobed species and correlates with the timing of the posterior lobe's formation.

In addition, considering that the developmental timing of *D. biarmipes* lags behind *D.*

melanogaster by approximately 4 hours (Figure 1 - supplement 1), this suggests that *dumpy*

expression becomes restricted during an earlier developmental period in the non-lobed species

D. biarmipes.

Although, it appears that the expression of *dumpy* has expanded in *D. melanogaster,* Dumpy is

an extracellular protein, and cells expressing its mRNA may not correlate with its ultimate

protein abundance or localization. Since an antibody for Dumpy is not available, we adapted

lectin staining protocols which can detect glycosylated proteins like Dumpy in order to compare

the distribution of aECM in species which lack posterior lobes. We found that fluorescein

conjugated *Vicia villosa* lectin (VVA), which labels *N*-acetylgalactosamine (Tian & Ten Hagen,

2007), approximately recapitulated Dumpy:YFP in *D. melanogaster.* VVA strongly associates

with the posterior lobe, shows trace association with the lateral plate, and roughly mirrors the

complex three-dimensional shape of the Dumpy aECM network covering the center of the

genitalia (Figure 5C)*.* When we examined VVA in the non-lobed species *D. biarmipes*, we

observed strong VVA signal over the center of the genitalia with weak connections to the tip of

the lateral plate, similar to what we observe in *D. melanogaster* (Figure 5 C-D)*.* In contrast, we

only found a weak strand-like structure emanating from the clasper and connecting to the

crevice between the lateral plate and clasper where the presumptive posterior lobe would form

(Figure 5D). These results correlate with our *in situ* results, where we observe high expression

at the center of the genitalia and weak expression of *dumpy* at the base between the clasper

and lateral plate in *D. biarmipes*, which may be responsible for forming the weak aECM

connection from the clasper to the crevice. Further, we observed similar staining patterns in an

additional non-lobed species, *D. ananassae* (Figure 5 - supplement 2). Collectively, these data

suggest that an ancestral aECM network was associated with the central genital structures,

including the phallus, sheath, and clasper, and a weak association in the crevice next to

prospective posterior lobe cells. During the course of evolution, expression of *dumpy* has

expanded to integrate cells of the posterior lobe, creating a prominent connection to the aECM

network.

Dumpy is required for proper posterior lobe formation

Thus far, we observed a strong association of the aECM with cells that form the posterior lobe,

a trait which is much less pronounced in non-lobed species. To determine if Dumpy plays a role

in posterior lobe formation, we next employed transgenic RNAi to knock down its expression.

Previous studies of *dumpy* characterized a VDRC RNAi line that is effective at reducing its

function (Ray et al., 2015). We used a driver from the *Pox neuro* gene (Boll & Noll, 2002) to

reduce *dumpy* levels in the posterior lobe. This resulted in a drastic decrease in the size and

alterations to the shape of the posterior lobe compared to a control RNAi (Figure 6). In *dumpy*

knockdown individuals, we observe a variable phenotype, and even within single individuals, the

severity of phenotype differs between left and right posterior lobes (Figure 6A; Figure 6 -

supplement 1). Knockdown was completed at both 25°C and 29°C, as higher temperatures

increase the efficacy of the Gal4/UAS system (Duffy, 2002). At higher temperatures, the *dumpy*

knockdown phenotype trended towards more severe defects (Figure 6B). Together, these

results suggest that posterior lobe development is sensitive to levels of *dumpy*, and that *dumpy*

plays a vital role in shaping the posterior lobe.

Correlation of Dumpy deposition and cell height in the posterior lobe

We next sought to determine when during development *dumpy* knockdown influences the

morphogenetic progression of the posterior lobe. This was important because we observed both

a slow and a fast phase of lobe development (Figure 1F), and also reasoned that posterior lobe

cells secrete cuticle once they have adopted their final adult conformations, of which any of

these phases could represent a critical Dumpy-dependent stage of development. We found that

dumpy knockdown individuals manifest phenotypes very early on (Figure 7A) and continue to

show abnormal lobe development through the end of its formation (Figure 7B). Interestingly,

differences in the height of cells on the ventral side of the posterior lobe are not observed

between control and *dumpy* knockdown treatments, instead defects in cell height are observed

in the more dorsally-localized cells of the posterior lobe (Figure 7A-B). This correlates with the

phenotypes of the adults in the *dumpy* knockdown in which the ventral tip is usually of normal

height with defects observed towards the dorsal side (Figure 6A). However, this phenotype

appears counterintuitive, as Dumpy protein normally associates along the entire posterior lobe,

so why does the tip of the posterior lobe develop to normal height when Dumpy is absent? To

better understand this phenotype, we examined Dumpy:YFP localization in the *dumpy*

knockdown background. We observed weak association of Dumpy with the tallest cells on the

ventral side of the posterior lobe both in early (Figure 7D n=5/5 samples) and late (Figure 7F

n=4/5 samples) stages compared to control animals. In contrast, no Dumpy was observed in

contact with the short cells on the dorsal side (Figure 7D & F). Together, this highlights a

correlation between the height of posterior lobe cells and presence of dumpy. One of our late

samples lacks a Dumpy connection to the ventral cells, correlating with our observation that not

all adult samples are fully extended on the ventral side (Figure 7 - supplement 1). This suggests

that ventral cell connections to the Dumpy aECM network may be lost late in development,

ultimately causing a shortening of these cells. In addition, we observed more severe phenotypes

of *dumpy* knockdown in the *dumpy-yfp* background (compared to the *dumpy*-WT background

alone), suggesting that Dumpy:YFP is a mild hypomorph (not shown). We also observed at

early time points highly variable strands of Dumpy in the middle of the lobe (between the ventral

and dorsal sides) (Figure 7 -supplement 2). These strands visually resembled the weak strands

of VVA observed in *D. biarmipes* (Figure 5D), in that they emanate from the clasper and

connect to the crevice between the posterior lobe and clasper. Overall, the most pronounced

phenotypic defects manifest in regions with the strongest reduction in Dumpy aECM deposition,

implying that Dumpy's presence is required for posterior lobe cells to elongate and project from

the lateral plate.

Discussion

Here, we determined how a morphological novelty forms at the cellular level, and in doing so,

revealed distinctive cell and aECM interactions underlying its development and evolution. We

identified how an extreme change in the shape of cells in the developing posterior lobe accounts

for its novel morphology. While intrinsic cytoskeletal components may contribute to this process,

our results highlight the critical role played by a vast extrinsic network of ECM on the apical side

of the epithelium. It was unexpected that such an elaborate supercellular matrix structure would

participate in the evolution of a seemingly simple novelty. Below, we consider the potential roles

played by the aECM in posterior lobe development and diversification, and discuss how studies

of morphogenesis can illuminate the simple origins of structures that might otherwise seem

impossibly complex to evolve.

Mechanisms for aECM-mediated control of cell height in the posterior lobe

Our work demonstrates an important role for the aECM protein, Dumpy, in the growth of the

posterior lobe, as exhibited by the dramatic phenotypes in the *dumpy* RNAi background and the

strong association of Dumpy:YFP with only the tallest cells in these experiments. Our data is

consistent with three possible mechanisms. First, Dumpy could serve as a structural support

while autonomous cell mechanical processes drive apico-basal elongation. Second, the cells of

the posterior lobe could be pulled mechanically through their connection to the Dumpy aECM.

This process could operate passively, deforming cells of the lobe, but could also drive changes

in the cytoskeleton in response to external tensions. Finally, the aECM could play a direct role

by altering cell signaling dynamics, as has been exhibited by the basal ECM (Kirkpatrick et al.,

2004; Kreuger et al., 2004; Wang et al., 2008). Previous research has shown that the JAK/STAT

pathway is important for posterior lobe development (Glassford et al., 2015), and their ability to

signal to the correct cells could be altered in the absence of Dumpy. Of course, these models

are not mutually exclusive and some combination of these mechanisms may be integrated to

shape the posterior lobe. Our observations of increased cytoskeletal components in posterior

lobe cells and the reduced height of cells that lack Dumpy in our knockdown experiments are

consistent with all three mechanisms, which are difficult to differentiate experimentally. When

we examine morphogenesis in non-lobed species, we observed that the lateral plate drops

below the clasper (Figure 1 - supplement 1). Assuming this ancestral process still occurs in

lobed species, it is quite possible that the aECM 'holds' cells of the posterior lobe during the

early stages of posterior lobe development while the lateral plate is pulled down, causing cells of

the posterior lobe to elongate to relieve the stress. Future manipulative biomechanical studies

will be required to explore these possibilities.

The role of aECM in the diversification of genital structures

Genitalia represent some of the most rapidly diversifying structures in the animal kingdom, and

our results suggest the aECM may participate in the modification of *Drosophila* genital

structures. The shape of the posterior lobe is extremely diverse among species of the

melanogaster clade (Coyne, 1993). Our results demonstrate that reducing the levels of Dumpy

can affect the shape of the posterior lobe, with extreme knockdown phenotypes approximating

the posterior lobe of *D. mauritiana*. Furthermore, the clasper and phallus show dense deposits

of Dumpy, suggesting that the aECM could play important roles in diversifying these remarkably

variable structures. During the course of evolution, one could imagine that by altering which

cells are connected to the aECM, the strength of those connections, or the forces acting on

those connections could lead to changes in morphological shape. Hence identifying causative

genes that differentiate these structures could uncover novel mechanisms for genetically

controlling the behavior of this aECM and behaviors of cells bound to this dynamic scaffold.

Integrating cells into a pre-existing aECM network to generate morphological

novelty

In comparing the morphogenesis of a novel structure to close relatives which lack it

(representing a proxy for the ancestral state), we identified a likely path by which the aECM

became associated with the posterior lobe. The aECM, while understudied, has been implicated
in the morphogenesis of many structures (Diaz-de-la-Loza et al., 2018; Etournay et al., 2015;
Ray et al., 2015; Fernandes et al., 2010; Dong et al., 2014; Heiman et al., 2009; Low et al.,
2019), and yet, its role during the evolution of novel structures is largely unexplored. We find a
conserved aECM network associated with central genital structures (clasper, sheath, and
phallus) in both lobed and non-lobed species. In non-lobed species, *dumpy* is expressed weakly
at the base between the lateral plate and clasper resulting in a thin connection of aECM from
clasper to the crevice (Figure 8). By contrast, lobed species express high levels of *dumpy*
between the presumptive posterior lobe and clasper, resulting in large amounts of aECM in the
crevice. We hypothesize that this increase in aECM allows cells at the base of the lateral plate
to be integrated into this ancestral aECM network (Figure 8), a step which was likely significant
to the evolution of the posterior lobe. Overall, this suggests that the aECM could be an
unexpected target for generating novel anatomical structures.
422
The expanded *dumpy* expression we observed caused us to consider how the posterior lobe
gained this aECM attachment. Interestingly, our previous work found a gene regulatory network
(GRN) that regulates development of an ancestral embryonic structure, the posterior spiracles,
which was co-opted during the evolution of the posterior lobe and regulates its development
(Glassford et al., 2015). Previous work has shown that *dumpy* is expressed in the posterior
spiracles (Wilkin et al., 2000), and we have observed a thin tether of Dumpy:YFP connecting the
posterior spiracles to the surrounding embryonic cuticle (Figure 8 - Supplement 1). This is
consistent with previously identified roles for Dumpy in epithelia-cuticle attachment in the wing
(Etournay et al., 2015; Ray et al., 2015) and hypothesized role in the muscle, leg, and antenna
(Wilkin et al., 2000; Ray et al., 2015). Identification of regulatory elements which activate *dumpy*
in the posterior lobe will be necessary to determine whether its role in the posterior spiracle was
relevant to the evolution of expanded genital expression.
435
Evolution is thought to act through the path of least resistance. When confronted with the
remarkable diversity of genital morphologies present in insects, one must wonder how the
intricate projections, bumps, and divots form in its underlying epithelia. Models of co-option have
been appealing because they establish pre-existing mechanisms in place that can be rapidly
ported to new locations to generate massive changes in a tissue. Our examination of the cellular
processes during posterior lobe morphogenesis highlights a different way that co-option may
work. Here, the aECM mechanism we uncovered appears to be a path of least resistance

because this tissue already uses a vast network of aECM to potentially pattern other structures,

such as the phallus and its multiple elaborations (Rice et al., 2019; Peluffo, et al. 2015;

Kamimura, 2010). Because this network of aECM represents a pre-existing condition, it is easy

to appreciate how cells of the posterior lobe could evolve novel extracellular connections to this

network to generate a new protrusion. On the other hand, tissues which lack such an ancestral

network may well be less likely to evolve projections through this mechanism. While the aECM

is required for this morphogenetic process, we envision that additional networks and processes

must be contributing to the full morphogenesis of the posterior lobe. Determining genetic

changes which underlie such remarkable cellular responses represents a major looming

challenge in evo-devo research (Smith et al., 2018).

Materials and Methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional Information
Antibody	rat anti-alpha tubulin (tyrosinated)	MilliporeSigma	Millipore Cat# MAB1864-I	IHC (1:500)
Antibody	mouse anti-alpha tubulin (acetylated)	Sigma-Aldrich	Sigma-Aldrich Cat# T6793, RRID:AB_477585	IHC (1:500)
Antibody	rat anti-Ecadherin	DSHB	DSHB Cat# DCAD2, RRID:AB_528120	IHC (1:500)
Antibody	mouse anti-fasciclin III	DSHB	DSHB Cat# 7G10 anti-Fasciclin III, RRID:AB_528238	IHC (1:500)
Antibody	rabbit anti-histone H3 (phospho S10)	Abcam	Abcam Cat# ab5176, RRID:AB_304763	IHC (1:50)
Antibody	goat anti-GFP	Abcam	Abcam Cat# ab6662, RRID:AB_305635	IHC (1:300)

Antibody	fluorescein Vicia Villosa Lectin (VVA)	Vector Laboratories	Vector Laboratories Cat# FL-1231, RRID:AB_2336856	IHC (1:200)
Chemical compound, drug	rhodamine phalloidin	Thermo Fisher Scientific	Thermo Fisher Scientific Cat# R415, RRID:AB_2572408	IHC (1:200)
Strain, strain background (*Drosophila melanogaster*)	$y^1 w^1$ *Drosophila melanogaster*	Bloomington Drosophila Stock Center	BDSC Cat# 1495, RRID:BDSC_1495	
Strain, strain background (*Drosophila biarmipes*)	wild type	National Drosophila Species Stock Center (NDSSC)	NDSSC Stock #: 14023-0361.10 RRID:FlyBase_FBst0203870	
Strain, strain background (*Drosophila ananassae*)	wild type	National Drosophila Species Stock Center (NDSSC)	NDSSC Stock #: 14024-0371.13 RRID:FlyBase_FBst0201380	No longer available
Strain, strain background (*Drosophila pseudoobscura*)	wild type	National Drosophila Species Stock Center (NDSSC)	NDSSC Stock #: 14011-0121.87 RRID:FlyBase_FBst0200074	No longer available
Genetic reagent (*Drosophila melanogaster*)	*UAS*-Raeppli-CAAX	Bloomington Drosophila Stock Center (BDSC)	BDSC Cat# 55084, RRID:BDSC_55084	
Genetic reagent (*Drosophila melanogaster*)	*pox neuro-Gal4*	(Boll & Noll, 2002)	Construct #13	
Genetic reagent (*Drosophila melanogaster*)	*D. simulans pox neuro*-Gal4	This paper	N/A	
Genetic reagent (*Drosophila melanogaster*)	hs – flippase[122]	Gift from Erika A. Bach	Flybase: FBtp0001101	
Genetic reagent (*Drosophila melanogaster*)	*armadillo-GFP*	Bloomington Drosophila stock center	BDSC Cat# 8556, RRID:BDSC_8556	
Genetic reagent (*Drosophila*	Dumpy:YFP	Drosophila Genomics and	DGGR Cat# 115238,	

melanogaster)		Genetic Resources	RRID:DGGR_115238	
Genetic reagent (*Drosophila melanogaster*)	E-cadherin:mCherry	Bloomington Drosophila stock center	BDSC Cat# 59014, RRID:BDSC_59014	
Genetic reagent (*Drosophila melanogaster*)	*UAS*-dumpyRNAi	Vienna Drosophila Resource Center	VDRC Cat#44029, RRID:FlyBase_FBst0465370	
Genetic reagent (*Drosophila melanogaster*)	*UAS*-mCherryRNAi	Bloomington Drosophila stock center	BDSC Cat# 35785, RRID:BDSC_35785	
Recombinant DNA reagent	pS3aG4	Gift from Benjamin Prud'homme	N/A	Gal4 vector used to make *D. simulans pox neuro* gal4 line
Sequence-based reagent	GCCACTAACAATCCATGCGGTT	This paper	N/A	*dumpy* probe forward primer
Sequence-based reagent	TAATACGACTCACTATAGGGAGAAATAGCCCTGTCCTTGGAATCC	This paper	N/A	*dumpy* probe reverse primer with T7 primer
Sequence-based reagent	TTCCGGGCGCGCCTCGGTGGCTTAACACGCGCATT	This paper	N/A	*D. simulans pox neuro* forward primer for gal 4 line
Sequence-based reagent	TTGCCCCTGCAGGATCGCTGATTCCATGGCCCAGT	This paper	N/A	*D. simulans pox neuro* reverse primer for gal 4 line
Software algorithm	Fiji (ImageJ v2.0)	(Schindelin et al., 2012)	RRID:SCR_002285	
Software algorithm	GenePalette	(Rebeiz & Posakony, 2004; Smith et al., 2017)	N/A	
Software algorithm	Leica Application Suite X	Leica	RRID:SCR_013673)	
Software algorithm	Microsoft Excel	Microsoft	RRID:SCR_016137	
Software algorithm	MorphoGraphX	(Barbier de Reuille et al., 2015)	N/A	

Software algorithm	Prism 8	GraphPad	N/A	

Fly stocks and genetics

Fly stocks were reared using standard culture conditions. Wild type species used in this study were obtained from the University of California, San Diego *Drosophila* Stock Center (now known as The National Drosophila Species Stock Center at Cornell University)(*Drosophila biarmipes* #14024-0361.10, *Drosophila ananassae* #14024-0371.13, *Drosophila pseudoobscura* #14011-0121.87) and from the Bloomington Drosophila Stock Center (*Drosophila melanogaster* [y^1w^1] #1495). *pox neuro*-Gal4 (construct #13) was obtained from Werner Boll (Boll & Noll, 2002). The following were obtained from the Bloomington Drosophila stock center: *UAS*-Raeppli-CAAX (#55084), *armadillo-GFP* (#8556), Ecadherin:mCherry (#59014), and *UAS*-mCherryRNAi (control for RNAi experiments, as mCherry is not present in the *Drosophila* genome)(35785). *UAS*-dumpyRNAi was obtained from the Vienna Drosophila Resource Center (#44029) and Dumpy:YFP was obtained from the Drosophila Genomics and Genetic Resources (#115238).

For the Raeppli experiments, stable lines of hs-flippase;;*UAS*-Raeppli-CAAX/*UAS*-Raeppli-CAAX and *D. simulans pox neuro*-gal4/*D. simulans pox neuro*-gal4;*UAS*-Raeppli-CAAX/*UAS*-Raeppli-CAAX were generated. *D. simulans pox neuro*-gal4 was used as opposed to *pox neuro*-gal4 because a gal4 driver on the second chromosome was required. Virgin females from the first line were crossed to males from the second line to ensure hs-flippase was inherited by all offspring. Offspring were collected and grown as normal, heat shocked at 37°C for 1 hour around 24 to 28 hours APF, and allowed to finish development at 25°C.

Sample Preparation

Pupal samples were prepared following the protocol in Glassford, et al., 2015. Briefly, samples were incubated at 25°C unless otherwise noted. Dissections were performed in cold PBS, pupae were cut in half, removed from their pupal cases, and fat bodies removed by flushing. Larval samples were dissected in cold PBS by cutting the larva in half, and flipping the posterior end of the larva inside out. All samples were fixed for 30 minutes at room temperature in PBS with 0.1% Triton-X and 4% paraformaldehyde. Samples stained with phalloidin had Triton-X concentrations increased to 0.3%. Samples used for VVA staining were removed from pupal cuticle before being fixed in PBS with 0.1% Triton-x, 4% paraformaldehyde, and 1% trichloroacetic acid on ice for 1 hour followed by 30 minutes at room temperature. The

trichloroacetic acid method causes some slight tissue distortion, as the precipitation treatment

utilized to refine the VVA signal causes the posterior lobe to become slightly deformed and

curve in towards the clasper. However, similar defects were not observed in the other structures

such as the lateral plate or in *D. biarmipes*. Samples were stored in PBT for immunostaining at

4°C for up to two days. For *in situ* hybridization, samples were rinsed twice in methanol and

rinsed twice in ethanol. Samples were stored at -20°C in ethanol.

Immunostaining and *in situ* hybridization

For immunostaining, genital samples were removed from the surrounding pupal cuticle and

incubated overnight at 4°C with primary antibodies diluted in PBS with 0.1% Triton-X (PBT).

VVA and phalloidin samples were placed on a rocker. The following primary antibodies were

used: rat anti-alpha tubulin (tyrosinated) 1:500 (MAB 1864-I, MilliporeSigma), mouse anti-alpha

tubulin (acetylated) 1:500 (T6793, Sigma-Aldrich), rat anti-Ecadherin 1:500 (DCAD2, DSHB),

mouse anti-fasciclin III 1:500 (7G10, DSHB), rabbit anti-histone H3 (phospho S10) 1:50

(ab5176, Abcam), goat anti-GFP 1:300 (ab6662, Abcam), fluorescein Vicia Villosa Lectin (VVA)

1:200 (FL-1231, Vector Laboratories). The goat anti-GFP was used to increase signal of

Dumpy:YFP in the knockdown experiments only. Primary antibody was removed by performing

two quick rinses and two long washes (at least 5 minutes) in PBT. Samples were incubated

overnight at 4°C in secondary antibodies diluted in PBT. The following secondary antibodies

were used: donkey anti-rat Alexa 594 1:500 (A21209, Invitrogen), donkey anti-mouse Alexa 488

1:500 (A21202, Thermo Fisher Scientific), donkey anti-rat Alexa 488 1:500 (A21208, Thermo

Fisher Scientific), goat anti-mouse Alexa 594 1:500 (A-11005, Thermo Fisher Scientific), goat

anti-rabbit Alexa 594 1:500 (A-11012, Thermo Fisher Scientific), donkey anti-goat Cy2 1:500

(705-225-147, Jackson ImmunoResearch). Rhodamine phalloidin (R415, Thermo Fisher

Scientific) stain was performed with secondary antibody. Samples were washed out of

secondary antibody by performing two quick rinses and two long washes (at least 5 minutes) in

PBT. Samples were then incubated in 50% PBT/50% glycerol solution for at least 5 minutes.

Pupal samples were mounted on glass slides coated with Poly-L-Lysine Solution. Glass slides

had 1 to 2 layers of double side tape with a well cut out in which the sample was placed and

covered with a cover slip

in situ hybridization was performed following the protocol in Rebeiz et al., 2009 with

modifications to perform *in situs* in the InsituPro VSi robot (Intavis Bioanalytical Instruments) as

done by Glassford et al., 2015.

Microscopy and live imaging

Cuticles of adult posterior lobes and in situ hybridization samples were imaged on Leica
DM2000 with a 40x objective for cuticles and a 10x objective for in situ samples. Samples with
fluorescent antibodies and fluorescently tagged proteins were imaged using a Leica TCS SP5
Confocal microscope using either a 40x or 63x oil immersion objective.

To live image genital development, a 2% agar solution was poured into a small petri dish filling
the dish half way. A 0.1-10µL pipette tip was used to make small wells in the agar for pupal
samples. Timed pupal samples were inserted head first into the small well and a 5-300µL
pipette tip was used to push sample into agar by placing the tip around the posterior spiracles
on the pupal case. To better image the developing genitalia the pupal case at the posterior end
was removed with forceps. Deionized water was used to cover the samples and imaged on a
Leica TCS SP5 Confocal microscope using a 63x water objective.

To live image embryos, Dumpy:YFP flies were grown in egg-laying chamber with grape agar
plates (Genesee Scientific). Embryos were removed from plates using forceps and rolled on a
piece of double sided tape to remove the chorion. Embryos then were positioned on a glass
coverslip coated with embryo glue. A glass slide was covered with double sided tape and a well
was made and filled with halocarbon 27 oil. The cover slip with the embryos was then placed on
the glass slide, submerging the embryos in halocarbon oil. Embryos were imaged on a Leica
TCS SP8 confocal with a 63x oil objective.

Image analysis

Images were processed with Fiji (Schindelin et al., 2012) and Photoshop. Three-dimensional
views were completed in MorphoGraphX (Barbier de Reuille et al., 2015) or Leica Application
Suite X. Movies were processed in Fiji and cell rearrangements were tracked using the manual
tracking plugin. Tissue thickness/cell height during development was measured in cross-section
view by drawing a line centered between the two sides (based on apical membrane) of the lobe
until the basal side was reached. Area of adult posterior lobe cuticles and height of the adult
lobe were measured by using the lateral plate as a guide for determining the bottom boundary
of the posterior lobe. To prevent any possible bias for one lobe vs the other (i.e. left vs right)
which lobe was used in statistical analysis was randomly decided, except for Figure 6 -
supplement 1 where both sides of the posterior lobe were considered.

Transgenic Constructs

To make the *D. simulans pox neuro*-gal4 driver, the posterior lobe enhancer for *pox neuro* in *D.*
simulans, identified in Glassford et al., 2015, was cloned using primers listed in key resources
table using genomic DNA purified with the DNeasy Blood and Tissue Kit (QIAGEN). Primers
were designed using sequence conservation with the GenePalette software tool (Rebeiz and
Posakony 2004; Smith et al., 2017). The cloned sequence was inserted into the pS3aG4 (Gal4)
using *AscI* and *SbfI* restriction sites. The final construct was inserted into the 51D landing site
on the second chromosome (Bischof et al., 2007).

Acknowledgements

571
The authors thank the members of the M.R. laboratory for comments and discussion on the
manuscript. We thank Werner Boll and Markus Noll for the *pox neuro*-gal4 line, the Bloomington
stock center, VDRC, and DGGR stock centers for fly stocks, Benjamin Prud'homme for the
s3aG4 vector, Erika A, Bach for the hs-flippase line, and Winslow Johnson for the *D. simulans*
pox neuro-gal4 line. This work was supported by the National Institutes of Health (GM107387 to
M.R. and HD044750 to L.A.D.).

Author details

580
Sarah Jacquelyn Smith
Department of Biological Sciences, University of Pittsburgh
Contributions: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data
curation, Writing-original draft, Writing-review and editing, Visualization
Competing interests: No competing interests declared
ORCID: 0000-0002-1469-1821

587
Lance A. Davidson
Department of Bioengineering University of Pittsburgh
Contributions: Conceptualization, Methodology, Writing-review and editing, Funding acquisition
Competing interests: No competing interests declared

ORCID: 0000-0002-2956-0437

Mark Rebeiz

Department of Biological Sciences, University of Pittsburgh

Contributions: Conceptualization, Methodology, Writing-review and editing, Supervision,

Funding acquisition

Competing interests: No competing interests declared

ORCID: 0000-0001-5731-5570

References

Atallah, J., Liu, N.H., Dennis, P., Hon, A., & Larsen, E.W. (2009). Developmental constraints

and convergent evolution in Drosophila sex comb formation. *Evolution & Development*,

11(2), 205–218. https://doi.org/10.1111/j.1525-142X.2009.00320.x

Bainbridge, S.P., & Bownes, M. (1981). Staging the metamorphosis of Drosophila

melanogaster. *Journal of Embryology and Experimental Morphology*, *66*, 57–80.

Barbier de Reuille, P., Routier-Kierzkowska, A.L., Kierzkowski, D., Bassel, G.W., Schüpbach,

T., Tauriello, G., Bajpai, N., Strauss, S., Weber, A., Kiss, A., Burian, A., Hofhuis, H., Sapala,

A., Lipowczan, M., Heimlicher, M.B., Robinson, S., Bayer, E.M., Basler, K., Koumoutsakos,

P., Roeder, A.H., Aegerter-Wilmsen, T., Nakayama, N., Tsiantis, M., Hay, A., Kwiatkowska,

D., Xenarios, I., Kuhlemeier, C., Smith, R.S. (2015). MorphoGraphX: A platform for

quantifying morphogenesis in 4D. *eLife*, *4*, 05864. https://doi.org/10.7554/eLife.05864

Bischof, J., Maeda, R.K., Hediger, M., Karch, F., & Basler, K. (2007). An optimized transgenesis

system for Drosophila using germ-line-specific phiC31 integrases. *Proceedings of the

National Academy of Sciences of the United States of America*, *104*(9), 3312–3317.

Boll, W., & Noll, M. (2002). The Drosophila Pox neuro gene: control of male courtship behavior

and fertility as revealed by a complete dissection of all enhancers. *Development*, *129*(24),

5667–5681. https://doi.org/10.1242/dev.00157

Brown, N.H. (2011). Extracellular matrix in development: insights from mechanisms conserved

between invertebrates and vertebrates. *Cold Spring Harbor Perspectives in Biology*, *3*(12),

a005082. https://doi.org/10.1101/cshperspect.a005082

Coyne, J.A. (1993). The genetics of an isolating mechanism between two sibling species of

Drosophila. *Evolution, 47*(3), 778-788.

Daley, W.P., & Yamada, K.M. (2013). ECM-modulated cellular dynamics as a driving force for

tissue morphogenesis. *Current Opinion in Genetics & Development*, *23*(4), 408–414.

https://doi.org/10.1016/j.gde.2013.05.005

Diaz-de-la-Loza, M.D., Ray, R.P., Ganguly, P.S., Alt, S., Davis, J.R., Hoppe, A., Tapon, N.,

Salbreux, G., & Thompson, B.J. (2018). Apical and basal matrix remodeling control epithelial

morphogenesis. *Developmental Cell*, *46*(1), 23-39.e5.

https://doi.org/10.1016/j.devcel.2018.06.006

Dong, B., Hannezo, E., & Hayashi, S. (2014). Balance between apical membrane growth and

luminal matrix resistance determines epithelial tubule shape. *Cell Reports*, *7*(4), 941–950.

https://doi.org/10.1016/j.celrep.2014.03.066

Duffy, J.B. (2002). GAL4 system in Drosophila: a fly geneticist's Swiss army knife. *Genesis*,

34(1–2), 1–15. https://doi.org/10.1002/gene.10150

Eberhard, W.G. (1985). *Sexual selection and animal genitalia*. Harvard University Press.

Etournay, R., Popović, M., Merkel, M., Nandi, A., Blasse, C., Aigouy, B., Brandl, H., Myers, G.,

Salbreux, G., Jülicher, F., Eaton, S. (2015). Interplay of cell dynamics and epithelial tension

during morphogenesis of the Drosophila pupal wing. *eLife*, *4*, e07090.

https://doi.org/10.7554/eLife.07090

Fernandes, I., Chanut-Delalande, H., Ferrer, P., Latapie, Y., Waltzer, L., Affolter, M., Payre, F.,

& Plaza, S. (2010). Zona pellucida domain proteins remodel the apical compartment for

localized cell shape changes. *Developmental Cell*, *18*(1), 64–76.

https://doi.org/10.1016/j.devcel.2009.11.009

Frazee, S.R., & Masly, J.P. (2015). Multiple sexual selection pressures drive the rapid evolution

of complex morphology in a male secondary genital structure. *Ecology and Evolution*, *5*(19),

4437–4450. https://doi.org/10.1002/ece3.1721

Glassford, W.J., Johnson, W.C., Dall, N.R., Smith, S.J., Liu, Y., Boll, W., Noll, M., & Rebeiz, M.

(2015). Co-option of an Ancestral Hox-Regulated Network Underlies a Recently Evolved

Morphological Novelty. *Developmental Cell*, *34*(5), 520–531.

https://doi.org/10.1016/j.devcel.2015.08.005

Green, J.E., Cavey, M., Médina Caturegli, E., Aigouy, B., Gompel, N., & Prud'homme, B.

(2019). Evolution of ovipositor length in drosophila suzukii is driven by enhanced cell size

expansion and anisotropic tissue reorganization. *Current Biology, 29*(12), 2075-2082.

https://doi.org/10.1016/j.cub.2019.05.020

Guirao, B., & Bellaïche, Y. (2017). Biomechanics of cell rearrangements in Drosophila. *Current

Opinion in Cell Biology*, *48*, 113–124. https://doi.org/10.1016/j.ceb.2017.06.004

Heiman, M.G., & Shaham, S. (2009). DEX-1 and DYF-7 establish sensory dendrite length by

anchoring dendritic tips during cell migration. *Cell, 137*(2), 344-355.

https://doi.org/10.1016/j.cell.2009.01.057

Heisenberg, C.P., & Bellaïche, Y. (2013). Forces in tissue morphogenesis and patterning. *Cell*,

153(5), 948–962. https://doi.org/10.1016/j.cell.2013.05.008

Irvine, K.D., & Wieschaus, E. (1994). Cell intercalation during Drosophila germband extension

and its regulation by pair-rule segmentation genes. *Development*, *120*(4), 827–841.

https://doi.org/10.1007/s00435-010-0109-5

Jagadeeshan, S., & Singh, R.S. (2006). A time-sequence functional analysis of mating

behaviour and genital coupling in Drosophila: role of cryptic female choice and male sex-

drive in the evolution of male genitalia. *Journal of Evolutionary Biology*, *19*(4), 1058–1070.

https://doi.org/10.1111/j.1420-9101.2006.01099.x

Kamimura, Y. (2010). Copulation anatomy of Drosophila melanogaster (Diptera: Drosophilidae):

wound-making organs and their possible roles. *Zoomorphology, 129*(3), 163-174.

Kanca, O., Caussinus, E., Denes, A.S., Percival-Smith, A., & Affolter, M. (2014). Raeppli: a

whole-tissue labeling tool for live imaging of Drosophila development. *Development*, *141*(2),

472–480. https://doi.org/10.1242/dev.102913

Kirkpatrick, C.A., Dimitroff, B.D., Rawson, J.M., & Selleck, S.B. (2004). Spatial regulation of

Wingless morphogen distribution and signaling by Dally-like protein. *Developmental Cell*,

7(4), 513–523. https://doi.org/10.1016/j.devcel.2004.08.004

Kopp, A., & True, J.R. (2002). Evolution of male sexual characters in the oriental Drosophila

melanogaster species group. *Evolution & Development*, *4*(4), 278–291.

https://doi.org/10.1046/j.1525-142X.2002.02017.x

Kreuger, J., Perez, L., Giraldez, A.J., & Cohen, S.M. (2004). Opposing activities of Dally-like

glypican at high and low levels of Wingless morphogen activity. *Developmental Cell*, *7*(4),

503–512. https://doi.org/10.1016/j.devcel.2004.08.005

Lecuit, T., & Lenne, P.F. (2007). Cell surface mechanics and the control of cell shape, tissue

patterns and morphogenesis. *Nature Reviews. Molecular Cell Biology*, *8*(8), 633–644.

https://doi.org/10.1038/nrm2222

LeVasseur-Viens, H., Polak, M., & Moehring, A.J. (2015). No evidence for external genital

morphology affecting cryptic female choice and reproductive isolation in Drosophila.

Evolution, *69*(7), 1797–1807. https://doi.org/10.1111/evo.12685

Linde-Medina, M., & Marcucio, R. (2018). Living tissues are more than cell clusters: The

extracellular matrix as a driving force in morphogenesis. *Progress in Biophysics and*

Molecular Biology, *137*, 46–51. https://doi.org/10.1016/j.pbiomolbio.2018.01.009

Li, L., Tang, Q., Nakamura, T., Suh, J.G., Ohshima, H., & Jung, H.S. (2016). Fine tuning of

Rac1 and RhoA alters cuspal shapes by remolding the cellular geometry. *Scientific Reports*,

6, 37828. https://doi.org/10.1038/srep37828

Liu, Y., Ramos-Womack. M., Han, C., Reilly, P., Brackett, K.L., Rogers, W., Williams, T.M.,

Andolfatto, P., Stern, D.L. and Rebeiz, M. (2019). Changes throughout a Genetic network

mask the contribution of hox gene evolution. *Current Biology 29*(13), 2157-2166.36.

https://doi.org/10.1016/j.cub.2019.05.074

Loganathan, R., Rongish, B.J., Smith, C.M., Filla, M.B., Czirok, A., Bénazéraf, B., & Little, C.D.

(2016). Extracellular matrix motion and early morphogenesis. *Development*, *143*(12), 2056–

2065. https://doi.org/10.1242/dev.127886

Low, I.I.C., Williams, C.R., Chong, M.K., McLachlan, I.G., Wierbowski, B.M., Kolotuev, I., &

Heiman, M.G. (2019). Morphogenesis of neurons and glia within an epithelium.

Development, 146, dev171124. https://doi.org/10.1242/dev.171124

Martin, A.C., & Goldstein, B. (2014). Apical constriction: themes and variations on a cellular

mechanism driving morphogenesis. *Development*, *141*(10), 1987–1998.

https://doi.org/10.1242/dev.102228

Moczek, A.P. (2008). On the origins of novelty in development and evolution. *BioEssays*, *30*(5),

432–447. https://doi.org/10.1002/bies.20754

Osterfield, M., Schüpbach, T., Wieschaus, E., & Shvartsman, S.Y. (2015). Diversity of epithelial

morphogenesis during eggshell formation in drosophilids. *Development*, *142*(11), 1971–

1977. https://doi.org/10.1242/dev.119404

Paluch, E., & Heisenberg, C.P. (2009). Biology and physics of cell shape changes in

development. *Current Biology*, *19*(17), R790-799. https://doi.org/10.1016/j.cub.2009.07.029

Peluffo, A.E., Nuez, I., Debat, V., Savisaar, R., Stern, D.L., & Orgogozo, V. A Major Locus

Controls a Genital Shape Difference Involved in Reproductive Isolation Between Drosophila

yakuba and Drosophila santomea. *G3: Genes, Genomes, Genetics, 5*(12), 2893-2901.

https://doi.org/10.1534/g3.115.023481

Peter, I.S., & Davidson, E.H. (2015). *Genomic Control Process: Development and Evolution*.

Academic Press.

Ray, R.P., Matamoro-Vidal, A., Ribeiro, P.S., Tapon, N., Houle, D., Salazar-Ciudad, I., &

Thompson, B.J. (2015). Patterned anchorage to the apical extracellular matrix defines tissue

shape in the developing appendages of drosophila. *Developmental Cell*, *34*(3), 310–322.

https://doi.org/10.1016/j.devcel.2015.06.019

Rebeiz, M., Patel, N.H., & Hinman, V.F. (2015). Unraveling the tangled skein: the evolution of

transcriptional regulatory networks in development. *Annual Review of Genomics and Human*

Genetics, *16*, 103–131. https://doi.org/10.1146/annurev-genom-091212-153423

Rebeiz, M., Pool, J.E., Kassner, V.A., Aquadro, C.F., & Carroll, S.B. (2009). Stepwise

modification of a modular enhancer underlies adaptation in a Drosophila population.

Science, *326*(5960), 1663–1667. https://doi.org/10.1126/science.1178357

Rebeiz, M., & Posakony, J.W. (2004). GenePalette: a universal software tool for genome

sequence visualization and analysis. *Developmental Biology*, *271*(2), 431–438.

https://doi.org/10.1016/j.ydbio.2004.04.011

Renvoisé, E., Kavanagh, K.D., Lazzari, V., Häkkinen, T.J., Rice, R., Pantalacci, S., Salazar-

Ciudad, I., Jernvall, J. (2017). Mechanical constraint from growing jaw facilitates mammalian

dental diversity. *Proceedings of the National Academy of Sciences of the United States of*

America, *114*(35), 9403–9408. https://doi.org/10.1073/pnas.1707410114

Rice, G., David, J., Kamimura, Y., Masly, J., Mcgregor, A., Nagy, O., Noselli, S., Nunes, M.D.S.,

O'Grady, P., Sánchez-Herrero, E., Siegal, M., Toda, M., Rebeiz, M., Courtier-Orgogozo, V.,

& Yassin, A. (2019). A Standardized Nomenclature and Atlas of the Male Terminalia of

Drosophila melanogaster. *Preprints,* 2019060071.

https://doi.org/10.20944/preprints201906.0071.v1

Roll-Mecak, A. (2019). How cells exploit tubulin diversity to build functional cellular microtubule

mosaics. *Current Opinion in Cell Biology*, *56*, 102–108.

https://doi.org/10.1016/j.ceb.2018.10.009

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,

S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V., Eliceiri,

K., Tomancak, P., Cardona, A. (2012). Fiji: an open-source platform for biological-image

analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schnatwinkel, C., & Niswander, L. (2013). Multiparametric image analysis of lung-branching

morphogenesis. *Developmental Dynamics*, *242*(6), 622–637.

https://doi.org/10.1002/dvdy.23961

Smith, A.F., Posakony, J.W., & Rebeiz, M. (2017). Automated tools for comparative sequence

analysis of genic regions using the GenePalette application. *Developmental Biology, 429*(1),

158-164. https://doi.org/10.1016/j.ydbio.2017.06.033

Smith, S.J., Rebeiz, M., & Davidson, L. (2018). From pattern to process: studies at the interface

of gene regulatory networks, morphogenesis, and evolution. *Current Opinion in Genetics &*

Development, *51*, 103–110. https://doi.org/10.1016/j.gde.2018.08.004

Tada, M., & Heisenberg, C.P. (2012). Convergent extension: using collective cell migration and

cell intercalation to shape embryos. *Development*, *139*(21), 3897–3904.

https://doi.org/10.1242/dev.073007

Tanaka, K., Barmina, O., & Kopp, A. (2009). Distinct developmental mechanisms underlie the

evolutionary diversification of Drosophila sex combs. *Proceedings of the National Academy*

of Sciences of the United States of America, *106*(12), 4764–4769.

https://doi.org/10.1073/pnas.0807875106

Tian, E., & Ten Hagen, K.G. (2007). O-linked glycan expression during Drosophila

development. *Glycobiology*, *17*(8), 820–827. https://doi.org/10.1093/glycob/cwm056

Wagner, G.P. (2014). *Homology, Genes, and Evolutionary Innovation*. Princeton, NJ: Princeton

University Press.

Wagner, G.P. & Lynch, V.J. (2010). Evolutionary novelties. *Current Biology*, *20*(2), R48-52.

https://doi.org/10.1016/j.cub.2009.11.010

Walck-Shannon, E., & Hardin, J. (2014). Cell intercalation from top to bottom. *Nature Reviews.*

Molecular Cell Biology, *15*(1), 34–48. https://doi.org/10.1038/nrm3723

Wang, X., Harris, R. E., Bayston, L. J., & Ashe, H. L. (2008). Type IV collagens regulate BMP

signalling in Drosophila. *Nature*, *455*(7209), 72–77. https://doi.org/10.1038/nature07214

Webster, D.R., Gundersen, G.G., Bulinski, J.C., & Borisy, G.G. (1987). Differential turnover of

tyrosinated and detyrosinated microtubules. *Proceedings of the National Academy of*

Sciences of the United States of America, *84*(24), 9040–9044.

https://doi.org/10.1073/pnas.84.24.9040

Wilkin, M. B., Becker, M. N., Mulvey, D., Phan, I., Chao, A., Cooper, K., Chung H.J., Campbell,

I.D., Baron, M., MacIntyre, R. (2000). Drosophila dumpy is a gigantic extracellular protein

required to maintain tension at epidermal-cuticle attachment sites. *Current Biology*, *10*(10),

559–567. https://doi.org/10.1016/S0960-9822(00)00482-6

Xu, Z., Schaedel, L., Portran, D., Aguilar, A., Gaillard, J., Marinkovich, M.P. Théry, M., &

Nachury, M.V. (2017). Microtubules acquire resistance from mechanical breakage through

intralumenal acetylation. *Science*, *356*(6335), 328–332.

https://doi.org/10.1126/science.aai8764

Figure 1. The posterior lobe a novel protrusion from the lateral plate of *D. melanogaster*.

(A) Phylogenetic tree with representative bright-field images of adult cuticle of the lateral plate

and posterior lobe (arrow). (B-E) Illustration, (B'-E') maximum projection, and (B''-E'') three-

dimensional projection of early (28 hours APF) and late (52 hours APF) developing genitalia

showing the posterior lobe projecting form the lateral plate of *D. melanogaster* (D''), but absent

in *D. biarmipes* (E''). Relevant structures are labeled: posterior lobe (PL), lateral plate (LP),

clasper (C), sheath (S), phallus (P), anal plate (AP), and hypandrium (H). All max projections

are oriented with ventral side towards to top and dorsal sides towards the bottom. (F) Zoomed in

illustration of posterior lobe and (G) a cross-sectional/lateral view of the posterior lobe. The

highest point of the lobe is the apex and the invagination between the lobe and the clasper is

termed the crevice (G). Scale bar, 20μm.

Figure 1 – supplement 1

Supplement 1 - figure 1 . Developmental timing of lobed vs non-lobed genitalia.

Developmental time course of the lobed species *D. melanogaster* (A-D) and the non-lobed

species *D. biarmipes* (E-H) with E-cadherin label. Location of respective cross sections

indicated in yellow for lateral plate and blue for posterior lobe (*D. melanogaster)* or equivalent

location in non-lobed species (*D. biarmipes*). Relevant structures are labeled: posterior lobe

(PL), lateral plate (LP), clasper (C), sheath (S), and phallus (P). Scale bar, 20µm. At 28 hours

APF the genitalia looks relatively similar between *D. melanogaster* (A-A2) and *D. biarmipes* (E-

E2). At 32 hours APF in *D. melanogaster* the clasper and lateral plate have fully begun to cleave

(B1-2 red arrowhead=cleavage), the lateral plate is lower than the clasper (B1), and the

hypandrium, sheath, and phallus have fully everted and are neighboring the clasper and lateral

plate (B1-2). *D. biarmipes* lags behind approximately 4 hours. At 32 hours APF there is slight

cleavage near the dorsal side of the lateral plate and clasper (F2 red arrowhead), but no

cleavage has occurred at the ventral side (F1). In addition, the sheath, hypandrium, and phallus

have not everted yet (F1-2). At 36 hours APF in *D. biarmipes,* cleavage has begun along the full

length of the lateral plate and clasper (G1-2 red arrowhead), the lateral plate is lower than the

clasper (G1-2), and the hypandrium, sheath, and phallus have everted and are next to the

lateral plate and clasper (G1-2). As development proceeds later at 52 hours APF the lateral

plate and clasper fully separate at the ventral side of the genitalia in both *D. melanogaster* (D1

green arrow) and *D. biarmipes* (H1 green arrow). Full cleavage does not span the length of the

lateral plate and clasper (D2 and H2) and stops right before the posterior lobe forms (D2) and

also stops before reaching the very dorsal side of the lateral plate and clasper in *D. biarmipes*

(H2).

See supporting file.

Supplement 1 - video 1. The posterior lobe protrudes from the lateral plate.

Three-dimensional projections of *D. biarmipes* (left) and *D. melanogaster* (right) samples at 52

hours APF labeled with E-cadherin.

Figure 2. Posterior lobe cells increase in height to project out from the lateral plate.

(A) A single cell in the posterior lobe labeled with Raeppli-mTFP1 (green) spans the height of the tissue labeled with lateral membrane marker fasciclin III (Fas3, magenta). Apical side of posterior lobe identified with dotted line. Sample is 44h after pupal formation (APF), but was heat shocked for 1 hour at 24h APF causing it to develop faster and more closely resembles a 48h APF sample. Scale bar, 10µm. n=4 (B-E) Maximum projections of early (36h APF) and late (52h APF) genital samples labeled with Fas3 (lateral membranes, green) and E-Cadherin (apical membranes, magenta). Location of respective cross sections indicated in yellow for lateral plate (B1-E1) and blue for posterior lobe (*D. melanogaster)* (B2-C2) or equivalent location in non-lobed species (*D. biarmipes*) (D2-E2). Scale bar, 20µm. (F) Quantification of tissue thickness of the lateral plate (light blue) and posterior lobe (dark blue). Illustration represents approximate location of cross-section that was used for tissue height measurement. Individual data points a presented; n=10 per each time point. (G) Quantification of tissue thickness of the posterior lobe in *D. melanogaster* (dark blue) and equivalent location in non-lobed species *D. biarmipes* (orange). Illustration represents approximate location of cross-section that was used for tissue thickness measurement. Individual data points presented; n≥9 per each time point. Statistical significance is indicated (unpaired t-test; ****p≤0.0001; n.s.=not significant p≥0.05). *D. melanogaster* tissue height measures in (G) are replotted from (F) to facilitate direct comparisons with *D. biarmipes.*

 Figure 2 – supplement 1

Figure 2 - supplement 1. Cell division dynamics do not differ between lobed and non-

lobed species.

Developmental time course with Phospho-Histone H3 (Ser 10) (PH3; green) labeling actively dividing

cells and Ecad (magenta) labeling the apical membrane of the tissue. Only superficial slices are shown to

avoid fat body signals beneath lateral plate and clasper. n ≥ 3 per each time point. Scale bar, 20µm. In

both *D. melanogaster* and *D. biarmipes* cell division is widespread at 24 hours APF (A & D). Cell division

is decreased by 32 hours APF (B & E). By 40 hours APF no cell division is occurring (C & F).

 Figure 2 – supplement 2

Figure 2 - supplement 2. Extended time course of tissue thickness in lobed and non-

lobed species.

Extended time course for samples quantified in Figure 2F-G. (A-C) Max and cross-section view of 44

hours APF (A & C) and 48 hours APF (B) genital samples with lateral membrane labeled with Fas3

(green) and apical membrane labeled with Ecad (magenta). Location of respective cross sections

indicated in yellow for lateral plate (A1-C1) and blue for posterior lobe (*D. melanogaster)* or equivalent

location in non-lobed species (*D. biarmipes*) (A2-C2). n≥ 9 per experiment. Scale bar, 20μm.

See supporting file.

Figure 2 - video 1. Minor cell rearrangement during posterior lobe development.

Live imaging of posterior lobe development with GFP tagged armadillo (apical membrane

marker) illustrating a cell dropping from the apical surface and a neighboring cell filling in the

gap. Imaging starts at approximately 36 hours APF. Due to uncontrolled temperatures during

imaging that were cooler than normal growing conditions, the posterior lobe develops slower

and the time indicated is not comparable to other images in the manuscript which were all

grown under controlled settings. Based on the thickness of the posterior lobe at the end of the

movie the posterior lobe is between 48 to 52 hours APF. Cells were tracked manually and

indicated with colored dots. Some dots disappear towards the end of the movie as they become

difficult to track due to the signal from cells on the medial side of the posterior lobe.

Figure 3. Cytoskeletal components are concentrated in posterior lobe cells.

(A-D) Maximum projection, and respective cross-sections of late (48h APF) genital samples of

the lobed species *D. melanogaster* labeled with F-actin/phalloidin and Ecad (A), acetylated

tubulin (B,C), and tyrosinated tubulin (B,D). Location of respective cross sections indicated in

yellow for lateral plate (A1-D1) and blue for posterior lobe (A2-D2). Cross-sections are

maximum projections of a restricted 5.434µm thick section to provide a complete view of

cytoskeletal components along the apico-basal axis. All cross-sections are oriented with apical

side at the top and basal side at the bottom. Asterisk identifies bristles which have high levels of

F-actin and tubulin. Bright basal signal in A1 and A2 are fat bodies. Bottom layers were

removed in panel A to remove fat body signal which overwhelmed other details. (B-D2) Panels

C and D show separate channels of panel B. Relevant structures labeled: Posterior lobe (PL),

lateral plate (LP), clasper (C), and sheath (S). Scale bar, 20µm. n≥ 3 per experiment.

Figure 3 – supplement 1

Figure 3 - supplement 1. Uniform distribution of cytoskeletal components in non-lobed

species.

(A-D) Max projections of late (48h APF) genital samples of non-lobed species *D. biarmipes*

labeled with F-actin/phalloidin and Ecad (A), acetylated tubulin (B,C), and tyrosinated tubulin

(B,D). Location of respective cross sections indicated in blue for presumptive posterior lobe cells

(A1-D1). Cross-sections are maximum projection of a restricted 5.434μm thick section to display

the full view of the cytoskeleton along the apico-basal axis. All cross-sections are oriented with

apical side at the top and basal side at the bottom. Asterisk identifies bristles which have high

levels of F-actin and tubulin. Bright basal signal in A1 are fat bodies. Bottom layers were

removed in panel A to avoid fat body signal which masked other details. Panels C and D show

separate channels of panel B. Relevant structures labeled: Lateral plate (LP), clasper (C), and

sheath (S) labeled. Scale bar, 20μm. n≥ 3 per experiment.

Figure 4. Dumpy deposition is correlated with posterior lobe development.

(A-D) Maximum projection and (A'-B'') respective zoom, indicated with pink box, labeled with

Dumpy:YFP (green) and Ecad (magenta) for each time point. Location of respective cross

sections indicated in yellow for lateral plate (A1-D1) and blue for posterior lobe (A2-D2).

Arrowhead in (A2) indicates future posterior lobe cells. Cross-sections are oriented with apical

side at the top and basal side at the bottom. Relevant structures labeled: Posterior lobe (PL),

lateral plate (LP), clasper (C), sheath (S), and phallus (P). Scale bar, 20μm. n≥ 4 per

experiment. Images were independently brightened to show relevant structures.

 Figure 4 – supplement 1

Figure 4 - supplement 1. Limited basal ECM present during posterior lobe

morphogenesis.

Basal ECM markers Collagen IV (Viking:GFP; green)(A & C) and Perlecan (Perlecan:GFP;

green) (B & D) in L3 larval genital disc (A & B) and in 44 hours APF genitalia (C & D). Image

settings were the same for each marker between larval and pupal samples. Sporadic dots

observed are fat bodies (white arrows in cross section), which fill the basal lumen of the pupal

genital epithelium. Location of respective cross sections indicated in white. Cross-sections for

larval samples are oriented basal sides out, as the disc has not yet everted. Pupal samples are

oriented with apical side at the top and basal side at the bottom. Higher amounts of basal ECM

are observed in larvae compared to 44 hour APF genital samples. Relevant structures labeled:

Posterior lobe (PL) and clasper (C). Scale bar, 20µm.

 Figure 4 – supplement 2

Figure 4 - supplement 2. Dumpy extends above the apical surface of the phallus.

(A) Projection of Dumpy:YFP (green) and Ecad:mCherry (magenta) imaged live at 48 hours

APF. Location of respective cross sections indicated in orange. (A1) Cross section showing

extent of Dumpy:YFP observed above the surface of the genitalia. Relevant structures labeled:

Posterior lobe (PL), lateral plate (LP), clasper (C), sheath (S), and phallus (P). Scale bar, 20µm.

n=3.

 Figure 4 – supplement 3

Figure 4 - supplement 3. A tether of Dumpy connects the genitalia to the pupal cuticle

membrane that encases the developing pupa.

(A-B) Live imaging of Dumpy:YFP (green) and Ecad:mCherry (magenta) at respective time

points. Location of respective cross sections indicated in orange. (A1-B1) Cross-sections are

max projection of a 4.94μm (A1) and 1.73μm (B1) thick section to show full tether (arrow) and

its connection to the cuticle (arrowhead) and anal plate. All cross-sections are oriented with

apical side at the top and basal side at the bottom. Relevant structures labeled: Posterior lobe

(PL), lateral plate (LP), clasper (C), sheath (S), phallus (P), and anal plate (AP). Scale bar,

20μm. n=1 per each time point.

 Figure 4 – supplement 4

Figure 4 - supplement 4. The Dumpy aECM network extends weak connection to lateral

plate.

(A-D) Brightened images of respective cross sections from Figure 2 of lateral plate (A1-D1) in

yellow and posterior lobe in blue (A2-D2). Cross-sections are oriented with apical side at the top

and basal side at the bottom. Relevant structure labeled: Posterior lobe (PL), lateral plate (LP),

clasper (C), sheath (S), and phallus (P). Scale bar, 20μm. n≥ 4 per experiment. Images were

overexposed to show relevant structures.

See supporting file.

Figure 4 - video 1. Three-dimensional structure of the genital Dumpy aECM network.

A 52 hour APF genital sample with Dumpy:YFP (green) and E-cadherin (magenta) is shown.

Part 1 of the movie shows a cross-sectional view starting at the ventral side of the posterior lobe

and moving towards the dorsal side of the posterior lobe and part 2 shows the same view but

starting at the ventral tip of the lateral plate and moving towards the ventral side of the posterior

lobe. In the upper-right corner there is a guide that roughly depicts the running location of the

cross section. Cross-sections are oriented with apical side at the top and basal side at the

bottom. Relevant structures labeled: Posterior lobe (PL), lateral plate (LP), clasper (C), sheath

(S), and phallus (P).

See supporting file.

Figure 4 – video 2. A tether of Dumpy connects the genitalia to the surrounding cuticle.

3D rotation of Dumpy:YFP (green) and Ecad:mCherry (magenta) imaged live at 44 hours APF.

Relevant structures labeled: Posterior lobe (PL), lateral plate (LP), clasper (C), sheath (S),

phallus (P), and anal plate (AP).

Figure 5. aECM is spatially expanded in lobed species compared to non-lobed species.

(A-B) *in situ* hybridization for *dumpy* mRNA in the lobed species *D. melanogaster* (A) and the

non-lobed species *D. biarmipes* (B). Pink box outlines location of zoom in for A1 and B1.

Posterior lobe associated expression highlighted with arrow (purple/white) for strong expression,

asterisk for weak expression, and arrowhead for clasper-specific expression. Expression

observed in *D. melanogaster* at 44 hours APF is not present in all samples (see Figure 5 –

supplement 1). (C-D) aECM is labeled with *Vicia villosa* lectin (VVA; green) and apical

membrane labeled with Ecad (magenta) at 44 hours APF in *D. melanogaster* (C) and *D.*

biarmipes (D). Location of respective cross sections indicated in yellow for lateral plate (C2-D2)

and blue for posterior lobe in *D. melanogaster* (C1) and corresponding position in *D. biarmipes*

(D1). All cross-sections are oriented with apical side at the top and basal side at the bottom.

White arrows highlight the crevice localization between the lateral plate and clasper, which the

aECM fills in *D. melanogaster* (C1), but only a weakly stained strand-like structure of aECM

appears in *D. biarmipes* (D1). Tendrils of aECM can also be observed connecting to the lateral

plate in both species (red arrowheads). Relevant structures labeled: Posterior lobe (PL), lateral

plate (LP), clasper (C), sheath (S), and phallus (P). Scale bar, 20µm. n=at least 5 per

experiment.

Figure 5 – supplement 1

Figure 5 - supplement 1. *dumpy* **expression is spatially expanded in lobed species**

compared to non-lobed species.

(A-B) Additional *in situ* hybridization samples for *dumpy* mRNA in lobed species *D.*

melanogaster (A) and non-lobed species *D. biarmipes* (B) to show full range of expression

observed in experiment. Samples without outlines on one side are due to the tissue being

damaged on that side. Green circle in first image highlights relevant location at the base of the

lateral plate, but not included in the remaining images to leave images unobstructed. Asterisk

indicates the expression is deep in the sample and not expressed in lateral plate or clasper

cells. n= 4 per experiment.

1003

 Figure 5 – supplement 2

Figure 5 - supplement 2. aECM not expanded in non-lobed species *D. ananassae*

(A-C) aECM labeled with VVA (green) and apical membrane labeled with Ecad (magenta) at 40

hours APF in non-lobed species *D. ananassae*. Location of respective cross-sections indicated

in blue. Top cross-section displayed with normal brightness to show details and bottom cross-

section has been brightened to show where all populations of aECM are located. All cross-

sections are oriented with apical side at the top and basal side at the bottom. White arrow

highlights the 'crevice' between the lateral plate and clasper, which is not pronounced at 40

hours APF in *D. ananassae*. Relevant structures labeled: lateral plate (LP) and clasper (C)

labeled. Scale bar, 20µm. n=at least 2 per experiment.

 Figure 6

Figure 6. Dumpy is required for proper posterior lobe shape.

(A) Range of adult posterior lobe phenotypes produced by control (*mCherry* RNAi) and *dumpy*

RNAi animals. Phenotypic classes defined from wild type (I) to most severe (V). Scale bar,

20µm. (B) Percentage of posterior lobes in each class for control, *dumpy* RNAi at 25°C, and

dumpy RNAi at 29°C. (C) Quantification of area of adult posterior lobes of *mCherry* RNAi

(control) and *dumpy* RNAi at 25°C and 29°C. Statistical significance between each temperature

indicated (unpaired t-test; ****p≤0.0001).

 Figure 6 – supplement

Figure 6 - supplement 1. Increased left-right variability of posterior lobe phenotype upon

***dumpy* knockdown.**

(A) Comparison of *dumpy* knockdown (purple circles) and control knockdown (green squares) of

left and right adult posterior lobes in single individuals grown at 29°C measuring height at the

ventral side of the posterior lobe (single individual represented as a single dot or square). Black

line represents perfect correlation in height. *dumpy* knockdown individuals stray more form

perfect correlation, indicating that the height of the posterior lobe varies more in the *dumpy*

knockdown. (B) Percentage of *dumpy* knockdown individuals plotted in (A) in which both

posterior lobes were classified as the same phenotype or different phenotypes (defined in

Figure 6).

 Figure 7

Figure 7. Correlation between the deposition of Dumpy and knockdown phenotype.

(A-B) Comparison of *mCherry* RNAi (control) and *dumpy* RNAi at 44 hours APF (A) and 52 hours APF (B). Images are rotated in 3D to visualize the full shape of the posterior lobe labeled with E-cadherin. Quantification of tissue height at the ventral tip (dark blue) and dorsal base (light blue) of the posterior lobe. Cartoon represents relative location of cross-section used for tissue thickness measurement. Individual data points presented; n=at least 10 per each time point. The ventral tip is defined as the location where the posterior lobe is at its max height. The base was determined by moving 19.76µm dorsally from the ventral tip. Statistical significance for each time point indicated (unpaired t-test; ***p≤0.001; n.s.=not significant p≥0.05). (C-F)

Comparison of *mCherry* RNAi (control) (C & E) and *dumpy* RNAi (D & F) at 44 hours APF and 52 hours APF with Dumpy:YFP (Green) and Ecad (Magenta). GFP antibody was used to increase YFP signal. All cross-sections are oriented with apical side at the top and basal side at the bottom. Relevant structures labeled: Lateral plate (LP) posterior lobe (PL), and clasper (C). Cross-sections are max projections of 5.434µm sections to show full Dumpy connection. Images were independently brightened to show relevant structures. Scale bar, 20µm. n=at least 5 per experiment.

 Figure 7 – supplement 1

1054   **Figure 7 - supplement 1. Variability in height of the adult posterior lobe in *dumpy***

1055   **knockdown.**

1056   Comparison of *mCherry* RNAi (control) and *dumpy* RNAi adults. Quantification of height of

1057   cuticle at the ventral side of the posterior lobe. (unpaired t-test; ***p≤0.001; ****p≤0.0001; n≥28).

1058

Figure 7 – supplement 2

Figure 7 - supplement 2. Strands of Dumpy in *dumpy* knockdown.

(A & B) *dumpy* RNAi at 44 hours APF with Dumpy:YFP showing strands of Dumpy connecting to the

crevice between the lateral plate and clasper (arrow). Relevant structures labeled: Lateral plate (LP)

posterior lobe (PL), and clasper (C). Cross-sections are max projection of 5.434µm section to show full

Dumpy connection. Scale bar, 20µm.

Figure 8. Expansion of apical extracellular matrix underlies the morphogenesis of a
recently evolved structure.

(Top) Illustration of non-lobed species, *D. biarmipes*, with ancestral aECM network covering

central genital structures (2B) including the clasper (C), sheath, and phallus. Weak connections

of aECM span from the clasper to the lateral plate (LP) during early development (1 & 2A - top).

(Bottom) Illustration of lobed species, *D. melanogaster*. The aECM network has expanded to fill

the crevice between the lateral plate and clasper (1-bottom) integrating these cells into the

ancestral aECM network (2-bottom). This aECM population is needed for cells to properly

project from the lateral plate, forming the posterior lobe.

 Figure 8 – supplement 1

1077   **Figure 8 - supplement 1. Dumpy anchors posterior spiracles to surrounding cuticle.**

1078   Live imaging of Dumpy:YFP in the embryonic posterior spiracles. Posterior spiracle (dotted line)

1079   is connected to the cuticle (arrowhead) via a tether of dumpy (arrow). Scale bar, 20µm.