Sexual selection does not increase the rate of compensatory adaptation to a

mutation influencing a secondary sexual trait in *Drosophila melanogaster*

Christopher H. Chandler[1,2], Anna Mammel[1,3], Ian Dworkin[1,4]

[1]Department of Integrative Biology, BEACON Center for the Study of Evolution in

Action, Michigan State University, East Lansing, MI, United States

[2]Department of Biological Sciences, State University of New York at Oswego, Oswego,

NY, United States

[3]Basic Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA, USA

[4]Department of Biology, McMaster University, Hamilton, Ontario, Canada

16 Data and scripts to reproduce analyses can be found at:

17 https://github.com/DworkinLab/Chandler_etal_G3_2020

Abstract

Theoretical work predicts that sexual selection can enhance natural selection, increasing the rate of adaptation to new environments and helping purge harmful mutations. While some experiments support these predictions, remarkably little work has addressed the role of sexual selection on compensatory adaptation—populations' ability to compensate for the costs of deleterious alleles that are already present. We tested whether sexual selection, as well as the degree of standing genetic variation, affect the rate of compensatory evolution via phenotypic suppression in experimental populations of *Drosophila melanogaster*. These populations were fixed for a spontaneous mutation causing mild abnormalities in the male sex comb, a structure important for mating success. We fine-mapped this mutation to an ~85 kb region on the X chromosome containing three candidate genes, showed that the mutation is deleterious, and that its phenotypic expression and penetrance vary by genetic background. We then performed experimental evolution, including a treatment where opportunity for mate choice was limited by experimentally enforced monogamy. Although evolved populations did show some phenotypic suppression of the morphological abnormalities in the sex comb, the amount of suppression did not depend on the opportunity for sexual selection. Sexual selection, therefore, may not always enhance natural selection; instead, the interaction between these two forces may depend on additional factors.

Introduction

Sexual selection has important impacts on many aspects of how organisms evolve, including on speciation rates and the degree of sexual dimorphism (e.g., Masta and Maddison 2002; Ellis and Oakley 2016). It was once thought that sexual selection may act independently or even antagonistically to other components of natural selection (e.g. viability and fecundity). However, sexual selection might also be concordant with, and represent a substantial portion of, the total selection on an allele. If so, sexual selection on males might also influence the overall mutation load or rate of adaptation, including in females. For instance, sexual selection may influence how organisms respond to selective pressures in the short term, influencing how quickly populations adapt to novel environments, in particular when the population begins at a distance from an optimum (Long *et al.* 2012). Additionally, some models predict that sexual selection should help populations filter out harmful mutations more rapidly than selection on other fitness components (viability and fecundity selection) alone (Agrawal 2001). This prediction is based partly on the observation that sexual displays are often correlated with overall condition. Any mutation that reduces an organism's nonsexual fitness is therefore also likely to affect its mating success (Rowe and Houle 1996; Chandler *et al.* 2013b) or even its success in sperm competition (Clark *et al.* 2012). In those cases, total selection against such mutations is stronger than it would be without sexual selection.

Empirical support for this scenario has been mixed. In some studies testing these predictions, evidence supported a role for sexual selection in purging deleterious mutations or accelerating adaptation (Radwan 2004; Sharp and Agrawal 2008; Hollis *et*

al. 2009; Jarzebowska and Radwan 2010; McGuigan *et al.* 2011; Long *et al.* 2012;

Almbro and Simmons 2014; Lumley *et al.* 2015; Grieshop *et al.* 2016; Jacomb *et al.*

2016). However, a handful of studies also contradict these, perhaps because of the

confounding effects of sexual conflict (Hollis and Houle 2011; Arbuthnott and Rundle

2012; Chenoweth *et al.* 2015), or because they used large-effect mutations or strong

mutagens not representative of natural variation (Plesnar *et al.* 2011; Cabral and

Holland 2014).

70   Of course, deleterious mutations are not always purged by selection; they can

increase in frequency and occasionally become fixed via drift, hitchhiking, or

antagonistic pleiotropy, especially if their effects on fitness are only mildly deleterious

(and in populations with a small effective population size). In those cases where the

deleterious alleles are difficult for selection to purge, alleles at other loci that

compensate epistatically for the fitness costs of these fixed deleterious alleles may be

favored by selection. There is evidence of compensatory adaptation in both microbial

and multicellular organisms. For instance, alleles conferring antibiotic resistance are

sometimes costly in the absence of antibiotics, but compensatory mutations can reduce

those costs (Reynolds 2000; Maisnier-Patin *et al.* 2002; Comas *et al.* 2012). In the

blowfly, diazinon resistance via alleles at the *Rop-1* gene had negative pleiotropic

effects, increasing fluctuating asymmetry, but these effects were ultimately

compensated by modifiers (McKenzie and Clarke 1988; Davies *et al.* 1996).

Additionally, sex chromosome dosage compensation could also be considered a form of

compensatory adaptation, having likely evolved in response to loss-of-function

mutations on Y or W chromosomes (Charlesworth 1978). In addition, the phenotypic

expression (penetrance and expressivity) of many mutations can be strongly influenced

by genetic background (e.g., Chandler *et al.* 2013a, 2017; Mullis *et al.* 2018; Hou *et al.*

2019). Thus, selection favoring suppressor alleles at other loci may also contribute to

compensatory adaptation by limiting the phenotypic expression of a deleterious

mutation.

Although sexual selection has received a lot of attention as a possible influence

on the rate of purging of deleterious mutations, the role of sexual selection in

compensatory evolution remains largely unexplored. Nevertheless, we might similarly

predict that sexual selection can also accelerate compensatory adaptation, especially if

sexual displays are condition dependent. In one study (Pischedda and Chippindale

2005), the *nub^1* mutation, which drastically reduces the size of the wing, resulting in an

inhibition of males' ability to generate courtship songs, was fixed in experimental

populations of *Drosophila melanogaster*. As predicted, this mutation had greater fitness

costs in males than it did in females, but males also showed greater compensatory

fitness recovery over 180 generations (albeit without compensating for the effects on

wing morphology directly; A. Chippindale, personal communication), providing some

support that sexual selection may enhance compensatory adaptation. However, this

study was not replicated (only a single lineage), since the *nub^1* populations were

originally generated for other purposes. Clearly, more study is needed on whether

sexual selection can speed up compensatory adaptation.

In this study, we address the question of whether sexual selection can impact the

rate of compensatory evolution (via phenotypic suppression) using experimental

evolution in *Drosophila melanogaster*. We chose a mutation in the *sex combs distal*

gene (scd^1) (Boube *et al.* 1997; Randsholt and Santamaria 2008), a spontaneous,

partially penetrant mutation affecting the development of the male sex comb, a structure

critical for male mating success (Ng and Kopp 2008) and rapidly evolving across

Drosophila species (Atallah *et al.* 2009, 2012; Kopp 2011; Malagón *et al.* 2014). First,

we further mapped the mutation and characterized its effects across different wild type

genetic backgrounds; because we found abundant genetic variation in natural

populations modifying its penetrance and expressivity, we next focused on

compensatory adaptation via phenotypic suppression in experimentally evolved

populations. Despite a general compensatory response, we observed no evidence that

sexual selection influenced the rate of compensatory evolution via phenotypic

suppression of the sex comb phenotypes.

Methods

Mapping scd^1—

sex combs distal^1 (scd^1) is a spontaneous X-linked allele resulting in ectopic sex comb

bristles on the second tarsal segment of the prothoracic leg in males (Boube *et al.*

1997). In the base stock (Bloomington Drosophila Stock Center strain #5070, y^1 scd^1

ras^1 v^1 f^1), it has incomplete penetrance, with only about 70% of males showing the

ectopic sex comb bristles (Figure 1), and no visible phenotype in homozygous or

heterozygous females. In our populations, males also sometimes exhibited minor

defects in the primary sex comb, such as a gap or partially untransformed bristles.

The identity of the gene and molecular lesion of this allele are unknown, although

some previous recombination mapping suggested it was near 1-30.6, and that a local

duplication of the 8C-9B region of the X chromosome onto the Y (*DP(1:Y)FF*), could

partially rescue the phenotype of *scd¹* (Santamaria and Randsholt 1995; Randsholt and

Santamaria 2008). We attempted to further fine map the gene through duplication

mapping. Virgin female flies of strain BDSC 5070 (*y¹ scd¹ ras¹ v¹ f¹*) were crossed to

males of strains carrying duplicated segments of the X chromosome translocated onto

the Y chromosome or chromosome III (Table 1; Cook *et al.* 2010; Venken *et al.* 2010),

and the male offspring were scored for the presence of mutant phenotypes, such as the

ectopic sex comb or disruptions in the primary sex comb. Assuming *scd¹* is a recessive

loss-of-function allele, if the duplicated segment contains a functional wild-type copy of

the *scd* gene, then the mutant phenotype would be rescued and no male offspring from

these crosses would show sex comb defects. Because *scd¹* is only partially penetrant,

we scored at least 20 male offspring from each cross.

Influence of genetic background on penetrance and expressivity—

To determine the extent of genetic variation for the penetrance and expressivity

of *scd¹*, we crossed virgin female *y¹ scd¹ ras¹ v¹ f¹* flies to males of a randomly chosen

subset of Drosophila Genetic Reference Panel lines (Mackay *et al.* 2012). When the

adult F1 offspring eclosed, we fixed specimens in 70% ethanol, and then mounted male

prothoracic legs in 70% glycerol and scored them for the presence of ectopic sex combs

on the second tarsal segment, abnormalities in the primary sex comb, and primary sex

comb tooth number. These crosses only examine each wild-type genetic background in

a heterozygous state, and thus it is expected that this will underestimate the actual

degree of background dependence, as recessive effects of alleles in each background

will not be captured.

To test for an effect of genetic background on penetrance, we fit a logistic model

testing for the effect of genotype on presence of an ectopic sex comb using glm() in

base R version 3.6.1. We also confirmed those results using glmer() in the lme4

package version 1.1-21.

Introgression of scd^1—

To generate populations for experimental evolution, we introgressed the *scd^1*

mutation into FVW, a domesticated lab-maintained population founded from flies

collected from Fenn Valley, MI in 2010. The FVW population was maintained in

population cages with 10 bottles for egg-laying each generation for approximately 10

generations prior to beginning introgressions (Chari *et al.* 2017).

To begin the introgression (Supplementary Figure 1A), virgin females of the 5070

progenitor strain (with the genotype *y^1 scd^1 ras^1 v^1 f^1*) were crossed to FVW males; this

stock strain carries visible X-linked genetic markers (*y^1* causes yellow body

pigmentation, *ras^1* and *v^1* affect eye color, and *f^1* produces forked bristles). The

heterozygous F1 females were then backcrossed to FVW males. From the F2 offspring,

we selected males showing the *scd^1* phenotype, which were thus hemizygous for *scd^1*,

but with phenotypically wild-type eyes and normal bristles to eliminate the *ras^1*, *v^1*, and

f^1 mutations, which are all located to the right of *scd^1* on the X chromosome. We then

crossed these males to virgin FVW females, to obtain female offspring heterozygous for

scd^1 in a partial FVW background. We then crossed these females to FVW males, and

selected males with ectopic sex combs, but not yellow bodies, to eliminate the *y^1*

mutation to the left of *scd^1*. Each male from these crosses thus carries an independently

derived X chromosome with *scd^1* in a random FVW background.

We obtained eight such males and crossed them to FVW females. From these

crosses, we obtained virgin females heterozygous for scd^1 in an FVW background. The

first of these virgin scd^1/scd^+ females to emerge were crossed with FVW males to obtain

more scd^1 males with a mostly FVW background. The later-emerging scd^1/scd^+ females

were kept isolated at cooler temperatures (18˚C) until the scd^1 males from the previous

cross emerged. We then crossed the scd^1/scd^+ females to the scd^1 males. Finally, we

set up sib matings among the offspring of these crosses, using only males hemizygous

for scd^1 (showing a sex comb phenotype) and females of unknown genotype (either

scd^1/scd^+ or scd^1/scd^1). Of those crosses, we kept those in which the mother was

inferred to be homozygous for scd^1, in which nearly all male progeny displayed scd^1

phenotypes. This allowed us to establish a homozygous scd^1 line with an FVW genetic

background, which we designated as scd^*. scd^* carried at least four independently

derived X chromosomes with scd^1 in an FVW background (Supplementary Figure 1A).

To introduce further genetic diversity (from the FVW population) into scd^*, scd^*

males were crossed to wild-type FVW females to obtain heterozygous scd^1 females with

additional genetic material from the FVW background. These females were then

backcrossed to FVW males. Five replicate backcrosses were set up in culture bottles,

each with 25-30 FVW males and 25-30 females, heterozygous for scd^1 and for alleles

from the FVW background. We then selected males with the scd^1 phenotype, and

backcrossed them to virgin scd^* females, in six replicate bottles each containing 20-25

males and 20-25 females, to maintain scd^1 while introducing additional genetic diversity

from the FVW population. This whole cycle was then repeated once to establish the

scd^{**} base population for experimental evolution (Supplementary Figure 1B).

Fitness effects of scd[1]—

To test whether the *scd[1]* allele was deleterious, we tracked changes in the

frequency of the *scd[1]* phenotype in polymorphic populations with the *scd[1]* allele at 0.7

initial frequency. We initiated four replicate populations, each consisting of 70 *scd[**]*

males, 70 *scd[**]* females, 30 FVW males, and 30 FVW females. Populations were

placed in population cages with four culture bottles for mating and oviposition for five

days, after which the flies were discarded and the bottles transferred to fresh cages at

24°C. After adult flies began emerging, they were allowed to mate for three to four days.

The old bottles were then removed, and fresh bottles were placed in the cage for egg

laying. After two days of egg laying, the flies were discarded and the bottles moved to

fresh cages. This cycle was repeated for a total of nine generations.

For each of the first five generations, and at generation nine, we scored male sex

comb phenotypes. 50 males were picked randomly, and the first prothoracic legs from

each male were mounted on glass slides in 70% glycerol/PBS to check for the presence

of an ectopic second sex comb and other abnormalities. While this does not give an

exact measurement of the frequency of the *scd[1]* allele because of this allele's

incomplete penetrance (though penetrance is almost complete in the FVW background;

see below), it should provide a reasonable proxy. Even though reductions in the

frequency of the mutant phenotype could also be driven by selection for suppressor

alleles, this should still give an indication of whether or not the *scd[1]* phenotype is

deleterious.

To test whether there was evidence that the *scd[1]* allele was deleterious (and

decreased in allele frequency) we fit a logistic regression tracking number of *scd[1]* and

wild- type males each generation. As the frequency of *scd^1* at generation 0 was set at

exactly 0.7, we utilized an offset and suppressed the model intercept. Additionally we

checked the results of this model using a logistic mixed model allowing for a variation in

the slope of the response by replicate lineage. Analyses were conducted in R using

glm() and glmer() from the lme4 package.

Experimental evolution—

To test whether sexual selection influences the rate of compensatory adaptation,

we set up two treatments. In the low sexual selection (LSS) treatment, we removed

sexual selection by enforcing monogamous mating. Each generation, we set up 100

vials, each containing one male and one virgin female. After a three-day interaction

period, we anesthetized the flies using CO_2, discarded males, and placed the females in

a population cage with four bottles containing culture media for egg laying. After four

days, the bottles were removed and incubated at 24˚C. When adult flies began eclosing,

we selected virgins for the next generation. Thus, while this treatment did preclude mate

choice, it still allowed for fecundity and viability selection (Arbuthnott and Rundle 2012).

In the high sexual selection (HSS) treatment, we followed a similar protocol

except allowed the opportunity for sexual selection. Each generation, 100 males and

100 virgin females were allowed to interact in a population cage, along with an open

culture bottle for food and moisture. After the three-day interaction period, we placed the

cage in a refrigerator to knock the flies out, and then we sorted males and females.

Males were discarded, and females were placed in fresh cages with four culture bottles

for a four-day egg-laying period. After egg laying, females were discarded, and the

bottles were placed in an environmental chamber at 24˚C until adults began emerging,

at which point we collected virgins for the next generation.

Additionally, we also set up a treatment with low levels of genetic variation (LV)

to test whether compensatory adaptation is limited when segregating genetic variation is

diminished; in other words, testing whether the mutational target size of the

compensatory response was large enough that *de novo* mutations could contribute in

the time frame of the experimental evolution regime. In this treatment, each population

was established from the offspring of a single-pair mating between a randomly chosen

virgin *scd*** female and a randomly chosen *scd*** male. These populations were kept

under the same regime as the HSS treatment.

Finally, we set up a wild-type control (WTC) treatment using wild-type FVW flies.

WTC populations were also maintained under the same regime as the HSS treatment.

These provide a control for lab domestication and unknown aspects of the experimental

protocol.

All populations were initiated using randomly selected *scd*** flies (see above),

except for the LV treatments as described. We set up three replicate populations of

each treatment except for WTC, in which we performed two replicates. Experimental

evolution was conducted for a total of 24 generations. We assayed male sex comb

phenotypes as described earlier at generations 1, 7, 13, 19, and 24, using 30 randomly

selected males from each population at each time point.

To test how male sex comb traits changed over the course of experimental

evolution, we fit generalized linear mixed models using the glmmTMB v0.2.3 (Hadfield

2010) and lme4 v1.1-21 (Bates *et al.* 2015) packages in R (v3.6.1). For sex comb tooth

number for both primary and ectopic/secondary combs, we assumed a Poisson

distribution and used a log link function. The model included generation, treatment, and

their interaction as fixed effects; we also included individual fly, and in some cases

replicate population nested within treatment, as random effects (some models failed to

converge when replicate nested within treatment was included as a random effect). We

also tested for lineage specific zero-inflation in the data, but found no evidence for this,

so excluded this to reduce number of parameters. To test whether the frequency of

defects in the primary sex comb changed over time, we fit a mixed logistic model (sex

comb defects present/absent), again with generation, treatment, and their interaction as

fixed effects, and replicate nested within treatment, as well as individual fly, as random

effects. For all of these models we examined whether a model with a common intercept

(using generation 1 data as the intercept) for each treatment altered estimates relative

to models with varying intercepts. In no case did it substantially alter the conclusions,

and we present both the common intercept model and varying intercept model in results

and supplements. Power simulations were performed using simr v1.0.5 (Green and

MacLeod 2016).

Data availability—

All data and scripts are available on Github

(https://github.com/DworkinLab/Chandler_etal_G3_2020).

Results

Mapping scd[1]—

Though we could not map scd^1 to a specific gene, we were able to further narrow

down its location to an ~85-kb (cytological region 8F8-9A1) region on the X

chromosome through duplication mapping (Table 1, Figure 2). This region contains only

two complete annotated protein-coding genes, *btd* and *Sp1* (both of which influence

aspects of leg development and morphogenesis), and three annotated long non-coding

RNAs, CR42657, CR44016, and CR53498. Interestingly, the two parent strains carrying

the duplications that rescued scd^1 had smaller than average sex combs in the absence

of the scd^1 mutation (Table 1), similar to a past study involving this mutation (Randsholt

and Santamaria 2008), suggesting that the *scd* gene product is a suppressor of sex

comb development.

Influence of genetic background on penetrance and expressivity—

When females of the original scd^1 stock strain were crossed to males of various

wild-type strains to generate males that were hemizygous for scd^1 and heterozygous for

different genetic backgrounds, the penetrance and expressivity of scd^1 varied widely

(Figure 3) demonstrating segregating variation for them. A logistic model using

penetrance (presence/absence of ectopic sex comb) as the response variable with the

genetic background as a fixed effect was a significantly better fit than a null model not

accounting for genetic background ($\chi^2 = 127.6$, df = 18, $p = 5.3 \times 10^{-19}$), and when we fit

a model including genetic background as a random effect, there was substantial among-

strain variance ($\sigma^2 = 4.19$ on the link scale). Similarly, a model with number of ectopic

sex comb teeth as the response variable (expressivity) and genetic background as a

fixed effect was a significantly better fit than the null model ($\chi^2 = 259.0$, df = 18, $p < 1.0 \times$

10^{-10}). The progenitor scd^1 strain from the Bloomington stock center had the lowest

penetrance (frequency of flies exhibiting an ectopic sex comb on the second tarsal

segment) and among the lowest expressivity (number of teeth in the ectopic sex comb).

Some of the other wild-type genetic backgrounds, even in a heterozygous state,

resulted in nearly complete penetrance for scd^1, including the outbred population (FVW)

used for experimental evolution (described below). Interestingly, the FVW outbred

population only showed intermediate levels of expressivity of the mutant phenotype,

consistent with segregating variation in this population. Overall this result suggests

genetic background has a strong impact on the phenotypic expression on scd^1. It also

suggests that the partial penetrance initially observed in the progenitor stock (strain

5070) may reflect the accumulation of suppressor/compensatory mutations in the base

stock center strain itself. These results suggest that a compensatory response could

potentially be due to the accumulation of segregating suppressor alleles.

Fitness effects of scd^1—

The frequency of male flies exhibiting the scd^1 phenotype (in an FVW genetic

background) decreased across five generations of experimental evolution in populations

polymorphic for scd^1 (Figure 4). With a starting allele frequency of 0.7 the frequency

decreased to an average frequency of 0.4 (across the multiple replicates) in males by

generation 9. To test this more rigorously, we fit a logistic model with an offset (starting

frequency of scd^1 = 0.7), and the effect of generation was significant (effect = -0.197 on

logit link scale, s.e. = 0.049, $p = 5.1 \times 10^{-5}$). These results are consistent with the

mutation having moderate deleterious effects.

Experimental evolution—

In the populations carrying the scd^1 allele, defects such as gaps in the primary

sex comb were observed occasionally, and at significantly higher frequencies in the

High Sexual Selection (HSS) populations, and marginally significant frequencies in the

Low Sexual Selection (LSS) populations, than in the wild-type populations. The

frequency of these gaps appeared to decrease in the HSS populations across the 24

generations of the experiment, although the interaction between generation and

treatment was not significant (Figure 5B; Table 2). In addition, the ectopic sex combs

induced by the scd^1 mutation became smaller on average in both the HSS and LSS

treatments (Figure 5C; Table 3), losing on average ~0.5 teeth across the 24 generations

of experimental evolution, though the effect of generation was only marginally significant

in the HSS treatment. This is consistent with some of the compensatory response being

the result of the increase in frequency of naturally occurring suppressor alleles. When

we allowed the intercepts to vary for the HSS and LSS treatments, results were similar,

with a significant main effect for generation, but no significant generation-by-treatment

interaction (Supplementary Tables 1 and 2). Thus we saw no evidence for differences in

rate of compensation between the HSS and LSS treatments, with the magnitude of the

interaction term (change in slope relative to LSS) being much smaller than the effect of

generation (i.e. overall compensatory response). Indeed, the slope of the compensatory

response is weaker for the HSS treatment (relative to LSS), counter to our predictions.

This suggests any additional compensatory effects of sexual selection (i.e. after

accounting for viability and fecundity selection) were relatively weak in this experimental

system. Using a power analysis, we confirmed that the power to detect such an effect

would be very small (Supplementary Figure 2) unless we used a large number of

independent replicate lineages (~30 per treatment), although the power to detect an

effect of this magnitude (assuming it was real) would be approximately 80% with three

replicates if the response continued for 35 or more generations of experimental

evolution (Supplementary Figure 3). As a thought experiment (as this variable is not

under experimental control, but was being estimated), we performed a power analysis

where we modified the magnitude of the slopes of compensatory response (between

LSS and HSS). Assuming the estimated effect is real (just with high degree of

uncertainty due to sampling) the power for our experimental design would be ~18%

(95% CIs 11 - 26.9%). If the difference between the LSS and HSS treatments is of

similar magnitude to that of the overall compensatory response the power increases to

~34% (CIs 24.8 – 44.1%), and reaches power of 76% (CIs 66.4 – 84.0 %) when the

magnitude of the difference between LSS and HSS is 100% greater than the observed

slope.

For the Low genetic Variation (LV) treatment no significant change in number of

ectopic sex comb teeth over time was observed, as expected (generation effect =

0.0017, SD = 0.0044, p = 0.70). No significant changes were observed in the primary

sex comb tooth number across 24 generations in any of the experimental treatments

(Figure 5A; Table 4).

Discussion

Some, but not all, previous work has found that sexual selection may facilitate

populations in purging deleterious mutations (Radwan 2004; Hollis *et al.* 2009;

Jarzebowska and Radwan 2010) or accelerating rate of adaptation (Jacomb *et al.* 2016;

Parrett and Knell 2018). Few studies, however, have addressed whether sexual

selection may facilitate compensatory adaptation, where populations evolve traits to

compensate for the fitness costs of deleterious mutations. While compensatory

adaptation itself is well documented in other systems (Reynolds 2000; Maisnier-Patin *et*

al. 2002; Estes *et al.* 2011; Chandler *et al.* 2012; Comas *et al.* 2012; Chari *et al.* 2017),

in this experiment, we found no effect of the sexual selection regime on the rate of

compensatory adaptation (at least with respect to the mutation's sex comb phenotypes).

This is perhaps surprising, because we found clear evidence of standing genetic

variation modulating the expression of this mutation, so genetic variation does not

appear to be a limiting factor here. Moreover, our experiments show that the *scd^1*

mutation is deleterious (Figure 4), and given the importance of the *Drosophila* sex comb

for male mating success (Ng and Kopp 2008), we expected that the fitness costs of this

mutation would involve male sexual fitness. Thus, we predicted that the costs of this

mutation would be higher in the HSS treatment, in which there was a high opportunity

for female mate choice, than in the LSS treatment, with reduced opportunity for sexual

selection. It is possible that this mutation has effects on other aspects of fitness in males

or females (viability, fecundity), but unfortunately our experiments did not directly

measure specific fitness components. Even so, theoretical work predicts that sexual

selection should act in concert with natural selection because of condition dependence

(Whitlock and Agrawal 2009); that is, mutations that reduce nonsexual fitness should

also reduce mating success, since sexual displays are often indicators of overall

condition. While some work has supported this prediction, our findings add to a growing

body of work suggesting that this is not always the case (Hollis and Houle 2011; Plesnar

et al. 2011; Arbuthnott and Rundle 2012, 2014; Cabral and Holland 2014; Power and

Holman 2015; Chenoweth *et al.* 2015). It is possible that the relatively minor degree of

phenotypic suppression observed here made it difficult to detect differences between

treatments.

Even though sexual selection did not impact the rate of compensatory evolution,

we did observe evidence of weak compensatory adaptation via phenotypic suppression

of the *scd^1* mutation in both the HSS and LSS treatments. On average, the ectopic sex

combs lost about half a tooth (starting with a mean of ~ 3.5 teeth) over the course of 24

generations in these populations; in other words, the expressivity of the mutation

declined slightly. One possible explanation for the similar response in both the high and

low sexual selection treatment is simply that the compensatory response (in terms of

phenotypic suppression) was sufficiently weak that any subtle difference between these

treatments would be difficult to detect given our design. However, the power analyses

(Supplementary Figures 2 and 3) suggest that if an effect of this magnitude were real, it

is sufficiently small that it would require ~25 replicate lineages of each treatment to

detect or a doubling of the number of generations of experimental evolution.

As expected, we did not observe any significant trend in the LV treatment, in

which populations experienced genetic bottlenecks prior to beginning the experiment

(LV populations were treated the same way as HSS populations). Combined with the

observation that genetic background has strong influences on the penetrance and

expressivity of this mutation (Figure 3), this suggests that compensatory adaptation by

phenotypic suppression relies heavily on the presence of standing genetic variation,

rather than rapid accumulation of new mutations. An interesting side note is that the

initial strain (obtained from the Drosophila stock center) carrying scd^1 has among the

lowest penetrance/expressivity for this mutation of all the genetic backgrounds that we

tested. This may suggest that the stock strain has already undergone compensatory

adaptation, and that alleles suppressing the phenotypic expression of scd^1 had become

fixed throughout the maintenance of this stock (which were subsequently removed

when we outcrossed the mutation), though unfortunately we do not have any data on

how long the scd^1 stock strain has been maintained.

While we were unable to map scd^1 to a specific gene, we were able to localize it

to an ~85 kb region containing only a few candidates. The only protein-coding candidate

genes in this region, *Sp1* and *btd*, both have known roles in leg development (Estella

and Mann 2010), but are not specifically known to influence sex comb development.

There are also three long non-coding RNAs in this region (CR42657, CR44016, and

CR43498). Interestingly, all three of these RNAs show evidence of male-specific

expression in modENCODE RNA-seq data available on FlyBase (Graveley *et al.* 2011;

Brown *et al.* 2014). However, CR44016 shows expression only at very low levels and

only in adult males, not pupae or larvae, suggesting it is unlikely to be involved in the

development of sex combs. CR42657 and CR43498 both show expression in pupae

and/or larvae, as well as adult males (but not adult females); however, these RNAs

seem to be expressed in the testis and accessory gland and not other tissues (though

expression in legs specifically was not assessed in the modENCODE dataset). This

suggests that these male-specific reproductive tissues may be driving these sex-specific

expression patterns, not a role in sex comb development. Further work is necessary to

identify the molecular nature of scd^1.

There are a number of important limitations to point out about our study. First,

much of the focus was on compensation by suppression of the phenotypic effects of the

scd^1 mutation on the sex combs directly. While we observed similar levels of phenotypic

compensation with both our high and low sexual selection treatments (LSS and HSS), it

is possible that compensatory evolution differed with respect to the fitness components

(viability, fecundity, and sexual/mating components), which were not evaluated. Thus

we limit our interpretation to the effects on morphological compensation/suppression,

recognizing that we cannot rule out differential patterns of compensatory response for

fitness *per se*. Indeed, this pattern has been observed previously (Pischedda and

Chippindale 2005; Chari *et al.* 2017). Additionally, this experiment was performed over a

relatively short time period (25 generations); if we continued the experiment over a

longer period, subtle differences in the rate of morphological compensation may have

become apparent, as indicated by our power analysis (Supplementary Figure 3).

In summary, we found evidence of moderate compensatory adaptation to a

deleterious mutation by selection for modifier alleles that suppress the mutation's

phenotypic effects. However, while compensatory adaptation did depend on the

presence of standing genetic variation, increased opportunity for sexual selection did

not increase the rate of compensatory response, in spite of the affected phenotype's

known role in mating. Indeed, the HSS treatment showed a slightly weaker (but not

significantly different) rate of compensatory response. However, as shown in the power

analysis, additional replicates, or following these lineages for a longer evolutionary

trajectory would be helpful to reduce the uncertainty due to sampling. Nonetheless, this

adds to a growing body of studies suggesting that sexual selection does not always

enhance natural selection. Future work should tease apart when and why sexual and

natural selection act in concert and when they are likely to operate differently (Martínez-

Ruiz and Knell 2017).

Acknowledgments

We thank Shelby Hemker and Cody Porter for assistance in data collection and fly

maintenance, as well as the editor and two anonymous reviewers for their feedback on

earlier versions of this manuscript. The BEACON Center for the Study of Evolution in

Action (NSF DBI-0939454) provided support for this work. ID was supported by funds

from NSERC (Canada) Discovery and Discovery Accelerator awards.

Figure Legends

Figure 1. (A) Wild-type *Drosophila melanogaster* leg showing a normal male sex comb (black arrow). (B) *scd^1* leg showing the normal primary sex comb (black arrow) as well as a smaller ectopic sex comb on the second tarsal segment (white arrow).

Figure 2. Duplication mapping of *scd^1*. Purple bars represent duplications that rescued the *scd^1* mutant phenotype; pink bars represent duplications that failed to rescue *scd^1*. (A) Entire region of the X chromosome in which duplications were tested. (B) Close-up of the putative *scd^1*-containing region (red box). Because the duplication carried by strain 30333 is sufficient to rescue the *scd^1* phenotype, we hypothesize that the *scd* gene must lie entirely within this region; at the same time, the neighboring duplications (30522 and 30334) did not rescue the *scd^1* phenotype, so *scd* must lie at least partially outside of these regions.

Figure 3. (A) Penetrance and (B) expressivity of the *scd^1* mutation varies among different wild type genetic backgrounds. In these experiments, stock *scd^1* females were crossed to males of different wild-type strains, and phenotypes were scored in F1 males (which were hemizygous for *scd^1* and heterozygous for a different wild genetic background). Penetrance was measured as the proportion of males showing an ectopic sex comb on the second tarsal segment of the prothoracic leg, while expressivity was measured as the number of sex comb teeth on the ectopic sex comb. Error bars indicate 95% confidence intervals.

Figure 4. The *scd^1* mutation is deleterious. Average frequency of *scd^1* over time, across 4 replicate populations, each initialized with 70% *scd^1* males. Error bars indicate 95%

confidence intervals. The shaded region indicates the 95% confidence interval for the

best-fit line, with the starting frequency fixed at 70%.

Figure 5. Plots of model fits for sex comb traits in experimental evolution populations.

(A) Number of teeth in the primary sex comb across 24 generations of experimental

evolution in all experimental treatments. (B) Proportion of male flies with defects in the

primary sex comb in the HSS and LSS treatments; we fit a model with a shared

intercept for these two treatments . (C) Number of teeth in ectopic/secondary sex combs

in males in the HSS and LSS treatments, again with a common intercept. Shaded

regions on plots indicate 95% confidence intervals for the predicted values.

References

Agrawal, A. F., 2001 Sexual selection and the maintenance of sexual reproduction.

Nature 411: 692–695.

Almbro, M., and L. W. Simmons, 2014 Sexual Selection Can Remove an Experimentally

Induced Mutation Load. Evolution 68: 295–300.

Arbuthnott, D., and H. D. Rundle, 2014 Misalignment of natural and sexual selection

among divergently adapted Drosophila melanogaster populations. Animal

Behaviour 87: 45–51.

Arbuthnott, D., and H. D. Rundle, 2012 Sexual Selection Is Ineffectual or Inhibits the

Purging of Deleterious Mutations in Drosophila Melanogaster. Evolution 66:

2127–2137.

Atallah, J., N. H. Liu, P. Dennis, A. Hon, and E. W. Larsen, 2009 Developmental

constraints and convergent evolution in Drosophila sex comb formation.

Evolution & Development 11: 205–218.

Atallah, J., H. Watabe, and A. Kopp, 2012 Many ways to make a novel structure: a new

mode of sex comb development in Drosophilidae. Evolution & Development 14:

476–483.

Bates, D., M. Mächler, B. Bolker, and S. Walker, 2015 Fitting Linear Mixed-Effects

Models Using lme4. Journal of Statistical Software 67: 1–48.

Boube, M., C. Benassayag, L. Seroude, and D. L. Cribbs, 1997 Ras1-Mediated

Modulation of Drosophila Homeotic Function in Cell and Segment Identity.

Genetics 146: 619–628.

Brown, J. B., N. Boley, R. Eisman, G. E. May, M. H. Stoiber *et al.*, 2014 Diversity and

dynamics of the *Drosophila* transcriptome. Nature 512: 393–399.

Cabral, L. G., and B. Holland, 2014 Courtship Song Does Not Increase the Rate of

Adaptation to a Thermally Stressful Environment in a Drosophila melanogaster

Laboratory Population. PLOS ONE 9: e111148.

Chandler, C. H., G. E. Chadderdon, P. C. Phillips, I. Dworkin, and F. J. Janzen, 2012

Experimental Evolution of the Caenorhabditis Elegans Sex Determination

Pathway. Evolution 66: 82–93.

Chandler, C. H., S. Chari, and I. Dworkin, 2013a Does your gene need a background

check? How genetic background impacts the analysis of mutations, genes, and

evolution. Trends in Genetics 29: 358–366.

Chandler, C. H., S. Chari, A. Kowalski, L. Choi, D. Tack *et al.*, 2017 How well do you

know your mutation? Complex effects of genetic background on expressivity,

complementation, and ordering of allelic effects. PLOS Genetics 13: e1007075.

Chandler, C. H., C. Ofria, and I. Dworkin, 2013b Runaway Sexual Selection Leads to

Good Genes. Evolution 67: 110–119.

Chari, S., C. Marier, C. Porter, E. Northrop, A. Belinky *et al.*, 2017 Compensatory

evolution via cryptic genetic variation: Distinct trajectories to phenotypic and

fitness recovery. bioRxiv 200725.

Charlesworth, B., 1978 Model for evolution of Y chromosomes and dosage

compensation. Proceedings of the National Academy of Sciences 75: 5618–

5622.

Chenoweth, S. F., N. C. Appleton, S. L. Allen, and H. D. Rundle, 2015 Genomic

Evidence that Sexual Selection Impedes Adaptation to a Novel Environment.

Current Biology 25: 1860–1866.

Clark, S. C. A., N. P. Sharp, L. Rowe, and A. F. Agrawal, 2012 Relative Effectiveness of

Mating Success and Sperm Competition at Eliminating Deleterious Mutations in

Drosophila melanogaster (S. R. Proulx, Ed.). PLoS ONE 7: e37351.

Comas, I., S. Borrell, A. Roetzer, G. Rose, B. Malla *et al.*, 2012 Whole-genome

sequencing of rifampicin-resistant *Mycobacterium tuberculosis* strains identifies

compensatory mutations in RNA polymerase genes. Nature Genetics 44: 106–

110.

Cook, R. K., M. E. Deal, J. A. Deal, R. D. Garton, C. A. Brown *et al.*, 2010 A New

Resource for Characterizing X-Linked Genes in Drosophila melanogaster:

Systematic Coverage and Subdivision of the X Chromosome With Nested, Y-

Linked Duplications. Genetics 186: 1095–1109.

Davies, A. G., A. Y. Game, Z. Chen, T. J. Williams, S. Goodall *et al.*, 1996 Scalloped

wings Is the Lucilia cuprina Notch Homologue and a Candidate for the Modifier of

Fitness and Asymmetry of Diazinon Resistance. Genetics 143: 1321–1337.

Ellis, E. A., and T. H. Oakley, 2016 High Rates of Species Accumulation in Animals with

Bioluminescent Courtship Displays. Current Biology 26: 1916–1921.

Estella, C., and R. S. Mann, 2010 Non-Redundant Selector and Growth-Promoting

Functions of Two Sister Genes, buttonhead and Sp1, in Drosophila Leg

Development. PLoS Genetics 6: e1001001.

Estes, S., P. C. Phillips, and D. R. Denver, 2011 Fitness Recovery and Compensatory

Evolution in Natural Mutant Lines of C. Elegans. Evolution 65: 2335–2344.

Graveley, B. R., A. N. Brooks, J. W. Carlson, M. O. Duff, J. M. Landolin *et al.*, 2011 The

developmental transcriptome of *Drosophila melanogaster*. Nature 471: 473–479.

Green, P., and C. J. MacLeod, 2016 simr: an R package for power analysis of

generalised linear mixed models by simulation. Methods in Ecology and

Evolution 7: 493–498.

Grieshop, K., J. Stångberg, I. Martinossi-Allibert, G. Arnqvist, and D. Berger, 2016

Strong sexual selection in males against a mutation load that reduces offspring

production in seed beetles. Journal of Evolutionary Biology 29: 1201–1210.

Hadfield, J. D., 2010 MCMC Methods for Multi-Response Generalized Linear Mixed

Models: The MCMCglmm R Package. Journal of Statistical Software 33: 1–22.

Hollis, B., J. L. Fierst, and D. Houle, 2009 Sexual Selection Accelerates the Elimination

of a Deleterious Mutant in Drosophila Melanogaster. Evolution 63: 324–333.

Hollis, B., and D. Houle, 2011 Populations with elevated mutation load do not benefit

from the operation of sexual selection. Journal of Evolutionary Biology 24: 1918–

1926.

Hou, J., G. Tan, G. R. Fink, B. J. Andrews, and C. Boone, 2019 Complex modifier

landscape underlying genetic background effects. PNAS 116: 5045–5054.

Jacomb, F., J. Marsh, and L. Holman, 2016 Sexual selection expedites the evolution of

pesticide resistance. Evolution 70: 2746–2751.

Jarzebowska, M., and J. Radwan, 2010 Sexual Selection Counteracts Extinction of

Small Populations of the Bulb Mites. Evolution 64: 1283–1289.

Kopp, A., 2011 Drosophila sex combs as a model of evolutionary innovations. Evolution

& Development 13: 504–522.

Long, T. A. F., A. F. Agrawal, and L. Rowe, 2012 The Effect of Sexual Selection on

Offspring Fitness Depends on the Nature of Genetic Variation. Current Biology

22: 204–208.

Lumley, A. J., Ł. Michalczyk, J. J. N. Kitson, L. G. Spurgin, C. A. Morrison *et al.*, 2015

Sexual selection protects against extinction. Nature 522: 470–473.

Mackay, T. F. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles *et al.*, 2012 The

Drosophila melanogaster Genetic Reference Panel. Nature 482: 173–178.

Maisnier-Patin, S., O. G. Berg, L. Liljas, and D. I. Andersson, 2002 Compensatory

adaptation to the deleterious effect of antibiotic resistance in Salmonella

typhimurium. Molecular Microbiology 46: 355–366.

Malagón, J. N., A. Ahuja, G. Sivapatham, J. Hung, J. Lee *et al.*, 2014 Evolution of

Drosophila sex comb length illustrates the inextricable interplay between

selection and variation. Proceedings of the National Academy of Sciences 111:

E4103–E4109.

Martínez-Ruiz, C., and R. J. Knell, 2017 Sexual selection can both increase and

decrease extinction probability: reconciling demographic and evolutionary

factors. Journal of Animal Ecology 86: 117–127.

Masta, S. E., and W. P. Maddison, 2002 Sexual selection driving diversification in

jumping spiders. Proceedings of the National Academy of Sciences 99: 4442–

4447.

McGuigan, K., D. Petfield, and M. W. Blows, 2011 Reducing Mutation Load Through

Sexual Selection on Males. Evolution 65: 2816–2829.

McKenzie, J. A., and G. Clarke, 1988 Diazinon Resistance, Fluctuating Asymmetry and

Fitnessin the Australian Sheep Blowfly, Lucilia cuprina. Genetics 120: 213–220.

Mullis, M. N., T. Matsui, R. Schell, R. Foree, and I. M. Ehrenreich, 2018 The complex

underpinnings of genetic background effects. Nat Commun 9: 1–10.

Ng, C. S., and A. Kopp, 2008 Sex Combs are Important for Male Mating Success in

Drosophila melanogaster. Behav Genet 38: 195.

Parrett, J. M., and R. J. Knell, 2018 The effect of sexual selection on adaptation and

extinction under increasing temperatures. Proceedings of the Royal Society B:

Biological Sciences 285: 20180303.

Pischedda, A., and A. Chippindale, 2005 Sex, mutation and fitness: asymmetric costs

and routes to recovery through compensatory evolution. Journal of Evolutionary

Biology 18: 1115–1122.

Plesnar, A., M. Konior, and J. Radwan, 2011 The role of sexual selection in purging the

genome of induced mutations in the bulb mite. Evolutionary Ecology Research

13: 209–216.

Power, D. J., and L. Holman, 2015 Assessing the alignment of sexual and natural

selection using radiomutagenized seed beetles. Journal of Evolutionary Biology

28: 1039–1048.

Radwan, J., 2004 Effectiveness of sexual selection in removing mutations induced with

ionizing radiation. Ecology Letters 7: 1149–1154.

Randsholt, N. B., and P. Santamaria, 2008 How Drosophila change their combs: the

Hox gene Sex combs reduced and sex comb variation among Sophophora

species. Evolution & Development 10: 121–133.

Reynolds, M. G., 2000 Compensatory Evolution in Rifampin-Resistant Escherichia coli.

Genetics 156: 1471–1481.

Rowe, L., and D. Houle, 1996 The lek paradox and the capture of genetic variance by

condition dependent traits. Proceedings of the Royal Society of London. Series

B: Biological Sciences 263: 1415–1421.

Santamaria, P., and N. B. Randsholt, 1995 Characterization of a region of the X

chromosome of Drosophila including multi sex combs (mxc), a Polycomb group

gene which also functions as a tumour suppressor. Molec. Gen. Genet. 246:

282–290.

Sharp, N. P., and A. F. Agrawal, 2008 Mating Density and the Strength of Sexual

Selection Against Deleterious Alleles in Drosophila Melanogaster. Evolution 62:

857–867.

Venken, K. J. T., E. Popodi, S. L. Holtzman, K. L. Schulze, S. Park *et al.*, 2010 A

Molecularly Defined Duplication Set for the X Chromosome of Drosophila

melanogaster. Genetics 186: 1111–1125.

Whitlock, M. C., and A. F. Agrawal, 2009 Purging the Genome with Sexual Selection:

Reducing Mutation Load Through Selection on Males. Evolution 63: 569–582.

Table 1. Duplication mapping of the scd^1 allele. Homozygous female scd^1 flies were crossed to males carrying a segment

of the X chromosome duplicated onto either the Y chromosome or chromosome III. Assuming scd^1 is a recessive loss-of-

function allele, if the male progeny of these crosses show a mutant phenotype, then the duplication does not rescue scd^1

and scd must lie at least partially outside the duplication. If none of the male offspring show a mutant phenotype, then

scd^1 is rescued and scd lies within the duplication.

Duplication Stock #	Symbol	Duplicated region	Rescues scd^1?
29759	Dp(1;Y)BSC35	X:8192725-8271204;X:9030055;Y & X:1;X:493529;Y	No; replicated 2x
30522	Dp(1;Y)BSC146	X:8714331-8897281;X:9686653;Y & X:1;X:493529;Y	No; replicated 2x
29782	Dp(1;Y)BSC58	X:9355691-9500067;X:10744934;Y & X:1;X:493529;Y	Yes; replicated 2x; note: duplication on its own has reduced sex comb teeth in primary comb (mean=8.2, sd=1.3, SE=0.4, N=11)
29783	Dp(1;Y)BSC59	X:10085520-10218380;X:10744934;Y & X:1;X:493529;Y	No
29784	Dp(1;Y)BSC60	X:10331363-10385547;X:10744934;Y & X:21572099-22456281;h28-h29;Y & X:1;X:493529;Y	No
29786	Dp(1;Y)BSC62	X:10460679-10601454;X:10744934;Y & X:1;X:493529;Y	No
29788	Dp(1;Y)BSC64	X:10601454-10738950;X:10744934;Y & X:1;X:493529;Y & X:21204834-21318903;h28-h29;Y	No
30333	Dp(1;3)DC212	X:9677341;X:9784700;3L:6442676 (r6, Dp)	Yes; note: duplication on its own has small sex combs for primary comb (3-5 teeth)
30334	Dp(1;3)DC213	X:9762229;X:9807047;3L:6442676 (r6, Dp)	No

8   **Table 2.** Effect estimates for logistic mixed model for gaps in the primary sex comb as response variable. This model had

9   included only the HSS and LSS treatments, and these two treatments were constrained to have a common intercept, but

10   not slope. The confidence intervals for the effect of generation in each treatment overlap.

	Estimate	Standard Error	P
Intercept	-2.30	0.25	$< 2.0 \times 10^{-16}$
generation-HSS	-0.034	0.021	0.106
generation-LSS	-0.026	0.020	0.188

Table 3. Effect estimates for generalized linear model (poisson) using number of teeth in the ectopic/secondary sex comb

as response variable. This model included only the HSS and LSS populations; WTC populations were excluded because

they did not display ectopic sex combs, and because the purpose of this model was to test for a difference between the

HSS and LSS treatments specifically, we considered the LV treatment on its own separately. The HSS and LSS

treatments were constrained to have a common intercept in this model, but were allowed to have different slopes. The

confidence intervals for the effect of generation in each treatment overlap.

	Estimate	Standard Error	P
Intercept	1.31	0.066	$< 2.0 \times 10^{-16}$
generation-HSS	-6.3×10^{-3}	3.7×10^{-3}	0.085
generation-LSS	-0.011	3.8×10^{-3}	0.005

21   **Table 4.** Effect estimates for generalized linear mixed model for primary sex comb tooth number as response variable;

22   generation, treatment, and their interaction as fixed effects; and individual fly as a random effect (models including

23   replicate population nested within treatment failed to converge).

	Estimate	Standard Error	P
Intercept	2.41	0.035	$< 2 \times 10^{-16}$
generation	4.4×10^{-4}	2.2×10^{-3}	0.85
treatment-HSS	4.2×10^{-4}	0.046	0.99
treatment-LSS	0.017	0.045	0.71
treatment-LV	6.9×10^{-3}	0.047	0.88
generation x treatment-HSS	5.2×10^{-4}	2.9×10^{-3}	0.86
generation x treatment-LSS	1.3×10^{-3}	2.9×10^{-3}	0.66
generation x treatment-LV	1.4×10^{-3}	3.0×10^{-3}	0.64

A

1.0 mm

B

1.0 mm

1 **Supplementary Table 1.** Effect estimates for logistic mixed model for gaps in the primary sex comb as response variable;

2 generation, treatment, and their interaction as fixed effects; and replicate nested within treatment, and individual fly as

3 random effects. This model included only the HSS and LSS treatments, as the WTC treatment was not expected to have

4 appreciable frequencies of sex comb defects, and the LV treatment was considered separately.

	Estimate	Standard Error	*P*
Intercept	-1.98	0.33	1.4×10^{-9}
generation	-0.054	0.026	0.037
treatment-LSS	-0.66	0.50	0.19
generation x treatment-LSS	0.048	0.036	0.18

7 **Supplementary Table 2.** Effect estimates for logistic mixed model for number of teeth in the ectopic sex comb as

8 response variable; generation, treatment, and their interaction as fixed effects; and replicate nested within treatment, and

9 individual fly as random effects. This model included only the HSS and LSS treatments, as the WTC treatment was not

10 expected to have ectopic sex combs, and the LV treatment was considered separately.

	Estimate	Standard Error	P
Intercept	1.37	0.089	$< 2 \times 10^{-16}$
generation	-0.0077	0.0038	0.0489
treatment-LSS	-0.12	0.13	0.34
generation x treatment-LSS	-1.5×10^{-3}	5.6×10^{-3}	0.79

Supplementary Figure 1. Crossing scheme for introgression of scd^1 into the FVW wild-

type genetic background. (A) Initial introgression of scd^1 into FVW to establish the scd^*

population. (B) Further backcrossing using larger numbers of flies to introduce

additional genetic diversity from FVW to establish the scd^{**} base population for

experimental evolution. See Methods section for detailed description of introgression

scheme.

A

y^1 scd^1 ras^1 v^1 f^1

y^1 scd^1 ras^1 v^1 f^1

y^1 scd^1 ras^1 v^1 f^1

Select male offspring with scd phenotype
and wild-type eyes

y^1 scd^1

y^1 scd^1

Select male offspring with scd phenotype
and wild-type body color

scd^1

First female offspring to emerge

Maintain late-emerging female
offspring at 18°C

scd^1

Select male offspring with
scd phenotype

scd^1

scd^1

Individual sibling pair
crosses between males
with scd phenotype and
females of unknown
scd genotype

Females in these crosses
are either:
scd^1/scd^1 or scd^1/scd$^+$

scd^1

scd^1

?

scd^1

scd^1

scd^1

Keep crosses that yield only scd males to
establish scd* population

B

scd** population

Supplementary Figure 2. Statistical power to detect a significant effect of sexual

selection regime on compensatory adaptation as a function of number of replicate

lineages in each treatment, holding all other aspects of the experiment constant. This

analysis assumes the magnitude of the effect of sexual selection regime is similar to

what we actually observed. At least 25 replicate lineages are necessary to detect such

an effect 80% of the time.

Supplementary Figure 3. Statistical power to detect a significant effect of sexual

selection regime on compensatory adaptation as a function of number of generations of

experimental evolution, holding all other aspects of the experiment constant. This

analysis assumes the magnitude of the effect of sexual selection regime is similar to

what we actually observed. Approximately 35 generations are necessary to detect such

an effect 80% of the time.

23